



Group connectivity and group coloring

Small groups versus large groups

Langhede, Rikke; Thomassen, Carsten

Published in:

Electronic Journal of Combinatorics

Link to article, DOI:

[10.37236/8984](https://doi.org/10.37236/8984)

Publication date:

2020

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Langhede, R., & Thomassen, C. (2020). Group connectivity and group coloring: Small groups versus large groups. *Electronic Journal of Combinatorics*, 27(1), Article P1.49. <https://doi.org/10.37236/8984>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Group connectivity and group coloring: small groups versus large groups

Rikke Langhede Carsten Thomassen*

Department of Applied Mathematics and Computer Science
Technical University of Denmark
DK-2800 Lyngby, Denmark

rimla@dtu.dk ctho@dtu.dk

Submitted: Aug 30, 2019; Accepted: Feb 4, 2020; Published: Mar 6, 2020
© The authors. Released under the CC BY-ND license (International 4.0).

Abstract

A well-known result of Tutte says that if Γ is an Abelian group and G is a graph having a nowhere-zero Γ -flow, then G has a nowhere-zero Γ' -flow for each Abelian group Γ' whose order is at least the order of Γ . Jaeger, Linial, Payan, and Tarsi observed that this does not extend to their more general concept of group connectivity. Motivated by this we define $g(k)$ as the least number such that, if G is Γ -connected for some Abelian group Γ of order k , then G is also Γ' -connected for every Abelian group Γ' of order $|\Gamma'| \geq g(k)$. We prove that $g(k)$ exists and satisfies for infinitely many k ,

$$(2 - o(1))k < g(k) \leq 8k^3 + 1.$$

The upper bound holds for all k . Analogously, we define $h(k)$ as the least number such that, if G is Γ -colorable for some Abelian group Γ of order k , then G is also Γ' -colorable for every Abelian group Γ' of order $|\Gamma'| \geq h(k)$. Then $h(k)$ exists and satisfies for infinitely many k ,

$$(2 - o(1))k < h(k) < (2 + o(1))k \ln(k).$$

The upper bound (for all k) follows from a result of Král', Pangrác, and Voss. The lower bound follows by duality from our lower bound on $g(k)$ as that bound is demonstrated by planar graphs.

Mathematics Subject Classifications: 05C15, 05C21

*Supported by Independent Research Fund Denmark, 8021-002498 AlgoGraph.

1 Introduction

Tutte's 5-Flow Conjecture states that any 2-edge-connected graph has a nowhere-zero 5-flow (see e.g. [1, 4, 6, 20]). Seymour [15] proved that every 2-edge-connected graph has a nowhere-zero 6-flow.

Jaeger et al. [5] introduced the concept of group-connectivity and proved that every 3-edge-connected graph is Γ -connected for any Abelian group Γ of size $|\Gamma| \geq 6$. This extends the 6-flow theorem since that theorem is easily reduced to the 3-connected case as pointed out by Seymour [15].

Tutte [18] (see also [20]) proved that if Γ is an Abelian group and G is a graph having a nowhere-zero Γ -flow, then G has a nowhere-zero Γ' -flow for each Abelian group Γ' whose order is at least the order of Γ . Jaeger et al. [5] observed that this does not extend to their more general concept of group connectivity. We prove that the statement becomes true if Γ' is sufficiently large compared to Γ , as explained in the Abstract.

If Γ is an Abelian group and G is a planar 2-edge-connected graph, then G is Γ -connected if and only if the dual graph G^* is Γ -colorable. This duality was part of the motivation for these concepts [5]. This suggests the definition of $g(k)$, $h(k)$ in the Abstract, and we prove:

$$(2 - o(1))k < g(k) \leq 8k^3 + 1, \tag{1}$$

and

$$(2 - o(1))k < h(k) < (2 + o(1))k \ln(k). \tag{2}$$

The lower bounds hold for infinitely many k , and we do not know if g, h are monotone. The upper bounds hold for all k . The upper bound on $h(k)$ follows from a result of Král' et al. [7]. The lower bound on $h(k)$ follows from the lower bound of $g(k)$ as that bound is demonstrated by planar graphs.

It was proved in [11] that $g(3) = 3$. We conjecture that $h(3) = 3$ and prove that $h(3) \leq 5$.

The *group chromatic number* (see e.g. [7, 10]) $\chi_g(G)$ is the smallest number k such that G is Γ -colorable for every Abelian group Γ of order at least k . We point out that there is another possible definition, namely the *weak group chromatic number* $\chi_{wg}(G)$ which is the smallest number k such that G is Γ -colorable for *some* Abelian group Γ of order k . Clearly

$$\chi(G) \leq \chi_{wg}(G) \leq \chi_g(G). \tag{3}$$

Thus our lower bound on $h(k)$ shows that $\chi_g(G)$ may be almost twice as big as $\chi_{wg}(G)$.

The *group connectivity number* (see e.g. [9]) $\Lambda_g(G)$ is the smallest number k such that G is Γ -connected for every Abelian group Γ of order at least k . Similarly, we define the *weak group connectivity number* $\Lambda_{wg}(G)$ which is the smallest number k such that G is Γ -connected for *some* Abelian group Γ of order k .

Our lower bound on $g(k)$ shows that $\Lambda_g(G)$ may be almost twice as big as $\Lambda_{wg}(G)$.

2 Group connectivity

We use essentially the terminology and notation in [1]. We allow a graph to have multiple edges but no loops. If v is a vertex in a directed graph, then $E^+(v)$ (respectively $E^-(v)$) denotes the set of edges going out from v (respectively going into v). Jaeger et al. [5] introduced the concept of group connectivity as follows.

Definition 1. Let Γ be an Abelian group. The graph G is said to be Γ -connected if the following holds: Given some orientation D of G and any function $\beta : V(G) \rightarrow \Gamma$ satisfying $\sum_{v \in V(G)} \beta(v) = 0$, there exists a function $f : E(G) \rightarrow \Gamma$ such that $\sum_{e \in E^+(v)} f(e) - \sum_{e \in E^-(v)} f(e) = \beta(v)$ for all $v \in V(G)$ and such that $f(e) \neq 0$ for all $e \in E(G)$.

Note that the direction of an edge is not important. Indeed, we may replace “some orientation” by “any orientation” in the definition because we may replace $f(e)$ by $-f(e)$, if necessary.

Jaeger et al. [5] gave the following criterion for group connectivity in terms of forbidden flow values.

Theorem 2. *The graph G is Γ -connected if and only if the following holds: Given any orientation D of G and any function $\varphi : E(G) \rightarrow \Gamma$, there exists a function $f : E(G) \rightarrow \Gamma$ which has $f(e) \neq \varphi(e)$ for all $e \in E(G)$ and $\sum_{e \in E^+(v)} f(e) - \sum_{e \in E^-(v)} f(e) = 0$ for all $v \in V(G)$.*

The function f in Theorem 2 is called a *flow*, and $f(e)$ is called a *flow value*. $\varphi(e)$ is called a *forbidden flow value*.

3 A lower bound for g

We shall use the following lemma which is an easy exercise. A proof can be found in [5].

Lemma 3. *Let P be a cyclic group of prime order, let S be a non-empty proper subset of P , and let T be a subset of P which contains at least two elements. Then $|S + T| > |S|$.*

Let q, k be natural numbers. We define $G_{q,k}$ as the graph consisting of two vertices s, t connected by q internally disjoint paths of length k .

Theorem 4. *Let k be a natural number, and let p be the smallest prime $> 2^{k-1} + 1$. Then $G_{q,2^{k-1}}$ is not \mathbb{Z}_2^k -connected for any odd q . $G_{q,2^{k-1}}$ is \mathbb{Z}_p -connected when $q \geq p$.*

Proof. We first prove that $G = G_{q,2^{k-1}}$ is not \mathbb{Z}_2^k -connected. By reversing directions, if necessary, we may assume that all edges are directed towards t . Let $\varphi : E(G) \rightarrow \mathbb{Z}_2^k$ be a function such that each path forbids all the 2^{k-1} elements having an even number of 1's. As the flow values on a directed path has to be the same on all edges of the path, the forbidden values imply that the flow value of any of the paths must have an odd number of 1's. Since G consists of an odd number of paths which all have a flow with an odd

number of 1's, the in-flow in t (or, similarly, out-flow in s) can never sum to 0. Thus, G is not \mathbb{Z}_2^k -connected.

To prove that G is \mathbb{Z}_p -connected we consider any function $\varphi : E(G) \rightarrow \mathbb{Z}_p$, and again, we let all edges be directed towards t . The goal is to find a flow $f : E(G) \rightarrow \mathbb{Z}_p$ such that $f(e) \neq \varphi(e)$ for all $e \in E(G)$. As noted above f must have the same value on all edges of a path between s and t , and as each path has length 2^{k-1} , it has at least $p - 2^{k-1} \geq 3$ possible flow values of \mathbb{Z}_p . It remains to check that we can choose the values of f on each path such that the sum is 0 in s and t . Given $q \geq p$ subsets of \mathbb{Z}_p of size at least 2, it follows from Lemma 3 that the sum of these contains all elements of \mathbb{Z}_p , in particular the sum contains 0. Thus we can choose f such that $f(e) \neq \varphi(e)$ for all $e \in E(G)$ and the sum in s and t is 0. \square

Corollary 5. *Given any $\epsilon > 0$ there exists an infinite number of graphs which are Γ -connected for some group Γ of prime order, but not Γ' -connected for some group Γ' satisfying $|\Gamma'| = (2 - \epsilon)|\Gamma|$. Hence $g(k) > (2 - o(1))k$ for infinitely many primes k .*

Proof. Let p_n denote the n 'th prime, and let $g_n = p_{n+1} - p_n$. The Prime Number Theorem implies that for any $\epsilon > 0$ there exists a natural number N such that for any $n \geq N$, $g_n < \epsilon p_n$. Now choose n, k such that $N < p_n < 2^k$, and furthermore, p_n is the largest prime $< 2^k$. Then

$$p_{n+1} < 2^k + g_n < (1 + \epsilon)2^k.$$

Put $p = p_{n+1}$. If q is any odd number $\geq p$ then, by Theorem 4, $G_{q,2^k}$ is \mathbb{Z}_p -connected, but not \mathbb{Z}_2^{k+1} -connected. It follows that $g(p) > 2(1 - \epsilon)p$. \square

There is a slight inaccuracy in the proof above, namely when $p = p_{n+1} = 2^k + 1$, that is, p is a Fermat prime. But this can happen only if k is a power of 2 which does not affect the correctness of Corollary 5.

4 The cyclicity of a graph

Definition 6. Let G be a 2-edge-connected graph. We say that two edges e_1, e_2 are *cycle-equivalent* if every cycle that contains one of e_1, e_2 also contains the other. It is easy to see that this defines an equivalence relation on $E(G)$ and that two distinct edges e_1, e_2 are cycle-equivalent if and only if the two edges form a *2-edge-cut*, that is, $G - e_1 - e_2$ is disconnected. We define the *cyclicity* of G , denoted $q(G)$, to be the size of a largest equivalence class. In particular, if G has no 2-edge-cuts (i.e. G is 3-edge-connected), then $q(G) = 1$.

The following result follows from Proposition 3.2 and Lemma 3.3 in [8]. For the sake of completeness we include a proof.

Proposition 7. *If G is Γ -connected, then $|\Gamma| > q(G)$.*

Proof. Suppose G is Γ -connected. Let $e_1, \dots, e_{q(G)}$ be the edges in a largest cycle-equivalence class. Let C be a cycle which contains one, and hence all of $e_1, \dots, e_{q(G)}$. By reversing directions, if necessary, we may assume that all edges in $e_1, \dots, e_{q(G)}$ have the same direction when we traverse C . For any flow, all edges in $e_1, \dots, e_{q(G)}$ have the same flow value. We now use Theorem 2. If $|\Gamma| \leq q(G)$ we can define $\varphi : E(G) \rightarrow \Gamma$ such that it is surjective on $e_1, \dots, e_{q(G)}$, that is, all elements in Γ are forbidden on the edges $e_1, \dots, e_{q(G)}$. Hence $|\Gamma| > q(G)$. \square

We shall prove the following (see Theorem 17).

Theorem 8. *If G is 2-edge-connected, then G is Γ -connected for any Abelian group Γ of order $|\Gamma| > 8q(G)^3$.*

We can now combine Proposition 7 and Theorem 8 to get:

Corollary 9. *If G is Γ -connected for some Abelian group Γ , then G is Γ' -connected for any Abelian group Γ' of order $|\Gamma'| > 8|\Gamma|^3$.*

5 Flows in 3-edge-connected graphs with multiple forbidden flow-values

We shall use the following definition and theorems by Jaeger et al. [5]:

Definition 10. A 2-constructible graph G is defined recursively as follows.

- (i) The graph with one vertex (and no edges) is 2-constructible.
- (ii) If G_1, G_2 are 2-constructible, then the disjoint union of G_1 and G_2 together with two new edges joining them is 2-constructible.

Jaeger et al. [5] proved the following.

Theorem 11. *Let G be a cubic 3-edge-connected graph and let v be a vertex in G . Define $H = G - v$. Then H has a spanning tree T such that the contraction of the edges of H which are not in T yields a 2-constructible graph.*

They used it to prove the following.

Theorem 12. *Let G be a 3-edge-connected graph and let v be a vertex of degree 3 in G . Then $G - v$ is Γ -connected for any Abelian group Γ of order $|\Gamma| \geq 6$.*

We shall also use the following definition and lemma.

Definition 13. Given a finite subset $\Pi = \{a_1, a_2, \dots, a_k\}$ of an Abelian group Γ , we define the *simple sum* Π' of Π to be the set of all elements on the form

$$\alpha_1 a_1 + \alpha_2 a_2 + \dots + \alpha_k a_k$$

where $\alpha_i \in \{0, \pm 1\}$ for $1 \leq i \leq k$. In particular, Π' contains 0.

Lemma 14. *Given a natural number k and an Abelian group Γ of order $|\Gamma| > k$, there exists a subset $\Pi \subseteq \Gamma$ which is closed under taking inverses and has $|\Pi| = k$ such that the simple sum Π' of Π satisfies $|\Pi'| \leq k^2$.*

Proof. For $k = 1$ we let Π consist of 0, so let $k > 1$. If some element a in Γ has order at least k , then we put $\Pi = \{\pm a, \pm 2a, \dots, \pm \frac{k}{2}a\}$ if k is even, and $\Pi = \{0, \pm a, \pm 2a, \dots, \pm \frac{k-1}{2}a\}$ if k is odd. So assume all elements have order $< k$.

Let Γ_0 be the largest subgroup of order $< k$. Let $a \in \Gamma \setminus \Gamma_0$. Then the subgroup generated by $\Gamma_0 \cup \{a\}$ has order greater than k but less than k^2 and any k -subset closed under taking inverses can play the role of Π . \square

We use these results to prove the following:

Theorem 15. *Let G be a 3-edge-connected graph and let v be a vertex of degree 3 in G . Define $H = G - v$. Assume each edge of H has a direction. Let $k \in \mathbb{N}$, and let Γ be any Abelian group of order $|\Gamma| > 8k^3$. Assume that for each edge e , F_e is a set of at most k elements in Γ . Then there exists a flow $f : E(H) \rightarrow \Gamma$ such that $f(e) \notin F_e$ for all $e \in E(G)$.*

We say that F_e is the set of *forbidden flow values* for e . If it becomes convenient to reverse the orientation of an edge e we replace F_e by $-F_e = \{-\gamma \mid \gamma \in F_e\}$.

Proof. By Lemma 14 there exist subsets $\Pi, \Pi' \subseteq \Gamma$ such that Π is closed under inverses and has $|\Pi| = 2k$ and Π' is the simple sum of Π and has $|\Pi'| \leq 4k^2$.

It suffices to prove Theorem 15 in the case where G is cubic. For if u is a vertex of degree $d > 3$, then we replace u by a cycle of length d such that all vertices in that cycle have degree 3 in the resulting graph. (If u is a cutvertex, we can make sure that no two edges of the cycle form a 2-edge-cut.) So assume that G is cubic and 3-edge-connected. The case $k = 1$ follows from Theorem 2 and Theorem 12, so we may assume that $k > 1$.

By Theorem 11, $H = G - v$ has a spanning tree T in H such that the contraction of the edges of H which are not in T yields a 2-constructible graph. We colour the edges of T red and colour the edges in $H - T$ blue. Let E_b be the set of blue edges.

As H/E_b is 2-constructible there exists a sequence H_0, H_1, \dots, H_t of graphs such that H_0 is the empty graph on $|V(H/E_b)|$ vertices, $H_t = H/E_b$, and each H_i is obtained from H_{i-1} by adding two red edges (which we denote by e_i and e'_i) between two disjoint connected components of H_{i-1} .

We first describe informally the method, which is similar to the method in [5]. We give H flow in two steps. In the first step we give all red edges flow values which are non-forbidden and which remain to be non-forbidden after Step 2. Step 1 also affects the blue edges but this is not important. In Step 2 we send flow through the blue edges such that the flow values of the blue edges are non-forbidden. There is already flow through the blue edges after Step 1. The additional flow values (added in Step 2) in the blue edges will be in Π . This will then affect the red edges in such a way that the additional flow through a red edge will be in Π' . In Step 1, the flow is chosen such that an additional flow value in Π' will keep the flow value in the red edge non-forbidden.

We now argue formally.

Step 1: Consider a cycle C'_t in $H_t = H/E_b$ through e_t and e'_t and let C_t be a cycle in H containing the edges of C'_t and no other red edges. If the orientations of e_t and e'_t do not agree on C_t then reverse the orientation of e_t and replace F_e by $-F_e$. Let $F = F_{e_t} \cup F_{e'_t}$. We pick flow γ_t to send through C_t such that:

$$(\gamma_t + \Pi') \cap F = \emptyset. \quad (4)$$

Since Π' has size at most $4k^2$, F has size at most $2k$, and γ_t can be chosen in more than $8k^3$ ways, this is indeed possible. Modify the set of forbidden flow values in each red edge e of $C_t - e_t - e'_t$ such that the new set of forbidden flow values is $F_e - \gamma_t = \{\alpha - \gamma_t \mid \alpha \in F_t\}$ if the orientation of e agrees with that of e_t , and $F_e + \gamma_t = \{\alpha + \gamma_t \mid \alpha \in F_t\}$ if not. We call this the *first iteration* of Step 1.

Next, we consider a cycle C'_{t-1} in $H/E_b - e_t - e'_t$ through e_{t-1} and e'_{t-1} and let C_{t-1} be a cycle in $H - e_t - e'_t$ containing the edges of C'_{t-1} and no other red edges. As above we find an appropriate flow γ_{t-1} to send through C_{t-1} . Note that e_t and e'_t are not in C_{t-1} (or any other C_i for $1 \leq i \leq t-1$) by the construction of H/E_b , so the flow in e_t, e'_t will not be changed in Step 1. We call this the *second iteration* of Step 1. We repeat this argument for e_{t-2} and e'_{t-2} and then e_{t-3} and e'_{t-3} , etc., until all red edges have received a flow value. Once e_i, e'_i have received a flow value in the $(t+1-i)$ 'th iteration of Step 1, that flow value will not be further changed in Step 1.

Step 2: If all blue edges have a non-forbidden flow after Step 1, the proof is complete and there will be no Step 2. So consider a blue edge e which has a forbidden flow value $f'(e)$, say, after Step 1. Since $|\Pi| = 2k$ and F_e has at most k elements, there exists a $\gamma \in \Pi$ such that $f'(e) + \gamma$ and $f'(e) - \gamma$ are both non-forbidden, that is, they are both in $\Gamma \setminus F_e$. We say that γ is *good for e* . Let E_γ be the set of all edges which currently have forbidden flow values and for which γ is good. For each edge in E_γ , let C_e be the unique cycle in $T + e$. Form the symmetric difference H_γ of C_e taken over all edges e in E_γ . Then H_γ is an even graph (that is, a graph where each component is Eulerian), and hence H_γ has a flow using only γ and $-\gamma$. We add this flow to the flow obtained after Step 1, and now all edges in E_γ have non-forbidden flow values.

Repeat this step as long as there are blue edges with forbidden flow values.

Consider now a red edge e after Step 2. Let $f'(e)$ be its flow value after Step 1. In Step 2 we add some elements in Π to $f'(e)$. Thus the final flow of e is of the form $f'(e) + \gamma'$ where γ' is a simple sum of elements from Π , that is $\gamma' \in \Pi'$. By the choice of $f'(e)$ it follows that the final flow value $f'(e) + \gamma' \notin F_e$, as required. (In Step 1 we modified the forbidden flow values in the red edges. Here in Step 2, F_e denotes the original forbidden flow values.) \square

Theorem 16 below follows from Theorem 15 by adding a vertex of degree 3 to G which may be removed again.

Corollary 16. *Let G be a 3-edge-connected graph. Let $k \in \mathbb{N}$, and let Γ be any Abelian group of order $|\Gamma| > 8k^3$. Given any orientation of G , if G has at most k forbidden flow values from Γ in each edge, then there exists a flow $f : E(G) \rightarrow \Gamma$ such that $f(e)$ is not forbidden for any $e \in E(G)$.*

6 Flows in 2-edge-connected graphs with forbidden flow-values

Now we can use Theorem 16 to prove a similar statement about 2-edge-connected graphs.

Theorem 17. *Let G be a 2-edge-connected graph. Let $k \in \mathbb{N}$, and let Γ be an Abelian group of order $|\Gamma| > 8(kq(G))^3$. Given any orientation of G , if G has at most k forbidden flow values from Γ in each edge, then there exists a flow $f : E(G) \rightarrow \Gamma$ such that $f(e)$ is not forbidden for any $e \in E(G)$.*

Proof. We may assume that the orientation of G is strongly connected by reversing the edges necessary and adjusting the forbidden sets in the edges accordingly. Then all edges in a cycle-equivalence class have the same direction in each cycle containing them. We form a new directed graph G' on the same edge set as G (but possibly with a different vertex set) such that

- (i): the edge set of any cycle in G is also the edge set of a cycle in G' and vice versa (in particular G and G' have the same cycle-equivalence classes).
- (ii): the orientations of all edges agree in any two cycles with the same edge set in G and G' , respectively. More precisely: If E is the edge set of a cycle in G and hence also a cycle in G' , then we can choose a clockwise orientation of C, C' such that an edge e is directed clockwise in C if and only if e is directed clockwise in C' .
- (iii): every Γ -flow in G is also a Γ -flow in G' and vice versa.
- (iv): the edges of each cycle-equivalence class in G' form a directed path such that each intermediate vertex has indegree 1 and outdegree 1.

Note that (iv) is equivalent with the following:

- (v): If we delete the edges of a cycle-equivalence class in G' , then the resulting graph has precisely one component with edges.

If G satisfies (iv), we put $G' = G$. Otherwise, there exists a cycle-equivalence class such that the deletion of its edges results in a graph with more than one component. Let G_1 be one component. Let G_2 consist of all other components containing edges together with those paths in the cycle-equivalence class that connect them. Then the edges in the cycle-equivalence class that are not in $G_1 \cup G_2$ form two directed paths P_1 with vertices v_1, v_2, \dots, v_s and directed edges e_1, e_2, \dots, e_{s-1} and P_2 with vertices u_1, u_2, \dots, u_t and directed edges $e'_1, e'_2, \dots, e'_{t-1}$ where $v_1, u_t \in V(G_1)$ and $u_1, v_s \in V(G_2)$. The two paths P_1, P_2 are disjoint except that possibly $v_1 = u_t$ and possibly $u_1 = v_s$. Only their

endvertices are in $G_1 \cup G_2$. Now we form a new graph H from $G_1 \cup G_2$ by first identifying v_1, v_s and then adding a directed path with edges $e_1, e_2, \dots, e_{s-1}, e'_1, e'_2, \dots, e'_{t-1}$ from u_1 to u_t . Then H also satisfies (i),(ii),(iii). Moreover, H has more vertices of indegree 1 and outdegree 1 than G . So, in a finite number of steps we obtain a graph satisfying (i),(ii),(iii),(iv).

It follows that G' is a subdivision of a 3-edge-connected graph G'' . An edge in G'' corresponds to a cycle-equivalence class in G and is therefore subdivided into at most $q(G)$ edges, by the definition of $q(G)$. Now we complete the proof by applying Theorem 16 to G'' . As each edge in G has at most k forbidden flow values, each edge in G'' has at most $kq(G)$ forbidden flow values. \square

Theorem 17 for $k = 1$ combined with Theorem 2 implies Theorem 8 as well as the upper bound $g(k) \leq 8k^3 + 1$.

7 Group coloring

Jaeger et al. [5] define group colourability as follows.

Definition 18. Let Γ be an Abelian group. The graph G is said to be Γ -colorable if the following holds: Given some orientation D of G and any function $\varphi : E(G) \rightarrow \Gamma$ there exists a vertex coloring $c : V(G) \rightarrow \Gamma$ such that $c(w) - c(u) \neq \varphi(uw)$ for each $uw \in E(D)$.

We say that φ allows c .

A graph is d -degenerate if every subgraph contains a vertex of degree at most d . The coloring number $Col(G)$ is the largest number such that G is $(d - 1)$ -degenerate. Equivalently, $Col(G) - 1$ is the maximum minimum degree where the maximum is taken over all subgraphs of G . Clearly

$$\chi(G) \leq \chi_{wg}(G) \leq \chi_g(G) \leq Col(G). \quad (5)$$

Also

$$\chi_l(G) \leq Col(G), \quad (6)$$

where $\chi_l(G)$ is the list-chromatic number.

Now let G be a graph with $\chi_{wg}(G) = k$, and let Γ be an Abelian group such that G is Γ -colorable. Král' et al. [7] proved that $k > \delta/2 \ln(\delta)$ where δ is the minimum degree of G . (Král' et al. formulated their result as one about the group chromatic number but the proof works for the weak group chromatic number as well.) As this holds for every subgraph of G it also holds for a subgraph of minimum degree $\delta = Col(G) - 1$. Hence

$$\chi_{wg}(G) = k > \frac{\delta}{2 \ln \delta} = \frac{Col(G) - 1}{2 \ln(Col(G) - 1)} \geq \frac{\chi_g(G) - 1}{2 \ln(\chi_g(G) - 1)}. \quad (7)$$

This implies, for each natural number k , the upper bound in the following:

$$(2 - o(1))k < h(k) < (2 + o(1))k \ln(k). \quad (8)$$

Since the graphs $G_{q,k}$ are planar, their dual graphs establish the lower bound (for infinitely many primes k) by the proof of Theorem 4.

The group chromatic number and weak group chromatic number have some similarity to the list-chromatic number. Indeed, the proof of the 5-list-color theorem for planar graphs [16] translates, word for word, to the result that every planar graph has group chromatic number at most 5. And the proof of the 3-list-color theorem for planar graphs of girth at least 5 [17] translates to the result that every planar graph has group chromatic number at most 3. Only a minor detail in the proof needs additional explanation, see [14].

The following conjecture was made by Král' et al. in [7] and, according to [7], independently by Margit Voigt.

Conjecture 19. For every graph G , $\chi_l(G) \leq \chi_g(G)$.

We propose the analogous conjecture for the weak group chromatic number.

Conjecture 20. For every graph G , $\chi_l(G) \leq \chi_{wg}(G)$.

In [5] it is shown that every graph with two edge-disjoint spanning trees is Γ -connected for every Abelian group Γ of order at least 4. This implies that $g(3) = 3$ as pointed out in [11]. For, if G is \mathbb{Z}_3 -connected we let $\beta(v) = 1 - d(v)$ (where $d(v)$ denotes the degree of v) for every vertex v , except possibly one, in Definition 1. We may choose the resulting f such that $f(e) = 1$ for every edge e , by reversing the direction of those edges having flow value 2. This gives an orientation of the edges such that all vertices, except possibly one, have outdegree $2 \pmod{3}$ and hence outdegree at least 2. Thus G has at least $2|V(G)| - 2$ edges. As this also holds for every graph obtained from G by identifying vertices (and removing the loops that may arise), G has two edge-disjoint spanning trees, by a fundamental result of Edmonds [2], Nash-Williams [13] and Tutte [19].

Note that the dual statement does not hold: A \mathbb{Z}_3 -colorable graph is not necessarily the union of two spanning trees. For example is $K_{3,5}$ \mathbb{Z}_3 -colorable (see e.g. [10]) but contains 8 vertices and 15 edges, thus it has too many edges to be the union of two spanning trees.

Conjecture 21. $h(3) = 3$.

Conjecture 21 clearly implies Conjecture 22 below.

Conjecture 22. If G is \mathbb{Z}_3 -colorable, then G is \mathbb{Z}_4 -colorable and also $\mathbb{Z}_2 \times \mathbb{Z}_2$ -colorable.

Since $g(3) = 3$, the answer is affirmative for planar graphs. Jaeger et al. [5] asked if a graph is \mathbb{Z}_4 -connected if and only if it is $\mathbb{Z}_2 \times \mathbb{Z}_2$ -connected. Hušek et al. [3] answered this in the negative for both implications. As some of the counterexamples are planar, their

dual graphs show that \mathbb{Z}_4 -colorability does not imply $\mathbb{Z}_2 \times \mathbb{Z}_2$ -colorability. These graphs have multiple edges. Examples without multiple edges can be obtained using Hajos' construction. Such examples can even be made planar. Their dual graphs are therefore 3-edge-connected planar graphs that are \mathbb{Z}_4 -connected, but not $\mathbb{Z}_2 \times \mathbb{Z}_2$ -connected. For details, see [12]. We do not know if \mathbb{Z}_4 -colorability is implied by $\mathbb{Z}_2 \times \mathbb{Z}_2$ -colorability.

We conclude the paper by showing that Conjecture 22 in fact is equivalent to Conjecture 21. This follows from the two propositions below.

Proposition 23. *If G is \mathbb{Z}_3 -colorable, then G is 5-degenerate, i.e., $\chi_g(G) \leq \text{Col}(G) \leq 6$.*

Proof. Consider the bipartite graph with vertices $A \cup B$ in which the vertices in A corresponds to all possible \mathbb{Z}_3 -colorings of the vertices in G , and the vertices in B correspond to all possible \mathbb{Z}_3 -colorings of the edges in G . We join two vertices by an edge if the edge-coloring allows the vertex-coloring.

Let $n = |V(G)|$ and $m = |E(G)|$. Note that there 3^n vertices in A and 3^m vertices in B . Furthermore, each vertex in A has degree 2^m . As each edge-coloring will have at least one allowed vertex-coloring, all vertices in B has degree at least 1. Thus $3^n \cdot 2^m \geq 3^m$. We conclude

$$m \leq n \cdot \frac{\log(3)}{\log(\frac{3}{2})} < 2.8n < 3n, \tag{9}$$

so G is 5-degenerate. Thus, G is Γ -colorable for every Abelian group Γ of order $|\Gamma| \geq 6$. \square

We do not know if every \mathbb{Z}_3 -colorable graph is even 3-degenerate.

Proposition 24. *If G is \mathbb{Z}_3 -colorable, then G is \mathbb{Z}_5 -colorable.*

Proof. Let G be a \mathbb{Z}_3 -colorable graph, let D be an orientation of G , and let $\varphi : E(D) \rightarrow \mathbb{Z}_5$ be given. We think of \mathbb{Z}_5 as the numbers $0, 1, -1, 2, -2$. We define $\varphi' : E(D) \rightarrow \mathbb{Z}_3$ by reducing these numbers modulo 3, where we think of \mathbb{Z}_3 as the numbers $0, 1, -1$. More precisely, for each edge $e \in E(D)$,

$$\varphi'(e) = \begin{cases} 0 & \text{if } \varphi(e) = 0, \\ 1 & \text{if } \varphi(e) = 1 \text{ or } \varphi(e) = -2, \\ -1 & \text{if } \varphi(e) = 2 \text{ or } \varphi(e) = -1. \end{cases} \tag{10}$$

As G is \mathbb{Z}_3 -colorable, there exists a vertex-coloring $c' : V(G) \rightarrow \mathbb{Z}_3$ such that $c'(v) - c'(u) \neq \varphi'(uv)$ for each $uv \in E(D)$. Consider the vertex-coloring $c : V(G) \rightarrow \mathbb{Z}_5$ defined by $c(v) = c'(v) \pmod{5}$ for each vertex v . We argue that c is a proper coloring. Let $e = uv$ be a directed edge. If $c'(v) - c'(u) = 0$, then $c(v) - c(u) = 0$. As $\varphi'(e) \neq 0$ we get $\varphi(e) \neq 0$ so $c(v) - c(u) \neq \varphi(e)$. If $c'(v) - c'(u) = 1$, then $c(v) - c(u) = 1$ or $c(v) - c(u) = -2$. As $\varphi'(e) \neq 1$ we get $\varphi(e) \neq 1$ and $\varphi(e) \neq -2$ so $c(v) - c(u) \neq \varphi(e)$. If $c'(v) - c'(u) = -1$, then $c(v) - c(u) = 2$ or $c(v) - c(u) = -1$. As $\varphi'(e) \neq -1$ we get $\varphi(e) \neq 2$ and $\varphi(e) \neq -1$ so $c(v) - c(u) \neq \varphi(e)$. \square

References

- [1] J. A. Bondy and U. S. R. Murty, Graph Theory with Applications. *The MacMillan Press Ltd.* (1976).
- [2] J. Edmonds, Minimum partition of a matroid into independent subsets. *J. Res. Nat. Bur. Standards Sect. B* **69B** (1965) 67–72.
- [3] R. Hušek, L. Mohelníková, R. Šámal, Group Connectivity: \mathbb{Z}_4 v. \mathbb{Z}_2^2 . (2017) [arXiv:1711.03895](https://arxiv.org/abs/1711.03895).
- [4] F. Jaeger, Nowhere-zero flow problems. In: Selected Topics in Graph Theory 3 edited by L. W. Beineke and R. J. Wilson. *Academic Press* (1988) 71–95.
- [5] F. Jaeger, N. Linial, C. Payan, and M. Tarsi, Group connectivity of graphs – a nonhomogeneous analogue of nowhere-zero flow properties. *J. Combinatorial Theory, Ser B* **56** (1992) 165–182.
- [6] T. Jensen and B. Toft, Graph Coloring Problems. *John Wiley* (1995).
- [7] D. Král', O. Pangrác, and H.-J. Voss, A note on group colorings of planar graphs. *J. Graph Theory* **50** (2005) 123–129.
- [8] H.-J. Lai, Group connectivity of 3-edge-connected chordal graphs. *Graphs and Combinatorics* **16** (2000) 165–176.
- [9] H.-J. Lai, and X. Yao, Group connectivity of graphs with diameter at most 2. *European Journal of Combinatorics* **27** (2006) 436–443.
- [10] H.-J. Lai, and X. Zhang, Group colorability of graphs. *Ars Comb.* **62** (2002) 299–317.
- [11] J. Li, H.-J. Lai, and R. Luo, Group connectivity, strongly \mathbb{Z}_m -connectivity, and edge disjoint spanning trees. *SIAM Journal on Discrete Mathematics* **31** (2017) 1909–1922.
- [12] R. Langhede, PhD thesis, Technical University of Denmark, *to appear*.
- [13] C. St. J. A. Nash-Williams, Edge-disjoint spanning trees of finite graphs. *J. London Math. Soc.* **36** (1961) 445–450.
- [14] R. B. Richter, C. Thomassen, and D.H. Younger, Group-colouring, group-connectivity, claw-decompositions, and orientations in 5-edge-connected planar graphs. *Journal of Combinatorics* **7** (2016) 219–232.
- [15] P.D. Seymour, Nowhere-zero 6-flows. *J. Combinatorial Theory, Ser B* **30** (1981) 130–135.
- [16] C. Thomassen, Every planar graph is 5-choosable. *J. Combinatorial Theory, Ser B* **62** (1994) 180–181.
- [17] C. Thomassen, A short list color proof of Grötzsch's theorem. *J. Combinatorial Theory, Ser B* **88** (2003) 189–192.
- [18] W. T. Tutte, On the embedding of linear graphs in surfaces, *Proc. London Math. SOC. Ser. 2.* **51** (1949) 474–489.
- [19] W. T. Tutte, On the problem of decomposing a graph into n connected factors. *J. London Math. Soc.* **36** (1961) 221–230.
- [20] C. Q. Zhang, Integer flows and cycle covers of graphs. *Marcel Dekker, Inc* (1997).