



## Protect the vulnerable from extreme heat during the COVID-19 pandemic

Martinez, G. S.; Linares, C.; de'Donato, F.; Diaz, J.

*Published in:*  
Environmental Research

*Link to article, DOI:*  
[10.1016/j.envres.2020.109684](https://doi.org/10.1016/j.envres.2020.109684)

*Publication date:*  
2020

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*  
Martinez, G. S., Linares, C., de'Donato, F., & Diaz, J. (2020). Protect the vulnerable from extreme heat during the COVID-19 pandemic. *Environmental Research*, 187, Article 109684.  
<https://doi.org/10.1016/j.envres.2020.109684>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Protect the vulnerable from extreme heat during the COVID-19 pandemic

Martinez, GS.<sup>1</sup>, Linares, C.<sup>2</sup>, De'Donato, F.<sup>3</sup>, Diaz, J.<sup>2</sup>

<sup>1</sup> Technical University of Denmark, Copenhagen, Denmark

<sup>2</sup> Instituto de Salud Carlos III, Madrid, Spain

<sup>3</sup> Department of Epidemiology Lazio Regional Health Service - ASL ROMA 1, Rome, Italy

There is a considerable body of research studying how viruses, including some coronaviruses and influenza, are affected by weather, in which evidence suggests that their transmission may decline with higher humidity or temperatures. The spread of COVID-19 is changing rapidly and there are still many unknown factors that still need to be understood on how the virus is evolving, the dominant way of spreading and how these may differ by geographical area or context. The evidence that has surfaced thus far on SARS-COV-2 has not conclusively determined whether the weather conditions will be a key modulating factor influencing the transmission of the virus (Bukhari & Jameel, 2020; Chen et al., 2020; Gunthe, Swain, Patra, & Amte, 2020; Gupta, Raghuwanshi, & Chanda, 2020; Liu et al., 2020; Luo et al., 2020; Ma et al., 2020; Şahin, 2020; Shi et al., 2020; Tobías & Molina, 2020; Tosepu et al., 2020; J. Wang, Tang, Feng, & Lv, 2020; M. Wang et al., 2020; Yao et al., 2020; Jüni et al., 2020).

However, as we approach the warm season in the Northern hemisphere still under a pandemic situation, there is reason for strong concern about a related cascade of impacts, namely how the pandemic may aggravate the health impacts of heat waves by hindering prevention efforts. Several health authorities at federal, national, subnational and local levels run prevention plans to respond to and reduce health impacts of heat. These plans, generically known as Heat-health Action Plans (WHO, 2008), typically comprise a series of interventions, including heat warning systems, advice and information on keeping safe from heat, specific outreach and care for vulnerable population groups, surveillance of heat-related mortality and illnesses, and local interventions to reduce heat exposure through cooling centers and cool recreational areas.

The physical distancing measures and common space use restrictions set in place by most countries in response to the COVID-19 pandemic may hamper the implementation of those core heat-health prevention activities and aggravate the population's vulnerability to extreme temperatures this summer. We will explore some possible unintended effects of such restrictions, using as a framework typical national/federal level Heat-Health Action Plans in Europe.

For example, the effectiveness and outreach of heat warnings and health protective advice could be diminished in a context of widespread health warnings and information related to COVID-19. Far from an academic digression, for vulnerable groups this is a life-or-death issue: thermal extremes (both heat and cold) are by far the deadliest climate exposure in Europe, well above storms and flooding for example (CRED, 2020). Extreme heat causes significant mortality in the region every summer, with periodical peaks; the summer heatwaves of 2003 killed over 70,000 (Fouillet et al., 2008) and the combination of heat and wildfire smoke killed over 55,000 in Russia in 2010 (Barriopedro, Fischer, Luterbacher, Trigo, & Garcia-Herrera, 2011). Yet the health risks of heat are systematically underestimated by the general public and even by those most vulnerable to them (Abrahamson et al., 2009; Akompab et al., 2013; Cuesta, van Loenhout, Colaço, & Guha-Sapir, 2017; Howe, Marlon, Wang, & Leiserowitz, 2019; Van Loenhout & Guha-Sapir, 2016; Bittner & Stöbel, 2012). Adequate public health communication and media coverage should be ensured, and the language of heat-health warnings should thus reflect the seriousness of the risks posed by extreme heat, even in the current context of a pandemic. Warnings should integrate concomitant risk factors amidst the epidemic and heat, and ensure information on response measures and adaptation is clear even under lockdown or social distancing phases.

Similarly, the ability to reach out to and care for vulnerable people (e.g. living alone, chronically ill and the elderly) may be severely impaired in the current context of overwhelmed health and social care systems at every level. Adequate engagement of local governments and NGOs in heat-health action plans – still uncommon throughout the region- can help in protecting the most vulnerable against heat, by ensuring they are checked upon and can receive adequate care and support. This additional support may prove critical for residents and workers of institutions like nursing homes, which have been hit particularly hard by the pandemic. Access to necessary healthcare both at primary and specialized levels will continue to be restricted. In addition, fear of contracting COVID-19 may prevent some patients from seeking care even when experiencing heat-related symptoms, for example related to pre-existing conditions or interactions with medications.

Recent studies have shown that among COVID-19 patients the most prevalent comorbidities are hypertension, cardiovascular diseases, diabetes mellitus, chronic obstructive pulmonary disease (COPD), malignancy, and chronic kidney disease and cases with these comorbidities are more severe (Emami, Javanmardi, Pirbonyeh, & Akbari, 2020; Hu et al., 2020; Yang et al., 2020). These same chronic diseases are risk factors during heatwaves (Benmarhnia, Deguen, Kaufman, & Smargiassi, 2015). Further studies have shown higher COVID-19 mortality rates among the elderly and subjects with multi-chronic conditions (Shahid et al., 2020), thus making the European elderly population at an even greater risk this summer.

Rapid surveillance systems are a core component of heat plans and have been introduced throughout Europe in recent years not only to monitor the health impacts of heatwaves but are considered a strategic tool for an effective public health response (Martinez et al., 2019; WHO, 2008). Data from surveillance systems, especially mortality data, has also been called upon in the context of COVID-19 monitoring as they provide unbiased estimates not affected by case classification and can be useful to monitor containment and re-opening strategies (Leon et al., 2020). With the coming of high summer temperatures, it is vital that these systems are not entirely devoted to COVID-19 activity and still have the bandwidth to detect health impacts related to heat waves in order to ensure an adequate and timely response. Furthermore, when evaluating increases in mortality the potential role of both factors (heat and covid-19) on how one may affect the other or vice versa need to be studied in detail.

The use of publicly available air-conditioned spaces as cooling centers may not be compatible with the current directives mandating to maintain physical distance and avoid gathering indoors. In addition to the closure of public facilities like air-conditioned libraries, swimming pools and others, typical cool spaces like shopping malls and cafes may also be closed or restricted to various extents, even during the re-opening phases. Restrictions on publicly accessible cooled spaces will hit hardest those who can least afford air conditioning. Even for some households who may have been able to pay for the equipment and installation, running costs of AC are sometimes unaffordable. Lacking the options to access cooled spaces, residents without effective protection against heat may flock to cooler outdoor recreational areas including parks and water bodies. If not adequately managed, increased attendance in these spots could undermine the effectiveness of measures towards physical distancing and non-gathering.

If the decision of opening cooling centres is taken, users and staff will have to abide by a clear set of rules supported by an adequate physical setup (separation marks, printed instructions, banisters, etc.), supplies (hand disinfectant dispensers, face masks, etc.) and protocols (enhanced cleaning and disinfection, one-way walking, etc.). Since public transport may be restricted, ensuring the accessibility of cooled shelters will require complex logistics. Moreover, since some of the people who are at highest risk of severe outcomes from COVID-19 (for example, the elderly and the chronically ill) are also the most vulnerable to health risks from heat, gathering them in air conditioned spaces is potentially risky and should be done only if adequate space and facility setup can be guaranteed. One

1 clear example of potentially risky gathering of vulnerable individuals in common air conditioned rooms  
2 are nursing homes and elderly residences. If no cooling protection through air conditioning can be  
3 provided to highly vulnerable patients, authorities may consider dispensing personal cooling devices,  
4 although their effectiveness has been thus far tested mainly on occupational settings, laboratory or  
5 healthy subjects (Rawal et al., 2020).  
6

7 Additional common cooling solutions that should be reviewed for safety include mist sprayers (until  
8 more is known about the aerosol potential for transmission), public drinking water fountains that  
9 require manual operation (due to risk of surface contamination), and home visits by volunteer  
10 networks (in which asymptomatic infected volunteers may inadvertently put at risk vulnerable  
11 individuals). In some countries through various stages of lockdown, time slots for outdoor activity are  
12 being allocated by age groups and may not be correlated with vulnerability to heat. In general, both  
13 general procedures within Heat Health Action Plans and specific solutions for cooling should be  
14 reviewed and modified if necessary under the current situation.  
15  
16

17 Any decisions will inevitably will have to be made with limited information and plagued by uncertainty.  
18 In some instances, air conditioning may aid droplet transmission of the virus (Lu et al., 2020; Correia,  
19 Rodrigues, Gameiro da Silva, & Gonçalves, 2020) thus casting doubt on its safe use for groups. Seasonal  
20 patterns of mortality may change significantly, since there is reason to believe that COVID-19 related  
21 mortality will have decimated a proportion of the most vulnerable to heat before the summer season  
22 starts. Research has observed that high respiratory, cardiovascular and influenza mortality in winter  
23 leads to lower temperature effects in the following summer (Rocklöv, Forsberg, & Meister, 2009).  
24 Though several climate-influenced exposures (e.g. air pollution, allergenic pollen, heat) tend to occur  
25 concurrently (Linares, Martinez, Kendrovski, & Diaz, 2020), the current situation will further hinder  
26 the integration of prevention efforts, thus affecting their effectiveness and reach.  
27  
28  
29  
30

31 Finding and implementing solutions to these conundrums is urgent. In 2019, the annual temperature  
32 for Europe was the highest on record, and 11 of the 12 hottest years on record have all occurred since  
33 the year 2000 (ECMWF, 2020a). Some forecasts predict that 2020 may also be one of the hottest, if  
34 not the hottest, year on record (ECMWF, 2020b; NOAA National Centers for Environmental  
35 Information, 2020). This discussion will inevitably lead to hard choices with far-reaching consequences.  
36 The WHO has warned that failing to adapt the COVID-19 response to the prevention and management  
37 of Non-Communicable Disease risks (among which is heat) will mean failing many people precisely at  
38 a time when their vulnerability is heightened (Kluge et al., 2020). Thus, it is paramount that we plan  
39 ahead and make such choices with enough lead time, based on the available evidence, with equity  
40 and respect to fundamental rights, and according to agreed upon ethical principles. Importantly, there  
41 is still enough time to issue national-level guidance to be trickled down to local authorities, so that  
42 implementation is homogeneous and controlled. Leaving decisions on heat-health protection to  
43 municipalities and responders in the field without guidance can create heterogeneity in  
44 implementation, confusion in the public and ultimately additional harm.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

- Abrahamson, V., Wolf, J., Lorenzoni, I., Fenn, B., Kovats, S., Wilkinson, P., ... Raine, R. (2009). Perceptions of heatwave risks to health: interview-based study of older people in London and Norwich, UK. *J.Public Health (Oxf)*, 31(1741-3850 (Electronic)), 119–126.
- Akompab, D., Bi, P., Williams, S., Grant, J., Walker, I., Augoustinos, M., ... Augoustinos, M. (2013). Heat Waves and Climate Change: Applying the Health Belief Model to Identify Predictors of Risk Perception and Adaptive Behaviours in Adelaide, Australia. *International Journal of Environmental Research and Public Health*, 10(6), 2164–2184. <https://doi.org/10.3390/ijerph10062164>
- Barriopedro, D., Fischer, E. M., Luterbacher, J., Trigo, R. M., & Garcia-Herrera, R. (2011). The Hot Summer of 2010: Redrawing the Temperature Record Map of Europe. *Science*, 332(6026), 220–224. <https://doi.org/10.1126/science.1201224>
- Benmarhnia, T., Deguen, S., Kaufman, J. S., & Smargiassi, A. (2015, October 1). Vulnerability to heat-related mortality: A systematic review, meta-analysis, and meta-regression analysis. *Epidemiology*. Lippincott Williams and Wilkins. <https://doi.org/10.1097/EDE.0000000000000375>
- Bittner, M. I., & Stöbel, U. (2012). Zur gesundheitlichen risikowahrnehmung von hitzewellen: Ergebnisse einer qualitativen interviewstudie mit älteren menschen und ihren pflegepersonen in Freiburg. *GMS Psycho-Social-Medicine*, 9, Doc05. <https://doi.org/10.3205/psm000083>
- Bukhari, Q., & Jameel, Y. (2020). Will Coronavirus Pandemic Diminish by Summer? *SSRN Electronic Journal*. <https://doi.org/10.2139/ssrn.3556998>
- Chen, B., Liang, H., Yuan, X., Hu, Y., Xu, M., Zhao, Y., ... Zhu, X. (2020). Roles of meteorological conditions in COVID-19 transmission on a worldwide scale. *MedRxiv*, 11, 2020.03.16.20037168. <https://doi.org/10.1101/2020.03.16.20037168>
- Correia, G., Rodrigues, L., Gameiro da Silva, M., & Gonçalves, T. (2020). Airborne route and bad use of ventilation systems as non-negligible factors in SARS-CoV-2 transmission. *Medical Hypotheses*, 141, 109781. <https://doi.org/10.1016/j.mehy.2020.109781>
- CRED. (2020). EM-DAT. The international disaster database. Retrieved from <http://www.emdat.be/>
- Cuesta, J. G., van Loenhout, J. A. F., Colaço, M. da C., & Guha-Sapir, D. (2017). General population knowledge about extreme heat: A cross-sectional survey in Lisbon and Madrid. *International Journal of Environmental Research and Public Health*, 14(2), 10.3390/ijerph14020122. <https://doi.org/10.3390/ijerph14020122>
- ECMWF. (2020a). *European State of the Climate 2019*. Reading, United Kingdom. Retrieved from [https://climate.copernicus.eu/sites/default/files/2020-04/ESOTC2019\\_summary.pdf](https://climate.copernicus.eu/sites/default/files/2020-04/ESOTC2019_summary.pdf)
- ECMWF. (2020b). *Surface air temperature for March 2020*. Reading, United Kingdom. Retrieved from <https://climate.copernicus.eu/surface-air-temperature-march-2020>
- Emami, A., Javanmardi, F., Pirbonyeh, N., & Akbari, A. (2020). Prevalence of Underlying Diseases in Hospitalized Patients with COVID-19: a Systematic Review and Meta-Analysis. *Archives of Academic Emergency Medicine*, 8(1), e35. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/32232218>
- Fouillet, A., Rey, G., Wagner, V., Laaidi, K., Empereur-Bissonnet, P., Le Tertre, A., ... Hémon, D. (2008). Has the impact of heat waves on mortality changed in France since the European heat wave of summer 2003? A study of the 2006 heat wave. *International Journal of Epidemiology*, 37(2), 309–317. <https://doi.org/10.1093/ije/dym253>
- Gunthe, S. S., Swain, B., Patra, S. S., & Amte, A. (2020). On the global trends and spread of the COVID-19 outbreak: preliminary assessment of the potential relation between location-specific temperature and UV index. *Journal of Public Health*. <https://doi.org/10.1007/s10389-020-01279-y>
- Gupta, S., Raghuwanshi, G. S., & Chanda, A. (2020). Effect of weather on COVID-19 spread in the US:

- A prediction model for India in 2020. *Science of The Total Environment*, 728, 138860.  
<https://doi.org/10.1016/j.scitotenv.2020.138860>
- Howe, P. D., Marlon, J. R., Wang, X., & Leiserowitz, A. (2019). Public perceptions of the health risks of extreme heat across US states, counties, and neighborhoods. *Proceedings of the National Academy of Sciences of the United States of America*, 116(14), 6743–6748.  
<https://doi.org/10.1073/pnas.1813145116>
- Hu, Y., Sun, J., Dai, Z., Deng, H., Li, X., Huang, Q., ... Xu, Y. (2020, June 1). Prevalence and severity of corona virus disease 2019 (COVID-19): A systematic review and meta-analysis. *Journal of Clinical Virology*. Elsevier B.V. <https://doi.org/10.1016/j.jcv.2020.104371>
- Jüni, P., Rothenbühler, M., Bobos, P., Thorpe, K. E., Costa, B. R. da, Fisman, D. N., ... Gesink, D. (2020). Impact of climate and public health interventions on the COVID-19 pandemic: A prospective cohort study. *CMAJ*. <https://doi.org/10.1503/CMAJ.200920>
- Kluge, H. H. P., Wickramasinghe, K., Rippin, H. L., Mendes, R., Peters, D. H., Kontsevaya, A., & Breda, J. (2020). Prevention and control of non-communicable diseases in the COVID-19 response. *The Lancet*, 0(0). [https://doi.org/10.1016/S0140-6736\(20\)31067-9](https://doi.org/10.1016/S0140-6736(20)31067-9)
- Leon, D. A., Shkolnikov, V. M., Smeeth, L., Magnus, P., Pechholdová, M., & Jarvis, C. I. (2020). COVID-19: a need for real-time monitoring of weekly excess deaths. *The Lancet*, 0(0).  
[https://doi.org/10.1016/S0140-6736\(20\)30933-8](https://doi.org/10.1016/S0140-6736(20)30933-8)
- Linares, C., Martinez, G., Kendrovski, V., & Diaz, J. (2020). A New Integrative Perspective on Early Warning Systems for Health in the Context of Climate Change. *Environ Res (In Press)*.
- Liu, J., Zhou, J., Yao, J., Zhang, X., Li, L., Xu, X., ... Zhang, K. (2020). Impact of meteorological factors on the COVID-19 transmission: A multi-city study in China. *Science of the Total Environment*, 726, 138513. <https://doi.org/10.1016/j.scitotenv.2020.138513>
- Lu, J., Gu, J., Li, K., Xu, C., Su, W., Lai, Z., ... Yang, Z. (2020). COVID-19 Outbreak Associated with Air Conditioning in Restaurant, Guangzhou, China, 2020. *Emerging Infectious Diseases*, 26(7).  
<https://doi.org/10.3201/eid2607.200764>
- Luo, W., Majumder, M. S., Liu, D., Poirier, C., Mandl, K. D., Lipsitch, M., & Santillana, M. (2020). The role of absolute humidity on transmission rates of the COVID-19 outbreak. *MedRxiv*, 2020.02.12.20022467. <https://doi.org/10.1101/2020.02.12.20022467>
- Ma, Y., Zhao, Y., Liu, J., He, X., Wang, B., Fu, S., ... Luo, B. (2020). Effects of temperature variation and humidity on the mortality of COVID-19 in Wuhan. *MedRxiv*, 2020.03.15.20036426.  
<https://doi.org/10.1101/2020.03.15.20036426>
- Martinez, G. S., Linares, C., Ayuso, A., Kendrovski, V., Boeckmann, M., & Diaz, J. (2019). Heat-health action plans in Europe: Challenges ahead and how to tackle them. *Environmental Research*, 176, 108548. <https://doi.org/10.1016/J.ENVRES.2019.108548>
- NOAA National Centers for Environmental Information. (2020). *State of the Climate: Global Climate Report for March 2020*. Silver Spring, MD. Retrieved from  
<https://www.ncdc.noaa.gov/sotc/global/202003/supplemental/page-2>
- Rawal, R., Schweiker, M., Kazanci, O. B., Vardhan, V., Jin, Q., & Duanmu, L. (2020, May 1). Personal comfort systems: A review on comfort, energy, and economics. *Energy and Buildings*. Elsevier Ltd. <https://doi.org/10.1016/j.enbuild.2020.109858>
- Rocklöv, J., Forsberg, B., & Meister, K. (2009). Winter mortality modifies the heat-mortality association the following summer. *European Respiratory Journal*, 33(2), 245–251.  
<https://doi.org/10.1183/09031936.00037808>
- Şahin, M. (2020). Impact of weather on COVID-19 pandemic in Turkey. *Science of the Total Environment*, 728, 138810. <https://doi.org/10.1016/j.scitotenv.2020.138810>
- Shahid, Z., Kalayanamitra, R., McClafferty, B., Kepko, D., Ramgobin, D., Patel, R., ... Jain, R. (2020). COVID-19 and Older Adults: What We Know. *Journal of the American Geriatrics Society*, jgs.16472. <https://doi.org/10.1111/jgs.16472>
- Shi, P., Dong, Y., Yan, H., Li, X., Zhao, C., Liu, W., ... Xi, S. (2020). The impact of temperature and absolute humidity on the coronavirus disease 2019 (COVID-19) outbreak - evidence from China.

MedRxiv, 2020.03.22.20038919. <https://doi.org/10.1101/2020.03.22.20038919>  
 Tobías, A., & Molina, T. (2020, July 1). Is temperature reducing the transmission of COVID-19 ?  
*Environmental Research*. Academic Press. <https://doi.org/10.1016/j.envres.2020.109553>  
 Tosepu, R., Gunawan, J., Effendy, D. S., Ahmad, L. O. A. I., Lestari, H., Bahar, H., & Asfian, P. (2020).  
 Correlation between weather and Covid-19 pandemic in Jakarta, Indonesia. *Science of the Total  
 Environment*, 725, 138436. <https://doi.org/10.1016/j.scitotenv.2020.138436>  
 Van Loenhout, J. A. F., & Guha-Sapir, D. (2016). How resilient is the general population to  
 heatwaves? A knowledge survey from the ENHANCE project in Brussels and Amsterdam. *BMC  
 Research Notes*, 9(1), 1–5. <https://doi.org/10.1186/s13104-016-2305-y>  
 Wang, J., Tang, K., Feng, K., & Lv, W. (2020). High Temperature and High Humidity Reduce the  
 Transmission of COVID-19. *SSRN Electronic Journal*. <https://doi.org/10.2139/ssrn.3551767>  
 Wang, M., Jiang, A., Gong, L., Luo, L., Guo, W., Li, C., ... Li, H. (2020). Temperature significant change  
 COVID-19 Transmission in 429 cities. *MedRxiv*, 2020.02.22.20025791.  
<https://doi.org/10.1101/2020.02.22.20025791>  
 WHO. (2008). *Heat-health action plans: a guidance*. (F. Matthies, G. Bickler, N. Cardeñosa, & S.  
 Hales, Eds.). Copenhagen, Denmark: World Health Organization Regional Office for Europe.  
 Yang, J., Zheng, Y., Gou, X., Pu, K., Chen, Z., Guo, Q., ... Zhou, Y. (2020). Prevalence of comorbidities  
 and its effects in coronavirus disease 2019 patients: A systematic review and meta-analysis.  
*International Journal of Infectious Diseases*, 94, 91–95.  
<https://doi.org/10.1016/j.ijid.2020.03.017>  
 Yao, Y., Pan, J., Liu, Z., Meng, X., Wang, W., Kan, H., & Wang, W. (2020). No Association of COVID-19  
 transmission with temperature or UV radiation in Chinese cities. *European Respiratory Journal*,  
 2000517. <https://doi.org/10.1183/13993003.00517-2020>