Intersecting xenobiology and neo-metabolism to bring novel chemistries to life

Nieto Domínguez, Manuel José; Nickel, Pablo Ivan

Published in:
ChemBioChem

Link to article, DOI:
[10.1002/cbic.202000091](https://doi.org/10.1002/cbic.202000091)

Publication date:
2020

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Nieto Domínguez, M. J., & Nickel, P. I. (2020). Intersecting xenobiology and neo-metabolism to bring novel chemistries to life. *ChemBioChem*, 21(18), 2551-2571. <https://doi.org/10.1002/cbic.202000091>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Article

Title: Intersecting xenobiology and neo-metabolism to bring novel chemistries to life

Authors: Manuel Nieto-Dominguez and Pablo Ivan Nikel

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *ChemBioChem* 10.1002/cbic.202000091

Link to VoR: <https://doi.org/10.1002/cbic.202000091>

Intersecting xenobiology and *neo*-metabolism to bring novel chemistries to life

by

Manuel Nieto-Domínguez and Pablo I. Nikel*

The Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, 2800 Kongens Lyngby, Denmark

* Correspondence to: *Pablo I. Nikel* (pabnik@biosustain.dtu.dk)
The Novo Nordisk Foundation Center for Biosustainability
Technical University of Denmark
Kgs Lyngby, Denmark
Tel: (+45 93) 51 19 18

Accepted Manuscript

Summary

The life diversity relies on a handful chemical elements (carbon, oxygen, hydrogen, nitrogen, sulphur and phosphorus) as part of essential building blocks, whereas other atoms are needed to a lesser extent and most of the remaining elements are excluded from biology. This circumstance limits the scope of biochemical reactions in extant metabolism — yet it offers a phenomenal playground for synthetic biology. Xenobiology aims at bringing novel bricks to life that could be exploited towards (xeno)metabolite synthesis. In particular, the assembly of novel pathways engineered to handle non-biological elements (*neo*-metabolism) will broaden the chemical space beyond the reach of natural evolution. In this review, xeno-elements that could be blended into Nature's biosynthetic portfolio are discussed together with their physicochemical properties and tools and strategies to incorporate them into biochemistry. We argue that current bioproduction methods can be revolutionized by intersecting xenobiology with *neo*-metabolism for the synthesis of new-to-Nature molecules, e.g. organohalides.

1. Introduction

There is little doubt that the XXI century has come to be the *age of biology*, in the same way as the XX century has been a prime time for physics and chemistry. Biotechnology and synthetic biology aim at developing sustainable alternatives for a number of crucial areas, e.g. energy supply^[1], synthesis of chemicals,^[2] food production,^[3] biomaterials,^[4] bioremediation,^[5] therapies^[6] or diagnostics.^[7] This overarching goal faces the challenge of alleviating our dependence on fossil fuels, non-renewable resources and conventional synthetic chemistry. As such, the biological realm is being increasingly mined by adopting core engineering principles to unleash the untapped potential of life diversity. Life is the main resource of biology, and its rich diversity on the Earth's biosphere has surpassed all expectations. From the first classifications by Linnaeus in the XVIII century to the latest estimations nowadays, the figures have grown from originally spanning 11,000 different animals and plants to currently cover more than 8 million eukaryotic organisms^[8] and as many as 10^{12} microbial species.^[9] The ability of life to expand and prosper even in extreme environments (e.g. hot springs, deep sea, Antarctic ice or industrial chemical wastes) is only possible due to the existence of versatile biochemical adaptations to harsh and often adverse conditions. The relatively new era of microbial «omics» approaches^[10] paved the way towards identifying genetic parts and pieces encoding the rich biochemical machinery underlying adaptation to such conditions, enabling their use for engineering purposes. However, at the very fundamental level, this extreme biochemical versatility is based on just a small number of chemical elements.^[11] Specifically, carbon (C), oxygen (O), hydrogen (H), nitrogen (N), sulphur (S) and phosphorous (P) are the main components of virtually every known lifeform. A myriad of other elements are, to a greater or lesser extent, necessary for life — but none of them is essential for the synthesis of the basic biomolecules of the cell (i.e. proteins, lipids, sugars and nucleic acids). This state of affairs means that, beyond the outstanding adaptability of life, the development of bio-based alternatives to production is confronted with the limited panoply of building blocks naturally used in biological processes.

Xenobiology is a specific sub-field of *synthetic biology* that deals with non-conventional forms of life differing from the canonical dogma of biology^[12] (which is based in DNA, RNA and the 20 natural amino acids — although the very concept has been since challenged by bringing forward the prominent role of metabolism in orchestrating the functional relationship between nucleic acids and proteins^[13]). From a wide and fundamental perspective, xenobiology seeks to answer one simple but key question: is life as

we know it the *only possible* one? During the past decades, scientists have largely speculated about this matter, and the question also reached the general public (mostly through science fiction — the extremely corrosive blood from the xenomorph in the *Alien* movie or the mineral body of the Crystalline Entity from *Star Trek* series are just a few noted examples on how alternative life chemistries have been imagined in popular culture). Most of the work done in the field of xenobiology has focused on incorporating new «letters» in both the genetic and protein alphabets.^[14] Apart from these fundamental aspects, the core tenets of xenobiology can be also applied to overcome limitations of natural metabolism by developing artificial routes (i.e. synthetic or *neo-metabolism*)^[15] to introduce *new-to-Nature molecules* and intermediates in the chemical diversity of cells. As such, *neo-metabolism* can be defined as a set of (novel) biochemical reactions and routes purposely assembled to integrate substrates, intermediates, products and chemical reactions unprecedented (or very rare) in Nature. The intersection of *neo-metabolism* with xenobiology embodies the efforts of blending abiotic atoms into the native biochemistry of the cells — not only by incorporating completely alien- (or seldom associated-) to-life chemical elements but also by rewriting their role to display central functions instead of being restricted to few specific routes in the secondary metabolism. On this background, this review focuses on biological mechanisms, both natural and engineered, for bringing novel elements from the inorganic realm into biology. Specifically, we present and discuss available information on enzymes and pathways towards re-defining the role of fluorine (F), chlorine (Cl), bromine (Br), iodine (I), silicon (Si) and arsenic (As) in the chemistry of life, with a focus on microbial hosts. A more comprehensive description is reserved to F owing to its differences with other halogens. On this background, we will separately address how these features are relevant for F incorporation into microbial biochemistry, and the present and future of biocatalysts and pathways towards organofluorine biosynthesis will be reviewed.

From a broader perspective, the choice of the chemical elements covered herein is not accidental: each one of them can be considered as the *dark twin* of atoms usually found in life, closely related in terms of physicochemical properties yet apparently rejected by the forces of natural evolution. In this regard, F and, to a lesser extent, Cl, Br and I, are chemical equivalents of H in C–H bonds, whereas Si mirrors C and As is a potential analogue of P. When present in biological molecules, the role of these atoms remains relegated to secondary metabolic pathways. **Fig. 1** displays a summary of the most relevant features of selected biological chemical elements and their xeno-analogues, including their abundance in the Earth crust and physicochemical properties. We argue that *neo-metabolism* can be deployed to handle abiotic

elements for the sake of whole-cell and cell-free biocatalysis by exploiting chemical similarities between these atoms. Finally, and based on these scenarios, we discuss the potential outcomes and benefits of incorporating novel chemical elements into biomolecules and synthetic pathways.

2. Chlorine, bromine and iodine

Cl, Br and I, together with F, are the main members of the group of elements called *halogens*. Halogens were firstly recognized for their ability to readily form salts with metals (*halo-*, derived from the Greek ἅλς, *salt*), although organohalides, compounds containing at least one C–X bond (where X represents a halogen atom), are referred to in scientific literature dating back to at least the XIX century.^[16] In fact, organohalides occupy a central position in modern organic chemistry, and innovative synthetic approaches for their preparation are continuously reported.^[17] However, not every halogenated compound possesses an artificial origin: in a recent review, the number of natural organohalides was established to be represented by more than 5,000 different compounds,^[18] most of them from marine organisms.^[19] In 2017, a fascinating report by Fayolle and co-workers^[20] revealed the existence of extraterrestrial organohalides, likely formed during a pre-planetary process.

Physicochemical features of each halogen determine the properties of the C–X bond. In particular, bond polarity depends on the halogen electronegativity, which follows the order $F > Cl > Br > I$ (**Fig. 1**). Therefore, the C–F bond shows the highest polarity and the C–I bond the lowest one.^[21] Reactivity is a direct result of bond strength, which displays an order equivalent to the polarity and, as such, C–F has the greatest strength and the least reactivity.^[21] Organohalides are valuable starting points for the synthesis of important molecules in well-known chemical reactions, e.g. the widely-used Grignard reagent,^[22] the Gilman reagent^[23] or the Simmons-Smith reagent.^[24] In Nature, halogenation seems to operate as a general mechanism to modify the physicochemical properties of certain (halo)metabolites.^[25] Neumann and co-workers^[26] went a step further and stated that the halogenation of natural product scaffolds allows the organisms that produce them to optimize the bioactivity of small molecules thus conferring an evolutionary advantage. Indeed, there are several reports on organohalides that lose or decrease their bioactivity in their dehalogenated form.^[27] Considering these aspects, the present section of the article will be focused on the available biological mechanisms and tools for the incorporation of Cl, Br and I into organic molecules and the relevance of halometabolites for xenobiology. Due to its

importance in countless industrial applications and its unique particularities, F will be considered in a separate section.

2.1. Mechanisms of enzymatic halogenation involving Cl, Br and I

Four classes of halogenating enzymes are known to catalyze the formation of C–Cl, C–Br and C–I bonds: haem haloperoxidases, vanadium (V) haloperoxidases, non-haem iron halogenases and non-metallo-haloperoxidases,^[28] with the latest group forming a category that includes flavin-dependent halogenases and S-adenosyl-L-methionine (SAM)-dependent halogenases.^[29] Both *haem haloperoxidases* and *vanadium haloperoxidases* contain a metal catalyst coordinated to their active site (Fe or V, respectively). The catalytic mechanisms rely on the oxidation of the metal by hydrogen peroxide (H₂O₂), which subsequently oxidizes the halide atom to the highly reactive OX[–] form. This species rapidly reacts with an appropriate organic substrate forming the halogenated organic product. *Non-haem iron halogenases* or *α-ketoglutarate-dependent, non-haem iron (Fe_{NH}) halogenases* receive their name because Fe is coordinated to α-ketoglutarate (which also acts as co-substrate of the reaction) instead of being attached to a porphyrin group. In this case, the reaction mechanism is more complex than that of haloperoxidases, and requires dioxygen (O₂) instead of H₂O₂ as the oxidizing agent. The organic substrate is converted into a radical because of the abstraction of a hydrogen atom in the active site, and reacts with a halogen atom bound to the Fe center to form the halogenated product. *Flavin-dependent halogenases*, in turn, catalyze the reaction of FADH₂ with O₂ to yield a peroxide-linked isoalloxazine ring, which then oxidizes the halide and generates OX[–]. This compound reacts with the organic substrate, leading to the synthesis of the organohalogen molecule. Yet another mechanism of halogenation relies on SAM as the organic substrate. Methyl halide transferases are the most studied class among the group of enzymes collectively known as *SAM-dependent halogenases*. These biocatalysts use the universal methyl donor SAM to methylate halides (thus yielding Me–X), and the catalytic mechanism of this type of enzyme involves the nucleophilic attack of SAM by the halide anion (X[–]) to form Me–X and releasing S-adenosyl-L-homocysteine (SAH) in the process. Another class of SAM-dependent halogenases executes a different (and distinct) reaction mechanism, and the so-called *fluorinases* have aroused the greatest attention among them (despite the fact that this group of enzymes includes some examples that are active on Cl[–], Br[–] and I[–]).^[29b,30] As such, fluorinase enzymes will be described in more detail in the section dealing with the incorporation of F into organic molecules. The enumeration of the diverse mechanisms mediating the

addition of halogens into organic molecules would point to the existence of a considerable portfolio of organohalides formed thereby, as elaborated below.

2.2. Natural and synthetic occurrence of organohalides

The biocatalysts described in the previous section mediate the incorporation of halogens in Nature, and Cl and Br are the atoms more frequently incorporated into organic structures, while iodination seems to be rarer occurrence.^[26] Halogenated natural compounds with known biological effects include molecules displaying antibacterial, antifungal, antiparasitic, antiviral, antitumor, antioxidant and antiinflammatory activities.^[31] Many of these properties rely on the capacity of halogenated compounds to inhibit key enzymes involved in the cognate biological processes, e.g. as kinases, phosphatases or proteases.^[19] A detailed description of the nature and characteristics of organohalides is beyond the purpose of this review, but several articles enumerate naturally-occurrences of these molecules, and the list is continuously expanding.^[32] Intriguingly enough, despite of the variety of effects and molecular targets they exhibit, naturally-occurring halogenated metabolites are largely restricted to secondary microbial metabolism and they are essentially absent in macromolecules. Natural halogenated polysaccharides have not been reported and, in general, halogenated carbohydrates are rare,^[33] although a few have been identified, e.g. two 4'-fluoro-3'-O- β -glucosylated metabolites isolated from *Streptomyces calvus*^[34] and a chloro-glycosylated antitumoral molecule synthesized by *Pseudomonas* sp. strain 2663.^[35] Natural halogenated polypeptides are scarce, although peroxidase-mediated chlorination and bromination of proteins have been reported in eosinophils^[36] and the thyroid gland of vertebrates contains natural iodinated proteins, of which thyroglobulin is the most extensively studied.^[37] Interestingly, a recent report suggests that the de-iodination of the thyroid hormone, a process catalyzed by a selenium (Se)-containing enzyme, might occur through a cooperative Se...I bond, which would be a noteworthy example of biochemical interaction between two xeno-elements.^[38] Additionally, small amounts of I and, to a lesser extent, Br, have been found in scleroproteins from different invertebrates. There are more examples of naturally-occurring halogenated fatty acids,^[19,39] the halogen atoms of which are incorporated mainly through the action of haloperoxidases after the core polymeric substrate of the lipid has been already synthesized.^[40]

1 In contrast to the scarcity of halogens in large proteins, there are several occurrences of natural
2 halogenated amino acids (HAAs) (**Fig. 2**) and a few naturally-occurring peptides incorporating different
3 halogen atoms.^[41] Interestingly, and in spite of the (apparent) rarity of natural pathways for iodination,
4 one of the first identified HAA has been 3,5-diiodotyrosine (used as a precursor in the production of the
5 thyroid hormone), isolated from a coral in the XIX century. Over time, a number of reports indicated that
6 aromatic amino acids seem to be the preferred structures subjected to natural halogenation. Thus, the
7 classic work of Hager and co-workers^[42] reported on the synthesis of Cl, Br and I derivatives of tyrosine
8 using a chloroperoxidase as the biocatalyst for halogen incorporation. Tryptophan is a preferred target
9 molecule of biohalogenation pathways, and halogen substituents in different positions in the aromatic
10 ring of this amino acid have been described in the primary literature.^[43]

11
12 From a different perspective, and complementary to the natural spectrum of HAAs, chemical synthesis
13 offers access to a wide repertoire of these molecules, both aromatic and aliphatic (non-polar, polar and
14 charged).^[44] Furthermore, in an attempt to overcome natural synthetic limitations, sophisticated
15 approaches for engineering the translational machinery of certain yeast and bacterial hosts have been
16 developed during the two last decades. These strategies allowed the site-specific incorporation of
17 different unnatural amino acids (including HAAs^[45]) into peptides and proteins. The most remarkable
18 methods of this sort will be discussed in the section dealing with fluorination, which, as indicated before,
19 displays potential to be introduced in the cellular proteome without compromising viability because of its
20 distinct physicochemical properties. We note that an outstanding attempt to engineer a microbial host to
21 incorporate a halogenated compound as a primary building block was not related to the proteome but
22 with the genome. Philippe Marlière and co-workers^[46] evolved an *Escherichia coli* strain unable to
23 synthesize thymine to replace these residues across the genome by the synthetic variant 5-chlorouracil
24 (**Fig. 2**), thereby reducing the genomic content of thymine below 2% — a strain dubbed *Escherichia*
25 *chlori*. Interestingly, the evolved *E. chlori* strain was shown to display a phenotype similar to that observed
26 in its wild-type counterpart, highlighting the flexibility of the biochemical machinery to recognize and
27 incorporate a halogenated nucleotide analogue. Yet, the bulkiness of Cl, Br and I may hinder broad
28 applications in engineering synthetic metabolism (or, for that matter, incorporation into the core
29 components of living cells) — a feature that sets F aside and highlights its potential applications as
30 disclosed in the next section.

3. Fluorine

The combination of a small size (close to that of H) and the high electronegativity of the F atom not only sets this element apart from the other halogens,^[47] but they also make it an essential element in the pharmaceutical, material and agrochemical industry.^[48] F can easily replace H in organic compounds causing minimal distortion of the molecular geometry,^[49] and incorporation of F into biomolecules considerably increases their hydrophobicity, exerting a positive impact on bioavailability.^[50] Such properties have been increasingly exploited in the pharma industry over the years, and approximately 25% of the licensed pharmaceuticals nowadays contain at least one F atom^[51] — ranking second only after N as the favorite heteroatom in drug discovery and development.^[52] F is also present in more complex molecules of industrial interest, widely distributed in daily applications. Fluoropolymers, for instance, have been successfully commercialized for more than 70 years because of their outstanding material properties (including TEFLON™, among many other examples of this sort).^[53] Low-molecular-weight fluorinated compounds, e.g. 3-fluoropyruvate (Pyr-3-F), serve as F donors for the synthesis of a wide variety of added-value products (as indicated in **Fig. 2**, showing the main F-containing organic compounds discussed in this review). In particular, Pyr-3-F is the key precursor for the enzymatic synthesis of fluorosugars,^[54] fluorinated amino acids^[55] and α -fluoro- β -hydroxy carboxylic esters.^[56] Although all these organofluorine molecules are essential building blocks in a number of practical applications, their chemical synthesis is still quite challenging and often requires the use of hazardous reagents (i.e. HF or F₂). The production of Pyr-3-F itself relies entirely on traditional chemistry,^[57] with no bio-based alternatives for fluorochemical production known to date. Yet, all applications of F-containing molecules continue pushing market requirements.^[58] Unfortunately, nature has barely evolved biochemical reactions involving F, and fluorinated compounds are still synthesized exclusively by chemical approaches nowadays. The isolation of the *fluorinase* from the actinomycete soil bacterium *Streptomyces cattleya*, the first enzyme identified to catalyze the formation of a C–F bond,^[59] constituted a breakthrough and a major step towards the development of a bio-based production of fluorochemicals as elaborated in the next section. The other known biological mechanism of (transient) bonding between C and F is catalyzed by mutant variants of a β -glucosidase from *Agrobacterium* sp. and a β -mannosidase from *Cellulomonas fimi* through nucleophilic halogenation of 2,4-dinitrophenyl- β -glucoside.^[60]

3.1. The fluorinase enzyme: challenges and opportunities for establishing *neo*-metabolism for fluorochemicals

At a first glance, it may seem surprising that only a few naturally-occurring fluorometabolites have been identified so far. F is the 13th element of the continental crust in terms of mass abundance, a value close to Cl and much higher than Br or I (**Fig. 1**). However, inorganic halogens are bioavailable mainly as anions in aqueous solutions, and the abundance of F⁻ in the oceans is 1.3 ppm — far below from Cl⁻ (20,058 ppm) and Br⁻ (70 ppm).^[61] Yet, this difference in abundance and bioavailability is not large enough to explain why F is almost excluded from the native biochemistry of many organisms, since I⁻ (0.03–0.001 ppm in sea water)^[62] is linked to a relatively high number of natural iodinated metabolites.^[63] Therefore, the extreme rarity of F in life seems to be associated to its physicochemical properties rather than its abundance.

There are two major reasons explaining why the oldest demonstration of enzymatic chlorination was reported in 1959^[64] whereas the first fluorinase was not isolated until 2002 by David O'Hagan and co-workers.^[65] On the one hand, F possesses an elevated redox potential (F⁻/F₂ = -2.87 eV), considerably higher than that of hydrogen peroxide (H₂O/H₂O₂ = -1.78 eV) and therefore its direct oxidation by haloperoxidases (i.e. the most common route to form C–X bonds in the case of Cl, Br, and I)^[29a] is largely precluded. On the other hand, F shows the highest hydration enthalpy among halogens (H_{hyd} = -506 kJ mol⁻¹), rendering the enzymatic mechanism of desolvation a critical step prior to accomplish nucleophilic catalysis in an aqueous environment. The fluorinase enzyme was found to be able to overcome these barriers by catalyzing the transfer of F from inorganic salts to the universal methyl donor SAM through an S_N2 substitution reaction.^[66] According to the currently accepted catalytic mechanism,^[67] the fluoride ion (F⁻) is desolvated by replacing water molecules for hydrogen bonding contacts within the active site of the fluorinase. The reaction then proceeds by establishing a C–F bond between the fluoride anion and the electropositive 5'-C of the ribose ring from a SAM molecule — thereby forming 5'-fluoro-5'-deoxyadenosine (5'-FDA) and simultaneously releasing L-methionine.^[68] The canonical biofluorination pathway in *S. cattleya* further uses 5'-FDA as the key fluorinated precursor to synthesize fluoroacetate or 4-fluorothreonine (**Fig. 3**).

Our current knowledge on the mechanistic insights of fluorinase are supported by structural,^[65] kinetic^[69] and quantum mechanics/molecular mechanics theoretical data,^[70] but some questions and catalysis details of the fluorination reaction remain open. For example, the underlying reasons for the preference of the enzyme for F⁻ over other halides are still to be unraveled and may be attributed to the presence of a specific pocket for the fluoromethyl group in 5'-FDA or other physicochemical factors, e.g. specific hydrogen-bonding patterns and the electrostatic environment of the active site.^[70] At the structural level, it would appear that adopting a trimeric structure is essential for the activity of fluorinase, since SAM binds at the interface of this trimer.^[65] However, the native quaternary structure of the enzyme has determined to be not a trimer but actually a dimer of trimers; whether this hexameric conformation is also required (or at all relevant) for activity is still unknown. Crystallization studies showed that adenosine binds tightly the active site and treatment with deaminase is necessary to obtain the apoenzyme.^[71] Thus, most of the available structures of the fluorinase depict the enzyme bound to either the organic substrate (SAM) or the products (i.e. 5'-FDA and L-methionine). Structural data indicate that SAM is buried in a *closed* conformation of the fluorinase, and no evidence has been gathered to indicate the presence of appropriate channels facilitating the entrance of substrates or the release of products. This feature led to the prediction of an alternative *open* conformation of the enzyme that has not been crystalized as of yet.^[65] Three molecules of SAM have been found per trimer, and each SAM molecule is bound by a combination of hydrogen bonds and van der Waals contacts with the N- and C-terminal domains of two different monomers. It has been suggested that these interactions are relevant to keep the trimer assembled. Along this line, insights on the buried SAM structure will be particularly useful to unravel mechanistic details of the fluorinase, since the conformation of the bound molecule has been defined as exceedingly rare.^[70] Recently, the catalytic mechanism proposed for the fluorinase has been challenged, questioning the role of anion solvation/desolvation by adopting a SAM-dependent chlorinase (supposed to operate through a mechanism to form C–Cl bonds analogous to that of the fluorinase) as the model.^[72] The authors reported theoretical calculations indicating that the solvation of the Cl⁻ anion is even higher in the protein environment than in the aqueous solution. Thus, the enhancement of Cl⁻ reactivity due to the desolvation effect is largely questioned by these figures. Reproducing and discussing these calculations for the specific case of F⁻ and the fluorinase enzyme would be of great interest towards its functional implementation as a source of fluorometabolites.

As it will be discussed later, fluoroacetate and 4-fluorothreonine are promising fluorometabolites towards the implementation of *neo*-metabolism. However, despite the huge step forward that the chemical mechanism brought about by the fluorinase represents for green synthesis of fluorochemicals, the potential of the native enzyme is constrained by some intrinsic limitations. On the one hand, the affinity of the enzyme for fluorine was determined to be very low, with a K_m about 8.5 mM, a value above the maximum concentration of fluorine salts tolerated by most of the traditional microbial hosts used for metabolic engineering. On the other hand, the intracellular concentration of SAM, determined to be in the μM range in *E. coli* cells,^[73] appears to be too low to sustain biosynthesis of fluorometabolites. Therefore, efforts are currently being targeted to surpass these constraints, including the identification of novel, more efficient fluorinase orthologues, deep protein engineering of the enzyme and manipulation of intracellular SAM levels via SAM transporter systems as disclosed below.

To date, only four different fluorinases have been isolated and fully characterized *in vitro*;^[74] all of them showing similar kinetic properties (**Table 1**) — thus displaying similar catalytic limitations. Directed evolution has been applied to enhance the properties of the archetypal fluorinase, and engineered versions of the enzyme have been isolated, displaying improved specificity towards 5'-chloro-5'-deoxyadenosine^[17c] and SAM analogues^[75] as alternative substrates. Despite advances in substrate diversification, further improvements, needed to reach catalytic efficiency high enough to be compatible with bioproduction of fluorochemicals, will face significant challenges. For instance, elegant approaches developed for *in vitro* SAM recycling^[76] are not appropriate to boost biofluorination, since these strategies only consider the typical role of SAM as methyl donor — whereas it functions as a nucleoside (5'-deoxyadenosyl) donor for the fluorinase, releasing L-methionine during the reaction (**Fig. 3**). Separate efforts have focused on enhancing the intracellular SAM content *in vivo*, e.g. by pathway engineering^[77] or by increasing ATP formation and cycling.^[73,78] SAM availability may be also increased by combining external addition of the cofactor with the expression of genes encoding the rare SAM transporters known to date.^[79] A related challenge towards efficient biofluorination is the presence of SAH, since this metabolite acts as a strong competitive inhibitor of the fluorinase.^[59] SAH is a product resulting from reactions where the methyl group of SAM is transferred to an acceptor,^[76a] therefore any strategy towards increasing SAM levels in a given microbial host should consider the enzymes that use the cofactor as methyl donor and how a SAM increase affects the intracellular SAH levels.

A complementary approach would be to boost the intracellular levels of fluoride to overcome the barrier imposed by the large K_m of the fluorinase, although this strategy is in obvious conflict with the well-known toxicity of fluoride for both bacteria and eukaryotic cells.^[80] Fluoride channels specialized in exporting any excess of this anion are widespread across bacteria, archaea and eukaryotes.^[81] Suppressing these systems and studying fluoride tolerance and toxicity mechanisms in the absence of secretion processes is thus of prime importance. The work by Baker and co-workers^[82] in *E. coli* showed that once the *crcB* transporter is removed, which encodes a putative fluoride exporter, the tolerance to the chemical dropped from a minimum inhibitory concentration of 200 mM to approximately 1 mM NaF. Unfortunately, mechanisms underlying fluoride tolerance in microorganisms are not well-understood and the secretion of the intracellular anion is the only one unequivocally established.^[82-83] Another major discovery in the overall bacterial response to the presence of fluoride are riboswitches, RNA molecules that sense fluoride, present in three branches of the tree of life. These sensors have been revealed to act as triggers of the expression of genes encoding either fluoride exporters or some enzymes inhibited by F^- . Riboswitches have been also associated with stress response genes and DNA repair genes.^[84] Liao and co-workers^[85] published several studies on the resistance of the cariogenic bacterial species *Streptococcus mutans* to fluoride, a typical additive in toothpaste to prevent cavities.^[86] Besides the presence of specific exporters, the research published by these authors also suggests single point mutations in two key enzymes of the glycolytic pathway, pyruvate kinase and enolase, to be potentially involved in fluoride tolerance.^[87]

Two cases of metabolic engineering approaches involving the fluorinase have been reported in the literature. Eustáquio and co-workers took advantage of the fact that *Salinispora* sp.^[88] encodes a SAM-dependent chlorinase, which operates through a mechanism analogue to the one of the fluorinase, as the first step in synthetic pathway of the anticancer agent salinosporamide A. Replacing the gene encoding this chlorinase by the fluorinase gene from *S. cattleya* led to biosynthesis of fluorosalinosporamide. More recently, the expression of the fluorinase gene in an *E. coli* strain resulted in the production of 5'-FDA, but this approach towards *in vivo* fluorination required the implementation of some of the strategies mentioned above. Specifically, removing the CrcB transporter and externally feeding the cells with SAM (upon expression of the gene encoding a SAM transporter from *Rickettsia prowazekii*) were essential manipulations to enable 5'-FDA synthesis.^[89] Further efforts are thus necessary to substantially increase the intracellular concentrations of SAM and fluoride and to improve

the catalytic properties of fluorinase enzymes towards establishing *neo*-metabolism for biosynthesis of fluorochemicals. While *de novo* biofluorination may be a challenging endeavor, several fluorometabolites and fluorinated compounds, discussed in the next section, could become attractive nodes to engineer novel pathways using them as substrates and intermediates.

3.2. Naturally-occurring and synthetic fluorochemicals as a starting point for *neo*-metabolism

Fluoroacetate and 4-fluorothreonine are the known end products of the canonical biofluorination route of *S. cattleya* (Fig. 3). Both compounds show potential to be starting points of synthetic pathways that override central metabolism towards production of added-value fluorinated products. On this premise, hypothetical and demonstrated applications for these fluorochemicals (and related molecules) are likewise discussed in the next two sections below.

3.2.1. Fluoroacetate and other low-molecular-weight fluorometabolites

Fluoroacetate is a promising low-molecular-weight organofluorine to be used as a substrate for *neo*-metabolism. As mentioned above, the small size of F atoms allow them to replace hydrogen in C–H bonds without distorting molecular structure and many enzymes are still able to recognize the fluorinated counterpart of their natural substrates (although the catalytic efficiency usually decreases).^[90] A recent report illustrates this fact, indicating how fluorinated derivatives of anthranilic acids can be accepted as substrates for the biosynthesis of aurachin, whereas the chlorinated versions decrease product formation to negligible levels.^[91] On this background, routes that recognize acetate as the native substrate are likely to accept the fluorinated version of the acid. Indeed, fluoroacetate is easily and rapidly activated to fluoroacetyl-coenzyme A (CoA) by acetyl-CoA synthetase. The toxicity of fluoroacetate is connected to its conversion to the CoA-activated form; this metabolite enters the tricarboxylic acid (TCA) cycle (since it is recognized by citrate synthase) and further metabolized into 2-fluorocitrate. In turn, 2-fluorocitrate strongly inhibits aconitase, interrupting the TCA cycle and causing acute toxicity in a variety of organisms.^[92] Natural resistance to fluoroacetate is known to rely on the absence of enzymes from the glyoxylate cycle,^[93] acetyl-CoA synthetase variants endowed with different specificities^[94] or the presence of specific fluoroacetate dehalogenases.^[5,95] Strategies for assembling biosynthetic pathways from fluoroacetate *in vivo* would thus require the activation of the substrate while avoiding defluorination and

1 creation of metabolic dead-ends. Besides by-passing the entrance of fluoroacetyl-CoA into the TCA cycle,
2 another interesting possibility would be to engineer an aconitase insensitive to the inhibition exerted by
3 2-fluorocitrate or a citrate synthase variant unable to fit fluoroacetyl-CoA in its active site. Indeed, a recent
4 report highlights the value of rationally designing fluorometabolites that bind to citrate synthase with
5 different degrees of affinity.^[96] The number of applications of fluoroacetyl-CoA as starting point for *neo*-
6 metabolism will increase once the toxicity of fluoroacetate is avoided (or, at least, attenuated). According
7 to the KEGG database,^[97] acetate participates directly in 139 biological reactions, and the figures reach
8 a staggering 215 reactions when acetyl-CoA is considered. The group of Michelle Chang reported an
9 elegant proof of concept on the incorporation of F into biomolecules starting from fluoroacetate.^[98] The
10 approach consisted on assembling a simple pathway to convert fluoroacetate into fluoromalonate *via*
11 activation into fluoroacetyl-CoA followed by a carboxylation step. Then, fluoromalonate was used by an
12 engineered polyketide synthase as the extender unit for the synthesis of fluorinated polyketides. In a
13 more recent work from the same group,^[99] the accumulation of fluorinated poly(3-hydroxybutyrate) was
14 achieved by feeding fluoromalonate to an engineered *E. coli* strain expressing the PHA biosynthesis
15 machinery.^[100] This strategy required assembling an heterologous pathway composed by malonate:CoA
16 ligase, acetoacetyl-CoA synthase, acetoacetyl-CoA reductase and PhaC polymerase, together with the
17 expression of a transporter to uptake the externally added fluoromalonate. While the field of
18 fluorometabolite biosynthesis is benefiting from sophisticated synthetic biology and metabolic
19 engineering strategies towards new-to-Nature compounds,^[101] the incorporation of F into proteins has
20 been reported since long ago as discussed in the next section.

3.2.2. Synthesis of fluoro-amino acids and their incorporation into peptides

Fluoro-amino acids (FAAs) are non-canonical HAAs containing one or more F atoms. As indicated previously, the only known naturally-produced FAA is 4-fluorothreonine (including not only the formation of this fluorometabolite by *Streptomyces* species, but also the assembly of a synthetic route *in vitro*^[102]), although many different synthetic FAAs have been chemically synthesized.^[103] These unnatural amino acids are particularly interesting for the purpose of xenobiology and the design of *neo*-metabolism. Besides, their use provides valuable hints on how information-based process allowed for the genetic information to emerge and propagate.^[104] Many FAAs can be directly incorporated into polypeptides by the native translational machinery of cells instead of their natural proteinogenic counterparts. On this background, biosynthesis of fluorinated proteins has been demonstrated *in vivo*, and the integration of FAAs into polypeptides can be accomplished by means of two different approaches: *residue-specific incorporation* and *site-specific incorporation*.

Residue-specific incorporation (**Fig. 4**) is based on the addition of the FAA to the culture medium of a yeast or bacterial strain rendered auxotrophic for the canonical, non-halogenated counterpart. When grown in a minimal medium (i.e. where the auxotrophic strain cannot synthesize the corresponding amino acid), the cells undergo a global replacement of the natural amino acid in the microbial proteome by its fluorinated analogue. A usually adopted variant of the technique consists of growing the auxotroph with a limited concentration of the natural amino acid and, when the residue is totally consumed, the FAA is added and the production of the protein of interest is concomitantly induced. By using this strategy, the effects of fluorination in a recombinant protein can be evaluated while minimizing the negative effects on cell viability due to misincorporation. The list of FAAs assimilated following this strategy is surprisingly long and not only includes monofluorinated residues (e.g. fluoroproline,^[105] fluorohistidine,^[106] fluoroleucine,^[107] fluorotyrosine,^[108] fluorophenylalanine^[109] and fluorotryptophan^[110]), but also polyfluorinated amino acids (e.g. trifluoroleucine,^[111] hexafluoroleucine,^[112] trifluoroisoleucine,^[113] and trifluorovaline^[114]). In fact, one of the first successful reports on the *in vivo* production of fluorinated proteins was achieved by Rennert and Anker^[115] by feeding trifluoroleucine to cultures of an *E. coli* strain auxotrophic for L-leucine. Most of the studies listed above have been developed in different *E. coli* strains; the adaptability of this bacterial host to FAA even allowed for the production of a recombinant lipase with simultaneous replacement of three canonical amino acids by their monofluorinated equivalents. In this

case, proline, phenylalanine and tryptophan were substituted by their synthetic counterparts 4-(S)-fluoroproline, 4-fluorophenylalanine and 6-fluorotryptophan. The fluorinated version of the lipase was subsequently purified and characterized, and the variant was shown to keep its enzymatic activity — although the triple-fluorinated enzyme was produced at a remarkably low levels when compared with the mono-substituted version.^[116]

Site-specific incorporation (**Fig. 4**) enables total control on the position where the FAA will be inserted, but it also requires engineering the translation machinery of the host by expressing an exogenous pair of aminoacyl-tRNA synthetase (aaRS) and tRNA. This pair should be completely orthogonal to the host machinery, unambiguously recognizing the FAA without showing (ideally, any) cross-reactivity with the canonical amino acids. This requirement dictates a major difference with the residue-specific approach; the former strategy relied on the similarity between the fluorinated and the canonical residue, but certain structural differences may prove beneficial to achieve orthogonality.^[117] To date, most of site-specific systems have been adapted to recognize the *UAG* amber codon, which is recorded in the desired positions of the target protein.^[118] This system is limited by the fact that it only allows for the incorporation of a single FAA species (or, for that matter, any single, non-canonical version of the residue). To overcome the disadvantage, more *blank* codons are necessary and new strategies have been developed based on quadruplet codons^[119] — that, despite of being functional, still require considerable improvement in their efficiency.^[120] The first FAA subjected to site-specific incorporation was *p*-fluorophenylalanine by expressing a tRNA^{Phe}(*Amber*)/phenylalanyl-tRNA synthetase pair from yeast in *E. coli*, as described in the pioneer work of Furter.^[121] Improved orthogonality was later achieved by Young and co-workers^[122] by applying an evolved *Methanococcus jannaschii* aaRS-tRNA pair that suppresses the amber stop codon, and enables the incorporation of *p*-fluorophenylalanine (among other unnatural amino acids) into peptides. Similarly, another evolved aaRS-tRNA pair from *M. jannaschii* was engineered for the site-specific incorporation of F_n-tyrosine.^[123] Such strategy has been also applied to the incorporation of multifluorinated FAAs (e.g. trifluoromethyl-phenylalanine^[124]) and even more heavily fluorinated phenylalanine derivatives.^[125] As indicated for the residue-specific strategies, most of the former studies were developed in engineered *E. coli* strains. Thus far, most of the site-specific developments for FAAs have been focused on derivatives of phenylalanine and tyrosine with different degrees of fluorination and other substituents.^[126] Nevertheless, site-specific incorporation has been also achieved for aliphatic FAAs (e.g. N ϵ -acetyl-L-lysine), and trifluoroacetyl-L-lysine has been demonstrated

1 to be incorporated in proteins produced by *Saccharomyces cerevisiae* using the pyrrolysyl-tRNA
2 synthetase/tRNA pair from *Methanosarcina barkeri*.^[127]

3
4 The incorporation of FAAs into enzymes is often performed to gain insight into mechanistic^[128] or
5 structural^[129] details. However, from the perspective of xenobiology, the idea of adapting a living form to
6 include a FAA as one of their fundamental building blocks is highly suggestive. In this regard, the early
7 work of Wong^[130] on mutational adaptation to fluorotryptophan has been groundbreaking. In these
8 experiments, *Bacillus subtilis* strain QB928, a tryptophan auxotroph, was subjected to a sequential
9 mutagenic and selection process to identify mutants efficiently growing in 4-fluorotryptophan. In the
10 parental QB928 strain, 4-fluorotryptophan is incorporated to a much lesser extent than its natural
11 analogue. A major achievement in these experiments was the isolation of a strain (termed HR23) that not
12 only showed an enhanced fitness for the FAA, but also preferred the fluorinated variant over the natural
13 counterpart. The growth of the evolved strain HR23 was even inhibited by tryptophan in an agar plate
14 assay, and new mutants from QB928 were selected to grow in the presence of 5- and 6-
15 fluorotryptophan.^[131] However, the 20 canonical amino acids still constitute the core list of bricks for the
16 natural proteome (and they have been playing such a role for approximately 3 billion years), which
17 provides an idea of how tight should the adjustment of the cellular machinery be in order to recognize
18 and use orthogonal building blocks. Replacing a canonical amino acid by its FAA counterpart is thus
19 expected to be accompanied by some degree of loss in biological fitness. In the case of *B. subtilis* strains
20 adapted to grow in the presence of 4-fluorotryptophan, Mat and co-workers^[131] reported the detection of
21 a derived mutant restoring the initial preference for non-halogenated tryptophan. Thus, the global
22 replacement of a canonical amino acid by a FAA probably introduces a strong evolutionary stimulus to
23 recover the natural residue.

24
25 An attractive alternative to generate life forms stably integrating FAAs may proceed through a site-specific
26 approach. Instead of replacing *one* of the canonical amino acids, the idea here is expanding the natural
27 portfolio of building blocks by including a 21st (and beyond) FAA residue. Engineering specific targets of
28 the host proteome to integrate FAAs might have a beneficial effect sufficient to turn the course of the
29 evolutionary stream towards fixing F atoms in (*neo*)metabolism. **Fig. 5** schematizes this overall approach
30 and shows the possibilities of a potential future generation of F-containing enzymes. The strategy has
31 some precedents beyond the incorporation of this specific element, e.g. the results reported by Ma and

co-workers.^[132] In this proof-of-concept study, a single host was engineered to incorporate both a pathway for the biosynthesis of an unnatural amino acid and the site-specific machinery for its addition to a target enzyme, although no biocatalytic advantage was described for the emergent protein. Although the evidence reported thus far is relatively scarce, some studies dealing with *fluorinated enzymes* indicate that F addition results in enhanced catalytic properties. About four decades ago, Ring and co-workers^[133] studied the role of a tyrosine residue in the active site of the archetypal β -galactosidase enzyme of *E. coli* by globally replacing tyrosine residues by *m*-fluorotyrosine. Interestingly, the resulting «fluoroenzyme» showed a several-fold activity enhancement on phenyl- β -D-galactopyranoside and *p*-nitrophenyl- β -D-galactopyranoside as compared to the non-fluorinated version. Similarly, the replacement of the catalytic tyrosine of a Δ^5 -3-ketosteroid isomerase of *Comamonas (Pseudomonas) testosteroni* by 2-fluorotyrosine and the incorporation of 3-fluorophenylalanine into the *PvuII* endonuclease (both heterologously produced in *E. coli*), led to 2-fold increase in their specific activities.^[134] More recently, all phenylalanine residues in the S5 phosphotriesterase were replaced by *p*-fluorophenylalanine by expressing the gene encoding the esterase in a phenylalanine-auxotroph derivative of *E. coli*, resulting in a substantial improvement in catalytic efficiency against two different substrates.^[135]

As interesting as catalytic improvements in fluorinated proteins are, they are also hardly predictable, requiring detailed knowledge on structure and mechanism of the target enzyme. In contrast, the «fluorous effect», i.e. fluorophilicity towards highly fluorinated organic molecules, has been proposed by Budisa and co-workers^[136] as a robust way to enhance enzyme stability. The approach consists in replacing the hydrophobic core of a certain protein by a *fluorous core* formed by perfluorinated amino acids. The trifluoromethyl group ($-\text{CF}_3$) is almost twice as hydrophobic as the methyl group ($-\text{CH}_3$), suggesting that a protein with a fluorous core would be subjected to hydrophobic collapse and therefore to proper folding driven by exclusion from both aqueous and organic phases.^[136] Enhancements in stability associated to this rationale have been confirmed by multiple studies that explored the effects of including FAAs on proteins. Some of the reported improvements affected refoldability and thermoactivity,^[135] thermostability and organic solvent tolerance,^[137] thermal and chemical stability,^[111] folding efficiency^[138] and global stability.^[139] Indeed, Buer and co-workers^[140] expanded on the molecular basis of the stabilization effect of a perfluorinated protein core and found an upgraded packing efficiency of side chains that increases the buried hydrophobic surface area. The same study lists cases where FAAs incorporation had no

positive effect on protein stability — i.e. when F atoms have no impact on the relation between the volume occupied by the peptide chains and the total volume of the core.

Despite the potential benefits brought about by the presence of F on protein properties, the bioproduction of fluorinated polypeptides is severely hampered by the dearth of FAAs stemming from biological routes. Currently, the broad repertoire of FAAs mostly encompasses amino acids exclusively produced by chemical synthesis.^[141] As indicated above, FAAs should be externally added to the culture to force their incorporation into the proteome, which is considered a disadvantage in terms of cost and sustainability.^[142] Deciphering regulatory details of the 4-fluorothreonine biosynthetic pathway in *S. cattleya* could be a major breakthrough to overcome this limitation, but this is not the only route towards *in vivo* production of FAA via *neo*-metabolism. In this sense, the use of tyrosine phenol-lyases is remarkable because of its simplicity. Fluorinated tyrosine can be synthesized in a single step from a fluorophenol derivative, pyruvate and NH₃. VonTersch and co-workers^[143] described the synthesis of different mono-, di- and tri-fluorinated derivatives of L-tyrosine using a tyrosine phenol-lyase from *Citrobacter freundii*, although the yields ranged between 10%-42% in comparison to the synthesis of the natural amino acid. Interestingly, in a later work, the same tyrosine phenol-lyase mediated 99% conversion of 2-fluorophenol into 3-fluorotyrosine, whereas the yield of tyrosine on phenol was limited to 40%.^[144]

The canonical reaction of phenylalanine dehydrogenases is the interconversion between phenylpyruvate with NH₃ and phenylalanine with NADH as cofactor. However, protein engineering has proven to be an effective strategy to expand the promiscuity of these enzymes to accept different 2-oxoacids as substrates.^[145] Based on this potential, Busca and co-workers^[146] used three different mutant versions of phenylalanine dehydrogenase from *Bacillus sphaericus* to synthesize 2-, 3- and 4-fluorophenylalanine from the corresponding fluorophenylpyruvate derivatives and NH₃. Another engineered version of the enzyme from *B. sphaericus* displayed the ability to synthesize 4-fluorophenylalanine together with enhanced stability against organic solvents.^[147] Paradisi and co-workers^[148] established a more sophisticated system, combining mutant versions of the orthologue from *B. sphaericus* with an alcohol dehydrogenase to oxidize ethanol to acetaldehyde — a cost-effective NADH regeneration strategy that enabled millimole-scale synthesis of *p*-fluorophenylalanine and *p*-CF₃-phenylalanine.

Fluorotryptophan can be also synthesized in a one-pot reaction from serine and a fluoroindole derivative using a tryptophan synthase as the biocatalyst. Goss and co-workers^[149] developed a method to produce fluorotryptophan using a cell lysate of *E. coli* expressing the gene encoding a tryptophan synthase from *Salmonella enterica*. By following this strategy, production of 5-, 6- and 7-L-fluorotryptophan was scaled up to gram quantities with yields on indole ranging from 40 to 80%. Biosynthesis of 5-fluorotryptophan by the tryptophan synthase from *S. enterica* has been also performed using immobilized enzyme and in catalytic biofilms,^[150] the latter with high conversion yields above 90%.^[151] In another recent and very suggestive study, Fang and co-workers^[152] discussed the potential of type II aldolases for the synthesis of valuable fluorinated compounds. The approach proposed by the authors relies on the ability of aldolases to form C–C bonds, coupling Pyr-3-F to different aldehydes with very high selectivity. Two different β -fluoroamino acids (among other fluorinated structures that can be obtained by adopting this approach) were synthesized by combining the activities of an aldolase from *E. coli* or *Sphingomonas wittichi* with a transaminase from *Vibrio fluvialis*. β -Amino acids, with or without F, cannot be directly used for the synthesis of natural polypeptides, but β -fluoroamino acids are endowed with physicochemical properties that make them attractive for medical applications.^[153]

Pyr-3-F is also a key substrate for the synthesis of fluoroalanine. Early works showed that (*R*)-3-fluoroalanine can be easily produced from Pyr-3-F and NH_3 in a one-pot reaction using alanine dehydrogenase, although this method consumes the expensive cofactor NADH in stoichiometric quantities. Ohshima and co-workers^[154] overcame this drawback by coupling the synthesis of (*R*)-3-fluoroalanine with enzymatic oxidation of formate to CO_2 , a process catalyzed by a NAD^+ -dependent formate dehydrogenase. The system was successfully transferred to a continuous reactor using alanine dehydrogenase from *B. sphaericus* and a formate dehydrogenase from yeast. More recently, a different scheme was developed to completely circumvent redox cofactor requirements. In this one-step reaction, the synthesis of (*R*)-3-fluoroalanine was catalyzed by the ω -transaminase from *V. fluvialis* using Pyr-3-F and (*S*)- α -methylbenzylamine as substrates.^[155] Interestingly, the synthesis of (*S*)-3-fluoroalanine is more challenging and it is performed through a chemoenzymatic approach. The first step consists of reductive amination of Pyr-3-F to form *rac*-fluoroalanine, a mixture of the *R*- and *S*-isomers. (*R*)-3-fluoroalanine is then converted into Pyr-3-F and NH_3 by alanine dehydrogenase — now catalyzing the reverse reaction and reducing (rather than oxidizing) NAD^+ . The last step is catalyzed by L-lactate dehydrogenase, forming (*R*)-3-fluorolactic acid and oxidizing NADH.^[155] Unlike (*R*)-3-fluoroalanine, the (*S*)-isomer cannot replace

1 L-alanine in biosynthesized polypeptides, but this chemical species is considered as an antibiotic because
2 of its capacity to irreversibly inactivate bacterial racemases.^[156]

3
4 Despite using different biocatalysts and substrates, one essential aspect of *in vitro* biosynthetic
5 approaches distinguishes them from the pathway towards 4-fluorothreonine synthesis. All of these
6 reaction sequences require a fluorinated organic substrate, whereas 4-fluorothreonine synthesis begins
7 with inorganic fluorine *via* the fluorinase reaction. Fluorosubstrates needed for the (partially bio-based)
8 synthesis of FAAs are obtained by chemical synthesis in ways that are more cost-effective, simpler and/or
9 more environmentally friendly when compared to FAAs production by purely chemical means. From the
10 perspective of xenobiology and *neo*-metabolism, connecting these biosynthetic strategies with inorganic
11 fluorine would be a major achievement and the canonical biosynthetic route of *S. cattleya* is key to this
12 purpose. The end-metabolite of the route is fluoroacetate and, as mentioned above, fluoroacetate and its
13 activated derivative fluoroacetyl-CoA may be subjected to a number of metabolic conversions towards
14 fluorometabolites. In this context, the reactions within the TCA cycle, which uses acetyl-CoA as the
15 starting C2 unit and whose intermediates are channeled into the synthesis of several amino acids,^[157]
16 could be exploited for incorporation of F into different chemical structures. This strategy is a promising
17 source of reactions to engineer a fluorinated proteome, yet the inhibition of aconitase by 2-fluorocitrate is
18 still an unsolved problem. Additionally, metabolic adjustments and engineering of key enzymes are likely
19 necessary to efficiently produce a certain FAA to be part of a specific protein without exerting a negative
20 impact on fitness. While these challenges are being addressed, fluorotyrosine is probably the most
21 promising FAA for xenobiology developments in the short term.

22
23 Unlike other FAAs, fluorotyrosine is not only enzymatically synthesized with comparatively high yields and
24 in one-step reactions from fluorophenol, a cost-effective substrate,^[144] but there is also an orthogonal
25 system for site-specific incorporation of F_n-tyrosine in reassigned amber codons.^[123] In this sense, its
26 potential for xenobiology is superior to that of 4-fluorothreonine itself, which lacks an available system for
27 site-specific incorporation. In *S. cattleya*, natural producer of 4-fluorothreonine, this FAA is
28 misincorporated in proteins in lieu of L-threonine, and a fluorothreonyl-tRNA deacylase that hydrolyzes
29 fluorothreonyl-tRNAs is needed to limit the potential toxicity of this FAA.^[158] On the contrary, and as
30 mentioned above, incorporation of fluorotyrosine has been demonstrated to improve the catalytic
31 properties of certain enzymes. A strategy to exploit such advantages might consist of developing a

biocatalyst, either by rational design^[159] or by directed evolution,^[160] in which the presence of fluorotyrosine causes a significant catalytic enhancement or confers the ability to mediate a non-natural reaction.^[161] Such an engineered enzyme should be expressed in a microbial host carrying both the site-specific system for FAA incorporation and the biosynthetic machinery recognizing fluorotyrosine. By following this strategy, it may be even possible to revert the normal evolutionary stream, switching from the natural tendency of avoiding the presence of F (i.e. through the mechanisms indicated in the previous sections, including fluoride transporters,^[81] dehalogenases^[162] or fluoroaminoacyl-tRNA deacylases^[158]) to the positive selection of mutants displaying enhanced ability to use fluorine and fluorometabolites. In other words, a change from fluorine-tolerance to fluorine-dependence would constitute a giant leap forward for xenobiology and the implementation of *neo*-metabolism for biofluorination.

3.2.3. Integrating F into other macromolecules and alternative F-based life forms

Proteins are just one example of polymers that could accept fluorinated monomers. XNAzymes are a remarkable example of macromolecules where the presence of F could bring about emergent properties. Xenonucleic acids (XNAs) are genetic polymers with backbone structures different from those associated to DNA and RNA.^[163] Similarly, XNAzymes are the xeno-analogues of ribozymes and DNAzymes^[164] — folded polymers build of unnatural nucleic acid units able to catalyze a certain reaction. Currently, the number of XNAzymes is limited, but, interestingly, one of them is the group of 2'-fluoroarabino nucleic acid enzymes or FANAzymes. These fluorinated biocatalysts have been reported to cleave RNA and, more recently, evolved versions have proven to attack chimeric DNA substrates.^[165]

The wide variety of fluorinated structures (both monomeric and polymeric) poses an exciting question for xenobiologists: could F serve as the chemical basis of (alternative) biological systems? The suggestive observations of Budisa and co-workers^[166] highlighted the possibility that life could develop in a F-rich planetary environment. The authors based their analysis on the physicochemical traits of this element, e.g. its potential to be incorporated into cellular proteomes and the fact that perfluorinated molecules form their own «fluorous phase», different from both aqueous and hydrocarbon phases. The evidence would indicate that F could be a main constituent of (alternative) life forms, provided that it is sufficiently abundant. The study also points that aqueous HF could be a suitable solvent for the emergence of

biologically-relevant reactions, although both inorganic and organic biochemistry in such a chemical environment would be quite different from the ones we know on Earth.

4. Silicon

Historically, Si is the element that has aroused more fascination than any other atom — not only from the perspective of xeno- but also exobiology. Several scientific reports have discussed its potential role in alternative life forms,^[167] and the very expression *in silico* alludes to silicon and it was coined in 1987 to refer to artificial living systems.^[168] This interest is based on its similarities to C, the essential backbone element in biomolecules. C and Si are the first and second members of Group 14 in the periodic table, respectively (**Fig. 1**). As deduced by their positions in the periodic table, both elements share the same number of valence electrons, although there is a crucial difference between the electron configurations of C ($1s^2 2s^2 2p^2$) and Si ($1s^2 2s^2 2p^6 3s^2 3p^2$), implying that the available *p* orbitals of Si free to form double or triple bonds are located in the third shell. This occurrence results in a poor overlap of the orbitals, causing weaker π bonds and making unstable Si=Si, Si=C and Si=O double bonds.^[169] On the contrary, Si mirrors C regarding formation of single chemical bonds, being able to establish four covalent links, including Si–Si, Si–C, Si–H and Si–O. Despite the fact that there are no organosilicone compounds (i.e. compounds containing C–Si) produced biologically,^[170] there is a huge number of different Si-containing molecules chemically synthesized (as displayed in **Fig. 6**). Many of these organosilicones are endowed with different bioactivities,^[171] some of which will be mentioned below. Remarkably, polysilanes, molecules composed by a Si–Si backbone, exhibit an amphiphilic nature that determines their self-aggregation in water, forming vesicles and micelles that resemble cellular membranes.^[172]

In spite of the versatility of Si chemistry, this element is still an outsider of natural biochemistry, and biological mechanisms to introduce Si into organic molecules are scarce. Biologists have gained insight into the natural utilization of Si through the study of marine organisms able to produce large quantities of silicified structures. Different organisms, e.g. sponges or diatoms, possess silica skeletons formed by sequestering the orthosilicic acid $[\text{Si}(\text{OH})_4]$ present in seawater (**Fig. 6**). Interestingly, the process through which these skeletons are built up occurs at physiological conditions, it is genetically controlled and several proteins are involved.^[173] The marine demosponge *Tethya aurantia* is known to produce large quantities of silica, and it has become a model organism to decipher the process of biosilicification.^[174]

Demosponges contain specialized cells, called sclerocytes, which synthesize spicules, the basic structural element of the silica skeleton. Spicules are structures composed by an axial protein filamentous core included within the biosilica. The protein component is formed by three subunits named silicateins α , β and γ . Homologues of these proteins have been identified in other demosponges, e.g. *Suberites domuncula*, *Petrosia ficiformis*, *Lubomirskia baicalensis* and *Ephydatia fluviatilis*. Initially, silicateins were believed to possess a purely scaffolding function, acting as a template for the silica structure to grow, but the finding of a close relation between silicatein and the enzyme cathepsin L prompted the investigation of a possible biocatalytic role by these proteins.^[174] Experiments performed using the model substrate tetraethyl orthosilicate and a set of mutants in key residues of the active site unequivocally demonstrated the biocatalytic properties of silicatein and allowed to propose a general mechanism for the enzyme activity. According to the proposed model, silicatein works through a nucleophilic attack by a catalytic serine on the silicon center of the substrate causing the disruption of an ethyl radical and the formation of an enzyme–O–Si intermediate. The intermediate is broken due to a second nucleophilic attack by a water molecule, thereby releasing a silanol (R–Si–OH) group. Silanol is amenable to condensation with the silicon center of tetraethyl orthosilicate (**Fig. 6**), initiating the polymerization of silica. The final deposition of the formed silica is directed by the silicatein filaments.^[175] In a recent report, Tabatabaei Dakhili and co-workers^[176] provided mechanistic insight into the ability of silicateins to hydrolyze and condense Si–O bonds. The study highlighted the capacity of recombinant Sil α from *Suberites domuncula* in promoting hydrolysis, condensation and transesterification of a range of Si-containing substrates. The preference of the enzyme for aromatic rather than aliphatic hydroxyl groups while catalyzing their condensation was identified as a significant difference in comparison to non-enzymatic silylations. Interestingly, these properties of silicatein have proven to be adaptable to metalloids other than Si. In this way, a simple metalloid oxide could be converted into nanocrystalline polymorphs in a process carried out under mild conditions, whereas the chemical approach would require high temperatures and/or extreme pH values. Some examples of hydrolysis and polycondensation reactions catalyzed by silicatein are the formation of titanium dioxide and gallium oxide.^[174] Yet, what is the role Si in the biological synthesis of organic molecules?

4.1. Enzymatic synthesis of organosilicon compounds

The remarkable catalytic properties of silicatein still keep Si as a collateral element for biochemistry, as it mostly part of skeletons instead of being a component of metabolites. The situation changed drastically in 2016, when the first enzyme engineered to mediate the formation of C–Si bonds was reported.^[170] In the groundbreaking research carried out in the group of Frances Arnold, it was hypothesized that some heme proteins may be able to catalyze the formation of C–Si bonds. After assaying different bacterial and eukaryotic cytochrome *c* proteins, one isolate from *Rhodothermus marinus* (*Rma* cyt *c*) emerged as a candidate for this purpose — not only because of its ability to catalyze the reaction but also due to its stereocontrol, far above that in any other enzyme tested thus far. Mutagenesis of the key residue of the active site was followed by sequential site-saturation mutagenesis and led to the identification of a triple-mutant with a 15,000-fold enhancement of the initial turnover and >99% enantiomeric specificity. In the same study, the evolved *Rma* cyt *c* was successfully used to synthesize 20 silicon-containing organic products.

Little is known about the metabolic processing of organosilicon compounds (if at all relevant), but several articles describing different biotransformations of these unnatural chemicals demonstrate the existence of natural enzymes active on silano molecules despite the presence of the C–Si bond. Frampton and Zelisko,^[177] for instance, listed organisms (comprising multiple eukaryotes, especially yeast, and several bacterial species) known to execute biotransformations on organosilicon molecules. Most of these organisms were identified in an early screening of strains able to enantioselectively reduce acetyldimethylphenylsilane,^[178] although the yeasts *Kloeckera cortices* and *Trigonopsis variabilis* are the most studied organisms displaying catalytic activities on organosilicones.^[177]

In addition to whole-cell transformations mediated by these organisms, some enzymes are capable of catalyzing the conversion of organosilicon substrates. A lipase from *Candida cylindracea* uses *rac*-1,1-dimethyl-1-sila-cyclohexan-2-ol as substrate for transesterification, and the OF 360 lipase from the same yeast also employs silyl alcohols as acyl acceptors. Trimethylsilylmethanol can be esterified by lipases from *C. cylindracea*, *Rhizopus japonicus* and hog pancreas. Other lipases able to esterify specific silylated alcohols are the enzymes isolated from *Chromobacterium viscosum* and *Pseudomonas fluorescens*.^[177] Abbate and co-workers^[179] identified several commercial hydrolases that could mediate

condensation of silanols in aqueous environments through the formation of siloxane bonds (Si–O–Si). These enzymes, belonging to diverse functional groups, encompass the phytases from *Aspergillus ficuum* and *A. niger*, chicken egg white lysozyme, porcine gastric mucosa pepsin and a lipase from *Rhizopus oryzae*. The well-known penicillin G acylase enzyme was shown to selectively hydrolyze a specific phenylsilane from a racemic mixture.^[180] An alcohol dehydrogenase from horse liver also displays the ability of oxidizing silylated alcohols.^[181] Besides natural biocatalysts, enzyme engineering expanded the scope of enzymatic functionalization of organosilanes. For instance, the cytochrome P450 monooxygenase P450BM3 from *Bacillus megaterium* was recently submitted to directed evolution, and the engineered enzyme could selectively oxidize a series of organosilanes to the corresponding silanols.^[182] These examples expose the role of Si as a rare occurrence that is tolerated (rather than preferred) by selected microbial enzymes, but the presence of this atom in key metabolic intermediates (e.g. amino acids) deserves special attention because of its biological relevance.

4.2. Amino acids containing C–Si bonds as key building blocks

A particularly relevant family of organosilicon molecules for xenobiology is formed by Si-containing amino acids (SiAAs), broadly defined as residues displaying a stable C–Si bond. Unlike HAAs, no examples are known of naturally-occurring SiAAs, and no biosynthetic methods for their preparation have been reported to date. However, there is a wide variety of SiAAs produced by chemical synthesis, not only comprising α -, β - and γ -silyl- α -amino acids, but also other varieties such as *p*-(trimethylsilyl)phenylalanine or silaproline.^[183] Some of these molecules are known to possess similar physicochemical properties as those of proteinogenic amino acids, and β -trimethylsilyl alanine is considered to be a bioisostere for L-phenylalanine.^[184]

For a detailed description on synthetic methods of an extensive list of SiAAs, we recommend the recent review of Rémond and co-workers.^[183] To the best of our knowledge, there are no reports of Si-containing amino acids incorporated into proteins or large polypeptides, but there are several bioactive peptides containing SiAAs. In the same way that C–H can be replaced by C–F with minimum structural distortion, the similarities between C and Si led researchers to think that it would be possible to introduce the latter without loss of active conformation. However, according to Sieburth^[185] some intrinsic limitations should be taken into account; e.g. that Si–H bonds are easily oxidized, multiple bonds to silicon are remarkably

unstable (mainly because of the weak π bonds, as mentioned above), the presence of Si in 3- and 4-member rings makes the structure too strained, an unsaturation close to a Si atom introduces molecular instability and Si–heteroatom bonds are highly susceptible to hydrolysis. As a general consequence of these considerations, quaternary carbons in bioactive molecules are the ideal target for the C-to-Si swap.

Some reported bioactive SiAA-containing peptides include a variant of the renin inhibitor, where trimethylsilyl alanine (TMSAla) replaces a key phenylalanine residue, thereby increasing the resistance to α -chymotrypsin treatment.^[184] Septide, a version of substance P, is a neuropeptide associated with inflammatory processes and pain, a specific variety of this biopeptide containing β -trimethylsilyl alanine and proline showed reduced affinity compared to the natural peptide but, remarkably, the replacement of proline by silaproline changed pharmacological properties and fully restored the affinity.^[186] Cetrorelix is a gonadotropin-releasing hormone receptor (GnRH receptor) antagonist and it shows strong binding to GnRH receptor and enhanced duration effect when the peptide is modified by incorporating trimethylsilyl alanine into its structure.^[173] Another example of this sort is octadecaneuropeptide and its minimum sequence retaining activity (OP), which belong to the endozepin family and contain a proline important for the active conformation. The replacement of this proline residue by silaproline led to the synthesis of an analogue with a higher bioactivity than octadecaneuropeptide in terms of inducing the release of Ca^{2+} in rat astrocytes.^[183] The incorporation of SiAAs in a target peptide usually increases its lipophilicity, and this feature has been exploited to enhance the membrane permeability of some pharmacologically relevant peptides, e.g. alamethicin^[187] or the cell penetrating peptide CF-(VRLPPP)₃.^[188]

In conclusion, a Si-based biochemistry is at an early stage of development compared with, for instance, F and other halogens. However, the key elements for the development of xenobiology approaches to incorporate Si into life are already available (or their development is considerably advanced) — enzymes capable of transferring inorganic Si to organic molecules,^[170] the identification of Si-containing bioisosteres to natural molecular building blocks and a number of experimental reports on the incorporation of Si to bioactive compounds to enhance physicochemical properties.

5. Arsenic

In the same way that F and Si can be considered analogues of H and C, respectively, As is similar to P. Both chemical elements belong to Group 15 of the periodic table and occupy contiguous positions, which implies similar physicochemical properties, although the huge abundance of P in comparison to As confers a great advantage to the former to be preferred for life (**Fig. 1**). Speculations on alternative life development replacing P by As had an (apparent) major step forward in 2010 with the isolation of *Halomonadaceae* GFAJ-1, a bacterial strain supposedly replacing P by As in macromolecules in the absence of the former chemical element.^[189] However, critical assessment of the contents of this study soon led to the general opinion that the organism was simply an isolate displaying high As tolerance. Particularly enlightening contributions led to this conclusion. Reaves and co-workers^[190] carefully replicated Wolfe-Simon's culture conditions but, unlike the original work, no arsenate was detected in DNA extracted from GFAJ-1 cells. Another study put the focus on the fact that the original culture conditions contained 2.7 to 3.2 μM P from impurities. When this concentration was decreased by one order of magnitude, the strain lost the ability to grow on arsenate.^[191] Furthermore, Basturea and co-workers^[192] demonstrated that massive ribosome breakdown can provide enough P to keep cell viability. From a more theoretical perspective, Tawfik and Viola^[193] remarked the fact that arsenate esters are orders of magnitude more susceptible to hydrolysis than the P counterparts, therefore it would be highly unlikely that arsenates can replace phosphates in life macromolecules in the Earth's biosphere.^[194] Although there are enzymes known to accept the arsenate oxyanion in place of phosphate, the arsonylated products rapidly decompose because of their intrinsic instability. Interestingly, arsonates ($-\text{C}-\text{AsO}_3^{2-}$) act as analogues of phosphonates ($-\text{C}-\text{PO}_3^{2-}$) with no major differences in chemical stability. Despite the fact that C–P bonds are not involved in most biochemical processes, analogue substrates containing C–As are reported to be recognized by enzymes, and the cognate reactions can proceed as long as the transformation of the C–As bond itself is not involved.^[193] Nevertheless, As does serve a major role in microbial metabolism. Arsenate [As(V)] acts as the respiratory oxidant for certain bacteria. This phenomenon was described in *Sulfurospirillum arsenophilum* and *Sulfurospirillum barnesii*, which couple the oxidation of lactate to the reduction of As(V) to arsenite [As(III)]. The opposite process has been also reported for a number of prokaryotic species. Oxidation of As(III) to As(V) has been traditionally interpreted as a detoxification mechanism,^[195] although this process is involved in CO_2 fixation in some chemolithoautotrophs.^[196] In addition, As has been suggested to play an essential role in life,^[197] related

to methionine metabolism.^[198] These metabolic implications are likely to be the origin of naturally-occurring organoarsenic compounds that will be reviewed next.

5.1. Enzymatic synthesis of organoarsenic compounds

All the considerations listed in the preceding section seem to relegate As to a secondary role for xenobiology purposes. Nevertheless, early reported facts on biological processing of As strongly suggest that much of the biochemistry of this element and the (potential) role of As-metabolites is still unknown. The first enzymatic transformation of As experimentally identified has been methylation, a reaction by means of which inorganic As enters the organic realm, frequently in the form of a volatile metabolite. As early as 1892, Bartolomeo Gosio demonstrated the conversion of inorganic As into a volatile metabolite^[199] by the fungus *Scopulariopsis (Microascus) brevicaulis*. In 1932, this compound was identified as trimethylarsine [As(CH₃)], which became the first identified As-containing metabolite.^[200] It was not until the 1990s that arsenite methyltransferases were purified and characterized, exposing SAM as the key methyl donor for the reaction.^[201] Initially, these biocatalysts were interpreted to be part of detoxification processes, but this hypothesis has been dismissed because of the fact that biologically formed (mono)methylarsonic acid and dimethylarsonic acid are by far more cytotoxic than inorganic arsenite.^[202] Besides arsenite methyl transferases, studies on As biotransformations have identified two classes of enzymes involved in the oxidation/reduction of arsenocompounds, i.e. arsenate reductases,^[203] necessary to convert As(V) into As(III), and arsenite oxidases that transform As(III) into As(V).^[204] However, the existence of these three families of biocatalysts are insufficient to explain the synthesis of all arsenometabolites discovered up to date, which includes arsenosugars,^[205] arsenolipids^[206] and other compounds such as arsenobetaine.^[207] **Fig. 7** displays the main As-containing organocompounds indicated in this section. Despite the fact that some pathways have been tentatively proposed, involving enzymatic and non-enzymatic steps (e.g. direct oxidation by H₂O₂^[208]), evidence fully supporting the existence (and relevance) of these routes is still missing. In this context, new As-related biocatalysts may be identified in the future, providing new tools for xenobiology to exploit incorporation of As into biomolecules *via neo*-metabolism. As-based drugs (i.e. arsanilic acid) have been shown to act as effective treatments against highly relevant diseases (e.g. cancer, trypanosomiasis or syphilis),^[209] highlighting the importance of developing bio-based alternatives to chemical synthesis for As-containing chemicals. Biological parts towards such compounds could be actually hiding in plain sight: Danchin and

co-workers^[210] have explored scenarios where bacteria living in an As-replete environment could recruit biological processes involving Se and sulfur to detoxify As via the formation of innocuous monothioarsenate.

6. Other chemical elements

Apart from the chemical elements listed in the previous sections, other relevant atoms can be bio-incorporated into organic molecules — although the biological relevance of the cognate metabolites (if any) is still obscure. A detailed description of all of them is beyond the scope of this review, but some remarkable occurrences (i.e. boron [B], tellurium [Te] and metals) are summarized in the following sections.

6.1. Boron

Boron-containing compounds are scarce in nature and no biological organoboranes (i.e. molecules with at least one stable C–B bond) have been reported thus far.^[211] The few known organisms able to produce them appear to exploit the reactivity of some metabolites to spontaneously react with chemicals containing B in the environment, in a synthetic process that does not seem to be enzymatically catalyzed. The work of Kan and co-workers^[212] completely changed this state of affairs by using directed evolution approaches to engineer the cytochrome *c* from *R. marinus* (i.e. the same enzymatic scaffold used for synthesizing organosilicons, discussed in previous sections) to catalyze B incorporation into organic structures. The evolved enzyme was determined to form C–B from borane–Lewis base complexes, presenting strong enhancements both in turnover and enantiomeric ratio compared to the wild-type biocatalyst. The role of B as a constituent of proteins has also witnessed major advances. Chemical synthesis provides access to a variety of B-containing amino acids^[213] and a site-specific approach has been developed to insert *p*-boronophenylalanine into polypeptides.^[214] Interestingly, B incorporation allows engineering mechanisms for carbohydrate recognition that would be challenging for natural proteins.^[215]

6.2. Tellurium

The group of chalcogens comprises four stable elements: O, S, Se and Te. Among them, O and S are part of the canonical elements of life, and Se is intricately intertwined to biochemistry as well. Reich and Hondal^[216] published a review on the role of Se in life, explained on the basis of its physicochemical properties and reactivity; the review includes the known biotransformations of selenochemicals and Se-containing enzymes. The situation with tellurium, another chalcogen, is very different since this element is almost completely orthogonal to life despite its proximity (and similarities) to Se. To date, most efforts on Te biochemistry have focused on studying its effects on bacteria, e.g. mechanisms by which Te salts enter the cell, insights on the toxicity of the tellurite oxyanion and identification of enzymes involved in its reduction (and detoxification).^[217] From a practical perspective, biologically synthesized Te-nanoparticles have been recognized as valuable products with a number of interesting applications, and several bacterial species have been characterized as bioproducers of such nanoparticles.^[218] Nevertheless, a major interest on Te relies on the existence of a few examples of Te-containing amino acids. Telluromethionine,^[219] 2-tellurienylalanine (an isostere of L-phenylalanine)^[220] and tellurocysteine^[221] could be incorporated into proteins by the cell machinery through residue-specific approaches, without causing any significant perturbation of their native conformations. This capacity has been exploited as an approach for X-ray structure analysis of proteins, combining the individual advantages of Se and heavy metals incorporation.^[222] The study of Liu and co-workers,^[221] also relying on bioisosterism, is particularly attractive because it describes the development of a new enzymatic activity by the rational incorporation of tellurocysteine into a glutathione transferase. Interestingly, fungi tolerant to Te salts have been reported to form telluromethionine and tellurocystine in the presence of tellurite,^[219] suggesting the possibility of developing biosynthetic approaches for these residues.

6.3. Metallic elements

Artificial metalloenzymes have emerged as a powerful tool to develop non-biological reactivities. The fundamental reasoning here is that inclusion of non-native prosthetic groups into appropriate protein scaffolds can change an enzyme scope beyond what would be possible with conventional directed evolution or rational mutagenesis.^[223] Apart from the novel catalytic possibilities brought about by this strategy, such an approach can be harnessed for xenobiology. On the one hand, unnatural amino acids

have been reported as useful linkages between cofactor(s) and the protein scaffold.^[224] On the other hand, some artificial metalloenzymes carry metal atoms that are not part of the natural biochemistry — yet another example of the potential of bringing abiotic chemical elements into life. Some examples are iridium,^[225] ruthenium,^[226] palladium,^[227] osmium^[228] and rhodium.^[229] While the incorporation of metallic elements into living cells *via neo-metabolism* awaits further development, critically revisiting the role of alkali metallic ions (e.g. potassium,^[230] widespread in life) indicates that they have been selected because of some of their idiosyncratic properties rather than by an accidental evolutionary occurrence. These principles provide fundamental guidance towards engineering routes for integration of non-biological atoms into the biochemistry of microbial cells.

7. Summary and Outlook

Current demands for developing green production alternatives able to match the versatility and the cost-effectiveness of conventional organic chemistry are unfortunately not complemented by bio-based options that could fulfil these needs. We argue that intersecting xenobiology with the design of *neo-metabolism* could pave the way towards addressing this problem, and the last two decades have witnessed several examples of exogenous, non-biological elements incorporated into the chemistry of life. More members of the periodic table are expected to join the biological agenda in the years to come, and the role of new atoms is expected to be expanded as long as protein engineering, synthetic biology and metabolic engineering strategies becomes more and more sophisticated.

In this article, we have discussed the state-of-the-art for the incorporation of Cl, Br, I, F, Si and As into organic molecules, with varying degrees of development depending on the element considered. The occurrence and knowledge on the incorporation of halogens into organic compounds is deeper than for other xeno-elements, and includes several classes of enzymes for their incorporation into organic structures together with thousands of identified halogenated metabolites. Yet, why are halogens largely restricted to secondary metabolism? Their absence from naturally occurring proteins, nucleic acids and metabolites in central biochemical pathways may imply the existence of a strong evolutionary barrier — probably because, unlike F, the remaining halogens are not ideal replacements of H in C–H bonds in terms of molecular distortion. Additionally, the low abundance of all the mentioned elements with the exception of Cl and Si should be also considered and properly addressed. Although the potential of F in

neo-metabolism is hampered by the inefficiency of the fluorinase enzyme, the capacity of F to mimic H in size and bond geometry, together with the fact that most tested enzymes are able to catalyze the conversion of fluorinated substrates (despite some degree of efficiency loss), renders F one of the most promising element for xenobiology. Mechanisms for site-specific incorporation of some FAAs and studies on fluorinated proteins are particularly encouraging to engineer a host where, in a suitable environment, F confers an evolutionary advantage. A major obstacle is the difficulty of accurately predicting the effects of incorporating F to a substrate (or to the protein itself) on enzyme activity. As more information is collected and new advances are made in the field of protein structure and function, the rational assembly of a *neo*-metabolism based on the F presence will become feasible in the near future. Particularly, the introduction of modern machine-learning in the field of protein engineering holds the potential to be disruptive.^[231] In the meantime, directed evolution represents a powerful, semi-rational approach to adapt enzymes to fluorinated *neo*-metabolism and, in an even more ambitious objective, enhancing catalytic properties or developing non-natural catalysis through the incorporation of FAAs. As an example, the combination of site-specific FAA incorporation and saturated mutagenesis of an enzyme active site may lead to a library of mutants composed by at least 21 different amino acids — thereby allowing for the identification of enhanced variants carrying one or several FAAs.

The situation with pnictogens is radically different. No suitable replacement for P has been found in life, mostly due to the high instability of arsenates. In addition, the formation of C–As bonds (together with their intrinsic physicochemical properties) is still a poorly understood process, although natural enzymes involved in these transformations are expected to be isolated and characterized in the near future. The potential of Si, on the other hand, has been boosted by engineering an enzyme capable of forming C–Si bonds, although little is known about the role (if any) of silicone containing organo-compounds in biological systems.

A complementary perspective on xenobiology has been proposed by Antoine Danchin in a recent and inspiring review.^[232] Besides the incorporation of xeno-elements into biochemistry via *neo*-metabolism as discussed herein, the article proposes a role for stable CHONS isotopes, with a focus on deuterium (i.e. ^2H). The review analyzes known and potential effects of ^2H on biochemical pathways based on its physicochemical properties, together with reports on the toxicity and safety of ^2H enrichment in different species. These occurrences are epitomized in the recent article by Kampmeyer and co-workers,^[233]

1 identifying the multi-levels adaptations undergone by *Schizosaccharomyces pombe* when exposed to
2 D₂O. Other isotopes are likewise considered, and some enzymes have been suggested to differentiate
3 between ¹²C and ¹³C-containing compounds. The existence of biocatalysts displaying such a subtle
4 discrimination is particularly exciting for establishing *neo*-metabolism towards isotopic enrichment. From
5 a broader perspective, it should be noted that the current developments of xenobiology mostly rely on
6 physicochemical analogies between canonical chemical elements and their alternative partners when it
7 comes to introduce abiotic atoms into life. The overarching ambition of creating *xeno-life* is presently
8 limited to substantially modify the known biochemistry and metabolism, rather than creating an entirely
9 novel one. Incorporation of xeno-elements in enzymes and metabolites may lead to catalytic possibilities
10 completely absent in canonical life, paving the way towards fulfilling this ambitious objective.

11
12 Key to all these developments will be the adoption of robust microbial hosts that could support *neo*-
13 metabolism involving orthogonal parts and metabolites. As such, environmental bacteria, e.g.
14 *Pseudomonas putida*,^[234] emerge as attractive *chassis* for these developments. The advantages of this
15 non-pathogenic soil bacterium for handling xeno-elements have been recently reviewed.^[235] The well-
16 known capacity of *P. putida* of growing in harsh environments (including industrial wastes)^[236] and its
17 tolerance to many organic solvents are traits that can be exploited to overcome the toxicity of compounds
18 that can be exploited from *neo*-metabolism (e.g. fluoroacetate). Moreover, efficient mechanisms for
19 organohalide dehalogenation^[237] have been identified in this species and phylogenetically-related
20 isolates, suggesting that the biochemistry of *P. putida* has been evolutionarily shaped to deal with
21 halogenated structures. In addition, the robustness, interconnected and versatile central carbon
22 metabolism of *Pseudomonas*^[238] constitutes a treasure trove of enzymatic activities that can be
23 harnessed to build synthetic routes involving non-biological atoms. Using *P. putida* (and other species)
24 for this purpose will require the (rationale) design and testing of *neo*-metabolism *in vitro* followed by their
25 implantation into the extant biochemical network, combining *de novo* design with evolutionary trajectories
26 that environmental bacteria have followed for recognition of novel substrates.^[239] Further developments
27 in these areas in the near future are expected to bridge this gap towards consolidating the production of
28 new-to-Nature molecules.

Acknowledgements

The authors are indebted to Antoine Danchin and Philippe Marlière for fruitful and stimulating discussions. The work in the authors' laboratory is generously supported by The Novo Nordisk Foundation (NNF10CC1016517 and NNF18OC0034818), the European Union's *Horizon 2020* Research and Innovation Programme under grant agreement No. 814418 (*SinFonia*), and the Danish Council for Independent Research (*SWEET*, DFF-Research Project 8021-00039B) to P.I.N. M.N.D. is supported by the European Union's *Horizon 2020* Research and Innovation Programme under the Marie Skłodowska-Curie grant agreement No. 713683 (*COFUNDfellowsDTU*).

Conflict of interest

The authors declare no conflict of interest.

Keywords: Xenobiology · Metabolic Engineering · Synthetic Biology · Metabolism · *Pseudomonas putida*

References

- [1] A. P. Zeng, *Biotechnol. Adv.*, **2019**, 37, 508-518.
- [2] M. J. Smanski, H. Zhou, J. Claesen, B. Shen, M. A. Fischbach, C. A. Voigt, *Nat. Rev. Microbiol.*, **2016**, 14, 135-149.
- [3] E. T. Wurtzel, C. E. Vickers, A. D. Hanson, A. H. Millar, M. Cooper, K. P. Voss-Fels, P. I. Nikel, T. J. Erb, *Nat. Plants*, **2019**, 5, 1207-1210.
- [4] S. Y. Choi, M. N. Rhie, H. T. Kim, J. C. Joo, I. J. Cho, J. Son, S. Y. Jo, Y. J. Sohn, K. A. Baritugo, J. Pyo, Y. Lee, S. Y. Lee, S. J. Park, *Metab. Eng.*, **2020**, 58, 47-81.
- [5] P. Dvořák, P. I. Nikel, J. Damborský, V. de Lorenzo, *Biotechnol. Adv.*, **2017**, 35, 845-866.
- [6] D. Braff, D. Shis, J. J. Collins, *Adv. Drug Deliv. Rev.*, **2016**, 105, 35-43.
- [7] S. Slomovic, K. Pardee, J. J. Collins, *Proc. Natl. Acad. Sci. USA*, **2015**, 112, 14429.
- [8] C. Mora, D. P. Tittensor, S. Adl, A. G. B. Simpson, B. Worm, *PLoS Biol.*, **2011**, 9, e1001127.
- [9] K. J. Locey, J. T. Lennon, *Proc. Natl. Acad. Sci. USA*, **2016**, 113, 5970.
- [10] a) H. Machado, R. N. Tuttle, P. R. Jensen, *Curr. Opin. Microbiol.*, **2017**, 39, 136-142; b) M. Fondi, P. Liò, *Microbiol. Res.*, **2015**, 171, 52-64; c) K. Yugi, H. Kubota, A. Hatano, S. Kuroda, *Trends Biotechnol.*, **2016**, 34, 276-290.
- [11] a) K. B. Reed, H. Alper, *Synth. Syst. Biotechnol.*, **2018**, 3, 20-33; b) W. Kaim, B. Schwederski, A. Klein, *Bioinorganic chemistry—Inorganic elements in the chemistry of life: An introduction and guide*, 2 ed., Wiley, Chichester, UK, 2013.
- [12] a) C. G. Acevedo-Rocha, N. Budisa, *Microb. Biotechnol.*, **2016**, 9, 666-676; b) M. Schmidt, L. Pei, N. Budisa, *Adv. Biochem. Eng. Biotechnol.*, **2018**, 162, 301-315; c) K. Hamashima, M. Kimoto, I. Hirao, *Curr. Opin. Chem. Biol.*, **2018**, 46, 108-114; d) M. Schmidt, *BioEssays*, **2010**, 32, 322-331; e) M. Schmidt, V. de Lorenzo, *Curr. Opin. Biotechnol.*, **2016**, 38, 90-96.
- [13] V. de Lorenzo, *BioEssays*, **2014**, 36, 226-235.
- [14] a) J. W. Chin, *Annu. Rev. Biochem.*, **2014**, 83, 379-408; b) D. A. Malyshev, K. Dhami, T. Laverne, T. Chen, N. Dai, J. M. Foster, I. R. Corrêa, F. E. Romesberg, *Nature*, **2014**, 509, 385-388; c) D. A. Malyshev, F. E. Romesberg, *Angew. Chem. Internatl. Ed.*, **2015**, 54, 11930-11944; d) J. W. Chin, *Nature*, **2017**, 550, 53-60; e) J. W. Chin, *Nat. Chem. Biol.*, **2006**, 2, 304-311; f) F. Agostini, J. S. Voller, B. Koks, C. G. Acevedo-Rocha, V. Kubyshev, N. Budisa, *Angew. Chem. Internatl. Ed.*,

- 2017, 56, 9680-9703; g) K. Lang, J. W. Chin, *Chem. Rev.*, **2014**, 114, 4764-4806; h) J. C. Chaput, P. Herdewijn, M. Hollenstein, *ChemBioChem*, **2020**, In press, DOI: 10.1002/cbic.201900725.
- [15] a) T. J. Erb, P. R. Jones, A. Bar-Even, *Curr. Opin. Chem. Biol.*, **2017**, 37, 56-62; b) P. I. Nikel, *Trends Biotechnol.*, **2019**, 37, 1036-1038.
- [16] a) F. S. Kipping, W. J. Pope, *J. Chem. Soc. Trans.*, **1895**, 67, 371-398; b) A. R. Ling, J. L. Baker, *J. Chem. Soc. Trans.*, **1892**, 61, 589-593.
- [17] a) E. He, Y. Lu, Z. Zheng, F. Guo, S. Gao, L. Zhao, Y. Zhang, *J. Mat. Chem.*, **2020**, 8, 139-146; b) S. Liang, T. Kumon, R. A. Angnes, M. Sanchez, B. Xu, G. B. Hammond, *Org. Lett.*, **2019**, 21, 3848-3854; c) H. Sun, W. L. Yeo, Y. H. Lim, X. Chew, D. J. Smith, B. Xue, K. P. Chan, R. C. Robinson, E. G. Robins, H. Zhao, E. L. Ang, *Angew. Chem. Internatl. Ed.*, **2016**, 55, 14277-14280.
- [18] J. A. Field, Natural production of organohalide compounds in the environment, in: L. Adrian, F.E. Löffler (Eds.) *Organohalide-respiring bacteria*, Springer, Berlin, Germany, 2016, pp. 7-29.
- [19] G. W. Gribble, *Marine Drugs*, **2015**, 13, 4044-4136.
- [20] E. C. Fayolle, K. I. Öberg, J. K. Jørgensen, K. Altwegg, H. Calcutt, H. S. P. Müller, M. Rubin, M. H. D. van der Wiel, P. Bjerkeli, T. L. Bourke, A. Coutens, E. F. van Dishoeck, M. N. Drozdovskaya, R. T. Garrod, N. F. W. Ligterink, M. V. Persson, S. F. Wampfler, H. Balsiger, J. J. Berthelie, J. De Keyser, B. Fiethe, S. A. Fuselier, S. Gasc, T. I. Gombosi, T. Sémon, C. Y. Tzou, T. R. team, *Nat. Astron.*, **2017**, 1, 703-708.
- [21] L. Mázor, Properties, preparation and reactions of the halogens and of organic halogen compounds, in: L. Mázor (Ed.) *Analytical chemistry of organic halogen compounds*, Pergamon Press, Oxford, United Kingdom, 1975, pp. 19-52.
- [22] D. A. Shirley, *Org. React.*, **1954**, 8, 28-58.
- [23] H. Gilman, R. G. Jones, L. A. Woods, *J. Org. Chem.*, **1952**, 17, 1630-1634.
- [24] H. E. Simmons, R. D. Smith, *J. Am. Chem. Soc.*, **1958**, 80, 5323-5324.
- [25] W. J. Chung, C. D. Vanderwal, *Angew. Chem. Internatl. Ed.*, **2016**, 55, 4396-4434.
- [26] C. S. Neumann, D. G. Fujimori, C. T. Walsh, *Chem. Biol.*, **2008**, 15, 99-109.
- [27] a) C. M. Harris, R. Kannan, H. Kopecka, T. M. Harris, *J. Am. Chem. Soc.*, **1985**, 107, 6652-6658; b) M. K. Renner, P. R. Jensen, W. Fenical, *J. Org. Chem.*, **1998**, 63, 8346-8354; c) E. Rodrigues Pereira, L. Belin, M. Sancelme, M. Prudhomme, M. Ollier, M. Rapp, D. Severe, J.-F. Riou, D. Fabbro, T. Meyer, *J. Med. Chem.*, **1996**, 39, 4471-4477.

- [28] a) A. V. Fejzagić, J. Gebauer, N. Huwa, T. Classen, *Molecules*, **2019**, *24*; b) V. Weichold, D. Milbredt, K. H. van Peé, *Angew. Chem. Internatl. Ed.*, **2016**, *55*, 6374-6389; c) K. H. van Peé, *Arch. Microbiol.*, **2001**, *175*, 250-258.
- [29] a) A. Butler, M. Sandy, *Nature*, **2009**, *460*, 848-854; b) D. C. Lohman, D. R. Edwards, R. Wolfenden, *J. Am. Chem. Soc.*, **2013**, *135*, 14473-14475; c) J. W. Schmidberger, A. B. James, R. Edwards, J. H. Naismith, D. O'Hagan, *Angew. Chem. Internatl. Ed.*, **2010**, *49*, 3646-3648; d) C. Wagner, M. El Omari, G. M. König, *J. Nat. Prod.*, **2009**, *72*, 540-553.
- [30] A. S. Eustáquio, F. Pojer, J. P. Noel, B. S. Moore, *Nat. Chem. Biol.*, **2008**, *4*, 69-74.
- [31] a) S. Brown, S. O'Connor, *ChemBioChem*, **2015**, *16*, 2129-2135; b) K. H. van Pée, S. Unversucht, *Chemosphere*, **2003**, *52*, 299-312.
- [32] a) G. W. Gribble, *Acc. Chem. Res.*, **1998**, *31*, 141-152; b) G. W. Gribble, *ArkivoC*, **2018**, *2018*, 39.
- [33] G. W. Gribble, *Environ. Chem.*, **2015**, *12*, 396-405.
- [34] X. Feng, D. Bello, P. T. Lowe, J. Clark, D. O'Hagan, *Chem. Sci.*, **2019**, *10*, 9501-9505.
- [35] H. Nakajima, Y. Hori, H. Terano, M. Okuhara, T. Manda, S. Matsumoto, K. Shimomura, *J. Antibiot.*, **1996**, *49*, 1204-1211.
- [36] a) Y. Kato, Y. Kawai, H. Morinaga, H. Kondo, N. Dozaki, N. Kitamoto, T. Osawa, *Free Rad. Biol. Med.*, **2005**, *38*, 24-31; b) S. G. Venkatesh, V. Deshpande, *Comp. Biochem. Physiol. C Pharmacol. Toxicol. Endocrinol.*, **1999**, *122*, 13-20.
- [37] J. Roche, R. Michel, Natural and artificial iodoproteins, in: M.L. Anson, J.T. Edsall, K. Bailey (Eds.) *Advances in Protein Chemistry*, Academic Press 1951, pp. 253-297.
- [38] S. Mondal, D. Manna, K. Raja, G. Mugesh, *ChemBioChem*, **2020**, *21*, 911-923.
- [39] V. M. Dembitsky, M. Srebnik, *Prog. Lipid Res.*, **2002**, *41*, 315-367.
- [40] R. Goss, C. Wilhelm, Lipids in algae, lichens and mosses, in: H. Wada, N. Murata (Eds.) *Lipids in photosynthesis: Essential and regulatory functions*, Springer, Dordrecht, Netherlands, 2009, pp. 117-137.
- [41] a) A. Broberg, A. Menkis, R. Vasiliauskas, *J. Nat. Prod.*, **2006**, *69*, 97-102; b) R. Fujii, H. Yoshida, S. Fukusumi, Y. Habata, M. Hosoya, Y. Kawamata, T. Yano, S. Hinuma, C. Kitada, T. Asami, M. Mori, Y. Fujisawa, M. Fujino, *J. Biol. Chem.*, **2002**, *277*, 34010-34016; c) A. Pohanka, A. Menkis, J. Levenfors, A. Broberg, *J. Nat. Prod.*, **2006**, *69*, 1776-1781; d) M. Sjögren, U. Göransson, A.-L. Johnson, M. Dahlström, R. Andersson, J. Bergman, P. R. Jonsson, L. Bohlin, *J. Nat. Prod.*, **2004**, *67*, 368-372.

- [42] L. P. Hager, D. R. Morris, F. S. Brown, H. Eberwein, *J. Biol. Chem.*, **1966**, *241*, 1769-1777.
- [43] a) M. Hölzer, W. Burd, H.-U. Reißig, K.-H. van Pée, *Adv. Synth. Catal.*, **2001**, *343*, 591-595; b) S. A. Shepherd, B. R. K. Menon, H. Fisk, A.-W. Struck, C. Levy, D. Leys, J. Micklefield, *ChemBioChem*, **2016**, *17*, 821-824; c) S. Zehner, A. Kotzsch, B. Bister, R. D. Sussmuth, C. Méndez, J. A. Salas, K. H. van Pee, *Chem. Biol.*, **2005**, *12*, 445-452.
- [44] M. Strickland, C. L. Willis, Synthesis of halogenated α -amino acids, in: A.B. Hughes (Ed.) *Amino acids, peptides and proteins in organic chemistry: Origins and synthesis of amino acids*, Wiley, Weinheim, Germany, 2010, pp. 441-471.
- [45] I. Kwon, P. Wang, D. A. Tirrell, *J. Am. Chem. Soc.*, **2006**, *128*, 11778-11783.
- [46] P. Marlière, J. Patrouix, V. Döring, P. Herdewijn, S. Tricot, S. Cruveiller, M. Bouzon, R. Mutzel, *Angew. Chem. Internatl. Ed.*, **2011**, *50*, 7109-7114.
- [47] D. O'Hagan, *Chem. Soc. Rev.*, **2008**, *37*, 308-319.
- [48] A. Harsanyi, G. Sandford, *Green Chem.*, **2015**, *17*, 2081-2086.
- [49] M. Braun, J. Eicher, The fluorine atom in health care and agrochemical applications: A contribution to life science, in: H. Groult, F.R. Leroux, A. Tressaud (Eds.) *Modern synthesis and reactivity of fluorinated compounds*, Elsevier, London, UK, 2017, pp. 7-25.
- [50] J. C. Biffinger, H. W. Kim, S. G. DiMugno, *ChemBioChem*, **2004**, *5*, 622-627.
- [51] a) J. Wang, M. Sánchez-Roselló, J. L. Aceña, C. del Pozo, A. E. Sorochinsky, S. Fustero, V. A. Soloshonok, H. Liu, *Chem. Rev.*, **2014**, *114*, 2432-2506; b) Y. Zhou, J. Wang, Z. Gu, S. Wang, W. Zhu, J. L. Aceña, V. A. Soloshonok, K. Izawa, H. Liu, *Chem. Rev.*, **2016**, *116*, 422-518.
- [52] a) K. Haranahalli, T. Honda, I. Ojima, *J. Fluorine Chem.*, **2019**, *217*, 29-40; b) P. Richardson, *Exp. Opin. Drug Discov.*, **2016**, *11*, 983-999; c) T. Fujiwara, D. O'Hagan, *J. Fluorine Chem.*, **2014**, *167*, 16-29.
- [53] H. Teng, *Appl. Sci.*, **2012**, *2*, 496.
- [54] a) H. A. Chokhawala, H. Cao, H. Yu, X. Chen, *J. Am. Chem. Soc.*, **2007**, *129*, 10630-10631; b) J. Stockwell, A. D. Daniels, C. L. Windle, T. A. Harman, T. Woodhall, T. Lebl, C. H. Trinh, K. Mulholland, A. R. Pearson, A. Berry, A. Nelson, *Org. Biomol. Chem.*, **2016**, *14*, 105-112.
- [55] H.-S. Bea, S.-H. Lee, H. Yun, *Biotechnol. Bioprocess Eng.*, **2011**, *16*, 291-296.
- [56] J. K. Howard, M. Müller, A. Berry, A. Nelson, *Angew. Chem. Internatl. Ed.*, **2016**, *55*, 6767-6770.
- [57] C. Ding, Y. Pan, G. Zhang, J. Lv, Preparation method of 3-fluoro pyruvic acid, in: F.Z. Shenqing (Ed.), *Zhejiang University of Technology*, China, 2017.

- [58] L. Martinelli, P. I. Nikel, *Microb. Biotechnol.*, **2019**, 12, 187-190.
- [59] C. Schaffrath, H. Deng, D. O'Hagan, *FEBS Lett.*, **2003**, 547, 111-114.
- [60] a) D. L. Zechel, S. P. Reid, O. Nashiru, C. Mayer, D. Stoll, D. L. Jakeman, R. A. J. Warren, S. G. Withers, *J. Am. Chem. Soc.*, **2001**, 123, 4350-4351; b) O. Nashiru, D. L. Zechel, D. Stoll, T. Mohammadzadeh, R. A. J. Warren, S. G. Withers, *Angew. Chem. Internatl. Ed.*, **2001**, 40, 417-420.
- [61] A. Dickson, C. Goyet, Handbook of methods for the analysis of the various parameters of the carbon dioxide system in sea water, in: U.S.D.o. Energy (Ed.), 1994, pp. 187.
- [62] K. Ito, *Anal. Chem.*, **1997**, 69, 3628-3632.
- [63] a) F. Borrelli, C. Campagnuolo, R. Capasso, E. Fattorusso, O. Tagliatela-Scafati, *Eur. J. Org. Chem.*, **2004**, 2004, 3227-3232; b) C. Leblanc, C. Colin, A. Cosse, L. Delage, S. La Barre, P. Morin, B. Fiévet, C. Voiseux, Y. Ambroise, E. Verhaeghe, D. Amouroux, O. Donard, E. Tessier, P. Potin, *Biochimie*, **2006**, 88, 1773-1785; c) S. Lavoie, D. Brumley, T. S. Alexander, C. Jasmin, F. A. Carranza, K. Nelson, C. L. Quave, J. Kubanek, *J. Org. Chem.*, **2017**, 82, 4160-4169; d) P. G. Williams, W. Y. Yoshida, R. E. Moore, V. J. Paul, *Org. Lett.*, **2003**, 5, 4167-4170.
- [64] P. D. Shaw, L. P. Hager, *J. Am. Chem. Soc.*, **1959**, 81, 1011-1012.
- [65] C. Dong, F. Huang, H. Deng, C. Schaffrath, J. B. Spencer, D. O'Hagan, J. H. Naismith, *Nature*, **2004**, 427, 561-565.
- [66] K. K. Chan, D. O'Hagan, *Methods Enzymol.*, **2012**, 516, 219-235.
- [67] D. O'Hagan, H. Deng, *Chem. Rev.*, **2015**, 115, 634-649.
- [68] D. O'Hagan, H. Deng, *Chem. Rev.*, **2015**, 115, 634-649.
- [69] X. Zhu, D. A. Robinson, A. R. McEwan, D. O'Hagan, J. H. Naismith, *J. Am. Chem. Soc.*, **2007**, 129, 14597-14604.
- [70] H. M. Senn, D. O'Hagan, W. Thiel, *J. Am. Chem. Soc.*, **2005**, 127, 13643-13655.
- [71] S. L. Cobb, H. Deng, A. R. McEwan, J. H. Naismith, D. O'Hagan, D. A. Robinson, *Org. Biomol. Chem.*, **2006**, 4, 1458-1460.
- [72] E. Araújo, A. H. Lima, J. Lameira, *Phys. Chem.*, **2017**, 19, 21350-21356.
- [73] Y. Chen, H. Zhou, M. Wang, T. Tan, *RSC Adv.*, **2017**, 7, 22409-22414.
- [74] S. A. Sooklal, C. De Koning, D. Brady, K. Rumbold, *Protein Expr. Purif.*, **2020**, 166, 105508.

- [75] a) P. T. Löwe, S. L. Cobb, D. O'Hagan, *Org. Biomol. Chem.*, **2019**, 17, 7493-7496; b) W. L. Yeo, X. Chew, D. J. Smith, K. P. Chan, H. Sun, H. Zhao, Y. H. Lim, E. L. Ang, *Chem. Commun.*, **2017**, 53, 2559-2562.
- [76] a) C. Liao, F. P. Seebeck, *Nat. Catal.*, **2019**, 2, 696-701; b) S. Mordhorst, J. Siegrist, M. Müller, M. Richter, J. N. Andexer, *Angew. Chem. Internatl. Ed.*, **2017**, 56, 4037-4041; c) J. Micklefield, *Nat. Catal.*, **2019**, 2, 644-645.
- [77] a) H. Chen, Z. Wang, H. Cai, C. Zhou, *World J. Microbiol. Biotechnol.*, **2016**, 32, 153; b) G. Han, X. Hu, T. Qin, Y. Li, X. Wang, *Enzyme Microb. Technol.*, **2016**, 83, 14-21; c) M. Kanai, M. Masuda, Y. Takaoka, H. Ikeda, K. Masaki, T. Fujii, H. Iefuji, *Appl. Microbiol. Biotechnol.*, **2013**, 97, 1183-1190; d) C. Xu, Z. Shi, J. Shao, C. Yu, Z. Xu, *World J. Microbiol. Biotechnol.*, **2019**, 35, 185.
- [78] a) M. Li, X. Meng, E. Diao, F. Du, X. Zhao, *J. Chem. Technol. Biotechnol.*, **2012**, 87, 1379-1384; b) J. Zhang, X. Wang, Y. Zheng, G. Fang, D. Wei, *Bioprocess Biosyst. Eng.*, **2008**, 31, 63-67.
- [79] a) R. Binet, R. E. Fernandez, D. J. Fisher, A. T. Maurelli, *mBio*, **2011**, 2, e00051-00011; b) A. M. Tucker, H. H. Winkler, L. O. Driskell, D. O. Wood, *J. Bacteriol.*, **2003**, 185, 3031.
- [80] S. Flisfisch, J. Meyer, J. Meurman, T. Waltimo, *Oral Dis.*, **2008**, 14, 296-301.
- [81] C. B. Macdonald, R. B. Stockbridge, *Curr. Opin. Struct. Biol.*, **2017**, 45, 142-149.
- [82] J. L. Baker, N. Sudarsan, Z. Weinberg, A. Roth, R. B. Stockbridge, R. R. Breaker, *Science*, **2012**, 335, 233-235.
- [83] a) X. Men, Y. Shibata, T. Takeshita, Y. Yamashita, *PLoS One*, **2016**, 11, e0165900; b) T. Murata, N. Hanada, *FEMS Microbiol. Lett.*, **2016**, 363.
- [84] R. R. Breaker, *Caries Res.*, **2012**, 46, 78-81.
- [85] Y. Liao, B. W. Brandt, J. Li, W. Crielaard, C. Van Loveren, D. M. Deng, *J. Oral Microbiol.*, **2017**, 9, 1344509.
- [86] P. Loskill, C. Zeitz, S. Grandthyll, N. Thewes, F. Müller, M. Bischoff, M. Herrmann, K. Jacobs, *Langmuir*, **2013**, 29, 5528-5533.
- [87] Y. Liao, J. Yang, B. W. Brandt, J. Li, W. Crielaard, C. van Loveren, D. M. Deng, *Front. Microbiol.*, **2018**, 9.
- [88] A. S. Eustáquio, D. O'Hagan, B. S. Moore, *J. Nat. Prod.*, **2010**, 73, 378-382.
- [89] K. Markakis, P. T. Lowe, L. Davison-Gates, D. O'Hagan, S. J. Rosser, A. Elfick, *ChemBioChem*, **2020**, In press, DOI: 10.1002/cbic.202000051.

- [90] a) E. M. Gal, *Arch. Biochem. Biophys.*, **1960**, *90*, 278-287; b) C. L. Tober, P. Nicholls, J. D. Brodie, *Arch. Biochem. Biophys.*, **1970**, *138*, 506-514.
- [91] A. Sester, K. Stür-Patowsky, W. Hiller, F. Kloss, S. Lütz, M. Nett, *ChemBioChem*, **2020**, In press, DOI: 10.1002/cbic.202000166.
- [92] H. Lauble, M. C. Kennedy, M. H. Emptage, H. Beinert, C. D. Stout, *Proc. Natl. Acad. Sci. USA*, **1996**, *93*, 13699.
- [93] C. Ingram-Smith, S. R. Martin, K. S. Smith, *Trends Microbiol.*, **2006**, *14*, 249-253.
- [94] a) F. Huang, S. F. Haydock, D. Spiteller, T. Mironenko, T.-L. Li, D. O'Hagan, P. F. Leadlay, J. B. Spencer, *Chem. Biol.*, **2006**, *13*, 475-484; b) J. E. Turner, K. Greville, E. C. Murphy, M. A. Hooks, *J. Biol. Chem.*, **2005**, *280*, 2780-2787.
- [95] a) C. K. Davis, R. I. Webb, L. I. Sly, S. E. Denman, C. S. McSweeney, *FEMS Microbiol. Ecol.*, **2012**, *80*, 671-684; b) T. Koudelakova, S. Bidmanova, P. Dvořák, A. Pavelka, R. Chaloupkova, Z. Prokop, J. Damborský, *Biotechnol. J.*, **2013**, *8*, 32-45.
- [96] D. Bello, M. G. Rubanu, N. Bandaranayaka, J. P. Götz, M. Bühl, D. O'Hagan, *ChemBioChem*, **2019**, *20*, 1174-1182.
- [97] M. Kanehisa, S. Goto, Y. Sato, M. Furumichi, M. Tanabe, *Nucleic Acids Res.*, **2012**, *40*, D109-D114.
- [98] M. C. Walker, B. W. Thuronyi, L. K. Charkoudian, B. Lowry, C. Khosla, M. C. Y. Chang, *Science*, **2013**, *341*, 1089.
- [99] B. W. Thuronyi, M. C. Y. Chang, *Accounts Chem. Res.*, **2015**, *48*, 584-592.
- [100] N. I. López, M. J. Pettinari, P. I. Nikel, B. S. Méndez, *Adv. Appl. Microbiol.*, **2015**, *93*, 93-106.
- [101] a) E. Kim, B. S. Moore, Y. J. Yoon, *Nat. Chem. Biol.*, **2015**, *11*, 649-659; b) H. U. Kim, P. Charusanti, S. Y. Lee, T. Weber, *Nat. Prod. Rep.*, **2016**, *33*, 933-941; c) M. Sulzbach, A. M. Kunjapur, *Trends Biotechnol.*, **2020**, In press, DOI: 10.1016/j.tibtech.2019.1012.1014.
- [102] H. Deng, S. M. Cross, R. P. McGlinchey, J. T. Hamilton, D. O'Hagan, *Chem. Biol.*, **2008**, *15*, 1268-1276.
- [103] a) X.-L. Qiu, W.-D. Meng, F.-L. Qing, *Tetrahedron*, **2004**, *60*, 6711-6745; b) R.-Y. Zhu, K. Tanaka, G.-C. Li, J. He, H.-Y. Fu, S.-H. Li, J.-Q. Yu, *J. Am. Chem. Soc.*, **2015**, *137*, 7067-7070.
- [104] a) A. Danchin, *Beilstein J. Org. Chem.*, **2017**, *13*, 1119-1135; b) G. Boël, O. Danot, V. de Lorenzo, A. Danchin, *Microb. Biotechnol.*, **2019**, *12*, 210-242; c) V. de Lorenzo, A. Sekowska, A. Danchin, *FEMS Microbiol. Rev.*, **2015**, *39*, 96-119.

- [105] M. D. Crespo, M. Rubini, *PLoS One*, **2011**, 6, e19425.
- [106] J. F. Eichler, J. C. Cramer, K. L. Kirk, J. G. Bann, *ChemBioChem*, **2005**, 6, 2170-2173.
- [107] J. Feeney, J. E. McCormick, C. J. Bauer, B. Birdsall, C. M. Moody, B. A. Starkmann, D. W. Young, P. Francis, R. H. Havlin, W. D. Arnold, E. Oldfield, *J. Am. Chem. Soc.*, **1996**, 118, 8700-8706.
- [108] B. D. Sykes, H. I. Weingarten, M. J. Schlesinger, *Proc. Natl. Acad. Sci. USA*, **1974**, 71, 469-473.
- [109] C. Minks, R. Huber, L. Moroder, N. Budisa, *Anal. Biochem.*, **2000**, 284, 29-34.
- [110] J. F. Parsons, G. Xiao, G. L. Gilliland, R. N. Armstrong, *Biochemistry*, **1998**, 37, 6286-6294.
- [111] Y. Tang, G. Ghirlanda, W. A. Petka, T. Nakajima, W. F. DeGrado, D. A. Tirrell, *Angew. Chem. Internatl. Ed.*, **2001**, 40, 1494-1496.
- [112] Y. Tang, D. A. Tirrell, *J. Am. Chem. Soc.*, **2001**, 123, 11089-11090.
- [113] P. Wang, Y. Tang, D. A. Tirrell, *J. Am. Chem. Soc.*, **2003**, 125, 6900-6906.
- [114] P. Wang, A. Fichera, K. Kumar, D. A. Tirrell, *Angew. Chem. Internatl. Ed.*, **2004**, 43, 3664-3666.
- [115] O. M. Rennert, H. S. Anker, *Biochemistry*, **1963**, 2, 471-476.
- [116] L. Merkel, M. Schauer, G. Antranikian, N. Budisa, *ChemBioChem*, **2010**, 11, 1505-1507.
- [117] C. Odar, M. Winkler, B. Wiltshi, *Biotechnol. J.*, **2015**, 10, 427-446.
- [118] A. K. Antonczak, J. Morris, E. M. Tippmann, *Curr. Opin. Struct. Biol.*, **2011**, 21, 481-487.
- [119] A. Chatterjee, M. J. Lajoie, H. Xiao, G. M. Church, P. G. Schultz, *ChemBioChem*, **2014**, 15, 1782-1786.
- [120] a) E. D. Hankore, L. Zhang, Y. Chen, K. Liu, W. Niu, J. Guo, *ACS Synth. Biol.*, **2019**, 8, 1168-1174;
b) H. Neumann, K. Wang, L. Davis, M. Garcia-Alai, J. W. Chin, *Nature*, **2010**, 464, 441-444.
- [121] R. Furter, *Prot. Sci.*, **1998**, 7, 419-426.
- [122] Y.-J. Lee, M. J. Schmidt, J. M. Tharp, A. Weber, A. L. Koenig, H. Zheng, J. Gao, M. L. Waters, D. Summerer, W. R. Liu, *Chem. Commun.*, **2016**, 52, 12606-12609.
- [123] E. C. Minnihan, D. D. Young, P. G. Schultz, J. Stubbe, *J. Am. Chem. Soc.*, **2011**, 133, 15942-15945.
- [124] J. C. Jackson, J. T. Hammill, R. A. Mehl, *J. Am. Chem. Soc.*, **2007**, 129, 1160-1166.
- [125] S. J. Miyake-Stoner, C. A. Refakis, J. T. Hammill, H. Lusic, J. L. Hazen, A. Deiters, R. A. Mehl, *Biochemistry*, **2010**, 49, 1667-1677.
- [126] L. Merkel, N. Budisa, *Org. Biomol. Chem.*, **2012**, 10, 7241-7261.
- [127] S. M. Hancock, R. Uprety, A. Deiters, J. W. Chin, *J. Am. Chem. Soc.*, **2010**, 132, 14819-14824.
- [128] B. R. Howard, J. A. Endrizzi, S. J. Remington, *Biochemistry*, **2000**, 39, 3156-3168.

- [129] a) R. P. Gibson, C. A. Tarling, S. Roberts, S. G. Withers, G. J. Davies, *J. Biol. Chem.*, **2004**, 279, 1950-1955; b) R. D. Gordon, P. Sivarajah, M. Satkunarajah, D. Ma, C. A. Tarling, D. Vizitiu, S. G. Withers, J. M. Rini, *J. Mol. Biol.*, **2006**, 360, 67-79.
- [130] J. T. Wong, *Proc. Natl. Acad. Sci. USA*, **1983**, 80, 6303.
- [131] W.-K. Mat, H. Xue, J. T.-F. Wong, *PLoS One*, **2010**, 5, e12206.
- [132] Y. Ma, H. Biava, R. Contestabile, N. Budisa, L. M. Di Salvo, *Molecules*, **2014**, 19.
- [133] M. Ring, I. M. Armitage, R. E. Huber, *Biochem. Biophys. Res. Commun.*, **1985**, 131, 675-680.
- [134] a) B. Brooks, R. S. Phillips, W. F. Benisek, *Biochemistry*, **1998**, 37, 9738-9742; b) M. A. Domínguez., K. C. Thornton, M. G. Melendez, C. M. Dupureur, *Prot. Struct. Funct. Bioinform.*, **2001**, 45, 55-61.
- [135] P. J. Baker, J. K. Montclare, *ChemBioChem*, **2011**, 12, 1845-1848.
- [136] N. Budisa, O. Pipitone, I. Siwanowicz, M. Rubini, P. P. Pal, T. A. Holak, M. L. Gelmi, *Chem. Biodivers.*, **2004**, 1, 1465-1475.
- [137] a) K. Deepankumar, M. Shon, S. P. Nadarajan, G. Shin, S. Mathew, N. Ayyadurai, B.-G. Kim, S.-H. Choi, S.-H. Lee, H. Yun, *Adv. Synth. Catal.*, **2014**, 356, 993-998; b) J.-C. Horng, D. P. Raleigh, *J. Am. Chem. Soc.*, **2003**, 125, 9286-9287.
- [138] T. H. Yoo, A. J. Link, D. A. Tirrell, *Proc. Natl. Acad. Sci. USA*, **2007**, 104, 13887.
- [139] S. K. Holmgren, L. E. Bretscher, K. M. Taylor, R. T. Raines, *Chem. Biol.*, **1999**, 6, 63-70.
- [140] B. C. Buer, J. L. Meagher, J. A. Stuckey, E. N. G. Marsh, *Proc. Natl. Acad. Sci. USA*, **2012**, 109, 4810.
- [141] a) J. Hao, T. Milcent, P. Retailleau, V. A. Soloshonok, S. Onger, B. Crousse, *Eur. J. Org. Chem.*, **2018**, 2018, 3688-3692; b) F. Mansour, L. Hunter, Synthesis and applications of backbone-fluorinated amino acids, in: G. Haufe, F.R. Leroux (Eds.) Fluorine in life sciences: Pharmaceuticals, medicinal diagnostics, and agrochemicals, Academic Press, London, United Kingdom, 2019, pp. 325-347.
- [142] H. D. Biava, *ChemBioChem*, **2020**, In press, DOI: 10.1002/cbic.201900756.
- [143] R. L. VonTersch, F. Secundo, R. S. Phillips, M. G. Newton, Preparation of fluorinated amino acids with tyrosine phenol lyase: Effects of fluorination on reaction kinetics and mechanism of tyrosine phenol lyase and tyrosine protein kinase Csk, in: I. Ojima, J.R. McCarthy, J.T. Welch (Eds.) Biomedical frontiers of fluorine chemistry, American Chemical Society, Washington D.C., USA, 1996, pp. 95-104.

- [144] B. Seisser, R. Zinkl, K. Gruber, F. Kaufmann, A. Hafner, W. Kroutil, *Adv. Synth. Catal.*, **2010**, 352, 731-736.
- [145] S. Y. K. Seah, K. Linda Britton, P. J. Baker, D. W. Rice, Y. Asano, P. C. Engel, *FEBS Lett.*, **1995**, 370, 93-96.
- [146] P. Busca, F. Paradisi, E. Moynihan, A. R. Maguire, P. C. Engel, *Org. Biomol. Chem.*, **2004**, 2, 2684-2691.
- [147] S. Chen, P. C. Engel, *Enzyme Microb. Technol.*, **2007**, 40, 1407-1411.
- [148] F. Paradisi, S. Collins, A. R. Maguire, P. C. Engel, *J. Biotechnol.*, **2007**, 128, 408-411.
- [149] R. J. M. Goss, P. L. A. Newill, *Chem. Commun.*, **2006**, 4924-4925.
- [150] a) D. C. Volke, P. I. Nikel, *Adv. Biosyst.*, **2018**, 2, 1800111; b) I. Benedetti, V. de Lorenzo, P. I. Nikel, *Metab Eng*, **2016**, 33, 109-118.
- [151] A. N. Tsoligkas, M. Winn, J. Bowen, T. W. Overton, M. J. H. Simmons, R. J. M. Goss, *ChemBioChem*, **2011**, 12, 1391-1395.
- [152] J. Fang, D. Hait, M. Head-Gordon, M. C. Y. Chang, *Angew. Chem. Internatl. Ed.*, **2019**, 58, 11841-11845.
- [153] a) M. A. T. Blaskovich, *J. Med. Chem.*, **2016**, 59, 10807-10836; b) L. Kiss, F. Fülöp, *Chem. Record*, **2018**, 18, 266-281.
- [154] T. Ohshima, C. Wandrey, D. Conrad, *Biotechnol. Bioeng.*, **1989**, 34, 394-397.
- [155] L. P. B. Gonçalves, O. A. C. Antunes, G. F. Pinto, E. G. Oestreicher, *Tetrahedron*, **2000**, 11, 1465-1468.
- [156] J. Kollonitsch, L. Barash, F. M. Kahan, H. Kropp, *Nature*, **1973**, 243, 346-347.
- [157] a) J. Cronan, D. Laporte, *EcoSal Plus Online*, **2006**, (DOI:10.1128/ecosalplus.1123.1125.1122); b) R. Gerstmeir, V. F. Wendisch, S. Schnicke, H. Ruan, M. Farwick, D. Reinscheid, B. J. Eikmanns, *J. Biotechnol.*, **2003**, 104, 99-122.
- [158] J. L. McMurtry, M. C. Y. Chang, *Proc. Natl. Acad. Sci. USA*, **2017**, 114, 11920.
- [159] I. N. Ugwumba, K. Ozawa, Z.-Q. Xu, F. Ely, J.-L. Foo, A. J. Herlt, C. Coppin, S. Brown, M. C. Taylor, D. L. Ollis, L. N. Mander, G. Schenk, N. E. Dixon, G. Otting, J. G. Oakeshott, C. J. Jackson, *J. Am. Chem. Soc.*, **2011**, 133, 326-333.
- [160] a) E. M. Brustad, F. H. Arnold, *Curr. Opin. Chem. Biol.*, **2011**, 15, 201-210; b) M. T. Reetz, *Acc. Chem. Res.*, **2019**, 52, 336-344; c) M. Pott, T. Hayashi, T. Mori, P. R. E. Mittl, A. P. Green, D. Hilvert, *J. Am. Chem. Soc.*, **2018**, 140, 1535-1543; d) S. Notonier, Ł. Gricman, J. Pleiss, B. Hauer,

- 1 *ChemBioChem*, **2016**, 17, 1550-1557; e) M. H. Barley, N. J. Turner, R. Goodacre, *ChemBioChem*,
2 **2017**, 18, 1087-1097.
- 3 [161] M. Jeschek, R. Reuter, T. Heinisch, C. Trindler, J. Klehr, S. Panke, T. R. Ward, *Nature*, **2016**, 537,
4 661-665.
- 5 [162] a) C. Donnelly, C. D. Murphy, *Biotechnol. Lett.*, **2008**, 31, 245; b) T. Kurihara, T. Yamauchi, S.
6 Ichiyama, H. Takahata, N. Esaki, *J. Mol. Catal. B Enzym.*, **2003**, 23, 347-355.
- 7 [163] K. Morihiro, Y. Kasahara, S. Obika, *Mol. Biosyst.*, **2017**, 13, 235-245.
- 8 [164] L. Ma, J. Liu, *iScience*, **2020**, 23, 100815.
- 9 [165] Y. Wang, A. Vorperian, M. Shehabat, J. C. Chaput, *ChemBioChem*, **2020**, 21, 1001-1006.
- 10 [166] N. Budisa, V. Kubyskhin, D. Schulze-Makuch, *Life*, **2014**, 4, 374-385.
- 11 [167] a) G. C. Pimentel, K. C. Attwood, H. Gaffron, H. K. Hartline, T. H. Jukes, E. C. Pollard, C. Sagan,
12 Exotic biochemistry in exobiology, in: C.S. Pittendrigh, W. Vishniac, J.P.T. Pearman (Eds.) *Biology*
13 *and the exploration of Mars*, National Academy of Sciences Press, Washington, D. C., USA, 1966;
14 b) M. S. Kondratyev, A. V. Kabanov, A. A. Samchenko, V. M. Komarov, N. N. Khechinashvili, *J.*
15 *Biomol. Struct. Dyn.*, **2013**, 31, 10-11.
- 16 [168] a) D. T. Jacob, *Silicon*, **2016**, 8, 175-176; b) D. J. Norris, *Nature*, **2007**, 446, 146-147; c) S. R.
17 Hameroff, *Ultimate computing: biomolecular consciousness and nanotechnology*, 1st ed., Elsevier,
18 North Holland, 1987.
- 19 [169] J. B. Lambert, *Silicon and life*, in: V.M. Kolb (Ed.) *Handbook of Astrobiology*, Taylor & Francis
20 Group, Boca Raton, 2018, pp. 866.
- 21 [170] S. B. J. Kan, R. D. Lewis, K. Chen, F. H. Arnold, *Science*, **2016**, 354, 1048.
- 22 [171] a) V. D'yakov, *Organosilicon compounds in medicine and cosmetics*, in: J. Weis, N. Auner (Eds.)
23 *Organosilicon Chemistry V: From molecules to materials*, Wiley, Darmstadt, Germany, 2008, pp.
24 348-351; b) A. K. Franz, S. O. Wilson, *J. Med. Chem.*, **2013**, 56, 388-405.
- 25 [172] S. A. Benner, A. Ricardo, M. A. Carrigan, *Curr. Opin. Chem. Biol.*, **2004**, 8, 672-689.
- 26 [173] R. Tacke, *Angew. Chem. Internatl. Ed.*, **1999**, 38, 3015-3018.
- 27 [174] R. L. Brutchey, D. E. Morse, *Chem. Rev.*, **2008**, 108, 4915-4934.
- 28 [175] J. N. Cha, K. Shimizu, Y. Zhou, S. C. Christiansen, B. F. Chmelka, G. D. Stucky, D. E. Morse, *Proc.*
29 *Natl. Acad. Sci. USA*, **1999**, 96, 361-365.
- 30 [176] S. Y. Tabatabaei Dakhili, S. A. Caslin, A. S. Faponle, P. Quayle, S. P. de Visser, L. S. Wong, *Proc.*
31 *Natl. Acad. Sci. USA*, **2017**, 114, E5285-E5291.

- [177] M. B. Frampton, P. M. Zelisko, *Silicon*, **2009**, *1*, 147-163.
- [178] C. Sylatk, A. Stoffregen, F. Wuttke, R. Tacke, *Biotechnol. Lett.*, **1988**, *10*, 731-734.
- [179] V. Abbate, A. R. Bassindale, K. F. Brandstadt, P. G. Taylor, *J. Inorg. Biochem.*, **2011**, *105*, 268-275.
- [180] H. Hengelsberg, R. Tacke, K. Fritsche, C. Sylatk, F. Wagner, *J. Organometallic Chem.*, **1991**, *415*, 39-45.
- [181] K. Sonomoto, H. Oiki, Y. Kato, *Enzyme Microb. Technol.*, **1992**, *14*, 640-643.
- [182] S. Bähr, S. Brinkmann-Chen, M. Garcia-Borràs, J. M. Roberts, D. E. Katsoulis, K. N. Houk, F. H. Arnold, *Angew. Chem. Internatl. Ed.*, **2020**, In press, DOI: 10.1002/anie.202002861.
- [183] E. Rémond, C. Martin, J. Martínez, F. Cavelier, *Chem. Rev.*, **2016**, *116*, 11654-11684.
- [184] B. Weidmann, *Chimia*, **1992**, *46*, 312-313.
- [185] S. M. Sieburth, Bioactive amino acids, peptides and peptidomimetics containing silicon, in: P.M. Zelisko (Ed.) *Bio-inspired silicon-based materials*, Springer, Dordrecht, Netherlands, 2014, pp. 103-123.
- [186] F. Cavelier, D. Marchand, J. Martinez, S. Sagan, *J. Peptide Res.*, **2004**, *63*, 290-296.
- [187] J. L. H. Madsen, C. U. Hjørringgaard, B. S. Vad, D. Otzen, T. Skrydstrup, *Chemistry*, **2016**, *22*, 8358-8367.
- [188] S. Pujals, J. Fernández-Carneado, M. J. Kogan, J. Martinez, F. Cavelier, E. Giralt, *J. Am. Chem. Soc.*, **2006**, *128*, 8479-8483.
- [189] F. Wolfe-Simon, J. S. Blum, T. R. Kulp, G. W. Gordon, S. E. Hoefft, J. Pett-Ridge, J. F. Stolz, S. M. Webb, P. K. Weber, P. C. W. Davies, A. D. Anbar, R. S. Oremland, *Science*, **2011**, *332*, 1163.
- [190] M. L. Reaves, S. Sinha, J. D. Rabinowitz, L. Kruglyak, R. J. Redfield, *Science*, **2012**, *337*, 470.
- [191] T. J. Erb, P. Kiefer, B. Hattendorf, D. Günther, J. A. Vorholt, *Science*, **2012**, *337*, 467.
- [192] G. N. Basturea, T. K. Harris, M. P. Deutscher, *J. Biol. Chem.*, **2012**, *287*, 28816-28819.
- [193] D. S. Tawfik, R. E. Viola, *Biochemistry*, **2011**, *50*, 1128-1134.
- [194] M. Elias, A. Wellner, K. Goldin-Azulay, E. Chabriere, J. A. Vorholt, T. J. Erb, D. S. Tawfik, *Nature*, **2012**, *491*, 134-137.
- [195] A. D. Páez-Espino, P. I. Nikel, M. Chavarría, V. de Lorenzo, *Environ. Microbiol.*, **2020**, In press, DOI: 10.1111/1462-2920.14991.
- [196] R. S. Oremland, J. F. Stolz, *Science*, **2003**, *300*, 939.
- [197] E. O. Uthus, *Environ. Geochem. Health*, **1992**, *14*, 55-58.

- [198] E. O. Uthus, *J. Trace Elem. Exp. Med.*, **2003**, *16*, 345-355.
- [199] R. Bentley, Bartolomeo Gosio, 1863–1944: An appreciation, *Adv. Appl. Microbiol.*, Academic Press 2001, pp. 229-250.
- [200] F. Challenger, C. Higginbottom, L. Ellis, *J. Chem. Soc.*, **1933**, *1*, 95-101.
- [201] a) R. Zakharyan, Y. Wu, G. M. Bogdan, H. V. Aposhian, *Chem. Res. Toxicol.*, **1995**, *8*, 1029-1038; b) R. A. Zakharyan, F. Ayala-Fierro, W. R. Cullen, D. M. Carter, H. V. Aposhian, *Toxicol. Appl. Pharmacol.*, **1999**, *158*, 9-15.
- [202] D. J. Thomas, M. Styblo, S. Lin, *Toxicol. Appl. Pharmacol.*, **2001**, *176*, 127-144.
- [203] a) T. R. Radabaugh, H. V. Aposhian, *Chem. Res. Toxicol.*, **2000**, *13*, 26-30; b) T. R. Radabaugh, A. Sampayo-Reyes, R. A. Zakharyan, H. V. Aposhian, *Chem. Res. Toxicol.*, **2002**, *15*, 692-698; c) E. Sánchez-Bermejo, G. Castrillo, B. del Llano, C. Navarro, S. Zarco-Fernández, D. J. Martínez-Herrera, Y. Leo del Puerto, R. Muñoz, C. Cámara, J. Paz-Ares, C. Alonso-Blanco, A. Leyva, *Nat. Commun.*, **2014**, *5*, 4617; d) A. D. Páez-Espino, G. Durante-Rodríguez, V. de Lorenzo, *Environ. Microbiol.*, **2015**, *17*, 229-238.
- [204] a) G. L. Anderson, J. Williams, R. Hille, *J. Biol. Chem.*, **1992**, *267*, 23674-23682; b) J. M. Santini, R. N. van den Hoven, *J. Bacteriol.*, **2004**, *186*, 1614.
- [205] C. Niegel, F.-M. Matysik, *Anal. Chim. Acta*, **2010**, *657*, 83-99.
- [206] K. O. Amayo, A. Raab, E. M. Krupp, H. Gunnlaugsdottir, J. Feldmann, *Anal. Chem.*, **2013**, *85*, 9321-9327.
- [207] A. Popowich, Q. Zhang, X. C. Le, *Natl. Sci. Rev.*, **2016**, *3*, 451-458.
- [208] H. V. Aposhian, R. A. Zakharyan, M. D. Avram, A. Sampayo-Reyes, M. L. Wollenberg, *Toxicol. Appl. Pharmacol.*, **2004**, *198*, 327-335.
- [209] S. Gibaud, G. Jaouen, Arsenic-based drugs: From Fowler's solution to modern anticancer chemotherapy, in: G. Jaouen, N. Metzler-Nolte (Eds.) *Medicinal organometallic chemistry*, Springer, Berlin, Germany, 2010, pp. 1-20.
- [210] R. M. Couture, A. Sekowska, G. Fang, A. Danchin, *Environ. Microbiol.*, **2012**, *14*, 1612-1623.
- [211] S. Touchet, F. Carreaux, B. Carboni, A. Bouillon, J. L. Boucher, *Chem. Soc. Rev.*, **2011**, *40*, 3895-3914.
- [212] S. B. J. Kan, X. Huang, Y. Gumulya, K. Chen, F. H. Arnold, *Nature*, **2017**, *552*, 132-136.
- [213] G. W. Kabalka, M. L. Yao, S. R. Marepally, S. Chandra, *Appl. Radiat. Isot.*, **2009**, *67*, S374-379.

- [214] a) E. Brustad, M. L. Bushey, J. W. Lee, D. Groff, W. Liu, P. G. Schultz, *Angew. Chem. Internatl. Ed.*, **2008**, 47, 8220-8223; b) A. Schiefner, L. Nästle, M. Landgraf, A. J. Reichert, A. Skerra, *Biochemistry*, **2018**, 57, 2597-2600.
- [215] S. Edwardraja, A. Eichinger, I. Theobald, C. A. Sommer, A. J. Reichert, A. Skerra, *ACS Synth. Biol.*, **2017**, 6, 2241-2247.
- [216] H. J. Reich, R. J. Hondal, *ACS Chem. Biol.*, **2016**, 11, 821-841.
- [217] A. Presentato, R. J. Turner, C. C. Vasquez, V. Yurkov, D. Zannoni, *Environ. Chem.*, **2019**, 16, 266-288.
- [218] R. J. Turner, R. Borghese, D. Zannoni, *Biotechnol. Adv.*, **2012**, 30, 954-963.
- [219] L. Moroder, *J. Peptide Sci.*, **2005**, 11, 187-214.
- [220] a) J. Bassan, L. M. Willis, R. N. Vellanki, A. Nguyen, L. J. Edgar, B. G. Wouters, M. Nitz, *Proc. Natl. Acad. Sci. USA*, **2019**, 116, 8155; b) N. Vurgun, M. Nitz, *ChemBioChem*, **2020**, In press: DOI: 10.1002/cbic.201900635.
- [221] X. Liu, L. A. Silks, C. Liu, M. Ollivault-Shiflett, X. Huang, J. Li, G. Luo, Y.-M. Hou, J. Liu, J. Shen, *Angew. Chem. Internatl. Ed.*, **2009**, 48, 2020-2023.
- [222] N. Budisa, W. Karnbrock, S. Steinbacher, A. Humm, L. Prade, T. Neufeind, L. Moroder, R. Huber, *J. Mol. Biol.*, **1997**, 270, 616-623.
- [223] D. M. Upp, J. C. Lewis, *Curr. Opin. Chem. Biol.*, **2017**, 37, 48-55.
- [224] Y. Yu, C. Hu, L. Xia, J. Wang, *ACS Catal.*, **2018**, 8, 1851-1863.
- [225] a) H. M. Key, P. Dydio, D. S. Clark, J. F. Hartwig, *Nature*, **2016**, 534, 534-537; b) J. Huang, Z. Liu, B. Bloomer, D. S. Clark, A. Mukhopadhyay, J. D. Keasling, J. F. Hartwig, *ChemRxiv*, **2020**, DOI: 10.26434/chemrxiv.11955174.v11955171.
- [226] C. Mayer, D. G. Gillingham, T. R. Ward, D. Hilvert, *Chem. Commun.*, **2011**, 47, 12068-12070.
- [227] M. Filice, O. Romero, A. Aires, J. M. Guisan, A. Rumbero, J. M. Palomo, *Adv. Synth. Catal.*, **2015**, 357, 2687-2696.
- [228] N. Fujieda, T. Nakano, Y. Taniguchi, H. Ichihashi, H. Sugimoto, Y. Morimoto, Y. Nishikawa, G. Kurisu, S. Itoh, *J. Am. Chem. Soc.*, **2017**, 139, 5149-5155.
- [229] Q. Jing, R. J. Kazlauskas, *ChemCatChem*, **2010**, 2, 953-957.
- [230] A. Danchin, P. I. Nikel, *J. Mol. Evol.*, **2019**, 87, 271-288.
- [231] K. K. Yang, Z. Wu, F. H. Arnold, *Nat. Methods*, **2019**, 16, 687-694.
- [232] A. Danchin, *ChemBioChem*, **2020**, In press, DOI: 10.1002/cbic.202000060.

- [233] C. Kampmeyer, J. V. Johansen, C. Holmberg, M. Karlson, S. K. Gersing, H. N. Bordallo, B. B. Kragelund, M. H. Lerche, I. Jourdain, J. R. Winther, R. Hartmann-Petersen, *ACS Synth. Biol.*, **2020**, In press, DOI: 10.1021/acssynbio.1029b00376.
- [234] a) P. I. Nickel, M. Chavarría, A. Danchin, V. de Lorenzo, *Curr. Opin. Chem. Biol.*, **2016**, 34, 20-29; b) P. I. Nickel, M. Chavarría, T. Fuhrer, U. Sauer, V. de Lorenzo, *J. Biol. Chem.*, **2015**, 290, 25920-25932; c) P. I. Nickel, E. Martínez-García, V. de Lorenzo, *Nat. Rev. Microbiol.*, **2014**, 12, 368-379; d) D. C. Volke, P. Calero, P. I. Nickel, *Trends Microbiol.*, **2020**, In press, DOI: 10.1016/j.tim.2020.1002.1015; e) N. T. Wirth, E. Kozaeva, P. I. Nickel, *Microb. Biotechnol.*, **2020**, 13, 233-249; f) I. Poblete-Castro, C. Wittmann, P. I. Nickel, *Microb. Biotechnol.*, **2020**, 13, 32-53.
- [235] a) P. I. Nickel, V. de Lorenzo, *Metab. Eng.*, **2018**, 50, 142-155; b) P. Calero, P. I. Nickel, *Microb. Biotechnol.*, **2019**, 12, 98-124.
- [236] a) I. Poblete-Castro, J. M. Borrero de Acuña, P. I. Nickel, M. Kohlstedt, C. Wittmann, Host organism: *Pseudomonas putida*, in: C. Wittmann, J.C. Liao (Eds.) *Industrial Biotechnology: Microorganisms*, Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, 2017; b) P. I. Nickel, D. Pérez-Pantoja, V. de Lorenzo, *Environ. Microbiol.*, **2016**, 18, 3565-3582.
- [237] P. I. Nickel, V. de Lorenzo, *Metab. Eng.*, **2013**, 15, 98-112.
- [238] a) E. Belda, R. G. A. van Heck, M. J. López-Sánchez, S. Cruveiller, V. Barbe, C. Fraser, H. P. Klenk, J. Petersen, A. Morgat, P. I. Nickel, D. Vallenet, Z. Rouy, A. Sekowska, V. A. P. Martins dos Santos, V. de Lorenzo, A. Danchin, C. Médigue, *Environ. Microbiol.*, **2016**, 18, 3403-3424; b) J. Nogales, J. Mueller, S. Gudmundsson, F. J. Canalejo, E. Duque, J. Monk, A. M. Feist, J. L. Ramos, W. Niu, B. Ø. Palsson, *Environ. Microbiol.*, **2020**, 22, 255-269; c) A. Sánchez-Pascuala, V. de Lorenzo, P. I. Nickel, *ACS Synth. Biol.*, **2017**, 6, 793-805; d) A. Sánchez-Pascuala, L. Fernández-Cabezón, V. de Lorenzo, P. I. Nickel, *Metab. Eng.*, **2019**, 54, 200-211; e) S. Sudarsan, S. Dethlefsen, L. M. Blank, M. Siemann-Herzberg, A. Schmid, *Appl. Environ. Microbiol.*, **2014**, 80, 5292-5303; f) M. Chavarría, A. Goñi-Moreno, V. de Lorenzo, P. I. Nickel, *mSystems*, **2016**, 1, e00154-00116; g) M. Chavarría, P. I. Nickel, D. Pérez-Pantoja, V. de Lorenzo, *Environ. Microbiol.*, **2013**, 15, 1772-1785.
- [239] a) P. I. Nickel, D. Pérez-Pantoja, V. de Lorenzo, *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, **2013**, 368, 20120377; b) D. Pérez-Pantoja, P. I. Nickel, M. Chavarría, V. de Lorenzo, *PLoS Genet.*, **2013**, 9, e1003764; c) Ö. Akkaya, P. I. Nickel, D. Pérez-Pantoja, V. de Lorenzo, *Environ. Microbiol.*, **2019**,

- 1 21, 314-326; d) Ö. Akkaya, D. Pérez-Pantoja, B. Calles, P. I. Nickel, V. de Lorenzo, *mBio*, **2018**, 9,
2 e01512-01518.
- 3 [240] H. Deng, L. Ma, N. Bandaranayaka, Z. Qin, G. Mann, K. Kyeremeh, Y. Yu, T. Shepherd, J. H.
4 Naismith, D. O'Hagan, *ChemBioChem*, **2014**, 15, 364-368.
- 5 [241] Y. Wang, Z. Deng, X. Qu, *F1000Research*, **2014**, 3, 61.

TABLES

Table 1. Kinetic properties of the fluorinase enzymes characterized thus far.^a

Organism	K_m		k_{cat}	k_{cat}/K_m	K_i SAH	References
	SAM (μ M)	F ⁻ (mM)	(min ⁻¹) ^b	(mM ⁻¹ min ⁻¹) ^b	(mM)	
<i>Streptomyces cattleya</i>	29.2 $\pm$ 2.41	8.56 $\pm$ 0.52	0.083	2.84	0.029	[59, 240]
<i>Streptomyces</i> sp. MA37	82.4 $\pm$ 18.6	N.R.	0.262	3.18	N.R.	[240]
<i>Nocardia brasiliensis</i>	27.8 $\pm$ 4.23	4.15	0.122	4.40	N.R.	[240, 241]
<i>Actinoplanes</i> sp. N902-109	45.8 $\pm$ 7.91	N.R.	0.204	4.44	N.R.	[240]

^a A recent report includes yet another fluorinase enzyme from *Actinopolyspora mzabensis*,^[74] but kinetic data is not available for this isolate. N.R., not reported.

^b These k_{cat} and k_{cat}/K_m values correspond to assays considering SAM as the kinetic substrate in the presence of a F⁻ excess.

FIGURE CAPTIONS

Figure 1. Abundance in the Earth's continental crust and physicochemical properties of selected biotic chemical elements and their xeno-analogues. Elements are represented at scale based on their van der Waals radii. The (–) symbol indicates pairing of C with the same canonical element (i.e. C–H, C–C and C–P).

Canonical element	Potential analogue	Group	van der Waals radii (Å)	Abundance in Earth's crust (ppm)	Electro-negativity X	Target bond	Bond length (Å)	Bond dissociation energy (kJ/mol)
–	1	1.10	-	2.20	C–H	1.09	339
	17	1.47	585	3.98	C–F	1.35	514
	17	1.75	145	3.16	C–Cl	1.77	395
	17	1.85	2.4	2.96	C–Br	1.94	318
	17	1.98	0.45	2.66	C–I	2.14	253
–	14	1.70	200	2.55	C–C	1.54	605
	14	2.10	282,000	1.90	C–Si	1.85	447
						Si–Si	2.33	310
–	15	1.80	1,050	2.19	P–O	1.63	589
						C–P	1.84	508
	15	1.85	1.8	2.01	As–O	1.78	484
						C–As	1.98	Unknown

Figure 2. Examples of natural and synthetic halogen-containing organocompounds. Selected examples of halocompounds containing F, Cl, Br and I are discussed in the text. Note that some of these chemical species can be used as building blocks for polymers (e.g. halo-amino acids), whereas others could be exploited as biochemical precursors for *neo*-metabolism.

Figure 3. Canonical biofluorination pathway in *Streptomyces cattleya*. The biofluorination route described by David O'Hagan and co-workers^[68] starts from inorganic fluoride (F^-) and SAM, and finally yields either fluoroacetate or 4-fluorothreonine. All relevant enzymes are indicated along the reaction they are predicted to catalyze, including the fluorinase, which mediates the formation of a C–F bond. Abbreviations used in the scheme are as follows: SAM, S-adenosyl-L-methionine; 5'-FDA, 5'-fluoro-5'-deoxyadenosine; 5-FDRP, 5'-fluoro-5'-deoxy-D-ribose-1-phosphate; 5-FDRibulP: 5'-fluoro-5'-deoxy-D-ribulose-1-phosphate.

Figure 4. Residue- and site-specific incorporation of unnatural amino acids into nascent polypeptides. The residue-specific incorporation relies on the similarity between an unnatural amino acid and its canonical counterpart, which is incorporated in the newly born protein by the corresponding natural tRNA/aaRS (aminoacyl-tRNA synthetase) pair. This approach requires a higher abundance of the amino acid analogue in comparison to the natural version, which is generally attained by using auxotrophic strains. In this case, the unnatural amino acid is incorporated replacing the canonical residue in every position of the polypeptide. The site-specific incorporation strategy is based on the engineering of a suitable host equipped with an orthogonal tRNA/aaRS pair specific for the selected unnatural amino acid. The unnatural residue is inserted in the newly born protein in response to a recoded blank codon (usually the *amber* codon). Note that, in this case, there is no interference by the natural pool of canonical amino acids, and the xeno-analogue is inserted in selected positions of the polypeptide.

Figure 5. General strategy to expand the catalytic scope of an enzyme by combining directed-evolution and the incorporation of xeno-elements towards *neo*-metabolism. A conventional mutant library is firstly generated from the gene encoding the selected enzyme. Some of the enzyme variants stemming from the library might display altered kinetic properties, or amber codons can be randomly introduced in the coding sequence (which allows for the incorporation of unnatural amino acids). The gene library can be transformed in both a non-modified and an engineered microbial host. The latter possesses the machinery to specifically incorporate a certain fluoro-amino acid (FAA) in response to the amber codon. In the presence of this unnatural (FAA) amino acid, clones of the library carrying the *UAG* triplet will produce fluorinated versions of the enzyme. On the contrary, the non-engineered host will lead to variants limited to the 20 canonical amino acids. The scheme further illustrates how adding a novel element to the natural building blocks (based on C, H, N, O and P) expands the reaction space that is enzymatically attainable towards the production of new-to-Nature molecules.

Figure 6. Si-containing organocompounds and related chemicals relevant for xenobiology. Due to its similarity to C, several organocompounds can likewise incorporate Si. Selected examples indicated in the figure comprise small building blocks [e.g. $\text{Si}(\text{OH})_4$] and also peptides containing silaproline (Sip) and trimethylsilyl alanine (TMSAla).

Figure 7. Key examples of As-containing organocompounds. Both metabolites (e.g. sugar derivatives) and polymers (e.g. lipids) are known to incorporate As in their structures. Other naturally occurring arsenometabolites are listed.

Table of contents

Bringing xeno-elements into the chemistry of microbial cells by combining several approaches. Engineered hosts are programmed through *neo*-metabolism to accept non-biological atoms, incorporated into xeno-metabolites. Some of these molecules are added-value chemicals themselves, whereas others (e.g. modified amino acids) can be integrated into proteins to explore emergence of novel functions.