Impact of ohmicsonication treatment on pectinmethylesterase in not-from-concentrate orange juice

Abdelmaksoud, Tarek Gamal; Mohsen, Sobhy Mohamed; Duedahl-Olesen, Lene; Elnikeety, Mohamed Mohamed; Feyissa, Aberham Hailu

Published in:
Journal of Food Science and Technology

Link to article, DOI:
[10.1007/s13197-019-03834-2](https://doi.org/10.1007/s13197-019-03834-2)

Publication date:
2019

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Abdelmaksoud, T. G., Mohsen, S. M., Duedahl-Olesen, L., Elnikeety, M. M., & Feyissa, A. H. (2019). Impact of ohmicsonication treatment on pectinmethylesterase in not-from-concentrate orange juice. *Journal of Food Science and Technology*, 56(8), 3951-3956. <https://doi.org/10.1007/s13197-019-03834-2>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Impact of ohmicsonication treatment on pectinmethylesterase in not-from-concentrate orange juice

Tarek Gamal Abdelmaksoud^{1,2} · Sobhy Mohamed Mohsen¹ · Lene Duedahl-Olesen² · Mohamed Mohamed Elnikeety¹ · Aberham Hailu Feyissa²

Revised: 9 May 2019 / Accepted: 14 May 2019
© Association of Food Scientists & Technologists (India) 2019

Abstract The present study investigates the application of ohmicsonication (OS) as a new hurdle technology for pasteurization of Not-from-concentrate orange juice (NFCOJ). OS process parameters to inactivate pectinmethylesterase (PME) activity in NFCOJ were optimized using response surface methodology. The influence of Sonication (S), Thermosonication (TS), Ohmic heating (OH) and OS on inactivation of PME were compared to conventional heat (CH) treatment. Their effects on physical, chemical and microbiological contents were included. In comparison to fresh orange juice, the inactivation of PME was 96%, 95%, 89%, 90% and 29% for OS, OH, TS, CH and S treatments, respectively. Highest cloud value was obtained for OS (1.240 A) treatment. OS treatment gave a lower vitamin C loss compared to TS, OH and CH treatments. A significant increase in the total phenolic content were obtained in the following order OS > TS > OH > CH. OS treated juice also contained the lowest value of hydroxymethyl furfural (0.90 mg/L) compared to OH (0.95 mg/L), TS (1.37 mg/L) and CH (2.72 mg/L) treated samples. Overall, the results indicated that OS can be integrated as a substitute to pasteurization of NFCOJ.

Keywords Ohmic heating · Ohmic-ultrasonic heating · Pectinmethylesterase · Phenolic content

Introduction

Orange juice (OJ) is a good source of bioactive compounds (i.e. vitamin C, phenolics and carotenoids) (Galaverna and Dall'Asta 2014). During storage, OJ can be subject to deteriorative reactions including enzymatic activities, microbial spoilage, vitamin C degradation, cloud loss and changes in flavor and color, all reactions that lead to loss of product quality. Due to the perishable nature of juices, several technologies have been used to prolong the shelf life (Polydera et al. 2005). Today, recent technologies in food processing either thermal or non-thermal are designed to meet the consumer demands (Williams 1994). Not-From-Concentrate (NFC) juice is the fruit juice extract without concentration or dilution. Insoluble pulp, skin and seeds are removed before heat treatment of the juice for controlling the microbial load and enzymatic activities (Abdelmaksoud et al. 2018a).

The activity of pectinmethylesterase (PME) results in unwanted layer separation during processing. Conventional heat (CH) treatment reduce PME activity as well as microbial load, however, this treatment can cause loss of nutritional value and produce undesirable flavor in the juice product, especially at temperatures higher than 80 °C (Giner et al. 2013). Hence, the juice industry searches for mild thermal technologies, which are able to inactivate both enzymes and microorganisms with retention of nutritional value (Chemat et al. 2011).

Hurdle technology is one of the promising technologies in food processing to improve the quality, safety, and stability of food products. Sonication (S) is one of the

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s13197-019-03834-2>) contains supplementary material, which is available to authorized users.

✉ Tarek Gamal Abdelmaksoud
tareekgamal_88@agr.cu.edu.eg

¹ Food Science Department, Faculty of Agriculture, Cairo University, Giza, Egypt

² National Food Institute, Technical University of Denmark, Søtofts Plads 227, 2800 Lyngby, Denmark

emerging green methods for food processing and preservation. S is defined as a potential technology acting on reduction of the microbial load in food by 5-log when S generates cavitation bubbles due to pressure changes (USFDA 2001). S as a substitute to the CH has the ability to decrease processing time, cost and energy as well as enhanced quality, shelf life and ensured safety of fruit juices (Chemat et al. 2011). However, S alone has limited applications in juices due to its insufficient enzymes and microorganisms inactivation. Therefore, a combination of S with other technologies with improved efficiencies has been investigated and considered due to the possibility for the application at the industrial scale (Leistner 2000). Previous literature however lack the combination of ohmic heating (OH) with S as an alternative to the CH for production and evaluation of NFC orange juice (NFCOJ). Therefore, the main objectives of this study was to optimize the OS parameters (OH temperature and S time) for inactivation of PME and evaluate the effects on the final NFCOJ. Effects on quality characteristics of the final OJ product at optimum OS conditions were compared to production by other technologies such as S, Thermosonication (TS), OH and CH.

Materials and methods

Chemicals

All chemicals in this study were purchased from (Sigma-Aldrich Chemical Co., Denmark).

Raw material

Orange fruits (*Citrus sinensis*, cv. *Navel*) purchased from a local supermarket in Copenhagen, Denmark were rinsed and cut into halves. The juice was extracted (Extractor, Krups Citrus Juicer, Spain) and filtrated using a double-layered muslin cloth. The extracted juice was divided into six groups and subjected to S, OS, TS, OH, CH or kept fresh and all groups were rapidly cooled to 4 °C and stored at − 18° C for further analysis.

Processing methods

Conventional heating

Orange juice (OJ) (150 mL) was heated at 95 °C for 60 s using a shaker water bath (Julabo, SW22, made in Germany) according to Abdelmaksoud et al. (2018b).

Sonication and Thermosonication (S and TS)

Samples of 150 mL OJ was treated with the S processor of 550 W at 20 kHz with a 0.5-inch probe (Sonifier SFX550 Model, Mexico). Treatments at 25 °C using 100% power (550 W) for 8 min at pulse durations of 5 s was used. TS was done at 60 °C at conditions similar to S.

Ohmic heating (OH)

The treatment of OJ (150 mL) by OH (at 42 V/cm, 69 °C and held for 60 s) was conducted according to Demirdöven and Baysal (2014).

Ohmicsonication (OS)

OS was done by a combination of S and OH treatments at the obtained optimum parameters, firstly treated by S for 8 min at 25 °C and then directly followed by OH at 40 V/cm, to 68 °C for a holding time of 60 s.

Experimental design and statistical methods

Response surface methodology (RSM) was used for optimization of OS parameters (OH temperature and S time). Two factors (OH temperature and S time) with three levels (− 1, 0, + 1), the factorial design (3^2) was used. The OH temperature and S time range were 60, 65, 70 °C and 2, 5, 8 min, respectively. To describe the effect of parameters, the second-order polynomial model was used (Eq. 1).

$$Y = a_0 + a_1x_1 + a_2x_2 + a_{12}x_1x_2 + a_{11}x_1^2 + a_{22}x_2^2 \quad (1)$$

where Y is the % of PME inactivation, x_1 is OH temperature and x_2 is S time, $a_0, a_1, a_2, a_{11}, a_{22}$ and a_{12} are regression coefficients for intercept, the linear, the quadratic and interaction term, respectively.

The analysis of variance (ANOVA) for the response was used to find the significant terms in the models (Table S1, supplementary data). Design Expert Version 10.0.6 software was used for the analysis. To optimize the OS parameters, the desirability function method was used. The objective function was to maximize the PME inactivation using desirability function as in Abdelmaksoud et al. (2018b).

To evaluate the variances among different treatments at significance levels ($p \leq 0.05$), data in Tables 2 was statistically analyzed with one-way ANOVA (Duncan test), using SPSS 13 software (SPSS Inc., Chicago IL, USA).

Physical analysis

Electric conductivity (EC), cloud value and color values (L, a and b) were determined using methods described by

Abdelmaksoud et al. (2018a). Size distribution particles were detected with the Mastersizer (Model 2000, Malvern, UK).

Chemical analysis

Vitamin C content was determined using a 2,6-dichlorophenol indophenol (DCPIP) visual titration method (Ranganna 1986).

Total phenolic content was calorimetrically measured applying Folin-Ciocalteu reagent as described by Abdullakassim et al. (2007) with modifications for microplate reader in Abdelmaksoud et al. (2018b). Results were given as mg of Gallic acid/100 mL OJ.

Total carotenoids was measured according to Lee and Castle, (2001) with some modifications based on 5 mL of OJ (Abdelmaksoud et al. 2018a). Absorbance at 450 nm of the final supernatant was measured and total carotenoid contents were calculated according to Ritter & Purcell (1981) using an extinction coefficient of β -carotene ($\mu\text{g/g}$), $E^{1\%} = 2505$.

Total flavonoids was measured based on an assay developed by Kim et al. (2003) and the results were expressed as mg catechin equivalents/100 g OJ.

Hydroxymethyl furfural (HMF) was determined according to a method of Vorlova et al. (2006) using a Vortex Genie II (Scientific Industries, Bohemia, USA) for mixing 1 mL methanol and 0.5 mL juice. The centrifuged and filtered (0.45 μm) extract was injected (20 μL) onto HPLC (Alliance, Waters Company) equipped with a Zorbax Eclipse XDB-C8, 4.6 $\times$ 150 mm, 5 μm column (Waters, Milford, USA) at 30 $^{\circ}\text{C}$ using a flow rate of 1 mL/min and an isocratic mobile phase (10% methanol in water). UV detection at 285 (2996 diode array detector) using external standard method for quantification of HMF (retention time 3.17 min) with a linearity concentration range of 0.01–200 mg/L based on Empower software (Waters).

Pectinmethylesterase activity was determined according to the method described by Rouse and Atkins (1955) and Ting and Rouseff (1986).

Microbial load

Total plate count and mold and yeast were determined as in Andrews (1992).

Results and discussion

Optimization of ohmicsonication (OS) conditions

Pre-experiments determined optimal OH temperatures in the range 60–70 $^{\circ}\text{C}$ and S time in the range of 2–8 min, where S time $\geq$ 8 min would result in adverse color and vitamin C changes for OJ (data not included). Also, increasing the temperature more than 80 $^{\circ}\text{C}$ caused deterioration of the color and increased juice bubbling leading to juice loss. The voltage gradient of each OH treatments was selected to be 42 V/cm according to optimization of OH conditions by Demirdöven and Baysal (2014). RSM set up resulted in PME inhibitions (%) presented in Table 1.

Optimization of the conditions of OS by applying second order polynomial equation and multiple regression analysis were used to obtain the regression coefficients for independent variables (Eq. 1).

The effect of OH temperature and S time on PME activity at 95% confidence interval (Table S1, supplementary data). The experimental data was fitted and significant with the used model. Insignificant difference between adj- R^2 value (0.968) and R^2 for PME—this means high degree of correlation between the predicted and experimental values with insignificant lack-of-fit. The model was suitable for describing the % inactivation of PME within tested experimental ranges.

The positive linear effect of OH temperature (χ_1), S time (χ_2) were found to be significant for the response variable (Y : % inactivation of PME). Also, the interaction ($\chi_1 \chi_2$) and the quadratic effect of OH temperature (χ_1^2) on PME were found to be significant. However, the quadratic of S time (χ_2^2) had an insignificant effect on the PME. The fitted second order polynomial equation are presented as (Eq. 2):

$$Y = +93.35 + 3.94\chi_1 + 1.76\chi_2 - 1.00\chi_1\chi_2 - 2.37\chi_1^2 \quad (2)$$

where χ_1 : OH temperature ($^{\circ}\text{C}$) and χ_2 : S time (min). According to the obtained second order polynomial models, the optimum conditions for OS of NFCOJ was obtained at the maximum PME inactivation by applying desirability function. The obtained optimum parameters were OH at 68 $^{\circ}\text{C}$ for 60 s combined with S at 550 W for 8 min, with a PME inactivation of 96% in NFCOJ (Fig. 1).

Physical, chemical and microbiological contents of fresh and treated NFC orange juice

Table 2 shows the effects of OS, S, CH, TS and OH on physical, chemical and microbial load contents of NFCOJ. Compared to fresh orange juice (FOJ), an increase in the EC was observed for all treatments of OJ in the following

Table 1 Experimental design of ohmicsonication (OS) and PME activity in NFC orange juice

Run order	OH temperature (°C), χ_1	Sonication time (min), χ_2	% inhibition of PME
1	65 (0)	5 (0)	93.53
2	65 (0)	5 (0)	93.40
3	70 (+ 1)	5 (0)	94.93
4	70 (+ 1)	2 (− 1)	93.88
5	60 (− 1)	5 (0)	87.98
6	65 (0)	2 (− 1)	92.04
7	70 (+ 1)	8 (+ 1)	95.94
8	60 (− 1)	2 (− 1)	83.53
9	60 (− 1)	8 (+ 1)	89.61
10	65 (0)	8 (+ 1)	94.44

In the 2nd and 3rd column: the coded values of the test parameters are in parenthesis and the real (un-coded) values are outside the parenthesis, *PME* pectinmethylesterase

Fig. 1 Effect of Ohmicsonication (OS) parameters (OH temperature and Sonication time) on the PME activity of orange juice (U/mL/min)—response surface and contour plots

order: OS > TS > S > OH > CH. Increased EC of juice, especially for S (S, TS and OS) might be attributed to the agitated enhanced release of minerals and bioactive components as well as increased ionic mobility (Zou and Jiang 2016).

A lower ADS, indicating a more homogenous juice, was observed in the sonicated samples (OS, S, TS) compared to non-sonicated samples (FOJ, CH and OH) (Table 2). This is due to the effect of cavitation collapse by S and the resulting particle size reduction (Franco et al. 2004).

For color values (L^* , a^* and b^*) the L^* -value was increased by S, TS and OS, while insignificantly changed by CH and

OH treatments. The a^* and b^* were significantly increased for all treatments (OS > TS > S > OH > CH) compared to FOJ, attributed to chemical changes by thermal treatments (Bhale 2004). Increased color values by S could be due to less enzymatic color changes and an increased phenolic and carotenoid condensation reducing their oxidation as explained by Tiwari et al. (2008).

Vitamin C were significantly decreased for all treatments when compared to FOJ (52.77 mg/100 mL) and followed the order S > OS > OH > TS > CH. S alone was not efficient in the inactivation of PME and microorganisms (Table 2) and therefore not accepted for

Table 2 Physical, chemical and microbiological contents of fresh and processed NFC orange juice during different treatments of NFC orange juice

Parameters	FOJ	S	CH	TS	OH	OS
EC (s/m)	0.272 ± 0.003c	0.293 ± 0.004ab	0.282 ± 0.004bc	0.297 ± 0.007ab	0.286 ± 0.004abc	0.303 ± 0.009a
Cloud value (A)	0.253 ± 0.07c	0.274 ± 0.003c	0.970 ± 0.005b	0.946 ± 0.008b	0.994 ± 0.003b	1.240 ± 0.04a
ADS (µm)	3.970 ± 0.04b	0.304 ± 0.01d	2.096 ± 0.02c	0.290 ± 0.01f	4.191 ± 0.02a	0.323 ± 0.01e
L*	42.57 ± 0.02d	44.71 ± 0.09c	42.63 ± 0.03d	46.03 ± 0.08b	42.77 ± 0.04d	46.95 ± 0.29a
a*	− 6.66 ± 0.02e	− 5.15 ± 0.04b	− 6.08 ± 0.02d	− 4.96 ± 0.03 cd	− 5.83 ± 0.02 cd	− 4.73 ± 0.01a
b*	9.65 ± 0.02e	10.36 ± 0.14c	9.95 ± 0.06f	11.33 ± 0.02a	10.15 ± 0.05d	10.76 ± 0.03b
ΔE	—	2.72 ± 0.27c	0.65 ± 0.02e	4.21 ± 0.04a	0.99 ± 0.03d	4.91 ± 0.28b
Vitamin C (mg/100 mL)	52.77 ± 0.14a	48.22 ± 0.22b	43.58 ± 0.25f	46.08 ± 0.16e	47.22 ± 0.18d	47.62 ± 0.19c
Total carotenoids (µg / 100 g)	1270 ± 13.5b	1308 ± 3.22a	982 ± 8.16f	1107 ± 7.62e	1126 ± 5.54d	1183 ± 8.73c
Total flavonoids (mg/ 100 mL)	20.01 ± 0.39c	21.80 ± 0.47b	20.32 ± 0.62c	22.27 ± 0.24a	20.58 ± 1.38c	22.83 ± 0.31a
Total phenolic (mg/ 100 mL)	37.47 ± 0.13e	42.06 ± 0.22a	37.61 ± 0.26e	39.64 ± 0.10c	38.22 ± 0.09d	40.73 ± 0.69b
HMF (mg/L)	nd	0.73 ± 0.02c	2.72 ± 0.02a	1.37 ± 0.18b	0.95 ± 0.21c	0.90 ± 0.08c
PME (U/mL/min)	47.57 ± 0.6a	33.55 ± 0.4b	5.38 ± 0.3c	4.62 ± 0.2d	2.21 ± 0.1e	1.93 ± 0.3f
Total plate count (log cfu/ mL)	2.27 ± 0.5	nd	nd	nd	nd	nd
Mold and yeast (log cfu/ mL)	1.92 ± 0.3	nd	nd	nd	nd	nd

Different letters (a, b, c) mean statistical significant difference ($p < 0.05$); the results represent the mean $\pm$ SD
FOJ fresh orange juice, S sonication, CH conventional heating, TS thermosonication, OH ohmic heating, OS ohmicsonication, EC electric conductivity, ADS average droplet sizes (µm), HMF hydroxymethylfurfural, PME pectinmethylesterase

processing of OJ. The vitamin C reduction was attributed to chemical decomposition due to both temperature and time of processing correlating with the findings by Demirdöven and Baysal (2014), who reported a decrease of vitamin C in the OH and CH treatments compared to the FOJ.

A significant increase in the total carotenoids of OJ for S (1308 µg/100 g) correlates with cell wall disruption causing more free carotenoids in the juice stated by Plaza et al. (2011). Also, Abid et al. (2014) observed increased carotenoid levels with S. Oxygen, light, metals and enzyme availability as well as heat application led to loss of carotenoids for OS, TS, OH and CH (Table 2), which is in agreement with previous reports on carotenoid stability (Rawson et al. 2011; Esteve et al. 2009).

Total phenolic content (TPC) of OJ were significantly increased in S, OS, TS and OH treatments ($S > OS > TS > OH$) compared to FOJ, while insignificant increase was found for CH treated sample (Table 2). Heating (during CH and OH treatments) might increase the extractability of TPC due to breakdown of the interaction between proteins and polyphenols (Girgin and El 2015). Previous reports explain TPC release during OH induced by the alternating current (Roy et al. 2009), while the increased release of TPC during S is due to the cavitation

phenomenon resulting in breakdown of the cell wall based on liquid pressure changes during S treatment, thus increasing the availability of phenols in the juice (Abid et al. 2014).

OS treated juice contained the lowest value of HMF (0.90 mg/L) followed by OH (0.95 mg/L) and TS (1.37 mg/L) with highest values in CH treated juice (2.72 mg/L). The presence of HMF in foods (containing carbohydrates in an acidic environment) is a result of high heat treatment ($T > 80\text{ }^{\circ}\text{C}$) and inappropriate and long-term storage. As expected, levels for HMF varies e.g. from not detected to 27 mg/L in fruit juices (Vorlova et al. 2006).

All treatments showed a significant increase in cloud value with the highest value indicating highest cloud stability and PME inhibition obtained for OS (Table 2). Increased parameters were attributed to the combined effects of heat and voltage gradient (which might remove the metallic prosthetic groups present in the PME) (Castro et al. 2004) and cavitation on enzyme activity, causing a decreased layer separation.

After treatment of OJ sample by S, CH, TS, OH and OS no microbial growth neither total plate count nor mold or yeast were detected (Table 2).

307 Conclusion

308 The use of OS result in an improved quality of OJ compared to other treatments. The highest inactivation of PME
309 activity with no microbial load as well as highest retention
310 of vitamin C, carotenoids, phenolics, and flavonoids were
311 obtained with OS compared to the other treatments (OH,
312 TS, S and CH). In addition, OS treatment resulted in
313 increased EC, cloud value and color values. S treatment
314 alone was not sufficient for inactivation of PME at lower
315 temperature. Overall, OS improved the quality of OJ in
316 laboratory scale compared to other treatments and can be a
317 potential technology for pasteurization of juice. An appli-
318 cation in a pilot plant or large scale could be interesting and
319 needs to be considered for further studies.

321 **Acknowledgements** Tarek Abdelmaksoud would like to thank The
322 Danish Agency for Higher Education for a research grant as a guest
323 Ph.D. Student for one year in Food Production Engineering Research
324 Group, Technical University of Denmark. Authors would like to
325 thank Anette Bysted and Inge Holmberg at DTU for support on
326 Sugars, HMF and carotenoids analysis.

327 References

- 328 Abdullakassim P, Songchitsomboon S, Techagumpuch M et al (2007)
329 Antioxidant capacity, total phenolics and sugar content of
330 selected Thai health beverages. *Int J Food Sci Nutr* 58:77–85
331 Abdelmaksoud TG, Mohsen SM, Duedahl-Olesen L et al (2018a)
332 Effect of ohmic heating parameters on inactivation of enzymes
333 and quality of not-from-concentrate mango juice. *Asian J Sci*
334 *Res* 11(3):383–392
335 Abdelmaksoud TG, Mohsen SM, Duedahl-Olesen L et al (2018b)
336 Optimization of ohmic heating parameters for polyphenoloxi-
337 dase inactivation in not-from-concentrate elstar apple juice using
338 RSM. *J Food Sci Technol* 55(7):2420–2428
339 Abid M, Jabbar S, Wu T et al (2014) Sonication enhances
340 polyphenolic compounds, sugars, carotenoids and mineral ele-
341 ments of apple juice. *Ultrason Sonochemistry* 21:93–97
342 Andrews W (1992) Manuals of food quality control, microbiological
343 analysis, (chapter 4). Food and Drug Administration Washing-
344 ton, DC, USA. (FAO Food and Nutrition Paper, 14/4 Rev. 1,
345 FAO Consultant), M-82. ISBN: 92-5-103189-4
346 Bhale SD (2004) Effect of ohmic heating on color, rehydration and
347 textural characteristics of fresh carrot cubes. LSU Master's
348 Theses. p 3918. https://digitalcommons.lsu.edu/gradschool_theses/3918
349
350 Castro I, Macedo B, Teixeira JA, Vicente AA (2004) The effect of
351 electric field on important food-processing enzymes: comparison
352 of inactivation kinetics under conventional and ohmic heating.
353 *J Food Sci* 69:696–701
354 **AQ4** Chemat F, Zill-e-Huma, Khan MK (2011) Applications of ultrasound
355 in food technology: processing, preservation and extraction.
356 *Ultrason Sonochemistry* 18:813–835
357 Demirdöven A, Baysal T (2014) Optimization of ohmic heating
358 applications for pectin methylesterase inactivation in orange
359 juice. *J Food Sci Technol* 51:1817–1826
360 Esteve MJ, Barba FJ, Palop S, Frígola A (2009) The effects of non-
361 thermal processing on carotenoids in orange juice. *Czech J Food*
362 *Sci* 27:S304–S306

- Franco F, Pérez-Maqueda LA, Pérez-Rodríguez JL (2004) The effect
of ultrasound on the particle size and structural disorder of a
well-ordered kaolinite. *J Colloid Interface Sci* 274:107–117
Galaverna G, Dall'Asta C (2014) Production processes of orange
juice and effects on antioxidant components. In: Preedy VR (ed)
Processing and impact on antioxidants in beverages. Elsevier,
Amsterdam, pp 203–214
Giner MJ, Hizarci Ö, Martí N et al (2013) Novel approaches to reduce
Brown Pigment Formation and Color Changes in Thermal
Pasteurized Tomato Juice. *Euro Food Res Technol* 236:507–515
Girgin N, El SN (2015) Effects of cooking on in vitro sinigrin
bioaccessibility, total phenols, antioxidant and antimutagenic
activity of cauliflower (*Brassica oleraceae* L. var. Botrytis).
J Food Compos Anal 37:119–127
Kim DO, Jeong SW, Lee CY (2003) Antioxidant capacity of phenolic
phytochemicals from various cultivars of plums. *Food Chem*
81:321–326
Lee HS, Castle WS (2001) Seasonal change of carotenoid pigments
and color in Hamlin, Earlygold, and Budd Blood orange juices.
J Agric Food Chem 49:877–882
Leistner I (2000) Basic aspects of food preservation by hurdle
technology. *Inter J Food Micro* 55:181–186
Plaza L, Sánchez-Moreno C, De Ancos B et al (2011) Carotenoid and
flavanone content during refrigerated storage of orange juice
processed by high-pressure, pulsed electric fields and low
pasteurization, LWT – Food Sci Technol 44:834–839
Polydera AC, Stoforos NG, Taoukis PS (2005) Quality degradation
kinetics of pasteurised and high pressure processed fresh navel
orange juice: nutritional parameters and shelf life. *Technol*
Emerg Sci Food Innov 6(1):1–9
Ranganna S (1986) Handbook of analysis and quality control for fruit
and vegetable products. 2nd. Chapter 5 vitamins, pp 105–107.
ISBN-13: 978-0-07-451861-9
Rawson AB, Tiwari A, Patras N et al (2011) Effect of thermosoni-
cation on bioactive compounds in watermelon juice. *Food Res*
Inter 44:1168–1173
Ritter E, Purcell AE (1981) Carotenoid analytical methods. In:
Bavernfeind JC (ed) Carotenoids as colorants and vitamin A
precursors. Academic Press, New York, pp 815–883
Rouse AH, Atkins CD (1955) Pectinesterase and pectin in commer-
cial citrus juices as determined by methods used at the Citrus
Experiment Station. *Calif Agric Exp Stn Bull* 570:1–9
Roy MK, Juneja LR, Isobe S, Tshushida T (2009) Steam processed
broccoli (*Brassica oleracea*) has higher antioxidant activity in
chemical and cellular assay systems. *Food Chem* 114:263–269
Ting SV, Rouseff RL (1986) Measurement of quality for grades and
standards. Citrus fruits and their products: analysis and technol-
ogy. Marcel Dekker, Inc., New York, pp 35–65
USFDA (2001) Hazard analysis and critical control point (HACCP)
procedures for the safe and sanitary processing and importing of
juices: final rule. *Fed Reg* 66:6138–6202
Vorlova L, Borkovcová I, Kalábová K, Večerek V (2006) Hydrox-
ymethylfurfural contents in foodstuffs determined by HPLC
method. *J Food Nutr Res* 45: 34–38
Williams A (1994) New technologies in food preservation and
processing: part II. *Nutr Food Sci* 94(1):20–23
Zou Y, Jiang A (2016) Effect of ultrasound treatment on quality and
microbial load of carrot juice. *Food Sci Technol Campinas*
36:111–115

Publisher's Note Springer Nature remains neutral with regard to
jurisdictional claims in published maps and institutional affiliations.

Journal : **13197**

Article : **3834**

Author Query Form

Please ensure you fill out your response to the queries raised below and return this form along with your corrections

Dear Author

During the process of typesetting your article, the following queries have arisen. Please check your typeset proof carefully against the queries listed below and mark the necessary changes either directly on the proof/online grid or in the 'Author's response' area provided below

Query	Details Required	Author's Response
AQ1	Please confirm if the author names are presented accurately and in the correct sequence (given name, middle name/initial, family name). Author 1 Given name: [Tarek Gamal] Last name [Abdelmaksoud]. Author 2 Given name: [Sobhy Mohamed] Last name [Mohsen]. Author 4 Given name: [Mohamed Mohamed] Last name [Elnikeety]. Author 5 Given name: [Aberham Hailu] Last name [Feyissa]. Also, kindly confirm the details in the metadata are correct.	
AQ2	Reference Tiwari et al. (2008) is cited in text but not provided in the reference list. Please provide references in the list or delete these citations.	
AQ3	Please confirm the section headings are correctly identified.	
AQ4	Please provide author initials for the reference Chemat et al. (2011).	