Geomagnetic jerks and rapid hydromagnetic waves focusing at Earth's core surface

Aubert, Julien; Finlay, Christopher C.

Published in:
Nature Geoscience

Link to article, DOI:
[10.1038/s41561-019-0355-1](https://doi.org/10.1038/s41561-019-0355-1)

Publication date:
2019

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Aubert, J., & Finlay, C. C. (2019). Geomagnetic jerks and rapid hydromagnetic waves focusing at Earth's core surface. *Nature Geoscience*, 12(5), 393–398. <https://doi.org/10.1038/s41561-019-0355-1>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Geomagnetic jerks and rapid hydromagnetic waves focusing at Earth's core surface

Julien Aubert, Christopher Finlay

► To cite this version:

Julien Aubert, Christopher Finlay. Geomagnetic jerks and rapid hydromagnetic waves focusing at Earth's core surface. Nature Geoscience, Nature Publishing Group, 2019, 12 (5), pp.393-398. 10.1038/s41561-019-0355-1 . hal-02129842

HAL Id: hal-02129842

<https://hal.archives-ouvertes.fr/hal-02129842>

Submitted on 15 May 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Geomagnetic jerks and rapid hydromagnetic waves focusing at Earth's core surface

Julien Aubert^{1*} and Christopher C. Finlay²

¹*Institut de Physique du Globe de Paris, Sorbonne Paris Cité, Université Paris Diderot, CNRS UMR 7154 CNRS, F-75005 Paris, France. (*) : aubert@ipgp.fr*

²*Division of Geomagnetism, National Space Institute, Technical University of Denmark, Elektrovej, 2800 Kgs. Lyngby, Denmark*

Geomagnetic jerks are abrupt changes in the second time derivative - the secular acceleration- of Earth's magnetic field that punctuate ground observatory records. They are presently the major obstacle to the prediction of geomagnetic field behaviour years to decades ahead. Recent jerks have been linked to short-lived, temporally alternating and equatorially localised pulses of secular acceleration observed in satellite data, associated with rapidly alternating flows at Earth's core surface. The dynamical origin of jerks has been unclear but can now be investigated in numerical models of the geodynamo that realistically simulate the interaction between slow core convection and rapid hydromagnetic waves. Using one such model, here we show that the observed jerk patterns can be explained by the arrival of localised Alfvén wave packets radiated from sudden buoyancy releases inside the core. As they reach the core surface, the waves focus their energy towards the equatorial plane and along lines of strong magnetic flux, creating sharp interannual changes in core flow and producing geomagnetic jerks through the induced variations in field acceleration. The ability to numerically repro-

duce jerks offers a new way to probe the physical properties of Earth's deep interior.

The geomagnetic field displays temporal variations on a broad range of time scales. Through a self-sustained dynamo process (the geodynamo), slow convective motion in Earth's electrically conducting and liquid core is believed to maintain the field and drive its changes over centuries and longer. At the other end of the range, geomagnetic jerks with typical time scales a few years or less¹ represent the fastest observed features of the internally-generated field. They were initially identified as 'V-shaped' patterns (see examples in Fig. 1a) in time series of the magnetic field rate-of-change at ground observatories^{2,3} (the secular variation), indicating an abrupt change in the field acceleration amidst periods where this acceleration is otherwise approximately constant. Explaining the time scale disparity between rapid jerks and slow convection is a theoretical challenge that has recently spurred significant progress, both in observational geomagnetism and in numerical geodynamo simulations. In combination with an improving network of ground observatories, satellite magnetic field observations now provide a global and continuous view of the geomagnetic secular acceleration over the past two decades^{4,5}, with horizontal spatial resolution of approximately 2000 km at the core surface (spherical harmonic degree 9) and temporal resolution down to about a year on the largest length scales. This has dramatically enhanced our empirical knowledge of jerks, most notably by revealing⁶⁻⁹ their links to short-lived, temporally alternating pulses of geomagnetic acceleration at the surface of the Earth (Fig. 1c,e), that at the core surface are most prominent at low latitudes and localised in longitude (Fig. 2a,b). In the wake of these results, the earlier historical jerks of the twentieth century have also been linked to considerably less resolved, but similarly alternating pulses with low-latitude foci¹⁰. It has long been suspected that

jerks could somehow represent the signature of hydromagnetic waves¹¹. This prompted an earlier explanation¹² for jerks in terms of time-varying zonal flows that are kinematically consistent with torsional Alfvén waves occurring between concentric, magnetically-coupled, axial cylinders in the core. Though torsional waves have later been successfully identified in Earth’s core¹³ and in self-consistent numerical simulations of the geodynamo^{14–17}, they have however been found to occur at interannual periods shorter than earlier decadal estimates, and with an amplitude that is too weak to account for the geomagnetic secular acceleration signal associated with jerks^{17,18}. Furthermore, the complex patterns of magnetic acceleration found in satellite observations require localised (i.e. non-axisymmetric), rapidly alternating flows beneath the core surface^{5,19–21}. These discount an explanation in terms of torsional waves, but provide valuable new constraints on the rapid dynamics taking place in Earth’s core.

Rapid hydromagnetic waves in advanced numerical geodynamo simulations

Numerical simulations of convective core magnetohydrodynamics have been successful at describing the detailed morphology of the geomagnetic field^{22,23}, its temporal variations and the underlying core flows²⁴. To achieve this, the magnetic Reynolds number comparing the magnetic diffusion and convective core overturn time scales τ_η and τ_U (see definitions in Methods) needs to be $Rm = \tau_\eta/\tau_U \approx 1000$, such that a realistic $\tau_U \approx 100$ yr is achieved when adopting an Earth-like value $\tau_\eta \approx 10^5$ yr as the fundamental time scale for casting the dimensionless model results back to the dimensional world (Methods). However, most existing simulations remain unrealistic when considering time scales significantly shorter than τ_U . In Earth’s core, the dynamics of geomagnetic

jerks will involve strong rotational constraints from the Coriolis force, because the jerk's interan-
 nual time scale is much longer than Earth's planetary rotation period $2\pi\tau_\Omega = 1$ day. Jerk dynamics
 will also be affected by hydromagnetic waves given the proximity of the Alfvén time¹³ $\tau_A \approx 2$ yr.
 However, because of computational limitations^{16,25}, and despite continuous advances^{14–16,26,27}, nei-
 ther of these processes are correctly rendered in standard simulations where $2\pi\tau_\Omega$ and τ_A remain
 much too long, and not sufficiently separated from τ_U . For instance, in our previous Coupled
 Earth dynamo model²⁴ $2\pi\tau_\Omega \approx 10$ yr and $\tau_A \approx \tau_U \approx 100$ yr. In order to remedy these problems,
 we have recently introduced a suite of numerical simulations^{17,25} following a well-defined path
 through control parameter space that connects the original coupled Earth model to Earth's core
 conditions. Our new simulations (Methods and Supplementary Table 1) involve a reasonably ac-
 curate large-scale approximation that enables the exploration of parameters significantly beyond
 current computational limits for direct numerical simulations. Along this path, the Earth-like field
 morphology and kinematics of the coupled Earth model used as starting point are preserved, as are
 the values of τ_U and τ_η , but the dynamics gradually evolves as $2\pi\tau_\Omega$ and τ_A decrease to become
 realistic and increasingly separated from τ_U and τ_η . An asymptotic regime of strong rotational
 and magnetic control pertaining to Earth's core conditions is reached¹⁷ at path positions beyond
 30%. In addition to the slow background convection at time scale τ_U that is present throughout
 the path, models in this regime additionally feature rapid magneto-inertial wave dynamics at time
 scale τ_A . This dynamics includes geostrophic torsional Alfvén waves of weak amplitude, and also
 non-axisymmetric, quasi-geostrophic Alfvén waves¹⁷. The relevance of the latter waves, that were
 previously unexpected²⁸, has only recently been released^{17,29}. They offer a promising explana-

tion to jerks since they have been linked¹⁷ to the occurrence of intermittent magnetic acceleration pulses occurring at low latitudes. In our most advanced Midpath model (50% of the path, Methods and Supplementary Table 1), the Alfvén time scale value $\tau_A = 14.3$ yr implies that the waves have interannual periods at wavelengths a fraction of the core size, well separated from convective processes. Since $\tau_A \gg 2\pi\tau_\Omega = 0.19$ yr, such periods are also well within the rotationally-dominated range where the Coriolis force plays a crucial role. Finally, a high ratio $\tau_\eta/\tau_A \approx 10^4$ of the magnetic diffusion and Alfvén time scales indicates that wave attenuation will be weak on large length scales.

Observed and simulated geomagnetic jerks

Short-lived, intermittent pulses in the magnetic acceleration energy at Earth’s surface (Fig. 1d) are observed in numerical simulations throughout the parameter space path, but migrate to low latitudes (Fig. 1f) only once the model conditions enter the rapid rotation regime¹⁷. In order to highlight the link between such pulses and jerks, and to facilitate comparison with geomagnetic field models of limited temporal resolution (Fig. 1c,e), we define jerk energy (Methods) as the mean-squared difference between time averages of Earth’s surface magnetic acceleration taken within two consecutive and non-overlapping 3-year time windows. With this definition, the timing of jerks in the simulation (Fig. 1d) can be properly defined from jerk energy pulses, and is found to either shortly precede or follow that of magnetic acceleration pulses, as observed with well-documented recent geomagnetic jerks⁶ (Fig. 1c). The intensity and duration of the events also match rather well the observations. Abrupt slope changes in the magnetic variation time se-

103 ries are observed at specific locations (Fig. 1b), with approximately constant acceleration away
 104 from the event, similar to the classic 'V-shaped' jerk signatures seen at ground observatories¹⁻³
 105 (Fig. 1a). Maps of the radial magnetic acceleration before and after the events feature alternating
 106 patterns (Fig. 1f) and indicate that simulated jerks are often visible over a large area (from Amer-
 107 ica to Indonesia for the event shown here) at low and mid-latitudes, comparable to observations
 108 of the well-characterised 2007.5 geomagnetic jerk (Fig. 1e) and to a number of earlier events^{30,31}.
 109 Descending to the core surface (Fig. 2c,d), the corresponding structures are series of intense and
 110 oppositely-signed patches of radial magnetic acceleration generated close to the equator and in a
 111 narrow longitudinal band, beneath westward-drifting patches of intense radial magnetic flux (see
 112 Fig. 3e) localised in the Atlantic hemisphere^{17,24}. The field acceleration patches alternate rapidly
 113 in time for a few years (Supplementary Movie 1 from time -10 yr to 10 yr) before fading away.
 114 The spatially localised morphology, interannual alternation time scale and the amplitude (approx-
 115 imately 2,000 nT.yr⁻² up to degree 9) of the simulation output reproduce well the core surface
 116 signature of geomagnetic jerks^{5,8} (Fig. 2a,b, see the events in 2007.5, 2011 and 2014.5).

117 **The origin of geomagnetic jerks and the role of hydromagnetic waves**

118 At the large scales accessible to observations (spherical harmonic degree up to 9), and in
 119 the rapid rotation regime, the magnetic acceleration pulses in the simulations result from the ac-
 120 tion of accelerating azimuthal core surface flows¹⁷ rather than from diffusive processes related to
 121 flux expulsion that are common at the start of the parameter space path. In our Midpath model
 122 sequence, a localised, intense and temporally alternating pulse of azimuthal flow acceleration is

indeed observed in the vicinity of the jerk time (Fig 2e,f, Supplementary Movie 2), resembling the localised alternating flows that have been inferred from geomagnetic variations⁵ associated with the 2007.5 geomagnetic jerk. The source of this perturbation can be traced back to a sudden buoyancy release from an isolated density anomaly at mid-depth in the core, 25 years before the event (Supplementary Movies 3,4). This release triggers strong azimuthal fluid flow accelerations that are entrained within the associated convective plume towards the core surface. The plume stalls at a cylindrical radius $s_c \approx 2950$ km (Supplementary Movie 3, Supplementary Fig. 1) where its decreasing radial velocity is overcome by the global westward drift. At cylindrical radii above s_c , we identify quasi-geostrophic Alfvén waves^{17,29} through the adherence of their trajectories to propagation at the locally variable, theoretical Alfvén speed and the deviation of their paths from that of material upwellings (Methods and Supplementary Fig. 1). In the upper outer core, material upwelling is indeed much slower than Alfvén waves, or even directed inwards. The perturbation energy however propagates further towards the core surface in well-defined, azimuthally extended, alternating wavefronts (Fig. 3a,b, Supplementary Movie 4) of columnar structure characteristic of rotationally-dominated dynamics¹⁷. The waves have a radial wavelength $d \approx D/4$ that is in line with the size of the density anomaly that initiated the event. Their energy becomes spatially concentrated as they approach the core-mantle boundary (Fig. 3c), yielding the intense, localised and temporally alternating surface flow acceleration signature (Fig. 2e,f) that causes the jerk, on a time scale comparable to the Alfvén wave period for these structures, $\sqrt{3}\tau_A d/D \approx 6$ yr. The energy concentration mechanism can be understood by noting that quasi-geostrophic Alfvén wavefronts are both guided along, and bounded by a strongly heterogeneous distribution of magnetic

field lines¹⁷. Beneath the jerk location, these field lines are arranged in an approximately axially-invariant funnel-like structure (Fig. 3d) that is shaped by the slow convection and remains approximately static during the event. This causes the waves to be longitudinally focused towards a pair of intense radial magnetic flux patches (see arrows in Fig. 3e) at the core surface. At the same time, latitudinal focusing towards the equator occurs because of the effect of the spherical core-mantle boundary on flow columns that tend to preserve their angular momentum as their height decreases (Fig. 3b, Supplementary Video 4). Finally, the wave speed decreases close to the core-mantle boundary (see curved green tracks in Supplementary Fig. 1) because the magnetic field is weaker at the surface than at depth²⁵. To preserve the energy flux, the amplitude of wavefronts increases, and preservation of the wave period also implies a reduction of the radial wavelength (Fig. 3c), similarly to a shoaling process for water waves³². This three-dimensional energy focusing mechanism is crucial in amplifying the weak quasi-geostrophic Alfvén waves so as to produce localised and temporally alternating disturbances in the core surface flow acceleration that are significant enough to cause jerks.

Implications for geomagnetism and global geodynamics

Since our models are in the dynamical regime of rapid rotation and strong magnetic control relevant to Earth’s core^{17,25}, their results can be extrapolated to natural conditions. From the mechanism described here, the duration and alternation time scale of jerk events are expected to scale with the Alfvén time τ_A , which is about 7 times shorter in Earth’s core¹³ than in our Midpath simulation (Supplementary Table 1). Yet the observed geomagnetic acceleration changes are only two

164 to three times faster than those simulated by the Midpath model (Fig. 2). This discrepancy is likely
 165 related to the limited temporal resolution of present geomagnetic field models, which prevents the
 166 true, potentially sub-annual³³ variations associated with jerks from being retrieved. In the upcom-
 167 ing years, further insight will be obtained from jerk events that will be imaged with improved
 168 resolution using data collected by the *Swarm* satellite mission. As we move along the parameter
 169 space path, our models also indicate that energetic jerks occur more frequently (Fig. 4a). It is pos-
 170 sible to construct statistical relationships between jerk energy and recurrence time (Fig. 4b), and
 171 derive a scaling relationship for the evolution along the path of jerk energy at a given recurrence
 172 time (Supplementary Fig. 2) in reasonable agreement with a theoretical prediction (Methods). The
 173 extrapolation of this relationship to the end of the path (Fig. 4b) also agrees with the observed sub-
 174 decadal to decadal jerk recurrence rates observed in the geomagnetic field^{1,5}. Jerk energy is also
 175 found to decrease with increasing lower mantle conductance, because of the associated additional
 176 Ohmic losses, and with increasing levels of upper outer core stratification (Supplementary Fig. 3).
 177 This latter effect is due to changes in the geometry and amplitude of the background magnetic field
 178 rather than to the wave mechanism itself, which is not sensitive to stratification. Finally, exam-
 179 ining simulated records of the length of day (Methods, Supplementary Fig. 4) in the vicinity of
 180 jerk events, we also observe signatures of the wave's arrival at the core surface. Rapid inflexions
 181 in the rate of change of the length-of-day similar to those observed for Earth^{31,34} are caused by
 182 pulses in the acceleration of the electromagnetic torque felt by the mantle. All these results high-
 183 light the potential importance of the numerical reproduction of jerks, as it may lead to an improved
 184 geomagnetic^{35,36} and geodetic³⁷ sounding of important, but poorly known physical properties such

as the lower mantle electrical conductivity and upper outer core thermal conductivity.

The integration of geomagnetic data into numerical geodynamo simulations through data assimilation has significantly advanced in the recent past³⁸, leading to inferences of the dynamical internal structure of the geodynamo and to predictions of the future geomagnetic field evolution³⁹ that have been integrated within the latest iteration^{40,41} of the International Geomagnetic Reference Field (IGRF). At interannual to decadal time scales, the accuracy of such predictions is currently hampered by the underlying dynamical model, which is located at the start of the parameter space path and hence does not account for wave dynamics. The availability of advanced numerical dynamo simulations that produce realistic rapid dynamics and jerks will significantly improve the quality of the prior information on which the predictions are based (in particular regarding the time-dependence of the field), with subsequent gains in their accuracy.

196 Methods

197 **Model description.** The full description of our numerical models can be found in refs. ^{17,25}. We
198 solve for Boussinesq convection, thermochemical density anomaly transport and magnetic induc-
199 tion in the magnetohydrodynamic approximation within an electrically conducting and rotating
200 spherical fluid shell of thickness $D = r_o - r_i$ representing the outer core, with $r_i/r_o = 0.35$ as in the
201 Earth. Our unknowns are the velocity field $\mathbf{u}$, magnetic field $\mathbf{B}$ and density anomaly field C , and
202 we analyse the magnetic variation $\partial\mathbf{B}/\partial t$, magnetic acceleration $\partial^2\mathbf{B}/\partial t^2$ and the flow acceleration
203 $\partial\mathbf{u}/\partial t$. The fluid shell is electromagnetically coupled both to a solid inner core of radius r_i and to a
204 solid outer shell representing the mantle between radii r_o and $1.83r_o$. The inner core and mantle are
205 furthermore coupled together by a gravitational restoring torque. Both the inner core and mantle
206 feature a time-dependent axial differential rotation with respect to the outer core. The three regions
207 are assigned moments of inertia respecting the proportions²⁴ relevant to Earth’s mantle, inner and
208 outer core, and the ensemble has a constant angular momentum defining the planetary rotation rate
209 Ω .

210 The mechanical boundary conditions are of the stress-free type at both boundaries. In the low
211 viscosity regime where our models operate, these are undistinguishable from no-slip conditions²⁵
212 while alleviating the need to resolve the viscous boundary layers. Electrically conducting bound-
213 ary conditions are used at both boundaries. The electrical conductivity of the inner core is set
214 at the same value σ_c as that of the outer core. The mantle features an electrically conducting
215 region at its base, with thickness Δ and conductivity σ_m . In our four main model cases (Supple-
216 mentary Table 1) the dimensionless conductance has been set to a median geophysical estimate⁴²

217 $\Sigma = \Delta\sigma_m/D\sigma_c = 10^{-4}$. Two other models (Midpath-I and Midpath-H) explore the end-member
 218 values $\Sigma = 0$ (insulating mantle) and $\Sigma = 10^{-3}$. The thermochemical boundary conditions are
 219 of heterogeneous, fixed-flux type. The homogeneous part F of the density anomaly flux is pre-
 220 scribed at the inner boundary. In our four main model cases the homogeneous density anomaly
 221 flux vanishes at the outer boundary (neutral buoyancy). A volumetric sink term is then present
 222 in the density anomaly transport equation to conserve mass. Within the Boussinesq approxima-
 223 tion, this configuration models bottom-driven chemical convection originating from inner core
 224 solidification, a fully convective outer core and an exactly adiabatic heat flow at the core-mantle
 225 boundary. An additional model (Midpath-S) explores the effect of a possible stratification of the
 226 upper outer core⁴³ by prescribing an negative (adverse buoyancy) density anomaly flux at the core-
 227 mantle boundary (see Stratified Core section below). Spatial modulations of the density anomaly
 228 fluxes are prescribed at both boundaries¹⁷, with the same geometry as in the coupled Earth model²⁴.
 229 These are meant to model a spatially heterogeneous growth of the inner core, and thermal control
 230 from the heterogeneous lower mantle.

231 **Model parameters, parameter space path and time scales.** The four main control parameters
 232 of the model are the flux-based Rayleigh, Ekman, Prandtl and magnetic Prandtl numbers

$$Ra_F = \frac{g_o F}{4\pi\rho\Omega^3 D^4}, \quad (1)$$

$$E = \frac{\nu}{\Omega D^2}, \quad (2)$$

$$Pr = \frac{\nu}{\kappa}, \quad (3)$$

$$Pm = \frac{\nu}{\eta}. \quad (4)$$

Here g_o , ρ , ν , κ and η are respectively the gravity at the outer boundary of the model, the fluid density, viscosity, thermo-chemical and magnetic diffusivities ($\eta = 1/\mu\sigma_c$, with μ the fluid magnetic permeability). We have recently introduced²⁵ the concept of a unidimensional path in parameter space, by showing that the variations in these control parameters that are necessary to bridge the gap between our previous coupled Earth model²⁴ and Earth's core conditions can be represented as power laws of a single variable ϵ . Any model along the path is defined using the following rules:

$$Ra_F = \epsilon Ra_F(\text{CE}), \quad (5)$$

$$E = \epsilon E(\text{CE}), \quad (6)$$

$$Pr = 1, \quad (7)$$

$$Pm = \sqrt{\epsilon} Pm(\text{CE}). \quad (8)$$

Here $Ra_F(\text{CE}) = 2.7 \cdot 10^{-5}$, $E(\text{CE}) = 3 \cdot 10^{-5}$ and $Pm(\text{CE}) = 2.5$ are the control parameters of the coupled Earth dynamo model defining the start of the path ($\epsilon = 1$), and we have shown²⁵ that conditions relevant to Earth's core are reached at the end of path defined by $\epsilon = 10^{-7}$. Our models are defined in refs. ^{17,25} and in Supplementary Table 1 by the values $\epsilon = 10^{-2}$, $3.33 \cdot 10^{-3}$, 10^{-3} and $3.33 \cdot 10^{-4}$, respectively corresponding to 29%, 35%, 43%, and 50% of the path (the Midpath model). The parameters of the Midpath model are the closest to Earth's core conditions employed to date in a numerical dynamo simulation, at the expense of a large scale approximation (see Numerical Implementation section below).

The model outputs follow scaling laws²⁵ depending on ϵ that also closely approach the conditions expected in Earth's core as we progress along the path (Supplementary Table 1). Once the

249 magnetic diffusion time $\tau_\eta = D^2/\eta$ is set to an Earth-like value (see Rescaling section below), the
 250 end of path simultaneously matches the Earth's core rotational time $\tau_\Omega = 1/\Omega$, convective overturn
 251 time $\tau_U = D/U$, and Alfvén time $\tau_A = \sqrt{\rho\mu}D/B$ (here U and B are respectively the root-mean-
 252 squared velocity and dynamo-generated magnetic field in the fluid shell). Numerical models taken
 253 along this path can therefore be understood as continuously progressing from imperfect towards
 254 geophysically appropriate conditions in all relevant aspects of their inputs and outputs. The dimen-
 255 sional values of τ_Ω , τ_U , and τ_A reached in our models and at the end of path are listed together with
 256 Earth's core estimates in Supplementary Table 1 (see ref. ¹⁷ for a complete list of dimensionless
 257 time scale ratios achieved in the models).

258 **Stratified core case.** The Midpath-S model (Supplementary Table 1) explores the effects of a pos-
 259 sible upper outer core stratification⁴³ on the occurrence of simulated jerks. Within the Boussinesq
 260 approximation, stratification is modelled by adding an adverse density anomaly gradient⁴⁴ to the
 261 background gradient prescribed by the neutral buoyancy conditions described above:

$$\frac{dC}{dr} = \frac{-N^2\rho}{2g_o} (1 + \tanh((r - r_s)/\delta)). \quad (9)$$

262 Here N is the Brunt-Väisälä frequency pertaining to the stratification level at the core surface, r
 263 is radius, $r_s = 3340$ km is the radius at which stratification sets in, and $\delta = 10^{-2}D = 22.6$ km is
 264 the thickness of the stratified layer front. The thickness of the stratified layer is $r_o - r_s = 140$ km,
 265 as proposed in ref. ⁴³. In the Midpath-S model we set $N = 1/\tau_\Omega$, as also proposed in ref. ⁴³.
 266 The output of the Midpath-S model demonstrates the preservation of simulated jerks against core
 267 stratification, albeit at a reduced energy level given the modifications of the background magnetic
 268 field that guides the waves.

Dimensional rescaling of dimensionless model output. The dimensionless model length unit is adjusted to the thickness $D = 2260$ km of Earth's core. Time is rescaled by adjusting the magnetic diffusion time scale $\tau_\eta = D^2/\eta$ to the value $\tau_\eta = 135\,000$ yr, corresponding to a value $\eta = 1.2$ m²/s at the midpoint of current estimates²⁵. Given the invariance of the magnetic Reynolds number $Rm = \tau_\eta/\tau_U \approx 1\,000$ along the parameter space path, this rescaling choice ensures $\tau_U \approx 130$ yr and Earth-like convective geomagnetic variations¹⁷. The fluid and Alfvén wave velocities are rescaled by using these length and time units. The magnetic field amplitude is presented by setting the Elsasser magnetic field unit $\sqrt{\rho\mu\eta\Omega}$ to the value 0.9 mT. Given the approximate invariance of the Elsasser number $B^2/\rho\mu\eta\Omega \approx 20$ along the path²⁵, this amounts to setting the root-mean-squared field amplitude within the core to a value of about 4 mT, in agreement with Earth's core current estimate¹³. Note that concerning the time and magnetic field units our choices slightly differ (by less than 5%) from ref. ¹⁷, as we adopt here the same units across all simulations. This change is done in order to obtain a consistent comparison between the original path models and those with a modified setup (Midpath-S,I,H, Supplementary Table 1) introduced in this study. Finally, the density anomaly rescaling used in Supplementary Movie 3 follows from the velocity rescaling and from adjustment of the dimensionless, time average convective power in the shell to an estimate²⁵ $P = 3$ TW of the geodynamo power.

Jerk energy definition and scaling. In Fig. 1c,d, we present the energy E_{SA} of the magnetic acceleration, defined as a mean-squared average over Earth's surface S_E :

$$E_{SA} = \left\langle (\partial_t^2 \mathbf{B})^2 \right\rangle = \frac{1}{S_E} \int_{S_E} \left(\frac{\partial^2 \mathbf{B}}{\partial t^2} \right)^2 dS. \quad (10)$$

288 To facilitate comparison of the model output with geomagnetic field models of limited temporal
 289 resolution, the jerk energy E_J is defined as a sliding finite difference between consecutive 3-yr time
 290 windows rather than an instantaneous rate-of-change:

$$E_J(t) = \left\langle \left(\left[\partial_t^2 \mathbf{B} \right]_t^{t+3 \text{ yr}} - \left[\partial_t^2 \mathbf{B} \right]_{t-3 \text{ yr}}^t \right)^2 \right\rangle. \quad (11)$$

291 As introduced above, the angle brackets denote the average over Earth's surface, and the square
 292 brackets denote a time average. Jerk recurrence statistics in Fig. 4b are obtained from time series
 293 (Fig. 4a) of E_J , by dividing the duration of the model run with the number of samples reaching
 294 or exceeding a given jerk energy. In Supplementary Fig. 2, jerk energies at 10, 30 and 100 yr
 295 recurrence times are extracted from Fig. 4b and scaled with the path parameter ϵ , revealing a
 296 common dependency in $\epsilon^{-0.19 \pm 0.01}$. The end-of-path prediction in Fig. 4b is obtained by collapsing
 297 the jerk statistics onto a single master curve according to this scaling, and extrapolating the master
 298 curve to the end-of-path conditions corresponding to $\epsilon = 10^{-7}$.

299 The amplitude of a secular acceleration pulse scales with the magnetic field amplitude B
 300 times the wave-induced flow acceleration U/τ_A . Given that the pulse duration should also scale
 301 with τ_A , jerk energy then scales with

$$E_J \sim \frac{\tau_A}{3 \text{ yr}} (BU/\tau_A)^2 \sim \frac{(BU)^2}{\tau_A} \quad (12)$$

302 Along the parameter space path, the dimensional values of U and B are approximately preserved,
 303 and the above scaling suggests that E_J should be inversely proportional to τ_A . The Alfvén number
 304 $A = \tau_A/\tau_U$ has been shown²⁵ to scale like $\epsilon^{0.25}$ along the path. The invariance of τ_U along the path
 305 (Supplementary Table 1) then leads to $\tau_A \sim \epsilon^{0.25}$ and $E_J \sim \epsilon^{-0.25}$, close to the numerical result

306 $E_J \sim \epsilon^{-0.19}$. The residual discrepancy mainly stems from the value of B which slightly decreases
 307 along the path (see ref. ²⁵).

308 **Identification of quasi-geostrophic Alfvén waves.** In Supplementary Fig. 1 we repeat the analy-
 309 sis carried out in ref. ¹⁷ to identify hydromagnetic wave propagation. The flow acceleration patterns
 310 that we analyse have a columnar structure that derives from the dominant rotational constraint of
 311 the Coriolis force. At any given time t , cylindrical radius s and at a fixed analysis longitude φ_0 , we
 312 therefore first compute the columnar average $\partial u_c / \partial t$ of azimuthal flow acceleration:

$$\frac{\partial u_c}{\partial t}(s, \varphi_0, t) = \frac{1}{z_+ - z_-} \int_{z_-}^{z_+} \frac{\partial(\mathbf{u} \cdot \mathbf{e}_\varphi)}{\partial t}(s, \varphi_0, z, t) dz. \quad (13)$$

313 Here s, φ, z are cylindrical coordinates, $\mathbf{e}_\varphi$ is the unit vector in the azimuthal direction, and the
 314 vertical integral is evaluated between the lower and upper heights $z_{-,+}$ of an axial column parallel
 315 to the rotation vector $\mathbf{\Omega}$ at cylindrical radius s . We then represent time-cylindrical radius maps of
 316 $\partial u_c / \partial t$ and overplot ray-tracing theoretical propagation tracks obtained by integrating in time the
 317 column-averaged Alfvén velocity c_A and column-averaged cylindrical radial fluid velocity V_s :

$$c_A(s, \varphi_0, t) = \sqrt{\frac{1}{z_+ - z_-} \int_{z_-}^{z_+} \frac{(\mathbf{B} \cdot \mathbf{e}_s)^2}{\rho \mu}(s, \varphi_0, z, t) dz}, \quad (14)$$

$$V_s(s, \varphi_0, t) = \frac{1}{z_+ - z_-} \int_{z_-}^{z_+} \mathbf{u}(s, \varphi_0, z, t) \cdot \mathbf{e}_s dz. \quad (15)$$

318 Here $\mathbf{e}_s$ is the unit vector in the cylindrical radial direction. The adherence of $\partial u_c / \partial t$ to the Alfvén
 319 tracks and deviation from material upwelling tracks demonstrates Alfvén wave propagation.

320 **Length-of-day variations.** The numerical simulation solves for the deviations Ω_M of the mantle
 321 angular velocity from the background planetary rotation rate Ω (see ref. ⁴² for details):

$$I_M \frac{d\Omega_M}{dt} = \Gamma_M + \Gamma_G. \quad (16)$$

Here I_M is the Earth's mantle moment of inertia, and $\Gamma_{M,G}$ are respectively the magnetic and gravitational torques felt by the mantle. The corresponding rate of change $d(\text{LOD})/dt$ in the length of the day is then

$$\frac{d(\text{LOD})}{dt} = -\frac{2\pi}{\Omega^2} \frac{d\Omega_M}{dt}, \quad (17)$$

where we have used $\Omega_M \ll \Omega$. Time series of $d(\text{LOD})/dt$ in the vicinity of jerk events are presented in Supplementary Fig. 4. The magnetic acceleration pulses cause pulses in $d^2\Gamma_M/dt^2$, and hence rapid inflexions in $d(\text{LOD})/dt$ with shape similar to that observed in geodetic time series^{31,34}. Note though that the amplitude of the inflexions is significantly weaker in the numerical simulations than in Earth's core, because the inverse squared Alfvén number $1/A^2$ measuring the relative importance of magnetic forces and inertia is about 50 times weaker²⁵ in the Midpath model than in the core.

Numerical Implementation. Our numerical implementation involves a decomposition of the fields in spherical harmonics up to degree and order 133, and a discretisation in the radial direction on a second-order finite-differencing scheme (see ref. ²⁵ for numerical resolution details). We use the spherical harmonics transform library⁴⁵ SHTns freely available at <https://bitbucket.org/nschaeff/shtns>. Time stepping is of second-order, semi-implicit type. Angular momentum conservation is controlled at each time step. To handle an increasing hydrodynamic turbulence along the path that does however only weakly affect the large-scale solution²⁵, hyperdiffusion is implemented on the velocity and density anomaly fields, but not on the magnetic field which remains fully resolved. The details, physical justification and validation of this approximation are presented in ref. ²⁵. Each model on the path is initialised using the output of the previous step. Integration times after statistical equilibration are listed in Supplementary Table 1. In our main

342 models these represent at least 18% of a magnetic diffusion time and 75% of a dipole decay time
343 $r_o^2/\pi^2\eta$. Within this time all model outputs are in a statistically-steady state¹⁷ demonstrating self-
344 sustained dynamo action. In particular, all models produced an axial dipole-dominated magnetic
345 field that did not reverse polarity.

346 **Data availability.** The numerical code and the simulation datasets analysed during the current
347 study are available from the corresponding author on reasonable request.

- 348 1. Brown, W., Mound, J. & Livermore, P. Jerks abound: An analysis of geomagnetic observatory
350 data from 1957 to 2008. *Phys. Earth Planet. Int.* **223**, 62 – 76 (2013).
- 351 2. Courtillot, V., Ducruix, J. & Le Mouél, J.-L. Sur une accélération récente de la variation
352 séculaire du champ magnétique terrestre. *C.R. Acad. Sci. Paris. D* **287**, 1095–1098 (1978).
- 353 3. Malin, S. R. C., Hodder, B. M. & Barraclough, D. R. Geomagnetic secular variation: A jerk in
354 1970. In Cardús, J. O. (ed.) *Scientific Contributions in Commemoration of Ebro Observatory's*
355 *75th Anniversary* (1983).
- 356 4. Lesur, V., Wardinski, I., Hamoudi, M. & Rother, M. The second generation of the GFZ
357 Reference Internal Magnetic Model: GRIMM-2. *Earth Planets Space* **62**, 765–773 (2010).
- 358 5. Finlay, C. C., Olsen, N., Kotsiaros, S., Gillet, N. & Tøffner-Clausen, L. Recent geomag-
359 netic secular variation from Swarm and ground observatories as estimated in the CHAOS-6
360 geomagnetic field model. *Earth, Planets and Space* **68**, 112 (2016).
- 361 6. Chulliat, A., Thebault, E. & Hulot, G. Core field acceleration pulse as a common cause of the
362 2003 and 2007 geomagnetic jerks. *Geophys. Res. Lett.* **37** (2010).
- 363 7. Chulliat, A. & Maus, S. Geomagnetic secular acceleration, jerks, and a localized standing
364 wave at the core surface from 2000 to 2010. *J. Geophys. Res.* **119**, 1531–1543 (2014).
- 365 8. Chulliat, A., Alken, P. & Maus, S. Fast equatorial waves propagating at the top of the Earth's
366 core. *Geophys. Res. Lett.* **42**, 3321–3329 (2015).

- 367 9. Torta, J. M., Pavón-Carrasco, F. J., Marsal, S. & Finlay, C. C. Evidence for a new geomagnetic
368 jerk in 2014. *Geophys. Res. Lett.* **42**, 7933–7940 (2015).
- 369 10. Pais, M. A., Alberto, P. & Pinheiro, F. J. G. Time-correlated patterns from spherical harmonic
370 expansions: Application to geomagnetism. *J. Geophys. Res.* **120**, 8012–8030 (2015).
- 371 11. Braginsky, S. I. Short-period geomagnetic secular variation. *Geophys. Astrophys. Fluid Dyn.*
372 **30**, 1–78 (1984).
- 373 12. Bloxham, J., Zatman, S. & Dumberry, M. The origin of geomagnetic jerks. *Nature* **420**, 65–68
374 (2002).
- 375 13. Gillet, N., Jault, D., Canet, E. & Fournier, A. Fast torsional waves and strong magnetic field
376 within the Earth’s core. *Nature* **465**, 74–77 (2010).
- 377 14. Wicht, J. & Christensen, U. R. Torsional oscillations in dynamo simulations. *Geophys. J. Int.*
378 **181**, 1367–1380 (2010).
- 379 15. Teed, R. J., Jones, C. A. & Tobias, S. M. The dynamics and excitation of torsional waves in
380 geodynamo simulations. *Geophys. J. Int.* **196**, 724–735 (2014).
- 381 16. Schaeffer, N., Jault, D., Nataf, H.-C. & Fournier, A. Turbulent geodynamo simulations: a leap
382 towards Earth’s core. *Geophys. J. Int.* **211**, 1–29 (2017).
- 383 17. Aubert, J. Geomagnetic acceleration and rapid hydromagnetic wave dynamics in advanced
384 numerical simulations of the geodynamo. *Geophys. J. Int.* (2018, in press). Preprint available
385 at <http://www.arxiv.org/abs/1804.04865>.

- 386 18. Cox, G., Livermore, P. & Mound, J. The observational signature of modelled torsional waves
387 and comparison to geomagnetic jerks. *Phys. Earth Planet. Int.* **255**, 50 – 65 (2016).
- 388 19. Wardinski, I., Holme, R., Asari, S. & Manda, M. The 2003 geomagnetic jerk and its relation
389 to the core surface flows. *Earth Plan. Sci. Lett.* **267**, 468 – 481 (2008).
- 390 20. Silva, L. & Hulot, G. Investigating the 2003 geomagnetic jerk by simultaneous inversion of
391 the secular variation and acceleration for both the core flow and its acceleration. *Phys. Earth
392 Planet. Int.* **198–199**, 28 – 50 (2012).
- 393 21. Gillet, N., Jault, D. & Finlay, C. C. Planetary gyre, time-dependent eddies, torsional waves
394 and equatorial jets at the Earth’s core surface. *J. Geophys. Res.* **120**, 3991–4013 (2015).
- 395 22. Christensen, U. R., Aubert, J. & Hulot, G. Conditions for Earth-like geodynamo models.
396 *Earth. Plan. Sci. Lett.* **296**, 487–496 (2010).
- 397 23. Mound, J., Davies, C. & Silva, L. Inner core translation and the hemispheric balance of the
398 geomagnetic field. *Earth Plan. Sci. Lett.* **424**, 148 – 157 (2015).
- 399 24. Aubert, J., Finlay, C. C. & Fournier, A. Bottom-up control of geomagnetic secular variation
400 by the Earth’s inner core. *Nature* **502**, 219–223 (2013).
- 401 25. Aubert, J., Gastine, T. & Fournier, A. Spherical convective dynamos in the rapidly rotating
402 asymptotic regime. *J. Fluid. Mech.* **813**, 558–593 (2017).
- 403 26. Yadav, R. K., Gastine, T., Christensen, U. R., Wolk, S. J. & Poppenhaeger, K. Approaching a
404 realistic force balance in geodynamo simulations. *PNAS* **113**, 12065–12070 (2016).

- 405 27. Sheyko, A., Finlay, C. C. & Jackson, A. Magnetic reversals from planetary dynamo waves.
406 *Nature* **539**, 551–554 (2016).
- 407 28. Labbé, F., Jault, D. & Gillet, N. On magnetostrophic inertia-less waves in quasi-geostrophic
408 models of planetary cores. *Geophys. Astrophys. Fluid Dyn.* **109**, 587–610 (2015).
- 409 29. Bardsley, O. P. & Davidson, P. A. Inertial–Alfvén waves as columnar helices in planetary
410 cores. *J. Fluid. Mech.* **805** (2016).
- 411 30. Dormy, E. & Manda, M. Tracking geomagnetic impulses at the core–mantle boundary. *Earth*
412 *Plan. Sci. Lett.* **237**, 300 – 309 (2005).
- 413 31. Manda, M. *et al.* Geomagnetic Jerks: Rapid Core Field Variations and Core Dynamics.
414 *Space. Sci. Rev.* **155**, 147–175 (2010).
- 415 32. Dean, R. G. & Dalrymple, R. A. *Water wave mechanics for engineers and scientists*, vol. 2
416 (World Scientific Publishing Company, 1991).
- 417 33. Olsen, N. & Manda, M. Rapidly changing flows in the Earth’s core. *Nature Geosci.* **1**,
418 390–394 (2008).
- 419 34. Holme, R. & de Viron, O. Geomagnetic jerks and a high-resolution length-of-day profile for
420 core studies. *Geophys. J. Int.* **160**, 435–439 (2005).
- 421 35. Lesur, V., Whaler, K. & Wardinski, I. Are geomagnetic data consistent with stably stratified
422 flow at the core–mantle boundary? *Geophys. J. Int.* **201**, 929–946 (2015).

- 423 36. Schaeffer, N. & Jault, D. Electrical conductivity of the lowermost mantle explains absorption
424 of core torsional waves at the equator. *Geophys. Res. Lett.* **43**, 4922–4928 (2016).
- 425 37. Holme, R. & de Viron, O. Characterization and implications of intradecadal variations in
426 length of day. *Nature* **499**, 202–204 (2013).
- 427 38. Fournier, A. *et al.* An Introduction to Data Assimilation and Predictability in Geomagnetism.
428 *Space. Sci. Rev.* **155**, 247–291 (2010).
- 429 39. Aubert, J. Geomagnetic forecasts driven by thermal wind dynamics in the Earth’s core. *Geo-*
430 *phys. J. Int.* **203**, 1738–1751 (2015).
- 431 40. Fournier, A., Aubert, J. & Thébault, E. A candidate secular variation model for IGRF-12 based
432 on Swarm data and inverse geodynamo modelling. *Earth. Planets. Space.* **67**, 81 (2015).
- 433 41. Thébault, E. *et al.* International geomagnetic reference field: the twelfth generation. *Earth.*
434 *Planets. Space.* **67**, 79 (2015).
- 435 42. Pichon, G., Aubert, J. & Fournier, A. Coupled dynamics of Earth’s geomagnetic westward
436 drift and inner core super-rotation. *Earth Planet. Sci. Lett.* **437**, 114–126 (2016).
- 437 43. Buffett, B. A. Geomagnetic fluctuations reveal stable stratification at the top of the Earth’s
438 core. *Nature* **507**, 484–487 (2014).
- 439 44. Nakagawa, T. Effect of a stably stratified layer near the outer boundary in numerical simula-
440 tions of a magnetohydrodynamic dynamo in a rotating spherical shell and its implications for
441 Earth’s core. *Phys. Earth. Planet. Int.* **187**, 342 – 352 (2011).

- 442 45. Schaeffer, N. Efficient spherical harmonic transforms aimed at pseudospectral numerical sim-
443 ulations. *Geophys. Geochem. Geosystems*. **14**, 751–758 (2013).

Acknowledgements JA acknowledges support from French *Programme National de Planétologie* (PNP) of CNRS/INSU, and from the Fondation Del Duca of Institut de France. This work was granted access to the HPC resources of S-CAPAD, IPGP, France, and to the HPC resources of IDRIS and CINES under the allocation 2016-A0020402122 made by GENCI. The results presented in this work rely on data collected at magnetic observatories. The authors thank the national institutes that support them and INTERMAGNET for promoting high standards of magnetic observatory practice (www.intermagnet.org).

Author contributions J.A. designed the project, designed and carried out the numerical experiments, and wrote the manuscript. C.C.F. processed the geomagnetic data, constructed the CHAOS-6x5 geomagnetic field model and led its comparison with the simulation results. J.A. and C.C.F. processed the results and discussed the manuscript.

Author Information Correspondence and requests for materials should be addressed to J. Aubert (email: aubert@ipgp.fr). The authors declare no competing financial interests.

Figure 1 Signature of geomagnetic jerks at Earth's surface. **a,c,e**, Observatory geomagnetic data and output from the CHAOS-6x5 geomagnetic field model⁵ based on satellite and ground observations. **b,d,f**, Output from the Midpath numerical model (Methods and Supplementary Table 1). **a**, Time series of the downward geomagnetic secular variation (rate of change of the geomagnetic field, blue lines and crosses) from annual differences of revised monthly means at two selected observatories, Kourou and Tamanrasset (marked as green dots in **e**), plotted together with the CHAOS-6x5 output (black). **b**, Downward vertical secular variation time series in the Midpath numerical model, at two locations marked with green dots in **f**. **c,d**, Time series of the mean-squared secular acceleration E_{SA} (grey) and of jerk energy E_J (black, see Methods for definitions and Fig. 4a for a longer time series of E_J), showing how secular acceleration pulses relate to the strong acceleration changes that characterise jerks. As identified locally from secular variation time series and globally by peaks in E_J , vertical dashed lines in **a,c** mark geomagnetic jerks^{6–9} occurring near epochs 2007.5, 2011, 2014 and in **b,d** the synthetic jerk event used to define the simulation time origin. **e,f**, Hammer projections of the radial secular acceleration (orange is outwards) before and after a jerk event, showing patterns alternating in time over a large portion of Earth's surface.

Figure 2 Signature of geomagnetic jerks at Earth's core surface. **a,b**, Output from the CHAOS-6x5 geomagnetic field model⁵. **c-f**, Output from the Midpath numerical dynamo model. **a,c**, Time-longitude plots of the radial secular acceleration at the equator (orange is outwards), filtered at spherical harmonic degree 9. The horizontal dashed lines locate geomagnetic jerks epochs^{6–9} in **a** and the synthetic jerk time in **c**. The vertical lines respectively

477 locate the longitudes selected for analysis in ref. ⁵ for the real events, and in **f** for the synthetic
 478 event. **b,d**, Hammer projections of the radial secular acceleration (same spatial filtering as
 479 in **a,c**) before and after jerk events, showing localised patterns alternating in time (see also a
 480 numerical model temporal sequence in Supplementary Movie 1). Black curves again locate
 481 the longitudes selected for analysis. **e**, Miller map showing details of the azimuthal flow ac-
 482 celeration at the core surface (native model spatial resolution, blue is westwards) during the
 483 simulated jerk event (see also global map and temporal sequence in Supplementary Movie
 484 2). The black vertical lines locate the analysis longitude used in **f**. **f**, Temporal evolution of
 485 the core surface, equatorial azimuthal flow acceleration at the analysis longitude, showing the
 486 structure of the wave packet that causes the jerk.

487 **Figure 3 Hydromagnetic waves inside the core and magnetic field structure from the**
 488 **Midpath model.** **a**, Planform of the azimuthal flow acceleration (blue is westward) at time
 489 -3.22 yr before the jerk event, in a quarter of the equatorial plane between longitudes $0^\circ E$
 490 and $90^\circ E$. Also shown are the directions of the rotation vector Ω and the wave vector **k**. **b**,
 491 Meridional planform of azimuthal flow acceleration outside the axial cylinder tangent to Earth's
 492 core and at the analysis longitude located by a black line in **a**. See supplementary Movie 4 for
 493 the corresponding temporal sequences. **c**, closeup of equatorial azimuthal flow acceleration
 494 corresponding to the dashed box in **a**, showing the concentrated wave structures below the
 495 core-mantle boundary. **d**, Semi-transparent detail of **a** (see second dashed box), with a volu-
 496 metric rendering of the magnetic field lines (grey, thickness proportional to local magnetic field
 497 amplitude) that channel and focus the waves towards the core surface. **e**, Hammer projection

of the radial magnetic field at the core surface (native model spatial resolution, orange is out-wards) at time -3.22 yr. Arrows in **d,e** locate the core surface magnetic flux patches where the focused waves emerge.

Figure 4 Statistics of jerk recurrence time. **a**, Time series of jerk energy E_J (see Methods) in the 29% of path and Midpath models (Supplementary Table 1). The arrow locates the event at time 0 yr analysed in Figs. 1-3. **b**, Distribution of the average recurrence time of jerks reaching or exceeding a given energy, for models within the rapid rotation regime¹⁷ (solid lines). Also shown is an extrapolation (see Methods and Supplementary Fig. 2) of the recurrence time distribution for the Earth's core conditions (dashed line and light grey shaded uncertainty area). The dark grey line segment locates the output of CHAOS-6x5 as estimated from Fig. 1c (three jerks with $E_J \geq 80 \text{ nT}^2 \cdot \text{yr}^{-4}$ and two with $E_J \geq 90 \text{ nT}^2 \cdot \text{yr}^{-4}$ within 19 years).

Figure 1

Figure 2

Figure 3

Figure 4

Supplementary information for: Geomagnetic jerks and rapid hydromagnetic waves focusing at Earth's core surface

Julien Aubert¹ and Christopher C. Finlay²

¹*Institut de Physique du Globe de Paris, Sorbonne Paris Cité, Université Paris Diderot, CNRS UMR 7154 CNRS, F-75005 Paris, France.*

²*Division of Geomagnetism, National Space Institute, Technical University of Denmark, Elektrovej, 2800 Kgs. Lyngby, Denmark*

Contents:

Supplementary Table 1

Supplementary Figs. 1-4

Captions for Supplementary Movies 1-4

Label	Path parameter ϵ	Path position	$\frac{\Delta\sigma_m}{D\sigma_c}$	core surface buoyancy	τ_U (yr)	τ_A (yr)	$2\pi\tau_\Omega$ (yr)	Integration time (yr)
Midpath	10^{-2}	29%	10^{-4}	neutral	129	31.5	1.0	42 900
	$3.33 \cdot 10^{-3}$	36%	10^{-4}	neutral	126	24.0	0.6	34 600
	10^{-3}	43%	10^{-4}	neutral	123	18.2	0.3	24 900
	$3.33 \cdot 10^{-4}$	50%	10^{-4}	neutral	125	14.3	0.2	24 400
Midpath-I	$3.33 \cdot 10^{-4}$	50%	0	neutral	120	14.2	0.2	11 400
Midpath-H	$3.33 \cdot 10^{-4}$	50%	10^{-3}	neutral	128	14.6	0.2	11 300
Midpath-S	$3.33 \cdot 10^{-4}$	50%	10^{-4}	adverse	121	14.5	0.2	10 100
End of path	10^{-7}	100%			130	1.9	$3.2 \cdot 10^{-3}$	
Earth					≈ 140	≈ 2	$2.7 \cdot 10^{-3}$	

Supplementary Table 1: **Models along a parameter space path to Earth’s core.** Key parameters and corresponding dimensional time scale values for numerical models located along a parameter space path²⁵ towards Earth’s core conditions. See Methods for definitions and ref. ¹⁷ for complete parameter data. Dimensional time scales values are obtained from the dimensionless time scale ratios reported in ref. ¹⁷ and the magnetic diffusion time scale set to $\tau_\eta = 135\,000$ yr in this study. Also shown are the values, closely approaching Earth’s core estimates, obtained by extrapolating scaling laws determined along the path²⁵ to its end point.

Supplementary Figure 1: **Alfvén wave propagation in the upper outer core.** Time-cylindrical radius diagrams of the column-averaged azimuthal flow acceleration $\partial u_c / \partial t$ (blue is westwards, see Methods for definitions) evaluated at two analysis longitudes, 31.5°E (as in Fig. 3b) and 37.5°E. Similar to ref. ¹⁷, green and brown curves respectively represent the ray-tracing theoretical propagation tracks of hydromagnetic waves at the column-averaged Alfvén speed c_A , and of material advection at the column-averaged cylindrical radial fluid velocity V_s . The slanted black line on the left panel denotes upward propagation at a speed 10 km/yr.

Supplementary Figure 2: **Scaling of jerk energy along the parameter space path.** Evolution of jerk energy at recurrence times 10, 30 and 100 years, as extracted from Fig. 4b, with the path parameter ϵ (Methods and Supplementary Table 1).

Supplementary Figure 3: **Sensitivity of the jerk recurrence time distribution to physical conditions in the lower mantle and upper outer core.** Distribution of the average recurrence time of jerks reaching or exceeding a given energy (same as Fig. 4b), for the models Midpath-I and Midpath-H with variable lower mantle electrical conductance, and model Midpath-S with a stratified region in the upper outer core (Methods). The Midpath model result from Fig. 4b is also reproduced for reference.

Supplementary Figure 4: **Signature of simulated jerks in the length of the day.** **a**, Squared acceleration $(d^2\Gamma_M/dt^2)^2$ (see Methods for definitions) of the electromagnetic torque exerted on the mantle by the outer core, as a function of time in the vicinity of four jerk events of model Midpath-H. **b**, first time derivative $d(\text{LOD})/dt$ of the simulated length of the day, as a function of time during the same jerk events. Arrows locate the pulses in the torque accelerations, that correspond to rapid inflexions in the rate of change of the length-of-day.

Supplementary Movie 1: Hammer projection of the core surface radial secular geomagnetic acceleration (orange is outwards) from the Midpath model, filtered at spherical harmonic degree 9, in the vicinity of the jerk event occurring at time 0 yr.

Supplementary Movie 2: Hammer projection of the core surface azimuthal flow acceleration (blue is westwards) from the Midpath model, in the vicinity of the jerk event occurring at time 0 yr.

Supplementary Movie 3: Partial equatorial cut (left) and meridional cut outside the tangent cylinder (right) of the convective density anomaly (orange denotes lighter fluid) from the Midpath model in the vicinity of the jerk event occurring at time 0 yr. The meridional cut in the right panel is taken at the analysis longitude marked by a black line in the left panel.

Supplementary Movie 4: Partial equatorial cut (left) and meridional cut outside the tangent cylinder (right) of azimuthal flow acceleration (blue is westwards) from the Midpath model in the vicinity of the jerk event occurring at time 0 yr. The meridional cut in the right panel is taken at the analysis longitude marked by a black line in the left panel.