A fungal scent from the cheese

Kovács, Ákos T.

Published in:
Environmental Microbiology

Link to article, DOI:
[10.1111/1462-2920.15267](https://doi.org/10.1111/1462-2920.15267)

Publication date:
2020

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Kovács, Á. T. (2020). A fungal scent from the cheese. *Environmental Microbiology*, 22(11), 4524-4526.
<https://doi.org/10.1111/1462-2920.15267>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 *Highlight on Cosetta et al doi: 10.1111/1462-2920.15223*

2

3 **A fungal scent from the cheese**

4

5 Ákos T. Kovács*

6

7 Bacterial Interactions and Evolution Group, DTU Bioengineering, Technical University of Denmark,
8 2800 Kongens Lyngby, Denmark

9

10 * For correspondence. E-mail atkovacs@dtu.dk; Tel. (+45) 45252727

11

12 Running title: Fungal volatile affected bacterial growth

13 Keywords: bacteria-fungi interaction; cheese rind; microbiome; community assembly; volatiles

14

15

16 **Summary**

17 Assembly of microbial communities is shaped by various physical and chemical factors deriving
18 from their environment, including other microbes inhabiting the certain niche. In addition to direct
19 cell-cell contacts, primary and secondary metabolites impact the growth of microbial community
20 members. Metabolites might act as growth-promoting (e.g. cross-feeding), growth-inhibiting (e.g.
21 antimicrobials), or signaling molecules. In multi-species microbial assemblies, secreted metabolites
22 might influence specific members of the community, altering species abundances and therefore
23 the functioning of these microcosms. In the current issue, Cosetta and colleagues describe a
24 unique volatile metabolite-mediated cross-kingdom interaction that shapes the cheese rind
25 community assembly. The study paves the way of our understanding how fungus-produced
26 volatile compounds promote the growth of a certain bacterial genus, a principal connection
27 between community members of the cheese rind.

28

29

30 **Main text**

31

32 Since the first formulation of the microbiological tenet '*Everything is everywhere, but, the*
33 *environment selects*' by Baas Beeking motivated by Martinus Beijerinck's work (Beijerinck, 1913;
34 Baas-Becking, 1934; De Wit and Bouvier, 2006), microbial ecologists have been researching the
35 factors determining microbial community assemblies. Complex association of diverse microbes in a
36 certain niche is driven by numerous influences from the environment, physical, chemical and
37 biological factors including temperature, humidity, nutrient availability, and various metabolites
38 from other organisms. Microbes can inhibit the growth of each other by producing specific or
39 general antimicrobial compounds, they can gain by acquiring beneficial metabolites secreted by
40 surrounding companions, in specific cases resulting in complex cross-feeding networks (D'Souza *et*
41 *al.*, 2018). In addition, the secreted primary- and secondary-metabolites could convey signaling
42 function that affect development of other microbes (Pishchany and Kolter, 2020). To gain deeper
43 understanding how the 'environment selects', simplified microbial systems are inspected ranging
44 from two or very few members to complex communities.

45 Cheese rind provides a relatively simple, tractable system that repetitively assembles into defined
46 composition of communities dominated by key genera, therefore allows examining microbial
47 succession (Wolfe *et al.*, 2014). The cheese diversity, including smell and taste, around the world is
48 in principle defined by the microorganisms accompanying the fermentation process that that is
49 thought to be invested and used since for 10.000 years. Originating from both natural
50 'contaminating' and inoculated microorganisms, the cheese rind community is dominated by few
51 prokaryotic phyla (*Firmicutes*, *Actinobacteria*, *Proteobacteria* and *Bacteriodes*) and eukaryotes
52 (yeast and molds) that forms a dense, biofilm-like layer around the cheese. The composition of such

53 rind cheese community is dynamic during the fermentation process and defined by abiotic and
54 biotic factors. This exceptional experimental system offers a platform to study interaction among
55 fungi and bacteria. The intricate interplay between these two kingdoms of life has been recently
56 exploited to understand the influence of diverse metabolites in microbial community dynamics.
57 What determines the abundances of microbial genera during cheese rind maturation? How do
58 specific metabolites impact the growth of microbes during the cheese ripening?

59 Cosetta and colleagues (2020) answer these questions and reveal how volatile organic compounds
60 (VOCs) produced by widespread cheese rind fungi specifically stimulate the growth of specific
61 cheese rind bacteria (Fig. 1). Exploiting a novel 'microtiter-plate sandwich' system, in which volatile
62 producers and responders are physically separated in the wells of the bottom and top plates,
63 respectively, but having common airspace, the influence of VOCs produced by five different cheese
64 rind derived fungi was inspected on the growth of bacterial isolates inoculated on cheese curd agar.
65 Intriguingly, the growth of various *Vibrio* species was generally promoted by the VOCs produced by
66 cheese rind fungi, including *Debaryomyces*, *Penicillium*, and *Galactomyces* isolates, but not when
67 bacteria and fungi were co-cultured. The promotion of *Vibrio* growth was also observed when a
68 synthetic bacterial community, representing a typical rind bacteria composition, was exposed to the
69 VOCs of the fungi. Characterization of the transcriptional profile of *Vibrio* population revealed that
70 VOCs promote the expression of genes involved in glyoxylate shunt, which allows bacteria to utilize
71 acetate and other fatty acids as carbon source. Indeed, esters and fatty acids, like acetic acid, are
72 widely detected in ripening cheeses, contributing to the sweet, fruity perception of surface-ripened
73 cheeses (Urbach, 1997). In line with this general observation, acetate-containing fatty acids could
74 be detected using gas chromatography/mass spectrometry in the headspace of cheese rind fungi
75 (Cosetta *et al.*, 2020). Finally, acetic acid, as well as heptyl acetate and methyl isobutyrate

specifically increased *Vibrio* growth and shifted the abundance of *Vibrio* within the synthetic microbial community.

The work of Cosetta and colleagues provides a valuable insight on VOC-mediated microbial interaction. Moreover, it reveals how growth promotion of specific genus shifts the composition of a synthetic microbial consortium and pinpoints the direct stimulation of growth by acetate-containing VOCs through metabolism in the glyoxylate shunt. VOCs have diverse impact on microbial growth and development, acting as nutrients, environmental modulators or signal molecules (Tyc *et al.*, 2017). Acetate-containing VOCs have been detected in numerous microbial systems (Ryu *et al.*, 2020), and while acetate acts as a metabolic signal inside bacterial cells (Wolfe, 2005), among microbial cells, it can also act as a signaling compound, altering development and differentiation, for example stimulating biofilm formation (Chen *et al.*, 2015). While the broad impact of this small volatile molecule has been previously suggested, Cosetta and colleagues have uniquely demonstrated the impact of microbial community composition.

The creation of synthetic microbial communities provides a tractable system to dissect the influence of various VOCs or generally of secondary metabolites to reveal the true nature of these compounds in nature. The activity of secondary metabolites has been traditionally explored in pairwise interactions and only recently, the influence of these specialized metabolites examined on microbial communities (Kiesewalter *et al.*, 2020). Once thought to act as fierce antimicrobial compounds, the real ecological impact of secondary metabolites within complex microbial communities still to be explored. The scent of the cheese is not only appreciated by its consumers, but also by the microbial community during the anciently acquired food fermentation process.

97 **Acknowledgements**

98 ÁTK was supported by the Danish National Research Foundation (DNRF137) for the Center for
99 Microbial Secondary Metabolites (CeMiSt).

100

101 **Competing Interests**

102 The author has no competing interests to declare.

103

104 **References**

- 105 Baas-Becking, L.G.M. (1934) *Geobiologie of inleiding tot de milieukunde*. The Hague, the
106 Netherlands: W.P. Van Stockum & Zoon (in Dutch).
- 107 Beijerinck, M.W. (1913) *De infusies en de ontdekking der backteriën. Jaarboek van de Koninklijke*
108 *Akademie van Wetenschappen*. Amsterdam, the Netherlands: Müller.
- 109 Chen, Y., Gozz, K., Yan, F., and Chai, Y. (2015) Acetic acid acts as a volatile signal to stimulate
110 bacterial biofilm formation. *mBio* **6**: e00392.
- 111 Cosetta, C.M., Kfoury, N., Robbat, A., and Wolfe (2020) Fungal volatiles mediate cheese rind
112 microbiome assembly. *Environ Microbiol* doi: 10.1111/1462-2920.15223.
- 113 D’Souza, G., Shitut, S., Preussger, D., Yousif, G., Waschina, S., and Kost, C. (2018) Ecology and
114 evolution of metabolic cross-feeding interactions in bacteria. *Nat Prod Rep* **35**: 455–488.
- 115 Kiesewalter, H.T., Lozano-Andrade, C.N., Strube, M.L., and Kovács, Á.T. (2020) Secondary
116 metabolites of *Bacillus subtilis* impact soil-derived semi-synthetic bacterial community assembly.
117 *bioRxiv* doi: 10.1101/2020.08.20.259788.
- 118 Pishchany, G., and Kolter, R. (2020) On the possible ecological roles of antimicrobials. *Mol*
119 *Microbiol* **113**: 580–587.
- 120 Ryu, C.M., Weisskopf, L., and Piechulla, B. (eds) (2020) *Bacterial Volatile Compounds as Mediators*
121 *of Airborne Interactions*. Springer Nature Singapore Pte Ltd.
- 122 Tyc, O., Song, C., Dickschat, J.S., Vos, M., and Garbeva, P. (2017) The ecological role of volatile and
123 soluble secondary metabolites produced by soil bacteria. *Trends Microbiol* **25**: 280–292.
- 124 Urbach, G. (1997) The flavour of milk and dairy products: II. Cheese: contribution of volatile
125 compounds. *Int J Dairy Technol* **50**: 79–89.
- 126 de Wit, R., and Bouvier, T. (2006) “Everything is everywhere, but, the environment selects”; what

127 did Baas Becking and Beijerinck really say? *Environ Microbiol* **8**: 755–758.

128 Wolfe, A.J. (2005) The acetate Switch. *Microbiol Mol Biol Rev* **69**: 12–50.

129 Wolfe, B.E., Button, J.E., Santarelli, M., and Dutton, R.J. (2014) Cheese rind communities provide

130 tractable systems for in situ and in vitro studies of microbial diversity. *Cell* **158**: 422–433.

131

132

133

134

135 Fig. 1 VOCs (heptyl acetate and methyl isobutyrate) mediated interaction between fungi (left inlet)

136 and bacteria (*Vibrio*, right inlet). Inlets were prepared on BioRender.com