



Light-controlled out-of-equilibrium assembly of cyclodextrins in an enzyme-mediated dynamic system

Larsen, Dennis; Bjerre, Philip M.; Beeren, Sophie R.

Published in:
Chemical Communications

Link to article, DOI:
[10.1039/c9cc08452e](https://doi.org/10.1039/c9cc08452e)

Publication date:
2019

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Larsen, D., Bjerre, P. M., & Beeren, S. R. (2019). Light-controlled out-of-equilibrium assembly of cyclodextrins in an enzyme-mediated dynamic system. *Chemical Communications*, 55(100), 15037-15040. <https://doi.org/10.1039/c9cc08452e>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

COMMUNICATION

Light-controlled out-of-equilibrium assembly of cyclodextrins in an enzyme-mediated dynamic system

Received 00th January 20xx,
Accepted 00th January 20xx

Dennis Larsen, Philip M. Bjerre and Sophie R. Beeren*

DOI: 10.1039/x0xx00000x

We show that the selective enzymatic synthesis of specific cyclodextrins can be modulated using light. We use enzyme-mediated dynamic combinatorial chemistry to generate a mixture of interconverting linear and cyclic α -1,4-glucans, and employ an azobenzene photoswitch as a template. Using UV or blue light to switch between photostationary states with different azobenzene *cis/trans* isomeric ratios, we can promote the out-of-equilibrium assembly of either α -cyclodextrin or β -cyclodextrin.

Light-responsive systems in living organisms exploit photoswitchable co-factors to trigger the onset of enzyme-catalysed reaction cascades to modulate growth and control essential functions. For example, phototropic plants optimize their growth in response to sunlight via the judicious synthesis of specific poly- and oligosaccharides,¹ while the *cis-trans* photoisomerisation of retinal is responsible for visual transduction.² Inspired by Nature, chemists have developed artificial light-responsive systems wherein a photo-activated conformation change in a small molecule or molecular moiety leads to a specific output from a chemical network, or altered physical properties of a self-assembled material.³

A number of artificial systems have recently been reported in which enzyme activity is controlled using light, either by direct bioconjugation of enzymes with photo-responsive moieties, or by photoisomerisation-based activation of a co-factor or deactivation of an inhibitor.⁴ More complex systems use light-controlled self-assembly of supramolecular structures for compartmentalization or release of co-factors or substrates, thus stimulating enzyme activity.⁵ Here we apply a unique strategy to control the outcome of an enzymatic process using light, which relies upon the use of a photo-responsive template to select different products in an enzyme-mediated dynamic system. Specifically, we control the preferential enzyme-

catalysed synthesis of α - or β -cyclodextrin via the photoisomerisation of an azobenzene template (Fig. 1).

Dynamic combinatorial chemistry is a now well-established




Figure 1 Light-controlled selective, enzyme-mediated synthesis of cyclodextrins. (a) CGTase acts on an α -1,4-glucan source to generate a dynamic combinatorial library of cyclodextrins. Different isomers of the azobenzene template promote the selective synthesis of different cyclodextrins. (b) Azobenzenes switch from the *trans*- to the *cis*-isomer upon irradiation with UV light and switch back thermally or with blue light.

methodology for the templated synthesis of complex molecular architectures, and the assembly of dynamic polymers and responsive systems from small building blocks under thermodynamic control.⁶ A few examples of dynamic combinatorial libraries based on reversible organic reactions (e.g. disulfide and hydrazone exchange) that incorporate light-responsive building blocks⁷ and templates⁸ have been reported.

We recently described enzyme-mediated dynamic combinatorial chemistry, wherein enzyme-catalysed reactions are used to reversibly link molecular building blocks and generate mixtures of interconverting bio-oligomers.⁹ We employed cyclodextrin glucanotransferase (CGTase)¹⁰ to generate a dynamic mixture of linear α -1,4-glucans

Department of Chemistry, Technical University of Denmark, Kemitorvet 207, DK-2800 Kongens Lyngby, Denmark.


Electronic Supplementary Information (ESI) available: Experimental details, full list of photoswitches screened, chromatographic data, UV-vis spectra, NMR spectra. See DOI: 10.1039/x0xx00000x

(maltooligosaccharides) and cyclic α -1,4-glucans (cyclodextrins (CDs)). We outlined how CGTase catalyses the fast, reversible transglycosylation and slow, irreversible hydrolysis of α (1 \rightarrow 4)-glycosidic linkages, and thus generates a complex dynamic chemical system, wherein cyclodextrins assemble transiently out-of-equilibrium in a kinetically trapped subsystem. This subsystem operates under *pseudo*-thermodynamic control and by adding different templates, we were able to alter the distribution of the different cyclodextrins formed to obtain either α -CD, β -CD or γ -CD, with high selectivities ranging from 89% to 99%.⁹

Azobenzene moieties have been extensively utilised to achieve photo-responsive behaviour in functional molecular and biomolecular systems.¹¹ Irradiation with UV light (typically 340–380 nm) causes a photo-induced isomerisation from the most stable *trans* isomer to the more sterically encumbered and less stable *cis* isomer (Fig. 1b).¹² Return to the *trans* isomer occurs via a thermal back-reaction, or can be promoted by irradiation with visible light (typically 430–550 nm). Sparsely substituted azobenzenes are known to bind within the hydrophobic cavities of cyclodextrins (in aqueous solution) and the selective recognition of the *cis* or *trans* isomers has been exploited to control self-assembly of functional materials.¹³ For this study, we screened a number of different substituted azobenzenes (ESI Table S1) and identified a set of water soluble azobenzene photoswitches (**1**–**5**) exhibiting markedly different *cis/trans* isomeric ratios in ambient light and when irradiated with UV light at 365 nm (Table 1 and ESI Figs S2 and S3).

To investigate the possibility of modulating the selective synthesis of different cyclodextrins using light, we examined the action of CGTase on α -CD (2 mg/ml in 50 mM sodium phosphate buffer at pH 7.5) in the presence of azobenzenes **1**–**5** (2 mM), in the dark, or with UV-irradiation (365 nm, for 1 hour prior to addition of enzyme and continuously throughout the reaction). The composition of each library was analysed after 2 hours, using high performance liquid chromatography with evaporative light-scattering detection (HPLC-ELS), and compared with an untemplated reaction (Fig. 2 and ESI Fig. S4).

Table 1 Physicochemical properties of azobenzenes **1**–**5**^a



	Photoswitch		λ_{\max} (nm)		<i>cis/trans</i> ratio ^b	
	R	R'	<i>trans</i>	<i>cis</i>	No irr	UV
1	COOH	H	326	426	15:85	73:27
2	COOH	COOH	331	429	10:90	51:49
3	SO ₃ Na	OAc	325	426	12:88	63:27
4	SO ₃ Na	NHAc	354	420	1:99	74:26
5			320	423	18:82	70:30

^a At room temperature in 50 mM sodium phosphate buffer at pH 7.5. ^b By ¹H NMR spectroscopy when kept in ambient light (no irr.) or after irradiation with light (365 nm) for 1 hour (UV).




Figure 2 Screening of azobenzenes **1**–**5** as light-responsive templates for preferential synthesis of different cyclodextrins in a CGTase-mediated dynamic combinatorial library. (a) Representative chromatograms (HPLC-ELS) depicting the relative concentrations of α -, β -, and γ -CD formed when CGTase acts on α -CD in the absence of template, and with **1** (2 mM) in the dark or under UV-irradiation (365 nm). (b) Bar graph depicting the cyclodextrin distributions produced in similar reactions templated with azobenzenes **1**–**5**. Reaction time: 2 hours. Conditions: 2 mg/mL α -CD used as starting α -glucan in 50 mM sodium phosphate buffer at pH 7.5.

Pleasingly, in all cases, we observed different distributions of cyclodextrin products depending on whether or not the reaction was irradiated with UV-light. HPLC chromatograms showing the differing distributions of cyclodextrins that are formed in the absence of any template and in the presence of photoswitch **1**, in the dark and with UV irradiation, are shown in Fig. 2a. In each of these reactions, α -CD and β -CD were the major products (very little γ -CD is produced at low concentrations of α -1,4-glucan (2 mg/mL)). A clear increase in the production of α -CD and concurrent decrease in the production of β -CD was apparent upon addition of template **1** in the dark. Conversely, an increase in the production of β -CD was observed upon application of UV-light together with the template. In the absence of template, UV-irradiation did not influence the distribution of products.

For each of the azobenzenes **1**–**5**, addition of the template without irradiation generated higher concentrations of α -CD, indicating that in each case the *trans* isomer binds preferentially to α -CD (Fig. 2b). For photoswitches **2**–**5**, irradiation led to a reduction in the amplification of α -CD suggesting that either the *cis*-isomers have little interaction with either of the cyclodextrins, or there is a favourable interaction between the *cis*-isomers and β -CD, but this is outweighed by the interaction between α -CD and any remaining *trans*-isomer (note that the




Figure 3. Light-controlled changes in CD distribution in CGTase-mediated dynamic cyclodextrin systems generated starting from α -CD (2 mg/mL) and templated with photoswitch **1** (2 mM). a) Dynamic system kept in the dark before exposure to UV light (365 nm). b) Dynamic system started under UV light irradiation (365 nm), then irradiated with blue light (460 nm). c) Consecutive cycles of irradiation with UV light (365 nm, 20 minutes) and blue light (460 nm, 15 minutes).

photostationary state obtained under UV irradiation contains significant amounts of the *trans* isomers for all of the azobenzene templates, Table 1). Only photoswitch **1** caused the amplification of β -CD when irradiated, indicating that *cis*-**1** binds preferentially to β -CD, while *trans*-**1** binds more strongly to α -CD.

Determination of association constants for the binding of a guest that exists as a mixture of slowly interconverting isomers is not straightforward, as the isomeric ratio is altered by selective interaction with the host. As photoisomerization of azobenzenes does not give complete *trans* to *cis* conversion, the *cis* isomer is always contaminated with a minority of the *trans* isomer even after photoirradiation. This means that simple titrations, and fitting to a 1-to-1 binding model, cannot be used to accurately determine association constants for binding of *cis*-isomers, due to competitive binding by the *trans* isomer minority. We have previously described how an NMR titration can be used to simultaneously determine association constants for the binding of a host to a number of different guests present in a mixture.¹⁴ This methodology does not require the concentrations of the host and guests to be known, nor should one be kept constant. It requires only that the association constant (K_a) is known for one of the guests. Here we show that the same methodology can be used to determine the association constants for *cis*-**1** and *trans*-**1** binding to α -CD and β -CD. Solutions of photoswitch **1** (1 mM in 50 mM sodium phosphate buffer at pH 7.5) were titrated with α -CD and β -CD, with each titration performed under two different sets of conditions: (1) in the dark and after heating at 80 °C for 2 days to obtain exclusively *trans*-**1**; (2) in ambient light, such that a mixture of *cis*-**1** and *trans*-**1** were present. For the titrations performed in the dark, association constants, $K_{a(\text{trans})}$ (Table 2) were obtained by fitting to a regular 1:1 binding isotherm (ESI Fig. S4). For the titrations in ambient light, both *cis*-**1** and *trans*-**1** were present, and these interacted unequally with the cyclodextrins, leading to a change in the ratio of isomers present upon increasing the concentration of cyclodextrin. By plotting the changes in chemical shift observed for each isomer upon addition of cyclodextrin against one another ($\Delta\delta_{\text{cis}}$ vs. $\Delta\delta_{\text{trans}}$),

and fitting to the equation below, we could obtain $K_{a(\text{cis})}$ for each cyclodextrin (Table 2 and ESI Fig. S5).

$$\Delta\delta_{\text{cis}} = \frac{\Delta\delta_{\text{max}(\text{cis})}\Delta\delta_{\text{trans}}K_{a(\text{cis})}}{\Delta\delta_{\text{max}(\text{trans})}K_{a(\text{trans})} - \Delta\delta_{\text{trans}}K_{a(\text{trans})} + \Delta\delta_{\text{trans}}K_{a(\text{cis})}}$$

Table 2. Association constants, K_a (M^{-1}), for α - and β -CD with *trans* and *cis* isomers of **1**^a

Host	Guest	
	<i>trans</i> - 1	<i>cis</i> - 1
α -CD	5000 \pm 400	140 \pm 70
β -CD	2500 \pm 300	2200 \pm 300

^a In sodium phosphate buffer (pH 7.5, 50 mM) in D₂O at 25 °C.

The relative affinities of *cis*-**1** and *trans*-**1** for α -CD and β -CD can thus explain the different product distributions obtained when CGTase catalyses the formation of CDs in the absence or presence of UV irradiation. When the enzymatic reaction is carried out in the dark, *trans*-**1** predominates, and since it binds α -CD approximately twice as strongly as β -CD, the production of α -CD is amplified. When exposed to UV light, *cis*-**1** predominates, and higher production of β -CD results, because *cis*-**1** binds 15 times more strongly to β -CD than to α -CD.

To illustrate how the dynamic enzymatic synthesis of cyclodextrins can be coupled to the dynamic switching between photostationary states of photoswitch **1**, the CGTase-catalysed interconversion of CDs was monitored over time under varying light conditions. Firstly, a reaction was started in the dark and kept in darkness for the first 80 minutes before being submitted to UV-irradiation (365 nm) (Fig. 3a). The system adapted quickly to the UV light, switching readily from 80:20 α -CD/ β -CD ratio in the dark to the expected 40:60 ratio when under UV irradiation. A second reaction was begun under UV irradiation, and after 60 minutes, the irradiation was switched to blue light (460 nm) (Fig. 3b). Irradiation with blue light returns the mainly *cis*-**1** to *trans*-**1**, and thus the α -CD/ β -CD ratio returned to approximately 80:20. Consecutive cycles of irradiation with UV-light (365 nm, 20 mins) and blue light (460 nm, 15 mins) were performed, demonstrating the possibility to oscillate up to six times between CD mixtures of predominantly α -CD or β -CD (Fig. 3c).

Due to the irreversible background hydrolysis of glycosidic linkages, and the gradual accumulation of glucose (which cannot act as a glycosyl donor in CGTase-mediated glycosyl transfer), a steady decrease in the total cyclodextrin concentration is observed (Fig. 3 grey line and ESI Fig. S6-9). This explains the gradual rise in the α -CD concentration and concomitant descent in the β -CD concentration, as at lower building block concentrations, shorter oligomers are favoured in dynamic combinatorial libraries.¹⁵ Background hydrolysis appears faster during UV-irradiation but this is presumably due to the overall weaker binding by both cyclodextrins to *cis*-1 compared to *trans*-1, and as we have previously shown, hydrolysis of cyclodextrins is hindered by complexation.⁹

In our enzyme-mediated dynamic system, cyclodextrins are not formed under equilibrium conditions, but instead self-assemble transiently out-of-equilibrium and are slowly converted to glucose, the thermodynamic product.⁹ Surprisingly, templates can nevertheless be exploited to select for specific cyclodextrin products, presumably because α -, β -, and γ -CD are separated only by low energy barriers and form a kinetically-trapped subsystem operating under *pseudo*-thermodynamic control. This azobenzene-templated system is doubly dynamic, as the equilibria for both the *cis/trans* isomerization and the CGTase-catalysed α -CD/ β -CD interconversion influence one another. The unconventional energy landscape for this synthetic dynamic system is depicted in Figure 4. Here we show that we can use UV light to push the system further out-of-equilibrium and transiently occupy an alternative, higher energy dissipative, but also kinetically trapped state.¹⁶ Using different wavelengths of light to switch between predominantly *trans*-1 or *cis*-1, we can control which kinetically trapped regime is occupied, and promote the selective formation of α -CD or β -CD, respectively.

Conclusions

In conclusion, we have demonstrated the use of light to artificially control an enzyme-driven process, not by activation of a substrate or co-factor, but by exploiting the molecular recognition of a light-responsive template to select specific products in an enzyme-mediated dynamic system.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- 1 E. Liscum, S. K. Askinosie, D. L. Leuchtman, J. Morrow, K. T. Willenburg and D. R. Coats, *Plant Cell*, 2015, **26**, 38-55.
- 2 K. Palczewski, *J. Biol. Chem.*, 2012, **287**, 1612-1619.
- 3 (a) Z. L. Pianowski, *Chem. Eur. J.*, 2019, **25**, 5128-5144; (b) A. K. Mandal, M. Gangopadhyay and A. Das, *Chem. Soc. Rev.*, 2015, **44**, 663-676; (c) J.-F. Gohy and Y. Zhao, *Chem. Soc. Rev.*, 2013, **42**, 7117-7129; (d) C. Zhu, C. Ninh and C. J. Bettinger, *Biomacromol.*, 2014, **15**, 3474-3494.
- 4 (a) T. Tian, Y. Song, J. Wang, B. Fu, Z. He, X. Xu, A. Li, X. Zhou, S. Wang and X. Zhou, *J. Am. Chem. Soc.*, 2016, **138**, 995-961;
- 5 (b) A. Ali, G. A. Bullen, B. Cross, T. R. Dafforn, H. A. Little, J. Manchester, A. F. A. Peacock and J. H. R. Tucker, *Chem. Commun.*, 2019, **55**, 5627-5630; (c) K. Aggarwal, M. Banik, B. Medellin and E. L. Que, *Biochemistry*, 2019, **58**, 48-53; (d) R. Mogaki, K. Okuro and T. Aida, *J. Am. Chem. Soc.*, 2017, **139**, 10072-10078; (e) W. Szymański, J. M. Beierle, H. A. V. Kistemaker, W. A. Velema and B. L. Feringa, *Chem. Rev.*, 2013, **8**, 6114-6178.
- 6 (a) P. Frei, R. Hevey and B. Ernst, *Chem. Eur. J.*, 2019, **25**, 61-73; (b) F. B. L. Cougnon and J. K. M. Sanders, *Acc. Chem. Res.*, 2012, **45**, 2211-2221.
- 7 (a) M. Kathan, F. Eisenreich, C. Jurissek, A. Dallmann, J. Gurke and S. Hecht, *Nat. Chem.*, 2018, **10**, 1031-1036; (b) L. A. Ingeman and M. L. Waters, *J. Org. Chem.*, 2009, **74**, 111-117; (c) S. Wang, L. Yue, Z.-Y. Li, J. Zhang, H. Tian and I. Willner, *Angew. Chem. Int. Ed.*, 2018, **57**, 8105-8109.
- 8 F. Ulatowski, K. Dabrowa and J. Jurczak, *Tet. Lett.*, 2016, **57**, 1820-1824.
- 9 D. Larsen, S. R. Beeren, *Chem. Sci.* **2019**, DOI: 10.1039/c9sc03983j.
- 10 (a) Y. B. Tewari, R. N. Goldberg and M. Sato, *Carbohydr. Res.*, 1997, **301**, 11-22; (b) J. C. M. Uitdehaag, B. A. van der Veen, L. Dijkhuizen and B. W. Dijkstra, *Enzyme Microb. Technol.*, 2002, **30**, 295-304; (c) J. C. M. Uitdehaag, K. H. Kalk, B. A. van der Veen, L. Dijkhuizen and B. W. Dijkstra, *J. Biol. Chem.*, 1999, **274**, 34868-34876.
- 11 (a) M. Zhu and H. Zhou, *Org. Biomol. Chem.*, 2018, **16**, 8434-8445. (b) A. A. Beharry and G. A. Wooley, *Chem. Soc. Rev.*, 2011, **40**, 4422-4437.
- 12 S. Crespi, N. A. Simeth and B. König, *Nat. Chem. Rev.*, 2019, **3**, 133-146.
- 13 D. Wang, W. Zhai, Q. Wei, C. Zhao and Y. Zheng, *ChemPhotoChem*, 2018, **2**, 403-415.
- 14 S. Meier and S. R. Beeren, *J. Am. Chem. Soc.*, 2014, **136**, 11284-11287.
- 15 (a) P. T. Corbett, S. Otto and J. K. M. Sanders, *Chem. Eur. J.*, 2004, **10**, 3139-3143; (b) P. T. Corbett, J. K. M. Sanders and S. Otto, *J. Am. Chem. Soc.*, 2005, **127**, 9390-9392.
- 16 M. Kathan, S. Hecht, *Chem. Soc. Rev.*, 2017, **46**, 5536-5550.




Figure 4 Schematic energy diagram summarizing the different energy levels populated in this photodynamic enzyme-mediated dynamic system. Photoactivation with UV or blue light shifts the cyclodextrin subsystem from one kinetically trapped minimum to another.