Interfacing MCNPX and McStas for simulation of neutron transport

Klinkby, Esben Bryndt; Lauritzen, Bent; Nonbøl, Erik; Willendrup, Peter Kjær; Filges, Uwe; Wohlmuther, Michael ; Gallmeier, Franz X.

Published in:

Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment

Link to article, DOI:

[10.1016/j.nima.2012.10.052](https://doi.org/10.1016/j.nima.2012.10.052)

Publication date:

2013

[Link back to DTU Orbit](#)

Citation (APA):

Klinkby, E. B., Lauritzen, B., Nonbøl, E., Willendrup, P. K., Filges, U., Wohlmuther, M., & Gallmeier, F. X. (2013). Interfacing MCNPX and McStas for simulation of neutron transport. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 700, 106-110. <https://doi.org/10.1016/j.nima.2012.10.052>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Interfacing MCNPX and McStas for simulation of neutron transport

Esben Klinkby^{*1,3}, Bent Lauritzen^{1,3}, Erik Nonbøl^{1,3}, Peter Kjær
Willendrup^{2,3}, Uwe Filges^{4,5}, Michael Wohlmuther^{4,5}, Franz X. Gallmeier⁶

1) DTU Nutech, Technical University of Denmark, DTU Risø Campus,
Frederiksborgvej 399, DK-4000 Roskilde

2) DTU Physics, Technical University of Denmark, DTU Lyngby Campus, Anker
Engelunds Vej 1, DK-2800 Kgs. Lyngby

3) ESS design update programme - Denmark

4) Paul Scherrer Institute, CH-5232 Villigen PSI, Switzerland

5) ESS design update programme - Switzerland

6) Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA

1. Abstract

Simulations of target-moderator-reflector system at spallation sources are conventionally carried out using Monte Carlo codes such as MCNPX[1] or FLUKA[2, 3] whereas simulations of neutron transport from the moderator and the instrument response are performed by neutron ray tracing codes such as McStas[4, 5, 6, 7]. The coupling between the two simulation suites typically consists of providing analytical fits of MCNPX neutron spectra to McStas. This method is generally successful but has limitations, as it e.g. does not allow for re-entry of neutrons into the MCNPX regime. Previous work to resolve such shortcomings includes the introduction of McStas inspired supermirrors in MCNPX. In the present paper different approaches to interface MCNPX and McStas are presented and applied to a simple test case. The direct coupling between MCNPX and McStas allows for more accurate simulations of e.g. complex moderator geometries, backgrounds, interference between beam-lines as well as shielding requirements along the neutron guides.

Keywords: Neutron, Transport, Simulation, MCNPX, McStas, Interface

^{*}Corresponding author

2. Introduction

In the target-moderator-reflector system of a spallation source, neutrons are slowed down from being fast at the formation in the spallation target to thermal or cold in the beam extraction guides.

To model the interaction of a proton beam with a spallation target and to model the thermalization of the produced neutrons in moderators, the MCNPX code is a standard of its field [1]. Since mainly being developed for applications involving fast or thermal neutrons, however, the MCNPX code does lack in description of coherent scattering applicable to the cold/thermal range.

The transport of cold/thermal neutron through guides and optics and the simulation of scattering instruments on the other hand are well described using neutron ray-tracing codes such as McStas [4, 5, 6, 7]. To bridge the gap between MCNPX and McStas, the approach has generally been to use analytical distributions fitted to MCNPX event spectra as input for the McStas simulation. This decoupled approach causes phase space information to be lost, and is limited by the fact that it does not allow the re-entry of McStas-simulated cold neutrons into the MCNPX regime. To estimate shielding requirements along a neutron guide or to calculate the gamma backgrounds relevant at neutron scattering instruments, it is necessary to apply MCNPX in calculating neutron absorption and gamma production¹.

In order to resolve this issue, a more direct coupling between MCNPX and McStas is required. Below, various possibilities for such MCNPX-McStas coupling are described. Based on experience gained during implementation and tests of the interfaces, the feasibility and usefulness of the individual approaches are evaluated.

While the present paper is a pure computational study describing the various interfaces between MCNPX and McStas, experiments at the BOA beam-line² are planned to validate against real measurements.

3. Concepts for automated interfacing of MCNPX and McStas

3.1. Tally option - the present default

This approach is based on fitting MCNPX neutron distributions e.g. at the moderator surface, allowing to model neutron states on a statistical basis. In short, a detailed MCNPX simulation of a target, reflector and moderator system of a given neutron facility is performed and the resulting neutron fluxes and energy spectra at the moderator surface are approximated by several Maxwellians. McStas then sample random neutron states from these distributions. A challenge faced when using this approach is to correctly describe the correlations between the parameters constituting a neutron state. For example, non-trivial

¹Normally in a McStas guide simulation un-reflected neutrons are simply discarded.

²One of the beam-lines at the SINQ spallation source at the Paul Scherrer Institute (PSI), Switzerland.

1 phase space correlations could exist e.g. between the neutron location and mo-
2 mentum at the moderator surface. Quantifying correlations is thus an important
3 part of employing the *Tally* method.

4 The advantage of the *Tally* method, as seen from a user perspective is, that
5 the time consuming MCNPX simulation step is decoupled from McStas and can
6 be carried out once-and-for-all. This makes subsequent McStas simulations fast
7 and therefore this method is very useful for e.g. instrument design.

8 3.2. *Ptrac* option

9 This approach utilises an intermediate step of event files, so that MCNPX
10 at a given user-defined surface writes a file containing the state of the individual
11 neutrons (position, momentum, time and Monte Carlo weight). An appropriate
12 McStas interface exists to read in the neutron events. An advantage of using the
13 *Ptrac* option with respect to the *Tally* approach is that all correlations between
14 neutron state parameters are automatically conserved. Apart from the sizable
15 intermediate files, a drawback by this approach is that the MCNPX code is
16 unable to re-import data in the *Ptrac* format. I.e., this approach can only be a
17 one way interface. Moreover the method is limited by the fact that MCNPX only
18 allows particles crossing *one* surface to be written to file, and that the *Ptrac*
19 option is unavailable under MPI³. For these reasons, relying on intermediate
20 *Ptrac* files is inadequate as a general solution to the problem faced.

21 3.3. *Source Surface Write/Read (SSW/SSR)* option

22 *SSW/SSR* is an MCNPX feature that allows to stop a simulation at a given
23 surface, and restart it later. It is not intended to be used as a switch for external
24 programs to be linked with an MCNPX simulation and the intermediate data
25 files have undocumented MCNPX version dependent binary formats. A new
26 interpreter has been developed, allowing McStas to run based on a *SSW/SSR*
27 file input, and to produce a *SSW/SSR* output once the McStas simulation is
28 complete. The main advantage of this approach compared to the *Ptrac* option
29 is that MCNPX can run based on the *SSW/SSR* input files. In this way one can
30 first perform an MCNPX simulation of e.g. thermal neutron moderation. Once
31 the neutrons enter the beam extraction region the neutron states are handed
32 to the *SSW/SSR* interface, and based on this a McStas simulation is carried
33 out, e.g. involving mirrors and coherent scattering (which is not possible in
34 MCNPX). The scattered and/or the non-scattered neutrons can then be handed
35 back to MCNPX using the same interface.

36 The corresponding McStas components to read/write from/to the *SSW/SSR*
37 format are called: *Virtual_MCNP_ss_input* and *Virtual_MCNP_ss_output*, and
38 are included as official McStas components starting from McStas version 2.0.

³MPI: Message Passing Interface, which is a method of parallelising computer processing.
For additional information, see: <http://www.mcs.anl.gov/research/projects/mpi/>.

3.4. Compile option

Closely resembling the above approach, the *Compile* option represents an even more direct coupling of the MCNPX and McStas codes. Rather than writing out intermediate files using the *SSW/SSR* interface, the McStas and MCNPX codes are compiled together so that once a neutron arrives at a *McStas surface*, a McStas simulation is launched from within MCNPX given a neutron state as input. After completion the updated neutron state is returned to MCNPX which proceeds the simulation. For an illustration see figure 1. The *Compile* method is very flexible since all McStas functionalities are available from within MCNPX, but there are drawbacks: Firstly, the above relies on changes to the MCNPX code. Changes that would need to be repeated, if one would want to upgrade to later versions of MCNPX⁴. Secondly, there is a licensing issue at hand when merging the codes: McStas is licensed under GNU GPL v2⁵, whereas MCNPX requires individual personal certification, something which many potential users may not be able to obtain. Thirdly, given the substantial difference in the CPU time consumption between typical MCNPX and McStas simulations (several orders of magnitude), there is an advantage in being able to separate them. If not, those e.g. designing/simulating neutron experiments at the end of the beam-line, would have to cope with very long simulation times since a full MCNPX simulation would have to be launched for each neutron. In many cases the lengthy simulations could be avoided with insignificant loss of precision if the McStas simulations were bootstrapped using the *Tally*, *SSW/SSR* or *Ptrac* interface.

Figure 1: The McStas surface interface in MCNPX - illustration of coherent scattering from atomic lattice.

3.5. Supermirror option

Similar to the *Compile* option, the *Supermirror* option is based on modifying the MCNPX source code [8, 9]. In this case, however, the idea is not to launch a McStas simulation from within MCNP, but rather to extend MCNPX, with

⁴Presently implemented in a development version of MCNPX 2.7. The cross release maintenance could be avoided if the changes were ported into the MCNPX development branch - this will be attempted in the future.

⁵A special internal DTU and ESS project license for this usage was applied.

1 functionality inspired from McStas. The first and most important shortcom-
 2 ing when using MCNPX for cold neutron applications is the lack of coherent
 3 scattering. Coherent scattering can be described as a neutron wave interacting
 4 with a lattice, while MCNPX only considers scattering on single particles. The
 5 process gives rise to wavelength-dependent reflection and can for the present
 6 purposes be well-approximated by the following expression [7]:

$$\begin{aligned}
 R &= \frac{R_0}{2} \left(1 - \tanh \frac{Q - m \cdot Q_c}{W} \right) \times (1 - \alpha(Q - Q_c)) & \text{for } Q > Q_c \\
 R &= R_0 & \text{otherwise}
 \end{aligned} \quad (1)$$

7 where Q is the scattering vector, Q_c is the critical scattering vector, R_0 is
 8 the low angle reflectivity constant, W is the width of supermirror cut-off, α is
 9 the reflectivity slope, and m is the m -value of the material. The wavelength-
 10 dependent reflectivity is depicted in figure 2.

11 As for the *Compile* option, maintenance across MCNPX releases is prob-
 12 lematic for the supermirror approach. Also McStas functionality other than su-
 13 permirrors may need to be implemented, potentially requiring significant code
 14 development.

Figure 2: Supermirror reflectivity as a function of the scattering vector Q . The parameters used correspond to the McStas defaults: $Q_c = 0.0219$, $m = 2$, $W = 0.003$, $R_0 = 0.99$, $\alpha = 6.07$.

15 4. Interface validation results

16 In order to validate the performance of the interfaces presented above we con-
 17 sider a test scenario consisting of a source plane, that emits 10^6 20 meV neutrons
 18 at a 45° angle toward a mirror (i.e. a neutron guide), which then reflects the
 19 neutrons to the end wall for detection (surface current tally in MCNPX). The
 20 geometry is shown in figure 3 along with an example neutron trajectory. Focus
 21 below is put on the three interfaces presented here for the first time: *SSW/SSR*,
 22 *Compile* and *Supermirror*.

23 The *SSW/SSR* and *Compile* approaches are developed in MCNPX release
 24 2.7 and to allow for direct comparison the existing *Supermirror* approach [8]
 25 was ported to the same release.

Figure 3: Geometry and example neutron simulation of the test setup used for MCNP-McStas interface validation. Illustrated using *Vised*[10].

Figure 4: Example event of the SSW/SSR approach as visualized by McStas' *mcdisplay* and MCNPX's *Vised*[10].

Figure 5: McStas PSD at the guide entrance (left) and exit (right).

Figure 6: Spectrum and angular distributions at the guide exit using the MCNPX mirror. As expected both angle and energy are conserved.

1 We consider the MCNPX perfect specular mirror. Rather than choosing
2 realistic settings for the guide (super)mirror, we choose the parameters of the
3 guide such that nearly total reflection is expected, hereby enabling direct com-
4 parison against the MCNPX specular mirror. By also setting mass densities to
5 zero, any bias due to lack of material effects in McStas is suppressed and thus,
6 all methods should give identical neutron yield at the far end of the guide (apart
7 from statistical fluctuations). To achieve this we set the following supermirror
8 parameters: $R_0 = 0.99$, $Q_c = 20$, $m = 2$, $W = 0.003$, $\alpha = 6.07$ (see Eq. 1) - note
9 that Q_c is a factor ~ 1000 above that of nickel (typical supermirror coating) to
10 ensure maximum reflection, even at an angle of 45° .

11 To picture the process consider the *SSW/SSR* interface. Immediately after
12 the generation at the guide entrance, the neutrons are exported to McStas
13 through the *SSW/SSR* interface. Using the visualisation capabilities of McStas
14 (*mcdisplay*) an example neutron is traced in figure 4(left). At the guide exit
15 ($z = 0.04\text{m}$) it is returned to MCNPX and its final path through the tally surface
16 is visualised using *Vised*[10] in figure 4(right). In figure 5 McStas Position
17 Sensitive Detectors (PSD) placed at the guide entrance and exit of the guide
18 show close to identical distributions, as expected given that the guide parameters
19 are set to near total reflection. The result of a surface current tally at the far
20 end of the guide (see figure 3) is shown in figure 6.

21 4.1. Cross Comparison

22 After reentering in MCNPX, figure 7 shows the *SSW/SSR*, *Compile* and
23 *Supermirror* performance in terms of spectrum and angular distributions at the
24 guide exit (surface current tally). For comparison, also the MCNPX built-in
25 mirror results are shown. The distributions agree very well both in terms of
26 peak position and size (neutron yield).

27 Another check is presented in figure 8 which based on the same test geometry
28 as figure 7, but the vacuum in the guide is replaced by air⁶. Given that no

⁶Dry air according to standard composition provided by the MCNPX group.

Figure 7: Spectrum (top) and incident angle comparison (bottom) of built-in specular MCNPX mirror, *SSW/SSR*, *Compile* and *Supermirror* approaches. The surface current tally is placed at the exit - see figure 3. The inserts below the main plots show the absolute differences of the various interface with respect to the built-in MCNPX specular mirror - the offset from zero is explained by the low angle reflectivity constant being different from unity. As expected all entries in the upper plot fall in the bin: $[19.6;20.6]\text{meV}$. Likewise, all entries in the lower plot fall in the bin covering the angular range: $[43.7;45.4]^\circ$.

1 attempt is made to simulate the effects of air in McStas, the non-zero bins in
 2 the histogram corresponding to *SSW/SSR* may seem surprising at first glance.
 3 They are in fact due to neutrons back-scattering off air molecules after the guide.
 4 Thus the various interfaces are considered to be validated and the McStas plug-
 5 ins to use them are made publicly available (under GNU licencing) via the
 6 McStas homepage: www.mcstas.org.

Figure 8: Logarithmic plots of spectra (top) and incident angle comparison (bottom) of built-in specular MCNPX mirror, *SSW/SSR*, *Compile* and *Supermirror* approaches using air-filled guide. The surface current tally is placed at the exit - see figure 3. The inserts below the main plots show the absolute differences of the various interface with respect to the built-in MCNPX specular mirror.

1 5. Prospects

2 The introduction of supermirrors in MCNPX [8] has already proved very
3 useful, and numerous simulations performed over the last years are based here-
4 upon. Our goal is that the combination of MCNPX and McStas will extent the
5 usability and become a new standard for detailed simulation of cold/thermal
6 neutron moderators. Besides being directly applicable to the simulation of the
7 target-moderator-reflector system of the spallation source, it will enable McStas-
8 based descriptions of e.g. reflecting material and crystals to be included in the
9 design and optimisation of advanced moderators. Examples include the recently
10 proposed Si-crystal vanes [11] or nano-diamond coatings [12], which by using the
11 directly coupled MCNPX-McStas interface could be simulated to a level beyond
12 what is possible with the MCNPX or McStas codes alone. Also we foresee that
13 the combination of MCNPX and McStas will enable more accurate calculation of
14 photon production along neutron guides, and thus ultimately yield better shield-
15 ing calculations. Finally, existing spallation sources have experienced problems
16 with crosstalk between neutron guides. Given that e.g. the beam-lines at ESS
17 are expected to be more closely spaced than at existing facilities, it is important
18 already before the construction phase to start studying these effects, and we
19 intend to do this using the coupled MCNPX McStas interface.

20 6. References

- 21 [1] L. S. Waters, G. W. McKinney, J. W. Durkee, M. L. Fensin, J. S. Hendricks,
22 et al., The MCNPX Monte Carlo radiation transport code, AIP Conf.Proc.
23 896 (2007) 81–90. doi:10.1063/1.2720459.
- 24 [2] G. Battistoni, S. Muraro, P. R. Sala, F. Cerutti, A. Ferrari, et al., The
25 FLUKA code: Description and benchmarking, AIP Conf.Proc. 896 (2007)
26 31–49. doi:10.1063/1.2720455.
- 27 [3] A. Ferrari, P. R. Sala, A. Fasso, J. Ranft, FLUKA: A multi-particle trans-
28 port code (Program version 2005).
- 29 [4] K. Lefmann, K. Nielsen, McStas, a General Software Package for Neutron
30 Ray-tracing Simulations, Neutron News 10 (1999) 20.
- 31 [5] P. Willendrup, E. Farhi, K. Lefmann, McStas 1.7 a new version of the
32 flexible Monte Carlo neutron scattering package, Physica B 350 (2004)
33 E735.
- 34 [6] P. Willendrup, E. Knudsen, E. Farhi and K. Lefmann, User and Program-
35 mers Guide to the Neutron Ray-Tracing Package McStas, Version 1.12c,
36 Risø–R–1416(rev.ed.)(EN) (2011).
- 37 [7] P. Willendrup, E. Knudsen, K. Lefmann and E. Farhi, Component
38 Manual for the Neutron Ray-Tracing Package McStas, Version 1.12,
39 Risø–R–1538(rev.ed.)(EN) (2011).

- 1 [8] F. X. Gallmeier, M. Wohlmuther, U. Filges, D. Kiselev, G. Muhrer, Im-
2 plementation of neutron mirror modeling capability into mcnp and its
3 demonstration in first applications, Nuclear Technology 168(3) (2009) 768–
4 772.
- 5 [9] D. Baxtor, A. Crabtree, P. Ferguson, F. Gallmeier, E. Iverson, W. Lu,
6 G. Muhrer, Spallation neutron source moderator overview. Presented at
7 IAEA Advanced Moderator Workshop, Tsukuba, Japan, November 22-25,
8 2011.
- 9 [10] <http://www.mcnpvised.com/>.
- 10 [11] E. Iverson, Contribution 336, ICANS XX, March 4 - 9, 2012. Bariloche,
11 Rio Negro, Argentina.
- 12 [12] V. Nesvizhevsky, Nuclear Instruments and Methods in Physics, A 595
13 (2008) 631.