Antimicrobial effect of nisin in processed cheese - Quantification of residual nisin by LC-MS/MS and development of new growth and growth boundary model for *Listeria monocytogenes*

Martinez Rios, Veronica; Pedersen, Mikael; Pedrazzi, Monica; Gkogka, Elissavet; Smedsgaard, Jørn; Dalgaard, Paw

Published in:
International Journal of Food Microbiology

Link to article, DOI:
[10.1016/j.ijfoodmicro.2020.108952](https://doi.org/10.1016/j.ijfoodmicro.2020.108952)

Publication date:
2021

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Martinez Rios, V., Pedersen, M., Pedrazzi, M., Gkogka, E., Smedsgaard, J., & Dalgaard, P. (2021). Antimicrobial effect of nisin in processed cheese - Quantification of residual nisin by LC-MS/MS and development of new growth and growth boundary model for *Listeria monocytogenes*. *International Journal of Food Microbiology*, 338, [108952]. <https://doi.org/10.1016/j.ijfoodmicro.2020.108952>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Antimicrobial effect of nisin in processed cheese - quantification of residual nisin by LC-MS/MS and development of new growth and growth boundary model for *Listeria monocytogenes*

Veronica Martinez-Rios, Mikael Pedersen, Monica Pedrazzi, Elissavet Gkogka, Jørn Smedsgaard, Paw Dalgaard

PII: S0168-1605(20)30446-3

DOI: <https://doi.org/10.1016/j.ijfoodmicro.2020.108952>

Reference: FOOD 108952

To appear in: *International Journal of Food Microbiology*

Received date: 3 June 2020

Revised date: 25 August 2020

Accepted date: 25 October 2020

Please cite this article as: V. Martinez-Rios, M. Pedersen, M. Pedrazzi, et al., Antimicrobial effect of nisin in processed cheese - quantification of residual nisin by LC-MS/MS and development of new growth and growth boundary model for *Listeria monocytogenes*, *International Journal of Food Microbiology* (2020), <https://doi.org/10.1016/j.ijfoodmicro.2020.108952>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Antimicrobial effect of nisin in processed cheese - Quantification of residual nisin by LC-MS/MS and development of new growth and growth boundary model for *Listeria monocytogenes*

Veronica Martinez-Rios^{1*}, Mikael Pedersen¹, Monica Pedrazzi^{1,3}, Elissavet Gkogka², Jørn Smedsgaard¹, Paw Dalgaard¹

¹ National Food Institute (DTU Food), Technical University of Denmark, Kgs. Lyngby, Denmark

² Arla Innovation Centre, Arla Foods a.m.b.a., Århus N, Denmark

* Corresponding author: Food Microbiology and Hygiene, National Food Institute, Technical University of Denmark, Kemitorvet, Building 204, DK-2800, Kgs. Lyngby, Denmark. E-mail: veri.food.dtu.dk (V. Martinez-Rios)

³ Present address: Via dell'industria 3, Verona, Italy.

ABSTRACT

This study tested the hypothesis that growth of *Listeria monocytogenes* in processed cheese with added nisin can be predicted from residual nisin A concentrations in the final product after processing. A LC-MS/MS method and a bioassay were studied to quantify residual nisin A concentrations and a growth and growth boundary model was developed to predict the antilisterial effect in processed cheese. 278 growth rates were determined in broth for 11 *L. monocytogenes* isolates and used to determine 13 minimum inhibitory concentration (MIC) values for nisin between pH 5.5 and 6.5. To supplement these data, 67 MIC-values at different pH-values were collected from the scientific literature. A MIC-term was developed to describe the effect of pH on nisin MIC-values. An available growth and growth boundary model ([doi: 10.1016/j.fm.2019.103255](https://doi.org/10.1016/j.fm.2019.103255)) was expanded with the new MIC-term for nisin to predict growth in processed cheese. To generate data for model evaluation and further model development, challenge tests with a total of 45 growth curves, were performed using processed cheese. Cheese were formulated with 11.2 or 12.0 ppm of nisin A and heat treated to obtain residual nisin A concentrations ranging from 0.56 to 5.28 ppm. Below 15°C, nisin resulted in extended lag times. A global regression approach was used to fit all growth curves determined in challenge tests. This was obtained by combining the secondary growth and growth boundary model including the new term for the inhibiting effect of nisin on μ_{max} with the primary logistic growth model with delay. This model appropriately described the growth inhibiting effect of residual nisin A and showed that relative lag times depended on storage temperatures. With residual nisin A concentrations, other product characteristics and storage temperature as input the new model correctly predicted all observed growth and no-growth responses for *L. monocytogenes*. This model can support development of nisin A containing recipes for processed cheese that prevent growth of *L. monocytogenes*. Residual nisin A concentrations in processed cheese were accurately quantified by the developed LC-MS/MS method with recoveries

of 83 to 110 % and limits of detection and quantification being 0.04 and 0.13 ppm, respectively. The tested bioassay was less precise and nisin A recoveries varied for 53% to 94%.

Keywords: Cardinal parameter model, global regression, growth boundary, ψ -value, product development

1. Introduction

Nisin A is an antimicrobial peptide belonging to the lantibiotic class of bacteriocins (Class I), composed of 34 aminoacids and with a molecular mass of 3.5 kDa. Nisin has antimicrobial activity against a wide range of Gram-positive food-borne pathogens and it influences both cell membrane pore formation and cell wall synthesis (Cotter et al., 2005). Commercial nisin preparations have been used for over 30 years and are allowed in foods as an added preservative (FDA, 1988; Thomas and Delves-Broughton, 2005). Nisin is used in processed cheese to control potential growth of spore-forming bacteria. The anaerobic environment, high moisture and pH of 5.4 to 6.0 usually found in processed cheese may allow germination of spores and growth of *Clostridium* spp. or facultative anaerobic *Bacillus* spp. if present in the product and not controlled by means of formulation (Delves-Broughton, 2008; Glass et al., 2017). Though nisin has a well-documented use for the control of these spore-forming bacteria in processed cheese formulations, little information is available with regards to its protective effect against vegetative pathogens such as *L. monocytogenes*, that may contaminate these products during consumer handling (Redmond and Griffith, 2003).

The antimicrobial activity of nisin against *L. monocytogenes* has been studied in broth or agar by determination of minimum inhibitory concentration (MIC) (Benkerroum and Sandine, 1988; Ferreira and Lund 1996; Martínez and Rodríguez, 2005). Furthermore, several nisin

bioassays have been developed over the years to measure the antimicrobial activity of nisin. The most widely used bioassay is the horizontal agar diffusion method relying on a test microorganism (Fowler et al., 1975; Papagianni et al., 2006; Tramer and Fowler, 1964). When estimating residual nisin A in foods one has to use a method describing the extraction of nisin A from foodstuffs. However, no information is available about recovery and precision for bioassays when used to estimate residual nisin A in foods (Pongtharangkul and Demirci, 2004; Rogers and Montville, 1991).

In the last years, more advance analytical methods have been developed to quantify nisin A present in foods. An ISO standard has been published for the quantification of nisin A in cheese by liquid chromatography/mass spectrometry (LC-MS) and LC-MS/MS (ISO, 2009). This LC-MS/MS method has been used to quantify nisin A in different types of cheeses (Fuselli et al., 2012; Ko et al., 2016; Schneider et al., 2011). However, the relation between concentrations of nisin measured by LC-MS/MS and its antimicrobial activity has not been studied.

The objectives of the present study were to develop and validate a LC-MS/MS method for quantification of residual nisin A in different cheeses and to determine the inhibitory effect of residual nisin A concentrations on growth responses of *L. monocytogenes* in cheese. The antilisterial effect of nisin A was studied by using a predictive food microbiology approach for comparison of observed and predicted growth. Firstly, a nisin MIC-term was developed including the effect of pH on nisin MIC-values and this term was added to an available cardinal parameter growth and growth boundary model. Secondly, processed cheese with added nisin A was studied by challenge testing to quantify growth of *L. monocytogenes* at different constant storage temperatures and product characteristics (pH, a_w , lactic-, acetic-, citric acids, orthophosphate salt and nisin A concentrations). These data was used to developed a simple secondary model for relative lag time (RLT). Lastly, the expanded growth and growth boundary model was evaluated by comparison of

observed and predicted growth for *L. monocytogenes* in cheeses when (a) assuming absence of nisin A, (b) using residual nisin A concentrations measured by LC-MS/MS and (c) using concentrations of nisin A added to the cheeses at formulation.

2. Materials and methods

2.1. Nisin and stock solutions

Nisin stock solutions (50 ppm and 500 ppm) were prepared with Nisaplin (2.5% nisin A, Danisco, batch number 4263501621) by adding 0.020 g or 0.200 g to 10 ml of 0.02N HCl. These stock solutions were stored at 4°C for a maximum of 24 hours.

2.2. Nisin quantification by LC-MS/MS

2.2.1. Nisin extraction method for LC-MS/MS analysis

Methanol:water (1:1 v/v) (HPLC grade, VWR chemicals, Søborg, Denmark) with 0.1% formic acid (695076, Sigma-Aldrich, St. Louis, MO, USA) adjusted with 5 M HCl to pH 2.0 was used as solvent to extract nisin A from cheese. Cheese (2.0 g) was homogenized in 30 mL extraction solvent with a T25 Ultra Turrax (IKA, Staufen, Germany). The suspension was sonicated for 10 min. at room temperature and then centrifuged (Eppendorf centrifuge 5810 R, Hørsholm, Denmark) at 4500 rpm for 20 min at 4°C. The supernatant was transferred to a volumetric flask and the precipitate was washed/extracted twice with 10 mL extraction solvent. Supernatants were combined and supplemented as necessary with extraction solvent to 50 ml and then filtered (0.2 µm Ministart®, Sartorius Stedim Biotech GmbH) prior to LC-MS/MS analysis.

2.2.2. LC-MS/MS analysis

For the LC-MS/MS analysis (Table 1), an EVOQ Elite^{TM} triple quadrupole mass spectrometer (Bruker, Frederikssund, Denmark), equipped with an electrospray ionization (ESI) source was coupled to an UltiMate 3000 ultra-high-performance liquid chromatography system (UHPLC) from Dionex Thermo scientific (Waltham, MA, USA). The UHPLC consisted of a pump, autosampler and column compartment. Chromeleon Xpress (Thermo Fisher Scientific, Waltham, USA) and MS Workstation 8 (Agilent, Santa Clara, USA) were used to perform system control and data acquisition on the UHPLC and mass spectrometer, respectively. Chromatographic separation was performed using a Zorbax 300 SB-C18 column (Table 1).

The mobile phase consisted of an aqueous solution with 0.1% formic acid (mobile phase A) and 100% acetonitrile (mobile phase B; Chromasolv^{TM} LC-MS, Honeywell, Seelze, Germany). A linear gradient was used as follows: 0-2 min 80% A; 2-6 min 20% A; 6-7 min 80% A, and held for 3 min to equilibrate the column. The column was maintained at 40°C. The flow rate was 0.3 mL/min and the injection volume for the main standards (0.1, 0.5, 1, 2.5, 5, 10 ppm) or samples was 10 µl. The diverter valve was used to eliminate NaCl interferences, directing the flow into the discharge for the first 3 minutes of each run, avoiding contamination of the ion source. The parent ion $[M+4H]^{4+}$ was identified as 839.0 (average mass, 3354). The fragment ions of nisin A were selected as 1081 (quantitative ion) and 811 (qualitative ion). Diagnostic ions were found using an infusion experiment. A standard solution of nisin A (1 ppm) was infused directly into the ion source without applying collision gas and collision energy to identify parent ions. Fragment ions were likewise identified with infusion of standard solution and collision gas and collision energy applied.

2.2.3. Calculation of nisin A

Content of nisin A in cheese samples was calculated using eq. (1).

$$\text{Nisin } A \text{ (ppm)} = \frac{H_t \cdot \rho_{nA,s} \cdot V_t}{H_s \cdot m_t}$$

(1)

where H_t is the test sample peak area, $\rho_{nA,s}$ is the concentration in ppm of the standard solution used to construct the calibration curve, V_t is the test sample volume (50 ml), H_s is the peak area of $\rho_{nA,s}$ and m_t is the mass (g) of cheese sample.

2.3. Nisin quantification by bioassay

The method described in this section combined a simple nisin extraction method and quantification by a plate diffusion assay with an indicator organism (Tramer and Fowler, 1964).

2.3.1. Nisin extraction for the bioassay

Cheese (2.0 g) was macerated in a 50 ml glass beaker using 10 mL of 0.02 N HCl with pH 2.0 and magnet stirring for 30 min. Thereafter the mixture was placed on a hot plate (pre-heat to 100 °C) with magnetic stirring and boiled for 5 min. 40 ml of 0.02N HCl was used to wash the beaker and the sample was transferred to a 50 ml falcon tube and centrifuged at 4500 rpm for 20 min at 4°C (Eppendorf centrifuge 5810 R, Hørsholm, Denmark). The supernatant with nisin was analysed as described below.

2.3.2. Indicator organism and pre-culture conditions

Preliminary studies with both *Micrococcus luteus* DSM 1790 and *Lactobacillus sakei* ATCC 15521 as indicator organisms were performed. Quantification of nisin did not differ between the two indicator organisms and *M. luteus* was retained as its yellow colonies produced a more clear inhibition zone. *M. luteus* DSM 1790 was transferred from storage at -80°C to Nutrient broth (NB) (CM001, Oxoid, Hampshire, UK) and incubated for 24h at 30°C. Subsequently, the culture with *M.*

luteus was streaked on NB agar (1.3%) plates and incubated for 48h at 30°C. 200 ml NB containing a magnet was inoculated with a single colony and incubated at 30°C with stirring (200 rpm) for 24h. The cell concentration was then determined by direct phase contrast microscopy at 1000x magnification considering that one cell per field of view corresponded to a concentration around 10^6 cfu/ml (Adams and Moss, 2016).

2.3.3 Agar diffusion bioassay

Molten NB (1.3%) agar tempered at 48°C was supplemented with 2% of a 1:1 mixture of Tween 20 (P9416, Sigma-Aldrich, St. Louis, USA) and sterile water, previously held for 30 minutes at 48°C, and thoroughly mixed. The medium agar (200 ml bottles) were held for 30 minutes at 46°C and then inoculated with the indicator organism to reach a final concentration of 10^6 cfu/ml. 20 ml of inoculated medium was placed in each sterile Petri dish (100 x 20 mm). After agar solidification at room temperature, wells of 5 mm diameter were made with a cork borer. 30 µl of nisin standard solutions (0.1, 0.5, 1, 2.5, 5, 10 ppm; pH 2.0) were placed in each well and the plates were incubated at 30°C for 72 h. The diameter of inhibition zones around wells with nisin were measured on images of the plates by using an automated image analyser software (ImageJ Basics, v. 1.38, 2008). Four zones were measured for each nisin concentration and assays were performed in triplicate on three different days ($n = 4 \times 3 \times 3$).

2.4. Evaluation of methods for quantification of nisin

The quantification methods were evaluated for linearity, specificity, recovery and precision (NMKL, 2009). The only difference in the evaluation of the methods studied was the used of two matrixes (PC and SPC) for LC-MS/MS or exclusively one (SPC) for the evaluation of the bioassay. Standard curves with four concentrations (including zero) (x-axis) and the corresponding peak areas

determined by the LC-MS/MS method (y-axis) were developed for both PC and SPC. Likewise, standard curves for the bioassay were developed with four concentrations and the corresponding inhibition zone but exclusively for SPC. Linearity was evaluated using two replicates for each nisin concentration, in three different days. LC-MS/MS linearity of the response was further studied by plotting residuals against concentrations of standard curves performed for the two types of cheeses. LC-MS/MS specificity was studied by analysing samples of each matrix, to evaluate the possibility for interference of some endogenous compounds. The results were assessed by checking the presence of interfering substances with the same retention time as nisin A, in comparison to cheese samples fortified with nisin A. The recovery and precision (repeatability and reproducibility) of both, the LC-MS/MS and bioassay method were determined by spiking experiments where PC and SPC samples were fortified with 0.1, 0.5 and 1 ppm of nisin A before and after the extraction procedure. Precision with limits of acceptability $\leq 16\%$ for CV and 80-110% for recovery were based on duplicate analysis on three different days for each concentration level. Precision was the degree of agreement between independent analysis results obtained in different days and was calculated by the standard deviation of the method's repeatability and reproducibility. Data analyses were performed using the Microsoft Excel software. The limit of detection (LOD) was calculated as three times the within sample standard deviation for reproducibility (S_R) and the limit of quantification (LOQ) was calculated as 10 times S_R . Matrix effects (suppression or enhancement of signal due to other substances in the sample) were calculated by comparing slopes for two types of calibration curves. Curve type I, was prepared by spiking extraction solvent with nisin A to reach 0.1, 0.5 and 1 ppm. Curve type II, was prepared by spiking cheese samples with nisin A to reach 0.1, 0.5 and 1 ppm. Matrix effect was calculated by division of the slope of curve type II with the slope of curve type I.

2.5 Development and evaluation of growth and growth boundary model for *Listeria monocytogenes*

2.5.1. Bacterial strains and pre-culture conditions

Seven dairy related strains of *L. monocytogenes* were provided by Arla Foods or Martinez and Rodríguez (2005). The strains were used as a cocktail (*Lm-mix* including SLU 92, 612, LM 19 and 6) or individually (LM 13, LM 4, LM 4R) to determine μ_{max} -values in broth and for inoculation of challenge tests. Strain LM 4R was a spontaneous nisin resistant derivative from LM 4. Each strain was transferred from storage at -80°C to Brain Heart Infusion (BHI) broth (CM1135, Oxoid, Hampshire, UK) and incubated for 24h at 25°C. Subsequently, for broth studies the strains were pre-cultured at 7°C or 20°C in BHI broth adjusted with HCl to pH 5.5, 6.0 or 6.5. For challenge tests the individual strains (SLU 92, 612, LM 19, 6) were pre-cultured at a temperature ranging from 8°C to 20°C in BHI broth with 1% NaCl and pH 5.5 (HCl) to simulate conditions in processed cheese. Pre-cultures were grown to a relative increase in absorbance (540 nm) of 0.05 to 0.2 (Novaspec II, Pharmacia Biotech, Allerød, Denmark). *Lm-mix* used in broth and challenge tests was obtained by mixing different volumes of individually pre-cultured strains depending in the absorbance measurement. The *Lm-mix* concentration was determined by direct phase contrast microscopy at 1000x magnification considering that one cell per field of view corresponded to a concentration around 10^6 cfu/ml (Adams and Moss, 2016).

2.5.2. Cardinal parameter term for nisin and expansion of growth and growth boundary model

The effect of 24 different nisin A concentrations (0-10 ppm nisin, Nisaplin®, Danisco, Braband, DK in 0.02N HCl) on μ_{max} -values of *Lm-mix* or individual strains were determined in duplicate by automated absorbance measurements at 540 nm (BioScreen C, Labsystems, Helsinki,

Finland) at 7°C or 20°C using BHI broth adjusted to pH 5.5, 6.0 or 6.5 with HCl and sterilized by filtration (0.2 µm, Minisart®, Sartorius Stedim Biotech GmbH, Goettingen, Germany). Detection times, defined as the incubation time necessary to observe an increase in absorbance of 0.05 from the lowest absorbance measured in the beginning of incubation were determined for each absorbance growth curve. μ_{max} -values were determined from absorbance detection times for serially diluted inoculation levels of 10^2 , 10^3 , 10^4 , 10^5 and 10^6 cfu/ml as previously described (Dalgaard and Koutsoumanis, 2001). A total of 278 μ_{max} -values of *L. monocytogenes* were determined experimentally in BHI-broth. Nisin MIC-values, μ_{ref*} , n1 and n2 were estimated by fitting eq. (2) to square root transformed μ_{max} -values obtained for the studied nisin concentrations.

$$\sqrt{\mu_{max}} = \sqrt{\mu_{ref*} \cdot \left(1 - \left(\frac{nisin}{MIC_{nisin}}\right)^{n_1}\right)^{n_2}} \quad (2)$$

where μ_{max} is the maximum specific growth rate (h^{-1}) and μ_{ref*} is the reference maximum specific growth rate for the studied temperature. When fitting eq. (2), n1 was set to 0.5 or 1 and n2 was set to 1 or 2 (Dalgaard, 2009) in order to describe data most appropriately and this was determined from root mean square error (RMSE) values. The “n” values are specific for the combination of the microorganism and the inhibitory compound. A simple equation was used to model and predict the effect of pH on log-transformed nisin MIC-values, $\log MIC_{nisin} = a + b * pH$.

A total of 67 nisin MIC-values, for different *L. monocytogenes* strains and determined in broth or agar with different pH values (relevant for processed cheese), were extracted from literature. Four MIC-values were obtained at pH 5.0, 39 MIC-values at pH 5.5 and 24 MIC-values at pH 6.8 for a total of 29 different strains (Ferreira and Lund 1996; Taylor, 2009).

The growth and growth boundary model of Martinez-Rios et al. (2019a) (Supplementary Table 1) previously validated for processed cheese and containing terms for the inhibitory effect of temperature, a_w , pH, lactic, acetic and citric acids as well as phosphate melting salts was expanded

with a nisin-term (eq. (2)) to predict growth of *L. monocytogenes* in processed cheese containing nisin (Eq. S3 in Supplementary Table 4)

2.5.3. *L. monocytogenes* growth in processed cheese with added nisin

To generate data for model development and evaluation, growth of *L. monocytogenes* in commercial PC and laboratory produced SPC was determined in 15 challenge tests resulting in 45 maximum specific growth rates, lag times and initial and maximum cell concentrations at constant temperatures from 3.6°C to 19.5°C.

2.5.3.1. Production, inoculation and microbiological analysis of processed cheese

Arla Foods supplied PC cheese formulated with 11.2 ppm of added nisin A and SPC cheese formulated with 12 ppm of nisin A was produced in the laboratory. SPC was produced by mixing and cooking the ingredients (Supplementary Table 2) in a Themomixer (TM5, Vorwerk, USA). The butter was melted at 40°C for four minutes, then the water was added and the mixture was blended for four more minutes at 40°C. Subsequently, the dry ingredients were added and blended for an additional four minutes at 40°C. At this point, the pH of SPC was adjusted (5.4-5.9) with 50% (w/v) citric acid monohydrate. Lastly the mixture was blended at 2000 rpm and cooked with different temperature/time profiles (Supplementary Table 3) as recorded by data loggers (TinytagPlus, Gemini Data Loggers Ltd, Chichester, UK). To increase the creaminess of SPC, the mixture was blended at 3100 rpm for 3 min. Both PC and SPC cheeses were inoculated 24h after production with 0.1% (v/w) of *Lm-mix* appropriately diluted in chilled saline water (0.85% NaCl) to obtain an initial concentration between 1 and 2 log cfu/g. After inoculation, cheeses were packaged and stored in containers with 50 ± 1 g cheese. Storage temperature was recorded by data loggers (TinytagPlus). At each time of sampling, 10 g of cheese from three different containers were diluted 10-fold with

chilled physiological saline (0.85% NaCl and 0.10% Bacto-peptone), homogenized for 30 s at normal speed in a Stomacher 400 (Seward Medical, London, UK) and then 10-fold serial dilutions were performed. Viable counts of *L. monocytogenes* were determined by surface plating on PALCAM agar base (CM0877, Oxoid, Hampshire, UK) with PALCAM selective supplement (SR0150, Oxoid, Hampshire, UK) and incubation at 37°C for 48 h.

To estimate the kinetic growth parameters for each growth curve the primary integrated and log transformed logistic model with delay (four parameter model) or without delay (three parameter model) (eq. (3)); Rosso et al., 1996) was fitted to all individual growth curves of *L. monocytogenes* obtained in challenge tests at constant temperatures. Fitted parameter values for initial cell concentration (Log N_0 , Log cfu/g), lag time (t_{lag} , h), maximum specific growth rate (μ_{max} , h⁻¹) and maximum population density (Log N_{max} , Log cfu/g) were determined using the MS-Excel Solver Add-in (Microsoft Corp. Redmond, WA., USA). Data was reported as average $\pm$ standard deviation for each challenge tests. An F-test was used to determine if the lag time was significant.

$$\text{Log } (N_t) = \text{Log } (N_0) \quad \text{if } t < t_{lag}$$

$$\text{Log } (N_t) = \text{Log} \left(\frac{\frac{N_{max}}{N_0}}{1 + \left(\left(\frac{N_{max}}{N_0} \right) - 1 \right) \exp(-\mu_{max}(t - t_{lag}))} \right) \quad \text{if } t \geq t_{lag} \quad (3)$$

where t is the storage time (h) and N_t is the cell concentration (cfu/g) at time t . Other parameters were indicated above.

2.5.3.2. Product characteristics

Product characteristics of cheeses were determined in triplicate at the start of each challenge test. In addition, pH was measure every day of microbiological analysis by a PHC725 SN probe (Hach, Brønshøj, Denmark) after 1h stirring of 5 g sample in 25 ml of distilled water. NaCl was

quantified by automated potentiometric titration (785 DMP Titrino, Metrohm, Hesisau, Switzerland) and a_w was measured by a water activity meter (Aqua Lab model CX-2, Decagon devices Inc., Pullman, US). Concentrations of lactic-, acetic- and citric acids were determined by HPLC using external standards for identification and quantification (Østergaard et al., 2014). Concentration of orthophosphate salt was determined by Eurofins (test method QA02S) using ion chromatography. Nisin concentrations were measured by LC-MS/MS (see section 2.2).

2.5.4. Evaluation of existing predictive models including the effect of nisin

Three predictive models including the inhibiting effect of nisin were evaluated to assess their ability to predict growth responses of *L. monocytogenes* in processed cheese containing nisin. The studied models included the models of Bouttiford et al. (2000), Boziaris and Nychas (2006) and Parente et al. (1998). These models were developed to predict the effect of nisin on survival or growth inhibition of *L. monocytogenes* in broth and did not consider the distinction between residual and added nisin A. Nevertheless, these models were evaluated by using the concentrations of added nisin A and the residual concentrations of nisin A in the studied cheeses. Evaluation was performed by comparison of predicted and observed growth and no-growth responses with calculation of the percentage of all samples that were correctly predicted. Incorrect predictions were described as fail-safe (growth predicted when no growth was observed) or as fail-dangerous (no growth predicted when growth was observed).

2.6. New growth and growth boundary model for *L. monocytogenes* in processed cheese containing nisin

The effect of temperature on lag times of *L. monocytogenes* in processed cheese containing nisin (see section 2.5.3.1.) was assessed by applying the relative lag time (RLT) concept ($RLT = t_{lag}$

· $\mu_{max}/\ln(2)$) (Mellefont and Ross, 2003). It was evaluated by using an F-test if RLT-values were constant ($RLT = k_1$) or dependent on storage temperature (eq. 4).

$$RLT = k_1 + k_2/T^3 \quad (4)$$

In a similar way, it was evaluate if $\log(N_{max})$ was constant (k_3) or dependent on storage temperature (eq. (5))

$$\text{Log}(N_{max}) = k_3 + k_4 \cdot T \quad (5)$$

Values of the parameters μ_{ref} in eq. (S3), k_1 and k_2 (eq. (4)), as well as k_3 and k_4 (eq. (5)) were estimated by using both a classical two-step predictive microbiology approach and a global regression procedure (Hereu et al., 2014; Jewel, 2012). For the classical two-step approach μ_{ref} in eq. (S3) was calibrated using μ_{max} -values from challenge tests (Table 4). Calibration allowed finding a μ_{ref} -value so that the model predictions corresponded to observed data as well as possible. Likewise, k_1 and k_2 in eq. (4) as well as k_3 and k_4 in eq. (5) were independently estimated from RLT - and $\log(N_{max})$ -values observed in challenge tests with growth of *L. monocytogenes* in processed cheese containing nisin A. Global (one-step) regression was used to determine values of μ_{ref} , k_1 , k_2 , k_3 and k_4 by combining the growth and growth boundary model (eq. (S3)) with the RLT model (eq. (4)), the $\text{Log}(N_{max})$ model (eq. (5)) and the primary logistic growth model with delay (eq. (6)). Eq. (6) was fitted in one regression to the entire data set with 45 growth curves and including, respectively, 157 and 417 cell concentrations ($\log$ cfu/g) for PC and SPC. Statistical performance of the classical two-step approach and the global regression procedure were compared by calculation of RMSE-values.

$$\text{Log}(N_t) = \text{Log}(N_0) \quad \text{if } t < t_{lag}$$

$$\text{Log } (N_t) = \text{Log} \left(\frac{10^{k_3+k_4 \cdot T}}{1 + \left(\left(\frac{10^{k_3+k_4 \cdot T}}{N_0} \right) - 1 \right) \cdot \exp \left(-(\mu_{\max, \text{Eq.}(S3)}) \cdot \left(t - \left(\frac{(k_1+k_2/T^3) \cdot \ln(2)}{\mu_{\max, \text{Eq.}(S3)}} \right) \right) \right)} \right) \quad \text{if } t \geq t_{\text{lag}} \quad (6)$$

2.7. Evaluation of new growth and growth boundary model

Performance of the new growth and growth boundary model including the effect of nisin (eq. (S3)) was evaluated by comparing observed and predicted growth and no-growth responses for the 45 growth responses determined in challenge tests with *L. monocytogenes* (See 2.5.3). The percentage of correct, fail-safe and fail-dangerous predictions were calculated (See 2.5.4). For each experiment with corresponding product characteristics and storage temperature, growth of *L. monocytogenes* was predicted for three scenarios: (a) assuming absence of nisin, (b) using the measured residual nisin concentration and (c) using the concentration of nisin added to the cheeses at formulation.

3. Results and discussion

3.1. Quantification of nisin in cheese by LC-MS/MS and bioassay

A clearly distinguishable peak with retention time of 4.48 min for nisin A was observed (Fig. 1). The LC-MS/MS method showed linear responses for nisin A concentrations ranging from 0.1 to 10 ppm ($r^2 > 0.98$). Residuals of LC-MS/MS responses at different nisin concentrations were randomly distributed, indicating variance homogeneity (Results not shown). The LOD for nisin A in cheese was 0.04 ppm, and the LOQ was 0.13 ppm. No matrix interference was observed for nisin A detection in the tested cheeses (PC and SPC). Precision was $< 16\%$ and nisin A recoveries from

the two studied cheeses ranged from 83 to 110 % (Table 2). Recovery of nisin A in SPC tended to increase with nisin concentrations (Table 2).

The bioassay showed linear responses for nisin A concentrations ranging from 0.1 to 1 ppm with a correlation coefficient $r^2 > 0.88$. Residuals were randomly distributed, indicating variance homogeneity (Results not shown). The LOD for nisin A in cheese was 0.07 ppm, and the LOQ was 0.25 ppm. Precision was $>16\%$ and nisin A recoveries from SPC ranged from 53 to 94 % (Table 2).

Clearly, LC-MS/MS measurement of residual nisin A in processed cheese was reproducible and provided a more precise quantifications of nisin A compared to the studied bioassay. The LC-MS/MS method developed in the present study provided recovery values similar to those previously obtained by Ko et al. (2016); however the present study used an extraction solvent without NaCl as it can interfere with ionization, cause ion suppression and broader peaks in the LC-MS/MS (Sterling et al., 2010). Residual concentrations of nisin A quantified in the present study using LC-MS/MS (0.56-5.28 ppm) were lower than the added concentrations used in the cheese formulations (11.2-12.0 ppm). These differences are in agreement with other studies on the stability of nisin in broth or foods after heat treatments (Davies et al., 1998). For instance, Schneider et al. (2011) found the nisin A concentrations determined by LC-MS/MS in processed cheese to be reduced more than 92 % by a heat treatment at 120 °C during 30 min. Ko et al. (2016) also quantified low nisin A concentrations in processed cheese by LC-MS/MS (0.34-0.64 mg/kg) but no information on nisin A concentrations recipes were provided.

3.2. Effect of nisin on growth rates of *L. monocytogenes* in laboratory media

The experimentally determined nisin MIC-values at pH 5.5 and at 7°C or 20°C ranged between 0.55 ± 0.05 ppm and 3.3 ± 0.1 ppm. At pH 6 and at 20°C the nisin MIC-value was 8.8 ± 1.2 ppm and at pH 6.5 the values were between 6.4 ± 1.0 and 24.9 ± 3.1 (Table 3; Fig. 2). Markedly,

higher MIC-values were determined for *L. monocytogenes* LM 4R compared to the other strains studied, including its wild type LM4 (Table 3). In order to best fit the μ_{max} -values by eq. (2) both n1 and n2 were found to be 1 for all strains studied. Nisin MIC-values for *L. monocytogenes* increased significantly ($p < 0.05$) with increase of pH and this was observed for data from the present study as well as for data from the scientific literature (Fig. 3). To predict MIC-values of nisin depending on pH a nisin MIC-term based on a simple regression model was developed (eq. 7)

$$\log(MIC_{\text{nisin}}, \text{ppm}) = -(3.08 \pm 0.32) + (0.64 \pm 0.05 \cdot \text{pH}) \quad R^2 = 0.65 \quad (7)$$

Several studies found nisin to be more effective in acidic environments (Khan et al., 2015; Thomas and Wimpenny, 1996) and this is in agreement with eq. (7). This simple model was developed including both experimental and literature nisin A MIC-values and reflected variability among 29 *L. monocytogenes* strains; an aspect also previously discussed by Katla et al. (2003). This simple model (eq. (7)), can estimate nisin A MIC-values for products with pH values outside the average values (5.4-6.0) for processed cheese and therefore support new formulations with altered pH values. Furthermore, it can support the development of other foods products with added nisin A such as dairy, meat and fish products with higher pH than processed cheese (Davies et al., 1997; Nilsson et al., 1997; Oshima et al., 2014; Ruiz et al., 2009; Schillinger et al., 2001). Importantly, this simple model predicts with reasonable precision (RMSE of 0.54) the MIC-values of nisin A (32-37 ppm) at pH 7.2-7.3 when compared to 53 MIC-values (average of 17 ppm) extracted from the literature for various isolates of *L. monocytogenes* (Campion et al., 2013; Hara et al., 2009; Martínez and Rodríguez, 2005; Neetoo et al., 2008; Popowska et al., 2006; Smith et al., 2016; Taylor et al., 2009).

3.3. Product characteristics of processed cheese and *L. monocytogenes* growth kinetics

Commercial PC and laboratory produced SPC showed some variation in initial pH (5.4-6.1) and a_w (0.970-0.983) (Table 4). More variability was observed for water phase concentrations of lactic acid (869-9869 ppm), acetic acid (<LOD-2384 ppm), citric acid (3766-20864 ppm) and concentration of orthophosphate (0.8-2.00%). The variability observed in residual nisin A concentrations among challenge test (CT7-15) (Table 4) was expected as a result of the different heat treatments used during preparation (Supplementary Table 3).

L. monocytogenes grew in both PC and SPC from 7 to 20°C, and μ_{max} -values were influenced by storage conditions and product characteristics (Table 4). Residual nisin A concentrations and pH had a pronounced effect on *L. monocytogenes* μ_{max} -values, as suggested by eq. (6). This is for example seen from the challenge tests 7 to 12, which were all formulated with 12 ppm nisin A. However, due to its heat treatment, challenge test 10 had lowest residual nisin A concentration (0.88 ppm) and also highest pH of 5.9 and this resulted a μ_{max} -value of 0.023 h⁻¹ compared to 0.000 h⁻¹ for the other challenge tests (Table 4). The observed long lag times at low temperatures for processed cheese containing nisin were in agreement with previous studies where researchers reported extended lag times for *L. monocytogenes* and other microorganisms in products containing nisin (Davies et al., 1997; Nilsson et al., 1997; Oshima et al., 2014; Ruiz et al., 2009; Schillinger et al., 2001). For example, Davies et al. (1997) showed that the incorporation of 2.5 ppm of nisin A in ricotta-cheese after production extended the lag times of *L. monocytogenes* by 44 days compared with the same product supplemented with 1.25 ppm when stored between 6-8°C. In the same way, nisin A was found to extend the lag times of *L. monocytogenes* in tofu stored at 10°C (Schillinger et al., 2001).

3.4. Evaluation of available predictive models including the growth inhibiting effect of nisin A

The effect of nisin on growth of *L. monocytogenes* in combination with other environmental factors including temperature, pH and NaCl/ a_w has previously been studied in broth and predictive models were suggested by Bouttefroy et al. (2000), Boziaris and Nychas (2006) and Parente et al. (1998). These models did not consider the distinction between residual and added nisin and they were not previously evaluated for their ability to predict growth responses of *L. monocytogenes* in foods containing nisin. Interestingly, the performance of two out of the three models improved when residual nisin A concentrations rather than added nisin A concentrations in processed cheese (Table 5) were used as model input together with other measured product characteristics and storage conditions (Table 4). Most notable, the percentage of correct predictions increased from 14% to 100% for the model of Boziaris and Nychas (2006) when nisin A concentrations measured by LC-MS/MS were used as model inputs rather than added nisin A concentrations (Table 5). These results suggest residual nisin A concentrations may be more useful than added nisin A concentrations when the objective is to predict the growth responses of *L. monocytogenes* in processed cheese containing nisin A.

3.6. Modelling growth of *L. monocytogenes* in processed cheese containing nisin

RLT-values significantly ($p < 0.05$) increased at low storage temperatures and this effect was appropriately described by eq. (4) (Fig. 4). In contrast, storage temperature had no significant effect ($p > 0.05$) on $\log(N_{max})$.

The model of Martinez-Rios et al. (2019a) was expanded with a nisin term that included MIC_{nisin} -values as described by eq. (7). For this new growth and growth boundary model (eq. (S3)), the μ_{ref} -value of 0.419 in the original model, increased markedly (Table 6) when the model was calibrated to describe the data for growth of *L. monocytogenes* in processed cheese containing nisin

A (Table 4). Importantly, all parameter values in eq. (S3), other than μ_{ref} , were kept constant during this product calibration of the growth and growth boundary model. The global (one-step) regression approach described the growth data (log cfu/g) slightly better than the two-step approach and resulted in the highest calibrated μ_{ref} -value of 0.629 (Table 6).

The long lag times (Table 4) and corresponding high RLT-values at low storage temperatures (Fig. 4; Fig 5b) observed together with increased growth rates (μ_{ref} -value; Table 6) of *L. monocytogenes* growing in processed cheese containing nisin A are in agreement with previous studies. In fact, increased μ_{max} -values in different foods were reported when the pathogen resumed growth after extended lag times (Davies et al., 1997; Nilsson et al., 1997; Ruiz et al., 2009). As shown in Figure 5b, the model of Martinez-Ros et al. (2019a) was unable to describe the RLT-values observed at low temperatures in challenge tests (Table 4). However, combining simple models for the effect of pH on MIC_{nisin} -values (eq. (7)) and for the effect of temperature on RLT-values (eq. (4)) with an extensive growth and growth boundary model (eq. S3) allowed this growth pattern to be described (Table 6) as shown in Figure 5b.

3.7. Evaluation of the new growth and growth boundary model

The new growth and growth boundary model (eq. S3) correctly predicted 100% of the growth or no growth responses for *L. monocytogenes* in processed cheese with nisin A. This positive result was obtained when using residual nisin A concentrations, determined by the developed LC-MS/MS method, as model input (Table 7). It is interesting that the same model was less performant both when nisin A concentrations added to products and when no nisin were used as model input together with other measured product characteristics and storage conditions (Table 4, Table 7). This supports the approach of the present study that accurate measurement of residual nisin A concentrations is important to predict growth of *L. monocytogenes* in foods that contain

nisin. Furthermore, these results represent a successful product validation of the new growth boundary model (eq. (S3)) for processed cheese with nisin A. However, when used to predict growth of *L. monocytogenes* in processed cheese with nisin, further evaluation of this model with additional data is needed. This is the case as model parameters related to lag time and growth rates were calibrated to data in Table 4, and these data therefore cannot be used to evaluate and validate the same model. The range of applicability, where the new model has been successfully validated for prediction of growth and no-growth responses include processed cheese with residual nisin A concentration of from 0.56 ppm to 5.28 ppm together with other product characteristics and storage condition as shown in Table 4.

86% correct and 14% fail-safe predictions of growth or no-growth responses were obtained by assuming no added nisin A in challenge tests (Table 7). These predictions correspond to using the model of Martinez-Rios et al. (2019a) that does not include a nisin term (eq. (S1)). This model however, include terms for the melting salts di-phosphate (P2) and tri-phosphate (P3) and these terms may be included in the new growth and growth boundary model to expand its range of applicability to processed cheese containing mixtures of different phosphate melting salts. As the developed nisin A term (eq. (7)) reasonably predicted MIC_{nisin} -values at pH as high as 7.2-7.3 (see section 3.3) it is also likely that the new model can be validated above the pH of 5.4-6.1 currently studied for processed cheese (Table 4). The logit model of Boziaris and Nychas (2006), including the effect of temperature, a_w , pH and nisin, also correctly predicted 100% of the observed growth and no-growth responses (Table 5). Compared to that model, the new model has the advantage of including the effect of more environmental factors relevant for processed cheese such as ortho-phosphate salt and organic acids as discuss below. Ψ is a quantitative measurement for the distance between specific environmental conditions and the growth boundary of *L. monocytogenes* ($\psi = 1$) (see Supplementary Table 5 for details about calculation of the ψ -value). Compared with logit

models where a probability response of growth/no-growth is obtained; the new model provides a mathematical value (ψ) representing how far the developed formulation is from the growth boundary of *L. monocytogenes*. The new model (Eq. (S3)) included the combined effect of eight environmental factors on the growth boundary of *L. monocytogenes* in processed cheese. As previously described and discussed in several studies such extensive models can be developed by variations of the cardinal parameter approach, which is much less data demanding than for example polynomial models or logistic regression (logit) models. Due to their demand for data, polynomial and logit models most often are limited to include the combined effect of between 4 to 7 environmental factors even when studied in liquid laboratory media and the calibration of these model for specific food remains a challenge (Dalgaard and Mejlholm, 2019; Le Marc et al., 2002; Ross and Dalgaard, 2004; Razavilar and Genigeorgis, 1998).

The developed extensive growth boundary model (Eq. (S3)) included the combined effect of eight environmental factors which will facilitate its use for product development and re-formulation. As an example if a processed cheese is contaminated by a consumer after opening a package, the new growth and growth boundary model will predict growth of *L. monocytogenes* with a ψ -value of 0.4 at 5 °C when the cheese contain 1.0 ppm of residual nisin A, pH 6.0, a_w of 0.980 and water phase concentrations of lactic acid (4,000 ppm), acetic acid (1,000 ppm), citric acid (10,000 ppm) and orthophosphate (0.8 %). By lowering the pH of this product to 5.75 with water phase concentrations of lactic acid (6,000 ppm) and acetic acid (2,000 ppm), *L. monocytogenes* will be predicted to be at its growth boundary with a ψ -value of 1.0. However, if the residual nisin A concentration in the processed cheese with pH 5.75 is increased from 1.0 to 2.5 ppm then the predicted ψ -value will become 2.2 and the product will be stabilized against growth of *L. monocytogenes* during the open shelf-life with a solid margin to its growth boundary (Dalgaard and Mejlholm, 2019). Clearly, practical application of the new growth and growth boundary model to

support product development and re-formulation of processed cheeses, stabilized against growth of *L. monocytogenes*, during the open shelf-life requires information about product characteristics. Residual nisin A concentrations in processed cheese can be measured by LC-MS/MS (see 3.1) and further information about added nisin A in formulations, applied heat processing and stability of nisin A during storage is likely to be required to obtain desired levels of residual nisin A (Supplementary Table 3). In conclusion, the present study quantified and modelled the effect of pH on nisin A MIC-values for *L. monocytogenes* and included the effect in a cardinal parameter growth and growth boundary model that can predict the effect of residual nisin A concentrations on growth/no-growth responses in cheese. The new cardinal parameter growth and growth boundary model is useful to predict growth/no-growth and consequently can support product development when used together with LC-MS/MS to determine residual nisin A concentrations. Further studies to determine the precision and range of applicability for the growth part of the new growth boundary model is desirable.

Acknowledgements

The Danish Dairy Research Foundation funded the present study as part of the project ‘Predictive food microbiology tool for risk assessment and documentation of food safety’. We thank Dr. Beatriz Martinez Fernández and Dr. Ana Rodríguez González from the Instituto de Productos Lácteos de Asturias (IPLA-CSIC) for providing *L. monocytogenes* strains LM4 and LM4R. We also thank Trine Bundesen from DuPont for providing Nisaplin®. Lastly, we thank laboratory technicians Rannvá Høhnadóttir Houmann, Else Petrine Celie Berthelsen and Peter Kristian Kristoffersen for their assistance with challenge tests and development of the nisin bioassay.

References

- Adams, M.R., Moss, M.O., McClure, P.J., 2016. Methods for the microbiological examination of foods. In: Food Microbiology. The Royal Society of Chemistry, Croydon, UK, pp. 413–446.
- Benkerroum, N., Sandine, W.E., 1988. Inhibitory action of nisin against *Listeria monocytogenes*. J. Dairy Sci. 71, 3237–3245. [https://doi.org/10.3168/jds.S0022-0302\(88\)79929-4](https://doi.org/10.3168/jds.S0022-0302(88)79929-4)
- Boziaris, I.S., Nychas, G.J.E., 2006. Effect of nisin on growth boundaries of *Listeria monocytogenes* Scott A, at various temperatures, pH and water activities. Food Microbiol. 23, 779–784. <https://doi.org/10.1016/j.fm.2006.03.003>
- Bouttefroy, A., Mansour, M., Linder, M., Milliere, J-B., 2000. Inhibitory combinations of nisin, sodium chloride, and pH on *Listeria monocytogenes* ATCC 15313 in broth by an experimental design approach. Int. J. Food Microbiol. 54, 109–115. [https://doi.org/10.1016/S0168-1605\(99\)00171-3](https://doi.org/10.1016/S0168-1605(99)00171-3)
- Campion, A., Casey, P.G., Field, D., Cotter, P.D., Hill, C., Ross, R.P., 2013. In vivo activity of nisin A and nisin V against *Listeria monocytogenes* in mice. BMC Microbiol. 13:23.
- Cotter, P.D., Hill, C., Ross, R.P., 2005. Bacteriocins: developing innate immunity for food. Nat. Rev. Microbiol. 3, 777–788. <https://doi.org/10.1038/nrmicro1273>
- Dalgaard, P., 2009. Modelling of microbial growth. Bulletin of the International Dairy Federation 433, 45–60.
- Dalgaard, P., Koutsoumanis, K., 2001. Comparison of maximum specific growth rates and lag times estimated from absorbance and viable count data by different mathematical models. J. Microbiol. Methods. 43, 183–196. [https://doi.org/10.1016/S0167-7012\(00\)00219-0](https://doi.org/10.1016/S0167-7012(00)00219-0)
- Dalgaard, P., Mejlholm, O., 2019. Modelling growth of *Listeria* and lactic acid bacteria in food environments. In: Brieler, A., (Ed.), Foodborne bacterial pathogens, Springer, New York, USA, 247–264.
- Davies, E.A., Bevis, H.E., Potter, R., Harris, J., Williams, G.C., Delves-Broughton, J., 1998. The effect of pH on the stability of nisin solution during autoclaving. Lett. Apply. Microbiol. 27, 186–187.
- Davies, E.A., Bevis, H.E., Delves-Broughton, J., 1997. The use of the bacteriocin nisin, as a preservative in ricotta-type cheeses to control the food-borne pathogen *Listeria monocytogenes*. Lett. Apply. Microbiol. 24, 343–346.
- Delves-Broughton, J., 2008. Use of the natural food preservatives, nisin and natamycin, to reduce detrimental thermal impact on product quality. In: Richardson, P. (Ed.), In-pack processed foods: Improving quality. Woodhead Publishing, Sawston, UK, pp. 319–337.

<https://doi.org/10.1533/9781845694692.4.319>

- FDA , 1988. Nisin preparation: Affirmation of GRAS status as a direct human food ingredient. Food and Drug Administration (Federal Register April 6, 1988), 53, 11247.
- Ferreira, M.A.S.S., Lund, B.M., 1996. The effect of nisin on *Listeria monocytogenes* in culture medium and long-life cottage cheese. Lett. Appl. Microbiol. 22, 433–438. <https://doi.org/10.1111/j.1472-765X.1996.tb01197.x>
- Fowler, G.G., Jarvis, B., Tramer, J., 1975. The assay of nisin in foods. In: Board, R.G., Lovelock, D.W. (Eds.), Some Methods for Microbiological Assay. Academic Press, New York, pp. 91–105.
- Fuselli, F., Guarino, C., La Mantia, A., Longo, L., Faberi, A., Marianella, R.M., 2012. Multi-detection of preservatives in cheeses by liquid chromatography-tandem mass spectrometry. J. Chromatogr. B Anal. Technol. Biomed. Life Sci. 906, 4–12. <https://doi.org/10.1016/j.jchromb.2012.07.035>
- Glass, K., A., Mu, M., LeVine., B., Rossi, F., 2017. Inhibition of *Clostridium botulinum* in model reduced-sodium pasteurized prepared cheese products. J. Food Prot. 80, 1478-1488. <https://doi.org/10.4315/0362-028X.JFP-17-027>
- Hara, H., Ohashi, Y., Sakurai, T., Yagi, K., Fujisawa, T., Igimi, S., 2008. Effect of nisin (Nisaplin) on the growth of *Listeria monocytogenes* in karashi-mentaiko (red-pepper seasoned cod roe). J. Food hyg. Soc. Japan, 4 173-177.
- Hereu, A., Dalgaard, P., Garriga, M., Aymerich, T., Bover-Cid, S., 2014. Analysing and modelling the growth behaviour of *Listeria monocytogenes* on RTE cooked meat products after a high pressure treatment at 400 MPa. Int. J. Food Microbiol. 186, 84–94. <https://doi.org/10.1016/j.ijfoodmicro.2014.06.020>
- ISO, 2009. Cheese - Determination of nisin A content by LC-MS and LC-MS-MS. ISO/TS 27106 IDF/RM 217 2002, 1–14.
- Jewell, K., 2012. Comparison of 1-step and 2-step methods of fitting microbiological models. Int. J. Food Microbiol. 160, 145–161.
- Katla, T., Naterstad, K., Vancanneyt, M., Swings, J., Axelsson, L., 2003. Differences in susceptibility of *Listeria monocytogenes* strains to sakacin P., sakacin A, pediocin PA-1 and nisin. Appl. Environ. Microbiol. 4431-4437. <https://doi.org/10.1128/AEM.69.8.4431-4437.2003>
- Khan, A., Vu, K.D., Riedl, B., Lacroix, M., 2015. Optimization of the antimicrobial activity of

- nisin, Na-EDTA and pH against gram-negative and gram-positive bacteria. *LWT - Food Sci. Technol.* 61, 124–129. <https://doi.org/10.1016/j.lwt.2014.11.035>
- Ko, K.Y., Park, S.R., Lim, H.S., Park, S.J., Kim, M., 2016. Improved pretreatment method for determination of nisins A and Z and monitoring in cheese using liquid chromatography-tandem mass spectrometry. *Food Anal. Methods* 9, 122–130. <https://doi.org/10.1007/s12161-015-0158-y>
- Le Marc Y., Huchet V., Bourgeois C.M., Guyonnet J.P., Mafart P., Thuault D., 2002. Modelling the growth kinetics of *Listeria* as a function of temperature, pH and organic acid concentration. *Int J Food Microbiol* 73:219–237. [https://doi.org/10.1016/S0168-1605\(01\)00640-7](https://doi.org/10.1016/S0168-1605(01)00640-7)
- Martinez-Rios, V., Gkogka, E., Dalgaard, P., 2019a. New term to quantify the effect of temperature on pH_{min} -values used in cardinal parameter growth models for *Listeria monocytogenes*. *Frontiers in Microbiology*, 84, 103255. <https://doi.org/10.3389/fmicb.2019.01510>
- Martinez-Rios, V., Jørgensen, M.Ø., Koukou, I., Gkogka, E., Dalgaard, P., 2019b. Growth and growth boundary model with terms for melting salts to predict growth responses of *Listeria monocytogenes* in spreadable processed cheese. *Food Microbiol.*, 10, 1510. <https://doi.org/10.1016/j.fm.2019.103255>
- Martínez, B., Rodríguez, A., 2005. Antimicrobial susceptibility of nisin resistant *Listeria monocytogenes* of dairy origin. *FEMS Microbiol. Lett.* 252, 67–72. <https://doi.org/10.1016/j.femsle.2005.08.025>
- Neetoo, H., Ye, M., Chen, H., 2008. Potential antimicrobials to control *Listeria monocytogenes* in vacuum-packaged cold-smoked salmon pâté and fillets. *Int. J. Food Microbiol.* 123, 220–227. <https://doi.org/10.1016/j.ijfoodmicro.2008.02.001>
- Nilsson, L., Huss, H.H., Gran, L., 1997. Inhibition of *Listeria monocytogenes* on cold-smoked salmon by nisin and carbon dioxide atmosphere. *Int. J. Food. Microbiol.* 38, 217-227.
- NMKL, 2009. Validation of chemical analytical methods (NMKL Procedure No. 4, 2009).
- Østergaard, N.B., Eklöw, A., Dalgaard, P., 2014. Modelling the effect of lactic acid bacteria from starter- and aroma culture on growth of *Listeria monocytogenes* in cottage cheese. *Int. J. Food Microbiol.* 188, 15–25. <https://doi.org/10.1016/j.ijfoodmicro.2014.07.012>
- Oshima, S., Hirano, A., Kamikado, H., Nishimura, J., Kawai, Y., Saito, T., 2014. Nisin A extends the shelf life of high-fat chilled dairy dessert, a milk-based pudding. *J. Appl. Microbiol.* 116, 1218-1228. <https://doi.org/10.1111/jam.12454>
- Papagianni, M., Avramidis, N., Filioussis, G., Dasiou, D., Ambrosiadis, I., 2006. Determination of bacteriocin activity with bioassays carried out on solid and liquid substrates: assessing the factor "indicator microorganism". *Microb. Cell Fact.* 5, 1-14. <https://doi.org/10.1186/1475->

2859-5-30.

- Parente, E., Giglio, M.A., Ricciardi, A., Clementi, F., 1998. The combined effect of nisin, leucocin F10, pH, NaCl, EDTA on the survival of *Listeria monocytogenes* in broth. Int. J. Food Microbiol. 40, 65-75. [https://doi.org/10.1016/S0168-1605\(98\)00021-X](https://doi.org/10.1016/S0168-1605(98)00021-X)
- Pongtharangkul, T., Demirci, A., 2004. Evaluation of agar diffusion bioassay for nisin quantification. Appl. Microbiol. Biotechnol. 65, 268-272.
- Popowska, M., Olszak, M., Markiewicz, Z., 2006. Susceptibility of *Listeria monocytogenes* strains isolated from dairy products and frozen vegetables to antibiotics inhibiting murein synthesis and to disinfectants. Polish J. Microbiol. 4, 279-288.
- Razavilar, V. and Genigeorgis, C., 1998. Prediction of *Listeria* spp. growth as affected by various levels of chemicals, pH, temperatures and storage time in a model broth. Int. J. Food Microbiol., 40, 149-157. [https://doi.org/10.1016/s0168-1605\(98\)00014-2](https://doi.org/10.1016/s0168-1605(98)00014-2)
- Redmond, E.C., Griffith, C.J., 2003. Consumer food handling in the home: A review of food safety studies. J. Food Prot. 66, 130–161. <https://doi.org/10.4315/0362-028X-66.1.130>.
- Ross, T., 1996. Indices for performance evaluation of predictive models in food microbiology. J. Appl. Bacteriol. 81, 501–508.
- Ross and Dalgaard, 2004. Chapter 3. Secondary Models. In: McKellar, R.C., Lu, X (Ed.), Modelling microbial responses in food. CRC Press, Boca Raton, pp. 110-119.
- Rogers, A.M., Montville, T.J., 1991. Improved agar diffusion assay for nisin quantification. Food Biotechnol. 5, 161-168.
- Rosso, L., Bajard, S., Flandrois, J.P., Lahellec, C., Fournaud, J., Veit, P., 1996. Differential growth of *Listeria monocytogenes* at 4 and 8°C: Consequences for the shelf life of chilled products. Journal of Food Protection 59, 944–949. <https://doi.org/10.4315/0362-028X-59.9.944>
- Ruiz, A., Williams, S.K., Djeri, N., Hinton, A., Rodrick, G.E., 2010. Nisin affects the growth of *Listeria monocytogenes* on ready-to-eat turkey ham stored at four degrees Celsius for sixty-three days. Poultry Science, 89, 353-358. <https://doi.org/10.3382/ps.2008-00503>
- Schillinger, U., Becker, B., Vignolo, G., Holzapfel, W.H., 2001. Efficacy of nisin in combination with protective cultures against *Listeria monocytogenes* Scott A in tofu. Int. J. Food Microbiol. 71, 159-168. [https://doi.org/10.1016/S0168-1605\(01\)00612-2](https://doi.org/10.1016/S0168-1605(01)00612-2)
- Schneider, N., Werkmeister, K., Pischetsrieder, M., 2011. Analysis of nisin A, nisin Z and their degradation products by LCMS/MS. Food Chem. 127, 847–854. <https://doi.org/10.1016/j.foodchem.2011.01.023>
- Smith, M.L., Draper, L., Hazelhoff, P.-J., Cotter, P.D., Ross, R.P., Hill, C., 2016. A bioengineered

nisin derivate, M21A, in combination with food grade additives eradicates biofilms of *Listeria monocytogenes*. Front. Microbiol. 7, 1939. <https://doi.org/10.3389/fmicb.2016.01939>

Sterling, H.J., Batchelor, J.D., Wemmer, D.E., Williams, E.R., 2010. Effects of buffer loading for electrospray ionization mass spectrometry of a noncovalent protein complex that requires high concentrations of essential salts. J. Am. Soc. Mass Spectrom. 21, 1045-1049. <https://doi.org/doi:10.1016/j.jasms.2010.02.003>.

Taylor, T.M., 2009. Efficacy of novel food antimicrobial combinations for control of *Listeria monocytogenes* for preservation of ready-to-eat (RTE) products. Research Report, Department of Animal Science, Texas A&M. <https://www.pork.org/research/efficacy-of-novel-food-antimicrobial-combinations-for-control-of-listeria-monocytogenes-for-preservation-of-ready-to-eat-rte-products/> (accessed 07.12.18).

Thomas, L.V., Delves-Broughton, J., 2005. Nisin. In: Davidson, P.M., Sofos, J.N., Branen, A.L. (Eds.), Antimicrobials in foods. CRC, Taylor & Francis Group, Boca Raton, pp. 237-274.

Thomas, L. V, Wimpenny, J.W.T., 1996. Investigation of the effect of combined variations in temperature, pH, and NaCl concentration on nisin inhibition of *Listeria monocytogenes* and *Staphylococcus aureus*. Appl. Environ. Microbiol. 52, 2006–2012.

Tramer, J., Fowler, G.G., 1964. Estimation of nisin in foods. J. Sci. Food Agric. 15, 522–528. <https://doi.org/http://0-dx.doi.org.lib.exeter.ac.uk/10.1111/j.1530-0277.2009.00957.x>

Table 1. Comparison of LC-MS/MS protocols from the present study and from ISO (2009).

	Present study	ISO (2009)
Equipment	UHPLC (Dionex Thermo scientific)	LC-MS/MS
	EVOQ Elite ^{TM} triple quadrupole MS(Bruker)	
Column	Zorbax 300 SB-C18 (2.1 x 50 mm)	PLRP-S (2.1 x 150 mm)
	(Agilent Technologies)	(Polymer Lab Ltd.)
Extraction solvent	0.1% formic acid, methanol/water (1:1), pH 2.0 with HCL	Formic acid/water (1:200)
Parent ion	839	672
Fragment ions	1081, 811	811

Table 2. Recovery and precision for determination of nisin A by LC-MS/MS and bioassay.

Method	Matrix	Conc. of nisin A (ppm)	Recovery (%)	CV (%)
LC-MS/MS	Processed cheese (PC)	0.1	99	13
		0.5	108	2
		1	110	4
	Spreadable processed cheese (SPC)	0.1	83	14
		0.5	95	11
		1	100	9
Bioassay	Spreadable processed cheese (SPC)	0.1	53 ^a	16 ^a
		0.5	94	35 ^a
		1	93	33 ^a

^a Result outside acceptability range being: CV < 16 % and recovery within 80 - 110%

Table 3. Experimentally determined minimum inhibitory concentration (MIC) values of nisin A for different *L. monocytogenes* strains

Strain	pH	Temp. (°C)	μ_{ref}^* , h ⁻¹ (AVG $\pm$ SE)	MIC, ppm (AVG $\pm$ SE)	Duration of experiments (days)
LM 4	6.5	7	0.205 $\pm$ 0.01	4.9 $\pm$ 0.4	20
LM 4R	6.5	7	0.081 $\pm$ 0.00	24.9 $\pm$ 3.1	14
LM13	6.5	7	0.193 $\pm$ 0.01	8.1 $\pm$ 0.7	14
<i>Lm-mix</i> ^a	6.5	7	0.225 $\pm$ 0.00	9.1 $\pm$ 0.3	5
LM4R	6.5	20	0.406 $\pm$ 0.01	27.0 $\pm$ 3.4	3
6	6.5	20	0.455 $\pm$ 0.02	6.4 $\pm$ 1.0	13
LM13	6.5	20	0.474 $\pm$ 0.01	14.1 $\pm$ 0.9	20
<i>Lm-mix</i> ^a	6.5	20	0.527 $\pm$ 0.01	15.5 $\pm$ 0.6	14
<i>Lm-mix</i> ^a	6.0	20	0.431 $\pm$ 0.01	8.8 $\pm$ 1.2	4
LM 4R	5.5	20	0.353 $\pm$ 0.01	3.3 $\pm$ 0.1	15
612	5.5	7	0.138 $\pm$ 0.00	0.55 $\pm$ 0.05	64
612	5.5	20	0.417 $\pm$ 0.02	0.75 $\pm$ 0.04	13
<i>Lm-mix</i> ^a	5.5	20	0.415 $\pm$ 0.02	0.55 $\pm$ 0.08	11

^aThe cocktail *Lm-mix* included the isolates SLU 92, 612, LM 19 and 6.

Table 4. Storage conditions, product characteristics and estimated growth parameters from fitting the logistic model with delay (eq. (3)) to the *L. monocytogenes* counts on processed cheese obtained from challenge tests.

CT _b	N _c	Storage Temp. (°C)	Product characteristics (Avg.±SD) ^a							Growth parameter values (Avg. ±SD) ^a				
			pH	a _w	Organic acids in water phase (ppm)			P1 ^d in water phase (%) ^d	Residual nisin A (ppm)	t _{lag} (h)	Log N ₀ (Log cfu/g)	Log N _{max} (Log cfu/g)	μ _{max} (h ⁻¹)	Duration of experiments (days)
					Lactic acid	Acetic acid	Citric acid							
1	3	10.0±0.	5.9±0.	0.981±0.00	9,415±29	2,471±60	5,030±26	0.9±N	0.68±0.0	63.1±16	0.8±0.	4.2±0.3	0.027±0.0	21
2	3	15.0±0.	6.0±0.	0.983±0.00	9,132±14	1,908±68	3,766±17	0.9±N	0.68±0.0	0.0±0.0	0.9±0.	7.1±0.1	0.050±0.0	19
3	3	19.5±0.	5.9±0.	0.982±0.00	9,345±	2,384±2	4,947±47	0.9±N	0.68±0.0	0.0±0.0	1.0±0.	6.5±0.1	0.074±0.0	16
4	3	10.0±0.	6.0±0.	0.983±0.00	9,869±	2,305±81	5,070±28	0.8±N	0.55±0.0	15.2±22	1.0±0.	4.6±0.2	0.024±0.0	21
5	3	15.0±0.	6.1±0.	0.981±0.00	9,364±	2,317±11	4,710±11	0.8±N	0.55±0.0	0.0±0.0	1.0±0.	5.4±0.1	0.045±0.0	21
6	3	19.5±0.	6.1±0.	0.982±0.00	9,679±40	2,397±10	5,234±44	0.8±N	0.59±0.0	0.0±0.0	1.1±0.	5.6±0.1	0.063±0.0	16
7	3	3.6±0.3	5.5±0.	0.973±0.00	1,101±33	< LOD	19,395±504	2.0±N	4.21±0.1	ND ^f	1.9±0.	<1.0±0.0	0.000±0.0	33
8	3	3.6±0.3	5.7±0.	0.972±0.00	1,109±33	< LOD	18,190±432	2.0±N	5.28±0.1	ND	2.0±0.	2.3±0.2	0.000±0.0	35
9	3	3.6±0.3	5.8±0.	0.975±0.00	1,152±90	< LOD	17,471±592	2.0±N	3.35±0.1	ND	2.1±0.	<1.0±0.0	0.000±0.0	33
10	3	7.1±0.3	5.9±0.	0.974±0.00	1,028±68	< LOD	17,585±1,1	2.0±N	0.88±0.6	211±6.3	1.4±0.	10.5±0.	0.023±0.	35
11	3	7.1±0.3	5.4±0.	0.973±0.00	877±17	< LOD	20,864±438	2.0±N	3.56±0.1	ND	1.2±0.	<1.0±0.0	0.000±0.0	32
12	3	7.1±0.3	5.7±0.	0.976±0.00	958±9	< LOD	17,172±699	2.0±N	4.38±0.2	ND	1.6±0.	<1.0±0.0	0.000±0.0	35
13	3	11.3±0.	5.8±0.	0.970±0.00	1,067±73	< LOD	17,721±235	2.0±N	0.98±0.0	31.6±4.	1.7±0.	8.7±0.1	0.058±0.	18
14	3	11.3±0.	5.6±0.	0.971±0.00	1,251±47	< LOD	18,660±442	2.0±N	1.07±0.1	27.1±11	1.2±0.	8.5±0.1	0.046±0.	21
15	3	11.3±0.	5.6±0.	0.970±0.00	869±23	< LOD	18,340±710	2.0±N	0.56±0.0	0.0±0.0	0.8±0.	8.2±0.1	0.040±0.	21

^a Avg: average; SD: standard deviation.^b Challenge test. Challenge tests 1-6 performed using processed cheese (PC) and 7-15 with spreadable processed cheese (SPC).^c Number of growth curves per experiment.^d P1: orthophosphate salt.^e NA, not available. Information not provided by Eurofins.^f ND: not determine.

Table 5. Comparison of observed and predicted growth and no-growth responses of *L. monocytogenes* in processed cheese containing nisin^a.

Models and responses ^a	Predicted responses by three available models including the effect of nisin A					
	Parente et al. (1998); Probability of survival in 1 ml		Bouttefroy et al. (2000); Viable population log ₁₀ cfu/ml		Boziaris and Nychas (2006); Probability of growth	
	pH, NaCl, EDTA, nisin		pH, NaCl, nisin		T, a _w , pH, nisin	
	Added nisin A ^b	Residual nisin A ^c	Added nisin A ^b	Residual nisin A ^c	Added nisin A ^b	Residual nisin A ^c
Correct (%)	14	3	86	86	14	100
Fail-safe (%)	0	11	0	11	0	0
Fail-dangerous (%)	86	86	14	0	86	0

^a Observed responses included 45 kinetics of which 30 showed growth and 15 showed no-growth.

^b Added nisin A: 11.2-12.0 ppm.

^c Nisin A measured by LC-MS/MS: 0.59-5.28 ppm.

Table 6. Growth kinetic parameters of *L. monocytogenes* in processed cheese containing nisin A and goodness of fit indices for the two-steps (eq. (S3); eq. (4)) and the global (one-step, eq. (5)) regression approaches.

Modelling approaches	Growth and growth boundary model	RLT ^a secondary model		Log (N _{max} , cfu/g)	Goodness of fit	
	μ_{ref} (h ⁻¹)	<i>k1</i>	<i>k2</i>		RSS ^b	RMSE ^c
Classical two-step approach	0.510	0.0	1957	6.60	465	1.21
Global (one-step) regression	0.629	0.0	2087	6.60	393	0.97

^a RLT: relative lag time

^b RSS: residual sum of squares

^c RMSE: root mean square error

Table 7. Observed and predicted growth and no-growth responses for *L. monocytogenes*.

Observed			Predicted		
n ^a	Growth	No growth	Assuming no added Nisin A	Residual nisin A	Nisin A added to formulation
45	30	15	0 ppm	0.59-5.28 ppm	11.2-12 ppm
	Correct (%)		86	100	14
	Fail-safe (%)		14	0	0
	Fail-dangerous (%)		0	0	86

^aNumber of growth curves: a total of 15 challenge tests with triplicate measurements.

Figure 1. Chromatogram obtained after LC-MS/MS analysis of cheese with non-spiked (top) and spiked (bottom) with nisin A (0.5 mg/kg).

Figure 2. Effect of nisin A on maximum specific growth rates (μ_{max} , h^{-1}) of *L. nonocytogenes* LM13 (a) and LM4R (b) in BHI broth at 20°C and pH 6.5. MIC-values were determined by fitting eq. (2) to the observed data (●). Solid and dashed lines represent the fitted values and confidence intervals (95%), respectively.

Figure 3. Effect of pH on nisin A MIC-values for *L. monocytogenes* as determined in the present study and from literature. Data for a cocktail of *L. monocytogenes* strains [*Lm-mix*: ST U 22, 612, LM19, 6 (×)], individual strains [LM 4 (Δ), LM 4R (□), LM 13 (○), 6 (◇), 612 (+)], Taylor et al., (2002) (□) and Ferreira and Lund (1996) (○). Solid line (—) and dashed line (---) represent, respectively, the fitted model (eq. (5)) and confidence interval (95%)

Figure 4. Effect of storage temperature on observed and fitted relative lag time (RLT) values for *L. monocytogenes* growing in processed cheese containing nisin A. The two-step approach (solid line) and the global regression (dashed line) were used to fit RLT-values (Table 6).

Figure 5. Growth of *L. monocytogenes* (○) in commercial (a) and laboratory made (b) processed cheese containing nisin and stored at 19.5°C or 7.3°C, respectively (CT 3, CT 10; Table 4). Solid lines indicate predictions obtained with the model developed in the present study (eq. (6) and Table 6) and dashed lines indicate predictions obtained with the model of Martinez-Rios et al. (2019a) including (eq. (S1)).

Highlights

- New model to predict growth of *Listeria monocytogenes* in processed cheese containing nisin
- Model including the effect of temperature, pH, a_w , lactic acid, acetic acid, citric acid, orthophosphate and nisin
- LC-MS/MS method was developed and validated to quantify nisin A present in cheese