

Estimation of external infection pressure and salmon-louse population growth rate in Faroese salmon farms

Kragesteen, Tróndur J.; Simonsen, Knud; Visser, André W.; Andersen, Ken H.

Published in: Aquaculture Environment Interactions

Link to article, DOI: 10.3354/aei00386

Publication date: 2021

Document Version Peer reviewed version

Link back to DTU Orbit

Citation (APA):

Kragesteen, Ť. J., Simonsen, K., Visser, A. W., & Andersen, K. H. (2021). Estimation of external infection pressure and salmon-louse population growth rate in Faroese salmon farms. *Aquaculture Environment Interactions*, *13*, 21-32. https://doi.org/10.3354/aei00386

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Estimation of external infection pressure and salmon-louse population growth rate in Faroese salmon farms

Tróndur J. Kragesteen^{a,b}, Knud Simonsen^{c,a}, André W. Visser^b, Ken H. Andersen^b

^aFiskaaling - Aquaculture Research Station of the Faroes, við Áir, 430 Hvalvík, Faroe Islands ^bVKR Centre for Ocean Life, National Institute of Aquatic Resources, Technical

^c Fróðskaparsetur Føroya - University of the Faroe Islands, J.C. Svabos gøta 14, 100 Tórshavn, Faroe Islands

Abstract

Managing salmon lice (*Lepeophtheirus salmonis*) outbreaks is a crucial part of salmon aquaculture in sea cages. Treatment management strategies can be optimized with the aid of salmon-louse population dynamic models. These models, however, need to be calibrated and validated with biological meaningful parameters. Here, based on a time-series of lice data, we estimated two essential model parameters: The external infection pressure and the salmonlouse population growth rate for each active salmon farm site in the period 2011 to 2018 in the Faroe Islands. External infection pressure was found to vary between farm sites and ranged on average from 0.002 to 0.1 lice the total number of gravid lice in the Faroese farm network. Salmon-louse population growth rates were found to vary between farm sites and ranged on average from 1.7 to 5.4 % d⁻¹. These model parameter estimates are crucial in developing a salmon-louse population dynamic model for the Faroe Islands and the method to estimate these parameters may be applicable in other aquaculture regions.

Keywords: Salmon Aquaculture, Salmon lice, Modelling

Email address: trondurk@fiskaaling.fo (Tróndur J. Kragesteen)

1 1. Introduction

Managing sea lice on a regional or national scale is a crucial part of 2 modern sea-based salmon aquaculture. In the northern hemisphere the most 3 important sea lice impacting salmon aquaculture is the ectoparasite Lepeophtheirus salmonis, also know as salmon louse. The salmon louse is a naturally 5 occurring parasite on salmonid fish and feeds on the mucus, skin and blood 6 of it's host (Pike & Wadsworth 1999). At high densities, lice can cause phys-7 ical damage to their host and expose them to secondary infections as well as causing stress and osmotic regulatory imbalance (Pike & Wadsworth 1999). 9 As the salmon farming industry has grown, so has the number of salmon-10 louse host's providing favourable conditions for the parasite. At sufficiently 11 high host densities salmon lice can develop into an epidemic (Frazer et al. 12 2012) negatively affecting both salmon farms and wild salmonid stocks (Krkošek 13 et al. 2006, Kristoffersen et al. 2018). Understanding the dynamics of louse 14 population growth and outbreaks and how lice disperse has obvious potential 15 to reduce treatment frequency and fish mortality on both farmed and wild 16 salmon stocks, increasing economic output and decreasing ecological impact. 17 Numerical models describing salmon-louse population growth have been 18 developed based on Anderson et al. (1979) type host-parasite models (Krkošek 19 et al. 2010, Frazer et al. 2012) and delayed stage structured models (Revie 20 et al. 2005, Stien et al. 2005, Robbins et al. 2010, Gettinby et al. 2011, 21 Adams et al. 2015, Kragesteen et al. 2019). Louse population growth rate 22 is determined by two processes: external and internal infection. The exter-23 nal infection pressure is determent by lice arriving from other salmon farms 24 and wild salmonid stocks. The internal infection pressure is determent by 25 the production of lice on the farm. If internal infection is high enough it 26 will eventually lead to a exponentially growing salmon lice population. In 27 the initial phase of salmon production, the external infection pressure is the 28 dominating process of the population growth rate (Kristoffersen et al. 2014). 29 Internal infection becomes the dominating process as the lice population in-30 creases (Krkošek et al. 2010). 31

The external infection pressure has been estimated using sentinel cages in Norway, Scotland and Iceland (Bjørn et al. 2011, Pert et al. 2014, Sandvik et al. 2016, Karbowski et al. 2019). A sentinel cage is typically deployed 2-3 weeks at a time over a period of months and for each deployment lice are

recorded. External infection pressure reported in these studies was found to 36 vary significantly depending on the level of gravid lice in the area. Bjørn 37 et al. (2011) found the external infection to be up to 2.3 lice salmon⁻¹ in the 38 Romsdalsfjord system, Norway, with a 14 days deployment time. Sandvik 39 et al. (2016) found up to 20 lice salmon⁻¹ in Hardangerfjord, Norway, with 14-40 21 days deployment time and Pert et al. (2014) found close to 15 lice salmon⁻¹ 41 in Loch Shieldaig, Scotland with 7 days deployment time. In Arnarfjörður, 42 Iceland, which only contained 2 salmon farms Karbowski et al. (2019) found 43 only 0.022 lice salmon⁻¹ when sampling for ≈ 21 days. 44

Salmon-louse population growth rates have been reported for salmon farms in the Broughton Archipelago, Canada and the Faroe Islands (Krkošek et al. 2010, Patursson et al. 2017). These estimates assume exponential growth and Krkošek et al. (2010) reported two farms having 0.9 and 4.8 %/d while Patursson et al. (2017) reported growth rates to range from 0.9 to 4.1 $\% d^{-1}$ for several Faroese farms.

Here, we describe the general development of salmon lice abundance in Faroe Islands from 2011 to 2018 based on an extensive time series of sea lice counts. Further, we estimated external infection pressure and salmon-louse population growth rate on a per-farm basis using an alternative approach described in the method section. These parameter estimates are essential for the development of a salmon-louse population dynamic models which can aid in salmon lice management strategies.

58 2. Methods

59 2.1. Lice data and salmon number

Lice data is based on a the Faroese lice count program which started in 60 2009 where the Faroese government mandated farmers to count lice every 61 14 days from 1 May to 31 December and once a month from 1 January to 62 30 April. However, lice counts were not performed on all farms until after 63 2012. From 2009 to 2016 a minimum of 10 fish were counted from 4 cages. 64 The farmer choose 2 cages where one being the first stocked cage and the 65 second cage was, based on prior experience, estimated to have the highest sea 66 lice load. The two other cages were chosen at random between cages which 67 had not been counted previously. After 2016 a minimum of 10 fish were 68 counted for all cages at each farm site every 14 days all year (Anon 2016). In 69 year 2011 counting was not performed at 3 farms and at 3 farms lice counts 70 started 3-6 months into the year 2011. In addition ally salmon numbers for 71

each farm site was provided by the Faroese aquaculture companies for theperiod between 2011 and 2018.

Both L. salmonis and Calique elongatus have been recorded, however, 74 we will exclusively focus on salmon louse as they have by far the largest 75 economic impact on the salmon industry (Boxaspen 2006). The life cycle of 76 L. salmonis consists 8 stages (Schram 1993, Hamre et al. 2013). Non-feeding 77 planktonic nauplii larvae hatch from gravid female egg strings and moults 78 into nauplius II, and subsequently, into the third, and infective, stage, as a 79 copepodite. After attachment, copepodites moult into immobile chalimus I 80 and start feeding. The chalimus phase consists of 2 stages (chalimus I and 81 II), where the latter moults into the first mobile pre-adult stage. After the 82 pre-adult stages (pre-adult I and II), the louse moults into the adult and final 83 stage. 84

Salmon louse counts categorise these life-stages into three groups: immobile (chalimus I and II), mobile, and gravid female. *L. salmonis* and *C. elongatus* are not distinguished from each other in the immobile group and these counts are therefore discarded and only mobile and gravid female lice are considered. Mobile lice counts include the pre-adult stages (male and female) and adult male stage while gravid lice only include this stage.

91 2.2. Treatments

Treatment data was gathered from the active Faroese farming companies 92 since 2011. There have been several kinds of treatments performed, which 93 here are organized into four groups: Medical oral (SLICE and Diffuben-94 zuron), medicinal bath (Hydrogen peroxide, Salmosan, Alpbamax, Betamax, 95 Pyretroid and Azametiphos) and mechanical (freshwater bath, hydrolicer, 96 optilicer, termolicer or flushing). A "treatment event" was defined as a given 97 kind of treatment performed at a given farm which had not been performed 98 within the previous 7 days. Number of treated cages was not resolved. 99

100 2.3. Sea Water Temperature

Sea water temperature was measured by the Faroe Marine Research Institute and available at the online data service, www.envofar.fo. Sea water temperature is measured at 3 m depth at Oyrargjógv (62°07'N, 7°10'W) which is located in a tidally well mixed strait, and thus representative for a relative large geographical region (Fig. 1).

106 2.4. Calculating total lice

The total number of gravid and mobile lice was estimated for each farm is i by linear interpolating between the day of the counts to obtain the daily values $l_i(t)$. Knowing the number of salmon in each farm per day $F_i(t)$ the total number of lice in the region on a given day, t, is obtained from:

$$L_{\rm tot}(t) = \sum_{i=1}^{n} l_i(t) F_i(t).$$
 (1)

¹¹¹ where n is number of farms in the system.

112 2.5. The Salmon-louse Model

To estimate the external infection pressure and salmon-louse population growth rate, we used a series of delay differential equations (Revie et al. 2005):

$$\frac{\mathrm{d}\rho_1(t)}{\mathrm{d}t} = \beta(t) - \beta(t-t_1)e^{-\mu_1 t_1} - \mu_1\rho_1(t)$$
(2)

$$\frac{\mathrm{d}\rho_2(t)}{\mathrm{d}t} = \beta(t-t_1)e^{-\mu_1 t_1} - \beta(t-t_1-t_2)e^{-\mu_1 t_1-\mu_2 t_2} - \mu_2\rho_2(t) \tag{3}$$

$$\frac{\mathrm{d}\rho_3}{\mathrm{d}t}(t) = \beta(t - t_1 - t_2)e^{-\mu_1 t_1 - \mu_2 t_2} - \beta(t - t_1 - t_2 - t_3)e^{-\mu_1 t_1 - \mu_2 t_2 - \mu_3 t_3} - \mu_3 \rho_3(t) \quad (4)$$

$$\frac{\mathrm{d}\rho_4(t)}{\mathrm{d}t} = \beta(t - t_1 - t_2 - t_3)e^{-\mu_1 t_1 - \mu_2 t_2 - \mu_3 t_3} - \mu_4, \rho_4(t), \tag{5}$$

where ρ_{1-4} represent the male and female lice at the chalimus, pre-adult, adult and sexually mature stages, respectively, and μ_{1-4} and t_{1-4} represent mortality and development times. The amount of attached larvae $\beta(t)$ is defined in Kragesteen et al. (2019) as:

$$\beta(t) = q\eta \rho_4(t - t_e)s(\rho_4(t - t_e)) + L_0, \tag{6}$$

where q is the amount of viable larvae per day per sexually mature lice which includes connectivity and larvae production rate and η is the proportion of female lice. L_0 is the external infection pressure and t_e is the time it takes larvae to reinfect a host and is assumed to be 5 days (Stien et al. 2005). $s(\rho_4(t))$ is mate limitation or an Allee effect which states that fertilisation success is close to zero at near zero lice abundances and close to 100 % at around 2 gravid lice salmon⁻¹ (Krkošek et al. 2012, Stormoen et al. 2013,
Kragesteen et al. 2019). The initial conditions are given by:

$$\rho_1(0) = 0 \text{ for } t = 0 \tag{7}$$

$$\rho_{j+1}(t) = 0 \text{ for } t \le t_j \text{ where } j \in \{1, 2, 3\}.$$
(8)

where t_j is the development time for each stage and salmon are stocked in sea cages at t = 0.

130 2.6. External infection

The external infection pressure in this study includes, as mentioned, salmon louse larvae from other farms and from the environment or natural background infection. To calculate external infection pressure we rewrite Eqs. 2-6 by summing all mobile stages $(\rho_2(t)+\rho_3(t)+\rho_4(t))$, which corresponds to the mobile and gravid lice counts, ρ_m . Therefore:

$$\frac{\mathrm{d}\rho_m(t)}{\mathrm{d}t} = \frac{\mathrm{d}\rho_2(t)}{\mathrm{d}t} + \frac{\mathrm{d}\rho_3(t)}{\mathrm{d}t} + \frac{\mathrm{d}\rho_4(t)}{\mathrm{d}t} = \beta(t-t_1)e^{-\mu_1 t_1} - \mu_2\rho_2(t) - \mu_3\rho_3(t) - \mu_4\rho_4(t).$$
(9)

If we assume that $\mu_m \approx \mu_2 \approx \mu_3 \approx \mu_4$ and the practical implementation of this assumption is $\mu_m = \frac{(\mu_2 + \mu_3 + \mu_4)}{3}$ Eq. 9 can be written as:

$$\frac{\mathrm{d}\rho_m(t)}{\mathrm{d}t} = \beta(t-t_1)e^{-\mu_1 t_1} - \mu_m \rho_m(t).$$
(10)

We argue that in the first 150 days of the production cycle there is virtually 138 no internal dynamics, meaning we assume no internal viable larvae and/or no 139 females are fertilized and set $q \cdot s$ in Eq. 6 to zero. There are two reasons for 140 assuming this: First, at 11 °C (maximum Faroese shelf water temperature) 141 the first attached lice start releasing larvae after ≈ 50 d (= $t_1 + t_2 + t_3$) 142 and it takes another ≈ 20 days (= $t_e + t_1$) until these lice appear in the 143 lice counts (Table 1, Stien et al. (2005)). Second, due to the Allee effect, 144 where few female lice get fertilized at low lice abundances (Krkošek et al. 145 2012, Stormoen et al. 2013), s is close to zero. Further, gravid lice have been 146 shown to produce fewer eggs in their first pair of egg strings (Heuch et al. 147 2000). With the approximation that $q \cdot s$ is zero, Eq. 10 can be written as: 148

$$\frac{d\rho_m(t)}{dt} = L_0 e^{-\mu_1 t_1} - \mu_m \rho_t(t), \quad \text{for } t < 150 \text{ days}$$
(11)

which with the initial conditions ($\rho_m(t_1) = 0$) has the analytical solution:

$$\rho_m(t) = L_0 \frac{e^{-\mu_1 t_1}}{\mu_m} \left(1 - e^{-\mu_m (t-t_1)} \right)$$
(12)

for $t \ge t_1$. From here the external infection pressure, L_0 , may be isolated to:

$$L_0 = \frac{\mu_m \rho_m(t)}{e^{-\mu_1 t_1} \left(1 - e^{-\mu_m (t-t_1)}\right)}.$$
(13)

The stage duration and mortality are based on the estimates by Stien et al. (2005) (Table 1), where μ_m is the average of the minimum mortalities.

153 2.7. Salmon-louse population growth rate

Salmon-louse population growth rates can be calculated using Eq. 10 and dividing with $\rho_m(t)$:

$$\frac{1}{\rho_m(t)} \frac{\mathrm{d}\rho_m(t)}{\mathrm{d}t} = \frac{\beta_i(t-t_1)}{\rho_m(t)} e^{-\mu_1 t_1} - \mu_m.$$
(14)

¹⁵⁶ Writing Eq. 6 for $\beta(t)$ we get:

$$\frac{\mathrm{d}\ln\left(\rho_m(t)\right)}{\mathrm{d}t} = \left[q\eta s(\rho_4(t-t_e-t_1))\frac{\rho_4(t-t_e-t_1)}{\rho_m(t)} + \frac{L_0}{\rho_m(t)}\right]e^{-\mu_1 t_1} - \mu_m.$$
(15)

All lice counts 150 days after production start are included. The growth rate 157 is calculated by taking the log of salmon lice counts $(\rho_m(t))$ and estimating 158 the slope of a fitted line with 5 consecutive points or counts using a linear 159 regression model MATLAB (2020) with random slope and intercept. Unfor-160 tunately, lice counts are affected by treatments events and/or cleaner fish. 161 To omit these periods we have discarded periods where the slope is negative 162 or has an adjusted $R^2 < 0.6$. See Appendix, Fig. A1 for an example of how 163 external infection pressure and population growth rate was estimated at one 164 farm site. 165

166 3. Results

167 3.1. Total lice

The total number of gravid and mobile lice in the Faroese aquaculture has fluctuated a lot since 2011. The highest number of gravid and mobile lice was in December 2015 with over 35 million and 129 million, respectively
(Fig. 2). The lowest number of gravid lice was under 5 million and recorded
in June 2013 and the lowest for mobile lice was under 5 million in March
2011. The total number of gravid lice has generally been below 20 million
except in the winter of 2011, 2014, 2015 and 2016. Lice numbers in year 2011
may be underestimated as 6 farms were not counted regularly or not at all.

Number of salmon in Faroese waters has been relatively stable and close to 20 million since 2011, while production has increased significantly from 50·10³ tonnes in 2011 to 71·10³ tonnes in 2014 (www.hagstovan.fo), indicating a general increase in harvest or stocking fish size. The maximum recorded number of salmon was 23.4 million in November of 2017 and the minimum was 16.8 million in February 2011 (Fig. 2a).

Number of treatments events per year has steadily increased since 2011 and was over 90 treatment events per year in years 2016-2018. There is no consistent seasonal treatment pattern. Treatment type shifted from chemical to mechanical starting in 2016 and was almost exclusively mechanical in 2018 (Fig. 2b).

Average shelf sea temperature varies consistently between 10-11 °C in September to 6 °C in March (Fig. 2c). There is a significant correlation between temperature and total gravid and mobile lice with a lag of -95 and -74 days having a correlation of 0.49 and 0.32, respectively. In addition, a high correlation (0.95) was found between average annual gravid lice population growth rate and temperature (see Fig. 6).

Overall the total amount of gravid and mobile lice steadily increased 193 from 2011 to 2015 where after it has steadily decreased. This is consistent 194 with changes in regulation in 2016 where treatment threshold was decreased 195 from 2 gravid lice salmon⁻¹ to 1.5 (Faroese Ministry of Foreign Affairs and 196 Trade 2016). We also see a shift from chemical to mechanical treatments and 197 increased treatment frequency in this period. Total lice numbers are gener-198 ally lowest between May and August and highest in December and January. 199 Number of salmon (Table 2) and average temperature in the period 2011 to 200 2018 have stayed relatively constant (Fig. 2) and therefore the total number 201 of lice between years is likely tied to treatment frequency and efficiency. 202

203 3.2. External Infection Pressure

Based on all lice counts in the first 150 days in all production cycles at each farm from 2010-2018 the external infection pressure is estimated on average to range from 0.002 to 0.1 lice salmon⁻¹ d⁻¹. This corresponds to 1 lice per

salmon for every 500 to 10 days (Fig. 3). A total of 30 farms were investigated 207 (Fig. 1). Farm 28 is an outlier and has the highest average external infection 208 pressure, however this estimate is only based on one production cycle in 209 2015. Farm 30 has clearly the lowest average external infection pressure with 210 0.002 lice salmon⁻¹ d⁻¹. In addition to the period of no internal dynamics of 211 150 days (discussed in section 2.6), the external infection is estimated using 212 periods of 75, 100, and 125 days. For many farms the external infection 213 pressure increases with number of days included while for other farms it 214 decreases and for a few farms external infection pressure stays constant, but 215 generally the effect is not significant (Fig. 3). 216

In highly connected farm networks such as Faroe Islands (Kragesteen 217 et al. 2018) we expect external infection pressure to increase with the total 218 amount of gravid lice within the farm network. Therefore, the relationship 219 between external infection pressure and the total number of gravid lice was 220 investigated by performing a linear regression model fit with random slope 221 and intercept for the estimated external infection pressure as a function of 222 the total gravid lice (MATLAB 2020). The mean level of gravid lice was 223 estimated from 15 to 45 days prior to each lice count. The results show that 224 an increase in total gravid lice significantly increased external infection pres-225 sure $(F_{29,844} = 12.7, p < 0.001)$, however, the R^2 was low (0.312) indicating 226 that total number of gravid lice does not explain the variability well (Fig. 4). 227 In addition, for the majority of farms, external infection pressure increases 228 roughly between 0.001 to 0.004 lice salmon⁻¹ d⁻¹ for every million gravid lice 229 in the farm network (Fig 4). Farm no. 30 is an exception having a significant 230 lower external infection pressure likely due to it's isolated location. 231

232 3.3. Population growth rate

Growth rates of lice per salmon was estimated to be between 1.7 and 5.4 %233 d^{-1} on average for all farms (Fig. 5). These estimates are slightly higher but 234 comparable to what Patursson et al. (2017) and Krkošek et al. (2010) found. 235 The estimated growth rates do not separate between internal and external 236 dynamics (Eq. 15), however, we expect that highly self-infectious farms will 237 have a higher growth rate and vise versa. For example, farm 4, 23 and 24 all 238 have relatively low population growth rates, which is expected because they 239 are located in areas with high ventilation due to the tidal currents. Farm 28 240 has the highest measured growth rate likely because the estimate is based on 241 only one production cycle between years 2015-2016 where the total number 242 of gravid lice also was high and/or the growth rate is positively influenced 243

by treatment events as growth rate is typically high for a period right after a treatment.

We also explored the average annual population growth rate of the total 246 number of gravid lice in Faroe Islands from 2011 to 2018 (Fig. 6). This was 247 done by calculating per day population growth rate by log transforming the 248 total number of gravid lice (Eq. 14) and finding the slope of a straight line 249 fitted with 14 and 90 consecutive days. Each calculated growth rate was 250 sorted into day of year and averaged between all years. We see that average 251 population growth rate is negative approximately the first 160-170 days of the 252 year where after the average growth rate turns positive until approximately 253 day 350 in the year. The average annual lice population growth rate over 90 254 d is highly correlated (0.95) with average annual temperature. The average 255 annual lice population growth rate over 14 d oscillates consistently and no 256 clear correlation with temperature is found. 257

258 4. Discussion

From an extensive time-series of lice counts we show how salmon louse 259 have developed in Faroese aquaculture from 2011 to 2018. Further, the av-260 erage external infection pressure and salmon lice growth rate for each active 261 farm site since 2009 was estimated. External infection pressure varies greatly 262 between farms (Fig. 3) which is also expected due to differences in hydrody-263 namic conditions (Patursson et al. 2017) and connectivity between farm sites 264 (Kragesteen et al. 2018). The external infection pressure within farms de-265 pends on the total amount of gravid lice present in the farm network (Fig. 4). 266 The total number of gravid and mobile lice in Faroe Islands is significantly 267 higher compared to a similar salmon aquaculture area (15-18 million salmon) 268 off the coast of mid-Norway (Jevne & Reitan 2019). This coastal area has 269 18 farm sites located between a group of islands and has a similar water 270 temperature range to the Faroe Islands. One difference is that this Nor-271 wegian aquaculture region has synchronized production cycles and the level 272 of salmon lice is therefore effectively reset between each cycle. Comparing 273 the number of treatments between these two areas is not straightforward as 274 treatments are reported differently (number of cages treated in Norway and 275 number of treatment events in the Faroe Islands). However, if we assume that 276 a treatment event on average represents 5 treated cages then about 450 cages 277 are treated per year (90.5) in Faroe Islands in the period from year 2016 to 278 2018 (Table 2). Jevne & Reitan (2019) reported 262, 550 and 102 cages being 279

treated in the 1, 2 and 3 production cycle, respectively, where a production 280 cycle is almost 2 years long. Production cycle 2 had the highest levels of lice 281 and also the highest number of cages treated but this is still a factor lower 2 282 compared to the Faroe Islands. A reason for the higher treatment frequency 283 in the Faroe Islands may be the overlapping production in contrast to syn-284 chronized production in the Norwegian area. This would contradict claims 285 that coordinated fallowing is ineffective (Guarracino et al. 2018). Another 286 reason could be the relatively low treatment threshold of 0.2 to 0.5 gravid lice 287 salmon⁻¹ in Norway (Anon 2012) in contrast to 1.5 to 2 gravid lice salmon⁻¹ 288 in the Faroe Islands which may seem counter-intuitive. However, farms in 289 Norway are forced to treat early to keep lice levels relatively low and thereby 290 earn the benefit of the Allee effect resulting in a overall lower larvae pro-291 duction rate and consequently a lower lice population growth rate (Krkošek 292 et al. 2012, Stormoen et al. 2013, Kragesteen et al. 2019). Third, there is a 293 mean current flow through the Norwegian area and therefore lice may be less 294 retained in contrast to Faroe Islands where shelf water is relatively retained 295 (Kragesteen et al. 2018). Last, we currently have insufficient data for cleaner 296 fish, which may be more widely used in the Norwegian island group resulting 297 in fewer treatments. 298

Salmon-louse levels are typically highest in December/January (Fig. 2). This is likely because the population growth rate of salmon-louse in Faroe Islands is positive until approximately 90 days after the highest sea temperature (September). Highest lice abundance is observed right before the net growth rate turns negative (Fig. 6).

We also speculate that chlorophyll could be a good indicator of lice population growth rate and lice levels. Because high levels of chlorophyll will subsequently lead to high levels of zooplankton which may lead to a higher mortality of planktonic lice larvae. However, we found no clear correlation, maybe due chlorophyll and zooplankton being out of phase or that the chlorophyll samples are from a single location not representative of the general chlorophyll concentration in the Faroe Islands.

If the sentinel cage external infection estimates are converted to lice salmon⁻¹ d⁻¹ using Eq. 2, we find up to 0.17 (Bjørn et al. 2011), 1.16 (Sandvik et al. 2016), 2.16 (Pert et al. 2014) and 0.001 lice salmon⁻¹ d⁻¹ (Karbowski et al. 2019), when minimum mortality of the chalimus stage is assumed (Table 1). The external infection pressure estimates in this study are therefore considerably lower compared to the sentinel cage estimates in Norway and Scotland (Bjørn et al. 2011, Pert et al. 2014, Sandvik et al. 2016) and while

slightly higher than estimates form Iceland in a fjord containing only two 318 salmon farms (Karbowski et al. 2019). One reason for the observed differ-319 ence could be the dilution effect reported by Samsing et al. (2014), because 320 there are relatively few salmon in the sentinel cages there are potentially a 321 lot more lice per salmon compared to a fully operational high salmon density 322 farm cage. External infection pressure estimates from sentinel cages may as a 323 result be much higher compared to our estimates. This dilution effect should 324 be investigated further. 325

External infection pressure was estimated based on the first 150 days of 326 a production cycle as we assume effects of salmon-louse internal dynamics to 327 be low or non-existing in this period. This assumption can be debated in par-328 ticular when water temperature is high (i.e. >11 °C). If internal dynamics are 329 significant in this period there will be an overestimation of the external in-330 fection pressure. However, the difference between including the first 75, 100, 331 125 and 150 days does not significantly affect the average estimated external 332 infection pressure (Fig. 3). An explanation for increasing external infection 333 pressure could be that the self-infection or internal dynamics starts before 334 the 150 days which would cause the estimated external infection pressure to 335 increase with time. Another explanation could be the increased surface area 336 of salmon e.g. if a salmon weighs 200 g when put out to sea, they will increase 337 their weight to about 900 g the first 150 days (Austreng et al. 1987). This 338 increase in weight will increase the salmon surface area from approximately 339 335 cm^2 to 810 cm^2 (O'Shea et al. 2006). Consequently infectious lice larvae 340 will have 2.4 time more area to attach on 150 day after sea stocking. There-341 fore, external infection pressure should be standardized with salmon size, 342 however, these data were not available. An issue with the external infection 343 pressure estimates is that treatments do occur in the first 150 days period 344 and in many farms especially after 2015 had cleaner fish present in their 345 sea cages. This would lead to an underestimation of the external infection 346 parameter L_0 and could cause the external infection pressure to decrease in 347 the 150 day period. In summary the external infection pressure estimates 348 are quite uncertain. Nevertheless, these estimates have a high applied value 349 as they are based on *in situ* lice counts from commercial farms reflecting the 350 actual infection pressure at a operational salmon farm. 351

Here we do not distinguish between external infection pressure caused by larvae production from neighboring farms and the natural background infection from wild salmonid stocks. There is a small wild salmon stock which was introduced and has been maintained since 1940's in four Faroese

rivers (www.laks.fo). The population size and level of infection of other 356 salmon-louse hosts like sea trout and Arctic charr are unknown. In Norway 357 the wild salmon stock is estimated to about 550.000 fish (Anon 2019), and 358 the Faroese wild stock is likely only a small fraction of this and consequently 359 the ratio between the wild stock and the 20 million salmon in the cages is 360 likely small, and therefore it seems reasonable to assume that the infection 361 load from the wild salmonids is low. As a result external infection should 362 predominately be determined by the total number of gravid lice in Faroese 363 salmon farms. Our study shows a significant but low R^2 although varying 364 highly between farms (Fig. 4). The reason for the relatively low correlation 365 could be the stochastic nature of the lice counts procedure, where only 10 366 fish per cage are counted. In addition, treatments and cleaner fish will also 367 negatively influence the correlation. Further, some farms may be strongly 368 connected to only a few farms and therefore the total amount of Faroese 369 gravid lice may not be representative of the external pressure at these farms 370 (Kragesteen et al. 2018). 371

The principles of measuring salmon-louse population growth rates are rel-372 atively simple as we assume exponential growth (Eq. 15) and fit a straight 373 line with a number of consecutive log transformed lice counts. Here, we have 374 decreased the effects of treatments by excluding negative growth rates and 375 badly correlated data ($R^2 < 0.6$, Fig. A1c). This approach differs from that 376 by Patursson et al. (2017) as we here consider all lice counts after 150 days 377 into a production cycle, while Patursson et al. (2017) discarded an initial pe-378 riod until the first treatment occurred. Population growth rates include both 379 internal and external dynamics and at low lice abundances external dynamics 380 are more dominant $(\frac{L_0 e^{-\mu_1 t_1}}{\rho_m(t)})$, while at higher lice abundances the internal dynamics will dominate the growth rate both due to decreased contribution 381 382 of external dynamics and low or absent Allee effect. A problem with this 383 method is that population growth rate is very high right after a treatment 384 event and may lead to an overestimation of the growth rate. Nevertheless, 385 we estimated the growth rate for each production cycle and each active farm 386 site with a relatively high number of growth rate estimates which makes the 387 estimates altogether robust and illustrates the variability between farm sites. 388

³⁸⁹ 5. Concluding remarks

In conclusion, our results provide estimates of vital salmon-louse population dynamic parameters based on biweekly monitoring lice counts. Ex-

ternal infection was found to vary between farm sites from 0.002 to 0.1 lice 392 and farm salmon⁻¹d⁻¹ (Fig. 3). Because of the likely small ratio between wild and farm 393 salmon we believe there is a negligible contribution from wild salmonid stocks 394 on external infection pressure compared to the infection within the Faroese 395 farm network. And we show an overall significant relationship between ex-396 ternal infection pressure and total number of gravid lice, which generally 397 increases around 0.001 to 0.004 lice salmon $^{-1}d^{-1}$ for every million gravid lice 398 (Fig. 4). The salmon-louse population growth rate was found to vary be-399 tween farms ranging from 1.7 - 5.4% d⁻¹ (Fig. 6). These growth rates are 400 comparable to other estimates (Krkošek et al. 2010, Patursson et al. 2017). 401 The estimated parameters can be used to fit a salmon-louse population 402

dynamic model allowing for robust predictions of salmon-louse development on a per farm basis. Further, such estimates can be used to calibrate and validate a bio-economic lice model (Kragesteen et al. 2019) forced by connectivity between farms based on hydrodynamic modelling, which could substantially improve lice management by identifying the most cost effective approach.

408 6. Acknowledgements

This study was financially supported by Danmarks Innovationsfond (Grant No. 5189-00032A) and by Granskingarráðið - Research Council Faroe Islands (Grant No. 0450). Thanks to the companies Bakkafrost, Luna and Mowi (Marine Harvest) for supplying data. Thanks to Ása Johannesen for help with statistical analysis, Tróndur T. Johannesen for help with the mathematics and thanks Gunnvør á Norði for scientific input. Constructive comments from the three anonymous reviewers are very much appreciated.

416 References

- Adams T, Proud R, Black KD (2015) Connected networks of sea lice populations: dynamics and implications for control. Aquacult Environ Interact 6(3):273–284
- Anderson RM, May RM, et al. (1979) Population biology of infectious diseases: Part i. Nature 280(5721):361-367

Anon (2012) Forskrift om bekjempelse av lakselus i akvakulturanlegg.
https://lovdatano/dokument/SF/forskrift/2012-12-05-1140#KAPI
TTEL_1 (accessed 16 Nov 2020)

Anon (2016) Kunnger um yvirtøku of tálming av lúsum á alifiski, kg. 75,
28 nov. 2016. https://logirfo/Kunngerd/75-fra-28-06-2016-um-yvirvokuog-talming-av-lusum-a-alifiski (accessed 16 Nov 2020)

- 428 Anon (2019) Status for norske laksebestander i 2018. 429 http://hdlhandlenet/11250/26198898 (accessed 16 Nov 2020)
- Austreng E, Storebakken T, Åsgård T (1987) Growth rate estimates for cultured atlantic salmon and rainbow trout. Aquaculture 60(2):157–160
- Bjørn PA, Sivertsgård R, Finstad B, Nilsen R, Serra-Llinares RM, Kristoffersen R (2011) Area protection may reduce salmon louse infection risk to
 wild salmonids. Aquacult Environ Interact 1(3):233–244
- Boxaspen K (2006) A review of the biology and genetics of sea lice. ICES J
 Mar Sci 63(7):1304–1316
- Faroese Ministry of Foreign Affairs and Trade (2016) Kunnger um yvirtøku
 of tálming av lúsum á alifiski, kg. 75, 28 nov. 2016.
- Frazer LN, Morton A, Krkošek M (2012) Critical thresholds in sea lice epidemics: evidence, sensitivity and subcritical estimation. Proc R Soc B
 279(1735):1950–1958
- Gettinby G, Robbins C, Lees F, Heuch PA, Finstad B, Malkenes R, Revie
 CW (2011) Use of a mathematical model to describe the epidemiology
 of lepeophtheirus salmonis on farmed atlantic salmon salmo salar in the
 hardangerfjord, norway. Aquaculture 320(3):164–170

Guarracino M, Qviller L, Lillehaug A (2018) Evaluation of aquaculture management zones as a control measure for salmon lice in norway. Dis Aquat
Org 130(1):1–9

Hamre LA, Eichner C, Caipang CMA, Dalvin ST, Bron JE, Nilsen F,
Boxshall G, Skern-Mauritzen R (2013) The salmon louse lepeophtheirus
salmonis (copepoda: Caligidae) life cycle has only two chalimus stages.
PloS one 8(9):e73539

Heuch P, Nordhagen J, Schram T (2000) Egg production in the salmon louse
[lepeophtheirus salmonis (krøyer)] in relation to origin and water temperature. Aquacult Res 31(11):805–814

Jevne LS, Reitan KI (2019) How are the salmon lice (lepeophtheirus salmonis krøyer, 1837) in atlantic salmon farming affected by different control efforts: A case study of an intensive production area with coordinated production cycles and changing delousing practices in 2013–2018. J Fish Dis 42(11):1573–1586

Karbowski CM, Finstad B, Karbowski N, Hedger RD (2019) Sea lice in iceland: assessing the status and current implications for aquaculture and
wild salmonids. Aquacult Environ Interact 11:149–160

Kragesteen TJ, Simonsen K, Visser AW, Andersen KH (2018) Identifying
salmon lice transmission characteristics between faroese salmon farms.
Aquacult Environ Interact 10:49–60

Kragesteen TJ, Simonsen K, Visser AW, Andersen KH (2019) Optimal
salmon lice treatment threshold and tragedy of the commons in salmon
farm networks. Aquaculture 512:734329

Kristoffersen AB, Jimenez D, Viljugrein H, Grøntvedt R, Stien A, Jansen
PA (2014) Large scale modelling of salmon lice (lepeophtheirus salmonis)
infection pressure based on lice monitoring data from norwegian salmonid
farms. Epidemics 9:31–39

474 Kristoffersen AB, Qviller L, Helgesen KO, Vollset KW, Viljugrein H, Jansen
475 PA (2018) Quantitative risk assessment of salmon louse-induced mortality
476 of seaward-migrating post-smolt atlantic salmon. Epidemics 23:19–33

- Krkošek M, Bateman A, Proboszcz S, Orr C (2010) Dynamics of outbreak and
 control of salmon lice on two salmon farms in the broughton archipelago,
 british columbia. Aquacult Environ Interact 1(2):137–146
- Krkošek M, Connors BM, Lewis MA, Poulin R (2012) Allee effects may slow
 the spread of parasites in a coastal marine ecosystem. Am Nat 179(3):401–
 412

Krkošek M, Lewis MA, Volpe JP, Morton A (2006) Fish farms and sea lice infestations of wild juvenile salmon in the broughton archipelago—a rebuttal
to brooks (2005). Rev Fish Sci 14(1-2):1–11

- 486 MATLAB (2020) 9.8.0.1323502 (R2020a). The MathWorks Inc., Natick,
 487 Massachusetts
- O'Shea B, Mordue-Luntz A, Fryer R, Pert C, Bricknell I (2006) Determination of the surface area of a fish. J Fish Dis 29(7):437–440

Patursson EJ, Simonsen K, Visser AW, Patursson Ø (2017) Effect of exposure
on salmon lice lepeophtheirus salmonis population dynamics in faroese
salmon farms. Aquacult Environ Interact 9:33–43

Pert CC, Fryer RJ, Cook P, Kilburn R, McBeath S, McBeath A, Matejusova
I, Urquhart K, Weir SJ, McCarthy U, et al. (2014) Using sentinel cages to
estimate infestation pressure on salmonids from sea lice in loch shieldaig,
scotland. Aquacult Environ Interact 5(1):49–59

- ⁴⁹⁷ Pike A, Wadsworth S (1999) Sealice on salmonids: their biology and control.
 ⁴⁹⁸ Adv Parasitol 44:233–337
- Revie CW, Robbins C, Gettinby G, Kelly L, Treasurer J (2005) A mathematical model of the growth of sea lice, lepeophtheirus salmonis, populations
 on farmed atlantic salmon, salmo salar l., in scotland and its use in the
 assessment of treatment strategies. J Fish Dis 28(10):603-613
- Robbins C, Gettinby G, Lees F, Baillie M, Wallace C, Revie CW (2010) Assessing topical treatment interventions on scottish salmon farms using a sea
 lice (lepeophtheirus salmonis) population model. Aquaculture 306(1):191–197

Samsing F, Oppedal F, Johansson D, Bui S, Dempster T (2014) High host
densities dilute sea lice lepeophtheirus salmonis loads on individual atlantic
salmon, but do not reduce lice infection success. Aquacult Environ Interact
6(1):81–89

Sandvik AD, Bjørn PA, Ådlandsvik B, Asplin L, Skarðhamar J, Johnsen IA,
 Myksvoll M, Skogen MD (2016) Toward a model-based prediction system
 for salmon lice infestation pressure. Aquacult Environ Interact 8:527–542

Schram TA (1993) Supplementary descriptions of the developmental stages of
lepeophtheirus salmonis (krøyer, 1837)(copepoda: Caligidae). Pathogens
of wild and farmed fish: sea lice 1:30–47

Stien A, Bjørn PA, Heuch PA, Elston DA (2005) Population dynamics of
salmon lice lepeophtheirus salmonis on atlantic salmon and sea trout. Mar
Ecol Prog Ser 290:263–275

Stormoen M, Skjerve E, Aunsmo A (2013) Modelling salmon lice, lepeophtheirus salmonis, reproduction on farmed atlantic salmon, salmo salar l. J
Fish Dis 36(1):25–33

Developmen	to or stage tengen wa	is calculated using.	$(I) = [P_1/(I)]$	$10 + p_1 p_2 $
Stage	β_1	β_2	$\mu (d^{-1})$	$\tau[11^{\circ}C]$ (d)
(ρ_1)	$74.70 (\pm 33.64)$	$0.246~(\pm 0.007)$	0.002 - 0.01	14.9 (t_1)
(ρ_2)	$67.47 (\pm 20.36)$	$0.177~(\pm 0.006)$	0.025 - 0.18	$27.2 (t_2)$
$(\rho_3 \& \rho_4)$	$41.98 \ (\pm 2.85)$	$0.338 \ (\pm 0.012)$	0.025 - 0.06	$7.6 (t_3)$

Table 1: Development and mortality for attached mobile lice stages (Stien et al. 2005). Development or stage length was calculated using: $\tau(T) = [\beta_1/(T-10+\beta_1\beta_2)]^2$.

Table 2: Number of salmons and treatments per year, and average number of gravid and mobile lice per year in the Faroe Islands for years 2011-2018.

	2011	2012	2013	2014	2015	2016	2017	2018
Salmon $\cdot 10^7$	1.81	1.87	1.98	1.94	2.01	2.16	2.02	1.92
Gravid lice $\cdot 10^7$	0.97	1.23	0.83	1.15	1.78	1.54	1.26	0.68
Mobile lice $\cdot 10^7$	1.92	4.93	3.52	4.10	5.80	4.81	3.57	2.03
Treatments $(\#)$	12	26	66	71	70	94	98	93



Figure 1: Faroese farm areas (black circles) and location of temperature measurements (black cross).



Figure 2: a) Total number of gravid lice (black line), other mobile lice (gray line) and salmon (green line) in Faroe Islands. b) Number of treatments in Faroe Islands per month shown as total (black line), mechanical (blue line), bath treatments (red line) and medicinal oral (green line). c) The Faroese shelf temperature (blue). Data is shown of the period from 2011 to 2018.



Figure 3: Average external infection pressure including first 75, 100, 125 and 150 days with 95 % CI error-bars for each farm site in the Faroe Islands.



Figure 4: External infection pressure estimates as a function of total number of gravid lice. Each panel is a farm site and the blue circles are averaged external infection pressures of each production cycle. Black error bars show 95 % confidence interval. Line (y = ax + b; gray line) is linear regression model fit on all data points (gray circle) with random slope and intercept. Bold number in the legend indicates farm number and * indicates p < 0.05. Overall linear regression model fit by farm: $\mathbb{R}^2 = 0.312$ and $\mathbb{F}_{29,844} = 12.7$ with a p < 0.001.



Figure 5: Average growth rate for each farm site in Faroe Island with 95 % confidence interval.



Figure 6: Salmon-louse growth rate in day of the year of total Faroese gravid lice averaged over 14 d (black line) and 90 d (black dotted line) from 2011 to 2018 and total gravid lice $\cdot 10^7$ (red line). Light gray and dark gray color indicate 95% confidence interval for population growth rate averaged over 14 and 90 days, respectively. Average annual temperature (blue line) shown on the right y-axis in day of year from 2011 to 2018.

523 Appendix

External infection pressure and salmon-louse population growth rate was 524 estimated for each active Faroese farm site and each production cycle since 525 2009. Calculation is exemplified with Fig. A1. External infection (L_0) pres-526 sure was estimated for each lice count performed the first 150 days after 527 sea-stocking (Fig. A1b) using Equation 13. Example: The third lice count at 528 58 days after sea-stocking has 0.45 gravid lice salmon⁻¹ and 1.25 mobile lice 529 salmon⁻¹ where the mean sea temperature the previous 30 days was 10.4 °C 530 (Fig. A1a). Using equation $t_1(T) = [\beta_1/(T-10+\beta_1\beta_2)]^2$ from (Stien et al. 531 2005) and Table 1. Then we find that $t_1(10.4^{\circ}C) = 14.8d$. When calculating 532 L_0 we use equation 13: 533

$$L_0 = \frac{0.025 \mathrm{d}^{-1} (0.45 + 1.25 \text{ lice salmon}^{-1})}{e^{-0.002 \mathrm{d}^{-1} \cdot 14.8d} (1 - e^{-0.025 \mathrm{d}^{-1} (58 \mathrm{d} - 14.8\mathrm{d})})} = 0.066 \text{ lice salmon}^{-1} \mathrm{d}^{-1}$$
(16)

⁵³⁴ Doing this calculation for all lice counts the first 150 days we get a mean ⁵³⁵ $L_0 = 0.06$ lice salmon⁻¹ d⁻¹.

Salmon-louse population growth rate (Eq. 15) was estimated by fitting a 536 straight line between 5 consecutive lice counts 150 days after sea-stocking and 537 excluding all lines with a negative slope and/or R^2 less than 0.6 (Fig. A1c). 538 Using the salmon-louse population model (Eq. 2-6) to simulate population 539 growth we find that the growth rate is relatively high for a short period after 540 a treatment, because a treatment only kills attached stages and therefore all 541 larvae produced before a treatment can re-infect the farm site resulting in a 542 higher percentage growth. 543



Figure A 1: Example of calculation of external infection pressure and population growth rate. a) Gravid and mobile lice shown as solid and dashed line, respectively. Dashed dotted line is the mean gravid lice salmon⁻¹ in farm network. Vertical lines indicate treatment events and blue line is temperature (right y-axis). b) External infection pressure, L_0 , (blue square) calculated for lice counts before 150 after sea-stocking. c) natural log of ρ_m where lines are fitted with 5 consecutive lice counts after 150 days after sea stocking. Red lines indicate positive slopes and/or $\mathbb{R}^2 > 0.6$ and the legend is the average slope of all red lines.