

Reply to Letten and Yamamichi: A rescue at the cost of falsifiability

Kiørboe, Thomas; Thomas, Mridul K.

Published in: Proceedings of the National Academy of Sciences of the United States of America

Link to article, DOI: 10.1073/pnas.2025720118

Publication date: 2021

Document Version Publisher's PDF, also known as Version of record

Link back to DTU Orbit

Citation (APA): Kiørboe, T., & Thomas, M. K. (2021). Reply to Letten and Yamamichi: A rescue at the cost of falsifiability. *Proceedings of the National Academy of Sciences of the United States of America, 118*(5), Article e2025720118. https://doi.org/10.1073/pnas.2025720118

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- · You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

REPLY TO LETTEN AND YAMAMICHI: A rescue at the cost of falsifiability

Thomas Kiørboe^{a,1} and Mridul K. Thomas^{a,b,c}

Letten and Yamamichi (1) make valuable comments about our recent paper (2). They point out that the gleaner-exploiter trade-off commonly refers to population growth rates and not prey consumption (the focus of our study). We agree and should have made our reasoning clearer. We assume that food consumption and population growth rates are positively associated; therefore, a positive association between maximum ingestion rate (I_{max}) and maximum clearance rate (C_{max}) reflects a positive association in population growth rates at high and low food concentrations. Letten and Yamamichi argue that we may have overlooked a tradeoff in population growth rates through two possible mechanisms.

The first mechanism is systematic variation in conversion efficiency. We concede this point, having included the same argument in the paper as a caveat; we also included data that lends support to our assumption.

The second mechanism is a negative association between mortality rate and I_{max} and/or C_{max} . Two aspects to this are worth distinguishing between:

- 1) The authors (1) focus on predation-induced mortality, the major source of mortality in nature. They argue that the foraging-predation risk trade-off that we implicate is a gleaner-exploiter tradeoff: More foraging leads to an increase in predation-induced mortality, which increases R* (lower is better), I_{max} , and C_{max} (higher is better). We think this argument is misguided for the following reasons:
- i) R* is best understood as the minimum required resource concentration in the absence of predation. This is consistent with prior theoretical usage (3, 4) and most empirical studies characterizing R*

(5, 6). If we were to include predation losses, R^* would no longer be a property of a population but a joint function of every population and its predation environment. Consequently, evaluating the gleaner–exploiter trade-off would require arbitrary choices of predation pressure for each species. This framing therefore makes the putative trade-off highly context dependent at best and unfalsifiable at worst.

- ii) Moreover, we already have a term for a function characterizing the minimum resource requirement across a gradient of predation pressure: It is a form of zero net growth isocline, or ZNGI (3, 4). Such a ZNGI intercepts the resource axis at the R* as we define it above (3, 4).
- iii) If the trade-off between foraging and predation risk is by definition a gleaner-opportunist tradeoff, as Letten and Yamamichi (1) argue, the latter concept has lost the specificity that makes it useful. It instead becomes a stand-in for a class of different trade-off mechanisms. We agree that a gleaner-opportunist trade-off may emerge from the more fundamental foraging-risk trade-off under certain conditions but prefer to retain separate terminology for the distinct mechanisms for clarity.
- 2) Background mortality captures losses due to factors other than predation; the simplest natural analog is death due to old age. If background mortality rates are strongly positively associated with I_{max} or C_{max} , then there could indeed be a gleaner–exploiter trade-off that we overlooked. This is a useful caveat to our results, and we thank the authors (1) for pointing it out. We believe such a relationship is unlikely, but this cannot be ruled out due to the lack of data.

2 T. Kiørboe, M. K. Thomas, Heterotrophic eukaryotes show a slow-fast continuum, not a gleaner–exploiter trade-off. Proc. Natl. Acad. Sci. U.S.A. 117, 24893–24899 (2020).

¹ A. D. Letten, M. Yamamichi, Gleaning, fast and slow: In defense of a canonical ecological trade-off. Proc. Natl. Acad. Sci. U.S.A., 10.1073/pnas.2022754118 (2021).

^aCentre for Ocean Life, DTU Aqua, Technical University of Denmark, DK-2800 Kgs Lyngby, Denmark; ^bDepartment F.-A. Forel for Environmental and Aquatic Sciences, University of Geneva, CH-1211 Geneva, Switzerland; and ^cInstitute for Environmental Sciences, University of Geneva, CH-1211 Geneva, Switzerland

Author contributions: T.K. and M.K.T. wrote the paper.

The authors declare no competing interest.

Published under the PNAS license.

¹To whom correspondence may be addressed. Email: tk@aqua.dtu.dk.

Published January 25, 2021.

M. A. Leibold, A graphical model of keystone predators in food webs: Diversity patterns in communities. Am. Nat. 147, 784–812 (1996).
T. Koffel, T. Daufresne, F. Massol, C. A. Klausmeier, Plant strategies along resource gradients. Am. Nat. 192, 360–378 (2018).

5 D. Tilman, M. Mattson, S. Langer, Competition and nutrient kinetics along a temperature gradient: An experimental test of a mechanistic approach to niche theory. Limnol. Oceanogr. 26, 1020–1033 (1981).

6 J. P. Grover, Phosphorus-dependent growth kinetics of 11 species of freshwater algae. Limnol. Oceanogr. 34, 341–348 (1989).