Biofilm dispersal for spore release in *Bacillus subtilis*

Kovács, Ákos T.; Stanley-Wall, Nicola R.

Published in:
Journal of Bacteriology

Link to article, DOI:
[10.1128/JB.00192-21](https://doi.org/10.1128/JB.00192-21)

Publication date:
2021

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Kovács, Á. T., & Stanley-Wall, N. R. (2021). Biofilm dispersal for spore release in *Bacillus subtilis*. *Journal of Bacteriology*, 23(14), [e00192-21]. <https://doi.org/10.1128/JB.00192-21>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Biofilm dispersal for spore release in *Bacillus subtilis*

Ákos T. Kovács¹, Nicola R. Stanley-Wall²

¹ Bacterial Interactions and Evolution Group, DTU Bioengineering, Technical University of Denmark, Kongens Lyngby, Denmark

² Division of Molecular Microbiology, School of Life Sciences, University of Dundee, Dundee, UK

Address correspondence to Ákos T. Kovács, atkovacs@dtu.dk

ABSTRACT

The dispersal of bacterial cells from a matured biofilm can be mediated either by active or passive mechanisms. In this issue of the *Journal of Bacteriology*, Nishikawa and Kobayashi demonstrate that the presence of calcium influences dispersal of spores from the pellicle biofilm of *Bacillus subtilis*. The authors propose that temporal heterogeneity in matrix production and chelation of calcium by dipicolinic acid in spores weakens the biofilm matrix and causes passive dispersal.

KEYWORDS

Bacillus subtilis, biofilm, dispersal, spore, development

COMMENTARY

Biofilm formation is a complex developmental process undertaken by microbes that is initiated by attachment or aggregation of cells, advanced by production of an extracellular matrix, and

24 generally finalized by disassembly of the biofilm, a process called dispersal (1). The specific biofilm
25 life cycle depends on the microorganism, its ecological niche, and the encoded regulatory
26 pathways. In addition, environmental factors, including intra- and interspecies compounds may
27 influence the different steps of biofilm development. A detailed understanding of the different
28 stages of biofilm formation and disassembly could help us to prevent deleterious microbial
29 communities and promote beneficial ones.

30 *Bacillus subtilis* became a model organism to study bacterial differentiation processes due to its
31 ability to create a dormant cell structure, called a spore, that has remarkably resistance to heat,
32 pressure, and chemicals in addition to its capability to take up extracellular DNA and incorporate it
33 via recombination into its genome (2). These features, alongside other biotechnologically
34 beneficial properties stimulated robust probing of the physiology and genetics of this species in
35 the last century. The study of *B. subtilis* biofilm formation was initiated about two decades ago,
36 and has created a plethora of understanding since the first publication (3) regarding how gene
37 expression connects to biofilm initiation and matrix production and the identity and function of
38 the main biofilm matrix components (4, 5). Interest in *B. subtilis* biofilms is further stimulated by
39 the species being more than a laboratory model: biofilms are important for plant growth
40 promotion, probiotic impact, and biotechnological applications (6, 7).

41 The molecular details of *B. subtilis* biofilm development have been predominantly explored in two
42 laboratory systems, air-liquid interface floating biofilms, known as pellicles, and architecturally
43 complex colonies formed on agar surface (6). Dissection of gene expression in colonies revealed
44 that *B. subtilis* biofilm population is phenotypically heterogeneous; distinct cell types inhabit a
45 biofilm, including motile cells, matrix producers, extracellular protease producers, and in the later
46 stages of development, spores are formed on the upper layer of the colonies (8, 9). Efficient

47 initiation of pellicle development requires motility (10) and establishment of the floating biofilm at
48 the air-medium interface proceeds through distinct morphological changes (11). Importantly,
49 matrix gene expression in the nascent pellicle is temporally and spatially heterogeneous, after a
50 highly heterogeneous matrix production during initiation of the pellicle, the majority of cells
51 express the genes for matrix production in the middle of biofilm development (around 24 hours)
52 (12). In the later stages, the population becomes heterogeneous again, only a fraction of the cells
53 will produce the matrix (12), while spores also appear (13, 14). Such temporal heterogeneity is
54 mirrored by physical heterogeneity; during the initial and later stages of pellicle development,
55 next to a robust, highly matrix-expressing population, a fragile fraction is also present, within
56 which the cell-cell aggregation can be easily disrupted (12). The dynamic transcriptional landscape
57 of the developing pellicle has also been associated with variation in metabolism of the cells (15).
58 Additionally, it has been proposed that during biofilm colony maturation, the evolutionary
59 younger and more diverged genes are increasingly expressed toward later timepoints of colony
60 development (16).

61 While initiation and maturation of pellicle biofilm development is extensively investigated,
62 dispersal mechanisms are less explored in *B. subtilis* and the literature still perpetuates errors in
63 the understanding of biofilm dispersal with respect to norspermidine and D-amino acids that have
64 since been corrected (17, 18). In this issue of *Journal of Bacteriology*, the publication by Nishikawa
65 and Kobayashi (19) reveals a novel mechanism of *B. subtilis* dispersal and highlights a connection
66 between emergence of spores and biofilm disassembly (Fig. 1). Interestingly, *B. subtilis* grown in
67 variety of commonly used synthetic and complex biofilm media (e.g. MSgg and 2×SGG) does not
68 display typical dispersal. The pellicle remains robust for up to a week at 30°C, during which time
69 only a minor and very slow decay in the thickness is observed in MSgg grown pellicles. However,

70 pellicles that were cultivated at 37°C in a modified LBGM medium (lysogeny broth supplemented
71 with glycerol and manganese (20), but containing reduced amount of manganese compared to
72 previous publications) showed a rapid establishment within a day, fragmented structure on the
73 second day, and strong dispersal after 3 days. While removal of manganese prevents biofilm
74 development (20, 21), supplementation at lower concentration creates conditions that allows
75 examination of the full pellicle biofilm life cycle in *B. subtilis*, including dispersal. These
76 observations highlight that biofilm dispersal might be more prevalent in the laboratory when slight
77 starvation is encountered, a condition that likely exists for microbes in nature. Interestingly, the
78 number of viable cells remained constant throughout the 3 days (19), suggesting the lack of active
79 lysis or cell death, but rather the presence of passive dispersal in *B. subtilis* succeeding the
80 previously reported reduced matrix gene expression at later stages of the pellicle development
81 (12, 15). Synthetic induction of genes involved in synthesis of the exopolysaccharides throughout
82 the cultivation and therefore prolonged exopolysaccharide production partially prevents biofilm
83 dispersal (19), thus the reduced matrix production only partially explains pellicle biofilm dispersal.
84 What could facilitate *B. subtilis* cells' dispersal from a biofilm in addition to reduced matrix
85 expression? *B. subtilis*, when colonizing plant roots under hydroponic conditions, first produces
86 the biofilm matrix followed by robust spore formation (22). Spore are anticipated to survive the
87 harsh conditions in the soil, including predation by protozoans, nematodes, or other microbes
88 (23–25). Therefore, pellicle dispersal could understandably be mediated by release of spores.
89 Nishikawa and Kobayashi (19) demonstrate that the induction of sporulation pathway, which
90 depends on a cascade of sigma factors activating specific gene expression either in the mother cell
91 (σ^E , and σ^K) or in the pre-spore (σ^F , σ^G), contributes to dispersal. Indeed, circumventing spore
92 formation by disrupting these sporulation-specific sigma factors, in addition to concomitantly

93 synthetically prolonging exopolysaccharide production, prevents pellicle dispersion. Systematic
94 disruption of σ^K -dependent genes, the last downstream sigma factor within the activation
95 cascade, revealed that *spoVFA*–*spoVFB* operon is sufficient to explain biofilm dispersal in *B.*
96 *subtilis*. The *spoVFA*–*spoVFB* operon encodes a dipicolinic acid synthase that creates dipicolinic
97 acid (DPA). DPA after being produced in the mother cells is transported to the forespore
98 compartment where it starts chelating calcium ions contributing to dehydration and
99 mineralization of the spore (26). The direct connection between DPA mediated chelation of
100 calcium, and pellicle dispersal could be verified by addition of calcium to the biofilm medium,
101 which prevented dispersal (19). The impact of calcium on biofilm colony development has been
102 previously observed (27). Consistently, both Nishikawa and Kobayashi (19) and Mhatre *et al.* (27)
103 could demonstrate that calcium does not impact the expression of matrix genes in established
104 pellicle biofilms and under biofilm inducing conditions, respectively, besides both studies reported
105 larger biofilm colony size in the absence of calcium. The larger colony size observed in the absence
106 of calcium is likely connected to passive surface spreading, termed sliding, as the influence of
107 calcium on colony size was only apparent in the presence of matrix components and surfactin that
108 are all necessary for sliding (27–29). Production of the secondary metabolite surfactin, while not
109 being essential for biofilm development, alters the architecture of biofilm colonies (30). Calcium
110 influences self-assembly of surfactin (31), which might explain a possible impact of calcium on
111 surface tension and therefore biofilm colony size (27). Nevertheless, it remains to resolve the
112 direct connection between calcium level and surfactin functioning in colonies.

113 Thus, calcium has an important role both in spore maturation and also influences dispersal.
114 Nishikawa and Kobayashi (19) offer a plausible explanation how spore formation indirectly
115 influences biofilm weakening. Besides lower production of the matrix, the onset of spore

116 maturation and accompanying DPA production depletes the calcium in the extracellular matrix,
117 resulting in biofilm dispersal (Fig. 1). Calcium seems to impact the biofilm matrix and/or influence
118 regulation of biofilm in numerous bacteria (32–34). It remains to demonstrate whether and how
119 calcium directly interacts with the biofilm matrix in *B. subtilis*. Nevertheless, the elegant work by
120 Nishikawa and Kobayashi describes an intriguing example of passive dispersal and connects spore
121 formation with its release from the biofilms.

122

123 **ACKNOWLEDGEMENT**

124 Biofilm related work in the group of Á.T.K. is supported by a DTU Alliance Strategic Partnership
125 PhD fellowship. Work in the laboratory of N.R.S.-W. is funded by the Biotechnology and Biological
126 Science Research Council (BBSRC) (BB/P001335/1 and BB/R012415/1).

127

128 **REFERENCES**

- 129 1. Watnick P, Kolter R. 2000. Biofilm, city of microbes. *J Bacteriol* 182:2675–2679.
- 130 2. Kovács ÁT. 2019. *Bacillus subtilis*. *Trends Microbiol* 27:724–725.
- 131 3. Branda SS, González-Pastor JE, Ben-Yehuda S, Losick R, Kolter R. 2001. Fruiting body
132 formation by *Bacillus subtilis*. *Proc Natl Acad Sci U S A* 98:11621–11626.
- 133 4. Mhatre E, Monterrosa RG, Kovács ÁT. 2014. From environmental signals to regulators:
134 Modulation of biofilm development in Gram-positive bacteria. *J Basic Microbiol* 54:616–
135 632.
- 136 5. Cairns LS, Hobley L, Stanley-Wall NR. 2014. Biofilm formation by *Bacillus subtilis*: New
137 insights into regulatory strategies and assembly mechanisms. *Mol Microbiol* 93:587–598.
- 138 6. Arnaouteli S, Bamford N, Stanley-Wall NR, Kovács ÁT. 2021. *Bacillus subtilis* biofilm

- 139 formation and social interactions. Nat Rev Microbiol [https://doi.org/10.1038/s41579-021-](https://doi.org/10.1038/s41579-021-00540-9)
140 00540-9.
- 141 7. Blake C, Christensen MN, Kovacs AT. 2021. Molecular aspects of plant growth promotion
142 and protection by *Bacillus subtilis*. Mol Plant-Microbe Interact 34:15–25.
- 143 8. Vlamakis H, Aguilar C, Losick R, Kolter R. 2008. Control of cell fate by the formation of an
144 architecturally complex bacterial community. Genes Dev 22:945–953.
- 145 9. Marlow VL, Cianfanelli FR, Porter M, Cairns LS, Kim Dale J, Stanley-Wall NR. 2014. The
146 prevalence and origin of exoprotease-producing cells in the *Bacillus subtilis* biofilm.
147 Microbiol (United Kingdom) 160:56–66.
- 148 10. Hölscher T, Bartels B, Lin Y-C, Gallegos-Monterrosa R, Price-Whelan A, Kolter R, Dietrich LEP,
149 Kovács ÁT. 2015. Motility, chemotaxis and aerotaxis contribute to competitiveness during
150 bacterial pellicle biofilm development. J Mol Biol 427:3695–3708.
- 151 11. Kobayashi K. 2007. *Bacillus subtilis* pellicle formation proceeds through genetically defined
152 morphological changes. J Bacteriol 189:4920–4931.
- 153 12. Otto SB, Martin M, Schäfer D, Hartmann R, Drescher K, Brix S, Dragoš A, Kovács ÁT. 2020.
154 Privatization of biofilm matrix in structurally heterogeneous biofilms. mSystems 5:e00425-
155 20.
- 156 13. Martin M, Dragoš A, Hölscher T, Maróti G, Bálint B, Westermann M, Kovács ÁT. 2017. *De*
157 *novo* evolved interference competition promotes the spread of biofilm defectors. Nat
158 Commun 8:15127.
- 159 14. Špacapan M, Danevčič T, Štefanič P, Porter M, Stanley-Wall NR, Mandić-Mulec I. 2020. The
160 ComX quorum sensing peptide of *Bacillus subtilis* affects biofilm formation negatively and
161 sporulation positively. Microorganisms 8:1131.

- 162 15. Pisithkul T, Schroeder JW, Trujillo EA, Yeesin P, Stevenson DM, Chaiamarit T, Coon JJ, Wang
163 JD, Amador-Noguez D. 2019. Metabolic remodeling during biofilm development of *Bacillus*
164 *subtilis*. MBio 10:e00623-19.
- 165 16. Futo M, Opašić L, Koska S, Corak N, Široki T, Ravikumar V, Thorsell A, Lenuzzi M, Kifer D,
166 Domazet-Lošo M, Vlahoviček K, Mijakovic I, Domazet-Lošo T. 2021. Embryo-like features in
167 developing *Bacillus subtilis* biofilms. Mol Biol Evol 38:31–47.
- 168 17. Hobley L, Kim SH, Maezato Y, Wyllie S, Fairlamb AH, Stanley-Wall NR, Michael AJ. 2014.
169 Norspermidine is not a self-produced trigger for biofilm disassembly. Cell 156:844–854.
- 170 18. Leiman SA, May JM, Lebar MD, Kahne D, Kolter R, Losick R. 2013. D-Amino acids indirectly
171 inhibit biofilm formation in *Bacillus subtilis* by interfering with protein synthesis. J Bacteriol
172 195:5391–5395.
- 173 19. Nishikawa M, Kobayashi K. 2021. Calcium prevents biofilm dispersion in *Bacillus subtilis*. J
174 Bacteriol.
- 175 20. Shemesh M, Chaia Y. 2013. A combination of glycerol and manganese promotes biofilm
176 formation in *Bacillus subtilis* via histidine kinase KinD signaling. J Bacteriol 195:2747–2754.
- 177 21. Mhatre E, Troszok A, Gallegos-Monterrosa R, Lindstädt S, Hölscher T, Kuipers OP, Kovács ÁT.
178 2016. The impact of manganese on biofilm development of *Bacillus subtilis*. Microbiology
179 162:1468–1478.
- 180 22. Charron-Lamoureux V, Beauregard PB. 2019. Arabidopsis thaliana seedlings influence
181 *Bacillus subtilis* spore formation. Mol Plant-Microbe Interact 32:1188–1195.
- 182 23. Laaberki MH, Dworkin J. 2008. Role of spore coat proteins in the resistance of *Bacillus*
183 *subtilis* spores to *Caenorhabditis elegans* predation. J Bacteriol 190:6197–6203.
- 184 24. Klobutcher LA, Ragkousi K, Setlow P. 2006. The *Bacillus subtilis* spore coat provides “eat

- 185 resistance" during phagocytic predation by the protozoan *Tetrahymena thermophila*. Proc
186 Natl Acad Sci U S A 103:165–170.
- 187 25. Seccareccia I, Kovács ÁT, Gallegos-Monterrosa R, Nett M. 2016. Unraveling the predator-
188 prey relationship of *Cupriavidus necator* and *Bacillus subtilis*. Microbiol Res 192:231–238.
- 189 26. Setlow P. 2006. Spores of *Bacillus subtilis*: Their resistance to and killing by radiation, heat
190 and chemicals, p. 514–525. In Journal of Applied Microbiology.
- 191 27. Mhatre E, Sundaram A, Hölscher T, Mühlstädt M, Bossert J, Kovács ÁT. 2017. Presence of
192 calcium lowers the expansion of *Bacillus subtilis* colony biofilms. Microorganisms 5:7.
- 193 28. Hölscher T, Kovács ÁT. 2017. Sliding on the surface: bacterial spreading without an active
194 motor. Environ Microbiol 19:2537–2545.
- 195 29. Grau RR, De Oña P, Kunert M, Leñini C, Gallegos-Monterrosa R, Mhatre E, Vileta D, Donato
196 V, Hölscher T, Boland W, Kuipers OP, Kovács ÁT. 2015. A duo of potassium-responsive
197 histidine kinases govern the multicellular destiny of *Bacillus subtilis*. MBio 6:e00581-15.
- 198 30. Thérien M, Kieseewalter HT, Auria E, Charron-Lamoureux V, Wibowo M, Maróti G, Kovács ÁT,
199 Beauregard PB. 2020. Surfactin production is not essential for pellicle and root-associated
200 biofilm development of *Bacillus subtilis*. Biofilm 2:100021.
- 201 31. Arutchelvi J, Sangeetha J, Philip J, Doble M. 2014. Self-assembly of surfactin in aqueous
202 solution: Role of divalent counterions. Colloids Surfaces B Biointerfaces 116:396–402.
- 203 32. Das T, Sehar S, Koop L, Wong YK, Ahmed S, Siddiqui KS, Manefield M. 2014. Influence of
204 calcium in extracellular DNA mediated bacterial aggregation and biofilm formation. PLoS
205 One 9:e91935.
- 206 33. Liu X, Zhang K, Liu Y, Zou D, Wang D, Xie Z. 2020. Effects of calcium and signal sensing
207 systems on *Azorhizobium caulinodans* biofilm formation and host colonization. Front

- 208 Microbiol 11:563367.
- 209 34. Tischler AH, Lie L, Thompson CM, Visick KL. 2018. Discovery of calcium as a biofilm-
- 210 promoting signal for *Vibrio fischeri* reveals new phenotypes and underlying regulatory
- 211 complexity. J Bacteriol 200:e00016-18.
- 212
- 213

214 **Figure 1**

215 **Schematic representation of the *B. subtilis* pellicle biofilm life-cycle.** After inoculation of
216 planktonic cells, oxygen depletion drives the motile cells to the air-medium interface, where
217 biofilm formation is initiated. At the start, part of the population produces the biofilm matrix.
218 During biofilm maturation, most cells expend energy making the biofilm matrix and calcium is
219 distributed across the biofilm, possibly stabilizing the matrix structure. Before dispersal, matrix
220 production diminishes, and spores are formed that chelate available calcium. Calcium depletion
221 and reduced matrix production allow passive dispersal of *B. subtilis* pellicle biofilms. Figure
222 created with BioRender.com