



Weierstrass semigroups on the Skabelund maximal curve

Beelen, Peter ; Landi, Leonardo; Montanucci, Maria

Published in:
Finite Fields and Their Applications

Link to article, DOI:
[10.1016/j.ffa.2021.101811](https://doi.org/10.1016/j.ffa.2021.101811)

Publication date:
2021

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Beelen, P., Landi, L., & Montanucci, M. (2021). Weierstrass semigroups on the Skabelund maximal curve. *Finite Fields and Their Applications*, 72, Article 101811. <https://doi.org/10.1016/j.ffa.2021.101811>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Weierstrass semigroups on the Skabelund maximal curve

Peter Beelen, Leonardo Landi, and Maria Montanucci

Abstract

In [14], D. Skabelund constructed a maximal curve over \mathbb{F}_{q^4} as a cyclic cover of the Suzuki curve. In this paper we explicitly determine the structure of the Weierstrass semigroup at any point P of the Skabelund curve. We show that its Weierstrass points are precisely the \mathbb{F}_{q^4} -rational points. Also we show that among the Weierstrass points, two types of Weierstrass semigroup occur: one for the \mathbb{F}_q -rational points, one for the remaining \mathbb{F}_{q^4} -rational points. For each of these two types its Apéry set is computed as well as a set of generators.

AMS: 11G20, 14H05, 14H55

Keywords: Finite field, maximal curve, Suzuki curve, Weierstrass semigroup, Weierstrass points.

1 Introduction

Let \mathcal{X} be a nonsingular, projective algebraic curve of genus g defined over a field \mathbb{K} . Let P be a rational point of \mathcal{X} . The Weierstrass semigroup $H(P)$ is defined as the set of natural numbers n for which there exists a function f on \mathcal{X} having pole divisor $(f)_\infty = nP$. In general, the semigroup $H(P)$ can be defined for any point $P \in \mathcal{X}$ by seeing \mathcal{X} as an algebraic curve over the algebraic closure of \mathbb{K} .

According to the Weierstrass gap Theorem, see [15, Theorem 1.6.8], the set $G(P) := \mathbb{N} \setminus H(P)$ contains exactly g elements called gaps. The structure of $H(P)$ in general varies as $P \in \mathcal{X}$ varies. However, it is known that generically the semigroup $H(P)$ is the same, but that there can exist finitely many points of \mathcal{X} , called Weierstrass points, with a different set of gaps. These points are of intrinsic theoretical interest, for example in Stöhr-Voloch theory [16] to obtain characterizing properties of the curve, but when \mathbb{K} is a finite field, they also occur in the study of algebraic-geometry (AG) codes [17].

In this direction, an intensively studied class of curves is the class of maximal curves, that is, algebraic curves defined over a finite field \mathbb{F}_q having as many rational points as possible according to the Hasse–Weil bound. More precisely, an algebraic curve \mathcal{X} of genus $g(\mathcal{X})$ defined over \mathbb{F}_q is \mathbb{F}_q -maximal if it has exactly $q + 1 + 2g(\mathcal{X})\sqrt{q}$ \mathbb{F}_q -rational points. Clearly, this can only be the case if the cardinality q of the finite field is a square or $g(\mathcal{X}) = 0$.

Important examples of maximal curves over suitable finite fields are the so-called Deligne-Lusztig curves; namely the \mathbb{F}_{q^2} -maximal Hermitian curve

$$\mathcal{H} : y^{q+1} = x^q + x,$$

the \mathbb{F}_{q^4} -maximal Suzuki curve

$$\mathcal{S} : y^q + y = x^{q_0}(x^q + x), \tag{1}$$

where $q_0 = 2^s$, $s \geq 1$ and $q = 2q_0^2$; and the \mathbb{F}_{q^6} -maximal Ree curve

$$\mathcal{R} : \begin{cases} z^q - z = x^{2q_0}(x^q - x), \\ y^q - y = x^{q_0}(x^q - x), \end{cases}$$

where $q_0 = 3^s$, $s \geq 1$ and $q = 3q_0^2$.

For a fixed g , the curve \mathcal{H} has the largest possible genus $g(\mathcal{H}) = q(q-1)/2$ that an \mathbb{F}_{q^2} -maximal curve can have. Also \mathcal{H} and \mathcal{S} are two of the four only curves of genus $g \geq 2$ having at least $8g^3$ automorphisms, see [11, Theorem 11.127] and [10]. The Weierstrass points of \mathcal{H} and \mathcal{S} as well as the precise structure of the Weierstrass semigroups at every point of these curves are known; see [6] and [1]. On the other hand nothing is known on the curve \mathcal{R} and the computation of Weierstrass semigroups seems to be a challenging task, see [4].

By a result commonly attributed to Serre, see [13, Proposition 6], any \mathbb{F}_{q^2} -rational curve which is covered by an \mathbb{F}_{q^2} -maximal curve is also \mathbb{F}_{q^2} -maximal. Apart from the Deligne-Lusztig curves, most of the known maximal curves are subcovers of the Hermitian curve.

Since 2009, examples of maximal curves that are not subcovers of the Hermitian curve have been constructed as Kummer extensions of the Deligne-Lusztig curves. The first known example $\tilde{\mathcal{H}}$ of a maximal curve which is not a subcover of the Hermitian curve was constructed by Giulietti and Korchmáros as

$$\tilde{\mathcal{H}} : \begin{cases} y^{q+1} = x^q + x, \\ z^{\frac{q^3+1}{q+1}} = y^{q^2} - y; \end{cases}$$

see [7]. This curve is \mathbb{F}_{q^6} -maximal and commonly called the Giulietti-Korchmáros (GK) curve. Other two families of maximal curves as generalizations of the GK curve and Kummer extensions of the Hermitian curve were constructed in [2] and [5].

Analogously, Skabelund [14] constructed Kummer extensions of the Suzuki and Ree curves as follows. Let $q_0 = 2^s$ with $s \geq 1$ and $q = 2q_0^2$. The curve

$$\tilde{\mathcal{S}} : \begin{cases} y^q + y = x^{q_0}(x^q + x), \\ t^{\frac{q^2+1}{q+2q_0+1}} = x^q + x, \end{cases} \quad (2)$$

is \mathbb{F}_{q^4} -maximal. Now let $q_0 = 3^s$ with $s \geq 1$ and $q = 3q_0^2$. The curve

$$\tilde{\mathcal{R}} : \begin{cases} y^q - y = x^{q_0}(x^q - x), \\ z^q - z = x^{2q_0}(x^q - x), \\ t^{\frac{q^3+1}{q+3q_0+1}} = x^q - x, \end{cases}$$

is \mathbb{F}_{q^6} -maximal. The Weierstrass points as well as the precise structure of the Weierstrass semigroups at every point of $\tilde{\mathcal{H}}$ was determined in [3]. Hence it is natural to ask whether the same can be done for the curve $\tilde{\mathcal{S}}$. In this paper, a complete answer to the aforementioned question is given. More precisely, we show the following theorem.

Theorem 1.1. *Let $q_0 = 2^s$ with $s \geq 1$ and $q = 2q_0^2$. Let $\tilde{\mathcal{S}}$ be the curve defined in Equation (2) and $P \in \tilde{\mathcal{S}}(\overline{\mathbb{F}}_q)$. As usual denote by $H(P)$ the Weierstrass semigroup of P . Then the following hold:*

If $P \in \tilde{\mathcal{S}}(\mathbb{F}_q)$, then $H(P) = \langle q^2 - 2q_0q + q, q^2 - q_0q + q_0, q^2 - q + 2q_0, q^2, q^2 + 1 \rangle$.

If $P \in \tilde{\mathcal{S}}(\mathbb{F}_{q^4}) \setminus \tilde{\mathcal{S}}(\mathbb{F}_q)$, then

$$H(P) = \langle q^2 - q + 1, q^2 - 2q_0 + 1, q^2 - q_0 + 1, q^2, q^2 + 1, f_i, h_j \mid i = 0, \dots, 2q_0 - 2, j = 0, \dots, q_0 - 2 \rangle,$$

where $f_i := (i + 1)q_0(q^2 - q + 1) - i(q^2 + 1) - 1$ and $h_j := (2j + 1)q_0(q^2 - q + 1) - j(q^2 + 1) - q_0$.

For integers a_1, a_2, a_3, a_4, f , we write $\sigma := a_1 + a_2 + a_3 + a_4 + f$ and $\nu := a_1 + a_2q_0 + a_32q_0 + a_4q + fq^2$. If $P \notin \tilde{\mathcal{S}}(\mathbb{F}_{q^4})$, then $H(P) = \mathbb{N} \setminus (F_1 \cup F_2 \cup F_3 \cup F_4 \cup F_5 \cup F_6)$, where

$$\begin{aligned} F_1 &= \{\nu + 1 \mid 0 \leq a_1 \leq q_0 - 1; 0 \leq a_2 \leq 1; 0 \leq a_3 \leq q_0 - 1; a_4 \geq 0; f \geq 0; \sigma \leq q - 2\}, \\ F_2 &= \{\nu + (n + 1)q_0q + 1 \mid 1 \leq n \leq 2q_0 - 2; 0 \leq a_1 \leq q_0 - 1; 0 \leq a_2 \leq 1; 0 \leq a_3 \leq q_0 - 1; \\ &\quad 0 \leq a_4 \leq q - q_0 - 1 - nq_0; f \geq 0; \sigma = q - q_0 - 2 - nq_0 + n\}, \\ F_3 &= \{\nu + (2n + 1)q_0q + n + 2 \mid 0 \leq n \leq q_0 - 2; 0 \leq a_1 \leq q_0 - 2 - n; 0 \leq a_2 \leq 1; 0 \leq a_3 \leq q_0 - 1; \\ &\quad 0 \leq a_4 \leq q_0 - 1; f \geq 0; \sigma = q - q_0 - 2 - 2nq_0 + n\}, \\ F_4 &= \{\nu + (2n + 2)q_0q + n + 3 \mid 0 \leq n \leq q_0 - 3; 0 \leq a_1 \leq q_0 - 3 - n; 0 \leq a_2 \leq 1; 0 \leq a_3 \leq q_0 - 1; \\ &\quad 0 \leq a_4 \leq q_0 - 1; f \geq 0; \sigma = q - 2q_0 - 2 - 2nq_0 + n\}, \\ F_5 &= \{\nu + cq_0(q + 1) + d(2qq_0 + 2q_0 + 1) + 1 \mid a_1 = 0; 0 \leq c \leq 1; 0 \leq a_2 \leq 1 - c; 1 - c \leq d \leq q_0 - 1; \\ &\quad 0 \leq a_3 \leq q_0 - 1 - d; 0 \leq a_4 \leq q_0 - 1; f \geq 0; \sigma = q - 2 - 2dq_0 - cq_0\}, \\ F_6 &= \{\nu + q_0 + (2n + 2)q_0q + n + 2 \mid a_1 = 0; a_2 = 0; 0 \leq n \leq q_0 - 2; 0 \leq a_3 \leq n; 0 \leq a_4 \leq q_0 - 1; \\ &\quad f \geq 0; \sigma = q - 2q_0 - 2 - 2nq_0 + n\}. \end{aligned}$$

As a result, we will also obtain the set of Weierstrass points of $\tilde{\mathcal{S}}$.

Corollary 1.2. *The set of Weierstrass points of $\tilde{\mathcal{S}}$ is equal to $\tilde{\mathcal{S}}(\mathbb{F}_{q^4})$.*

The paper is organized as follows: In the next section we give the necessary background on the curve $\tilde{\mathcal{S}}$ as well as some results on Weierstrass semigroups and their gaps that we will need later. In Section 3, we compute $H(P)$ for $P \in \tilde{\mathcal{S}}(\mathbb{F}_q)$. Here it should be mentioned that to a large extent this case was already treated in [14]. Our results complement those in [14] by proving a claim on the generators of the semigroup that was stated in [14] without proof. For $P \notin \tilde{\mathcal{S}}(\mathbb{F}_q)$, the corresponding Weierstrass semigroups are currently unknown. Our main results are the determination of these semigroups. More precisely, in Section 4 we compute the Weierstrass semigroup for $P \in \tilde{\mathcal{S}}(\mathbb{F}_{q^4}) \setminus \tilde{\mathcal{S}}(\mathbb{F}_q)$, while in Section 5 we deal with the generic case $P \notin \tilde{\mathcal{S}}(\mathbb{F}_{q^4})$.

2 The curve $\tilde{\mathcal{S}}$

Let $q_0 = 2^s$ with $s \in \mathbb{N}$ and let $q = 2q_0^2$. The curve $\tilde{\mathcal{S}}$ defined in Equation (2) was constructed by Skabelund in [14]. This curve has genus $g(\tilde{\mathcal{S}}) = \frac{q^3 - 2q^2 + q}{2}$, $q^5 - q^4 + q^3 + 1$ \mathbb{F}_{q^4} -rational points, and a unique point at infinity P_∞ , which is singular and \mathbb{F}_q -rational. The curve $\tilde{\mathcal{S}}$ has been introduced in [14], where it was proved that $\tilde{\mathcal{S}}$ is maximal over \mathbb{F}_{q^4} . As is clear from Equations (1) and (2), the curve $\tilde{\mathcal{S}}$ is a cyclic Galois cover of the Suzuki curve of degree $q - 2q_0 + 1$ by projecting (x, y, t) on (x, y) . In the remainder of this paper, we will denote this cover by $\text{pr} : \tilde{\mathcal{S}} \rightarrow \mathcal{S}$. The cover pr gives in the usual way rise to a map $\text{pr}^* : \text{Div}(\mathcal{S}) \rightarrow \text{Div}(\tilde{\mathcal{S}})$. If $Q \in \mathcal{S}$, then for any Q of \mathcal{S} , the divisor $\text{pr}^*(Q)$ is equal to the orbit of any point $P \in \text{pr}^{-1}(Q)$ under the cyclic group C_{q-2q_0+1} multiplied with the order of the stabilizer of P in C_{q-2q_0+1} . For an algebraic function

f on \mathcal{S} (resp. $\tilde{\mathcal{S}}$), we will write $(f)_{\mathcal{S}}$ (resp. $(f)_{\tilde{\mathcal{S}}}$) for its divisor. It is well known that for any function $f \in \mathbb{F}_q(\mathcal{S})$, it holds that $(f)_{\tilde{\mathcal{S}}} = \text{pr}^*((f)_{\mathcal{S}})$.

The automorphism group $\text{Aut}(\tilde{\mathcal{S}})$ of $\tilde{\mathcal{S}}$ is defined over \mathbb{F}_{q^4} and has size $q^2(q-1)(q^2+1)(q-2q_0+1)$. It has a normal subgroup H isomorphic to the Suzuki group $\text{Sz}(q)$ and $\text{Aut}(\tilde{\mathcal{S}}) = H \times C_{q-2q_0+1}$, where C_{q-2q_0+1} is the Galois group of the cyclic Galois-covering $\text{pr} : \tilde{\mathcal{S}} \rightarrow \mathcal{S}$. The set $\tilde{\mathcal{S}}(\mathbb{F}_{q^4})$ of the \mathbb{F}_{q^4} -rational points of $\tilde{\mathcal{S}}$ splits into two orbits under the action of $\text{Aut}(\tilde{\mathcal{S}})$: one orbit has size q^2+1 and equals $\mathcal{O}_1 = \tilde{\mathcal{S}}(\mathbb{F}_q)$. The other orbit is $\mathcal{O}_2 = \tilde{\mathcal{S}}(\mathbb{F}_{q^4}) \setminus \tilde{\mathcal{S}}(\mathbb{F}_q)$ and has size $q^5 - q^4 + q^3 - q^2$; see [8] for these and other details on $\tilde{\mathcal{S}}$. In particular, we can conclude that any point $P \in \mathcal{O}_1 = \tilde{\mathcal{S}}(\mathbb{F}_q)$ has the same Weierstrass semigroup and similarly for $P \in \mathcal{O}_2 = \tilde{\mathcal{S}}(\mathbb{F}_{q^4}) \setminus \tilde{\mathcal{S}}(\mathbb{F}_q)$.

Let $x, y, z, t \in \mathbb{F}_{q^4}(\tilde{\mathcal{S}})$ be the coordinate functions of the function field of $\tilde{\mathcal{S}}$, which satisfy $y^q + y = x^{q_0}(x^q + x)$ and $t^{q-2q_0+1} = x^q + x$. These functions have a pole at P_{∞} only. The same is true for the functions $z := y^{2q_0} + x^{2q_0+1}$ and $w := xy^{2q_0} + z^{2q_0}$. Now let $P \neq P_{\infty}$ be the point of $\tilde{\mathcal{S}}$ with (x, y, t) -coordinates (a, b, c) . Further let $Q := \text{pr}(P)$. Then Q has (x, y) -coordinates (a, b) . With $\Phi(P)$ (resp. $\Phi(Q)$) we denote the point $\Phi(P) = (a^q, b^q, c^q)$ (resp. $\Phi(Q) = (a^q, b^q)$). In the sequel we will frequently use the following functions and information on their divisors:

$$\tilde{x}_Q := x + a; \quad (\tilde{x}_Q)_{\mathcal{S}} = Q + E_x - qQ_{\infty}, \quad (3)$$

where $E_x \geq 0$ and $\text{Supp}(E_x) \cap \{Q, Q_{\infty}\} = \emptyset$,

$$\tilde{y}_Q := y + b + a^{q_0}(x + a); \quad (\tilde{y}_Q)_{\mathcal{S}} = q_0Q + \Phi(Q) + E_y - (q + q_0)Q_{\infty}, \quad (4)$$

where $E_y \geq 0$ and $\text{Supp}(E_y) \cap \{Q, \Phi(Q), Q_{\infty}\} = \emptyset$,

$$\tilde{z}_Q := a^{2q_0}x + z + b^{2q_0}; \quad (\tilde{z}_Q)_{\mathcal{S}} = 2q_0Q + \Phi(Q) + E_z - (q + 2q_0)Q_{\infty}, \quad (5)$$

where $E_z \geq 0$ and $\text{Supp}(E_z) \cap \{Q, \Phi(Q), Q_{\infty}\} = \emptyset$, and

$$\tilde{w}_Q := a^q\tilde{z}_Q + b^{2q_0}x + w + b^2 + a^{2q_0+2}; \quad (\tilde{w}_Q)_{\mathcal{S}} = qQ + 2q_0\Phi(Q) + \Phi^2(Q) - (q + 2q_0 + 1)Q_{\infty}. \quad (6)$$

Over \mathbb{F}_q , the Suzuki curve has L -polynomial $(1 + 2q_0T + qT^2)^{g(\mathcal{S})}$. By the Fundamental Equation [11, Proposition 10.6 (I)], the operator $q \text{id} + 2q_0\Phi + \Phi^2$ therefore acts trivially on linear equivalence classes of divisors of degree zero, for example on $Q - Q_{\infty}$. This means that the existence of a function \tilde{w}_Q with divisor $(q \text{id} + 2q_0\Phi + \Phi^2)(Q - Q_{\infty})$ is guaranteed. Similarly applying the Fundamental Equation to the maximal curve $\tilde{\mathcal{S}}$ (over \mathbb{F}_{q^4}), we see that there exists a function π_P on $\tilde{\mathcal{S}}$ with divisor:

$$(\pi_P)_{\tilde{\mathcal{S}}} = q^2P + \Phi^4(P) - (q^2 + 1)P_{\infty}. \quad (7)$$

In particular $(\pi_P)_{\tilde{\mathcal{S}}} = (q^2 + 1)(P - P_{\infty})$ if P is \mathbb{F}_{q^4} -rational. It is not hard to find the function π_P explicitly:

$$\pi_P = \tilde{w}_Q + A^q\tilde{z}_Q + A^{q+2q_0}\tilde{x}_Q + c^{q^2}(t + c), \text{ where } A := a^q + a.$$

Note that if P is \mathbb{F}_q -rational, then $A = 0$ and $\pi_P = \tilde{w}_Q$. This observation is consistent with the divisors given above. Indeed, if $\Phi(P) = P$, then also $\Phi(Q) = Q$ and since Q and Q_{∞} are totally ramified in the cover $\text{pr} : \tilde{\mathcal{S}} \rightarrow \mathcal{S}$:

$$(\pi_P)_{\tilde{\mathcal{S}}} = \text{pr}^*((\tilde{w}_Q)_{\mathcal{S}}) = (q + 2q_0 + 1)(q - 2q_0 + 1)(P - P_{\infty}) = (q^2 + 1)(P - P_{\infty}).$$

As we already announced, our aim is to determine the structure of the Weierstrass semigroup at every point P of the curve $\tilde{\mathcal{S}}$. First of all we observe that the orbit structure of $\text{Aut}(\tilde{\mathcal{S}})$ tells us already that the Weierstrass semigroup at $P \in \tilde{\mathcal{S}}(\mathbb{F}_q)$ will be the same as $H(P_\infty)$, and $H(R) = H(Q)$ for all $R, Q \in \tilde{\mathcal{S}}(\mathbb{F}_{q^4}) \setminus \tilde{\mathcal{S}}(\mathbb{F}_q)$.

Clearly, as we will do for $P \notin \tilde{\mathcal{S}}(\mathbb{F}_{q^4})$, computing $H(P)$ is equivalent to completely determine the gap structure at P , that is, the complement $G(P) = \mathbb{N} \setminus H(P)$. Hence to finish this section we state some facts that we will use to achieve this. We start with the following well-known proposition connecting regular differentials (i.e., differential forms having no poles anywhere on $\tilde{\mathcal{S}}$) and gaps of $H(P)$.

Proposition 2.1. *[18, Corollary 14.2.5] Let \mathcal{X} be an algebraic curve of genus g defined over a field \mathbb{K} . Let P be a point of \mathcal{X} and ω be a regular differential on \mathcal{X} . Then $v_P(\omega) + 1$ is a gap at P .*

This proposition has the following, for us very useful, consequence.

Corollary 2.2. *For any point P on the curve $\tilde{\mathcal{S}}$ distinct from P_∞ , and for any $f \in L((2g(\tilde{\mathcal{S}}) - 2)P_\infty)$, we have $v_P(f) + 1 \in G(P)$.*

Proof. First note that $(dx)_{\mathcal{S}} = (2g(\mathcal{S}) - 2)Q_\infty = (2q_0(q - 1) - 2)Q_\infty$. The set of points that branch in the cover $\text{pr} : \tilde{\mathcal{S}} \rightarrow \mathcal{S}$ is exactly $\mathcal{S}(\mathbb{F}_q)$, the set of \mathbb{F}_q -rational points of the Suzuki curve, all with ramification index $q - 2q_0 + 1$. Moreover, the points of $\tilde{\mathcal{S}}$ above $\mathcal{S}(\mathbb{F}_q)$ are precisely the \mathbb{F}_q -rational points of $\tilde{\mathcal{S}}$. Therefore, we immediately obtain that

$$\begin{aligned} (dx)_{\tilde{\mathcal{S}}} &= (q - 2q_0 + 1)(2q_0(q - 1) - 2)P_\infty + (q - 2q_0) \sum_{P \in \tilde{\mathcal{S}}(\mathbb{F}_q)} P \\ &= (2q_0q^2 - 2q^2 + q - 2)P_\infty + (q - 2q_0) \sum_{P \in \tilde{\mathcal{S}}(\mathbb{F}_q), P \neq P_\infty} P. \end{aligned}$$

Thus, from $t^{q-2q_0+1} = x^q + x$ and $(t)_{\tilde{\mathcal{S}}} = \sum_{P \in \tilde{\mathcal{S}}(\mathbb{F}_q), P \neq P_\infty} P - q^2P_\infty$ we get,

$$(dt)_{\tilde{\mathcal{S}}} = \left(\frac{dx}{t^{q-2q_0}} \right)_{\tilde{\mathcal{S}}} = (q^3 - 2q^2 + q - 2)P_\infty.$$

In particular a differential $f dt$ is regular if and only if $f \in L((q^3 - 2q^2 + q - 2)P_\infty) = L((2g(\tilde{\mathcal{S}}) - 2)P_\infty)$. The claim now follows by applying Proposition 2.1. \square

3 The Weierstrass semigroup for $P \in \tilde{\mathcal{S}}(\mathbb{F}_q)$

From Equations (3)-(6) and $(t)_{\tilde{\mathcal{S}}} = \sum_{P \in \tilde{\mathcal{S}}(\mathbb{F}_q), P \neq P_\infty} P - q^2P_\infty$, it is clear that the Weierstrass semigroup $H(P_\infty)$ contains the integers $q^2 - 2q_0q + q$, $q^2 - q_0q + q_0$, $q^2 - q + 2q_0$, q^2 , and $q^2 + 1$. This was already observed in [14], where in fact it was written that $H(P_\infty)$ is generated by these five numbers. To complete the results in [14], we give a formal proof of this claim in this section. Note that the fact that $\tilde{\mathcal{S}}(\mathbb{F}_q)$ forms one orbit under the action of the automorphism group of $\tilde{\mathcal{S}}$, directly implies that $H(P) = H(P_\infty)$ for any point $P \in \tilde{\mathcal{S}}(\mathbb{F}_q)$.

We will use the following standard terminology for numerical semigroups:

Definition. Let $Z \subset \mathbb{N}$ be a numerical semigroup. The set $G(Z) := \mathbb{N} \setminus Z$ is called the set of gaps of Z . The genus $g(Z)$, resp. multiplicity m_Z , resp. conductor c_Z of Z is defined to be

$$g(Z) := |\mathbb{N} \setminus Z|, \quad \text{resp.} \quad m_Z := \min\{z \in Z \mid z > 0\}, \quad \text{resp.} \quad c_Z := 1 + \max\{z \in \mathbb{N} \setminus Z\}.$$

Further, the Apéry set $\text{Ap}(Z)$ of Z is defined to be

$$\text{Ap}(Z) := \{z \in Z \mid z - m_Z \notin Z\}.$$

The Apéry set $\text{Ap}(Z)$ has cardinality m_Z and its elements form a complete set of representatives for the congruence classes of \mathbb{Z} modulo m_Z . Moreover, by definition of $\text{Ap}(Z)$, each representative is minimal among those lying in Z . As a consequence, the semigroup Z can conveniently be described as $Z = \{a + tm_Z \mid a \in \text{Ap}(Z), t \geq 0\}$. This also implies that the genus and conductor of Z can be deduced directly from m_Z and $\text{Ap}(Z)$:

$$g(Z) = \sum_{a \in \text{Ap}(Z)} \left\lfloor \frac{a}{m_Z} \right\rfloor \quad \text{and} \quad c_Z = 1 + \max\{z \in \text{Ap}(Z)\} - m_Z.$$

This implies that if $A \subset Z$ consists of m_Z elements that form a complete set of representatives for the congruence classes of \mathbb{Z} modulo m_Z , then $A = \text{Ap}(Z)$ if and only if $g(Z) = \sum_{a \in A} \left\lfloor \frac{a}{m_Z} \right\rfloor$. It is well known that $c_Z \leq 2g(Z)$. A numerical semigroup for which $c_Z = 2g(Z)$ is called a *symmetric* semigroup.

Now we return to the study of the semigroup $H(P_\infty)$. It will be convenient to introduce a notation for the five integers contained in it that we mentioned previously:

$$\mathfrak{g}_0 := q^2 - 2q_0q + q, \quad \mathfrak{g}_1 := q^2 - q_0q + q_0, \quad \mathfrak{g}_2 := q^2 - q + 2q_0, \quad \mathfrak{g}_3 := q^2, \quad \text{and} \quad \mathfrak{g}_4 := q^2 + 1.$$

Moreover, we write $H = \langle \mathfrak{g}_0, \mathfrak{g}_1, \mathfrak{g}_2, \mathfrak{g}_3, \mathfrak{g}_4 \rangle$ for the numerical semigroup generated by them. Since $\mathfrak{g}_0 < \mathfrak{g}_1 < \mathfrak{g}_2 < \mathfrak{g}_3 < \mathfrak{g}_4$, we see that $m_H = \mathfrak{g}_0$. Moreover, since $H \subseteq H(P_\infty)$, we have $g(H) \geq g(\tilde{\mathcal{S}})$. Using Apéry sets, we will actually show that $g(H) \leq g(\tilde{\mathcal{S}})$, which then immediately will imply that $H = H(P_\infty)$.

Lemma 3.1. Let $A := \{h\mathfrak{g}_1 + i\mathfrak{g}_2 + j\mathfrak{g}_3 + k\mathfrak{g}_4 \mid 0 \leq h \leq 1, 0 \leq i \leq q_0 - 1, 0 \leq j \leq q - 2q_0, 0 \leq k \leq q_0 - 1\}$. Then A is a complete set of representatives for the congruence classes of \mathbb{Z} modulo \mathfrak{g}_0 .

Proof. By definition of A , there are $2q_0^2(q - 2q_0 + 1) = \mathfrak{g}_0$ different four-tuples of parameters (h, i, j, k) describing the elements of A , so $|A| \leq \mathfrak{g}_0$. The lemma follows once we show that elements of the form $h\mathfrak{g}_1 + i\mathfrak{g}_2 + j\mathfrak{g}_3 + k\mathfrak{g}_4$ are pairwise distinct modulo \mathfrak{g}_0 for distinct four-tuples (h, i, j, k) satisfying

$$0 \leq h \leq 1, \quad 0 \leq i \leq q_0 - 1, \quad 0 \leq j \leq q - 2q_0, \quad 0 \leq k \leq q_0 - 1.$$

Now let $a = h\mathfrak{g}_1 + i\mathfrak{g}_2 + j\mathfrak{g}_3 + k\mathfrak{g}_4$ and $a' = h'\mathfrak{g}_1 + i'\mathfrak{g}_2 + j'\mathfrak{g}_3 + k'\mathfrak{g}_4$ be elements of A and suppose that $a \equiv a' \pmod{\mathfrak{g}_0}$. Recall that $q = 2q_0^2$ and that q divides \mathfrak{g}_0 . Hence the congruence $a \equiv a' \pmod{\mathfrak{g}_0}$ implies that $a \equiv a' \pmod{q_0}$. This in turn is equivalent to the congruence $k \equiv k' \pmod{q_0}$. Since $0 \leq k < q_0$ and $0 \leq k' < q_0$, we see that $k = k'$. Now working modulo $2q_0$, implies that $q_0h \equiv q_0h' \pmod{2q_0}$. Since $0 \leq h \leq 1$ and $0 \leq h' \leq 1$, we obtain $h = h'$. Similarly working modulo q , implies that $i = i'$. At this point, we know that $j\mathfrak{g}_3 \equiv j'\mathfrak{g}_3 \pmod{\mathfrak{g}_0}$. Dividing by q , we obtain that $j \equiv j' \pmod{q - 2q_0 + 1}$, whence $j \equiv j' \pmod{q - 2q_0 + 1}$ and therefore $j = j'$. \square

Theorem 3.2. For any $P \in \tilde{\mathcal{S}}(\mathbb{F}_q)$, we have $H(P) = \langle q^2 - 2q_0q + q, q^2 - q_0q + q_0, q^2 - q + 2q_0, q^2, q^2 + 1 \rangle$.

Proof. As before, let H denote the semigroup $\langle q^2 - 2q_0q + q, q^2 - q_0q + q_0, q^2 - q + 2q_0, q^2, q^2 + 1 \rangle$. As observed before, we know that $H(P)$ contains H and hence $g(H) \geq g(\tilde{\mathcal{S}})$. From Lemma 3.1, we see that $g(H) \leq \sum_{a \in A} \lfloor \frac{a}{\mathfrak{g}_0} \rfloor$. Note that equality holds if and only if $A = \text{Ap}(H)$.

For simplicity let us denote $m = 2q_0^2 - 2q_0 + 1$, so that $\mathfrak{g}_0 = 2q_0^2m$. Then:

$$\begin{aligned} \sum_{a \in A} \left\lfloor \frac{a}{\mathfrak{g}_0} \right\rfloor &= \sum_{a \in A} \frac{a - a \bmod \mathfrak{g}_0}{\mathfrak{g}_0} = \frac{1}{\mathfrak{g}_0} \sum_{a \in A} a - \frac{1}{\mathfrak{g}_0} \sum_{r=0}^{\mathfrak{g}_0-1} r \\ &= \frac{1}{\mathfrak{g}_0} \sum_{h=0}^1 \sum_{i=0}^{q_0-1} \sum_{j=0}^{m-1} \sum_{k=0}^{q_0-1} (h\mathfrak{g}_1 + i\mathfrak{g}_2 + j\mathfrak{g}_3 + k\mathfrak{g}_4) - \frac{1}{\mathfrak{g}_0} \cdot \frac{\mathfrak{g}_0(\mathfrak{g}_0-1)}{2} \\ &= \frac{1}{\mathfrak{g}_0} \left(q_0^2 m \mathfrak{g}_1 + 2q_0 m \frac{q_0(q_0-1)}{2} \mathfrak{g}_2 + 2q_0^2 \frac{m(m-1)}{2} \mathfrak{g}_3 + 2q_0 m \frac{q_0(q_0-1)}{2} \mathfrak{g}_4 \right) - \frac{\mathfrak{g}_0-1}{2} \\ &= g(\tilde{\mathcal{S}}). \end{aligned}$$

Hence $g(H) \leq g(\tilde{\mathcal{S}})$. Combining all the above, we see $g(H) = g(\tilde{\mathcal{S}})$, which implies that $H(P) = H$. \square

Corollary 3.3. *Let $P \in \tilde{\mathcal{S}}(\mathbb{F}_q)$ and as before, let $A := \{h\mathfrak{g}_1 + i\mathfrak{g}_2 + j\mathfrak{g}_3 + k\mathfrak{g}_4 \mid 0 \leq h \leq 1, 0 \leq i \leq q_0 - 1, 0 \leq j \leq q - 2q_0, 0 \leq k \leq q_0 - 1\}$. Then $\text{Ap}(H(P)) = A$. Further, $H(P)$ is a symmetric semigroup.*

Proof. The statement $\text{Ap}(H(P)) = A$ follows from the proof of Theorem 3.2. In particular, the largest gap of $H(P_\infty)$ is

$$\max\{z \in A\} - \mathfrak{g}_0 = \mathfrak{g}_1 + (q_0 - 1)\mathfrak{g}_2 + (q - 2q_0)\mathfrak{g}_3 + (q_0 - 1)\mathfrak{g}_4 - \mathfrak{g}_0 = 2g(\tilde{\mathcal{S}}) - 1.$$

This means that the conductor of $H(P)$ is $2g(\tilde{\mathcal{S}})$ and hence that the semigroup is symmetric. \square

We will later see that any $P \in \tilde{\mathcal{S}}(\mathbb{F}_q)$ is a Weierstrass point by computing the Weierstrass semigroups for all points on $\tilde{\mathcal{S}}$. However, using the symmetry of $H(P_\infty)$ that we just proved and [12, Proposition 50], we can already conclude this now.

4 The Weierstrass semigroup for $P \in \tilde{\mathcal{S}}(\mathbb{F}_{q^4}) \setminus \tilde{\mathcal{S}}(\mathbb{F}_q)$

Let P be a point in $\tilde{\mathcal{S}}(\mathbb{F}_{q^4}) \setminus \tilde{\mathcal{S}}(\mathbb{F}_q)$ with (x, y, t) -coordinates (a, b, c) . The aim of this section is to prove the following theorem.

Theorem 4.1. *The Weierstrass semigroup $H(P)$ is generated by the following $3q_0 + 3$ elements:*

$$\begin{aligned} \mathfrak{g}_0 &:= q^2 - q + 1, \\ \mathfrak{g}_1 &:= q^2 - 2q_0 + 1, \\ \mathfrak{g}_2 &:= q^2 - q_0 + 1, \\ \mathfrak{g}_3 &:= q^2, \\ \mathfrak{g}_4 &:= q^2 + 1, \\ \mathfrak{f}_i &:= (i+1)q_0g_0 - ig_4 - 1 \quad i = 0, \dots, 2q_0 - 2, \\ \mathfrak{h}_j &:= (2j+1)q_0g_0 - jg_4 - q_0 \quad j = 0, \dots, q_0 - 2. \end{aligned}$$

The strategy of the proof will be to first show that the $3q_0 + 3$ elements mentioned in Theorem 4.1 are in $H(P)$, then showing that the semigroup generated by them has $g(\tilde{\mathcal{S}})$ gaps. As before we write $Q = \text{pr}(P)$, the point of \mathcal{S} lying under P in the cover $\text{pr} : \tilde{\mathcal{S}} \rightarrow \mathcal{S}$.

Lemma 4.2. *There exists a divisor $D \geq 0$ on \mathcal{S} satisfying $\Phi^i(Q) \notin \text{Supp}(D)$ for $i = 0, \dots, 3$ such that:*

- a) *The divisor $q_0 \sum_{i=0}^3 \Phi^i(Q) + D - (q + 2q_0 + 1)Q_\infty$ is principal.*
- b) *The divisor $((2i + 1)qq_0 + q_0)Q + (iq + q_0)\Phi(Q) + (q_0 - 1 - i)\Phi^3(Q) + D - ((2i + 1)q_0 - i)(q + 2q_0 + 1)Q_\infty$ is principal for all $i \geq 0$.*

Proof. a) Using the transitivity of the automorphism group of \mathcal{S} on $\mathcal{S}(\mathbb{F}_{q^4}) \setminus \mathcal{S}(\mathbb{F}_q)$, we may assume without loss of generality that $a \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$. Define $\alpha := (a^q + a)^{q_0} \in \mathbb{F}_q$ and let

$$F_0 := \alpha(a^{q_0}x + y + a^{q_0+1} + b) + x(x + a)^{2q_0} + (y + b)^{2q_0}. \quad (8)$$

Observe that F_0 is defined over \mathbb{F}_q , since Equation (8) can also be expressed as

$$F_0 = a^{(q+1)q_0}x + \alpha y + y^{2q_0} + x^{2q_0+1} + (\alpha(a^{q_0+1} + b) + b^{2q_0}),$$

where $(\alpha(a^{q_0+1} + b) + b^{2q_0})$ lies in \mathbb{F}_q because

$$\begin{aligned} (\alpha(a^{q_0+1} + b) + b^{2q_0})^q + (\alpha(a^{q_0+1} + b) + b^{2q_0}) &= \alpha(a^{(q_0+1)q} + a^{q_0+1}) + \alpha(b^q + b) + (b^q + b)^{2q_0} = \\ \alpha(a^{(q_0+1)q} + a^{q_0+1}) + \alpha a^{q_0}(a^q + a) + (a^{q_0}(a^q + a))^{2q_0} &= \alpha^2 a^q + a^q \alpha^2 = 0. \end{aligned}$$

Recalling Equation (4), we have

$$v_Q(F_0) \geq \min\{v_Q(a^{q_0}x + y + a^{q_0+1} + b), v_Q(x(x + a)^{2q_0}), v_Q((y + b)^{2q_0})\} = v_Q(\tilde{y}_Q) = q_0. \quad (9)$$

Inequality (9) is in fact an equality, since $x(x + a)^{2q_0}$ and $(y + b)^{2q_0}$ have valuation at Q larger than or equal to $2q_0$. The valuation of F_0 at $\Phi^i(Q)$, for $i = 0, \dots, 3$ can be obtained from Inequality (9) and from the fact that F_0 is defined over \mathbb{F}_q , as

$$v_{\Phi^i(Q)}(F_0) = v_Q(\Phi^{-i}(F_0)) = v_Q(F_0) = q_0.$$

Finally, observe that Q_∞ is the only pole of F_0 and that

$$v_{Q_\infty}(F_0) = v_{Q_\infty}(x^{2q_0+1} + y^{2q_0}) = v_{Q_\infty}(z) = -(q + 2q_0).$$

Therefore there exists an effective divisor \tilde{D} with support not containing $\Phi^i(Q)$ for $i = 0, \dots, 3$ such that

$$(F_0)_S = q_0 \sum_{i=0}^3 \Phi^i(Q) + \tilde{D} - (q + 2q_0)Q_\infty.$$

The conclusion follows after setting $D := \tilde{D} + Q_\infty$.

b) Let us write

$$D_i := ((2i + 1)qq_0 + q_0)Q + (iq + q_0)\Phi(Q) + (q_0 - 1 - i)\Phi^3(Q) + D - ((2i + 1)q_0 - i)(q + 2q_0 + 1)Q_\infty.$$

Using part a) and Equation (6), we observe that $D_i = (F_0)_S + (\tilde{w}_Q^{(2i+1)q_0})_S - (\tilde{w}_{\Phi^i(Q)}^{i+1})_S$. \square

Proposition 4.3. *The integers $g_0, g_1, g_2, g_3, g_4, f_0, \dots, f_{2q_0-2}, h_0, \dots, h_{q_0-2}$ are elements of $H(P)$.*

Proof. The first five values g_0, g_1, g_2, g_3, g_4 can be obtained as pole orders at P of the functions $\gamma_0 = \tilde{w}_Q \cdot \pi_P^{-1}$, $\gamma_1 = \tilde{w}_{\Phi^3(Q)} \cdot \pi_P^{-1}$, $\gamma_2 = \tilde{y}_Q \cdot \pi_P^{-1}$, $\gamma_3 = \tilde{x}_Q \cdot \pi_P^{-1}$, $\gamma_4 = \pi_P^{-1}$ respectively. It can be easily seen from Equations (4), (6), and (7) that such functions are regular outside P .

Let us prove now that f_i is in $H(P)$ for $i = 0, \dots, 2q_0 - 2$. For each $i = 0, \dots, 2q_0 - 2$ define:

$$\alpha_i := \frac{\tilde{w}_Q^{(i+1)q_0} \cdot \tilde{w}_{\Phi^2(Q)}}{\tilde{w}_{\Phi(Q)}^{(i+1)}}.$$

By Equation (6), the divisor of α_i in \mathcal{S} is

$$(\alpha_i)_{\mathcal{S}} = ((i+1)qq_0 + 1)Q + (q - (i+1)q_0)\Phi^2(Q) + (2q_0 - (i+1))\Phi^3(Q) - ((i+1)q_0 - i)(q + 2q_0 + 1)Q_{\infty}.$$

Note that the assumption $i \leq 2q_0 - 2$ implies that $(q - (i+1)q_0) \geq q_0 > 0$ and $(2q_0 - (i+1)) \geq 1$. Since P is unramified and P_{∞} is totally ramified in the cover $\text{pr} : \tilde{\mathcal{S}} \rightarrow \mathcal{S}$, we have:

$$\begin{cases} v_P(\alpha_i) = v_Q(\alpha_i) = (i+1)qq_0 + 1, \\ v_{P_{\infty}}(\alpha_i) = v_{Q_{\infty}}(\alpha_i) \cdot (q - 2q_0 + 1) = -((i+1)q_0 - i)(q^2 + 1), \\ v_R(\alpha_i) \geq 0 \end{cases} \quad \text{for } R \neq P, P_{\infty}.$$

Therefore the function $\beta_i := \alpha_i / \pi_P^{(i+1)q_0 - i}$ satisfies

$$\begin{cases} v_P(\beta_i) = ((i+1)qq_0 + 1) - ((i+1)q_0 - i)(q^2 + 1) = -f_i, \\ v_{P_{\infty}}(\beta_i) = -((i+1)q_0 - i)(q^2 + 1) + ((i+1)q_0 - i)(q^2 + 1) = 0, \\ v_R(\beta_i) = v_R(\alpha_i) \geq 0 \end{cases} \quad \text{for } R \neq P, P_{\infty}.$$

Hence, β_i is a function having a unique pole in P of order f_i .

Let us finally prove that h_i is in $H(P)$ for $i = 0, \dots, q_0 - 2$. By Lemma 4.2 part b), there exists a function $\delta_i \in \mathbb{F}_{q^4}(\mathcal{S})$ and some effective divisor D such that

$$(\delta_i)_{\mathcal{S}} = D_i := ((2i+1)qq_0 + q_0)Q + (iq + q_0)\Phi(Q) + (q_0 - 1 - i)\Phi^3(Q) + D - ((2i+1)q_0 - i)(q + 2q_0 + 1)Q_{\infty}.$$

The assumption $i \leq q_0 - 2$ guarantees that δ_i is regular outside Q_{∞} . Recalling Equation (7), the function $\eta_i := \delta_i / \pi_P^{(2i+1)q_0 - i}$ satisfies

$$\begin{cases} v_P(\eta_i) = ((2i+1)qq_0 + q_0) - ((2i+1)q_0 - i)(q^2 + 1) = -h_i, \\ v_{P_{\infty}}(\eta_i) = -((2i+1)q_0 - i)(q^2 + 1) + ((2i+1)q_0 - i)(q^2 + 1) = 0, \\ v_R(\eta_i) = v_R(\delta_i) \geq 0 \end{cases} \quad \text{for } R \neq P, P_{\infty}.$$

Hence, η_i is a function having a unique pole in P of order h_i . □

A consequence of Proposition 4.3 is that the numerical semigroup

$$H := \langle g_0, g_1, g_2, g_3, g_4, f_i, h_j \mid i = 0, \dots, 2q_0 - 2 \text{ and } j = 0, \dots, q_0 - 2 \rangle$$

is contained in $H(P)$. To show that equality $H = H(P)$ holds, from which Theorem 4.1 follows, it suffices to prove that H has the same genus of $H(P)$, namely that $g(H) = g(H(P)) = \frac{1}{2}q(q-1)^2$.

Observe that g_0 is the smallest generator of H ; in particular g_0 is the multiplicity of H and the Apéry set $\text{Ap}(H)$ consists of g_0 distinct elements. We now want to find a map $\varphi : \{0, \dots, g_0 - 1\} \rightarrow \mathbb{N}$ such that any $a \in \text{Ap}(H)$ can be expressed as $a = \varphi(i)g_0 + i$ for some $i \in \{0, \dots, g_0 - 1\}$.

Lemma 4.4. Let $i \in \{0, \dots, \frac{q(q-2)}{2}\}$ and write $i = lq + kq_0 + j$ with $j \in \{0, \dots, q_0 - 1\}$, $k \in \{0, \dots, 2q_0 - 1\}$ and $l \geq 0$. Define:

$$\varphi_1(i) = \begin{cases} l & \text{if } j = 0 \text{ and } k = 0, \\ l + 1 + \max\{q - q_0(k + 2l + 2), 0\} & \text{if } j = 0 \text{ and } k \neq 0, \\ l + 1 + \max\{q - q_0(\lceil \frac{k}{2} \rceil + j + l + 1), 0\} & \text{if } j \neq 0. \end{cases}$$

Then $\varphi_1(i)g_0 + i$ belongs to H .

Proof. Clearly, if $j = 0$ and $k = 0$, then $\varphi_1(i)g_0 + i = lg_0 + lq = lg_4$ belongs to H .

We now consider the case of $j = 0, k \neq 0$ and $\max\{q - q_0(k + 2l + 2), 0\} > 0$. Note that this last condition implies that $2q_0 - k - 2l - 2 > 0$. Then:

$$\begin{aligned} \varphi_1(i)g_0 + i &= (l + 1 + q - q_0(k + 2l + 2))g_0 + lq + kq_0 \\ &= (l + 1)g_1 + (l + 1)(2q_0 - q) + (q - q_0(k + 2l + 2))g_0 + lq + kq_0 \\ &= (l + 1)g_1 + (2q_0 - k - 2l - 2)h_0. \end{aligned}$$

Hence $\varphi_1(i)g_0 + i$ belongs to H .

Let us consider now the case of $j = 0, k \neq 0$ and $\max\{q - q_0(k + 2l + 2), 0\} = 0$. This last condition can be expressed equivalently as $k \geq 2q_0 - 2l - 2$. Define $c := 2q_0 - k$, so that $1 \leq c \leq 2l + 2$. Then

$$\varphi_1(i)g_0 + i = (l + 1)g_0 + lq + kq_0 = (l + 1)g_4 - cq_0. \quad (10)$$

We can distinguish three cases.

- If $c \leq l + 1$, then Equation (10) can be written as $(l + 1 - c)g_4 + cg_2$.
- If $c > l + 1$ and c is even, then Equation (10) can be written as $(l + 1 - \frac{c}{2})g_4 + \frac{c}{2}g_1$.
- If $c > l + 1$ and c is odd (and in particular $c \leq 2l + 1$), then Equation (10) can be written as $(l - \frac{c-1}{2})g_4 + \frac{c-1}{2}g_1 + g_2$.

In all the three cases $\varphi_1(i)g_0 + i$ belongs to H .

Let us now assume $j \neq 0$ and $\max\{q - q_0(\lceil \frac{k}{2} \rceil + j + l + 1), 0\} > 0$; this last condition implies $2q_0 - \lceil \frac{k}{2} \rceil - j - l - 1 > 0$. Also,

$$\begin{aligned} \varphi_1(i)g_0 + i &= \left(l + 1 + q - q_0 \left(\left\lceil \frac{k}{2} \right\rceil + j + l + 1 \right) \right) g_0 + lq + kq_0 + j \\ &= f_{2q_0 - \lceil \frac{k}{2} \rceil - j - l - 2} + 2q_0q^2 + 2q_0 - \left\lceil \frac{k}{2} \right\rceil (q^2 + 1) - jq^2 - q - q^2 + kq_0. \end{aligned} \quad (11)$$

If k is even, then Equation (11) is equal to

$$f_{2q_0 - \frac{k}{2} - j - l - 2} + \left(q_0 - 1 - \frac{k}{2} \right) g_1 + (q_0 - 1 - j)g_3 + g_2.$$

If k is odd, then Equation (11) is equal to

$$f_{2q_0 - \frac{k+1}{2} - j - l - 2} + \left(q_0 - \frac{k+1}{2} \right) g_1 + (q_0 - 1 - j)g_3.$$

In both cases $\varphi_1(i)g_0 + i$ belongs to H .

Let us finally consider the case of $j \neq 0$ and $\max\{q - q_0(\lceil \frac{k}{2} \rceil + j + l + 1), 0\} = 0$; this last condition implies $k \geq 4q_0 - 2j - 2l - 3$. Define $c := 2q_0 - 1 - k$, so that $0 \leq c \leq 2(l + 1 + j - q_0)$. Then

$$\varphi_1(i)g_0 + i = (l + 1)g_0 + lq + kq_0 + j = (l + 1 + j - q_0)g_4 + (q_0 - j)g_3 - cq_0. \quad (12)$$

Let us distinguish three cases.

- If $c \leq l + 1 + j - q_0$, then Equation (12) can be written as $(l + 1 + j - q_0 - c)g_4 + (q_0 - j)g_3 + cg_2$.
- If $c > l + 1 + j - q_0$ and c is even, then Equation (12) can be written as

$$\left(l + 1 + j - q_0 - \frac{c}{2}\right)g_4 + (q_0 - j)g_3 + \frac{c}{2}g_1.$$

- If $c > l + 1 + j - q_0$ and c is odd (and in particular $1 \leq c \leq 1 + 2(l + j - q_0)$), then Equation (12) can be written as

$$\left(l + j - q_0 - \frac{c - 1}{2}\right)g_4 + (q_0 - j)g_3 + \frac{c - 1}{2}g_1 + g_2.$$

In all the three cases $\varphi_1(i)g_0 + i$ belongs to H . □

Lemma 4.5. *Let $i \in \{\frac{q(q-2)}{2} + 1, \dots, g_0 - 1\}$. Denote $i' = g_0 - 1 - i$ and write $i' = lq + kq_0 + j$ with $j \in \{0, \dots, q_0 - 1\}$, $k \in \{0, \dots, 2q_0 - 1\}$ and $l \geq 0$. Define:*

$$\varphi_2(i) = \begin{cases} q - l - 1 - \max\{q - q_0(k + 2l + 1), 0\} & \text{if } j = q_0 - 1, \\ q - l - 1 - \max\{q - q_0(\lceil \frac{k}{2} \rceil + j + l + 1), 0\} & \text{if } j \neq q_0 - 1. \end{cases}$$

Then $\varphi_2(i)g_0 + i$ belongs to H .

Proof. We first prove that if $\varphi_2(i) = q - l - 1$, then $\varphi_2(i)g_0 + i$ belongs to H .

$$\begin{aligned} \varphi_2(i)g_0 + i &= (q - l - 1)g_0 + g_0 - 1 - i' \\ &= (q - l - 1)g_4 + lq - (lq + kq_0 + j) \\ &= (q - l - k - j - 1)g_4 + kg_2 + jg_3. \end{aligned}$$

By assumption on i , the condition $l \leq \frac{q}{2} - 1$ holds, so

$$\begin{aligned} q - l - k - j - 1 &\geq q - \left(\frac{q}{2} - 1\right) - (2q_0 - 1) - (q_0 - 1) - 1 \\ &= \frac{q}{2} - 3q_0 + 2 = (q_0 - 1)(q_0 - 2) \geq 0. \end{aligned}$$

Therefore $\varphi_2(i)g_0 + i$ is in H .

We now consider the case of $j = q_0 - 1$ and $\max\{q - q_0(k + 2l + 1), 0\} > 0$. Note that this last condition implies $l < q_0$. Define $h_{q_0-1} := (2q_0 - 1)q_0g_0 - (q_0 - 1)g_4 - q_0$, coherently with the definition of h_j for $j = 0, \dots, q_0 - 2$, and observe that $h_{q_0-1} = f_{q_0-1} + f_{q_0-2} + (q_0 - 2)g_3 \in H$. Then:

$$\begin{aligned} \varphi_2(i)g_0 + i &= (q_0(k + 2l + 1) - l - 1)g_0 + g_0 - 1 - i' \\ &= (q_0(k + 2l + 1) - l)g_0 - 1 - (lq + kq_0 + j) \\ &= h_l + kh_0. \end{aligned}$$

Hence $\varphi_2(i)g_0 + i$ belongs to H .

We finally prove that if $j \neq q_0 - 1$ and $\max\{q - q_0(\lceil \frac{k}{2} \rceil + j + l + 1), 0\} > 0$, then $\varphi_2(i)g_0 + i$ belongs to H . Note that the condition on the maximum implies $\lceil \frac{k}{2} \rceil + j + l < 2q_0 - 1$. Then:

$$\begin{aligned}\varphi_2(i)g_0 + i &= \left(q_0 \left(\left\lceil \frac{k}{2} \right\rceil + j + l + 1 \right) - l - 1 \right) g_0 + g_0 - 1 - i' \\ &= \left(q_0 \left(\left\lceil \frac{k}{2} \right\rceil + j + l + 1 \right) - l \right) g_0 - 1 - (lq + kq_0 + j) \\ &= f_{\lceil \frac{k}{2} \rceil + j + l} + \left(\left\lceil \frac{k}{2} \right\rceil + j \right) q^2 - kq_0 + \left\lceil \frac{k}{2} \right\rceil.\end{aligned}\tag{13}$$

If k is even, then Equation (13) is equal to $f_{\frac{k}{2} + j + l} + \frac{k}{2}g_1 + jg_3$. If k is odd (and in particular $k \geq 1$), then Equation (13) is equal to $f_{\frac{k+1}{2} + j + l} + \frac{k-1}{2}g_1 + jg_3 + g_2$. In both cases $\varphi_2(i)g_0 + i$ belongs to H . \square

Define the map $\varphi : \{0, \dots, g_0 - 1\} \rightarrow \mathbb{N}$ by

$$\varphi(i) = \begin{cases} \varphi_1(i) & \text{if } i \in \{0, \dots, \frac{q(q-2)}{2}\}, \\ \varphi_2(i) & \text{if } i \in \{\frac{q(q-2)}{2} + 1, \dots, g_0 - 1\}.\end{cases}$$

Combining Lemma 4.4 and Lemma 4.5, it follows that $\varphi(i)g_0 + i$ belongs to H for all $i \in \{0, \dots, g_0 - 1\}$.

Lemma 4.6. *The Apéry set of $H(P)$ is $\text{Ap}(H(P)) = \{\varphi(i)g_0 + i \mid i = 0, \dots, g_0 - 1\}$.*

Proof. It is sufficient to show that $\sum_{i=0}^{g_0-1} \varphi(i) = g(\tilde{S})$. Indeed, the number of gaps of H is then shown to be at most $g(\tilde{S})$, which implies that $H = H(P)$, since we already know that $H \subseteq H(P)$. But in that case, the mentioned set is the Apéry set of $H(P)$, since otherwise $g(H(P))$ would be strictly less than $\sum_{i=0}^{g_0-1} \varphi(i)$.

First, let us prove that

$$\varphi(i) + \varphi((q-1)^2 - i) = q - 1 \quad \text{for } i \in \{0, \dots, (q-1)^2\}.\tag{14}$$

Observe that i is in range $0, \dots, \frac{q(q-2)}{2}$ if and only if $(q-1)^2 - i$ is in range $\frac{q(q-2)}{2} + 1, \dots, (q-1)^2$. Therefore it is enough to prove Equation (14) for $i \in \{0, \dots, \frac{q(q-2)}{2}\}$ only; the case $i \in \{\frac{q(q-2)}{2} + 1, \dots, (q-1)^2\}$ follows by symmetry.

For $i \in \{0, \dots, \frac{q(q-2)}{2}\}$ we have $\varphi(i) = \varphi_1(i)$ where φ_1 is defined as in Lemma 4.4 and $\varphi((q-1)^2 - i) = \varphi_2((q-1)^2 - i)$ where φ_2 is defined as in Lemma 4.5. Write $i = lq + kq_0 + j$ with $j \in \{0, \dots, q_0 - 1\}$, $k \in \{0, \dots, 2q_0 - 1\}$ and $l \geq 0$. Then:

$$g_0 - 1 - ((q-1)^2 - i) = q - 1 + i = l'q + k'q_0 + j'$$

with

$$\begin{cases} l' = l, k' = 2q_0 - 1, j' = q_0 - 1 & \text{if } j = 0 \text{ and } k = 0, \\ l' = l + 1, k' = k - 1, j' = q_0 - 1 & \text{if } j = 0 \text{ and } k \neq 0, \\ l' = l + 1, k' = k, j' = j - 1 & \text{if } j \neq 0, \end{cases}$$

satisfying $j' \in \{0, \dots, q_0 - 1\}$, $k' \in \{0, \dots, 2q_0 - 1\}$ and $l' \geq 0$. Consequently:

$$\begin{aligned}
q - 1 - \varphi((q - 1)^2 - i) &= q - 1 - \varphi_2((q - 1)^2 - i) \\
&= \begin{cases} l' + \max\{q - q_0(k' + 2l' + 1), 0\} & \text{if } j' = q_0 - 1, \\ l' + \max\{q - q_0(\lceil \frac{k'}{2} \rceil + j' + l' + 1), 0\} & \text{if } j' \neq q_0 - 1. \end{cases} \\
&= \begin{cases} l + \max\{-2q_0l, 0\} & \text{if } j = 0 \text{ and } k = 0, \\ l + 1 + \max\{q - q_0(k + 2l + 2), 0\} & \text{if } j = 0 \text{ and } k \neq 0, \\ l + 1 + \max\{q - q_0(\lceil \frac{k}{2} \rceil + j + l + 1), 0\} & \text{if } j \neq 0. \end{cases} \\
&= \varphi_1(i) = \varphi(i).
\end{aligned}$$

Equation (14) implies that

$$\sum_{i=0}^{(q-1)^2} \varphi(i) = \sum_{i=0}^{\frac{q(q-2)}{2}} (q - 1) = \left(\frac{q(q-2)}{2} + 1 \right) (q - 1). \quad (15)$$

Now let us assume $i \in \{(q - 1)^2 + 1, \dots, g_0 - 1\}$. Since $i' = g_0 - 1 - i$ ranges between 0 and $q - 2$, we can express i' as $i' = kq_0 + j$ with $j \in \{0, \dots, q_0 - 1\}$ and $k \in \{0, \dots, 2q_0 - 1\}$. In particular:

$$\varphi(i) = \varphi_2(i) = \begin{cases} q_0(k + 1) - 1 & \text{if } j = q_0 - 1, \\ q_0(\lceil \frac{k}{2} \rceil + j + 1) - 1 & \text{if } j \neq q_0 - 1. \end{cases}$$

Then:

$$\sum_{i=(q-1)^2+1}^{g_0-1} \varphi(i) = \sum_{k=0}^{2q_0-2} (q_0(k + 1) - 1) + \sum_{k=0}^{2q_0-1} \sum_{j=0}^{q_0-2} \left(q_0 \left(\lceil \frac{k}{2} \rceil + j + 1 \right) - 1 \right) = \frac{q^2}{2} - \frac{3}{2}q + 1. \quad (16)$$

Combining Equations (15) and (16) we obtain:

$$\sum_{i=0}^{g_0-1} \varphi(i) = \left(\frac{q(q-2)}{2} + 1 \right) (q - 1) + \left(\frac{q^2}{2} - \frac{3}{2}q + 1 \right) = \frac{1}{2}q(q - 1)^2 = g(\tilde{\mathcal{S}}).$$

□

Note that from the proof of the above lemma, we immediately conclude that $H = H(P)$, proving Theorem 4.1.

Corollary 4.7. *Let $P \in \tilde{\mathcal{S}}(\mathbb{F}_{q^4}) \setminus \tilde{\mathcal{S}}(\mathbb{F}_q)$. Then the Weierstrass semigroup $H(P)$ is symmetric.*

Proof. For $i = (q - 1)^2$ we have $a := \varphi(i)g_0 + i = q^2(q - 1) = 2g(\tilde{\mathcal{S}}) - 1 + g_0$. Since a is an element of $\text{Ap}(H(P))$, it follows that $a - g_0 = 2g(\tilde{\mathcal{S}}) - 1$ is a gap of H . By the Weierstrass gap theorem $2g(\tilde{\mathcal{S}}) - 1$ is the largest possible gap of $H(P)$, hence $H(P)$ has conductor $2g(\tilde{\mathcal{S}})$. □

As for $P \in \tilde{\mathcal{S}}(\mathbb{F}_q)$, we can now conclude from [12, Proposition 50] that any $P \in \tilde{\mathcal{S}}(\mathbb{F}_{q^4}) \setminus \tilde{\mathcal{S}}(\mathbb{F}_q)$ is a Weierstrass point of $\tilde{\mathcal{S}}$. In the next section, we will show that any $P \notin \tilde{\mathcal{S}}(\mathbb{F}_{q^4})$ has the same Weierstrass semigroup. This will imply that the set of Weierstrass points of $\tilde{\mathcal{S}}$ is equal to $\tilde{\mathcal{S}}(\mathbb{F}_{q^4})$.

5 The Weierstrass semigroup for $P \notin \tilde{\mathcal{S}}(\mathbb{F}_{q^4})$

Let now $P \in \tilde{\mathcal{S}}(\overline{\mathbb{F}}_q) \setminus \tilde{\mathcal{S}}(\mathbb{F}_{q^4})$. Let $a, b, c \in \overline{\mathbb{F}}_q$ be the affine coordinates of P and as before denote with Q , the point of \mathcal{S} with affine coordinates a and b . It will also be convenient to use the expression $A := a^q + a$.

From Corollary 2.2, the gap sequence $G(P)$ at P can be computed by constructing $g = g(\tilde{\mathcal{S}})$ functions f_1, \dots, f_g having pairwise distinct valuations at P and such that $f_i \in L((2g(\tilde{\mathcal{S}}) - 2)P_\infty)$ for all $i = 1, \dots, g$.

To this aim note first that $\tilde{x}_Q := x - a$ is a local parameter at P and $\tilde{x}_Q, \tilde{y}_Q, \tilde{z}_Q, \tilde{w}_Q \in L((q^3 - 2q^2 + q - 2)P_\infty)$ by Equations (3)-(6). Also, recall from Equation (7), that there exists a function $\pi_P \in \overline{\mathbb{F}}_q(\tilde{\mathcal{S}})$ such that $(\pi_P)_{\tilde{\mathcal{S}}} = q^2P + \Phi^4(P) - (q^2 + 1)P_\infty$.

In the following, the local power series expansions of \tilde{y}_Q, \tilde{z}_Q and \tilde{w}_Q at P (with respect to the local parameter \tilde{x}_Q) is computed. First of all note that from $Q \in \mathcal{S}$ and $y^q + y = x^{q_0}(x^q + x)$ we have

$$(y + b)^q + (y + b) = a^{q_0}(a^q + a) + x^{q_0}(x^q + x) = a^{q_0}\tilde{x}_Q + (a^q + a)\tilde{x}_Q^{q_0} + \tilde{x}_Q^{q_0+1} + a^{q_0}\tilde{x}_Q^q + \tilde{x}_Q^{q+q_0},$$

so that

$$(y + b) = a^{q_0}\tilde{x}_Q + A\tilde{x}_Q^{q_0} + \tilde{x}_Q^{q_0+1} + A^{q_0}\tilde{x}_Q^q + \tilde{x}_Q^{q+q_0} + A^q\tilde{x}_Q^{q_0q} + \tilde{x}_Q^{qq_0+q} + A^{q_0q} + \tilde{x}_Q^{q^2} + O(\tilde{x}_Q^{q^2+1}). \quad (17)$$

Hence, from $\tilde{y}_Q = (y + b) + a^{q_0}\tilde{x}_Q$ and Equation (17) we get

$$\tilde{y}_Q = A\tilde{x}_Q^{q_0} + \tilde{x}_Q^{q_0+1} + A^{q_0}\tilde{x}_Q^q + \tilde{x}_Q^{q+q_0} + A^q\tilde{x}_Q^{q_0q} + \tilde{x}_Q^{qq_0+q} + A^{q_0q} + \tilde{x}_Q^{q^2} + O(\tilde{x}_Q^{q^2+1}). \quad (18)$$

Since $A \neq 0$ from $P \notin \tilde{\mathcal{S}}(\mathbb{F}_q)$, while \tilde{x}_Q , seen as a function on \mathcal{S} , is also a local parameter at Q , we get that $v_P(\tilde{y}_Q) = v_Q(\tilde{y}_Q) = q_0$ as anticipated in Equation (4).

Now, $\tilde{z}_Q = a^{2q_0}(\tilde{x}_Q + a) + (\tilde{x}_Q + a)^{2q_0+1} + (y + b)^{2q_0} = a\tilde{x}_Q^{2q_0} + \tilde{x}_Q^{2q_0+1} + (y + b)^{2q_0}$, which combined with Equation (17) gives

$$\tilde{z}_Q = A\tilde{x}_Q^{2q_0} + \tilde{x}_Q^{2q_0+1} + A^{2q_0}\tilde{x}_Q^q + \tilde{x}_Q^{q+2q_0} + A^q\tilde{x}_Q^{2q_0q} + \tilde{x}_Q^{2q_0q+q} + A^{2q_0q}\tilde{x}_Q^{q^2} + O(\tilde{x}_Q^{q^2+1}). \quad (19)$$

The computation above yields that $v_P(\tilde{z}_Q) = v_Q(\tilde{z}_Q) = 2q_0$, which is again consistent with Equation (5). Finally, from $z = \tilde{z}_Q + a^{2q_0}x + b^{2q_0}$, $b^q + b = a^{q_0}(a^q + a)$, Equations (17) and (19), we have

$$\begin{aligned} \tilde{w}_Q &= a^q\tilde{z}_Q + b^{2q_0}(\tilde{x}_Q + a) + (\tilde{z}_Q + a^{2q_0}x + b^{2q_0})^{2q_0} + xy^{2q_0} + b^2 + a^{2q_0+2} \\ &= a^q\tilde{z}_Q + \tilde{z}_Q^{2q_0} + a^{2q_0}\tilde{x}_Q^{2q_0} + \tilde{x}_Q(y + b)^{2q_0} + a(y + b)^{2q_0} \\ &= a^q(A\tilde{x}_Q^{2q_0} + \tilde{x}_Q^{2q_0+1} + A^{2q_0}\tilde{x}_Q^q + \tilde{x}_Q^{q+2q_0} + A^q\tilde{x}_Q^{2q_0q} + \tilde{x}_Q^{2q_0q+q} + A^{2q_0q}\tilde{x}_Q^{q^2}) \\ &\quad + (A^{2q_0}\tilde{x}_Q^{2q} + \tilde{x}_Q^{2q+2q_0} + A^{2q}\tilde{x}_Q^{2q_0q} + \tilde{x}_Q^{2q_0q+2q}) \\ &\quad + a^{2q}\tilde{x}_Q^{2q_0} + \tilde{x}_Q(a^q\tilde{x}_Q^{2q_0} + A^{2q_0}\tilde{x}_Q^q + \tilde{x}_Q^{q+2q_0} + A^q\tilde{x}_Q^{2q_0q} + \tilde{x}_Q^{2q_0q+q}) \\ &\quad + a(a^q\tilde{x}_Q^{2q_0} + A^{2q_0}\tilde{x}_Q^q + \tilde{x}_Q^{q+2q_0} + A^q\tilde{x}_Q^{2q_0q} + \tilde{x}_Q^{2q_0q+q} + A^{2q_0q}\tilde{x}_Q^{q^2}) + O(\tilde{x}_Q^{q^2+1}), \end{aligned}$$

which yields,

$$\begin{aligned} \tilde{w}_Q &= A^{2q_0+1}\tilde{x}_Q^q + A^{2q_0}\tilde{x}_Q^{q+1} + A\tilde{x}_Q^{q+2q_0} + \tilde{x}_Q^{q+2q_0+1} + A^{2q_0}\tilde{x}_Q^{2q} + \tilde{x}_Q^{2q+2q_0} + A^q(A^q + A)\tilde{x}_Q^{2q_0q} \\ &\quad + A^q\tilde{x}_Q^{2q_0q+1} + A\tilde{x}_Q^{2q_0q+q} + \tilde{x}_Q^{2q_0q+q+1} + \tilde{x}_Q^{2q_0q+2q} + A^{2q_0q+1}\tilde{x}_Q^{q^2} + O(\tilde{x}_Q^{q^2+1}). \end{aligned} \quad (20)$$

Hence $v_P(\tilde{w}_Q) = v_Q(\tilde{w}_Q) = q$ as in Equation (6).

Remark 5.1. Fixing $i \geq 1$ and replacing P with $\Phi^i(P)$, one can obtain the power series expansions of $\tilde{x}_{\Phi^i(Q)}$, $\tilde{y}_{\Phi^i(Q)}$, $\tilde{z}_{\Phi^i(Q)}$ and $\tilde{w}_{\Phi^i(Q)}$. Clearly, they will be given as in Equations (18)-(20) simply replacing \tilde{x}_Q with $\tilde{x}_{\Phi^i(Q)}$ and A with A^q .

The following lemmas show that the computations carried out in Equations (18)-(20) for P (and for $\Phi^i(P)$ with $i \geq 1$ as in Remark 5.1), as well as the divisor in Equation (6) allow us to construct other functions in $L((2g(\tilde{S}) - 2)P_\infty)$ that we can use to obtain families of gaps according to Corollary 2.2.

Lemma 5.2. Let $n = 1, \dots, 2q_0 - 2$. Then there exists a function h_n on \mathcal{S} that is regular outside Q_∞ and such that

$$(h_n)_S = (n+1)q_0qQ + q_0(2q_0 - n - 1)\Phi^2(Q) + (2q_0 - n - 1)\Phi^3(Q) + \Phi^4(Q) - [(n+1)q_0 - n](q + 2q_0 + 1)Q_\infty.$$

In particular $v_P(h_n) = (n+1)q_0q$, $v_{P_\infty}(h_n) = -[(n+1)q_0 - n](q^2 + 1)$ and $h_n \in L((2g(\tilde{S}) - 2)P_\infty)$.

Proof. It is sufficient to define

$$h_n = \tilde{w}_{\Phi^2(Q)} \cdot \left(\frac{\tilde{w}_Q^{q_0}}{\tilde{w}_{\Phi(Q)}} \right)^{n+1}.$$

Indeed from Equation (6) we get

$$(h_n)_S = (n+1)q_0qQ + q_0(2q_0 - n - 1)\Phi^2(Q) + (2q_0 - n - 1)\Phi^3(Q) + \Phi^4(Q) - [(n+1)q_0 - n](q + 2q_0 + 1)Q_\infty.$$

The claim follows by observing that $2q_0 - n - 1 > 0$ since $n \leq 2q_0 - 2$. \square

Lemma 5.3. There exists a function f_1 on \mathcal{S} that is regular outside Q_∞ and such that

$$v_Q(f_1) = q_0q + q_0, \quad \text{and} \quad v_{Q_\infty}(f_1) \geq -q_0(q + 2q_0 + 1).$$

In particular $v_P(f_1) = q_0q + q_0$, $v_{P_\infty}(f_1) \geq -q_0(q^2 + 1)$ and $f_1 \in L((2g(\tilde{S}) - 2)P_\infty)$.

Proof. We have the following linear equivalence of divisors on \mathcal{S} ,

$$\begin{aligned} (q_0q + q_0)Q - q_0(q + 2q_0 + 1)Q_\infty &\cong (q_0q + q_0)Q - q_0(q + 2q_0 + 1)Q_\infty - (\tilde{w}_Q^{q_0})_S + (\tilde{w}_{\Phi(Q)})_S \\ &= q_0Q + q_0\Phi^2(Q) + \Phi^3(Q) - (q + 2q_0 + 1)Q_\infty. \end{aligned}$$

Hence if we find a function \tilde{f}_1 such that $v_Q(\tilde{f}_1) = q_0$, $v_{\Phi^2(Q)}(\tilde{f}_1) \geq q_0$, $v_{\Phi^3(Q)}(\tilde{f}_1) \geq 1$ and $v_{Q_\infty}(\tilde{f}_1) \geq -(q + 2q_0 + 1)$ then we can simply define $f_1 = \tilde{f}_1 \cdot \frac{\tilde{w}_Q^{q_0}}{\tilde{w}_{\Phi(Q)}}$, to complete the proof.

To construct \tilde{f}_1 we first observe that if $f_{\alpha,\beta,\gamma}$ denotes a linear combination $f_{\alpha,\beta,\gamma} := \alpha\tilde{w}_{\Phi(Q)} + \beta\tilde{y}_{\Phi^2(Q)} + \gamma\tilde{w}_{\Phi^2(Q)}$ with $\alpha, \beta, \gamma \in \overline{\mathbb{F}}_{q^4}$, then

$$v_{\Phi^2(Q)}(f_{\alpha,\beta,\gamma}) \geq \min\{v_{\Phi^2(Q)}(\tilde{y}_{\Phi^2(Q)}), v_{\Phi^2(Q)}(\tilde{w}_{\Phi^2(Q)}), v_{\Phi^2(Q)}(\tilde{w}_{\Phi(Q)})\} = \min\{q_0, q, 2q_0\} = q_0$$

and similarly,

$$v_{\Phi^3(Q)}(f_{\alpha,\beta,\gamma}) \geq \min\{v_{\Phi^3(Q)}(\tilde{y}_{\Phi^2(Q)}), v_{\Phi^3(Q)}(\tilde{w}_{\Phi^2(Q)}), v_{\Phi^3(Q)}(\tilde{w}_{\Phi(Q)})\} = \min\{1, 2q_0, 1\} = 1.$$

Also, from Equations (4) and (6), $f_{\alpha,\beta,\gamma}$ is regular outside Q_∞ and $v_{Q_\infty}(f_{\alpha,\beta,\gamma}) \geq -(q + 2q_0 + 1)$. Hence we give a local description of the functions $\tilde{y}_{\Phi^2(Q)}$, $\tilde{w}_{\Phi^2(Q)}$ and $\tilde{w}_{\Phi(Q)}$ at Q (equivalently P) and check whether

we can find a linear combination $f_{\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}}$ such that $v_Q(f_{\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}}) = q_0$. Doing so, it will be enough to define $\tilde{f}_1 := f_{\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}}$. From Equation (17) we have that

$$\begin{aligned}\tilde{y}_{\Phi^2(Q)} &= y + b^{q^2} + a^{q_0 q^2} (x + a^{q^2}) = (y + b) + (b^{q^2} + b) + a^{q_0 q^2} (\tilde{x}_Q + a + a^{q^2}) \\ &= (a^{q_0} \tilde{x}_Q + A \tilde{x}_Q^{q_0} + O(\tilde{x}_Q^{q_0+1})) + (b^q + b) + (b^q + b)^q + a^{q_0 q^2} (\tilde{x}_Q + a + a^{q^2}) \\ &= T(A) + (A^q + A)^{q_0} \tilde{x}_Q + A \tilde{x}_Q^{q_0} + O(\tilde{x}_Q^{q_0+1}),\end{aligned}\tag{21}$$

where using $a^{q_0} A = b^q + b$,

$$T(A) = (b^q + b) + (b^q + b)^q + a^{q_0 q^2} (A^q + A) = a^{q_0} A + a^{q_0 q} A^q + a^{q_0 q^2} (A^q + A) = A^{q_0+1} + A^{q q_0+1} + A^{q q_0+q}.$$

From Equation (5), $z = \tilde{z}_Q + a^{2q_0} (\tilde{x}_Q + a) + b^{2q_0}$ and this combined with Equation (19) gives

$$\begin{aligned}\tilde{z}_{\Phi^2(Q)} &= a^{2q_0 q^2} (\tilde{x}_Q + a) + z + b^{2q_0 q^2} \\ &= a^{2q_0 q^2} (\tilde{x}_Q + a) + a^{2q_0} (\tilde{x}_Q + a) + b^{2q_0} + b^{2q_0 q^2} + \tilde{z}_Q \\ &= (A^q + A) A^{2q_0 q} + A^{2q_0+1} + (A^q + A)^{2q_0} \tilde{x}_Q + O(\tilde{x}_Q^{2q_0}).\end{aligned}\tag{22}$$

Combining Equation (22) with Equations (6) and (20) one gets

$$\begin{aligned}\tilde{w}_{\Phi^2(Q)} &= a^{q^3} \tilde{z}_{\Phi^2(Q)} + b^{2q_0 q^2} x + w + b^{2q^2} + a^{q^2(2q_0+2)} \\ &= a^{q^3} \tilde{z}_{\Phi^2(Q)} + b^{2q_0 q^2} (\tilde{x}_Q + a) + (\tilde{w}_Q + a^q \tilde{z}_Q + b^{2q_0} (\tilde{x}_Q + a) + b^2 + a^{2q_0+2}) + b^{2q^2} + a^{q^2(2q_0+2)} \\ &= a^{q^3} ((A^q + A) A^{2q_0 q} + A^{2q_0+1} + (A^q + A)^{2q_0} \tilde{x}_Q) + b^{2q_0 q^2} (\tilde{x}_Q + a) \\ &\quad + b^{2q_0} (\tilde{x}_Q + a) + b^2 + a^{2q_0+2} + b^{2q^2} + a^{q^2(2q_0+2)} + O(\tilde{x}_Q^{2q_0}) \\ &= P(A) + T(A)^{2q_0} \tilde{x}_Q + O(\tilde{x}_Q^{2q_0}),\end{aligned}\tag{23}$$

where, $P(A) = A^{q^2+2q_0 q+q} + A^{q^2+2q_0 q+1} + A^{q^2+2q_0+1} + A^{q+2q_0+1} = A \cdot T(A)^{2q_0} + A^{q^2+2q_0 q+q}$. Finally,

$$\begin{aligned}\tilde{z}_{\Phi(Q)} &= a^{2q_0 q} (\tilde{x}_Q + a) + z + b^{2q_0 q} = a^{2q_0 q} (\tilde{x}_Q + a) + a^{2q_0} (\tilde{x}_Q + a) + b^{2q_0} + b^{2q_0 q} + \tilde{z}_Q \\ &= A^{2q_0+1} + A^{2q_0} \tilde{x}_Q + O(\tilde{x}_Q^{2q_0}),\end{aligned}$$

and hence

$$\begin{aligned}\tilde{w}_{\Phi(Q)} &= a^{q^2} \tilde{z}_{\Phi(Q)} + b^{2q_0 q} x + w + b^{2q} + a^{q(2q_0+2)} \\ &= a^{q^2} A^{2q_0+1} + a^{q^2} A^{2q_0} \tilde{x}_Q + b^{2q_0 q} (\tilde{x}_Q + a) + (b^{2q_0} (\tilde{x}_Q + a) + b^2 + a^{2q_0+2}) + b^{2q} + a^{q(2q_0+2)} + O(\tilde{x}_Q^{2q_0}) \\ &= A^{q+2q_0+1} + A^{q+2q_0} \tilde{x}_Q + O(\tilde{x}_Q^{2q_0}).\end{aligned}\tag{24}$$

Equations (21), (23) and (24) show that finding $\tilde{\alpha}$, $\tilde{\beta}$ and $\tilde{\gamma}$ such that $v_Q(f_{\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}}) = q_0$ is equivalent to find a solution to the following system of linear equations (corresponding to the coefficients of \tilde{x}_Q^0 , \tilde{x}_Q and $\tilde{x}_Q^{q_0}$ in the power series expansion of $f_{\alpha, \beta, \gamma}$ respectively),

$$\begin{cases} \alpha A^{q+2q_0+1} + \beta T(A) + \gamma [A \cdot T(A)^{2q_0} + A^{q^2+2q_0 q+q}] = 0, \\ \alpha A^{q+2q_0} + \beta (A^q + A)^{q_0} + \gamma T(A)^{2q_0} = 0, \\ \beta A \neq 0. \end{cases}\tag{25}$$

Since $A \neq 0$ and $(A^q + A)^{q_0} = (T(A) + A^{q_0q+q})/A$, System (25) can be rewritten as

$$\begin{cases} \beta = A^{q^2+q_0q}\gamma, \\ \alpha = \frac{\gamma(A^{q^2+q_0q} \cdot T(A) + P(A))}{A^{q+2q_0+1}}, \\ \beta \neq 0. \end{cases} \quad (26)$$

To conclude the proof it is now sufficient to define $\tilde{f}_1 = f_{\frac{A^{q^2+q_0q} \cdot T(A) + P(A)}{A^{q+2q_0+1}}, A^{q^2+q_0q}, 1}$. \square

Lemma 5.4. *There exists a function f_2 on \mathcal{S} that is regular outside Q_∞ and such that*

$$v_Q(f_2) = 2q_0q + 2q_0 + 1, \quad \text{and} \quad v_{Q_\infty}(f_2) \geq -2q_0(q + 2q_0 + 1).$$

In particular $v_P(f_2) = 2q_0q + 2q_0 + 1$, $v_{P_\infty}(f_2) \geq -2q_0(q^2 + 1)$ and $f_2 \in L((2g(\tilde{\mathcal{S}}) - 2)P_\infty)$.

Proof. We have the following linear equivalence of divisors on \mathcal{S} ,

$$\begin{aligned} (2q_0q + 2q_0 + 1)Q - 2q_0(q + 2q_0 + 1)Q_\infty &\cong (2q_0q + 2q_0 + 1)Q - 2q_0(q + 2q_0 + 1)Q_\infty - (\tilde{w}_Q^{2q_0})_{\mathcal{S}} + (\tilde{w}_{\Phi(Q)}^2)_{\mathcal{S}} \\ &= (2q_0 + 1)Q + 2q_0\Phi^2(Q) + 2\Phi^3(Q) - 2(q + 2q_0 + 1)Q_\infty. \end{aligned}$$

Hence if we find a function \tilde{f}_2 such that $v_Q(\tilde{f}_2) = 2q_0 + 1$, $v_{\Phi^2(Q)}(\tilde{f}_2) \geq 2q_0$, $v_{\Phi^3(Q)}(\tilde{f}_2) \geq 2$ and $v_{Q_\infty}(\tilde{f}_2) \geq -2(q + 2q_0 + 1)$ then we can simply define $f_2 = \tilde{f}_2 \cdot \frac{\tilde{w}_Q^{2q_0}}{\tilde{w}_{\Phi(Q)}^2}$, to complete the proof.

To construct \tilde{f}_2 , we first observe that

$$(\tilde{w}_{\Phi^2(Q)} \cdot \tilde{z}_Q)_{\mathcal{S}} = 2q_0Q + q\Phi^2(Q) + 2q_0\Phi^3(Q) + (E + \Phi^4(Q)) - [2(q + 2q_0 + 1) - 1]Q_\infty,$$

where E is effective and not containing Q , $\Phi^2(Q)$ and $\Phi^3(Q)$. Also, from the proof of Lemma 5.3 we have a function $\tilde{f}_1 = \frac{A^{q^2+q_0q} \cdot T(A) + P(A)}{A^{q+2q_0+1}} \tilde{w}_{\Phi(Q)} + A^{q^2+q_0q} \tilde{y}_{\Phi^2(Q)} + \tilde{w}_{\Phi^2(Q)}$, such that $(\tilde{f}_1^2) = 2q_0Q + 2q_0\Phi^2(Q) + 2\Phi^3(Q) + E_1 - 2(q + 2q_0 + 1)Q_\infty$, where E_1 is effective and not containing Q .

Hence if $h_{\alpha,\beta}$ denotes a linear combination $h_{\alpha,\beta} := \alpha \tilde{w}_{\Phi^2(Q)} \cdot \tilde{z}_Q + \beta \tilde{f}_1^2$ with $\alpha, \beta \in \overline{\mathbb{F}}_q$, then

$$v_{\Phi^2(Q)}(h_{\alpha,\beta}) \geq \min\{v_{\Phi^2(Q)}(\tilde{w}_{\Phi^2(Q)} \cdot \tilde{z}_Q), v_{\Phi^2(Q)}(\tilde{f}_1^2)\} = \min\{q, 2q_0\} = 2q_0$$

and similarly,

$$v_{\Phi^3(Q)}(h_{\alpha,\beta}) \geq \min\{v_{\Phi^3(Q)}(\tilde{w}_{\Phi^2(Q)} \cdot \tilde{z}_Q), v_{\Phi^3(Q)}(\tilde{f}_1^2)\} = \min\{2q_0, 2\} = 2.$$

Also, from Equations (4)-(6), $h_{\alpha,\beta}$ is regular outside Q_∞ and $v_{Q_\infty}(h_{\alpha,\beta}) \geq -2(q + 2q_0 + 1)$. Hence using the local description of the functions $\tilde{w}_{\Phi^2(Q)}$, \tilde{z}_Q and \tilde{f}_1 , we find a linear combination $h_{\tilde{\alpha},\tilde{\beta}}$ such that $v_Q(h_{\tilde{\alpha},\tilde{\beta}}) = 2q_0 + 1$ and doing so, it will be enough to define $\tilde{f}_2 := h_{\tilde{\alpha},\tilde{\beta}}$ to complete the proof.

Systems (25) and (26) yield, $\tilde{f}_1^2 = A^{2(q^2+q_0q+1)} \tilde{x}_Q^{2q_0} + O(\tilde{x}_Q^{2q_0+2})$, while from Equations (19) and (23) one has,

$$\begin{aligned} \tilde{w}_{\Phi^2(Q)} \cdot \tilde{z}_Q &= (P(A) + T(A)^{2q_0} \tilde{x}_Q + O(\tilde{x}_Q^{2q_0})) \cdot (A \tilde{x}_Q^{2q_0} + \tilde{x}_Q^{2q_0+1} + O(\tilde{x}_Q^q)) \\ &= A \cdot P(A) \tilde{x}_Q^{2q_0} + [P(A) + A \cdot T(A)^{2q_0}] \tilde{x}_Q^{2q_0+1} + O(\tilde{x}_Q^{2q_0+2}). \end{aligned}$$

So it is enough to define

$$\begin{aligned}\tilde{f}_2 &:= h_{1, AP(A)/A^{2(q^2+q_0q+1)}} = A \cdot P(A) \tilde{x}_Q^{2q_0} + [P(A) + A \cdot T(A)^{2q_0}] \tilde{x}_Q^{2q_0+1} + A \cdot P(A) \tilde{x}_Q^{2q_0} + O(\tilde{x}_Q^{2q_0+2}) \\ &= [P(A) + A \cdot T(A)^{2q_0}] \tilde{x}_Q^{2q_0+1} + O(\tilde{x}_Q^{2q_0+2}).\end{aligned}$$

Recalling that $P(A) + AT(A)^{2q_0} = A^{q^2+2q_0q+q} \neq 0$, we see that $v_Q(\tilde{f}_2) = 2q_0 + 1$ as desired. \square

Lemma 5.5. *For all $i = 0, \dots, q_0 - 2$ there exists a function g_i on \mathcal{S} that is regular outside Q_∞ and such that*

$$v_Q(g_i) = (2i+1)q_0q + i + 1, \quad \text{and} \quad v_{Q_\infty}(g_i) \geq -((2i+1)q_0 - i)(q + 2q_0 + 1).$$

In particular $v_P(g_i) = (2i+1)q_0q + i + 1$, $v_{P_\infty}(g_i) \geq -((2i+1)q_0 - i)(q^2 + 1)$ and $g_i \in L((2g(\tilde{\mathcal{S}}) - 2)P_\infty)$ for all $i = 0, \dots, q_0 - 2$.

Proof. Let $D_i := ((2i+1)q_0q + i + 1)Q - ((2i+1)q_0 - i)(q + 2q_0 + 1)Q_\infty$. Then,

$$\begin{aligned}D_i &\cong ((2i+1)q_0q + i + 1)Q - ((2i+1)q_0 - i)(q + 2q_0 + 1)Q_\infty - (\tilde{w}_Q^{(2i+1)q_0})_{\mathcal{S}} + (\tilde{w}_{\Phi(Q)}^{2i+1})_{\mathcal{S}} \\ &= (i+1)Q + (2i+1)q_0\Phi^2(Q) + (2i+1)\Phi^3(Q) - (i+1)(q + 2q_0 + 1)Q_\infty.\end{aligned}$$

Hence if we find a function \tilde{g}_i such that $v_Q(\tilde{g}_i) = i + 1$, $v_{\Phi^2(Q)}(\tilde{g}_i) \geq (2i+1)q_0$, $v_{\Phi^3(Q)}(\tilde{g}_i) \geq 2i+1$ and $v_{Q_\infty}(\tilde{g}_i) \geq ((2i+1)q_0 - i)(q + 2q_0 + 1)$ then we simply define $g_i = \tilde{g}_i \cdot \left(\frac{\tilde{w}_Q^{q_0}}{\tilde{w}_{\Phi(Q)}}\right)^{2i+1}$ to complete the proof.

To this end we first compute some local power series expansions at $\Phi^3(P)$ (equivalently $\Phi^3(Q)$) using the local parameter $\tilde{x}_{\Phi^3(Q)} = x + a^{q^3} = \tilde{x}_Q + a + a^{q^3} = \tilde{x}_Q + A + A^q + A^{q^2}$. Clearly, we can use the expressions we already obtained in Equations (17)-(20) to describe the local power series expansion of $y + b^{q^3}$, $\tilde{y}_{\Phi^3(Q)}$, $\tilde{z}_{\Phi^3(Q)}$, $\tilde{w}_{\Phi^3(Q)}$ at $\Phi^3(P)$ simply replacing a with a^{q^3} and hence A with A^{q^3} . From Equation (17) and $a^{q_0}A = b^q + b$, one has $y + b^{q^2} = (y + b^{q^3}) + (b^{q^3} + b^{q^2}) = a^{q_0q^2}A^{q^2} + a^{q_0q^3}\tilde{x}_{\Phi^3(Q)} + A^{q^3}\tilde{x}_{\Phi^3(Q)}^{q_0} + \tilde{x}_{\Phi^3(Q)}^{q_0+1} + O(\tilde{x}_{\Phi^3(Q)}^q)$. Hence

$$\begin{aligned}\tilde{y}_{\Phi^2(Q)} &= (y + b^{q^2}) + a^{q_0q^2}(x + a^{q^2}) = (y + b^{q^2}) + a^{q_0q^2}(\tilde{x}_{\Phi^3(Q)} + A^{q^2}) \\ &= A^{q_0q^2}\tilde{x}_{\Phi^3(Q)} + A^{q^3}\tilde{x}_{\Phi^3(Q)}^{q_0} + \tilde{x}_{\Phi^3(Q)}^{q_0+1} + O(\tilde{x}_{\Phi^3(Q)}^q).\end{aligned}\tag{27}$$

Similarly using Equation (19) and $z = \tilde{z}_{\Phi^3(Q)} + a^{2q_0q^3}x + b^{2q_0q^3}$ one gets,

$$\begin{aligned}\tilde{z}_{\Phi^2(Q)} &= a^{2q_0q^2}(\tilde{x}_{\Phi^3(Q)} + a^{q^3}) + z + b^{2q_0q^2} \\ &= a^{2q_0q^2}(\tilde{x}_{\Phi^3(Q)} + a^{q^3}) + \tilde{z}_{\Phi^3(Q)} + a^{2q_0q^3}x + b^{2q_0q^3} + b^{2q_0q^2} \\ &= A^{2q_0q^2}\tilde{x}_{\Phi^3(Q)} + \tilde{z}_{\Phi^3(Q)} = A^{2q_0q^2}\tilde{x}_{\Phi^3(Q)} + O(\tilde{x}_{\Phi^3(Q)}^{2q_0}).\end{aligned}\tag{28}$$

We define $H_1 := \tilde{z}_{\Phi^2(Q)} + A^{q_0q^2}\tilde{y}_{\Phi^2(Q)}$. Then using Equations (27) and (28),

$$\begin{aligned}H_1 &= A^{q_0q^2}(A^{q_0q^2}\tilde{x}_{\Phi^3(Q)} + A^{q^3}\tilde{x}_{\Phi^3(Q)}^{q_0} + \tilde{x}_{\Phi^3(Q)}^{q_0+1}) + A^{2q_0q^2}\tilde{x}_{\Phi^3(Q)} + O(\tilde{x}_{\Phi^3(Q)}^{2q_0}) \\ &= A^{q^3+q_0q^2}\tilde{x}_{\Phi^3(Q)}^{q_0} + A^{q_0q^2}\tilde{x}_{\Phi^3(Q)}^{q_0+1} + O(\tilde{x}_{\Phi^3(Q)}^{2q_0}).\end{aligned}$$

Since $A \neq 0$ this shows that $v_{\Phi^3(Q)}(H_1) = q_0$. Also note that

$$v_{\Phi^2(Q)}(H_1) = \min\{v_{\Phi^2(Q)}(\tilde{z}_{\Phi^2(Q)}), v_{\Phi^2(Q)}(\tilde{y}_{\Phi^2(Q)})\} = \min\{2q_0, q_0\} = q_0.$$

To compute the valuation of H_1 at P we see that from Equations (21) and (22),

$$H_1 = \tilde{z}_{\Phi^2(Q)} + A^{q_0 q^2} \tilde{y}_{\Phi^2(Q)} = P_1(A) + [(A^q + A)^2 + A^{q^2}(A^q + A)]^{q_0} \tilde{x}_Q + O(\tilde{x}_Q^{q_0}),$$

where $P_1(A) = (A^q + A)A^{2q_0 q} + A^{2q_0+1} + A^{q_0 q^2} T(A)$. Note that the coefficient of \tilde{x}_Q in H_1 is not zero as both $A^q + A = 0$ and $A^{q^2} + A^q + A = 0$ imply $P \in \tilde{\mathcal{S}}(\mathbb{F}_{q^4})$.

If $P_1(A) = 0$ then $v_Q(H_1) = 1$ and we define $N_1 := H_1$. Otherwise recalling that from Equation (23) we have $\tilde{w}_{\Phi^2(Q)} = P(A) + T(A)^{2q_0} \tilde{x}_Q + O(\tilde{x}_Q^{2q_0})$, we define $N_1 := P_1(A) \tilde{w}_{\Phi^2(Q)} + P(A) H_1$. Doing so,

$$\begin{aligned} N_1 &= P_1(A)(P(A) + T(A)^{2q_0} \tilde{x}_Q + O(\tilde{x}_Q^{2q_0})) + P(A)(P_1(A) + [(A^q + A)^2 + A^{q^2}(A^q + A)]^{q_0} \tilde{x}_Q + O(\tilde{x}_Q^{q_0})) \\ &= P_2(A) \tilde{x}_Q + O(\tilde{x}_Q^{q_0}). \end{aligned}$$

Using $P(A) = AT(A)^{2q_0} + A^{q^2+2q_0 q+q}$, $(A^q + A)^{q_0} = (T(A) + A^{q_0 q+q})/A$ and $P \notin \tilde{\mathcal{S}}(\mathbb{F}_{q^4})$, one obtains $P_2(A) = P_1(A)T(A)^{2q_0} + P(A)[(A^q + A)^2 + A^{q^2}(A^q + A)]^{q_0} = A^{q_0 q+q+q_0} P(A)^{q_0} \neq 0$.

Since N_1 is a linear combination of $\tilde{w}_{\Phi^2(Q)}$ and H_1 , both the valuations of N_1 at $\Phi^2(Q)$ and $\Phi^3(Q)$ are at least q_0 . This shows that, in both cases $P_1(A) = 0$ and $P_1(A) \neq 0$, one has

$$(N_1)_{\mathcal{S}} = Q + q_0 \Phi^2(Q) + q_0 \Phi^3(Q) + E_{N_1} - (q + 2q_0 + 1)Q_{\infty}, \quad (29)$$

where E_{N_1} is effective and with support not containing Q . Finally, we show that there exists a function N_0 on \mathcal{S} such that

$$(N_0)_{\mathcal{S}} = Q + 2q_0 \Phi^2(Q) + \Phi^3(Q) + E_{N_0} - (q + 2q_0 + 1)Q_{\infty}, \quad (30)$$

where E_{N_0} is effective and with support not containing Q . Indeed since we already obtained in Equations (22) and (23) that $\tilde{z}_{\Phi^2(Q)} = (A^q + A)A^{2q_0 q} + A^{2q_0+1} + (A^q + A)^{2q_0} \tilde{x}_Q + O(\tilde{x}_Q^{2q_0})$, and $\tilde{w}_{\Phi^2(Q)} = P(A) + T(A)^{2q_0} \tilde{x}_Q + O(\tilde{x}_Q^{2q_0})$, we can define $N_0 := P(A) \tilde{z}_{\Phi^2(Q)} + [(A^q + A)A^{2q_0 q} + A^{2q_0+1}] \tilde{w}_{\Phi^2(Q)}$. Doing so,

$$N_0 = (P(A)(A^q + A)^{2q_0} + [(A^q + A)A^{2q_0 q} + A^{2q_0+1}]T(A)^{2q_0}) \tilde{x}_Q + O(\tilde{x}_Q^{2q_0}),$$

where by direct checking $P(A)(A^q + A)^{2q_0} + [(A^q + A)A^{2q_0 q} + A^{2q_0+1}]T(A)^{2q_0} = A^{2q_0 q+2q+2q_0} \neq 0$, and hence $v_Q(N_0) = 1$. Also, $v_{\Phi^2(Q)}(N_0) = \min\{v_{\Phi^2(Q)}(\tilde{z}_{\Phi^2(Q)}), v_{\Phi^2(Q)}(\tilde{w}_{\Phi^2(Q)})\} = \min\{2q_0, q\} = 2q_0$, and $v_{\Phi^3(Q)}(N_0) = \min\{v_{\Phi^3(Q)}(\tilde{z}_{\Phi^2(Q)}), v_{\Phi^3(Q)}(\tilde{w}_{\Phi^2(Q)})\} = \min\{1, 2q_0\} = 1$, as desired. Using Equations (29) and (30) it is now enough to define $\tilde{f}_i = N_0^i \cdot N_1$. Indeed one has,

$$(\tilde{f}_i)_{\mathcal{S}} = (i+1)Q + (2i+1)q_0 \Phi^2(Q) + (q_0 + i) \Phi^3(Q) + (iE_{N_0} + E_{N_1}) - (i+1)(q + 2q_0 + 1)Q_{\infty},$$

where $q_0 + i > 2i + 1$ as $i \leq q_0 - 2$. □

To construct gaps using Corollary 2.2 we will look at functions of the form

$$\tilde{x}_Q^{a_1} \cdot \tilde{y}_Q^{a_2} \cdot \tilde{z}_Q^{a_3} \cdot \tilde{w}_Q^{a_4} \cdot \prod_{n=1}^{2q_0-2} h_n^{b_n} \cdot f_1^c \cdot f_2^d \cdot \prod_{n=0}^{q_0-2} g_n^{e_n} \cdot \pi_P^f \quad (31)$$

for suitable choices of exponents $a_1, a_2, a_3, a_4, b_1, \dots, b_{2q_0-2}, c, d, e_0, \dots, e_{q_0-2}, f$.

The six families of natural numbers $F_1, F_2, F_3, F_4, F_5, F_6$ defined in Theorem 1.1 correspond to sets of valuations of functions as in Equation (31). Indeed our aim is to show that $G(P) = F := \cup_{i=1}^6 F_i$. To this end we proceed with the following two steps: first, we prove that F contains exactly $g(\tilde{\mathcal{S}})$ elements. Then we prove that the functions as in Equation (31) whose valuations are contained in F , are in $L((2g(\tilde{\mathcal{S}}) - 2)P_{\infty})$.

Proposition 5.6. *The set F consists of $g(\tilde{\mathcal{S}})$ distinct natural numbers.*

Proof. If $s = 1, 2$ the claim can be checked directly using a computer. Table 1 collects the cardinalities of the families F'_i s in these two cases.

Table 1: The cases $s = 1$ and $s = 2$.

s	$ F_1 $	$ F_2 $	$ F_3 $	$ F_4 $	$ F_5 $	$ F_6 $	$ F $	$g(\tilde{\mathcal{S}})$
1	146	31	8	0	9	2	196	196
2	12584	2393	192	96	87	24	15376	15376

So in the following, we assume $s > 2$. We first show that for all $i = 1, \dots, 6$ different choices of coefficients within the same family F_i give rise to distinct natural numbers.

- For $i = 1$, assume by contradiction that there exists element $v \in F_1$ having two different expressions, namely

$$a_1 + a_2q_0 + 2a_3q_0 + a_4q + fq^2 + 1 = a'_1 + a'_2q_0 + 2a'_3q_0 + a'_4q + f'q^2 + 1. \quad (32)$$

Considering Equation (32) modulo q_0 we obtain $a_1 \equiv a'_1 \pmod{q_0}$ and therefore $a_1 = a'_1$, since $0 \leq a_1, a'_1 \leq q_0 - 1$. So, Equation (32) can be simplified as $a_2 + 2a_3 + 2a_4q_0 + 2fq_0 = a'_2 + 2a'_3 + 2a'_4q_0 + 2f'q_0$. Repeating the same procedure, firstly considering the equality modulo 2, then modulo q_0 again and finally modulo q , we obtain $a_2 = a'_2, a_3 = a'_3, a_4 = a'_4$ respectively. As a consequence, $f = f'$. This yields a contradiction.

- As before, for $i = 2$ let

$$a_1 + a_2q_0 + 2a_3q_0 + a_4q + fq^2 + (n+1)qq_0 + 1 = a'_1 + a'_2q_0 + 2a'_3q_0 + a'_4q + f'q^2 + (n'+1)qq_0 + 1.$$

With a similar argument used for $i = 1$, we can reduce the above equality modulo q_0 , then modulo 2, then modulo q_0 again and finally modulo q to obtain $a_1 = a'_1, a_2 = a'_2, a_3 = a'_3$ and $a_4 + nq_0 = a'_4 + n'q_0$ respectively. As a consequence $f = f'$. The conditions $a_1 + a_2 + a_3 + a_4 + f = q - q_0 - 2 - nq_0 + n$ and $a'_1 + a'_2 + a'_3 + a'_4 + f' = q - q_0 - 2 - n'q_0 + n'$ now imply that $n = n'$ and $a_4 = a'_4$.

- The argument used for $i = 3, \dots, 6$ is exactly the same as for $i = 1, 2$ and hence it is omitted.

Next, one needs to prove that $F_i \cap F_j = \emptyset$, for all $i, j = 1, \dots, 6, i \neq j$. To this end, one can use exactly the same method used in the first part of this proof. For this reason just the first two cases are proven here in full details. Let $\sigma = a_1 + a_2 + a_3 + a_4 + f$.

- $F_1 \cap F_2 = \emptyset$: suppose by contradiction that there exists $v \in F_1 \cap F_2$. Then:

$$\begin{aligned} v &= a_1 + a_2q_0 + 2a_3q_0 + a_4q + (\sigma - a_1 - a_2 - a_3 - a_4)q^2 + 1 \\ &= a'_1 + a'_2q_0 + 2a'_3q_0 + a'_4q + (n+1)q_0q + (q - q_0 - 2 - nq_0 + n - a'_1 - a'_2 - a'_3 - a'_4)q^2 + 1. \end{aligned}$$

Considering the equality above modulo q_0 , then modulo 2 and finally modulo q_0 we get $a_1 = a'_1, a_2 = a'_2, a_3 = a'_3$. This yields

$$a_4 + (\sigma - a_4)q = a'_4 + (n+1)q_0 + (q - q_0 - 2 - nq_0 + n - a'_4)q.$$

Since $a_4 \leq q - 2$ and $a'_4 + (n+1)q_0 \leq q - q_0 - 1 - nq_0 + nq_0 + q_0 = q - 1$, the equality above modulo q gives $a_4 = a'_4 + (n+1)q_0$ and $\sigma = q - 2 + n$, a contradiction to $\sigma \leq q - 2$ as $n \geq 1$.

- $F_1 \cap F_3 = \emptyset$: suppose by contradiction that there exists $v \in F_1 \cap F_3$. Then we can write

$$\begin{aligned} v &= a_1 + a_2q_0 + 2a_3q_0 + a_4q + (\sigma - a_1 - a_2 - a_3 - a_4)q^2 + 1 \\ &= a'_1 + a'_2q_0 + 2a'_3q_0 + a'_4q + (2n+1)q_0q + n+1 + (q - q_0 - 2 - 2nq_0 + n - a'_1 - a'_2 - a'_3 - a'_4)q^2 + 1. \end{aligned}$$

Arguing as in the previous case one gets $a_1 = a'_1 + n + 1$, $a_2 = a'_2$ and $a_3 = a'_3$, so that

$$a_4 + (\sigma - n - a_4)q = a'_4 + (2n+1)q_0 + (q - q_0 - 1 - 2nq_0 + n - a'_4)q.$$

Note that $a'_4 + (2n+1)q_0 \leq q_0 - 1 + (2q_0 - 3)q_0 < q$ so that considering the equality modulo q one gets $a_4 = a'_4 + (2n+1)q_0$. Hence $\sigma = q - 1 + 2n \geq q - 1$, a contradiction.

Finally, we need to prove that F consists of $g(\tilde{\mathcal{S}})$ distinct elements. From the previous step it is enough to check that there are exactly $g(\tilde{\mathcal{S}})$ possible distinct choices of such coefficients.

For a given $\sigma \in \mathbb{N}$, we denote the number of combinations of five natural numbers a_1, a_2, a_3, a_4, f such that $a_1 + a_2 + a_3 + a_4 + f = \sigma$ with $\mathcal{B}(\sigma) = \binom{\sigma+4}{4}$. Observe that from [9, Equation 5.10] for any $n \in \mathbb{N}$ it holds:

$$\sum_{\sigma=0}^n \mathcal{B}(\sigma) = \sum_{\sigma'=4}^{n+4} \binom{\sigma'}{4} = \binom{n+5}{5} = \frac{n(n+1)}{120}(n^3 + 14n^2 + 71n + 154) + 1.$$

It is also useful to denote the number of combinations of four natural numbers a_2, a_3, a_4, f such that $a_2 + a_3 + a_4 + f = \sigma$ with $\mathcal{B}'(\sigma) = \binom{\sigma+3}{3}$ and, similarly, the number of combinations of three natural number a_3, a_4, f such that $a_3 + a_4 + f = \sigma$ with $\mathcal{B}''(\sigma) = \binom{\sigma+2}{2}$.

- Let us count the number of elements in F_1 . By the inclusion-exclusion principle, this is given by:

$$\begin{aligned} &\sum_{\sigma=0}^{q-2} \mathcal{B}(\sigma) - \left(\sum_{\sigma_1=0}^{q-2-q_0} \mathcal{B}(\sigma_1) + \sum_{\sigma_2=0}^{q-2-2} \mathcal{B}(\sigma_2) + \sum_{\sigma_3=0}^{q-2-q_0} \mathcal{B}(\sigma_3) \right) \\ &+ \left(\sum_{\sigma_{1,2}=0}^{q-2-q_0-2} \mathcal{B}(\sigma_{1,2}) + \sum_{\sigma_{2,3}=0}^{q-2-2-q_0} \mathcal{B}(\sigma_{2,3}) + \sum_{\sigma_{1,3}=0}^{q-2-q_0-q_0} \mathcal{B}(\sigma_{1,3}) \right) - \sum_{\sigma_{1,2,3}=0}^{q-2-q_0-2-q_0} \mathcal{B}(\sigma_{1,2,3}) \\ &= \frac{q^3}{2} - q^2q_0 + \frac{7}{24}q^2 - \frac{1}{12}q. \end{aligned}$$

- Let us now count the number of elements in F_2 . In this case $\sigma = q - q_0 - 2 - 2nq_0 + n$ in F_2 . Observe that: $\sigma - q_0, \sigma - 2$ are always non-negative, while $\sigma - (q - q_0 - nq_0)$ is non-negative only for $n > 1$; $\sigma - 2q_0, \sigma - 2 - q_0$ are non-negative as $s > 1$; $\sigma - (q - q_0 - nq_0) - 2$ and $\sigma - (q - q_0 - nq_0) - q_0$ are non-negative only for $n > 3$ and $n > q_0 + 1$, respectively. Both conditions can hold true as $s > 1$; $\sigma - 2q_0 - 2$ is non-negative since $s > 1$; $\sigma - (q - q_0 - nq_0) - 2 - q_0$ is non-negative only for $n > q_0 + 3$, which can hold true as $s > 2$; $\sigma - (q - q_0 - nq_0) - 2q_0$ is always negative; and $\sigma - (q - q_0 - nq_0) - 2 - 2q_0$

is always negative. Hence, the number of elements in F_2 is given by:

$$\begin{aligned}
& \sum_{n=1}^{2q_0-2} \mathcal{B}(\sigma) - \left(\sum_{n=1}^{2q_0-2} \mathcal{B}(\sigma-2) + \sum_{n=1}^{2q_0-2} 2\mathcal{B}(\sigma-q_0) + \sum_{n=2}^{2q_0-2} \mathcal{B}(\sigma-(q-q_0-nq_0)) \right) + \left(\sum_{n=1}^{2q_0-2} \mathcal{B}(\sigma-2q_0) \right. \\
& + \sum_{n=1}^{2q_0-2} 2\mathcal{B}(\sigma-2-q_0) + \sum_{n=4}^{2q_0-2} \mathcal{B}(\sigma-(q-q_0-nq_0)-2) + \sum_{n=q_0+2}^{2q_0-2} 2\mathcal{B}(\sigma-(q-q_0-nq_0)-q_0) \Big) \\
& - \left(\sum_{n=1}^{2q_0-2} \mathcal{B}(\sigma-2-2q_0) + \sum_{n=q_0+4}^{2q_0-2} 2\mathcal{B}(\sigma-(q-q_0-nq_0)-2-q_0) \right) \\
& = q^2 q_0 - \frac{43}{24} q^2 + qq_0 + \frac{1}{12} q + 1.
\end{aligned}$$

- For counting the elements of F_3 , observe that $\sigma - q_0, \sigma - 2, \sigma - (q_0 - 1 - n), \sigma - 2 - q_0, \sigma - 2q_0, \sigma - (q_0 - 1 - n) - 2, \sigma - (q_0 - 1 - n) - q_0$ are always non-negative, and also $\sigma - 2 - 2q_0, \sigma - 2 - q_0 - (q_0 - 1 - n), \sigma - 2q_0 - (q_0 - 1 - n), \sigma - 2 - 2q_0 - (q_0 - 1 - n)$ are non-negative as $s > 1$.

Taking into consideration the aforementioned observations we get that the cardinality of F_3 is equal to

$$\begin{aligned}
& \sum_{n=0}^{q_0-2} \mathcal{B}(\sigma) - \sum_{n=0}^{q_0-2} (\mathcal{B}(\sigma-2) + 2\mathcal{B}(\sigma-q_0) + \mathcal{B}(\sigma-(q_0-1-n))) \\
& + \sum_{n=0}^{q_0-2} (\mathcal{B}(\sigma-2q_0) + 2\mathcal{B}(\sigma-2-q_0) + \mathcal{B}(\sigma-(q_0-1-n)-2) + 2\mathcal{B}(\sigma-(q_0-1-n)-q_0)) \\
& - \sum_{n=0}^{q_0-2} (\mathcal{B}(\sigma-2-2q_0) + 2\mathcal{B}(\sigma-2-q_0-(q_0-1-n)) + \mathcal{B}(\sigma-2q_0-(q_0-1-n))) \\
& + \sum_{n=0}^{q_0-2} \mathcal{B}(\sigma-2-2q_0-(q_0-1-n)) = \frac{1}{4} q^2 - \frac{1}{2} qq_0.
\end{aligned}$$

- The number of elements of F_4 is given by:

$$\begin{aligned}
& \sum_{n=0}^{q_0-3} \mathcal{B}(\sigma) - \sum_{n=0}^{q_0-3} (\mathcal{B}(\sigma-2) + 2\mathcal{B}(\sigma-q_0) + \mathcal{B}(\sigma-(q_0-2-n))) \\
& + \sum_{n=0}^{q_0-3} (\mathcal{B}(\sigma-2q_0) + 2\mathcal{B}(\sigma-2-q_0) + \mathcal{B}(\sigma-(q_0-2-n)-2) + 2\mathcal{B}(\sigma-(q_0-2-n)-q_0)) \\
& - \sum_{n=0}^{q_0-3} (\mathcal{B}(\sigma-2-2q_0) + 2\mathcal{B}(\sigma-2-q_0-(q_0-2-n)) + \mathcal{B}(\sigma-2q_0-(q_0-2-n))) \\
& + \sum_{n=0}^{q_0-3} \mathcal{B}(\sigma-2-2q_0-(q_0-2-n)) = \frac{1}{4} q^2 - \frac{3}{2} qq_0 + q.
\end{aligned}$$

- For studying the number of elements of F_5 , let us consider the cases $c = 0$ and $c = 1$ separately, starting from the case $c = 0$.

The number of elements in F_5 for $c = 0$ can be computed as follows

$$\begin{aligned} & \sum_{d=1}^{q_0-1} \mathcal{B}'(\sigma) - \sum_{d=1}^{q_0-1} (\mathcal{B}'(\sigma-2) + \mathcal{B}'(\sigma-q_0) + \mathcal{B}'(\sigma-(q_0-d))) \\ & + \sum_{d=1}^{q_0-1} (\mathcal{B}'(\sigma-2-q_0) + \mathcal{B}'(\sigma-2-(q_0-d)) + \mathcal{B}'(\sigma-q_0-(q_0-d))) \\ & - \sum_{d=1}^{q_0-1} \mathcal{B}'(\sigma-2-q_0-(q_0-d)) = \frac{1}{2}qq_0 - \frac{1}{2}q. \end{aligned}$$

Let us consider now the case $c = 1$. Observe that $\sigma - q_0$ and $\sigma - q_0 - (q_0 - d)$ are negative only for $d = q_0 - 1$, while the quantity $\sigma - (q_0 - d)$ is always non-negative as $s > 2$. Hence the number of elements in F_5 with $c = 1$ is given by

$$\sum_{d=0}^{q_0-1} \mathcal{B}''(\sigma) - \left(\sum_{d=0}^{q_0-1} \mathcal{B}''(\sigma - (q_0 - d)) + \sum_{d=0}^{q_0-2} \mathcal{B}''(\sigma - q_0) \right) + \sum_{d=0}^{q_0-2} \mathcal{B}''(\sigma - q_0 - (q_0 - d)) = \frac{1}{4}qq_0 + \frac{1}{4}q - 1.$$

- For counting the number of elements of F_6 , observe that $\sigma - q_0 - (n + 1)$ is non-negative as $s > 1$. So, $|F_6|$ coincides with

$$\sum_{n=0}^{q_0-2} \mathcal{B}''(\sigma) - \sum_{n=0}^{q_0-2} (\mathcal{B}''(\sigma - q_0) + \mathcal{B}''(\sigma - (n + 1))) + \sum_{n=0}^{q_0-2} \mathcal{B}''(\sigma - q_0 - (n + 1)) = \frac{1}{4}qq_0 - \frac{1}{4}q.$$

The conclusion follows after adding up the quantities obtained for each of the six families separately. For all $s > 2$ the computation yields $|F| = \frac{1}{2}q(q-1)^2 = g(\tilde{\mathcal{S}})$. \square

A consequence of Proposition 5.6 is that we can assign to each $v \in F$ a unique function \mathfrak{f}_v on $\tilde{\mathcal{S}}$ of the form as in Equation (31) that is regular outside P_∞ and has valuation $v - 1$ at P in the following way: if v is expressed as

$$v = a_1 + a_2q_0 + 2a_3q_0 + a_4q + qq_0 \sum_{n=1}^{2q_0-2} b_n(n+1) + cq_0(q+1) + d(2q_0+1) + \sum_{n=0}^{q_0-2} e_n((2n+1)q_0q + n+1) + fq^2 + 1,$$

then we define

$$\mathfrak{f}_v := \tilde{x}_Q^{a_1} \cdot \tilde{y}_Q^{a_2} \cdot \tilde{z}_Q^{a_3} \cdot \tilde{w}_Q^{a_4} \cdot \prod_{n=1}^{2q_0-2} h_n^{b_n} \cdot f_1^c \cdot f_2^d \cdot \prod_{n=0}^{q_0-2} g_n^{e_n} \cdot \pi_P^f.$$

Proposition 5.6 guarantees that there are exactly $g(\tilde{\mathcal{S}})$ such functions having pairwise distinct valuations at P . The following proposition, together with Proposition 5.6 and Corollary 2.2, completes the proof of Theorem 1.1.

Proposition 5.7. \mathfrak{f}_v belongs to $L((2g(\tilde{\mathcal{S}}) - 2)P_\infty)$ for all $v \in F$.

Proof. It has been already shown that \mathfrak{f}_v is regular outside P_∞ . It remains to show that $-v_{P_\infty}(\mathfrak{f}_v) \leq 2g(\tilde{\mathcal{S}}) - 2$. As before, let $\sigma = a_1 + a_2 + a_3 + a_4 + f$.

- If $v \in F_1$, then $-v_{P_\infty}(\mathbf{f}_v)$ is equal to $\sigma(q^2 + 1) - a_1(2qq_0 - q + 1) - a_2(qq_0 - q_0 + 1) - a_3(q - 2q_0 + 1) \leq (q - 2)(q^2 + 1) = 2g(\tilde{\mathcal{S}}) - 2$.
- If $v \in F_2$, then $-v_{P_\infty}(\mathbf{f}_v)$ is equal to $(q - 2)(q^2 + 1) - a_1(2qq_0 - q + 1) - a_2(qq_0 - q_0 + 1) - a_3(q - 2q_0 + 1) \leq (q - 2)(q^2 + 1) = 2g(\tilde{\mathcal{S}}) - 2$.
- If $v \in F_3$, then $-v_{P_\infty}(\mathbf{f}_v)$ is at most $(q - 2)(q^2 + 1) - a_1(2qq_0 - q + 1) - a_2(qq_0 - q_0 + 1) - a_3(q - 2q_0 + 1) \leq (q - 2)(q^2 + 1) = 2g(\tilde{\mathcal{S}}) - 2$.
- If $v \in F_4$, then $-v_{P_\infty}(\mathbf{f}_v)$ is at most $(q - 2)(q^2 + 1) - a_1(2qq_0 - q + 1) - a_2(qq_0 - q_0 + 1) - a_3(q - 2q_0 + 1) \leq (q - 2)(q^2 + 1) = 2g(\tilde{\mathcal{S}}) - 2$.
- If $v \in F_5$, then $-v_{P_\infty}(\mathbf{f}_v)$ is at most $(q - 2)(q^2 + 1) - a_2(qq_0 - q_0 + 1) - a_3(q - 2q_0 + 1) \leq (q - 2)(q^2 + 1) = 2g(\tilde{\mathcal{S}}) - 2$.
- If $v \in F_6$, then $-v_{P_\infty}(\mathbf{f}_v)$ is at most $(q - 2)(q^2 + 1) - a_3(q - 2q_0 + 1) \leq (q - 2)(q^2 + 1) = 2g(\tilde{\mathcal{S}}) - 2$.

□

It is possible, though a bit technical, to determine the Apéry set and a set of generators of $H(P)$ for $P \notin \tilde{\mathcal{S}}(\mathbb{F}_{q^4})$. It turns out that for $s = 1$, one needs 19 generators, while for $s \geq 2$, $H(P)$ has $3q - 2q_0$ generators. More details will appear in the upcoming PhD thesis of the second author.

Acknowledgements

The first author would like to acknowledge the support from The Danish Council for Independent Research (DFF-FNU) for the project *Correcting on a Curve*, Grant No. 8021-00030B.

References

- [1] D. Bartoli, M. Montanucci and G. Zini, *Weierstrass semigroups at every point of the Suzuki curve*, Acta Arithmetica, to appear.
- [2] P. Beelen and M. Montanucci, *A new family of maximal curves*, J. Lond. Math. Soc. **98**, 573–592 (2018).
- [3] P. Beelen and M. Montanucci, *Weierstrass semigroups on the Giulietti-Korchmáros curve*, Finite Fields and Appl. **52**, 10–29 (2018).
- [4] A. Eid and I. Duursma, *Smooth Embeddings for the Suzuki and Ree Curves*, Algorithmic Arithmetic, Geometry, and Coding Theory **637**, 251–291 (2015).
- [5] A. Garcia, C. Güneri, and H. Stichtenoth, *A generalization of the Giulietti-Korchmáros maximal curve*, Adv. Geom. **10**, 427–434 (2010).
- [6] A. Garcia and P. Viana, *Weierstrass points on certain non-classical curves*, Arch. Math. **46**, 315–322 (1986).
- [7] M. Giulietti and G. Korchmáros, *A new family of maximal curves over a finite field*, Math. Ann. **343**, 229–245 (2009).

- [8] M. Giulietti, M. Montanucci, L. Quoos and G. Zini, *On some Galois covers of the Suzuki and Ree curves*, Journal of Number Theory **189**, 220–254 (2018).
- [9] R. Graham, D. E. Knuth, O. Patashnik, *Concrete mathematics*. A foundation for computer science. Second edition. Addison-Wesley Publishing Company, Reading, MA, xiv+657 pp, (1994).
- [10] H.W. Henn, *Funktionenkörper mit großer Automorphismengruppe*, J. Reine Angew. Math. **302**, 96–115 (1978).
- [11] J.W.P. Hirschfeld, G. Korchmáros and F. Torres, *Algebraic Curves over a Finite Field, Princeton Series in Applied Mathematics*, Princeton, (2008).
- [12] S. Karanikolopoulos and A. Kontogeorgis, *Automorphisms of Curves and Weierstrass semigroups*, preprint, arXiv:1005.2871.
- [13] G. Lachaud, *Sommes d’Eisenstein et nombre de points de certaines courbes algébriques sur les corps finis*, C.R. Acad. Sci. Paris **305** (Série I), 729–732 (1987).
- [14] D. Skabelund, *New maximal curves as ray class fields over Deligne-Lusztig curves*, Proceedings of the American Mathematical Society **146**, 525–540 (2017).
- [15] H. Stichtenoth, *Algebraic function fields and codes*, Graduate Texts in Mathematics **254**, Springer, Berlin, (2009).
- [16] K.O. Stöhr, J.F. Voloch, *Weierstrass points and curves over finite fields*, Proc. London Math. Soc. **52**, 1–19 (1986).
- [17] M.A. Tsfasman, G. Vladut, *Algebraic-geometric Codes*, Kluwer, Dordrecht, (1991).
- [18] G. D. Villa Salvador, *Topics in the theory of algebraic function fields*, Mathematics: Theory & Applications. Birkhäuser Boston, Inc., Boston, MA, (2006).

Peter Beelen, Maria Montanucci and Leonardo Landi Technical University of Denmark,
 Department of Applied Mathematics and Computer Science,
 Matematiktorvet 303B,
 2800 Kgs. Lyngby,
 Denmark,
 pabe@dtu.dk
 lelan@dtu.dk
 marimo@dtu.dk