Deletion of RapPhr systems in *Bacillus subtilis* influences in vitro biofilm formation and plant root colonization

Nordgaard, Mathilde; Mortensen, Rasmus Møller Rosenbek; Kirk, Nikolaj Kaae; GallegosMonterrosa, Ramses; Kovács, Ákos T.

Published in:
MicrobiologyOpen

Link to article, DOI:
[10.1002/mbo3.1212](https://doi.org/10.1002/mbo3.1212)

Publication date:
2021

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Nordgaard, M., Mortensen, R. M. R., Kirk, N. K., GallegosMonterrosa, R., & Kovács, Á. T. (2021). Deletion of RapPhr systems in *Bacillus subtilis* influences in vitro biofilm formation and plant root colonization. *MicrobiologyOpen*, 10(3), [e1212]. <https://doi.org/10.1002/mbo3.1212>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Deletion of Rap-Phr systems in *Bacillus subtilis* influences in vitro biofilm formation and plant root colonization

Mathilde Nordgaard¹ | Rasmus Møller Rosenbek Mortensen¹ | Nikolaj Kaae Kirk¹ |
Ramses Gallegos-Monterrosa² | Ákos T. Kovács¹

¹Bacterial Interactions and Evolution Group, DTU Bioengineering, Technical University of Denmark, Lyngby, Denmark

²Terrestrial Biofilms Group, Institute of Microbiology, Friedrich-Schiller-University Jena, Jena, Germany

Correspondence

Ákos T. Kovács, Bacterial Interactions and Evolution Group, DTU Bioengineering, Technical University of Denmark, 2800 Kongens Lyngby, Denmark.
Email: atkovacs@dtu.dk

Funding information

DTU; Novo Nordisk Foundation, Grant/Award Number: NNFOC0055625

Abstract

Natural isolates of the soil-dwelling bacterium *Bacillus subtilis* form robust biofilms under laboratory conditions and colonize plant roots. *B. subtilis* biofilm gene expression displays phenotypic heterogeneity that is influenced by a family of Rap-Phr regulatory systems. Most Rap-Phr systems in *B. subtilis* have been studied independently, in different genetic backgrounds and under distinct conditions, hampering true comparison of the Rap-Phr systems' impact on bacterial cell differentiation. Here, we investigated each of the 12 Rap-Phr systems of *B. subtilis* NCIB 3610 for their effect on biofilm formation. By studying single $\Delta rap-phr$ mutants, we show that despite redundancy between the cell-cell communication systems, deletion of each of the 12 Rap-Phr systems influences matrix gene expression. These Rap-Phr systems therefore enable fine-tuning of the timing and level of matrix production in response to specific conditions. Furthermore, some of the $\Delta rap-phr$ mutants demonstrated altered biofilm formation *in vitro* and colonization of *Arabidopsis thaliana* roots, but not necessarily similarly in both processes, indicating that the pathways regulating matrix gene expression and other factors important for biofilm formation may be differently regulated under these distinct conditions.

KEYWORDS

Bacillus subtilis, biofilm formation, matrix gene expression, Rap-Phr, root colonization

1 | INTRODUCTION

In nature, biofilms are the predominant lifestyle of bacteria and are known as surface-associated microbial communities embedded in a self-produced matrix (Flemming & Wingender, 2010; Hall-Stoodley et al., 2004; López et al., 2010). Biofilms have been widely studied since they represent a fascinating example of microbial development in response to environmental cues (O'Toole et al., 2000). Furthermore, studying biofilms is of special interest due to their detrimental impact in clinical and industrial settings (Di Pippo et al.,

2018; Stewart, 2002) as well as their promising potential within the biotechnology industry (Blake et al., 2021; Singh et al., 2006). Regarding the latter, the gram-positive bacterium *Bacillus subtilis* has in the last two decades gained interest due to its promising potential as a biocontrol agent within agriculture (Kiesewalter et al., 2021; Ongena & Jacques, 2008). In its natural habitat, the soil-dwelling bacterium colonizes plants by forming a biofilm on the root (Bais et al., 2004; Beauregard et al., 2013; Chen et al., 2012). After successfully colonizing the root, *B. subtilis* exerts its plant-beneficial properties, including directly promoting plant growth and protecting

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *MicrobiologyOpen* published by John Wiley & Sons Ltd.

the plant against diseases (Blake et al., 2021). Furthermore, *B. subtilis* forms spores that are highly resistant to extreme environments (Piggot & Hilbert, 2004) facilitating easy formulation (Ongena & Jacques, 2008).

Bacillus subtilis can easily be isolated from the rhizosphere of plants (Fall et al., 2004), and a study performed by Chen et al. (2013) showed that the majority of natural strains isolated from the rhizosphere formed architecturally complex biofilms under laboratory conditions, indicating that biofilm formation is an important trait for *B. subtilis* to thrive in its natural habitat. In the laboratory, *B. subtilis* has long been studied using different kinds of biofilm models including colonies at the air-agar interface and floating biofilms formed at the air-liquid interface, termed pellicles (Arnaouteli et al., 2021).

A prevalent feature of *B. subtilis* biofilms is that they display complex phenotypic heterogeneity, where genetically identical cells differentiate into distinct cell types in response to external cues (López & Kolter, 2010; Lopez et al., 2009). The variation in environmental conditions throughout the biofilm (Costerton et al., 1994) thereby leads to a heterogeneous population with different cell types performing distinct tasks and occupying different micro-niches. The extracellular signals triggering cell differentiation include quorum-sensing molecules, natural products, and nutrient availability that activate a set of sensor kinases (Arnaouteli et al., 2021; Mhatre et al., 2014). Once activated, the sensor kinases phosphorylate their respective master transcriptional regulators, Spo0A, DegU, and ComA, each of which activates different sets of genes (López & Kolter, 2010). The Spo0A pathway governs differentiation into matrix-producing cells and sporulating cells. In response to external cues, one or more of five histidine kinases, KinA-E, are activated which results in phosphorylation of Spo0F. Spo0F~P then transfers its phosphoryl group to Spo0B, which in turn transfers the phosphoryl group to and thereby activates Spo0A (Fujita et al., 2005; Jiang, Shao, et al., 2000). At low Spo0A~P levels, the genes involved in the synthesis of matrix components, exopolysaccharide (EPS) and TasA protein fiber, are expressed (Cairns et al., 2014; Fujita et al., 2005). These two matrix components are well known to be required for biofilm formation *in vitro* and on the plant root (Beauregard et al., 2013; Branda et al., 2006). When high levels of Spo0A~P are reached, genes involved in sporulation are expressed (Fujita et al., 2005). The DegU response regulator is phosphorylated by its cognate histidine kinase DegS. Studies have indicated that inhibition of flagellar rotation, as may take place upon contact with a surface, acts as a mechanical trigger to activate the DegS-DegU two-component signaling pathway (Cairns et al., 2013). At very low levels of DegU~P, genes related to swarming motility are expressed, while elevated levels of DegU~P induce exoprotease production and at the same time represses motility genes (Belas, 2013; Verhamme et al., 2007). Finally, the pheromone ComX activates the histidine kinase ComP, which phosphorylates ComA, resulting in the expression of genes involved in competence development and surfactin production (Comella & Grossman, 2005).

This regulatory network governing cell differentiation in *B. subtilis* is further regulated by a family of response regulator

aspartyl-phosphate (Rap) phosphatases and their associated phosphatase regulator (Phr) peptides (Perego, 2013). In the *B. subtilis* group, 80 distinct putative *rap-phr* alleles have been identified with a strain having on average 11 *rap* genes (Even-Tov et al., 2016). The abundance of Rap and Phr peptides is transcriptionally controlled in response to different cellular signals (Auchtung et al., 2005; Jarmer et al., 2001; Jiang, Grau, et al., 2000; Lazazzera et al., 1999; McQuade et al., 2001; Mueller et al., 1992; Ogura et al., 2001; Perego et al., 1994). The genes encoding the Rap-Phr pairs are found as gene cassettes with the *phr* gene immediately downstream of the *rap* gene and the expression of these being transcriptionally coupled, with some *phr* genes also being transcribed independently of their cognate *rap* genes from promoters controlled by σ^H (McQuade et al., 2001; Pottahil & Lazazzera, 2003; Reizer et al., 1997). Some exceptions to this exist, for example, the *rapB* gene is not followed by an active peptide encoding gene (Perego et al., 1996). Moreover, some Rap proteins are regulated by Phr peptides encoded in other cassettes, for example, RapB and J are both controlled by PhrC (Parashar, Jeffrey, et al., 2013). When expressed, the Rap phosphatases exert their effect within the cell by either dephosphorylating Spo0F~P (thus hindering Spo0A phosphorylation) or inhibiting the DNA-binding activity of ComA or DegU (Perego, 2013). In contrast, the product of the *phr* gene is secreted out of the cell through the Sec-dependent export pathway and processed into mature five to six amino acid signaling peptides. At high cell density, the Phr peptides reach threshold concentrations at which they are transported back into the cell by the oligopeptide permease (Opp) (Perego, 2013; Pottahil & Lazazzera, 2003). Once within the cell, the Phr peptides will inhibit their cognate Rap proteins, thereby relieving the inhibition of the master regulators resulting in altered gene expression (Perego, 2013; Pottahil & Lazazzera, 2003). The Rap-Phr systems thereby act as cell-cell signaling systems in *B. subtilis*, allowing the bacteria to respond to environmental changes only at sufficient cell densities.

As expected by the diversity and abundance of multiple Rap-Phr systems regulating the activity of these three master regulators, the Rap phosphatases show high redundancy in their regulatory function: RapA, B, E, H, I, J, and P have been shown to dephosphorylate Spo0F~P, RapC, D, F, H, K, and P regulate ComA, while RapG has been shown to regulate the activity of DegU (Auchtung et al., 2006; Ogura & Fujita, 2007; Perego, 2013; Omer Bendori et al., 2015) (Table A1). Furthermore, RapI is involved in the regulation of mobile genetic elements, as it activates the propagation of the mobile genetic element that encodes it (Auchtung et al., 2005). The regulation of the master regulators by multiple Rap phosphatases allows the integration of diverse signals to control cell differentiation in response to different conditions. However, this overall overview of the Rap-Phr signaling network in *B. subtilis* is based on studies where most Rap-Phr systems have been tested in different genetic backgrounds and under distinct cultivation conditions (Perego, 2013). Additionally, previous investigations in *B. subtilis* have directed their study toward certain targets of Rap-Phr regulation, with RapA and B being mostly studied

for their impact on sporulation (Perego & Hoch, 1996), while RapC and F are involved in competence development (Bongiorni et al., 2005; Core & Perego, 2003). So far, only RapP has been demonstrated to impact biofilm formation (Parashar, Konkol, et al., 2013; Omer Bendori et al., 2015). We previously studied all 12 Rap-Phr systems of the undomesticated strain *B. subtilis* NCIB 3610 in the same genetic background by following the relative abundance of all possible single and double $\Delta rap-phr$ mutants as well as the wild-type WT (79 strains) in populations subjected to different selective conditions. This study highlighted that the variability in Rap-Phr systems affected the ability to compete in diverse environments (Gallegos-Monterrosa et al., 2021).

In this study, we systematically investigated the contribution of each of the 12 Rap-Phr systems in *B. subtilis* 3610 to biofilm formation. We assessed wild type (WT) and the 12 single $\Delta rap-phr$ mutants for matrix gene expression and biofilm formation under different conditions. We found that all 12 mutants showed altered matrix gene expression compared with the WT. Furthermore, we observed that the Rap-Phr modules affect not only in vitro biofilm formation but also the colonization of plant roots that represents an ecologically relevant environment.

2 | MATERIALS AND METHODS

2.1 | Bacterial strains and cultivation methods

Strains used in this study are listed in Table 1. The *B. subtilis* DK1042 strain (a natural competent derivative of the undomesticated NCIB 3610) (Konkol et al., 2013) was used as WT. The 12 single $\Delta rap-phr$ mutants were previously created and contain an antibiotic resistance cassette in place of the *rap-phr* gene pair (except for the markerless $\Delta rapB$ mutant) (Gallegos-Monterrosa et al., 2021). For flow cytometry, the $\Delta rap-phr$ mutants were transformed with the plasmid phy_mKATE2 harboring the *mKATE* gene under the control of the hyper-spank promoter (which is constitutive due to the absence of *lacI*) and a chloramphenicol (Chl) resistance gene within the flanking regions of the *amyE* gene (van Gestel et al., 2014). Transformants were identified by selecting for Chl resistance, and double crossovers were verified by the loss of amylase activity. The resulting mKATE-labelled $\Delta rap-phr$ mutants were transformed with genomic DNA from *B. subtilis* TB373, which harbors the *P_{tapA}-gfp* reporter construct with a kanamycin (Km) resistance gene integrated at the *sacI* locus. Successful transformants with the reporter construct inserted into the *sacI* locus were identified by selecting for Km resistance. The resulting reporter strains were verified for reporter activity under the fluorescence stereomicroscope. Importantly, due to the presence of a Km resistance cassette in place of the corresponding *rap-phr* gene pair in $\Delta rapA$, C, D, I, J, and K, markerless versions of those *rap-phr* mutants were used to obtain the reporter strains for the *tapA-sipW-tasA* operon for the flow cytometry analysis. For all experiments, strains were grown overnight in Lysogeny broth (LB; Lennox, Carl Roth; 10 g·L⁻¹ tryptone, 5 g·L⁻¹

TABLE 1 Strains used in this study

Name	Description	Reference
DK1042	NCIB 3610 <i>comI</i> ^{Q121}	Konkol et al. (2013)
TB499	DK1042 <i>rapA-phrA</i> ::Km ^R	Gallegos-Monterrosa et al. (2021)
TB575	DK1042 $\Delta rapB$	
TB396	DK1042 $\Delta rapC-phrC$::Km ^R	
TB315	DK1042 $\Delta rapD$::Km ^R	
TB339	DK1042 $\Delta rapE-phrE$::Spec ^R	
TB341	DK1042 $\Delta rapF-phrF$::Spec ^R	
TB404	DK1042 $\Delta rapG-phrG$::Spec ^R	
TB405	DK1042 $\Delta rapH-phrH$::Spec ^R	
TB272	DK1042 $\Delta rapI-phrI$::Km ^R	
TB274	DK1042 $\Delta rapJ$::Km ^R	
TB557	DK1042 $\Delta rapK-phrK$::Km ^R	
TB435	DK1042 $\Delta rapP-phrP$::Mls ^R	
TB588	DK1042 $\Delta rapA-phrA$	
TB410.1	DK1042 $\Delta rapC-phrC$	
TB513	DK1042 $\Delta rapD$	
TB444	DK1042 $\Delta rapI-phrI$	
TB411.2	DK1042 $\Delta rapJ$	
TB587	DK1042 $\Delta rapK-phrK$	
DTUB189	DK1042 $\Delta rapA-phrA$, <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R	This study
DTUB165	DK1042 $\Delta rapB$, <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R	This study
DTUB190	DK1042 $\Delta rapC-phrC$, <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R	This study
DTUB191	DK1042 $\Delta rapD$, <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R	This study
DTUB159	DK1042 $\Delta rapE-phrE$::Spec ^R , <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R	This study
DTUB160	DK1042 $\Delta rapF-phrF$::Spec ^R , <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R	This study
DTUB166	DK1042 $\Delta rapG-phrG$::Spec ^R , <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R	This study
DTUB167	DK1042 $\Delta rapH-phrH$::Spec ^R , <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R	This study
DTUB192	DK1042 $\Delta rapI-phrI$, <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R	This study
DTUB193	DK1042 $\Delta rapJ$, <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R	This study

(Continues)

TABLE 1 (Continued)

Name	Description	Reference
DTUB194	DK1042 $\Delta rapK$ - <i>phrK</i> , <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R	This study
DTUB173	DK1042 $\Delta rapP$ - <i>phrP</i> ::Mls ^R , <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R	This study
TB34	DK1042 <i>amyE</i> ::P _{hyspank} - <i>gfp</i> Chl ^R	Mhatre et al. (2017)
TB35	DK1042 <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R	Hölscher et al. (2016)
TB865	DK1042 <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R , <i>sacA</i> ::P _{tapA} - <i>gfp</i> Km ^R	Dragoš et al. (2018)
DTUB284	DK1042 $\Delta rapA$ - <i>phrA</i> , <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R , <i>sacA</i> ::P _{tapA} - <i>gfp</i> Km ^R	This study
DTUB285	DK1042 $\Delta rapB$, <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R , <i>sacA</i> ::P _{tapA} - <i>gfp</i> Km ^R	This study
DTUB286	DK1042 $\Delta rapC$ - <i>phrC</i> , <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R , <i>sacA</i> ::P _{tapA} - <i>gfp</i> Km ^R	This study
DTUB287	DK1042 $\Delta rapD$, <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R , <i>sacA</i> ::P _{tapA} - <i>gfp</i> Km ^R	This study
DTUB288	DK1042 $\Delta rapE$ - <i>phrE</i> ::Spec ^R , <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R , <i>sacA</i> ::P _{tapA} - <i>gfp</i> Km ^R	This study
DTUB289	DK1042 $\Delta rapF$ - <i>phrF</i> ::Spec ^R , <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R , <i>sacA</i> ::P _{tapA} - <i>gfp</i> Km ^R	This study
DTUB290	DK1042 $\Delta rapG$ - <i>phrG</i> ::Spec ^R , <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R , <i>sacA</i> ::P _{tapA} - <i>gfp</i> Km ^R	This study
DTUB291	DK1042 $\Delta rapH$ - <i>phrH</i> ::Spec ^R , <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R , <i>sacA</i> ::P _{tapA} - <i>gfp</i> Km ^R	This study
DTUB292	DK1042 $\Delta rapI$ - <i>phrI</i> , <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R , <i>sacA</i> ::P _{tapA} - <i>gfp</i> Km ^R	This study
DTUB293	DK1042 $\Delta rapJ$, <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R , <i>sacA</i> ::P _{tapA} - <i>gfp</i> Km ^R	This study
DTUB294	DK1042 $\Delta rapK$ - <i>phrK</i> , <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R , <i>sacA</i> ::P _{tapA} - <i>gfp</i> Km ^R	This study
DTUB295	DK1042 $\Delta rapP$ - <i>phrP</i> ::Mls ^R , <i>amyE</i> ::P _{hyspank} - <i>mKATE</i> Chl ^R , <i>sacA</i> ::P _{tapA} - <i>gfp</i> Km ^R	This study
TB373	DK1042 <i>sacA</i> ::P _{tapA} - <i>gfp</i> Km ^R	Gallegos-Monterrosa et al. (2016)

yeast extract and 5 g·L⁻¹ NaCl) at 37°C while shaking (220 rpm). For transformation and stock preparation, antibiotics were used at the following working concentrations: Km: 5 µg ml⁻¹ and Chl: 5

µg·ml⁻¹. For analyzing biofilm formation and matrix gene expression, strains were grown in MSgg (5 mmol·L⁻¹ potassium phosphate [pH 7], 0.1 mol·L⁻¹ 3-(N-morpholino)propanesulfonic acid (MOPS) [pH 7], 2 mmol·L⁻¹ MgCl₂, 700 µmol·L⁻¹ CaCl₂, 100 µmol·L⁻¹ MnCl₂, 50 µmol·L⁻¹ FeCl₃, 1 µmol·L⁻¹ ZnCl₂, 2 µmol·L⁻¹ thiamine, 0.5 % glycerol, and 0.5 % K-glutamate). For root colonization assay, strains were grown in MSNg (5 mmol·L⁻¹ potassium phosphate buffer [pH 7], 0.1 mol·L⁻¹ MOPS [pH 7], 2 mmol·L⁻¹ MgCl₂, 50 µmol·L⁻¹ MnCl₂, 1 µmol·L⁻¹ ZnCl₂, 2 µmol·L⁻¹ thiamine, 700 µmol·L⁻¹ CaCl₂, 0.2 % NH₄Cl₂, and 0.05 % glycerol).

2.2 | Microscopy imaging

All images were acquired with an Axio Zoom V16 stereomicroscope (Carl Zeiss, Germany) equipped with a Zeiss CL 9000 LED light source and an AxioCam MRm monochrome camera (Carl Zeiss, Germany).

2.3 | Biofilm formation assay

For biofilm formation on a solid surface, 7 µl overnight cultures were spotted on MSgg supplemented with 1.5 % agar. For pellicle biofilm formation at the air-liquid interface, 15 µl inoculum of overnight cultures adjusted to OD₆₀₀ of 5 was added to 1.5 ml MSgg medium in 24-well plates, giving a starting OD₆₀₀ of 0.05. Plates were incubated under static conditions at 30°C for 48 h, thereafter images of the arisen colonies and pellicles were obtained using the stereomicroscope.

2.4 | Root colonization assay

Arabidopsis thaliana Col-0 plants were used as a host for *B. subtilis* root colonization. *A. thaliana* seeds were sterilized in 2 % (v/v) sodium hypochlorite (NaOCl) for 10 min with an orbital shaker. Following this, NaOCl was removed and the seeds were washed five times in sterile water. Sterilized seeds were placed in MS agar plates (Murashige and Skoog basal salts, Sigma) (2.2 g·L⁻¹) with approximately 20 seeds per petri dish. The Petri dishes were wrapped in parafilm and left for stratification at 4°C for 3 days to break seed dormancy and were then moved to the plant chamber (cycles of 16 h light at 24°C and 8 h dark at 20°C). After five to seven days, seedlings of 0.5–1.2 cm in size were placed in 48-well plates containing 270 µl MSNg medium per well. To each well, 30 µl of overnight culture adjusted to OD₆₀₀ of 0.2 was added resulting in a final starting OD₆₀₀ of 0.02. The plates were sealed with parafilm and incubated in the plant chamber while shaking at 90 rpm for 16 h. Seedlings were then washed in minimal salts nitrogen glycerol (MSNg) to remove non-attached cells from the root. The washed seedlings were placed in Eppendorf tubes containing 1 ml of NaCl (0.9 %) and subjected to standard sonication protocol to disperse the biofilm (Dragoš et al., 2018). The resulting

bacterial cell suspension was diluted and plated on LB agar plates for CFU counting. To acquire the CFU per mm root, the obtained CFU was divided by the length of the corresponding root.

2.5 | Growth profiling

To follow the growth of WT and mutants, two overnight cultures of each strain were independently inoculated into a 96-well plate containing MSgg or LB broth, at a starting OD₆₀₀ of 0.05. Growth was monitored in a plate reader (Infinite F200 PRO, TECAN, and BioTek Synergy HTX Multi-Mode Microplate Reader) every 10 min for 48 h at 30°C under linear shaking conditions.

2.6 | Matrix gene expression assays by flow cytometry

To measure expression from the matrix operon *tapA-sipW-tasA*, mKate-labelled WT and $\Delta rap-phr$ mutants harboring the reporter construct $P_{tapA}-gfp$ were incubated in 10 ml MSgg in 100 ml bottles with a starting OD₆₀₀ of 0.02. Lids were loosely on, so oxygen would not be depleted. Bottles were incubated at 37°C at 220 rpm for 6 h. After incubation, 1 ml of each sample was transferred to a 2 ml Eppendorf tube, and samples were run on the flow cytometer (MACSQuant® VYB, Miltenyibiotec). mKate-positive cells were detected by yellow laser (561 nm) and filter Y3 (661/20 nm). Green fluorescent cells, representing the cells expressing the *tapA-sipW-tasA* operon, were detected by the blue laser (488 nm) and filter B1 (525/50 nm). Strain TB35 with constitutive mKate expression was used as a negative control for GFP expression, while strain TB34 with constitutive GFP expression was used as a positive control for GFP expression. In addition, TB34 and a medium control were used to identify the red background fluorescence noise due to autofluorescence, and cells above this background fluorescence were identified as the mKate-positive cells representing *B. subtilis* cells producing the mKate protein. For each WT and mutant sample, single events were identified on the SSC-H vs SSC-A plot and gated into the mKate-A vs SSC-A plot, where mKate-positive cells were identified. These gated cells were exported and read into Excel where the green fluorescence (GFP-A) values were used for analysis. To get rid of negative values, 300 AU was added to all events in the samples. To obtain the distribution of GFP expression, data obtained from each replicate were subjected to binning with an identical bin size (of 50). Events with GFP expression from 0 to 10,000 were included as the majority of events were within this interval (>98%). Next, the number of events in each bin was divided by the total number of events in the given replicate, resulting in the normalized frequency. To obtain the mean distribution, a mean frequency for each bin was obtained by averaging the individual frequencies within this bin across the replicates, resulting in the mean distribution of single-cell-level expression. For statistics, the relative OFF (GFP values between 0 and 500) and ON (GFP

values between 500–10,000) populations were calculated for each replicate, as was the mean fluorescence of the ON population.

2.7 | Statistical analysis

Statistical analyses were performed in R Studio. For each root colonization assay, a one-way analysis of variance (ANOVA) was performed on the log10-transformed data. P-values were adjusted using the Benjamini & Hochberg procedure. When adjusted P-values were significant ($p < 0.05$), a Dunnett's multiple comparison test was performed. When data failed to meet parametric assumptions (normality and equal variance), a non-parametric analysis (Kruskal–Wallis test) was used. If adjusted P-values were significant ($p < 0.05$), treatment means were compared via a Dunn test. To test for significant differences in relative OFF/ON population and mean fluorescence of ON population between mutants and WT, an ANOVA followed by Dunnett's multiple comparison test was performed. The significance level for all tests was set at 5 %.

3 | RESULTS

3.1 | Single $\Delta rap-phr$ mutants show altered matrix gene expression compared with the WT

To study the role of Rap-Phr systems of *B. subtilis* in biofilm formation, we used the *B. subtilis* DK1042 strain (a natural competent derivative of the undomesticated NCBI 3610) (Konkol et al., 2013) as WT. In contrast to domesticated strains which may have acquired mutations to the Rap-Phr systems or even lost some of them, NCIB 3610 and the derived DK1042 (hereafter referred to as 3610) contain all 12 *rap-phr* modules (McLoon et al., 2011). Rap-Phr systems of *B. subtilis* have been studied using both overexpression and deletion mutants of the Rap-Phr systems as reviewed by Perego (2013). Interestingly, the absence of Rap modules in a natural isolate of *B. subtilis* was shown to influence sporulation timing (Serra et al., 2014). In this study, we employed single $\Delta rap-phr$ mutants of *B. subtilis* created in our previous work (Gallegos-Monterrosa et al., 2021) to study the impact of the Rap-Phr modules on biofilm formation.

The diversity in Rap-Phr regulatory systems and the reported role of several of the Rap phosphatases in regulating the activity of Spo0A (Perego, 2013; Omer Bendori et al., 2015), which controls matrix gene expression, prompted us to test WT and the 12 single $\Delta rap-phr$ mutants for matrix gene expression. To quantify matrix gene expression during biofilm formation, we used a previously established approach (Kearns et al., 2005). This method utilizes the observation that expression from the promoter of the *tapA-sipW-tasA* operon, responsible for the production of the protein component of the extracellular matrix (Branda et al., 2006), is highly induced during the late exponential growth phase under shaking conditions in MSgg, a medium known to induce biofilm formation. After 6 h of growth in MSgg under shaking conditions, the fluorescence intensities of WT and mutants harboring a $P_{tapA}-gfp$

reporter construct were measured using flow cytometry. Importantly, the side scatter (SSC-A) vs forward scatter (FSC-A) plots revealed no big changes in cell granularity or size of the mutants compared with the WT (Figure A1). Since RapA, B, E, H, I, J, and P have been reported to dephosphorylate Spo0F~P (Perego, 2013; Omer Bendori et al., 2015), we hypothesized that the mutants lacking one of these Rap phosphatases would have a larger relative ON population (i.e., cells in the high expression state), due to more cells committing to matrix production, and/or display a higher mean expression from P_{tapA} in the ON population due to earlier matrix gene expression, as compared to the WT. Alternatively, if redundancy operates between some of the Rap-Phr

systems, the absence of one Rap-Phr system may have only minor or insignificant effects due to the expression and function of another redundant system. However, $\Delta rapA$ showed a significantly smaller relative ON population ($p = 0.022$, $n = 3-8$) than the WT (Figure 1), while all other mutants showed a significantly higher mean expression of the ON population compared with the WT. The increased mean expression by the mutants could potentially be caused by enhanced growth in MSgg, as strains growing faster would reach the threshold density for biofilm formation earlier, and thus show increased matrix gene expression. However, when the optical density (OD) at 600 nm was measured after 6 h in minimal medium (MSgg) prior to flow cytometry analysis,

FIGURE 1 Expression of the *tapA-sipW-tasA* operon in *Bacillus subtilis* WT and $\Delta rap-phr$ mutants after growth in MSgg under shaking conditions. Flow cytometry analysis showing the average distributions of GFP expression of $\Delta rap-phr$ mutants and WT harboring the P_{tapA} -*gfp* construct ($n = 3-8$). The average WT distribution is shown in each graph for comparison. Orange = WT, green = mutant, blue = non-labelled control strain. Letters denote the corresponding $\Delta rap-phr$ mutants, that is, a = $\Delta rapA$, b = $\Delta rapB$ and so forth. AU indicates arbitrary units. The significant difference in the mean fluorescence intensity of the ON population (GFP values between 500 and 10,000) between mutants and WT were tested by an ANOVA followed by Dunnett's multiple comparison test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

the mutants showed either similar or reduced OD₆₀₀ compared with WT, thus excluding this explanation. Interestingly, the increased matrix gene expression was not limited to the mutants lacking one of the Rap phosphatases reported to dephosphorylate Spo0F~P activity. These results show that all 12 Rap-Phr regulatory systems influence matrix gene expression under these conditions.

3.2 | Absence of certain Rap-Phr modules alters biofilm formation *in vitro*

Since matrix production is required for proper biofilm development (Branda et al., 2006; Dragoš et al., 2018), and most Δrap -*phr* mutants showed increased matrix gene expression under shaking conditions in MSgg, we hypothesized that this would manifest in more complex and robust biofilm formation. To investigate this, WT and mutants were tested for their ability to form biofilm on a solid surface (MSgg agar) and robust pellicle biofilms at the MSgg air-liquid interface. A pellicle forms when oxygen in the medium is exhausted and *B. subtilis* moves toward higher oxygen concentrations, the air-liquid surface, where cells create a biofilm (Hölscher et al., 2015).

In accordance with previous work, *B. subtilis* WT produced a wrinkled colony biofilm, as well as a robust, wrinkled pellicle (Branda et al., 2001; Gallegos-Monterrosa et al., 2016) (Figure 2). While $\Delta rapA$, *B*, *J*, and *K* formed more wrinkled colonies, $\Delta rapI$ and *P* formed complex but very small colonies on MSgg agar compared with the WT. In contrast, $\Delta rapC$ formed a large, transparent colony with fewer wrinkles than WT. When testing for the development of pellicle biofilms, the $\Delta rapA$, *C*, *I*, and *P* mutants formed thin and/or non-homogenous pellicles, while the qualitative assessment revealed no mutants with more complex or robust pellicles than WT (Figure 2). Lastly, the $\Delta rapD$, *E*, *F*, *G*, and *H* mutants formed comparable colony and pellicle biofilms to the WT. These results show that despite most Δrap -*phr* mutants showed increased matrix gene expression under shaking conditions compared with the WT, this did not necessarily manifest in the development of more complex biofilms on agar or at the air-liquid interface.

3.3 | Certain Δrap -*phr* mutants are affected in the colonization of *Arabidopsis thaliana* roots

To reveal how the Rap-Phr modules affect biofilm formation in a more ecologically relevant environment, WT and mutants were

FIGURE 2 Biofilm formation of *Bacillus subtilis* WT and single Δrap -*phr* mutants. Overnight cultures of *B. subtilis* WT and Δrap -*phr* mutants were spotted on MSgg medium solidified with 1.5 % agar (left) or inoculated in MSgg with a starting OD₆₀₀ = 0.05 (right). Images were taken from above after 48 h of incubation at 30°C using a stereomicroscope. Letters denote the corresponding Δrap mutant, that is, a = $\Delta rapA$, b = $\Delta rapB$, and so forth, but l = $\Delta rapP$. Bar denotes 5 mm for the biofilm colony images

FIGURE 3 *Arabidopsis thaliana* root colonization by *Bacillus subtilis* WT and single $\Delta rap-phr$ mutants. To estimate the impact of each of the Rap-Phr modules on root colonization, WT and single $\Delta rap-phr$ mutants were inoculated onto five-day-old seedlings of *A. thaliana* ($n = 7-10$). After 16 h, CFU per mm root length was quantified. The log10-transformed value of CFU/mm root for each technical replicate was normalized to the mean of the WT from the same experiment. Each dot represents a root, while the mean value for each mutant is displayed as a red horizontal line. The black horizontal line represents the mean of the WT and the SD of the WT from each respective experiment is shown in shaded gray. For each assay, an ANOVA was performed on the log10-transformed values of CFU/mm root length. When significant ($p < 0.05$), means were compared via Dunnett's multiple comparison test with WT as the control. When data failed to meet parametric assumptions, a Kruskal-Wallis test was performed followed by a Dunn's test. ** $p < 0.005$, *** $p < 0.001$

tested for biofilm formation on the roots of the model plant organism *A. thaliana*. Similar to biofilm formation *in vitro* (Branda et al., 2006), previous studies have shown that matrix production is required for biofilm formation on *A. thaliana* roots (Beauregard et al., 2013; Dragoš et al., 2018). We, therefore, speculated that the increased matrix gene expression observed in most strains would enable enhanced biofilm formation on plant roots.

For this purpose, sterile *A. thaliana* seedlings with a root length of 0.5–1.2 cm were inoculated with one of the 12 $\Delta rap-phr$ mutants or WT and incubated in a plant chamber at 24°C (90 rpm) for 16 h, after which root colonization was quantified as a colony-forming unit (CFU) per mm root length. Of the 12 mutants tested, only the $\Delta rapD$, J , and P mutants showed significantly increased root colonization compared with the WT (Figure 3). In contrast, the $\Delta rapI$ mutant was significantly reduced in root colonization. The observation that $\Delta rapI$ and P showed similar biofilm formation *in vitro* (Figure 2) but performed oppositely during root colonization (Figure 3) might indicate that the effect of each Rap phosphatase on biofilm formation depends on the environment. Furthermore, similarly to biofilm formation *in vitro*, these results show that increased matrix gene expression under shaking conditions did not necessarily manifest in increased root colonization.

3.4 | Certain $\Delta rap-phr$ mutants show altered growth compared with WT

The timing of biofilm initiation *in vitro* and on the root depends on the cell density, as only at sufficiently high cell density the Phr peptides will reach threshold concentrations allowing them to be imported into the cell, where they will inhibit their cognate Rap

phosphatases. Consequently, a subset of the Rap phosphatases influences phosphorylation of Spo0A and the expression of biofilm genes. Biofilm formation is, therefore, affected by the growth rate, as strains growing faster will reach the threshold density for biofilm formation earlier. To test whether the observed changes in biofilm formation *in vitro* and on the root could be (partly) attributed to altered growth, WT and the 12 $\Delta rap-phr$ mutants were tested for growth in MSgg under shaking conditions. For clear visualization, the growth curves of the 12 mutants were separated into four plots (Figure 4). Of the 12 mutants tested, only the $\Delta rapA$, I , and P showed an altered growth profile compared with WT. $\Delta rapA$ and I showed a reduced growth rate, a delayed entry into the stationary phase, and reduced max OD₅₉₀. In addition, $\Delta rapI$ had a prolonged stationary phase and a slower decline during the late stationary phase (possibly death phase). $\Delta rapP$ showed slower growth during the exponential phase, which continued for about 10 h longer than the WT, but displayed a higher max OD₅₉₀ at the stationary phase. In LB medium, the same mutants were similarly or less affected in growth compared with WT (Figure A2). The impaired growth of $\Delta rapA$ and I is in accordance with these two mutants forming thin and/or non-homogeneous pellicles (Figure 2), and $\Delta rapI$ displaying reduced root colonization (Figure 3). In contrast, the prolonged exponential growth of $\Delta rapP$ is inconsistent with the thin pellicle formed by this mutant, but correlating with the increased root colonization observed for this mutant. Finally, $\Delta rapC$ and $\Delta rapJ$ showed similar growth profiles as the WT but $\Delta rapC$ formed a thin pellicle, while $\Delta rapJ$ was increased in root colonization. The differential biofilm-forming capacity *in vitro* and on the root observed for some of the mutants, interestingly, cannot be directly correlated with the growth rates and profiles of the mutants.

FIGURE 4 Growth of *Bacillus subtilis* WT and single $\Delta rap-phr$ mutants in MSgg. WT and mutants were inoculated into 96-well plates with a starting OD_{590} of 0.05. OD_{590} was measured every 10 min for 48 h at 30°C; each time point represents the mean of six technical replicates from two overnight cultures ($N = 6$). Error bars represent standard error (SE).

4 | DISCUSSION

For decades, the Rap-Phr regulatory systems which control the activity of the three master regulators governing cell differentiation in *B. subtilis* have been extensively studied (Perego, 2013; Pottahil & Lazazzera, 2003). However, since most studies have investigated the Rap-Phr networks of *B. subtilis* independently from each other, in different genetic backgrounds, and under distinct conditions, it is difficult to compare the results from these studies. Here, we investigated all 12 Rap-Phr systems found in *B. subtilis* 3610 for their effect on matrix gene expression and biofilm development *in vitro*. In addition, we examined the impact of Rap-Phr systems in the colonization of plant roots for the first time.

Several of the Rap phosphatases have been reported to dephosphorylate Spo0F~P (Perego, 2013; Omer Bendori et al., 2015) which is expected to influence matrix production, but only RapP has previously been demonstrated to affect matrix gene expression (Parashar, Konkol, et al., 2013; Omer Bendori et al., 2015). We

were, therefore, interested in testing the effect of each of the 12 Rap-Phr systems on matrix gene expression. Inspired by a previously established method (Kearns et al., 2005), expression from the promoter of the *tapA-sipW-tasA* operon was measured for WT and the 12 $\Delta rap-phr$ mutants in MSgg under shaking conditions. Only $\Delta rapA$ showed a significantly reduced relative ON population compared with the WT. Such reduced relative proportion of $\Delta rapA$ cells in the ON state may be due to more cells committing to activation of the sporulation pathway, and therefore, attenuating induction of matrix genes (Bischofs et al., 2009). Although only some of the Rap phosphatases have been reported to dephosphorylate Spo0F~P (Perego, 2013; Omer Bendori et al., 2015), we observed that all mutants except $\Delta rapA$ showed increased matrix gene expression. This indicates that despite the diversity of targets among Rap-Phr systems, and the seeming redundancy of several Rap phosphatases regulating the same master regulator (Perego, 2013), each of the 12 Rap-Phr systems has a regulatory role that affects matrix gene expression under the tested conditions. The involvement of 12 Rap-Phr systems

in influencing matrix gene expression suggests that the production of these costly public goods is under complex control, and allows the integration of multiple signals to fine-tune the timing of matrix production in response to different conditions (Auchtung et al., 2006; Dragoš et al., 2018).

Matrix production is well known to be required for the formation of architecturally complex biofilms under laboratory conditions (Arnaouteli et al., 2021; Branda et al., 2006). Moreover, matrix production and localized cell death are responsible for the formation of wrinkles during biofilm development (Asally et al., 2012; Branda et al., 2006; Gallegos-Monterrosa et al., 2017). We, therefore, speculated that the increased matrix gene expression observed for all mutants, except $\Delta rapA$, would manifest in these 11 mutants forming more wrinkled colonies and more complex, robust pellicles compared with the WT. However, only some of the mutants were affected in biofilm formation. In accordance with increased matrix gene expression, the $\Delta rapB$, J , and K mutants formed more wrinkled colonies, and $\Delta rapI$ and P formed complex, though very small colonies compared with the WT. Moreover, $\Delta rapC$ formed a larger, smoother, and more transparent colony. Furthermore, $\Delta rapA$, with the similar mean expression of the *tapA-sipW-tasA* operon and a smaller relative ON population compared with WT, also formed a more wrinkled colony. Surprisingly, none of the mutants displayed increased pellicle robustness or complexity compared with the WT. In contrast, $\Delta rapA$, C , I , and P formed thinner and/or non-homogenous pellicles.

Next, we were interested in studying how Rap-Phr systems affect biofilm formation of *B. subtilis* in a more ecologically relevant environment, that is, the plant root. Similar to biofilm formation *in vitro*, biofilm formation on the plant root by *B. subtilis* depends on matrix gene expression regulated by Spo0A (Beauregard et al., 2013; Chen et al., 2013). We, therefore, hypothesized that the increased matrix gene expression observed for most mutants would allow more bacterial cells to attach to and colonize the root. However, only $\Delta rapD$, J , and P showed increased root colonization, while $\Delta rapI$ was reduced in root colonization. The biofilm and root colonization experiments of the Δrap -*phr* mutants thus show that the magnitude of matrix gene expression under shaking conditions does not directly correlate with the ability to develop complex biofilms *in vitro* (on agar and at the air-liquid interface) or to colonize the plant root (Table A1). A lack of positive correlation between matrix gene expression and biofilm formation was observed in a previous study which showed that the magnitude of expression of *epsA-O* and *tasA-sipW-tasA* in *B. subtilis* 168 variants did not directly correlate with the formation of wrinkled biofilms (Gallegos-Monterrosa et al., 2016). These experiments could thus support that biofilm formation in *in vitro* and on plant roots is influenced by additional factors than just matrix gene expression. For example, surfactin production, which is regulated by ComA—a target of several Rap phosphatases (Perego, 2013)—was shown to influence the colony structure of *B. subtilis* NCIB 3610 on MSgg, though this secondary metabolite was not essential for pellicle formation and root colonization (Thérien et al., 2020). However, it has to be noted that matrix gene expression was measured under heavily shaking conditions (220 rpm), while colonies and pellicles

were developed under static conditions, and root colonization was assayed under mildly shaking conditions (90 rpm). An alternative explanation for the lack of correlation between matrix gene expression under shaking conditions and biofilm formation *in vitro* and on the root could be that the effect of the *rap-phr* deletions on *tapA* operon expression may vary between these different conditions. Further work is needed to fully explain the discrepancies observed in this study between *tapA* operon expression and biofilm formation.

Interestingly, several studies have reported a correlation between the ability of strains to form robust biofilms *in vitro* and to colonize the root—both within and among strains (Chen et al., 2013; Gallegos-Monterrosa et al., 2016). However, the ability of the Δrap -*phr* mutants to form biofilm *in vitro* did not necessarily reflect the ability to colonize the root (compare Figures 2 and 3 and the summary in Table A1). For example, $\Delta rapA$ and C formed thin and non-homogenous pellicles but were able to colonize the root to similar levels as the WT. $\Delta rapD$ displayed comparable biofilm *in vitro* to the WT but was significantly better in root colonization. In addition, $\Delta rapJ$ formed a highly wrinkled colony, but a pellicle similar to the WT, and was increased in root colonization. Finally, both $\Delta rapI$ and P showed reduced colony size and thin and/or non-homogenous pellicle formation, but while $\Delta rapI$ showed reduced root colonization, $\Delta rapP$ was increased in root colonization compared with the WT. These results indicate that the effect of the *rap-phr* deletions on biofilm formation varies between *in vitro* and root conditions. This was similarly shown for a $\Delta tagE$ mutant (deficient in glycosylating wall teichoic acid), which was affected in root colonization but displayed similar biofilm formation on agar and at the air-liquid interface as the WT (Tzipilevich & Benfey, 2021).

In the study by Gallegos-Monterrosa et al. (2016), showing that strains forming complex colonies and robust pellicles also efficiently colonize the root, the *B. subtilis* 168 stocks displayed genetic variation in distinct loci (e.g., *epsC* that encodes an enzyme that is directly involved in matrix production), resulting in large differences among the strains in their ability to form biofilm and colonize the root. In contrast, the Δrap -*phr* mutations studied here might only slightly modulate the regulatory pathways of *B. subtilis*; therefore, the ability of the mutants to form biofilm and colonize the root is less altered compared with WT. Nonetheless, the same study also demonstrated that biofilm development is influenced by medium composition (Gallegos-Monterrosa et al., 2016). Besides static vs mild agitation and a temperature difference (30 vs. 24°C), the media used for testing colony and pellicle formation, and for testing root colonization also slightly differ. First, the MSgg medium used for colony and pellicle formation contains a 10 times higher concentration of glycerol (0.5 %) compared with the MSNg medium used for plant root colonization (0.05%). During plant root colonization, the bacteria thus depend on plant polysaccharides and root exudates as carbon sources. In addition, *in vitro* biofilm development depends on the availability of iron and manganese (Kolodkin-Gal et al., 2013; Mhatre et al., 2016; Shemesh & Chai, 2013), while during plant root colonization in MSNg, biofilm formation is induced by plant polysaccharides and root exudates (Beauregard et al., 2013; Chen et al.,

2013). The presence of plant polysaccharides may, therefore, allow $\Delta rap-phr$ mutants that form weak pellicle biofilms *in vitro* to efficiently colonize the root ($\Delta rapA$, C and P, Figures 2 and 3). Thereby, the pathways regulating matrix gene expression and other factors important for biofilm formation may be differently regulated under distinct conditions.

Finally, the disparate results obtained in this study may be understood in light of the full set of 12 Rap-Phr systems with redundant functions: the influence of a single *rap-phr* deletion might be masked by the function of another redundant Rap-Phr system. Furthermore, if such potential redundancy varies between the different conditions employed in this study (e.g., if the *rap-phr* genes are differentially expressed under the distinct conditions tested), this could (partly) explain the observed discrepancy, for example, between biofilm formation *in vitro* and on the root.

To conclude, we here show that all 12 Rap-Phr systems have an impact on matrix gene expression in liquid culture. Thereby, the diversity in Rap-Phr systems in *B. subtilis* 3610 could function to integrate multiple signals to fine-tune the timing and level of matrix gene expression in response to new ecological niches, such as those it will encounter in soil. Furthermore, we show that the ability to form biofilm *in vitro* not necessarily reflects the ability to colonize the root under the tested conditions. These findings thus support that the pathways involved in matrix gene expression and other components important for biofilm establishment could be differently influenced under distinct conditions.

ACKNOWLEDGEMENTS

We thank Mikael Lenz Strube for his suggestions on statistics. The work was supported by a DTU Bioengineering start-up fund to ÁTK. Funding from Novo Nordisk Foundation (grant NNFOC0055625) for the infrastructure "Imaging microbial language in biocontrol (IMLiB)" is acknowledged.

CONFLICT OF INTERESTS

None declared.

AUTHOR CONTRIBUTIONS

Mathilde Nordgaard Christensen: Conceptualization (equal); Formal analysis (lead); Investigation (lead); Methodology (lead); Visualization (lead); Writing-original draft (equal). **Rasmus Møller Rosenbek Mortensen:** Investigation (supporting); Writing-review & editing (supporting). **Nikolaj Kaae Kirk:** Investigation (supporting); Writing-review & editing (supporting). **Ramses Gallegos-Monterrosa:** Methodology (supporting); Resources (supporting); Writing-review & editing (equal). **Ákos T. Kovács:** Conceptualization (equal); Funding acquisition (lead); Methodology (supporting); Project administration (lead); Resources (equal); Supervision (lead); Writing-original draft (equal).

ETHICS STATEMENT

None required.

DATA AVAILABILITY STATEMENT

Data are provided in the results section and appendices of this article.

ORCID

Mathilde Nordgaard <https://orcid.org/0000-0003-0648-8978>

Ramses Gallegos-Monterrosa <https://orcid.org/0000-0002-2366-0912>

Ákos T. Kovács <https://orcid.org/0000-0002-4465-1636>

REFERENCES

- Arnaouteli, S., Bamford, N., Stanley-Wall, N., & Kovács, Á. T. (2021). *Bacillus subtilis* biofilm formation and social interactions. *Nature Reviews Microbiology*. <https://doi.org/10.1038/s41579-021-00540-9>
- Asally, M., Kittisopikul, M., Rue, P., Du, Y., Hu, Z., Cagatay, T., Robinson, A. B., Lu, H., Garcia-Ojalvo, J., & Suel, G. M. (2012). Localized cell death focuses mechanical forces during 3D patterning in a biofilm. *Proceedings of the National Academy of Sciences of the United States of America*, 109, 18891–18896. <https://doi.org/10.1073/pnas.1212429109>
- Auchtung, J. M., Lee, C. A., & Grossman, A. D. (2006). Modulation of the ComA-dependent quorum response in *Bacillus subtilis* by multiple Rap proteins and Phr peptides. *Journal of Bacteriology*, 188, 5273–5285.
- Auchtung, J. M., Lee, C. A., Monson, R. E., Lehman, A. P., & Grossman, A. D. (2005). Regulation of a *Bacillus subtilis* mobile genetic element by intercellular signaling and the global DNA damage response. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 12554–12559.
- Bais, H. P., Fall, R., & Vivanco, J. M. (2004). Biocontrol of *Bacillus subtilis* against infection of *Arabidopsis* roots by *Pseudomonas syringae* is facilitated by biofilm formation and surfactin production. *Plant Physiology*, 134, 307–319.
- Beauregard, P. B., Chai, Y., Vlamakis, H., Losick, R., & Kolter, R. (2013). *Bacillus subtilis* biofilm induction by plant polysaccharides. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 1621–1630. <https://doi.org/10.1073/pnas.1218984110>
- Belas, R. (2013). When the swimming gets tough, the tough form a biofilm. *Molecular Microbiology*, 90, 1–5. <https://doi.org/10.1111/mmi.12354>
- Bischofs, I. B., Hug, J. A., Liu, A. W., Wolf, D. M., & Arkin, A. P. (2009). Complexity in bacterial cell-cell communication: quorum signal integration and subpopulation signaling in the *Bacillus subtilis* phosphorelay. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 6459–6464. <https://doi.org/10.1073/pnas.0810878106>
- Blake, C., Christensen, M. N., & Kovács, Á. T. (2021). Molecular aspects of plant growth promotion and protection by *Bacillus subtilis*. *Molecular Plant-Microbe Interactions*, 34, 15–25.
- Bongiorno, C., Ishikawa, S., Stephenson, S., Ogasawara, N., & Perego, M. (2005). Synergistic regulation of competence development in *Bacillus subtilis* by two Rap-Phr systems. *Journal of Bacteriology*, 187, 4353–4361.
- Branda, S. S., Chu, F., Kearns, D. B., Losick, R., & Kolter, R. (2006). A major protein component of the *Bacillus subtilis* biofilm matrix. *Molecular Microbiology*, 59, 1229–1238. <https://doi.org/10.1111/j.1365-2958.2005.05020.x>
- Branda, S. S., González-Pastor, J. E., Ben-Yehuda, S., Losick, R., & Kolter, R. (2001). Fruiting body formation by *Bacillus subtilis*. *Proceedings of*

- the National Academy of Sciences of the United States of America, 98, 11621–11626. <https://doi.org/10.1073/pnas.191384198>
- Cairns, L. S., Hobley, L., & Stanley-Wall, N. R. (2014). Biofilm formation by *Bacillus subtilis*: new insights into regulatory strategies and assembly mechanisms. *Molecular Microbiology*, 93, 587–598.
- Cairns, L. S., Marlow, V. L., Bissett, E., Ostrowski, A., & Stanley-Wall, N. R. (2013). A mechanical signal transmitted by the flagellum controls signalling in *Bacillus subtilis*. *Molecular Microbiology*, 90, 6–21.
- Chen, Y., Cao, S., Chai, Y., Clardy, J., Kolter, R., Guo, J. H., & Losick, R. (2012). A *Bacillus subtilis* sensor kinase involved in triggering biofilm formation on the roots of tomato plants. *Molecular Microbiology*, 85, 418–430. <https://doi.org/10.1111/j.1365-2958.2012.08109.x>
- Chen, Y., Yan, F., Chai, Y., Liu, H., Kolter, R., Losick, R., & Guo, J. H. (2013). Biocontrol of tomato wilt disease by *Bacillus subtilis* isolates from natural environments depends on conserved genes mediating biofilm formation. *Environmental Microbiology*, 15, 848–864.
- Comella, N., & Grossman, A. D. (2005). Conservation of genes and processes controlled by the quorum response in bacteria: characterization of genes controlled by the quorum-sensing transcription factor ComA in *Bacillus subtilis*. *Molecular Microbiology*, 57, 1159–1174. <https://doi.org/10.1111/j.1365-2958.2005.04749.x>
- Core, L., & Perego, M. (2003). TPR-mediated interaction of RapC with ComA inhibits response regulator-DNA binding for competence development in *Bacillus subtilis*. *Molecular Microbiology*, 49, 1509–1522. <https://doi.org/10.1046/j.1365-2958.2003.03659.x>
- Costerton, J. W., Lewandowski, Z., DeBeer, D., Caldwell, D., Korber, D., & James, G. (1994). Biofilms, the customized microniche. *Journal of Bacteriology*, 176, 2137–2142. <https://doi.org/10.1128/jb.176.8.2137-2142.1994>
- Di Pippo, F., Di Gregorio, L., Congestri, R., Tandoi, V., & Rossetti, S. (2018). Biofilm growth and control in cooling water industrial systems. *FEMS Microbiology Ecology*, 94, fiy044.
- Dragoš, A., Kieselwaller, H., Martin, M., Hsu, C.-Y., Hartmann, R., Wechsler, T., Eriksen, C., Brix, S., Drescher, K., Stanley-Wall, N., Kümmerli, R., & Kovács, Á. T. (2018). Division of labor during biofilm matrix production. *Current Biology*, 28, 1903–1913. <https://doi.org/10.1016/j.cub.2018.04.046>
- Even-Tov, E., Omer Bendori, S., Pollak, S., & Eldar, A. (2016). Transient duplication-dependent divergence and horizontal transfer underlie the evolutionary dynamics of bacterial cell-cell signaling. *PLoS Biology*, 14, e2000330. <https://doi.org/10.1371/journal.pbio.2000330>
- Fall, R., Kinsinger, R. F., & Wheeler, K. A. (2004). A simple method to isolate biofilm-forming *Bacillus subtilis* and related species from plant roots. *Systematic and Applied Microbiology*, 27, 372–379. <https://doi.org/10.1078/0723-2020-00267>
- Flemming, H. C., & Wingender, J. (2010). The biofilm matrix. *Nature Reviews Microbiology*, 8, 623–633. <https://doi.org/10.1038/nrmicr02415>
- Fujita, M., González-Pastor, J. E., & Losick, R. (2005). High- and low-threshold genes in the Spo0A regulon of *Bacillus subtilis*. *Journal of Bacteriology*, 187, 1357–1368.
- Gallegos-Monterrosa, R., Christensen, M. N., Barchewitz, T., Koppenhöfer, S., Priyadarshini, B., Bálint, B., Maróti, G., Kempen, P. J., Dragoš, A., & Kovács, Á. T. (2021). Impact of Rap-Phr system abundance on adaptation of *Bacillus subtilis*. *Commun Biol*, 4, 468. <https://doi.org/10.1038/s42003-021-01983-9>
- Gallegos-Monterrosa, R., Kankel, S., Götze, S., Barnett, R., Stallforth, P., & Kovács, Á. T. (2017). Lysinibacillus fusiformis M5 induces increased complexity in *Bacillus subtilis* 168 colony biofilms via hypoxanthine. *Journal of Bacteriology*, 199, e00204-17. <https://doi.org/10.1128/JB.00204-17>
- Gallegos-Monterrosa, R., Mhatre, E., & Kovács, Á. T. (2016). Specific *Bacillus subtilis* 168 variants form biofilms on nutrient-rich medium. *Microbiology*, 162, 1922–1932. <https://doi.org/10.1099/mic.0.000371>
- Hall-Stoodley, L., Costerton, J. W., & Stoodley, P. (2004). Bacterial biofilms: From the natural environment to infectious diseases. *Nature Reviews Microbiology*, 2, 95–108. <https://doi.org/10.1038/nrmicr0821>
- Hölscher, T., Bartels, B., Lin, Y.-C., Gallegos-Monterrosa, R., Price-Whelan, A., Kolter, R., Dietrich, L. E. P., & Kovács, Á. T. (2015). Motility, chemotaxis and aerotaxis contribute to competitiveness during bacterial pellicle biofilm development. *Journal of Molecular Biology*, 427, 3695–3708. <https://doi.org/10.1016/j.jmb.2015.06.014>
- Hölscher, T., Dragoš, A., Gallegos-Monterrosa, R., Martin, M., Mhatre, E., Richter, A., & Kovács, Á. T. (2016). Monitoring spatial segregation in surface colonizing microbial populations. *Journal of Visualized Experiments*, 116, e54752. <https://doi.org/10.3791/54752>
- Jarmer, H., Larsen, T. S., Krogh, A., Saxild, H. H., Brunak, S., & Knudsen, S. (2001). Sigma A recognition sites in the *Bacillus subtilis* genome. *Microbiology*, 147, 2417–2424. <https://doi.org/10.1099/00221287-147-9-2417>
- Jiang, M., Grau, R., & Perego, M. (2000). Differential processing of propeptide inhibitors of Rap phosphatases in *Bacillus subtilis*. *Journal of Bacteriology*, 182, 303–310.
- Jiang, M., Shao, W., Perego, M., & Hoch, J. A. (2000). Multiple histidine kinases regulate entry into stationary phase and sporulation in *Bacillus subtilis*. *Molecular Microbiology*, 38, 535–542. <https://doi.org/10.1046/j.1365-2958.2000.02148.x>
- Kearns, D. B., Chu, F., Branda, S. S., Kolter, R., & Losick, R. (2005). A master regulator for biofilm formation by *Bacillus subtilis*. *Molecular Microbiology*, 55, 739–749. <https://doi.org/10.1111/j.1365-2958.2004.04440.x>
- Kieselwaller, H. T., Lozano-Andrade, C. N., Wibowo, M., Strube, M. L., Maróti, G., Snyder, D., Jørgensen, T. S., Larsen, T. O., Cooper, V. S., Weber, T., & Kovács, Á. T. (2021). Genomic and chemical diversity of *Bacillus subtilis* secondary metabolites against plant pathogenic fungi. *mSystems* 6, e00770-20.
- Kolodkin-Gal, I., Elsholz, A. K. W., Muth, C., Girguis, P. R., Kolter, R., & Losick, R. (2013). Respiration control of multicellularity in *Bacillus subtilis* by a complex of the cytochrome chain with a membrane-embedded histidine kinase. *Genes & Development*, 27, 887–899. <https://doi.org/10.1101/gad.215244.113>
- Konkol, M. A., Blair, K. M., & Kearns, D. B. (2013). Plasmid-encoded ComI inhibits competence in the ancestral 3610 strain of *Bacillus subtilis*. *Journal of Bacteriology*, 195, 4085–4093. <https://doi.org/10.1128/JB.00696-13>
- Lazazzera, B. A., Kurtser, I. G., McQuade, R. S., & Grossman, A. D. (1999). An autoregulatory circuit affecting peptide signaling in *Bacillus subtilis*. *Journal of Bacteriology*, 181, 5193–5200.
- López, D., & Kolter, R. (2010). Extracellular signals that define distinct and coexisting cell fates in *Bacillus subtilis*. *FEMS Microbiology Reviews*, 34, 134–149.
- Lopez, D., Vlamakis, H., & Kolter, R. (2009). Generation of multiple cell types in *Bacillus subtilis*. *FEMS Microbiology Reviews*, 33, 152–163.
- López, D., Vlamakis, H., & Kolter, R. (2010). Biofilms. *Cold Spring Harbor Perspectives in Biology*, 2, a000398. <https://doi.org/10.1101/cshperspect.a000398>
- McLoon, A. L., Guttenplan, S. B., Kearns, D. B., Kolter, R., & Losick, R. (2011). Tracing the domestication of a biofilm-forming bacterium. *Journal of Bacteriology*, 193, 2027–2034. <https://doi.org/10.1128/JB.01542-10>
- McQuade, R. S., Comella, N., & Grossman, A. D. (2001). Control of a family of phosphatase regulatory genes (*phr*) by the alternate sigma factor sigma-H of *Bacillus subtilis*. *Journal of Bacteriology*, 183, 4905–4909.
- Mhatre, E., Monterrosa, R. G., & Kovács, Á. T. (2014). From environmental signals to regulators: modulation of biofilm development in Gram-positive bacteria. *Journal of Basic Microbiology*, 54, 616–632. <https://doi.org/10.1002/jobm.201400175>

- Mhatre, E., Sundaram, A., Hölscher, T., Mühlstädt, M., Bossert, J., & Kovács, Á. T. (2017). Presence of calcium lowers the expansion of *Bacillus subtilis* colony biofilms. *Microorganisms*, 5, 7. <https://doi.org/10.3390/microorganisms5010007>
- Mhatre, E., Troszok, A., Gallegos-Monterrosa, R., Hölscher, T., Kuipers, O. P., Lindst, S., & Kovács, Á. T. (2016). The impact of manganese on biofilm development of *Bacillus subtilis*. *Microbiology*, 162, 1468–1478. <https://doi.org/10.1099/mic.0.000320>
- Mueller, J. P., Bukusoglu, G., & Sonenshein, A. L. (1992). Transcriptional regulation of *Bacillus subtilis* glucose starvation-inducible genes: control of *gsiA* by the ComP-ComA signal transduction system. *Journal of Bacteriology*, 174, 4361–4373. <https://doi.org/10.1128/jb.174.13.4361-4373.1992>
- Ogura, M., & Fujita, Y. (2007). *Bacillus subtilis* RapD, a direct target of transcription repression by RghR, negatively regulates *srfA* expression. *FEMS Microbiology Letters*, 268, 73–80.
- Ogura, M., Yamaguchi, H., Yoshida, K., Fujita, Y., & Tanaka, T. (2001). DNA microarray analysis of *Bacillus subtilis* DegU, ComA and PhoP regulons: an approach to comprehensive analysis of *B. subtilis* two-component regulatory systems. *Nucleic Acids Research*, 29, 3804–3813.
- Omer Bendori, S., Pollak, S., Hizi, D., & Eldar, A. (2015). The RapP-PhrP quorum-sensing system of *Bacillus subtilis* strain NCIB3610 affects biofilm formation through multiple targets, due to an atypical signal-insensitive allele of RapP. *Journal of Bacteriology*, 197, 592–602.
- Ongena, M., & Jacques, P. (2008). *Bacillus* lipopeptides: versatile weapons for plant disease biocontrol. *Trends in Microbiology*, 16, 115–125. <https://doi.org/10.1016/j.tim.2007.12.009>
- O'Toole, G., Kaplan, H. B., & Kolter, R. (2000). Biofilm formation as microbial development. *Annual Review of Microbiology*, 54, 49–79. <https://doi.org/10.1146/annurev.micro.54.1.49>
- Parashar, V., Jeffrey, P. D., & Neiditch, M. B. (2013). Conformational change-induced repeat domain expansion regulates Rap phosphatase quorum-sensing signal receptors. *PLoS Biology*, 11, e1001512. <https://doi.org/10.1371/journal.pbio.1001512>
- Parashar, V., Konkol, M. A., Kearns, D. B., & Neiditch, M. B. (2013). A plasmid-encoded phosphatase regulates *Bacillus subtilis* biofilm architecture, sporulation, and genetic competence. *Journal of Bacteriology*, 195, 2437–2448. <https://doi.org/10.1128/JB.02030-12>
- Perego, M. (2013). Forty years in the making: understanding the molecular mechanism of peptide regulation in bacterial development. *PLoS Biology*, 11, e1001516. <https://doi.org/10.1371/journal.pbio.1001516>
- Perego, M., Glaser, P., & Hoch, J. A. (1996). Aspartyl-phosphate phosphatases deactivate the response regulator components of the sporulation signal transduction system in *Bacillus subtilis*. *Molecular Microbiology*, 19, 1151–1157. <https://doi.org/10.1111/j.1365-2958.1996.tb02460.x>
- Perego, M., Hanstein, C., Welsh, K. M., Djavakhishvili, T., Glaser, P., & Hoch, J. A. (1994). Multiple protein-aspartate phosphatases provide a mechanism for the integration of diverse signals in the control of development in *B. subtilis*. *Cell*, 79, 1047–1055. [https://doi.org/10.1016/0092-8674\(94\)90035-3](https://doi.org/10.1016/0092-8674(94)90035-3)
- Perego, M., & Hoch, J. A. (1996). Cell-cell communication regulates the effects of protein aspartate phosphatases on the phosphorelay controlling development in *Bacillus subtilis*. *Proceedings of the National Academy of Sciences of the United States of America*, 93, 1549–1553. <https://doi.org/10.1073/pnas.93.4.1549>
- Piggot, P. J., & Hilbert, D. W. (2004). Sporulation of *Bacillus subtilis*. *Current Opinion in Microbiology*, 7, 579–586. <https://doi.org/10.1016/j.mib.2004.10.001>
- Pottahil, M., & Lazazzera, B. A. (2003). The extracellular Phr peptide-Rap phosphatase signaling circuit of *Bacillus subtilis*. *Frontiers in Bioscience*, 8, 32–45. <https://doi.org/10.2741/913>
- Reizer, J., Reizer, A., Perego, M., & Saier, M. H. (1997). Characterization of a family of bacterial response regulator aspartyl-phosphate (RAP) phosphatases. *Microbial and Comparative Genomics*, 2, 103–111. <https://doi.org/10.1089/omi.1.1997.2.103>
- Serra, C. R., Earl, A. M., Barbosa, T. M., Kolter, R., & Henriques, A. O. (2014). Sporulation during growth in a gut isolate of *Bacillus subtilis*. *Journal of Bacteriology*, 196, 4184–4196. <https://doi.org/10.1128/JB.01993-14>
- Shemesh, M., & Chai, Y. (2013). A combination of glycerol and manganese promotes biofilm formation in *Bacillus subtilis* via histidine kinase KinD signaling. *Journal of Bacteriology*, 195, 2747–2754. <https://doi.org/10.1128/JB.00028-13>
- Singh, R., Paul, D., & Jain, R. K. (2006). Biofilms: implications in bioremediation. *Trends in Microbiology*, 14, 389–397. <https://doi.org/10.1016/j.tim.2006.07.001>
- Stewart, P. S. (2002). Mechanisms of antibiotic resistance in bacterial biofilms. *International Journal of Medical Microbiology*, 292, 107–113. <https://doi.org/10.1078/1438-4221-00196>
- Thérien, M., Kiewewalter, H. T., Auria, E., Charron-Lamoureux, V., Wibowo, M., Maróti, G., Kovács, Á. T., & Beaugerard, P. B. (2020). Surfactin production is not essential for pellicle and root-associated biofilm development of *Bacillus subtilis*. *Biofilm*, 2, 100021. <https://doi.org/10.1016/j.bioflm.2020.100021>
- Tzipilevich, E., & Benfey, P. N. (2021). Phage-resistant bacteria reveal a role for potassium in root colonization. *bioRxiv*. <https://doi.org/10.1101/2021.05.12.443821>
- van Gestel, J., Weissing, F. J., Kuipers, O. P., & Kovács, A. T. (2014). Density of founder cells affects spatial pattern formation and co-operation in *Bacillus subtilis* biofilms. *ISME Journal*, 8, 2069–2079. <https://doi.org/10.1038/ismej.2014.52>
- Verhamme, D. T., Kiley, T. B., & Stanley-Wall, N. R. (2007). DegU co-ordinates multicellular behaviour exhibited by *Bacillus subtilis*. *Molecular Microbiology*, 65, 554–568. <https://doi.org/10.1111/j.1365-2958.2007.05810.x>

How to cite this article: Nordgaard Christensen, M., Mortensen, R. M. R., Kirk, N. K., Gallegos-Monterrosa, R., & Kovács, Á. T. (2021). Deletion of Rap-Phr systems in *Bacillus subtilis* influences in vitro biofilm formation and plant root colonization. *MicrobiologyOpen*, 10, e1212. <https://doi.org/10.1002/mbo3.1212>

APPENDIX 1

FIGURE A1 Single *Δrap-phr* mutants are not majorly affected in cell granularity or size. Flow cytometry analysis showing the side scatter (SSC-A) vs forward scatter (FSC-A) plots of ungated WT and *Δrap-phr* mutant cells harboring the *PtapA-gfp* construct

FIGURE A2 Growth of *Bacillus subtilis* WT and single $\Delta rap-phr$ mutants in LB medium. WT and mutants were inoculated into 96-well plates with a starting OD₆₀₀ of 0.05. OD₅₉₀ was measured every 10 min for 48 h at 30°C, and each time point represents the mean of six technical replicates from two overnight cultures ($n = 6$). Error bars represent standard error (SE)

TABLE A1 Overview of phenotypes of the 12 single $\Delta rap-phr$ mutants compared with WT

Mutant	Known target(s) of respective Rap proteins	Matrix expr.	Colony	Pellicle	Root	Growth
<i>rapA-phrA</i>	Spo0F	↓	↑	↓	–	↓
<i>rapB</i>	Spo0F	↑	↑	–	–	–
<i>rapC-phrC</i>	ComA	↑	↓	↓	–	–
<i>rapD</i>	ComA	↑	–	–	↑	–
<i>rapE-phrE</i>	Spo0F	↑	–	–	–	–
<i>rapF-phrF</i>	ComA	↑	–	–	–	–
<i>rapG-phrG</i>	DegU	↑	–	–	–	–
<i>rapH-phrH</i>	Spo0F, ComA	↑	–	–	–	–
<i>rapI-phrI</i>	Spo0F, regulation of mobile genetic elements	↑	↑ (but small)	↓	↓	↓
<i>rapJ</i>	Spo0F	↑	↑	–	↑	–
<i>rapK-phrK</i>	ComA	↑	↑	–	–	–
<i>rapP-phrP</i>	Spo0F, ComA	↑	↑ (but small)	↓	↑	↑

Note: Up arrows indicate increased while down arrows indicate decreased features compared with WT. For colony and pellicle formation, the direction of the arrow is related to wrinkles and complexity.