The Value of Experimental Approaches in Migration Biology

Birnie-Gauvin, Kim; Lennox, Robert J.; Guglielmo, Christopher G.; Teffer, Amy K.; Crossin, Glenn T.; Norris, D. Ryan; Aarestrup, Kim; Cooke, Steven J.

Published in:
Physiological and Biochemical Zoology

Link to article, DOI:
[10.1086/708455](https://doi.org/10.1086/708455)

Publication date:
2020

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Birnie-Gauvin, K., Lennox, R. J., Guglielmo, C. G., Teffer, A. K., Crossin, G. T., Norris, D. R., Aarestrup, K., & Cooke, S. J. (2020). The Value of Experimental Approaches in Migration Biology. *Physiological and Biochemical Zoology*, 93(3), 210-226. <https://doi.org/10.1086/708455>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The value of experimental approaches in migration biology

In press in Physiological and Biochemical Zoology

Kim Birnie-Gauvin¹, Robert J. Lennox², Christopher G. Guglielmo³, Amy Teffer⁴, Glenn T. Crossin⁵, D. Ryan Norris⁶, Kim Aarestrup¹, Steven J. Cooke⁷

¹ Section for Freshwater Fisheries and Ecology, Technical University of Denmark, Denmark

² Laboratory of Freshwater Ecology and Inland Fisheries, NORCE Environment, Norway

³ Department of Biology, University of Western Ontario, Canada

⁴ Department of Forest and Conservation Sciences, University of British Columbia, Canada

⁵ Department of Biology, Dalhousie University, Canada

⁶ Department of Integrative Biology, University of Guelph, Canada

⁷ Fish Ecology and Conservation Physiology Laboratory, Department of Biology and Institute of Environmental and Interdisciplinary Science, Carleton University, Canada

Author for correspondence: K. Birnie-Gauvin

kbir@aqu.dtu.dk

Running title: Experimental migration biology

What is already known: Animal migrations are amongst the most incredible natural phenomena. In recent years, much effort has been dedicated to studying these behaviours, which, combined with recent advances in tracking technology, has greatly improved our understanding of when and where animals migrate.

What this study adds: Although there has been tremendous progress towards understanding the ultimate and proximate drivers of migration, there are still many unanswered questions. We submit that these questions could be explored by a more rigorous experimental approach to study migration biology. We review studies that exemplify and demonstrate *how* the field of migration biology, as well as the conservation of migratory animals, may benefit from new experimental work.

Abstract

The past several decades have ushered a golden age in the study of migration biology, leading to a wealth of descriptive papers that characterize various aspects of migration and its implications for individuals, populations, and ecosystems. However, relatively few studies have adopted an experimental approach to the study of migration, and fewer still have combined lab and field experiments to glean insights to the mechanisms underlying variation in migration behaviour and success. Understanding the proximate and ultimate causes of migration timing, energy allocation and optimization, migration success, and fitness is important to aid the conservation and management of wildlife populations by establishing appropriate protections or managing environmental conditions that influence migration. With recent technological advances and miniaturization of animal-borne electronic tracking devices, as well as ground-, water-, and space-based telemetry infrastructure, researchers have the tools necessary to experimentally test hypotheses central to the mechanics of migrations and individual variation therein. By pairing physiological measurements, molecular analyses, and other approaches within an experimental framework, there is the potential to understand not only how animal migrations function, but also what differentiates successful from failed migrations and the associated fitness implications. Experimental approaches to migration biology are particularly important, as they will help us to better comprehend, and hopefully predict, animal responses to environmental and anthropogenic change by isolating confounding variables that challenge inferences from observations.

Keywords: experiments, laboratory, migration, proximate causes, treatments, ultimate causes

Introduction

Animal migration is one of the great themes in organismal biology and ecology (Bowlin et al. 2010), and by understanding the proximate and ultimate drivers of migration behaviour, insights to many fundamental questions in biology can be gleaned (Mayr 1961; Schwenk et al. 2009). These include: the response of organisms to their environments; the functional diversity of migratory strategies and tactics; how physiological pathways underlie variation in migratory phenotypes; and how organisms cope with the trade-offs inherent to migration, e.g. between investment in locomotion versus reproduction. Migratory animals exhibit a high degree of taxonomic and behavioural diversity, ranging from short- to long-distance migrations, partial migrations, altitudinal migrations, and more (Bowlin et al. 2010). Animals must therefore integrate physiological, morphological, behavioural, and genetic aspects of their biology with environmental factors to which they are exposed (Åkesson and Hedenström 2007; Kunz et al. 2008). Migrations represent some of the most fascinating and awe-inspiring phenomena observed in the natural world.

Not surprisingly, some iconic migrations (think salmon, wildebeest, terns, sea turtles, and even some land crabs) have also captured the imagination of the public with them being routinely featured in documentaries and even popular movies (e.g., *Turtle Tale*, *The Lion King*). Migration is also a sufficiently important and interesting topic that it is covered (either as a chapter or through case studies) in almost every ecology, behaviour, and animal physiology textbook from past (e.g., the first major text on ecology; Allee et al. 1949) to present (e.g., the classic Eckert Animal Physiology series and Alcock Animal Behaviour series, among many others), and now even has a dedicated journal (i.e., *Movement Ecology*).

Migration biology reveals when, where, why, and how animals undertake their often synchronized, cyclic, and predictable movements across land- and sea-scapes (Alerstam et al. 2003; Wilcove 2008; Bowlin et al. 2010). Answering these questions can inform both the management and conservation of migratory species and their habitats by allowing researchers to predict how animals respond to disturbances (Newton 2008; Wilcove and Wikelski 2008; Altizer et al. 2011). From field observations of migrations at varying spatial and temporal scales, and among a diversity of species, the general nature of organismal biology and animal migration has been fairly well described. Data from such observations allow us to build predictive models, but these have limited power to explain the proximate and ultimate mechanisms underlying migration. Experimental approaches to the study of migration, in which specific variables can be controlled or manipulated in the lab or field, are therefore needed to isolate causative factors differentiating migrants from non-migrants and, from a fitness perspective, successful from unsuccessful migrations.

We contend that experimental approaches are needed if we are to provide answers to the key questions about migration biology, especially the “how’s” (mechanistically) and “why’s” (evolutionarily). These approaches must integrate multiple scientific disciplines, and include both traditional and novel techniques, in studies across multiple taxonomic groups that span laboratory and field-based settings (Bowlin et al. 2010). A search of the Web of Science Core Collection Database revealed that within the field of ‘zoology’, 6850 migration studies (search: Topic = Migrat* NOT Topic = cell*, to avoid false positives related to cell migration) were conducted between 1900 and 2019 (as of 8 April 2019). Within those, experimental studies (search: Topic = experiment* OR manipul* OR treat*) comprised just 13% (855 hits) of the studies, suggesting that experimental approaches within the realm of migration biology are

101 somewhat uncommon. Manual screening of a subset ($N = 20$) by abstract revealed that false
102 positives for experimental studies were common (7/20) such that this number of 13% is inflated.
103 Although we acknowledge that this search is not fully comprehensive, it supports the perspective
104 that experimental approaches, where either the organisms or environmental contexts are
105 manipulated, are relatively uncommon in migration studies.

106 Kennedy and others (Kennedy and Booth 1963, 1964; Dingle 1965, 1966) are credited
107 with some of the earliest research concerning the proximate drivers of migration biology.
108 Experiments with insects revealed physiological pathways stimulating the onset of migration and
109 the determinants of migration as a distinct behaviour. Such experiments have contributed to a
110 definition of migration to differentiate it from other movement behaviours observed in animals
111 (Dingle 2006; Dingle and Drake 2007). In this review, we draw representative examples from the
112 literature to highlight the potential that experimental approaches to migration biology have for
113 revealing the many drivers of animal migrations (Figure 1). Whether under controlled laboratory
114 conditions or in field contexts, experimental studies are useful for determining the proximate
115 drivers, or causes, of migratory behaviours (Underwood 1990; Cooke et al. 2017). Here, we
116 advocate for experimental approaches to migration biology, and explore several thematic areas
117 of migration biology research that should be testable with experimental methods. We then
118 discuss the aspects of migration where experiments are currently lacking, but which would
119 greatly benefit our understanding of how, why, when, and where animals migrate.

121 **What can experimental biology bring to the study of migration?**

122 Experimental studies can be conducted either in laboratory or field settings to reveal the
123 underpinnings of various aspects of migration. In both settings, environmental (e.g., temperature,

oxygen availability), behavioural (e.g., predation, competition), or physiological (e.g., hormones, energetic substrates, disease) variables can be isolated by physical interventions. Experiments related to interspecific interactions are more challenging to perform but are highly relevant to explore given that species or conspecific interactions may be an important force driving the evolution of migratory behaviours (Furey et al. 2018).

Field experiments are useful for generating in situ data related to the migration of animals at ecologically-relevant scales which are not easily simulated in a laboratory (though we acknowledge that some mesocosm experiments can simulate nature fairly accurately, though mostly on a small scale). Frequently, these field studies require some method of remote observation to track the spatial and temporal progress of animals in the migratory arena. Methods include passive techniques like photo identification and mark-re-sight/recapture, both of which rely on the identification of uniquely marked individuals. For example, photo identification techniques were used to track the movement patterns of individual white sharks and revealed site fidelity in at least 78% of individuals around Guadalupe Island, Mexico (*Carcharodon carcharias*; Domeier and Nasby-Lucas 2007). Active techniques include various forms of electronic tracking, like biotelemetry or biologging, and can include satellite, radio, or acoustic transmitters that communicate position (and other) data to receivers or log it onboard for later retrieval. These active methods provide accurate information about individual migratory progress, which can be used to define discrete experimental treatment groups (Cooke et al. 2008).

However, not all questions about migration biology can be addressed using field studies. For example, when addressing uncertainty about the potential migration behaviour of genetically engineered animals in nature, contained laboratory or mesocosm experiments must be used (e.g.

(Sundström et al. 2010) as the release such animals to the wild is legally prohibited by most countries (Devlin et al. 2006; Kapuscinski et al. 2008). In cases when animals are too small or too sensitive to be tagged, laboratory studies have proven quite useful. Furthermore, because most field studies require some form of tagging to identify individuals, there is no possibility for true (untagged) migratory controls. Laboratory approaches also allow more direct observation and control over the environment to which animals are exposed. This is especially important as behavioural observations can be lost during field studies when electronic tags are shed or if they alter behaviour in unexpected ways (Hazekamp et al. 2010). In such cases, preliminary studies in the lab may help to guide field approaches.

One powerful, yet seldom used, way to address the inherent limitations of both laboratory and field studies is to conduct both types of studies in parallel or in succession to address the same question. Proof of concepts developed in a laboratory, which provide evidence that manipulations can effectively isolate variables that influence animal behaviour, can then be scaled to field settings where behavioural manipulations can be tested in a natural context. Hellström et al. (2016) performed such an experiment to study the effects of a GABA_A receptor agonist on Atlantic salmon (*Salmo salar*) smolt migration, revealing that the agonist increased migration intensity of smolts first in a circular pool mesocosm, then in the Ume River in Sweden. These combined approaches are powerful for providing data at both relevant scales to isolate mechanisms and reveal applications, and can make significant contributions to our understanding of animal migration.

Understanding the natural limits and regulations imposed on migratory populations is of crucial importance to adequately conserve and manage migratory species. Estimates of density-dependence are especially relevant to develop predictive population models (Sutherland and

Norris 2002; Norris and Taylor 2006; Taylor and Norris 2010). This is particularly salient for partially migratory species that are sensitive to density-dependence and for which the expression of migration may depend on conspecifics (Griswold et al. 2010; Chapman et al. 2011). These estimates can be obtained from time-series studies, but lacks proper controls. As such, experiments are key to advance this area of study (Flockhart et al. 2012), and can help us understand the mechanisms underpinning various aspects of migration (Figure 1). In the following sections, we highlight key experimental approaches that have provided valuable insights for understanding various aspects of migration. We acknowledge that these sections are not exhaustive, but provide clear examples of the use of experiments in migration biology in various taxa, and highlight their utility for elucidating migration processes.

Abiotic challenges experiments

There are a large number of environmental cues such as temperature, photoperiod, river discharge, and lunar phase, as well as endogenous cycles, that can influence migratory activity of animals via effects on endocrine signalling (Bradshaw and Holzapfel 2008; Lennox et al. 2018). Experimental manipulation of the environment by controlling some of these abiotic factors has yielded insights into how these processes progress or change (e.g. Gwinner 2003; Farrell 2009; Müller et al. 2018, Table 1). By manipulating the release date of red deer (*Cervus elaphus*) in Germany, Rivrud et al. (2016) showed that late-released individuals could compensate for having missed the peak plant maturity by migrating faster and to higher elevations than early-released individuals. These compensations are likely to involve extra energetic costs, demonstrating the importance of adequate environmental conditions during migration, and highlight the consequences that a mismatch between these conditions and migration timing could lead to.

Environmental conditions can negatively impact the capacity for migratory animals to reach their breeding or non-breeding destinations. Energetic constraints imposed on migration by the environment can disrupt the timing and success of migrations. Experimental studies of the metabolic capabilities of migrants and the fuel conversion efficiencies can be applied to estimate challenges to migration. Clark et al. (2011) compared the cardiorespiratory capabilities of Pacific salmon by manipulating temperatures in laboratory trials of swimming in respirometry chambers and revealed that maintenance of aerobic scope of pink salmon (*Oncorhynchus gorbuscha*) at high temperatures would confer an advantage over thermally-sensitive congeneric Pacific migrants. In this way, experimental manipulations can be useful in understanding how changing migration conditions will impact future reproductive output, survival, and the evolution of fitness-related traits.

Rearing organisms under various environmental conditions may also help to understand the underlying mechanisms that allow organisms to migrate. For example, Lague et al. (2016) reared bar-headed geese (*Anser indicus*) – some of the highest flying migratory birds – at high altitude (3200m) in the Himalayas and at sea-level and found that those reared at high altitude had lower metabolic rates and faster ventilation rates in hypoxic conditions compared to those reared at sea-level. These findings suggest that high altitude rearing enabled better capacity for high elevation flight due to a reduced oxygen demand during resting and an increased oxygen uptake and delivery during exercise. This adaptation is specifically what allows bar-headed geese to migrate through the Himalayas. It is also possible to use the natural variability or seasonality of environments to test how they shape and affect movements in animals. In one study, Conenna et al. (2019) tracked yellow-winged bats (*Lavia frons*) in dry and rainy seasons and found that home ranges were larger during the dry season, when resources are scarce. These findings may

suggest the presence of a coping mechanism, where bats cover larger areas and spend more time actively foraging to find food. These types of experiment not only help us understand the drivers of migration, but may also enable us to predict how these movements will change under various climate scenarios.

Biotic challenges experiments

Disease and parasitism have the potential to negatively affect the expression or success of migration. As animals move through the environment, their metabolism can be influenced by exposure to pathogens as well as latent infections (Altizer et al. 2011). Experimental manipulations have identified several pathways through which infections can alter the physiology and movement of migrant individuals (Table 2). Bradley and Altizer (2005) experimentally infected monarch butterflies (*Danaus plexippus*) with a protozoan parasite (*Ophryocystis elektroscirrha*), which impaired flying capabilities and reduced the capacity for these insects to migrate. Sea louse (*Lepeophtheirus salmonis*) ectoparasitism of Atlantic salmon (*Salmo salar*) has similarly been demonstrated to decrease host swimming capabilities and cardiac performance compared to controls (Wagner et al. 2003). In songbirds, lipopolysaccharide injections used to simulate bacterial infection were shown by automated radio tracking to alter movements and prolong migratory stopovers (Hegemann et al. 2018). These experimental manipulations provide support for a role of infections in modulating the capacity of animals to migrate, which can alter host-pathogen co-evolution dynamics (Nuismer et al. 1999). As our ability to characterize infections among wild animals develops through the novel application of molecular tools (e.g. eDNA, gene sequencing), our understanding of how multiple infections and host responses interact to impact migration biology is growing rapidly (Satterfield et al. 2018).

Infectious agents have been shown to impact animals at multiple life stages, with effects relevant to the migrations occurring therein. For example, the long-distance migrations of Pacific salmon (*Oncorhynchus spp.*) can be physiologically and metabolically challenging but necessary for growth (e.g. juvenile migration to sea) and reproduction (adult spawning migration; Groot and Margolis 1991). Among juvenile sockeye salmon (*O. nerka*), viral response indices were associated with reduced migration success from freshwater rearing lakes to marine feeding grounds (Jeffries et al. 2014); viral infection (infectious haematopoietic necrosis virus) has also been linked to predation susceptibility of juvenile sockeye in freshwater (Furey 2016) and marine habitats (Miller et al. 2014). Among adult sockeye, severe parasitic kidney infections (*Parvicapsula minibicornis*) were associated with reduced exercise recovery and swim speeds compared to controls, but similar initial swim performance (Wagner et al. 2005). The conditions of the migration (e.g. water flow) may therefore regulate the role of infections on migration success. Collectively, these studies have identified behavioural associations that warrant further experimentation and also demonstrate strong regulatory roles of the environment (e.g. stressors) and host biology (e.g. species, population, sex) on the degree to which infections influence survival and migration success.

Pathogens and infectious agents are pervasive in nature, and can have deleterious effects on animal performance and migration biology (Dhabar and McEwen 1997). Warming temperatures caused by climate change are projected to increase the virulence of infectious agents by increasing rates of replication and development, as well as suppressing the immune resistance of hosts (Altizer et al. 2013). This is especially true among ectotherms because body temperature and metabolic rate are determined by environmental temperatures, which in a warming climate can lead to accelerated pathological processes (Altizer et al. 2013). In one

example, Crossin et al. (2008) intercepted homing adult sockeye salmon during their spawning migration in the Fraser River (Canada) and held them in experimental pools at different temperatures for several weeks. Sockeye were then released back into the river to complete migration. Via acoustic telemetry, the study revealed that fish held at ecologically realistic high temperatures (mimicking recent high temperatures in the river) had reduced migration success and enhanced *P. minibicornis* infections, highlighting the interacting effects of thermal stress and disease processes. Given that many animal populations are locally adapted to historic thermal conditions encountered at each life stage, including crucial migration periods (Eliason et al. 2011; Hoffman et al. 2012), temperature perturbations have the potential to alter host-pathogen relationships during migration. Our knowledge of host defences against novel pathogen exposure during migration is still in its infancy, but understanding these relationships are important for conservation efforts that aim to improve landscape connectivity and enhance migratory (and subsequently, transmission) potential (Satterfield et al. 2018). Experiments that define the mechanisms of host-pathogen dynamics and migration behaviour in the context of climate change and environmental perturbation are urgently needed (Miller et al. 2014).

Bioenergetic and biomechanical experiments

Migration allows individuals to exploit fluctuating productivity among habitats (Alerstam et al. 2003; Buehler and Piersma 2008). However, there are substantial costs associated with migration that must be balanced against the benefits conferred by these long-distance movements. Experimental approaches to measuring the energy stores of migrants and the energetic costs of activity provide insight into the expression of migration (Table 3). Rates of energy expenditure can be measured by doubly-labelled water (Wikelski et al. 2003), wind tunnels in birds (e.g.

Klaassen et al. 2000), or respirometry in small gill-breathing species (e.g. Lee et al. 2003; Clark et al. 2011). Experimental intervention to manipulate aerobic capabilities of animals have seldom been performed, but such studies could be important to evaluate how animals respond to energetic challenges prior to or during migration. One example of experimentally manipulating energetic effort on migration success and timing has been done via wing clipping or handicapping in migratory birds. Elliott et al. (2014) attached “handicaps” using tape to the lower back of thick-billed murres (*Uria lomvia*) to evaluate energy expenditure during a period of high energy demand (chick-rearing), and found that energy expenditure did not differ between handicapped and non-handicapped individuals due to behavioural adjustments, but this finding was dependent on age. In another experiment, Cohen et al. (2012) manipulated the energetic condition of migrating red-eyed vireos (*Vireo olivaceus*) and then radio-tracked them to elucidate the importance of body condition on migration. Upon release, migrating vireos in lower condition moved faster and further than those of higher condition, perhaps as a means of accessing better or more food sources to replenishing energetic fuels necessary to continue migration.

Experimental manipulation of feeding or food availability provides insight into energetic constraints on migration. Whitethroats (*Sylvia communis*) attracted to feeding sites weighed significantly more and showed higher rates of fuel deposition and energy stores just prior to migration than control individuals (Fransson 1998), suggesting that mass, fuel deposition, and fuel load were adjusted according to food availability in preparation for migration. Starvation of large milkweed bugs (*Oncopeltus fasciatus*) prior to migration influenced levels of juvenile hormone and the expression of migration-associated flight behaviour (Rankin and Riddiford 1977). By manipulating the fatty acid composition of the diet of red-eyed vireos to test whether

fatty acid composition enhances exercise performance, Pierce et al. (2005) found that the energetic cost of migration in birds is likely to be modulated by the fatty acid composition of the diet (see review in McWilliams et al. 2004). In brown trout (*Salmo trutta*), short-term food deprivation resulted in decreased expression of migration compared to control animals (Midwood et al. 2016). Food-deprived trout that did migrate were delayed by approximately one week, suggesting that food availability and energetics are involved in dictating whether an individual migrates or not in this partially migratory species.

Migration timing is essential for animals to reach their destinations at optimal times. The green wave hypothesis posits that terrestrial migrants ‘surf’ a wave-like dynamic of progressively budding plants to match their northward migration to appropriate weather and forage possibilities (Van der Graaf et al. 2006; Rivrud et al. 2016). However, climate change has led to measurably shorter migrations in birds (Visser et al. 2009), and has generated phenological mismatches at foraging stopover sites (Kellerman and van Riper 2015). What effects this change will have on population dynamics may dictate whether a population persists or grows as a result of climate change (Reed et al. 2013). An artificial selection experiment suggests that residency will likely evolve rapidly in populations of migratory birds if selection for shorter migrations persists (Pulido and Berthold 2010). Similarly, the green wave hypothesis (and other ‘surfing’ hypotheses) is supported by ungulate migrations in savannahs, where animals attempt to balance the seasonality of rainfall and soil fertility gradients. Even shorter migrations, like that of elephants in the Marsabit protected area of Kenya support this hypothesis (Bohrer et al. 2014). Given the occurrence of climate change and other human-induced changes such as habitat alteration, experimental approaches are particularly needed to investigate how animals respond

to changes and what consequences can be expected for migrants working to make it to the right place at the right time.

Endocrine and signalling pathways experiments

Migration is regulated by endocrine signals that respond to exogenous cues and stimulate or suppress the migratory state. Seasonal changes to hormone levels or sensitivity are responsible for timing and synchronizing migration with the environment and food sources. Direct alteration of circulating hormone concentrations by injections or implants, or blocking receptors to attenuate effects of putative hormones, are useful approaches to investigate the roles and relative importance of hormones or neurotransmitters (see Sopinka et al. 2014; e.g. Hedenstrom and Lindstrom 2017; Minter et al. 2018, Table 4).

Glucocorticoid manipulation is a classic experimental approach used to reveal mechanisms of migration behaviour. Glucocorticoids are metabolic regulators involved in energy mobilization, as well as components of the stress response (Hau et al. 2016) and thus can yield information on the homeostatic status of an individual (Sopinka et al. 2015). Glucocorticoid hormone concentrations can be manipulated indirectly (e.g. induced reproductive failure; Ramos et al. 2018) or directly via implantation or injection. Glucocorticoid effects can also be modified by blocking of the receptors (Sopinka et al. 2014). Downstream effects differ for acute and chronic stressors with chronic stressors yielding tertiary responses such as compromised growth, immunity, reproduction, or migratory behaviour, which can be observed in animals that have experimentally altered stress states.

A variety of other hormones have been implicated in migratory behaviour. In birds, experiments have demonstrated that gonadotropins are crucial for eliciting vernal migratory

restlessness, although other hormones, such as thyroid hormones, appear to be more important in autumn (Ramenofsky and Wingfield 2007). Gonadotropin-releasing hormone (GnRH) may be manipulated by implantation, and Sato et al. (1997) and Kitahashi et al. (1998) observed more rapid homing by treated sockeye (*Oncorhynchus nerka*); however, Crossin et al. (2009) observed no differences in migration of GnRH treated sockeye compared to controls (though testosterone was related to migration timing in males). In amphibians, thyroxin treatment leads to a preference for terrestrial environments even during the aquatic phase of their lifecycle (e.g., Tassava and Kuenzli 1979) and juvenile hormone stimulates migratory behaviour of insects (Rankin 1978). In fish and birds, melatonin secreted in response to photoperiodic changes exerts influence over the expression of migratory restlessness and the onset of migration at stopover sites (Bradshaw and Holzapfel 2007). Experimental manipulation of melatonin levels via manipulations of light cycles led to the expression of migratory restlessness in bramblings (*Fringilla montifringilla*) during “cool light” to “warm light” cycles but not during blue light to red light cycles (Phol 2000). The findings suggest that melatonin cycles are involved in regulating migratory restlessness by granting or inhibiting nocturnal activity (Phol 2000). Garden warblers (*Sylvia borin*) injected with ghrelin (a hormone regulating physiological and behavioural aspects of foraging) exhibited diminished food intake and increased migratory restlessness (Goymann et al. 2017). Food intake must therefore regulate at least some of the hormones that control migratory behaviour, an inference made possible by an experimental approach to investigate migration. There are likely still more hormonal pathways contributing to migration of different species (and taxa beyond fish and birds) that can be investigated by applying these experimental approaches.

Sensory, navigation and orientation experiments

There is a long history of experimental laboratory and field studies directed at understanding how migrating animals orient and navigate (e.g., Åkesson 2003). Animals orient using a variety of means from visible landmarks to cues associated with the sun, stars, and Earth's magnetic field (Åkesson and Hedenström 2007). Isolating the critical components of compasses is challenging and requires experimentation (Table 5). Compasses that depend on decoding geophysical features of the environment such as magnetic field polarity, inclination, or strength, or skylight polarization, require sensory organs and neural processing of the information to be accurate. Evolutionary analysis is needed to establish the origin of such adaptations, but evidently there is strong selection towards accuracy that has permitted the evolution of finely tuned sensory organs for orientation and navigation. Characterizing and understanding them remains a major challenge.

Sensory manipulations in the field have contributed to our understanding of animal navigation. Sensory manipulations may be applied by direct intervention of the animal's senses or by altering the individual's environment and monitoring its integration of the stimuli. By depriving juvenile loggerhead (*Caretta caretta*) and green turtles (*Chelonia mydas*) of visual cues under laboratory conditions, and subsequently displacing them between 30 to 167 km away from their site of capture, Avens and Lohmann (2004) found that turtles oriented toward feeding areas between May and September, but toward their migration route during October and November. The findings suggest that turtles can assess their position relative to their ultimate destination using local cues, and as such, can navigate using a sensory 'map'. Ablation of vision, smell, and magnetoreception have been used to isolate senses and compare the progress of fish towards their destination to reveal the phases of homing and the sensory foundation that these

animals rely upon for wayfinding (Barbin 1998; Ueda et al. 1998; Mitamura et al. 2005). In another example, Kishkinev et al. (2016) surgically ablated either the magnetic or olfactory sensory systems of experienced (individuals that have migrated successfully at least once) white-throated sparrows (*Zonotrichia albicollis*) in Ontario and radio-tracked them after the birds were displaced 2200 km away in Saskatchewan. Both the ablated and non-ablated individuals demonstrated an inability to reorient once translocated to an unknown area, suggesting that this species does not display compensatory behaviour required for true navigation. In an experiment altering the sensed environment of butterflies, monarch butterflies (*Danaus plexippus*) were exposed to simulated magnetic fields: under control conditions, the insects oriented southwest (typical for migrants); under a reversed magnetic field, individuals oriented in the opposite direction (i.e., northeast); lastly, under an amagnetic condition, monarchs showed no orientation at all (Etheredge et al. 1999). Similarly, in birds, experimental cue conflict and surgical manipulations followed by radio tracking have been used to study compass orientation and navigation mechanisms (e.g. Holland et al. 2009, Cochran et al. 2004, Kishkinev et al. 2016). These experiments have been powerful in exhibiting the importance of orientation senses as investigators work towards understanding how animals migrate over large distances accurately.

Translocation experiments

Rather than simulating an environment for migrants, they can be relocated into a new venue where their responses to a novel environment can be monitored. Experimental displacement of wild animals is a widespread approach to study navigation and orientation mechanisms (Table 6). One of the first experimental displacement studies ever performed was on green sea turtle hatchlings, in which the newly emerged turtles were translocated from the east coast to the west

coast of Central America (Carr and Ogren 1960). This was the first indication that hatchlings likely use lighting cues rather than an innate preference for direction to migrate to the ocean (i.e., their sea finding phase). Experimental displacements of adults between islands have highlighted that local cues are important in the final phase of homing during breeding (reviewed in Lohmann et al. 2008). In migrating manx shearwaters (*Puffinus puffinus*), clock shifting (where animals are exposed to artificial photoperiods in captivity) coupled with experimental displacement also revealed the importance of the sun as a navigation tool (Padget et al. 2018). In Scopoli's shearwaters (*Calonectris diomedea*), displacement and sensory ablation (magnetically-disturbed and anosmic groups) revealed the importance of olfaction and visual integration of topography to navigation (Pollonara et al. 2015). These Mediterranean seabirds appeared to use olfactory cues over open water, but then relied on topographical cues from the coastline to orient neritic zones. In another study, Skov et al. (2010) investigated the influence of environmental cues, migration history, and habitat familiarity on the roach (*Rutilus rutilus*), a partially migrant fish species. Roach from populations with different migratory routes to lakes were translocated into unfamiliar habitats. The study found that translocated fish abandoned their previous migration tactic (i.e., partial migration), and adopted similar migration patterns to local fish, suggesting that partial migration is a phenotypically plastic trait triggered by local environmental cues.

Translocation experiments are not only useful for understanding the mechanisms used to orient and navigate, but they can be used to understand energy allocation processes and migratory costs. In a common-garden rearing experiment, Kinnison et al. (2003) raised two populations of chinook salmon (*Oncorhynchus tshawytscha*), from two different river systems (i.e., spawning areas 17 km inland and 17 m elevation vs. 100 km and 430 m elevation), and then experimentally released the adults into the non-natal river. By so doing, each locally adapted

population experienced a new spawning migration, and upon completion differential energy allocations and costs were measured in spawning adults. The study showed that males exhibited different reproductive morphologies as a result of the new migration costs they experienced, which indicated reallocation of limited energy to a new locally adapted form: when displaced from the short distance migration to the longer migration had smaller dorsal humps and kypes, with reduced tissue energy reserves, suggesting that longer migrations reduce available energy for spawning competition. Similarly, female chinook salmon were found to have smaller ovarian mass translocated from the short to longer migration (Kinnison et al. 2001). These findings highlight the importance of migration to the evolution of locally adapted forms in species operating under a fixed energy budget (capital breeding).

Conservation-related experiments

Migrating animals are exposed to a gamut of stressors, including naturally-occurring ones as well as anthropogenic challenges (Boonstra 2013), which can include barriers to migration, fisheries interactions, noise disturbance, light pollution, and exposure to contaminants, among others. Understanding how anthropogenic activities are affecting migrations is a pressing issue for the conservation and management of many migratory species (Martin et al. 2007). By experimentally exposing migrant animals to relevant single and/or cumulative stressors, the mechanisms underlying individual variation in behaviour, success, and survival can be investigated. In some instances, the effects of anthropogenic activities or infrastructure on migrations can be investigated without experimental intervention per se, but rather as a result of the occurrence of anthropogenic stressors in nature (Table 7). One such example, by Gula et al. (2009), showed that wolves (*Canis lupus*) were able to migrate to find mating partners through

densely populated areas with high densities of roads. Although this example does not provide insight into the mechanisms that underlie migration, it suggests that wolves may face multiple non-natural threats during their migration, which may in turn affect their success in reproduction. Similarly in freshwater fish, dams act as barriers to migration. Using existing dam infrastructure, much can be learned on the costs of overcoming these human-made structures in a semi-experimental approach. A study using 18,286 radio-tagged adult chinook salmon and steelhead trout (*O. mykiss*) that had to migrate passed eight Columbia and Snake river dams to reach spawning areas found that unsuccessful migrants had longer passage times at nearly all the dams (Caudill et al. 2007). These findings may suggest the presence of delayed negative effects or condition-dependent effects of dam passage which may vary across individuals, and highlight the need for a mechanistic understanding of the factors that influence migration success.

Contaminants and other forms of pollution are ubiquitous in nature, even in the most remote locations on the planet. Even at sub-lethal levels of exposure, environmental contaminants may pose threats to migratory animals. For example, an experimental lab study showed that methylmercury impaired flight performance of migratory yellow-rumped warblers (*Setophaga coronata*) in a wind tunnel (Ma et al. 2018), and a subsequent dosing and field release showed that mercury exposure altered activity behaviour of warblers at a migratory stopover site causing them to depart sooner than controls (Seewagen et al. 2019). Similarly, exposure to neonicotinoid and organophosphate pesticides disrupted migratory orientation, and neonicotinoids also reduced body mass in short-term captive white-crowned sparrows (*Zonotrichia leucophrys*; Eng et al. 2017) and a field manipulation and radio tracking study showed that neonicotinoid insecticides suppress fuelling, reduce body mass, and delay migration in free-living sparrows (Eng et al. 2019). In western sandpipers (*Calidris mauri*), experimental

oiling of feathers reduced escape flight performance and increased endurance flight energy costs (Maggini et al. 2017a, b), effects that were predicted to impair migration ability. If sub-lethal effects of contaminants at environmentally relevant exposures are demonstrated in the laboratory setting, then experimental field studies should be used to determine fitness consequences for migrants under natural conditions. Given the increasing number of threats that migratory organisms face in the ‘Anthropocene’ (Wilcove 2008), there are growing opportunities to use experimental approaches to understand the consequences of threats and identify and test potential strategies for mitigating them (Lennox et al. 2016).

Seasonal carryover effects, the need for experimental approaches to address conservation of migrants, and prospects for the future

As individuals migrate, they may carry with them the experiences that occurred during previous periods of the annual cycle (Harrison et al. 2011; O’Connor et al. 2014). These may affect future survival and alter migration route and timing, as well as population dynamics (Norris and Marra 2007). Much of these so-called carryover effects have been ascribed to habitat and diet (Norris, 2005; Conklin and Battley 2012), both of which can be significantly impacted by humans (Birnie-Gauvin et al. 2016), but there is evidence that other factors may impose delayed consequences on migration (Midwood et al. 2014; Peiman et al. 2017). For example, how does variation in population density (i.e., number of individuals within a given area) in one season affect recruitment and reproduction rate the following season (Betini et al. 2013)? How do inter- and intraspecific interactions alter migratory behaviours? How does habitat loss affect habitat selection and migratory routes in subsequent periods? How do stressors influence variation in reproductive investment, and thus breeding decisions, breeding success, and population

processes (O'Connor et al. 2014)? How do stressors affect the number of migrant individuals from a partially migratory species? These are only a few of the questions that future research should aim to address, but we argue that experimental studies provide an important approach for doing so. Comparing migratory and non-migratory species in an experimental set up could provide one way to investigate some of these questions.

At present, anthropogenic structures and stressors are abundant and widespread. Experiments to understand applied issues and their potential impacts on animal migrations will help conserve and manage species by answering key questions, such as: How do dams affect migratory fish species? How do wind turbines affect birds and bats during migration? How do landscape characteristics, including urbanization and agriculture, affect migration patterns and success? Whereas the immediate and direct impacts of these human-originated issues have been studied (e.g., Jepsen et al. 1998; Norberg et al. 2002; Donaldson et al. 2011; Keefer et al. 2011; Silva et al. 2012; Minderman et al. 2012; Spoelstra et al. 2015), their carryover effects on future aspects of individual success have seldom been the focus of research. For example, wind turbines may affect bird feeding outside of their migratory period, which could influence migration success by limiting the energy storage and stress status of the animal. These long-term effects must emerge as a focal point of studies to develop a more holistic, lifetime scale perspective on animal migration.

Conclusions

Given the importance of migratory species to biodiversity, ecosystem function, and human culture and economies, an understanding of where, when, how and why animals migrate is critical for their conservation and for the sustainability of the ecosystem services they provide.

Observational studies, where animals are not manipulated, will continue to play a fundamental role in expanding our knowledge of animal migrations (e.g., identifying pathways and timing patterns), and the effects of various extrinsic and intrinsic factors on individuals and populations. Experimental approaches, either with animals in the laboratory, field, or in combination, provide an avenue through which many crucial observations can be tested to establish mechanism and cause. Indeed, experiments provide strong inference and may in some cases be the only way to definitively test hypotheses. Experimental approaches are not without limitations. For example, it is much easier to manipulate large numbers of insects and fish than birds and mammals. There is a noticeable dearth of experimental studies on mammals (likely for ethical and logistical reasons) and herptiles. Many long-distance migrants are also small (especially insects), making tracking with telemetry devices for long durations and distances a challenge (Wikelski et al. 2007). However, technological improvements and miniaturization of electronics, sensors, computer memory, and batteries, particularly driven by innovations for portable communications, personal computing, and gaming are rapidly expanding capabilities to deploy powerful telemetry and geolocation devices on small migratory animals (Bridge et al. 2011). Combining these new tagging devices with large-scale collaborative land-, sea-, and space-based tracking platforms, such as GLATOS (Krueger et al. 2017), the Ocean Tracking Network (Cooke et al. 2011), the European Tracking Network (Abecasis et al. 2018), the Motus Wildlife Tracking System (Taylor et al. 2018), ATLAS (Toledo et al. 2016), and ICARUS (Wikelski et al. 2007) will provide unparalleled opportunities for experimental migration biology.

558 **Acknowledgments**

559 Cooke was supported by the Canada Research Chairs program and an NSERC Discovery Grant.

560 Crossin, Guglielmo, and Norris were also supported by NSERC Discovery Grants. Cooke and

561 Lennox were also supported by Ocean Tracking Network based at Dalhousie University. Teffer

562 is supported by a Mitacs postdoctoral fellowship.

563

Author copy

Literature Cited

- Abecasis D., A. Steckenreuter, J. Reubens, K. Aarestrup, J. Alós, F. Badalamenti, L. Bajona, P. Boylan, K. Deneudt, L. Greenberg, et al. 2018. A review of acoustic telemetry in Europe and the need for a regional aquatic telemetry network. *Ani Biotelem* 6:12.
- Åkesson S. 2003. Avian long-distance navigation: experiments with migratory birds. In *Avian Migration*. Pp. 471-492 in P. Berthold, E. Gwinner and E. Sonnenschein, eds. Springer, Berlin.
- Åkesson S., and A. Hedenström. 2007. How migrants get there: migratory performance and orientation. *BioSci* 57:123-133.
- Åkesson S., G. Walinder, L. Karlsson, and S. Ehnborn. 2002. Nocturnal migratory flight initiation in reed warblers *Acrocephalus scirpaceus*: effect of wind on orientation and timing of migration. *J Avian Biol* 33:349-357.
- Alerstam T., A. Hedenström, and S. Åkesson. 2003. Long-distance migration: evolution and determinants. *Oikos* 103:247-260.
- Allee, W.C., A.E. Emerson, O. Park, T. Park, and K.P. Schmidt. 1949. *Principles of Animal Ecology*. W.B. Saunders Company, Philadelphia.
- Altizer S., R. Bartel, and B.A. Han. 2011. Animal migration and infectious disease risk. *Science* 331:296-302.
- Altizer S., R.S. Ostfeld, P.T. Johnson, S. Kutz, and C.D. Harvell. 2013. Climate change and infectious diseases: from evidence to a predictive framework. *Science* 341:514-519.
- Avens L., and K.J. Lohmann. 2004. Navigation and seasonal migratory orientation in juvenile sea turtles. *J Exp Biol* 207:1771-1778.
- Barbin G.P. (1998). The role of olfaction in homing and estuarine migratory behavior of yellow-phase American eels. *Can J Fish Aquat Sci* 55:564-575.
- Bass A.L., S.G. Hinch, A.K. Teffer, D.A. Patterson, and K.M. Miller. 2019. Fisheries capture and infectious agents are associated with travel rate and survival of Chinook salmon during spawning migration. *Fish Res* 209:156-166.
- Berchtold A., I. Nightingale, C. Vandermeer, and S.A. MacDougall-Shackleton. 2017. Experimental temperature manipulations alter songbird autumnal nocturnal migratory restlessness. *Anim Migr* 4:1-7.
- Betini G.S., C.G. Griswold, and D.R. Norris. 2013. Carry-over effects, sequential density dependence and the dynamics of populations in a seasonal environment. *Proc R Soc Lond Biol* 280:20130110.

- Birnie-Gauvin K., K.S. Peiman, A.J. Gallagher, R. de Bruijn, and S.J. Cooke. 2016. Sublethal consequences of urban life for wild vertebrates. *Env Rev* 24:416-425.
- Bohrer G., P.S. Beck, S.M. Ngene, A.K. Skidmore, and I. Douglas-Hamilton. 2014. Elephant movement closely tracks precipitation-driven vegetation dynamics in a Kenyan forest-savanna landscape. *Mov Ecol* 2:2-14.
- Boonstra R. 2013. The ecology of stress: a marriage of disciplines. *Funct Ecol* 27:7-10.
- Bowlin M.S., and M. Wikelski. 2008. Pointed wings, low wingloading and calm air reduce migratory flight costs in songbirds. *PLoS One* 3:e2154.
- Bowlin M.S., I.A. Bisson, J. Shamoun-Baranes, J.D. Reichard, N. Sapir, P.P. Marra, T.H. Kunz, D.S. Wilcove, A. Hedenström, C.G. Guglielmo et al. 2010. Grand challenges in migration biology. *Integr Comp Biol* 50:261-279.
- Bradley C.A., and S. Altizer. 2005. Parasites hinder monarch butterfly flight: implications for disease spread in migratory hosts. *Ecol Lett* 8:290-300.
- Bradshaw W.E., and C.M. Holzapfel. 2008. Genetic response to rapid climate change: it's seasonal timing that matters. *Mol Ecol* 17:157-166.
- Breuner C.W., T.P. Hahn. 2003 Integrating stress physiology, environmental change, and behavior in free-living sparrows. *Horm Behav* 43:115-123.
- Bridge E.S., K. Thorup, M.S. Bowlin, P.B. Chilson, R.H. Diehl, R.W. Fléron, P. Hartl, R. Keys, J.F. Kelly, W.D. Robinson et al. 2011. Technology on the move: recent and forthcoming innovations for tracking migratory birds. *BioSci* 61:689-698.
- Brown G.P., C. Kelehear, L. Pizzatto, and R. Shine. 2016. The impact of lungworm parasites on rates of dispersal of their anuran host, the invasive cane toad. *Biol Invasions* 18:103-114.
- Buehler D.M., and T. Piersma. 2007. Travelling on a budget: predictions and ecological evidence for bottlenecks in the annual cycle of long-distance migrants. *Phil Trans R Soc B* 363:247-266.
- Burgerhout E., C. Tudorache, S.A. Brittiijn, A.P. Palstra, R.P. Dirks, and G.E. van den Thillart. 2013. Schooling reduces energy consumption in swimming male European eels, *Anguilla anguilla* L. *J Exp Mar Biol Ecol* 448:66-71.
- Burnett N.J., S.G. Hinch, D.C. Braun, M.T. Casselman, C.T. Middleton, S.M. Wilson, and S.J. Cooke. 2014. Burst swimming in areas of high flow: delayed consequences of anaerobiosis in wild adult sockeye salmon. *Physiol Biochem Zool* 87:587-598.

- Caldwell R.L., and M.A. Rankin. 1972. Effects of a juvenile hormone mimic on flight in the milkweed bug, *Oncopeltus fasciatus*. *Gen Comp Endocrinol* 19:601-605.
- Carr A.F., and L.H. Ogren. 1960. The ecology and migrations of sea turtles. 4, The green turtle in the Caribbean Sea. *Bull Am Mus Nat Hist* 121:1.
- Caudill C.C., W.R. Daigle, M.L. Keefer, C.T. Boggs, M.A. Jepson, B.J. Burke, R.W. Zabel, T.C. Bjornn, and C.A. Peery. 2007. Slow dam passage in adult Columbia River salmonids associated with unsuccessful migration: delayed negative effects of passage obstacles or condition-dependent mortality? *Can J Fish Aquat Sci* 64:979-995.
- Chapman B.B., C. Brönmark, J.Å. Nilsson, and L.A. Hansson. 2011. The ecology and evolution of partial migration. *Oikos* 120:1764-1775.
- Clark T.D., K.M. Jeffries, S.G. Hinch, and A.P. Farrell. 2011. Exceptional aerobic scope and cardiovascular performance of pink salmon (*Oncorhynchus gorbuscha*) may underlie resilience in a warming climate. *J Exp Biol* 214:3074-3081.
- Cochran W.W., H. Mouritsen, and M. Wikelski. 2004. Migrating songbirds recalibrate their magnetic compass daily from twilight cues. *Science* 304:405-408.
- Cohen E.B., F.R. Moore, and R.A. Fischer. 2012. Experimental evidence for the interplay of exogenous and endogenous factors on the movement ecology of a migrating songbird. *PloS One* 7:e41818.
- Conenna I., A. López-Baucells, R. Rocha, S. Ripperger, and M. Cabeza. 2019. Movement seasonality in a desert-dwelling bat revealed by miniature GPS loggers. *Mov Ecol* 7:27-37.
- Conklin J.R., and P.F. Battley. 2012. Carry-over effects and compensation: late arrival on non-breeding grounds affects wing moult but not plumage or schedules of departing bar-tailed godwits *Limosa lapponica baueri*. *J Avian Biol* 43:252-263.
- Cooke S.J., S.G. Hinch, A.P. Farrell, D.A. Patterson, K. Miller-Saunders, D.W. Welch, M.R. Donaldson, K.C. Hanson, G.T. Crossin, M.T. Mathes et al. 2008. Developing a mechanistic understanding of fish migrations by linking telemetry with physiology, behavior, genomics and experimental biology: an interdisciplinary case study on adult Fraser River sockeye salmon. *Fisheries* 33:321-339.
- Cooke S.J., S.J. Iverson, M.J. Stokesbury, S.G. Hinch, A.T. Fisk, D.L. VanderZwaag, R. Apostle, and F. Whoriskey. 2011. Ocean Tracking Network Canada: a network approach to addressing critical issues in fisheries and resource management with implications for ocean governance. *Fisheries* 36:583-592.
- Cooke S.J., K. Birnie-Gauvin, R.J. Lennox, J.J. Taylor, T. Rytwinski, J.L. Rummer, C.E. Franklin, J.R. Bennett, and N.R. Haddaway. 2017. How experimental biology and

- ecology can support evidence-based decision-making in conservation: avoiding pitfalls and enabling application. *Conserv Physiol* 5:cox043.
- Crossin G.T., S.G. Hinch, S.J. Cooke, D.W. Welch, D.A. Patterson, S.R.M. Jones, A.G. Lotto, R.A. Leggatt, M.T. Mathes, J.M. Shrimpton et al. 2008. Exposure to high temperature influences the behaviour, physiology, and survival of sockeye salmon during spawning migration. *Can J Zool* 86:127-140.
- Crossin G.T., S.G. Hinch, D.W. Welch, S.J. Cooke, D.A. Patterson, J.A. Hills, Y. Zohar, U. Klenke, M.C. Jacobs, L.B. Pon et al. 2009. Physiological profiles of sockeye salmon in the Northeastern Pacific Ocean and the effects of exogenous GnRH and testosterone on rates of homeward migration. *Mar Fresh Behav Physiol* 42:89-108.
- Dailey T.V., and N.T. Hobbs. 1989. Travel in alpine terrain: energy expenditures for locomotion by mountain goats and bighorn sheep. *Can J Zool* 67:2368-2375.
- Devlin R.H., L.F. Sundstrom, and W.M. Muir. 2006. Interface of biotechnology and ecology for environmental risk assessments of transgenic fish. *Trends Biotechnol* 24:89-97.
- Dhabhar F.S., and B.S. McEwen. 1997. Acute stress enhances while chronic stress suppresses cell-mediated immunity in vivo: A potential role for leukocyte trafficking. *Brain Behav Immun* 11:286-306.
- Dingle H. 1965. The relation between age and flight activity in the milkweed bug, *Oncopeltus*. *J Exp Biol* 42:269-283.
- Dingle H. 1966. Some factors affecting flight activity in individual milkweed bugs (*Oncopeltus*). *J Exp Biol* 44:335-343.
- Dingle H. 2006. Animal migration: is there a common migratory syndrome?. *J Ornithol* 147:212-220.
- Dingle H., and V.A. Drake. 2007. What is migration?. *BioSci* 57:113-121.
- Domeier M.L. and N. Nasby-Lucas. 2007. Annual re-sightings of photographically identified white sharks (*Carcharodon carcharias*) at an eastern Pacific aggregation site (Guadalupe Island, Mexico). *Mar Biol* 150:977-984.
- Donaldson M.R., S.G. Hinch, D.A. Patterson, J. Hills, J.O. Thomas, S.J. Cooke, G.D. Raby, L.A. Thompson, D. Robichaud, K.K. English et al. 2011. The consequences of angling, beach seining, and confinement on the physiology, post-release behaviour and survival of adult sockeye salmon during upriver migration. *Fish Res* 108:133-141.
- Dukas R., and L. Dukas. 2011. Coping with nonrepairable body damage: effects of wing damage on foraging performance in bees. *Anim Behav* 81 :635-638.

- Dussaubat C., A. Maisonnasse, D. Crauser, D. Beslay, G. Costagliola, S. Soubeyrand, A. Kretzchmar, Y. Le Conte. 2013. Flight behavior and pheromone changes associated to *Nosema ceranae* infection of honey bee workers (*Apis mellifera*) in field conditions. *J Invert Pathol* 113:42-51.
- Duvall D., and D.O. Norris. 1980. Stimulation of terrestrial-substrate preferences and locomotor activity in newly transformed tiger salamanders (*Ambystoma tigrinum*) by exogenous or endogenous thyroxine. *Anim Behav* 28:116-123.
- Eliason E.J., T.D. Clark, M.J. Hague, L.M. Hanson, Z.S. Gallagher, K.M. Jeffries, M.K. Gale, D.A. Patterson, S.G. Hinch, and A.P. Farrell. 2011. Differences in thermal tolerance among sockeye salmon populations. *Science* 332:109-112.
- Elliott K.H., M. Le Vaillant, A. Kato, A.J. Gaston, Y. Ropert-Coudert, J.F. Hare, J.R. Speakman, and D. Croll. 2014. Age-related variation in energy expenditure in a long-lived bird within the envelope of an energy ceiling. *J Anim Ecol* 83:136-146.
- Eng M.L., B.J. Stutchbury, and C.A. Morrissey. 2017. Imidacloprid and chlorpyrifos insecticides impair migratory ability in a seed-eating songbird. *Sci Rep* 7:15176.
- Eng M.L., B.J.M. Stutchbury, and C.A. Morrissey. 2019. A neonicotinoid insecticide reduces fueling and delays migration in songbirds. *Science* 365:1177-1180.
- Etheredge J.A., S.M. Perez, O.R. Taylor, and R. Jander. 1999. Monarch butterflies (*Danaus plexippus* L.) use a magnetic compass for navigation. *Proc Nat Acad Sci* 96:13845-13846.
- Farrell A.P. 2009. Environment, antecedents and climate change: lessons from the study of temperature physiology and river migration of salmonids. *J Exp Biol* 212:3771-3780.
- Flockhart D.T., T.G. Martin, and D.R. Norris. 2012. Experimental examination of intraspecific density-dependent competition during the breeding period in monarch butterflies (*Danaus plexippus*). *PLoS One* 7:e45080.
- Fransson T. 1998. A feeding experiment on migratory fuelling in whitethroats, *Sylvia communis*. *Anim Behav* 55:153-162.
- Furey N.B. (2016). Migration ecology of juvenile Pacific salmon smolts: the role of fish condition and behaviour across landscapes. PhD thesis, University of British Columbia, British Columbia, Canada.
- Furey N.B., J.B. Armstrong, D.A. Beauchamp, and S.G. Hinch. 2018. Migratory coupling between predators and prey. *Nat Ecol Evol* 2:1846.

- Goymann W. S. Lupi, H. Kaiya, M. Cardinale, and L. Fusani. 2017. Ghrelin affects stopover decisions and food intake in a long-distance migrant. *Proc Nat Acad Sci* 114:1946-1951.
- Griswold C.K., C.M. Taylor, and D.R. Norris. 2010. The evolution of migration in a seasonal environment. *Proc R Soc B* 277:2711-2720.
- Gula R., R. Hausknecht, and R. Kuehn. 2009. Evidence of wolf dispersal in anthropogenic habitats of the Polish Carpathian Mountains. *Biodivers Cons* 18:2173-2184.
- Gwinner E. 2003. Circannual rhythms in birds. *Curr Opin Neurosci* 13:770-778.
- Hansen, M.J., D.E. Cocherell, S.J. Cooke, P.H. Patrick, M. Sills, and N.A. Fanguie. 2018. Behavioural guidance of Chinook salmon smolts: the variable effects of LED spectral wavelength and strobing frequency. *Cons Physiol* 6:coy032.
- Harrison X.A., J.D. Blount, R. Inger, D.R. Norris, and S. Bearhop. 2011. Carry-over effects as drivers of fitness differences in animals. *J Anim Ecol* 80:4-18.
- Hau M., S. Casagrande, J.Q. Ouyang, and A.T. Baugh. 2016. Glucocorticoid-mediated phenotypes in vertebrates: multilevel variation and evolution. *Adv Study Behav* 48:41-115.
- Hazekamp A.A., R. Mayer, and N. Osinga. 2010. Flow simulation along a seal: the impact of an external device. *Eur J Wildl Res* 56:131-140.
- Hedenström A., and Å. Lindström. 2017. Wind tunnel as a tool in bird migration research. *J Avian Biol* 48:37-48.
- Hegemann A., P. Alcalde Abril, S. Sjöberg, R. Muheim, T. Alerstam, J.Å. Nilsson, and D. Hasselquist. 2018. A mimicked bacterial infection prolongs stopover duration in songbirds—but more pronounced in short-than long-distance migrants. *J Anim Ecol* 87:1698-1708.
- Hellström G., J. Klaminder, F. Finn, L. Persson, A. Alanärä, M. Jonsson, J. Fick, and T. Brodin. 2016. GABAergic anxiolytic drug in water increases migration behaviour in salmon. *Nat Commun* 7:13460.
- Hoffmann A.A., S.L. Chown, and S. Clusella-Trullas. 2013. Upper thermal limits in terrestrial ectotherms: how constrained are they?. *Funct Ecol* 27:934-949.
- Holland R.A., K. Thorup, A. Gagliardo, I.A. Bisson, E. Knecht, D. Mizrahi, and M. Wikelski. 2009. Testing the role of sensory systems in the migratory heading of a songbird. *J Exp Biol* 212:4065-4071.
- Holland R.A. 2014. True navigation in birds: from quantum physics to global migration. *J Zool* 293:1-15.

- Jeffries K.M., S.G. Hinch, M.K. Gale, T.D. Clark, A.G. Lotto, M.T. Casselman, S. Li, E.L. Rechisky, A.D. Porter, D.W. Welch et al. 2014. Immune response genes and pathogen presence predict migration survival in wild salmon smolts. *Mol Ecol* 23:5803-5815.
- Jepsen N., K. Aarestrup, F. Økland, and G. Rasmussen. 1998. Survival of radiotagged Atlantic salmon (*Salmo salar* L.)–and trout (*Salmo trutta* L.) smolts passing a reservoir during seaward migration. *Hydrobiol* 371:347.
- Kapuscinski A.R., K.R. Hayes, S. Li, and G. Dana. 2008. Environmental Risk Assessment of Genetically Modified Organisms. Vol 3. Methodologies for Transgenic Fish. CAB International, Oxford.
- Keefer M.L., C.A. Peery, S.R. Lee, W.R. Daigle, E.L. Johnson, and M.L. Moser. 2011. Behaviour of adult Pacific lamprey in near-field flow and fishway design experiments. *Fish Manag Ecol* 18:177-189.
- Kellermann J.L., and C. van Riper. 2015. Detecting mismatches of bird migration stopover and tree phenology in response to changing climate. *Oecologia* 178:1227-1238.
- Kennedy J.S., and C.O. Booth. 1963. Free flight of aphids in the laboratory. *J Exp Biol* 40:67–85.
- Kennedy J.S., and C.O. Booth. 1964. Co-ordination of successive activities in an aphid Depression of settling after flight. *J Exp Biol* 41:805–824.
- Kinnison M.T. M.J. Unwin, A.P. Hendry, and T.P. Quinn. 2001. Migratory costs and the evolution of egg size and number in introduced and indigenous salmon populations. *Evol* 55:1656-1667.
- Kinnison M.T., M.J. Unwin, and T.P. Quinn. 2003. Migratory costs and contemporary evolution of reproductive allocation in male chinook salmon. *J Evol Biol* 16:1257-1269.
- Kishkinev D., D. Heyers, B.K. Woodworth, G.W. Mitchell, K.A. Hobson, and D.R. Norris. 2016. Experienced migratory songbirds do not display goal-ward orientation after release following a cross-continental displacement: an automated telemetry study. *Sci Rep* 6:37326.
- Kitahashi T., A. Sato, D. Alok, M. Kaeriyama, Y. Zohar, K. Yamauchi, A. Urano, and H. Ueda. 1998. Gonadotropin-releasing hormone analog and sex steroids shorten homing duration of sockeye salmon in Lake Shikotsu. *Zool Sci* 15:767-771.
- Klaassen M., A. Kvist, and Å. Lindström. 2000. Flight costs and fuel composition of a bird migrating in a wind tunnel. *Condor* 102:444-451.

- Krueger C.C., C.M. Holbrook, T.R. Binder, C.S. Vandergoot, T.A. Hayden, D.W. Hondorp, N. Nate, K. Paige, S.C. Riley, A.T. Fisk et al. 2017. Acoustic telemetry observation systems: challenges encountered and overcome in the Laurentian Great Lakes. *Can J Fish Aquat Sci* 75:1755-1763.
- Kunz T.H., S.A. Gauthreaux Jr, N.I. Hristov, J.W. Horn, G. Jones, E.K. Kalko, R.P. Larkin, G.F. McCracken, S.M. Swartz, R.B. Srygley et al. 2008. Aeroecology: probing and modeling the aerosphere. *Integr Comp Biol* 48:1-11.
- Lague S.L., B. Chua, A.P. Farrell, Y. Wang, and W.K. Milsom. 2016. Altitude matters: differences in cardiovascular and respiratory responses to hypoxia in bar-headed geese reared at high and low altitudes. *J Exp Biol* 219:1974-1984.
- Lee C.G., A.P. Farrell, A. Lotto, S.G. Hinch, and M.C. Healey. 2003. Excess post-exercise oxygen consumption in adult sockeye (*Oncorhynchus nerka*) and coho (*O. kisutch*) salmon following critical speed swimming. *J Exp Biol* 206:3253-3260.
- LeMaster M.P., I.T. Moore, R.T., and Mason. 2001. Conspecific trailing behavior of red-sided garter snakes, *Thamnophis sirtalis parietalis*, in the natural environment. *Anim Behav* 61:827-833.
- Lennox R.J., J.M. Chapman, C.M. Souliere, C. Tudorache, M. Wikelski, J.D. Metcalfe, and S.J. Cooke. 2016. Conservation physiology of animal migration. *Conserv Physiol* 4:cov072.
- Lennox R.J., E.B. Thorstad, O.H. Diserud, F. Økland, S.J. Cooke, I. Aasestad, and T. Forseth. 2018. Biotic and abiotic determinants of the ascent behaviour of adult Atlantic salmon transiting passable waterfalls. *Riv Res Appl* 34:907-917.
- Lesbarrères D, T. Lodé, and J. Merilä. 2004. What type of amphibian tunnel could reduce road kills?. *Oryx* 38:220-223.
- Lindström Å., A. Kvist, T. Piersma, A. Dekinga, and M.W. Dietz. 2000. Avian pectoral muscle size rapidly tracks body mass changes during flight, fasting and fuelling. *J Exp Biol* 203:913-919.
- Lohmann K.J., N.F. Putman, and C.M. Lohmann. 2008. Geomagnetic imprinting: a unifying hypothesis of long-distance natal homing in salmon and sea turtles. *Proc Nat Acad Sci* 105:19096-19101.
- Ma Y., C.R. Perez, B.A. Branfireun, and C.G. Guglielmo. 2018. Dietary exposure to methylmercury affects flight endurance in a migratory songbird. *Environ Poll* 234:894-901.
- Maggini I., L.V. Kennedy, K.H. Elliott, K.M. Dean, R. MacCurdy, A. Macmillan, C.A. Pritsos, and C.G. Guglielmo. 2017a. Trouble on takeoff: crude oil on feathers reduces escape performance of shorebirds. *Ecotox Env Safe* 141:171-177.

- 928
929 Maggini I., L.V. Kennedy, A. Macmillan, K.H. Elliott, K. Dean, and C.G. Guglielmo. 2017b.
930 Light oiling of feathers increases flight energy expenditure in a migratory shorebird. *J*
931 *Exp Biol* 220:2372-2379.
932
- 933 Martin T.G., I. Chadès, P. Arcese, P.P. Marra, H.P. Possingham, and D.R. Norris. 2007. Optimal
934 conservation of migratory species. *PloS one* 2:e751.
935
- 936 Matter S.F., and J. Roland. 2002. An experimental examination of the effects of habitat quality
937 on the dispersal and local abundance of the butterfly *Parnassius smintheus*. *Ecol Entomol*
938 27:308-316.
939
- 940 Mayr E. 1961. Cause and effect in biology. *Science* 134:1501-1506.
941
- 942 McWilliams S.R., C. Guglielmo, B. Pierce, and M. Klaassen. 2004. Flying, fasting, and feeding
943 in birds during migration: a nutritional and physiological ecology perspective. *J Avian*
944 *Biol* 35:377-393.
945
- 946 Midwood J.D., M.H. Larsen, M. Boel, N. Jepsen, K. Aarestrup, and S.J. Cooke. 2014. Does
947 cortisol manipulation influence outmigration behaviour, survival and growth of sea trout?
948 A field test of carryover effects in wild fish. *Mar Ecol Progr Ser* 496:135-144.
949
- 950 Midwood J.D., M.H. Larsen, K. Aarestrup, and S.J. Cooke. 2016. Stress and food deprivation:
951 linking physiological state to migration success in a teleost fish. *J Exp Biol* 219:3712-
952 3718.
953
- 954 Miller K.M., A. Teffer, S. Tucker, S. Li, A.D. Schulze, M. Trudel, F. Juanes, A. Tabata, K.H.
955 Kaukinen, N.G. Ginther et al. 2014. Infectious disease, shifting climates, and
956 opportunistic predators: cumulative factors potentially impacting wild salmon declines.
957 *Evol Appl* 7:812-855.
958
- 959 Minderman J., C.J. Pendlebury, J.W. Pearce-Higgins, and K.J. Park. 2012. Experimental
960 evidence for the effect of small wind turbine proximity and operation on bird and bat
961 activity. *PLoS One* 7:e41177.
962
- 963 Minter M., A. Pearson, K.S. Lim, K. Wilson, J.W. Chapman, and C.M. Jones. 2018. The tethered
964 flight technique as a tool for studying life-history strategies associated with migration in
965 insects. *Ecol Entomol* 43:397-411.
966
- 967 Mitamura H., N. Arai, W. Sakamoto, Y. Mitsunaga, H. Tanaka, Y. Mukai, K. Nakamura, M.
968 Sasaki, and Y. Yoneda. 2005. Role of olfaction and vision in homing behaviour of black
969 rockfish *Sebastes inermis*. *J Exp Mar Biol Ecol* 322:123-134.
970
- 971 Mouritsen H., R. Derbyshire, J. Stalleicken, O.Ø. Mouritsen, B.J. Frost, and D.R. Norris. 2013.
972 An experimental displacement and over 50 years of tag-recoveries show that monarch
973 butterflies are not true navigators. *Proc Nat Acad Sci* 110:7348-7353.

- Müller F., G. Rüppel, and H. Schmaljohann. 2018. Does the length of the night affect the timing of nocturnal departures in a migratory songbird?. *Anim Behav* 141:183-194.
- Navarro J., J. González-Solís, G. Viscor, and O. Chastel. 2008. Ecophysiological response to an experimental increase of wing loading in a pelagic seabird. *J Exp Mar Biol Ecol* 358:14-19.
- Newton I. 2008. *The migration ecology of birds*. Elsevier, New York.
- Nolan D.T., P. Reilly, and S.W. Bonga. 1999. Infection with low numbers of the sea louse *Lepeophtheirus salmonis* induces stress-related effects in postsmolt Atlantic salmon (*Salmo salar*). *Can J Fish Aquat Sci* 56:947-959.
- Norberg U., K. Enfjäll, and O. Leimar. 2002. Habitat exploration in butterflies—an outdoor cage experiment. *Evol Ecol* 16:1-14.
- Norris D.R. 2005. Carry-over effects and habitat quality in migratory populations. *Oikos* 109:178-186.
- Norris D.R., and C.M. Taylor. 2006. Predicting the consequences of carry-over effects in migratory animals. *Biol Lett* 2:148-151.
- Norris D.R., and P.P. Marra. 2007. Seasonal interactions, habitat quality, and population dynamics in migratory birds. *Condor* 109:535-547.
- Norton S., L.A.L. Corner, and R.S. Morris. 2005. Ranging behaviour and duration of survival of wild brushtail possums (*Trichosurus vulpecula*) infected with *Mycobacterium bovis*. *New Zeal Vet J* 53:293-300.
- Noyce K.V., and D.L. Garshelis. 2014. Follow the leader: social cues help guide landscape-level movements of American black bears (*Ursus americanus*). *Can J Zool* 92:1005-1017.
- Nuismer S.L., J.N. Thompson, and R. Gomulkiewicz. 1999. Gene flow and geographically structured coevolution. *Proc R Soc Lond B* 266:605-609.
- Nunes S., T.R. Duniec, S.A. Schweppe, and K.E. Holekamp. 1999. Energetic and endocrine mediation of natal dispersal behavior in Belding's ground squirrels. *Horm Behav* 35:113-124.
- O'Connor C.M., D.R. Norris, G.T. Crossin, and S.J. Cooke. 2014. Biological carryover effects: linking common concepts and mechanisms in ecology and evolution. *Ecosph* 5:1-11.
- Padget O., S.L. Bond, M.M. Kavelaars, E. van Loon, M. Bolton, A.L. Fayet, M. Syposz, S. Roberts, and T. Guilford. 2018. In situ clock shift reveals that the sun Compass contributes to orientation in a pelagic seabird. *Curr Biol* 28:275-279.

- Peiman K.S., K. Birnie-Gauvin, J.D. Midwood, M.H. Larsen, A.D.M. Wilson, K. Aarestrup, and S.J. Cooke. 2017. If and when: Intrinsic differences and environmental stressors influence migration in brown trout (*Salmo trutta*). *Oecol* 184:375-384.
- Perez S.M., and O.R. Taylor. 2004. Monarch butterflies' migratory behavior persists despite changes in environmental conditions. Pp. 85-89 in K.S. Oberhauser, and M.J. Solensky, eds. *The monarch butterfly: Biology and conservation*. Cornell University Press, London.
- Phol H. 2000. Circadian control of migratory restlessness and the effects of exogenous melatonin in the brambling, *Fringilla montifringilla*. *Chronobiol Int* 17:471-488.
- Pierce B.J., S.R. McWilliams, T.P. O'Connor, A.R. Place, and C.G. Guglielmo. 2005. Effect of dietary fatty acid composition on depot fat and exercise performance in a migrating songbird, the red-eyed vireo. *J Exp Biol* 208:1277-1285.
- Pollonara E., P. Luschi, T. Guilford, M. Wikelski, F. Bonadonna, and A. Gagliardo. 2015. Olfaction and topography, but not magnetic cues, control navigation in a pelagic seabird: displacements with shearwaters in the Mediterranean Sea. *Sci Rep* 5:16486.
- Prange H.D. 1976. Energetics of swimming of a sea turtle. *J Exp Biol* 64:1-12.
- Pulido F., and P. Berthold. 2010. Current selection for lower migratory activity will drive the evolution of residency in a migratory bird population. *Proc Nat Acad Sci* 107:7341-7346.
- Ramenofsky M., and J.C. Wingfield. 2007. Regulation of migration. *BioSci* 57:135-143.
- Ramos R., V. Llabrés, L. Monclús, M. López-Béjar, and J. González-Solís. 2018. Costs of breeding are rapidly buffered and do not affect migratory behavior in a long-lived bird species. *Ecol* 99:2010-2024.
- Rankin M.A. 1978. Hormonal control of insect migratory behavior. In *Evolution of insect migration and diapause*. Pp. 5-32 in H. Dingle, ed. Springer, New York.
- Rankin M.A., and L.M. Riddiford. 1977. Hormonal control of migratory flight in *Oncopeltus fasciatus*: the effects of the corpus cardiacum, corpus allatum, and starvation on migration and reproduction. *Gen Comp Endocrinol* 33:309-321.
- Reed T.E., V. Grøtan, S. Jenouvrier, B.E. Sæther, M.E. Visser. 2013. Population growth in a wild bird is buffered against phenological mismatch. *Science* 340:488-491.
- Rivrud I.M., M. Heurich, P. Krupczynski, J. Müller, and A. Mysterud. 2016. Green wave tracking by large herbivores: an experimental approach. *Ecol* 97:3547-3553.
- Robinson W.D., M.S. Bowlin, I. Bisson, J. Shamoun-Baranes, K. Thorup, R.H. Diehl, T.H. Kunz, S. Mabey, and D.W. Winkler. 2010. Integrating concepts and technologies to advance the study of bird migration. *Front Ecol Evol* 8:354-361.

- Satterfield D.A., J.C. Maerz, M.D. Hunter, D.T. Flockhart, K.A. Hobson, D.R. Norris, H. Streit, J.C. de Roode, and S. Altizer. 2018. Migratory monarchs that encounter resident monarchs show life-history differences and higher rates of parasite infection. *Ecol Lett* 21:1670-1680.
- Sato A., H. Ueda, M. Fukaya, M. Kaeriyama, Y. Zohar, A. Urano, and K. Yamauchi, K. 1997. Sexual differences in homing profiles and shortening of homing duration by gonadotropin-releasing hormone analog implantation in lacustrine sockeye salmon (*Oncorhynchus nerka*) in Lake Shikotsu. *Zool Sci* 14:1009-1014.
- Schwenk K., D.K. Padilla, G.S. Bakken, and R.J. Full. 2009. Grand challenges in organismal biology. *Integr Comp Biol* 49:7-14.
- Seewagen C.L., Y. Ma, Y.E. Morbey, and C.G. Guglielmo. 2019. Stopover departure behaviors and flight orientations of spring-migrant Yellow-rumped Warblers (*Setophaga coronata*) experimentally exposed to methylmercury. *J Ornithol* 160:617-624.
- Silva A.T., J.M. Santos, M.T. Ferreira, A.N. Pinheiro, and C. Katopodis. 2012. Passage efficiency of offset and straight orifices for upstream movements of Iberian barbel in a pool-type fishway. *Riv Res Appl* 28:529-542.
- Skov C., K. Aarestrup, H. Baktoft, J. Brodersen, C. Brönmark, L.A. Hansson, E.E. Nielsen, T. Nielsen, and P.A. Nilsson. 2010. Influences of environmental cues, migration history, and habitat familiarity on partial migration. *Behav Ecol* 21:1140-1146.
- Sopinka N.M., L.D. Patterson, J.C. Redfern, N.K. Pleizier, C.B. Belanger, J.D. Midwood, G.T. Crossin, and S.J. Cooke. 2015. Manipulating glucocorticoids in wild animals: basic and applied perspectives. *Conserv Physiol* 3:cov031.
- Spoelstra K., R.H. van Grunsven, M. Donners, P. Gienapp, M.E. Huigens, R. Slaterus, F. Berendse, M.E. Visser, and E. Veenendaal. 2015. Experimental illumination of natural habitat—an experimental set-up to assess the direct and indirect ecological consequences of artificial light of different spectral composition. *Phil Trans R Soc B* 370:20140129.
- Sundström L.F., M. Löhmus, and R.H. Devlin. 2010. Migration and growth potential of coho salmon smolts: implications for ecological impacts from growth-enhanced fish. *Ecol Appl* 20:1372-1383.
- Sutherland W.J. and K. Norris. 2002. Behavioural models of population growth rates: implications for conservation and prediction. *Phil Trans R Soc Lond B* 357:1273-1284.
- Tassava R.A., and C. Kuenzli. 1979. The effects of prolactin and thyroxine on tail fin height, habitat choice, and forelimb regeneration in the adult newt (*Notophthalmus viridescens*). *Ohio J Sci* 79:32-37.

- Taylor C.M., and D.R. Norris. 2010. Population dynamics in migratory networks. *Theoret Ecol* 3:65-73.
- Taylor P., T. Crewe, S. Mackenzie, D. Lepage, Y. Aubry, Z. Crysler, G. Finney, C.M. Francis, C.G. Guglielmo et al. 2017. The Motus Wildlife Tracking System: a collaborative research network to enhance the understanding of wildlife movement. *Avian Conserv Ecol* 12:8.
- Teffler A.K., A.L. Bass, K.M. Miller, D.A. Patterson, F. Juanes, and S.G. Hinch. 2018. Infections, fisheries capture, temperature, and host responses: multistressor influences on survival and behaviour of adult Chinook salmon. *Can J Fish Aquat Sci* 75:2069-2083.
- Thums M., S.D. Whiting, J. Reisser, K.L. Pendoley, C.B. Pattiaratchi, M. Proietti, Y. Hetzel, R. Fisher, and M.G. Meekan. 2016. Artificial light on water attracts turtle hatchlings during their near shore transit. *R Soc Open Sci* 3:160142.
- Toledo S., O. Kishon, Y. Orchan, A. Shohat, and R. Nathan. 2016. Lessons and experiences from the design, implementation, and deployment of a wildlife tracking system. *Proceedings of the IEEE International Conference on Software Science, Technology and Engineering*. <https://ieeexplore.ieee.org/abstract/document/7515410>
- Tsoar A., R. Nathan, Y. Bartan, A. Vyssotski, G. Dell'Omo, and N. Ulanovsky. 2011. Large-scale navigational map in a mammal. *Proc Nat Acad Sci* 108:E718-E724.
- Ueda H., M. Kaeriyama, K. Mukasa, A. Urano, H. Kudo, T. Shoji, Y. Tokumitsu, K. Yamauchi, and K. Kurihara. 1998. Lacustrine sockeye salmon return straight to their natal area from open water using both visual and olfactory cues. *Chem Senses* 23:207-212.
- Underwood A.J. 1990. Experiments in ecology and management: their logics, functions and interpretations. *Austral Ecol* 15:365-389.
- Van der Graaf A.J., J. Stahl, A. Klimkowska, J.P. Bakker, and R.H. Drent. 2006. Surfing on a green wave-how plant growth drives spring migration in the Barnacle Goose *Branta leucopsis*. *Ardea* 94:567.
- Visser M.E., A.C. Perdeck, J.H. van Balen, and C. Both. 2009. Climate change leads to decreasing bird migration distances. *Glob Change Biol* 15:1859-1865.
- Wagner G.N., R.S. McKinley, P.A. Bjørn, and B. Finstad. 2003. Physiological impact of sea lice on swimming performance of Atlantic salmon. *J Fish Biol* 62:1000-1009.
- Wagner G.N., S.G. Hinch, L.J. Kuchel, A. Lotto, S.R. Jones, D.A. Patterson, J.S. Macdonald, G. Van Der Kraak, M. Shrimpton, K.K. English et al. 2005. Metabolic rates and swimming performance of adult Fraser River sockeye salmon (*Oncorhynchus nerka*) after a controlled infection with *Parvicapsula minibicornis*. *Can J Fish Aquat Sci* 62:2124-2133.

- Walcott C. 1996. Pigeon homing: observations, experiments and confusions. *J Exp Biol* 199:21-27.
- Wikelski M., E.M. Tarlow, A. Raim, R.H. Diehl, R.P. Larkin, and G.H. Visser. 2003. Avian metabolism: costs of migration in free-flying songbirds. *Nature* 423:704.
- Wikelski M., R.W. Kays, N.J. Kasdin, K. Thorup, J.A. Smith, and G.W. Swenson. 2007. Going wild: what a global small-animal tracking system could do for experimental biologists. *J Exp Biol* 210:181-186.
- Wilcove D.S. (2008). No way home: the decline of the world's great animal migrations. Island Press, Washington, DC.
- Wilcove D.S., and M. Wikelski. 2008. Going, going, gone: is animal migration disappearing? *PLoS One* 6:1361-1364.

1174 **Tables**

1175 **Table 1. Abiotic challenges experiments.**

Taxa	Species	Experimental Approach	References
Birds	White-throated sparrow (<i>Zonotrichia albicollis</i>)	Sparrows did not exhibit migratory restlessness at the warmest temperature treatment (24°C), suggesting that migration phenotypes respond to temperature	Berchtold et al. 2017
Fish	Brown trout (<i>Salmo trutta</i>)	Trout exposed to a temperature challenge designed to instigate a stress response migrated earlier than control animals, suggesting that acute environmental stressors influence an individual's migration strategy	Peiman et al. 2017
Herptiles	Green turtle (<i>Chelonia mydas</i>)	Hatchling turtles were exposed to natural light or artificial light early in the marine migration, showing significant affinity for the artificial lights compared to individuals released on nights with only natural light	Thums et al. 2016
Invertebrates	Monarch butterfly (<i>Danaus plexippus</i>)	Monarchs were either held at control conditions or treated with longer photoperiod and warmer temperatures; treated monarchs displayed similar behaviour to control individuals, suggesting the manipulation had not altered their migratory preparedness	Perez and Taylor 2004
Mammals	Red deer (<i>Cervus elaphus</i>)	The forage maturation hypothesis was tested by manipulating the release date of red deer in Germany. Deer released late missed the peak plant maturity and migrated more rapidly to elevation to compensate whereas deer released early capitalized on peak vegetation production and migrated more slowly	Rivrud et al. 2016

1176
1177
1178
1179
1180
1181

1182 **Table 2. Biotic challenges experiments.**

Taxa	Species	Experimental Approach	References
Birds	Six species of passerine birds	Experimentally challenged the immune system of six passerine species during autumn migration; immune-challenged groups prolonged their stopover duration	Hegemann et al. 2018
Fish	Atlantic salmon (<i>Salmo salar</i>)	Experimental infection of post-smolt salmon with sea lice (<i>Lepeophtheirus salmonis</i>) impacted the ability of fish to maintain homeostasis and constrained migration success	Nolan et al. 1999
Herptiles	Cane toad (<i>Rhinella marina</i>)	Comparative field study of experimentally-infected toads (with a lungworm) and uninfected controls using radio telemetry; revealed that lungworms did not modify rates of invasive toad dispersal	Brown et al. 2016
Invertebrates	Honey bee (<i>Apis mellifera</i>)	Comparative study of the flight activity of experimentally-infected (with <i>Nosema ceranae</i>) and non-infected bees in small colonies using an electronic optic bee counter. Infected bees showed precocious and elevated flight activity	Dussaubat et al. 2013
Mammals	Brush-tail possum (<i>Trichosurus vulpecula</i>)	Possums were experimentally infected with <i>Mycobacterium bovis</i> and their movement and activity was compared to non-infected possums in New Zealand using radio telemetry	Norton et al. 2005

1197 **Table 3. Bioenergetic and biomechanical challenges**

Taxa	Species	Experimental Approach	References
Birds	Cory's shearwater (<i>Calonectris diomedea</i>)	Increased workload by adding 45g of extra weight to the birds which led to extended foraging trips, lower rate of mass gain, and indication of muscular damage	Navarro et al. 2008
Fish	European eel (<i>Anguilla anguilla</i>)	Male eels were swum in groups and individually, and oxygen consumption measured; swimming costs were reduced by 1/3 when in groups	Burgerhout et al. 2013
Herptiles	Green turtles (<i>Chelonia mydas</i>)	Turtles were swum at sustained speeds and oxygen consumption was measured; the cost of transport is lower for turtles than for birds but higher than for fish of similar size	Prange 1976
Invertebrates	Honey bee (<i>Apis mellifera</i>)	Bees with experimentally trimmed wings foraged at similar rates but had reduced food delivery; may affect bees with damaged wings during seasonal migrations	Dukas and Dukas 2011
Mammals	Mountain goat (<i>Oreamnos americanus</i>) and bighorn sheep (<i>Ovis canadensis</i>)	Terrain conditions were altered using a treadmill to assess the energetic costs and efficiency of locomotion under different conditions (variable activity level, slope and snow condition)	Dailey and Hobbs 1989

1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215

1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227

Table 4. Endocrine and signaling pathways experiments

Taxa	Species	Experimental Approach	References
Birds	Mountain white-crowned sparrow (<i>Zonotrichia leucophrys oriantha</i>)	Endogenous corticosterone manipulation and radiotelemetry used to examine interactions between locale climate, hypothalamo-pituitary-adrenal axis activation (via corticosterone manipulation), migration arrival timing, and pre-breeding behaviour at high elevation sites in the Sierra Nevada	Breuner and Hahn 2003
Fish	Brown trout (<i>Salmo trutta</i>)	Endogenous cortisol manipulation coupled with electronic tracking used to examine the timing and success of seaward migration in trout smolts	Midwood et al. 2014
Herptiles	Tiger salamanders (<i>Ambystoma tigrinum</i>)	Laboratory based experiments used thyroxine manipulations to investigate land- and water-drive behaviours; results explained known salamander migration behaviours in the wild	Duvall and Norris 1980
Invertebrates	Milkweed bug (<i>Oncopeltus fasciatus</i>)	Milkweed bug were given a ‘juvenile hormone’ mimic and flight-tested; more males and females receiving the treatment made long, presumably migratory, flights	Caldwell and Rankin 1972
Mammals	Belding’s ground squirrels (<i>Spermophilus beldingi</i>)	Used food provisioning experiments and testosterone manipulations to examine the dispersal behaviour of free-living juvenile male and female squirrels	Nunes et al. 1999

1228

Table 5. Sensory, navigation and orientation experiments

Taxa	Species	Experimental Approach	References
Birds	Homing pigeons (<i>Columba livia domestica</i>)	Pigeons were displaced and various sensory modalities (e.g., vision, smell, magnetic compass) were manipulated to assess return rates	Walcott 1996
Fish	Chinook salmon (<i>Oncorhynchus tshawytscha</i>)	Smolts were introduced into a large flume and exposed to LED lights in an effort to guide them to safe migratory paths	Hansen et al. 2018
Herptiles	Red-sided garter snakes (<i>Thamnophis sirtalis parietalis</i>)	Snakes were exposed to pheromones from conspecifics to investigate trailing behaviour; demonstrated that both sexes trail the other sex for mating	LeMaster et al. 2001
Invertebrates	Monarch butterfly (<i>Danaus plexippus</i>)	Monarchs were exposed to simulated magnetic fields (control, reversed and amagnetic); demonstrated that the species relies on magnetic cues to orient	Etheredge et al. 1999
Mammals	American black bears (<i>Ursus americanus</i>)	Used extensive tracking datasets from over 200 bears to test alternative hypotheses regarding migration path choice (e.g. landscape morphology, olfaction, instinct etc.); revealed social interactions mediated by chemical communication as likely driver	Noyce and Garshelis 2014

1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245

1246 **Table 6. Translocation experiments**

Taxa	Species	Experimental Approach	References
Birds	Gray catbird (<i>Dumetella carolinensis</i>)	Veteran migratory catbirds displaced from Illinois to New Jersey had similar migration orientation to controls except when their sense of smell was ablated, suggesting a role of smell interacting with experience in orientation	Holland et al. 2009
Fish	Roach (<i>Rutilus rutilus</i>)	Roach from populations with different migratory opportunities to a lake were translocated; translocated fish showed migration patterns more similar to local fish than their home location suggesting that partial migration is phenotypically plastic and triggered by local environmental cues	Skov et al. 2010
Herptiles	Sea turtle (<i>Chelonia mydas</i>)	Hatchlings were translocated from the east to the west coast; demonstrated that hatchlings use light and not innate preferences to find their way to sea	Carr and Ogren 1960
Invertebrates	Monarch (<i>Danaus plexippus</i>)	Monarch flight orientation was compared in their capture site to a site 2500 km away to show that they did not compensate for the change in position, suggesting they are not true navigators	Mouritsen et al. 2013
Mammals	Egyptian fruit bats (<i>Rousettus aegyptiacus</i>)	Even when translocated, Egyptian fruit bats have cognitive maps supported by visual cues that allow them to commute successful between foraging sites and home	Tsoar et al. 2011

1247
1248
1249
1250
1251
1252
1253
1254
1255

1256 **Table 7. Conservation-related experiments**

Taxa	Species	Experimental Approach	References
Birds	Western sandpipers (<i>Calidris mauri</i>)	Experimental oiling of feathers reduced escape flight performance and increased endurance flight energy costs	Maggini et al. 2017a, b
Fish	Sockeye salmon (<i>Oncorhynchus nerka</i>)	Salmon were implanted with accelerometer transmitters and released in different locations to quantify and contrast the impacts of dam water releases and fish passage in a fishway on anaerobic energy expenditure and migration success to inform dam operations*	Burnett et al. 2014
Herptiles	Three anuran species	Behavioural choice experiments were used to determine if several anurans had preferences for different tunnel types intended to serve as safe crossing sites at roads	Lesbarrères et al. 2004
Invertebrates	Rocky Mountain apollo butterfly (<i>Parnassius smintheus</i>)	Nectar abundance was manipulated in a meadow, which influenced the dispersal behaviour of butterflies assessed via mark-recapture, thus informing meadow restoration strategies	Matter and Roland 2002
Mammals	Wolf (<i>Canis lupus</i>)	Used trapped or dead wolves to demonstrate that they could migrate through densely populated areas with high road densities*	Gula et al. 2009

1257 * denotes studies where the approach was not purely experimental, but rather made use of the
 1258 presence of anthropogenic activities/infrastructure to evaluate its effects on migrations.
 1259
 1260
 1261
 1262
 1263
 1264
 1265
 1266
 1267

Figure 1. Conceptual framework of experimental biology in migration. Migration is the result of and/or affected by an organism’s physiology and the environment it lives in. Physiology (red) dictates an individual’s internal state, which can be impacted either directly or indirectly by the surrounding environment (green). This interaction between physiology and the environment affects individuals throughout the entirety of the ”migration timeline”; by determining when an individual is ready to initiate migration, how and where an individual will migrate, and finally why an individual ultimately migrates (yellow). The blue boxes represent the types of experiments that can be performed at different stages along the migration timeline to understand both the proximate and ultimate causes of migration.