

# Loricate choanoflagellates (Acanthoecida) from warm water seas. IX. Coronoeca gen. nov., Polyfibula Manton and spiny forms of Parvicorbicula Deflandre

Thomsen, Helge Abildhauge; Hara, Seiko; Østergaard, Jette Buch

Published in: European Journal of Protistology

Link to article, DOI: 10.1016/j.ejop.2021.125826

Publication date: 2021

Document Version Publisher's PDF, also known as Version of record

# Link back to DTU Orbit

Citation (APA):

Thomsen, H. A., Hara, S., & Østergaard, J. B. (2021). Loricate choanoflagellates (*Acanthoecida*) from warm water seas. IX. *Coronoeca* gen. nov., *Polyfibula* Manton and spiny forms of *Parvicorbicula* Deflandre. *European Journal of Protistology*, *81*, Article 125826. https://doi.org/10.1016/j.ejop.2021.125826

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.





Available online at www.sciencedirect.com



European Journal of **PROTISTOLOGY** 

European Journal of Protistology 81 (2021) 125826

www.elsevier.com/locate/ejop

# Loricate choanoflagellates (Acanthoecida) from warm water seas. IX. *Coronoeca* gen. nov., *Polyfibula* Manton and spiny forms of *Parvicorbicula* Deflandre

Helge Abildhauge Thomsen<sup>a,\*</sup>, Seiko Hara<sup>b</sup>, Jette Buch Østergaard<sup>c</sup>

<sup>a</sup> Technical University of Denmark, National Institute of Aquatic Resources (DTU Aqua), Kemitorvet, Bygning 201, DK 2800 Kgs, Lyngby, Denmark

<sup>b</sup> Ishikawauchi 368, Kijo-cho, Koyu-gun, Miyazaki 884-0104, Japan

<sup>c</sup> Nørrebrogade 52a 5th, 2200 Copenhagen N, Denmark

Received 3 May 2021; revised 6 July 2021; accepted in revised form 12 July 2021; Available online 21 July 2021

#### Abstract

The ambition to generate an overview of warm water loricate choanoflagellate biodiversity, based on a classic morphometric approach, is here completed by analyses of a range of tiny forms with anterior spines or projections and in most cases also a posterior pedicel. The warm water study complements previously obtained results from the more extensively studied temperate and polar regions of the world's oceans. It thus contributes to a significantly more balanced approach to global diversity patterns for these organisms. The current survey includes taxa such as *Polyfibula elatensis*, *Parvicorbicula pedicellata*, as well as a range of primarily undescribed and taxonomically challenging species, that are in an interim approach allocated to *Coronoeca* gen. nov. (*C. kosmaniae* sp. nov., *C. conicella* sp. nov., *C. superpositus* (Booth) comb. nov., *C. marchantii* sp. nov., *C. tongiae* sp. nov., and *C. patongiensis* sp. nov.). The analysis of warm water acanthoecid biodiversity has revealed in total 80 species from the six geographic regions sampled, corresponding to approximately 50% of all loricate species described. Nineteen species are previously undescribed forms. The Andaman Sea, Thailand, and West Australia are in a global context the most species-rich regions with 62 and 64 species respectively.

© 2021 The Authors. Published by Elsevier GmbH. This is an open access article under the CC BY license (http:// creativecommons.org/licenses/by/4.0/).

Keywords: Acanthoecida; Biogeography; Loricate choanoflagellates; Taxonomy; Ultrastructure

# Introduction

In an ongoing effort (Thomsen and Østergaard 2019a; Thomsen and Østergaard 2019b; Thomsen and Østergaard 2019c; Thomsen and Østergaard 2019d; Thomsen and Østergaard 2019e; Thomsen and Østergaard 2021; Thomsen et al. 2020a; Thomsen et al. 2020b) to provide a first comprehensive overview of warm water loricate choanoflagellate diversity, based on a traditional microscopical approach, we here primarily deal with a contingent of small, spiny and often pedicellated species. While a single species (i.e. *Polyfibula elatensis* (Thomsen, 1978) Manton in Manton and Bremer 1981), belongs to an unambiguously defined genus, the remaining taxa have unresolved phyloge-

https://doi.org/10.1016/j.ejop.2021.125826

0932-4739/© 2021 The Authors. Published by Elsevier GmbH. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

<sup>\*</sup>Corresponding author. E-mail address: hat@aqua.dtu.dk (H.A. Thomsen).

netic positions within the loricate choanoflagellate species matrix. It is in our opinion beneficial to future research, to be able to refer to a morphospecies by using a specific taxon name, rather than an ad hoc labelling of forms that will not to the same extent, systematically and rigorously, serve as a reference point for future observations of the specific taxon. We have therefore created the genus Coronoeca gen. nov., that can serve as an interim 'parking space' for these forms, while awaiting an in-depth analysis aided by state-of-the-art molecular tools. The generic type-species, C. kosmaniae sp. nov., has unique morphological features, that irrespective of the future fate of other species currently allocated to the genus (i.e. C. conicella sp. nov., C. superpositus (Booth, 1990) comb. nov., C. marchantii sp. nov., C. tongiae sp. nov., and C. patongiensis sp. nov.) will likely remain a phylogenetically valid unit. Parvicorbicula superpositus Booth, 1990, is transferred to Coronoeca (C. superpositus (Booth, 1990) comb. nov.) based on critical discrepancies, between this species and a core group species of Parvicorbicula as defined by Leadbeater (2015), relating to in particular mutual positions of transverse and longitudinal costae.

The current paper rounds off the survey of warm water loricate choanoflagellate diversity with a brief global perspective of the results obtained, and an inter-comparison of the geographic regions sampled.

## Material and methods

The material that constitutes the background for this and a series of papers on warm water acanthoecid choanoflagellates was collected over a period of 35 years. The geographic origin of samples is recorded in Fig. 1. See Thomsen and Østergaard (2019a) for information on each of the collection sites and sampling campaigns. In order to substantiate the analysis of *Coronoeca superpositus* (=*Parvicorbicula superpositus*) we have added material (Fig. 12) from Danish coastal waters. In order to further corroborate observations from our standard regions (Thomsen and Østergaard 2019a) we have appended to this paper wherever relevant, also historical data from Japan (Seto Inland Sea) to species descriptions, as well as the global analysis of loricate choanoflagellate biogeography (Figs. 16, 17).

The general protocol for processing water samples for light (LM) and (TEM/SEM) electron microscopy was according to Moestrup and Thomsen (1980) and Thomsen (1982). For details on sample processing, preparational issues and microscopes used see Thomsen and Østergaard (2019a). The Japanese material from the Shioya Coast (Seto Inland Sea, Kobe, Japan) comprises surface water samples collected from April 1979 to September 1982 (7.5–27.3 °C; 27–33 PSU). The processing of the Japanese samples for electron microscopy followed the basic principles outlined by Moestrup and Thomsen (1980).

The material examined here is dried, which means that the natural 3-D structures have collapsed to become 2-D structures, leading to a partial dislocation of costal strips, and an artefactual expansion of in particular the lorica width. While several structures can still be measured with confidence, e.g. lorica height and the length of spines and pedicels, it does imply that certain values such as lorica diameter, typically at the level of the transverse costa(e), cannot be measured directly but only calculated from measurements of the circumference. This approach was used in the species descriptions below.



**Fig. 1.** Map showing the approximate sampling sites for material reported here and MODIS sea surface temperatures (2003–2011 average). A circular dot refers to a single spot sampling, while a line or square indicates that samples were collected along extended transects. For further information see the materials and methods section in Thomsen and Østergaard (2019a).

Efforts were made when describing lorica features to follow the terminology standards that have developed in the course of dealing with these organisms; see e.g. Leadbeater (2015; loc. cit. chapter 4 and glossary p. 278) and Thomsen and Buck (1991).

The unfortunate existence of an electron microscope specific problem causing negatives to appear horizontally flipped, was discussed in Thomsen and Østergaard (2019d). The evidence in favour of dealing with this prob-

lem as a purely technical issue is overwhelming. Scans of the affected negatives have accordingly been flipped horizontally to produce non-inverted images. In the current publication this applies to: Fig. 2a; Fig. 3c; Fig. 5d-f; Fig. 6a; Fig. 8a, b; Fig. 14a, b; Fig. 15a, b, g.

The shade matrix (Fig. 17) is constructed using the PRI-MER software package (ver. 7) that has a built-in wizard that carries out the sequences of routines needed to perform the analysis (Clarke et al. 2014; Clarke and Gorley 2015).



**Fig. 2. a-m.** TEM (a, d-g) and LM whole mounts (b, c, h-m; phase contrast) of *Polyfibula elatensis* from the Gulf of California (a-c, e), the Andaman Sea, Thailand (d, f, g), the Caribbean Sea (h), the Sargasso Sea (i), West Australia (j, l, m), and the equatorial Pacific Ocean (k). (a, d) Complete cells; the arrow (a) points to a single facetted costal strip from the posterior transverse costa; (b, c, h-m) Selection of specimens from various geographic regions; (e-g) Detail (reversed printing) of distinctly (e) and less distinctly (f, g) facetted transverse costal strips; the arrows (f, g) point to median thickenings of costal strips. The scale bar (b) applies to all LM images.



**Fig. 3. a-j.** TEM (a-e) and LM whole mounts (f-j; phase contrast) of *Parvicorbicula pedicellata* from the Andaman Sea, Thailand (a, b, d, e), the Gulf of California (c, i, j), the equatorial Pacific (f), and West Australia (g, h). (a-c) Complete loricae; (d) Detail (reversed printing) of anterior transverse costa (from (b)) showing the exterior position of the longitudinal costae; (e) Detail of the mid-lorica transverse costa (from (a)) to highlight the interior position of the longitudinal costae; (f-j) Selection of specimens from various geographic regions. The scale bar (g) applies to all LM images.

#### **Results and discussion**

#### Polyfibula Manton in Manton and Bremer, 1981

Species of *Polyfibula* are characterized by small, barrelshaped loricae that comprise 6–8 longitudinal costae and three transverse costae. The transverse costae are all located inside the longitudinal costae. The individual species may have a posterior pedicel and short to medium-sized spines anteriorly. Most characteristically, the transverse costal strips that form the anterior and intermediate costal rings, are all furnished with centrally placed enlargements (facets) that anteriorly secures the fusion (T-joins) of longitudinal and transverse costal strips. A single costal strip from the posterior transverse costa may also be furnished with a median central facet (Fig. 2a; arrow). The protoplast is located posteriorly in the lorica. The genus is characterized by tectiform division (Fig. 2J; see also Manton and Bremer 1981; loc. cit. Figs. 2, 18, 31).

The genus currently comprises five formally described species (i.e. P. sphyrelata (Thomsen, 1973) Manton in Manton and Bremer, 1981; P. elatensis (Thomsen, 1978) Manton in Manton and Bremer, 1981; P. caudata (Leadbeater, 1975) Manton in Manton and Bremer, 1981; P. hexacostata Manton in Manton and Bremer, 1981, and P. stipitata Manton in Manton and Bremer, 1981). A comprehensive description of the genus is provided by Manton and Bremer (1981). Tong (1997a) advocated that P. caudata, P. hexacostata and P. stipitata should be relegated to synonyms of P. sphyrelata. The distinguishing lorica features are merely the number of longitudinal costae, and the presence or absence of a pedicel. Such lorica features are generally accepted to show some level of variability at the intra-specific level (P. elatensis (see below) typically has seven longitudinal costae, however occasionally eight;



Fig. 4. a-h. Drawings to approximate scale of *Coronoeca* spp. The drawings are based on dried material as observed in either LM or TEM, hence the protoplast position and dimensions are only indicative. (a) *C. kosmaniae*; (b, c) *C. conicella*; (d) *C. superpositus* (Thailand); (e) *C. superpositus* (Denmark); (f) *C. tongiae*; (g) *C. patongiensis*; (h) *C. marchantii*. The scale bar applies to all images.

*P. sphyrelata* usually has seven longitudinal costae (Thomsen 1973) but is also found with eight longitudinal costae (Thomsen et al. 1991)). It is thus likely that the species circumscriptions provided by Manton and Bremer (1981) are too narrow.

While three morphotypes (i.e. *P. caudata, P. hexa-costata, P. stipitata*) are sparingly recorded and almost exclusively from North Pacific and North Atlantic subarctic seas (Manton and Bremer 1981), the generic type species *P. sphyrelata* is more widely reported from in particular mid-latitude locations on both hemispheres (Thomsen 1973; Moestrup 1979; Hallegraeff 1983; Thomsen et al. 1991; Tong 1997a; Tong et al. 1998; Bérard-Therriault et al. 1999; Menezes 2005). The species also extends into brackish waters (Thomsen 1979; Ikävalko and Thomsen 1997), Antarctic circumpolar waters (Marchant 1985; Thomsen et al. 1990), and is further recorded from the Beagle Channel, Patagonia (Thomsen, unpublished).

# *P. elatensis* (Thomsen, 1978) Manton in Manton and Bremer, 1981 (Fig. 2)

*Polyfibula elatensis* is unambiguously distinguished from other species of *Polyfibula* based on the presence of prominent anterior spines (Fig. 2a, d), each one-half costal strip in length. The material examined here (Fig. 2; Table 1) is morphometrically in good agreement with the type material from the Gulf of Aqaba (Thomsen 1978). While the facets on transverse costal strips stand out clearly in material from the Gulf of California (Fig. 2a, e), these structures are, however, in material from the Andaman Sea (Fig. 2d, f, g) often very little pronounced, and appearing as just a minor median bulging of the transverse costal strip (Fig. 2f, g; arrows). Such specimens are hard to distinguish from *Parvicorbicula pedicellata* (Fig. 3; see further below). The vast majority of specimens examined have seven longitudinal costae. However, a few cells with eight longitudinal costae were also registered (e.g. Fig. 2h). While the type material (Thomsen 1978) was characterized by a posterior pedicel comprising just a single costal strip, it is evident that specimens examined here may occasionally, and most likely depending on environmental nuances, carry a considerably longer pedicel (Fig. 2j, 1) that is up to five costal strips long.

*Polyfibula elatensis* was frequently observed at all sites here examined (Table 2) with the exception of the East Mediterranean Sea. It is also known from Japanese coastal waters and has furthermore been recorded from the Gulf of Aqaba, Israel (Thomsen 1978), Galapagos and N. Alaska (Manton and Bremer 1981), California (Thomsen et al. 1991), Sydney, Australia (Tong et al. 1998), Pettaquamscutt River Estuary, Rhode Island, USA (Menezes 2005), Brazil (Bergesch et al. 2008), and also from Beagle Channel, Chile, and Friday Harbor, Puget Sound, USA (Thomsen, unpublished), and from New Zealand (Moestrup, unpublished).

#### Parvicorbicula Deflandre, 1960

Non-spiny warm water forms of *Parvicorbicula* were commented on in Thomsen et al. (2020a), and overall



Fig. 5. a-f. LM (a-c; phase contrast) and TEM (d-f; reversed printing (e)) whole mounts of *Coronoeca kosmaniae* sp. nov. from the Gulf of California, Mexico. (a-c) Selection of images to show the lorica as appearing from different angles; (d) Complete lorica (type specimen); (e, f) Detail of longitudinal costal strip swelling and crossing of transverse costa. The scale bar (a) applies to all LM images.

concerns with reference to a proper circumscription of the genus discussed at length. The generic type species (P. socialis (Meunier, 1910) Deflandre, 1960) forms, together with a small group of species of Parvicorbicula and Pleurasiga Schiller, 1925, i.e. Pa. quadricostata Throndsen, 1970; Pa. circularis Thomsen, 1976; Pa. corynocostata Thomsen, Garrison and Kosman, 1997; Pl. minima Throndsen, 1970; Pl. reynoldsii Throndsen, 1970, Pl. tricaudata Booth, 1990, what appears to constitute a phylogenetically coherent cluster of species, that in the future should be singled out as a separate genus. A feature that characterizes this group of species is first and foremost the fact that they have both the anterior and the mid-lorica transverse costae on the outer surface of the lorica. Additional shared morphological features are longitudinal costae (4-10) composed of three costal strips, two transverse costae separated by a single longitudinal costal strip, anterior T-joins between longitudinal and transverse costal strips, a flattened and slightly biforked termination of the tip of the anterior longitudinal costal strips, and the presence of a membrane, suspended from the free posterior tips of the middle layer of costal strips, and enveloping the posteriorly positioned protoplast. None of the species have anterior spines or projections, and a pedicel is only reported from *Pl. tricaudata*.

The genus *Parvicorbicula* additionally comprises eight morphologically diverse species, including *P. pedicellata* (see below), that in different ways do not conform to the morphological circumscription of the core group of taxa referred to above. Because of uncertainties with respect to selecting, within the loricate choanoflagellate species matrix at large, more appropriate taxonomic positions for these outlier species, Thomsen et al. (2020a) opted for maintain-



Fig. 6. a-d. *Coronoeca kosmaniae* sp. nov. TEM (a) and LM (b-d; phase contrast) whole mount micrographs from the Gulf of California, Mexico. (a) Detail of posterior lorica end to show the arched and often duplicated or triplicated (arrows) costal strips that terminate the lorica posterior end. Notice the only solitary longitudinal costal strip (arrowhead) and the attenuating single costal strip from the pedicel; (b-d) Selection of specimens to show the variability in appearance under the LM; notice (b) the marked accumulation of additional costal strips at the posterior lorica end, and (d) the vacated membrane that normally secures the protoplast into position. The scale bar (b) applies to all light micrographs.

ing status quo while awaiting in particular molecular data to support the decisions needed.

#### P. pedicellata Leadbeater, 1973 (Fig. 3)

There is overall good agreement between the *P. pedicellata* material examined here (Fig. 3; Table 1) and the type material from Bay of Kotor, Montenegro (Leadbeater 1973). The only noticeable difference is with respect to the number of longitudinal costae. Leadbeater (1973) reports that there are 6–9 longitudinal costae, whereas in our material there is consistently seven longitudinal costae. In *P. pedicellata* (Table 3) the anterior transverse costa is located inside the longitudinal costae (Fig. 3d) while the mid-lorica transverse costa (Fig. 3e) is exterior. The posterior transverse costa is interior.

The only *P. pedicellata* morphological feature that tiesup this species to the *Parvicorbicula* core group of species as circumscribed above, is the fact the mid-lorica transverse costa is located outside the longitudinal costae. Features that separate *P. pedicellata* from *Parvicorbicula* sensu stricto are the presence of three transverse costae and anterior spines, and also the fact that each longitudinal costa consists of four costal strips.

Parvicorbicula pedicellata shares features with Coronoeca spp. (see below), e.g. anterior spines and four costal strips in each longitudinal costa, but is still clearly differentiated because of the external position of the mid-lorica transverse costa. While awaiting supporting molecular evidence it appears most relevant to make no taxonomic changes with reference to *P. pedicellata*.

*Parvicorbicula pedicellata* is easily mistaken for *Poly-fibula elatensis* (Fig. 2). Decisive differences between these taxa, that can only be resolved using electron microscopy, are the presence in *Pol. elatensis* of facets in the middle of transverse costal strips, and the fact that all transverse costae are located inside the longitudinal costae in *Pol. elatensis* (Table 3). At the light microscopical level the best distinguishing feature is the overall lorica size (*Pa. pedicellata* is 25% smaller) and the shape of the lorica chamber, which is strictly conical in *Pa. pedicellata* (Fig. 3f-j) while anteriorly more parallel sided in *Pol. elatensis* (Fig. 2b, c, h-m).

*Parvicorbicula pedicellata* is here recorded from the Andaman Sea, Thailand, West Australia, the Equatorial Pacific Ocean, and the Gulf of California, Mexico (Table 2). The species is also observed in Japanese coastal samples and has previously been reported from Bay of Kotor, Montenegro, and Bay of Algiers (Leadbeater 1973), the Gulf of Aqaba, Israel (Thomsen 1978), equatorial Pacific (Vørs et al. 1995), Sydney, Australia (Tong et al. 1998), and the Pettaquamscutt River, Rhode Island, USA (Menezes 2005).

	Lorica height (excl. pedicel)	Diam. ant. transv. costa	Diam. 2nd transv. costa	Diam. 3rd transv. costa	Diam. 4th transv. costa	# longitu- dinal costae	<pre># long. costal strips (incl. projections/ excl. pedicel)</pre>	Projecting spines	Pedicel	Flagellum	n
Coronoeca conicella	$16.4 \pm 2.5$	$11.1 \pm 1.7$	6.8 ± 1.4			4-(5)	4	4-(5)	$12.5 \pm 6.7$	8.1 ± 1.0	63
C. kosmaniae	$25.5 \pm 1.2$	$15.5 \pm 0.6$	$15.2 \pm 0.7$			7	4	7	$7.9 \pm 0.4$		26
C. marchantii	$14.2 \pm 1.4$	$8.9\pm0.8$	$6.6 \pm 0.5$	$2.9 \pm 0.4$		(7)-8	4	(7)-8	$3.9 \pm 0.6$	$6.6 \pm 0.4$	27
C. patongiensis	15.2/16.5	12.4/15.9	8.0/10.2	3.3/		10	4	10			2
C. patongiensis	18.5	16.1	10.3	4.6		11	4	11		15.2	1
(Davis, pers. comm./Sargasso Sea)											
C. superpositus	13.7	10.0	7.3			8	4	7–8	4.2	7.9	2
(Thail.)											
<i>C. superpositus</i> (Denm.)	$18.8\pm1.9$	13.6 ± 1.1	$9.5\pm0.7$			10-(11)	4	4–8	$10.7\pm2.6$	$11.9 \pm 1.7$	19
C. tongiae	$14.6 \pm 0.9$	$12.0 \pm 1.1$	$9.9 \pm 0.5$	$7.4 \pm 0.8$	$3.1 \pm 0.5$	9–10	4	9–10			7
C. tongiae	13.5	11.0	10.4	8.4	3.4	9	4	9			1
(Leadbeater 1974; loc. cit. Pl. 2E)											
Parvicorbicula pedicellata	13.7 ± 1.1	$7.7\pm0.6$	6.1 ± 0.6	$2.9\pm0.3$		7	4	7	$3.6\pm0.5$	5.7 ± 1.8	15
Polyfibula elatensis	$18.0\pm1.7$	$9.8 \pm 1.4$	$9.0\pm1.0$	$3.9\pm0.4$		8	4	8	$9.6\pm 6.0$	$12.3\pm2.0$	26

Table 1.	Summary	of lorica	morphometric	features.

	Andaman Sea, Thailand	West Australia	Sargasso Sea	Caribbean Sea	Equatorial Pacific Ocean	Gulf of California, Mexico	Mediterranean Sea, Alexandria	Reference
Acanthocorbis apoda	х							Thomsen and
								Østergaard 2019c
A. camarensis	Х							Thomsen and
								Østergaard 2019c
A. campanula	Х	х	Х					Thomsen and
/ · · · · · · ·								Østergaard 2019c
A. conicella*		х	Х			Х	Х	Thomsen and
1 aladialla*								Østergaard 2019c
A. glaatella*	X	X	X	х	X			Antoneound 2010a
1 haunabiana	V	V	v					Themson and
Α. παυτακιαπα	х	х	х					Astergaard 2010c
Anheloecion articulatum	v	v					v	Thomsen and
Apheioecion articulatum	л	л					л	Østergaard 2019b
A eapacia*	x	x			x			Thomsen and
in eqpacta								Østergaard 2019b
A. pentacanthum	х	х			Х	х	Х	Thomsen and
1								Østergaard 2019b
A. quadrispinum	х	х	х	х	Х		Х	Thomsen and
								Østergaard 2019b
Bicosta minor (form A)	Х	х		х	Х			Thomsen and
								Østergaard 2019b
B. spinifera						х		Thomsen and
								Østergaard 2019b
Calliacantha magna*	Х	х		х	Х			Thomsen and
								Østergaard 2019b
C. natans						Х		Thomsen and
								Østergaard 2019b
C. simplex	Х	х	Х	х	Х	Х	Х	Thomsen and
								Østergaard 2019b
Calotheca alata	Х	х						Thomsen et al. 2020b
Campyloacantha	Х	х	х	х	Х			Thomsen and
imbricata								Østergaard 2019b
C. spinifera	X	x	x			X	X	I nomsen and
<b>C</b> :*								Distergaard 20196
Contoeca boonruangit*	Х	х		х	X			Antonesen and
Coronagea conicella*	v	v	v	v	v	v	v	This paper
C kosmaniae*	Λ	Λ	л	Λ	Λ	A V	Λ	This paper
C. nosmuniue C. marchantii*	v	v	v	v	v	A V		This paper
C. murchunill	Λ	Λ	Λ	Λ	Λ	Λ		rins paper

Table 2. Occurrence pattern of species discussed here and in Thomsen and Østergaard (2019a, 2019b, 2019c, 2019d, 2019e, 2021) and Thomsen et al. (2020a, 2020b; 2021). New species described are marked with \*.

C. patongiensis*	Х	х						This paper
C. superpositus	Х							This paper
C, tongiae*	Х	х				Х	Х	This paper
Cosmoeca ceratophora	Х	х	х	х	Х	Х	Х	Thomsen and
								Østergaard 2019d
C. norvegica	Х	х	х	х	Х	Х	Х	Thomsen and
								Østergaard 2019d
C. phuketensis	Х	х	х	х	Х		Х	Thomsen and
								Østergaard 2019d
C. subulata	Х	х			Х			Thomsen and
								Østergaard 2019d
<i>C.ventricosa</i> (incl. form A)	Х	х	х	х	Х	Х	Х	Thomsen and
								Østergaard 2019d
C. ventricosa (form B)	Х	х	х		Х	Х	Х	Thomsen and
								Østergaard 2019d
C. ventricosa (form C)	Х	х	х	х	Х			Thomsen and
								Østergaard 2019d
Crinolina aperta	Х	х	х	х	Х		Х	Thomsen and
								Østergaard 2021a
C. isefiordensis	Х	х				Х		Thomsen and
								Østergaard 2021a
Crucispina cruciformis	Х	х	х			Х	Х	Thomsen and
								Østergaard 2019b
Diaphanoeca cylindrica	Х	х						Thomsen and
								Østergaard 2021
D. grandis	Х	х				Х	Х	Thomsen and
								Østergaard 2021
D. multiannulata		Х						Thomsen and
								Østergaard 2021
D. pedicellata		Х			Х	Х		Thomsen and
								Østergaard 2021
D. pedicellata cfr.		х	х		Х		Х	Thomsen and
								Østergaard 2021
D. pseudoundulata*	Х	х			Х	Х		Thomsen and
D								Østergaard 2021
D. sargassoensis*			Х					Thomsen and
D : 1:C								Østergaard 2021
D. spiralifurca	X	x						Thomsen and
D 41								Ustergaard 2021
D. inronasenii*	Х	Х	Х			Х		I nomsen and
N								The second secon
wannoeca mexicana*		х				Х		Anterseend 2010-
								Ostergaard 2019a

N. minuta	х	Х	Х	Х	Х	х	х	Thomsen and Østergaard 2019a
<i>N</i> minuta (form A)	x				x			Thomsen and
	А				A			Østergaard 2019a
Parvicorbicula		х	Х	х		х	Х	Thomsen et al. 2020a
circularis								
P. pedicellata	х	х			Х	х		This paper
P. socialis		х						Thomsen et al. 2020a
Pleurasiga	х	х	х	х	Х	х	Х	Thomsen et al. 2020a
echinocostata								
P. echinocostata form A	х	х						Thomsen et al. 2020a
<i>P. echinocostata</i> form B	х				х			Thomsen et al. 2020a
P. minima	х	х	х	х	Х	х		Thomsen et al. 2020a
P. minutissima*	х	х	х	х	Х	х	Х	Thomsen et al. 2020a
P. quadrangiella*	х	х			х			Thomsen et al. 2020a
P. revnoldsii		х	х		х	х		Thomsen et al. 2020a
P. tricaudata						х		Thomsen et al. 2020a
Polyfibula elatensis	x	x	x	x	x	x		This paper
Saroeca attenuata			х		(x)			Thomsen and
								Østergaard 2019b
S. paucicostata	х	х	х			х	Х	Thomsen and
1								Østergaard 2019b
Savillea sp.			х					This paper
Stephanacantha					Х	х		Thomsen et al. 2020b
campaniformis								
S. dichotoma	х	х	х	х	Х	х	Х	Thomsen et al. 2020b
S. formosa	х	х	х	х	Х	х	Х	Thomsen et al. 2020b
S. oceanica*		х	х					Thomsen et al. 2020b
S. parvula	х	х	х	х	Х			Thomsen et al. 2020b
S. zigzag	х	х	х	х	Х	х	Х	Thomsen et al. 2020b
Stephanoeca	х	х						Thomsen and
andemanica*								Østergaard 2019c
S. apheles	х	х						Thomsen and
-								Østergaard 2019c
S. broomia*	х	х						Thomsen and
								Østergaard 2019c
S. diplocostata var.		х						Thomsen and
paucicostata								Østergaard 2019c
S. naja*	х	х	Х		Х	Х		Thomsen and
-								Østergaard 2019c
S. supracostata	х							Thomsen and
-								Østergaard 2019c
S. urnula							Х	This paper

<b>Syndetophyllum</b> pulchellum	Х	Х				Х	х	Thomsen et al. 2020b
Thomsenella acuta	х	х	х	х	х		х	Thomsen and Østergaard 2019e
T. cercophora	х	х			х		X	Thomsen and Østergaard 2019e
T. infundibuliformis			Х		Х	Х	X	Thomsen and Østergaard 2019e
T. perforata	x	х	Х	х	x	x	X	Thomsen and Østergaard 2019e

Table 3. Summary of lorica features. I: Transverse costa inside longitudinal costae'; (I): Transverse costa outside anterior tip of the penultimate longitudinal costal strip, but inside the anterior projections; O: Transverse costa outside longitudinal costae; T: T-joins anteriorly; E: E-joins anteriorly.

	Anterior transverse costa	2nd transverse costa	3rd transverse costa	4th transverse costa	Anterior costal strip joins
Coronoeca conicella	(I)	Ι			Т
C. kosmaniae	(I)	Ι			Т
C. marchantii	Ι	Ι	Ι		Е
C. patongiensis	0	Ι	Ι		Е
C. superpositus (Thail.)	(I)	Ι			Т
C. superpositus (Denm.)	(I)	Ι			Т
C. tongiae	Ι	Ι	Ι	Ι	Т
Parvicorbicula pedicellata	Ι	0	Ι		Т
Polyfibula elatensis	Ι	Ι	Ι		Т

12

#### Coronoeca gen. nov.

Although the species dealt with below under the heading '*Coronoeca*' do share certain basic morphological features, we are painfully aware of that a future in depth analyses of these taxa, including also a multifaceted molecular probing, will undoubtedly unveil a need for a redistribution of these within a number of separate genera. Realizing that none of these forms can be smoothly and convincingly allocated to an existing genus, and further in an attempt to avoid establishing too many monotypic genera created on an uncertain knowledge basis, we here opt for creating a pro forma genus that can at least temporarily harbour these forms (Fig. 4). Choosing *C. kosmaniae* as the generic type species reflects the fact, that this species has a large range of unique morphological features, that makes it a good long-term candidate for genus survival.

**Diagnosis**: Lorica conical or barrel-shaped and comprising 4– 10 longitudinal costae and 2–4 transverse costae. There are anterior projections or spines, and the lorica may have a compound pedicel, a simple pedicel or no pedicel. Each longitudinal costa comprises four costal strips (including anterior projection). Anterior junctions between longitudinal and transverse costal strips can be either T-junctions or E-junctions. All costal strips are narrow rods. Minor morphological modifications can apply to anterior projections or costal strips from the pedicel. The longitudinal costae are external relative to the mid-lorica transverse costa. Tectiform division presumed but not yet verified in the type species.

Generic type species: Coronoeca kosmaniae sp. nov.

**Etymology**: Genus name derived from 'corona' (Latin) meaning 'crown' and referring to the prominent anterior spines or projections, and 'oicos' (Greek) meaning 'house'.

#### Coronoeca kosmaniae sp. nov. (Fig. 4a, 5, 6)

**Diagnosis**: Lorica barrel- or bell-shaped (lorica height:  $25.5 \pm$ 1.2 µm incl. anterior projections and excl. posterior pedicel), constructed of rod-shaped costal strips, and with modified attenuating anterior projections (seven) that have a proximal swelling ca. 0.2 µm from the posterior tip (Fig. 5e, f). Anterior transverse costal strips attach to longitudinal costae at the level of the swellings creating T-junctions. The lorica chamber is composed of seven longitudinal costae and two transverse costae (Fig. 5a-d). The anterior transverse costa rests in a fork created by the overlapping longitudinal costal elements (Fig. 5e, f) and is exterior relative to the tip of the penultimate longitudinal costal strip, but interiorly located relative to the anterior projection. The mid-lorica transverse costa is inside the longitudinal costae. A longitudinal costa is composed of four costal strips (including the anterior projections). Longitudinal costae converge posteriorly, with the posteriormost longitudinal costal strips being noticeably curved and sometimes duplicated. Transverse costae each comprising seven costal strips, and of approximately the same size (anterior ring:  $15.5 \pm 0.6 \ \mu m/mi$ 

d-lorica ring:  $15.2 \pm 0.7 \,\mu$ m). Lorica with a compound pedicel, usually with 1–3 diverging costal strips. Both anterior and posterior projections tapering, and ca. 1.0  $\mu$ m longer than all other costal strips ( $7.2 \pm 0.6 \,\mu$ m versus  $6.2 \pm 0.6 \,\mu$ m). The protoplast is located at the posterior end of the lorica chamber, within a suspensory membrane (Figs. 5c, d; 6a, d). Tectiform division presumed but not verified from an accumulation of newly formed costal strips in the collar region.

**Holotype**: The specimen illustrated in Fig. 5d of the present work is fixed as holotype (ICZN 1999, Article 73.1.4).

**Type locality**: Surface water sample (ca. 18 °C/35 PSU) from a near coastal site at Bahia de los Ángeles, Gulf of California, Mexico, collected 7 January 1990.

**Etymology**: The species-group name is chosen to acknowledge significant contributions to the microscopical examination of the Gulf of California material made by Carol Kosman. Carol was during the period 1991–1992 working at the Univ. of Copenhagen as a Fulbright Research Scholar.

The spacious *C. kosmaniae* lorica has an uncomplicated and fairly standard lorica construction, but is nevertheless uniquely defined, in particular because of the costal strip swellings (Fig. 5e, f) at the base of the anterior projections, and at a point where the anterior transverse costa crosses. These localized swellings are reminiscent of the median facets on *Polyfibula* transverse costal strips (Manton and Bremer 1981; see also Fig. 2e) and also the less confined swellings in the posterior one third of anterior longitudinal costal strips in *Saroeca attenuata* (Thomsen 1979).

There is in C. kosmaniae a variable degree of costal strip duplication at the posterior lorica end. The TEM image (Fig. 6a) thus shows a lorica in which all posterior longitudinal costal strips are duplicated (or even triplicated; Fig. 6a arrows) except for one costal strip (Fig. 6a; arrowhead). Duplications are also easily detected from LM (Fig. 6b). The anterior projections and costal strips from the compound pedicel are basically similar and deviate from midlorica longitudinal costal strips by being larger and attenuating in width from one end to the other (Fig. 5d). This precludes both the possibility that the posterior lorica end costal strip duplication could in part be explained by a displacement of costal strip elements from the compound pedicel, and likewise that the compound pedicel could be interpreted as a simple displacement of costal strips from the posterior lorica end duplications. A compound pedicel is a feature that is otherwise shared among several species, e.g. Didymoeca tricyclica (Bergesh et al. 2008), Pleurasiga tricaudata (Booth 1990; Thomsen et al. 2020a), Thomsenella cercophora (Thomsen and Boonruang 1983; Thomsen and Østergaard 2019e), and Syndetophyllum pulchellum (Thomsen and Moestrup 1983; Thomsen et al. 2020b).

The costal strip pattern at the level of the anterior transverse costa, where the transverse ring rests in a fork created by an overlap between neighbouring longitudinal costal strips (the ring is external relative to the penultimate longitudinal costal strip, however, overlaid externally by the anterior projection), is basically similar to what is observed in species of *Diaphanoeca*, e.g. *Diaphanoeca grandis* (Leadbeater 2015; loc. cit. page 80 and Figs 4.29–4.33) and *Conion groenlandicum* (Thomsen 1982), and also mirrored in species discussed below (*C. conicella*, *C. superpositus*, and *C. marchantii*).

*Coronoeca kosmaniae* is superficially reminiscent of *Diaphanoeca throndsenii* (Thomsen and Østergaard 2021). However, in this species there is consistently eight longitudinal costae, anterior E-junctions between longitudinal and transverse costal elements, no swelling at the posterior end of a projection, a pedicel that comprises a single costal strip, and no indication of costal strip duplications at the posterior lorica end.

The combination of lorica features in *C. kosmaniae* as accounted for above calls for the description of a new genus.

*Coronoeca kosmaniae* was frequently observed in samples from the Gulf of California, Mexico, and has also been observed in samples from the Seto Inland Sea, Japan. The Japanese specimens are morphometrically exactly similar to the Mexican type material.

#### Coronoeca conicella sp. nov. (Figs. 4b, c, 7–9)

**Diagnosis**: Lorica funnel-shaped (lorica height:  $16.4 \pm 2.5 \ \mu m$  incl. anterior projections and excl. posterior pedicel) and con-



**Fig. 7. a-I.** Coronoeca conicella sp. nov. TEM (a-c) and LM (d-f; DIC/g-l; phase contrast) whole mounts from the Andaman Sea, Thailand (a-f) and West Australia (g-l). (a-c) Selected specimens to show the variability encountered including specimens with extended (a) and short pedicel (c). (d-f) Andaman Sea images of long (d) and short (e, f) pedicellate forms. (g-l) The variety encountered in samples from West Australia. The scale bar (g) applies to all LM images.

structed of rod-shaped costal strips organized in 4-(5) longitudinal costae and two transverse costae. There are anterior attenuating projections and a pedicel of highly variable length. The anterior transverse costa (diam.  $11.1 \pm 1.7 \mu$ m) comprises twice as many costal strips as there are longitudinal costae (Tjunctions). The mid-lorica transverse costa (diam.  $6.8 \pm 1.4 \mu$ m) comprises (4)-5-(6) costal strips. The anterior transverse costa rests in a fork created by overlapping longitudinal costal elements (Fig. 8b). It is exterior relative to the tip of the penultimate longitudinal costal strip, but interiorly located relative to the anterior projections (Fig. 7a, Fig. 8b). The mid-lorica transverse costa is inside the longitudinal costae. A longitudinal costa is composed of four costal strips (including the anterior projections). The protoplast is located at the posterior end of the lorica chamber, within a suspensory membrane (Fig. 7a, b, j; Fig. 8f, h). Tectiform division evidenced from an accumulation of newly formed costal strips in the collar region (Fig. 9b).

**Holotype**: The specimen illustrated in Fig. 7a of the present work is fixed as holotype (ICZN 1999, Article 73.1.4).

**Type locality**: Surface water sample (ca. 28 °C/35 PSU/2 m depth) from a near coastal site at Ao Patong, Phuket Island, Thailand (7° 53'56" N/98° 17'36" E), collected 8 September 1981.

**Etymology**: The species-group name refers to the perfect conical shape of the lorica; -ella = diminutive.

Ignoring the anterior projections and the pedicel, *C. conicella* does superficially appear similar to e.g. *Parvicorbicula circularis* (Thomsen 1976). Shared features include



**Fig. 8. a-k.** *Coronoeca conicella* sp. nov. TEM (a, b) and LM (c-i; phase contrast/j, k; DIC) whole mounts from the Gulf of California, Mexico (a-e), the Sargasso Sea (f), the Caribbean Sea (g), the equatorial Pacific Ocean (h, i), and the eastern Mediterranean Sea (j, k). (a) Complete lorica without protoplast; notice five longitudinal costae; (b) High magnification (reverse printing) to show anterior projections and connections between longitudinal and transverse costal elements; (c-k) Specimens from various sources selected to show the morphological variability encountered; notice (e-f) five longitudinal costae. The scale bar (e) applies to all light micrographs.



Fig. 9. a,b. *Coronoeca conicella* sp. nov. TEM whole mounts from the Seto Inland Sea, Japan. (a) Complete cell. Notice that the lorica has five longitudinal costae; (b) Dividing cell that verifies tectiform division in this species.

four longitudinal costae each comprising three costal strips, and two transverse costae where the anterior costa has eight costal strips with every second strip forming T-junctions with the longitudinal costae, while the posterior transverse costa is formed by 5–6 costal strips. The decisive factor when considering a possible phylogenetic linkage between the two species is, however, that while the mid-lorica transverse costa is located outside the longitudinal costae in *P. circularis* it is inside in *C. conicella* (Table 3). This is a fundamental difference that is intimately linked to processes being part of the lorica assembly following cell division. Despite a superficial resemblance, *P. circularis* and *C. conicella* must therefore be kept apart at the supra-generic level.

In *C. conicella* the anterior transverse costa is external relative to the tip of the penultimate longitudinal costal strip, however, overlaid by the anterior projections (Figs. 7b, 8a, b). A similar set up is observed in *C. kosmaniae* (see above) and *C. superpositus* (see below).

Coronoeca conicella is, as defined here, a highly variable taxon, and has accordingly been lavishly illustrated from various geographic regions. We have at all sites examined observed two morphotypes of the species, i.e. forms with a short pedicel comprising just a single costal strip (e.g. Fig. 7c, k; 8j), and others that have an extended pedicel comprising multiple costal strips (e.g. Fig. 7d, g, j; 8f). There is an obvious linear linkage (Fig. 10A;  $r^2 = 0.77$ ) between lorica height and length of pedicel in the material examined (>60 specimens), as well as a non-patchy distribution, indicating that this element of variability is likely an inherent characteristic of the species. It can be further noticed from Fig. 10A that there are no obvious regional differences, although specimens from the Gulf of California are much less variable in size than e.g. specimens from West Australia.

Somewhat surprisingly it has been noticed that approximately 10% of all loricae examined have five rather than four longitudinal costae (Fig. 7g; 8a, f). The Gulf of California (Mexico) has a marked overrepresentation of such forms with five longitudinal costae (Fig. 10B). In material from Japanese coastal sites the only morphotype observed has five longitudinal costae (Fig. 9a, 10B).

All costal strips are narrow rods and quasi-identical throughout the lorica. However, the anterior projections do differ in being anteriorly attenuating and pointed and markedly thickened posteriorly (Fig. 8b). The maximum diameter of the rod is at the crossing of the transverse costa. Costal strips from extended pedicels generally appear somewhat more compact than costal strips elsewhere.

*Coronoeca conicella* was observed at all sites sampled (Table 2) and the form with an extended pedicel also occurred in Japanese coastal samples. The species has previously been observed (specimens with five longitudinal costae) in samples from the Galapagos Islands, Peru (I. Manton (†), pers. com.) and Sydney, Australia (Tong et al. 1998, loc. cit. Fig. 21; referred to as *Parvicorbicula superpositus*).

# Coronoeca superpositus (Booth, 1990) comb. nov. (Figs. 4d, e, 11, 12)

#### Basionym: Parvicorbicula superpositus Booth, 1990

When first described (Booth 1990) this species was primarily discussed with reference to *Parvicorbicula socialis* (*Meunier*, 1910) Deflandre, 1960, with which it does share a striking resemblance (e.g. ten longitudinal costae, two transverse costae separated by a single longitudinal costal strip, anterior T-junctions), when ignoring the *C. superpositus* anterior projections and pedicel. However, a close inspection of the material from Danish coastal waters that have been included here to consolidate the *C. superpositus* 



**Fig. 10.** Diagrams (A) to show the relationship between lorica height and length of pedicel in *C. conicella*, and site-specific variability, and (B) the site-specific occurrence of loricae with five rather than four longitudinal costae.

species circumscription, clearly unveils that the mid lorica transverse costa is located inside the longitudinal costae (Table 3), while it is clearly exterior in *Parvicorbicula socialis* (see e.g. Buck 1981, loc. cit. Figs. 9, 11). This is perhaps even more evident when consulting the excellent SEM images of *C. superpositus* published by Bérard-Therriault et al. 1999; loc. cit. Pl. 142a, b). An exterior versus interior placement of the mid-lorica transverse costa is, as previously emphasized, a fundamental distinguishing characteristic, which is coupled to intrinsic processes of lorica assembly, and hence being a feature only applicable at the supra-generic level. The anterior transverse costa is exterior relative to the tips of the longitudinal costal strips from the tier below, however externally overlaid by the

anterior projections (Fig. 11a, 12c). The costal strip configuration at the level of the anterior transverse costa is thus similar to what is observed in both *C. kosmaniae* and *C. conicella*. The actual morphological elaboration of the anterior projection is also identical in the species *C. kosmaniae*, *C. conicella* and *C. superpositus*. The projection is anteriorly sharply attenuated while bulging posteriorly (Fig. 12c). The maximum width is attained at exactly the level of the anterior transverse costa.

Specimens resembling *C. superpositus* (Fig. 11) are only sparingly observed in our warm water samples (Andaman Sea, Thailand). While the mid-lorica transverse costa is also in these specimens internal relative to the longitudinal costae, they deviate from *C. superpositus* sensu stricto by



Fig. 11. a-c. *Coronoeca superpositus* TEM (a, b) and LM (c; phase contrast) whole mounts from the Andaman Sea, Thailand. (a, b) Complete loricae (a; reverse printing) both with eight longitudinal costae and a similar number of projections; the interior location of the mid-lorica transverse costa is evident (a); (c) Empty lorica.

having fewer longitudinal costae (eight) and by the fact that all and not just a sub-set of longitudinal costae are here terminated by anterior projections (Fig. 11a, b). For the time being the variability encountered in Thailand specimens, when compared with *C. superpositus* sensu stricto (Fig. 12), is considered to represent intraspecific variability. The Thailand morphotype of *Coronoeca superpositus* was sparingly observed also in samples from Japanese coastal waters.

Coronoeca superpositus sensu stricto has previously been reported from the subarctic North Pacific Ocean (Booth 1990; type locality), Kaikoura, New Zealand (Moestrup 1979, loc. cit. Fig. 46; referred to as Parvicorbicula sp.), Kilsfjorden, Norway (Espeland and Throndsen 1986, loc. cit. Fig. 36; referred to as Parvicorbicula socialis aff.), California, USA (Thomsen et al. 1991), Southampton, UK (Tong 1997a), Taiwan (Hara et al. 1997), Sydney, Australia (Tong et al. 1998; only loc. cit. Fig. 2k), St. Lawrence Estuary, Canada (Bérard-Therriault et al. 1999), and Danish coastal waters (Thomsen et al. 2016). Soto-Liebe et al. (2007) reports C. superpositus from the Chilean coast. However, the resolution of the image provided (loc. cit. Fig. 1d) does not allow for a positive identification. The overall picture of the distribution pattern is thus that C. superpositus sensu stricto has a preference for the temperate climate zone avoiding high and low latitude habitats.

#### Coronoeca marchantii sp. nov. (Figs. 4h, 13)

**Diagnosis**: Lorica funnel-shaped (lorica height:  $14.2 \pm 1.4 \mu m$  incl. anterior projections and excl. posterior pedicel) and con-

structed of rod-shaped costal strips organized in (7)-8 longitudinal costae and three transverse costae separated by single tiers of longitudinal costal strips. There are anterior attenuating projections and a pedicel comprising a single costal strip. A longitudinal costa is composed of four costal strips (including the anterior projections). The anterior transverse costa (diam. 8.9  $\pm$  0.8 µm) comprises as many costal strips as there are longitudinal costae (E-junctions). The mid-lorica transverse costa (diam.  $6.6 \pm 0.5 \,\mu\text{m}$ ) has the same number of costal strips with much larger overlap between neighbouring strips. The posterior transverse costa (diam.  $2.9 \pm 0.4 \mu m$ ) comprises 4–5 costal strips. All transverse costae are internal relative to the longitudinal costae. The anterior transverse costa rests in a fork created by overlapping longitudinal costal elements. It is exterior relative to the tip of the penultimate longitudinal costal strip, but interiorly located relative to the anterior projections (Fig. 13a). The protoplast is located at the posterior end of the lorica chamber, within a suspensory membrane (Fig. 13a). Tectiform division evidenced from the accumulation of costal strips in the collar region (Fig. 13e, h).

**Holotype**: The specimen illustrated in Fig. 13a of the present work is fixed as holotype (ICZN 1999, Article 73.1.4).

**Type locality**: Surface water sample (ca. 28 °C/35 PSU/2 m depth) from a near coastal site at Ao Patong, Phuket Island, Thailand ( $7^{\circ} 53'56'' \text{ N/98}^{\circ} 17'36'' \text{ E}$ ), collected 8 September 1981.

**Etymology**: The species-group name acknowledges the great contributions that Dr. Harvey Marchant (Australian Antarctic Division, Hobart) has made to loricate choanoflagellate research.

There is little to add to the formal description of *C*. *marchantii* which has a well-defined costal strip configuration of its tiny conical lorica. In most cases this species can be unambiguously identified also from LM of dried



**Fig. 12. a-g.** *Coronoeca superpositus* TEM (a-c) and LM (d-g; phase contrast) whole mounts of specimens from Danish coastal waters (Kulhuse) selected to supplement the limited image gallery available of this taxon. (**a**, **b**) Well preserved loricae with four and six anterior projections. (**c**) Sharply pointed anterior projection and detail of the connection between transverse and longitudinal costal strips (reverse printing). (**d-g**) LM images showing the variability typically encountered. The scale bar (d) applies to all light micrographs.

specimens (e.g. Fig. 13c-e, j, k). A few slightly deviant forms were encountered in samples from West Australia and the Sargasso Sea (Fig. 13f-i). While these specimens obviously have eight longitudinal costae and three transverse costae, the difference noted is a slight pushing forward of the anterior transverse costa. Arrows (Fig. 13f-i) indicate where to find the transition between the anterior projections and the first tier of longitudinal costal strips. The overall appearance of these forms is reminiscent of Parvicorbicula pedicellata (Fig. 3). However, the pattern of crossing between longitudinal costae and costal strips from the anterior transverse costa (T- versus E-junctions) and the mere number of longitudinal costae (seven versus eight) adequately separates the two forms. For now, it thus appears relevant to treat this small deviation in lorica morphology as merely C. marchantii intraspecific variability. However, the validity of this approach will need to be tested using molecular tools.

*Coronoeca marchantii* was observed at all sites sampled with the exception of the eastern Mediterranean Sea (Table 2). It is additionally observed from the Galapagos Islands, Peru (I. Manton (†), pers. comm.).

#### Coronoeca tongiae sp. nov. (Figs. 4f, 14a-g)

**Diagnosis:** Lorica funnel-shaped (lorica height:  $14.6 \pm 0.9 \mu m$ ) and constructed of rod-shaped costal strips organized in 9–10 longitudinal costae and four transverse costae. There are anterior spines up to one half longitudinal costal strip in length. A longitudinal costa is composed of four costal strips. The mid-lorica and posterior transverse costae are separated by single tiers of longitudinal costal strips. The three anterior transverse costae (diam.  $12.0 \pm 1.1 \mu m$ ,  $9.9 \pm 0.5 \mu m$ , and 7.  $4 \pm 0.8 \mu m$ ) comprise as many costal strips as there are longi-



**Fig. 13. a-k**. *Coronoeca marchantii* sp. nov. TEM (a), SEM (b) and LM (c-k; phase contrast) whole mounts from the Andaman Sea, Thailand (a-b), West Australia (c-e; g, h), the Sargasso Sea (f), the Caribbean Sea (i), the equatorial Pacific Ocean (j), and the Gulf of California, Mexico (k). (a) Type specimen; notice the membrane that secures the protoplast in position; (b) Empty lorica which clearly shows the interior location of all three transverse costae; (c-e, j, k) Selected specimens to show the variability encountered across all regions sampled; (f-i) Deviant forms where the anterior transverse costa is pushed forward. The arrows point to the overlap between anterior longitudinal costal strips. The scale bar (e) applies to all LM images.

tudinal costae. While longitudinal costal strips attach to the middle of costal strips from the anterior ring, transverse costal strips from the mid-lorica costae span the width between neighbouring longitudinal costae. The reduction in diameter is occasioned by increasing overlaps between neighbouring transverse costal strips. The posterior transverse costa (diam.  $3.1 \pm 0.5 \mu$  m) comprises 4–5 costal strips. All transverse costae are internal relative to the longitudinal costae. The protoplast is located at the posterior end of the lorica chamber, within a suspensory membrane. Tectiform division assumed but not yet confirmed. **Holotype**: The specimen illustrated in Fig. 14a of the present work is fixed as holotype (ICZN 1999, Article 73.1.4).

**Type locality**: Water sample (ca. 28 °C/35 PSU/45 m depth) from a continental shelf station (#31) in the Andaman Sea, Thailand (7° 24' N/97° 27' E), collected March 1996. **Etymology**: The species-group name acknowledges the great contributions that Dr. Susan M. Tong has made to loricate choanoflagellate research.

This species was first observed by Leadbeater (1974; loc. cit. Pl. 2E/referred to as *Parvicorbicula* sp.) in samples from the Mediterranean (Split, Croatia and the Bay of Algiers). The single specimen illustrated is unmistakably morphome-



**Fig. 14. a-h.** *Coronoeca tongiae* sp. nov. (a-g) and *C. patongiensis* sp. nov. (h, i) TEM (a, b, d-f, i) and LM (c, g, h; phase contrast) whole mounts from the Andaman Sea, Thailand (a, b, d, i), West Australia (c, h), the eastern Mediterranean Sea, Alexandria (e), the Seto Inland Sea, Japan (f), and the Gulf of California, Mexico (g). (a) Type specimen (reverse printing). (c, g, h) Images selected to demonstrate that species specific details can be resolved using LM; (b, d-f) Specimens selected to document the consistency in lorica construction; (i) Type specimen (reverse printing). The scale bar (h) also applies to (c).

trically identical to the material examined here (Table 1). The generic affiliation of this species was uncertain when first examined by Leadbeater (1974), and it has remained so ever since. It therefore seems appropriate at this stage, and while awaiting a molecular based unravelling of its phylogenetic connections, to at least formally describe this

form, and as an interim solution allocate it to the 'pro forma' genus *Coronoeca* described here. The possibility initially indicated by Leadbeater (1974) of a certain resemblance between this form and species of *Parvicorbicula* is no longer valid considering the recent focus on regrouping selected species of *Pleurasiga* and core group members of



**Fig. 15. a-h.** TEM (a, b, g, h) and LM (c-f; phase contrast) whole mounts of miscellaneous species from the equatorial Pacific Ocean (a, b), the eastern Mediterranean Sea, Alexandria (c, d, h), the Sargasso Sea (e, f), and the Andaman Sea, Thailand (g). (a) Choano sp. 1; (b) Chono sp. 2; (c, d, h) *Stephanoeca urnula*; (e, f) *Savillea* sp.; (g) Choano sp. 3. The scale bar (f) applies to all LM images.

*Parvicorbicula* species that are closely affiliated with the *Parvicorbicula* type species *P. socialis* (Leadbeater 2015, Thomsen et al. 2020a).

*Coronoeca tongiae* is here (Fig. 14a-g; Table 2) reported from the Andaman Sea, Thailand, West Australia, the eastern Mediterranean Sea, Alexandria, and the Gulf of California, Mexico.

## Coronoeca patongiensis sp. nov. (Figs. 4g, 14h, i)

**Diagnosis**: Lorica funnel-shaped (lorica height ca. 15  $\mu$ m) and constructed of rod-shaped costal strips organized in ten longitudinal costae and three transverse costae. There are anterior spines corresponding in length to two thirds of an anterior longitudinal costal strip. A longitudinal costa is composed of four costal strips. Both the anterior transverse costa (diam. 12.5  $\mu$ m) and the mid-lorica costa (diam. 8.0  $\mu$ m) comprise as many costal strips as there are longitudinal costae, yet with a significantly larger overlap between neighbouring strips in the mid-lorica transverse costa. Transverse costal strips from both the anterior and the mid-lorica costa span the width between neighbouring longitudinal costae. The posterior transverse costa (diam. ca.  $3 \ \mu m$ ) comprises 4–5 costal strips. The anterior transverse costa is external relative to the longitudinal costae while the midlorica and the posterior transverse costae are internally located. The protoplast is located at the posterior end of the lorica chamber. Tectiform division presumed but not yet evidenced from the accumulation of costal strips in the collar region.

**Holotype**: The specimen illustrated in Fig. 14i of the present work is fixed as holotype (ICZN 1999, Article 73.1.4).

**Type locality**: Surface water sample (ca. 28 °C/35 PSU/2 m depth) from a near coastal site at Ao Patong, Phuket Island, Thailand (7° 53'56'' N/98° 17'36'' E), collected 8 September 1981.

**Etymology**: The species-group name refers to the type locality Ao Patong, Phuket Island, Thailand.

While this species is superficially reminiscent of *C. tongiae* with reference to lorica shape, dimensions and costal strip configuration, there are, however, a number of distinguishing features, i.e. the pattern of junctions between transverse and longitudinal costal strips at the level of the anterior transverse costa (E-joins in *C. patongiensis* and T-joins in *C. tongiae*), and the placing of the anterior transverse costa (internal in *C. tongiae* and external in *C.*  *patongiensis*). It is thus credible that the two species, once further studied using also molecular tools, cannot reside within the same genus. However, in an interim period it is convenient to be able to refer to a well-defined morphotype such as *C. patongiensis* using a formally established scientific name.

*Coronoeca patongiensis* has also been observed in samples from West Australia (Fig. 14h) and from the Sargasso Sea (Paul Davis; pers. com.).

#### Miscellaneous forms (Fig. 15)

Under this heading we have assembled material of loricate choanoflagellate taxa that have either been observed too sparingly to allow for a formal species description, or unintentionally left out while dealing with the relevant genus in a previous paper (*Stephanoeca urnula*).

#### Choano sp. 1 (Fig. 15a)

Lorica funnel shaped (lorica height:  $10.1 \ \mu m$ ) comprising eight longitudinal costae and two transverse costae (diam. 7.6 and 4.8  $\mu m$ ). The anterior tier of longitudinal costal strips project as spines. Costal strips from the anterior ring span the width between neighbouring longitudinal costal strips. This taxon resembles *C. patongiensis* but differs in having fewer transverse and longitudinal costae. Only observed (single specimen) in a water sample from the equatorial Pacific Ocean.

#### Choano sp. 2 (Fig. 15b)

Lorica funnel shaped (lorica height: 14.8  $\mu$ m) comprising ten longitudinal costae and two transverse costae (i.e. an anterior ring (diam. 12.0  $\mu$ m) forming E-junctions with the longitudinal costae, and a posterior costa (diam. 4.6  $\mu$ m) located at the joints between longitudinal costal strip number two and three counted from the anterior lorica end). This taxon, which is only observed once in a sample from the equatorial Pacific Ocean, resembles *Cosmoeca phuketensis* (see Thomsen and Østergaard 2019d) but differs because of a different number of longitudinal costae (there are consistently nine in *C. phuketensis*) and also by having a more complex posterior transverse costa.

# Choano sp. 3 (Fig. 15g)

A single cell observed in a sample from the Andaman Sea, Thailand. The funnel-shaped lorica (lorica height: 19.5  $\mu$ m) is constructed from eight longitudinal costae and two transverse costae. The anterior ring (diam. 12  $\mu$ m) closes the lorica (E-junctions). The mid lorica transverse costa (diam. 9  $\mu$ m) crosses posterior to the joints between the first and second longitudinal costal strips (counted from the anterior lorica end). Notice that the lon-

gitudinal costal strips are exceptionally long ( $6-7 \mu m$ ). The lorica chamber is pedicellate and posteriorly terminated by a basket-like configuration of obliquely oriented costal strips.

#### Stephanoeca urnula Thomsen, 1973 (Fig. 15c, d, h)

This species was observed repeatedly in samples from the Eastern Mediterranean Sea outside Alexandria, Egypt. It is morphometrically (lorica height: 11 µm; diam. 3.7 µm (anterior opening), 7.2 µm (max. lorica width), and 4.5 µm (lorica waist)) in overall good agreement with the Danish type material (Thomsen 1973), and also material sampled from the innermost parts of the Baltic Sea (Thomsen, 1979, Vørs 1992, Ikävalko and Thomsen 1997, Ikävalko 1998). Stephanoeca urnula has additionally been recorded from Lake Saroma, Japan (Takahashi 1981) and the South Atlantic coastal zone. Brazil (Bergesh et al. 2008; 14-15 PSU). The majority of samples yielding S. urnula are from low-saline regions such as the Baltic Sea. The occurrence of this form in an eastern Mediterranean high salinity environment is thus surprising. The exact habitat sampled is perhaps impacted by the river Nile freshwater outflow creating brackish water niches that may allow S. urnula to thrive, or is it rather that the Egyptian material, despite an overall similarity, does represent a separate taxon with markedly different habitat demands. It is in this connection relevant to refer to recently described nudiform Stephanoeca-like species (Enibas tolerabilis and E. thessalia; Schiwitza et al. 2019; Schiwitza and Nitsche 2021) that are morphologically reminiscent of S. urnula yet with a significantly different autecological signature. Based on laboratory tests is was found (Schiwitza et al. 2019) that E. tolerabilis is able to survive within a salinity range of 0-70 PSU.

#### Savillea sp. (Fig. 15e, f)

Two cells observed in samples from the Sargasso Sea. They differ from *Savillea parva* (Ellis, 1929) Loeblich III, 1967, by having a significantly more open-meshed pattern of spiralling and longitudinal costae. The lorica is 13–15  $\mu$ m high, and the diameter at the orifice 4–5  $\mu$ m and in the mid-lorica region 8–9  $\mu$ m. Similar material has previously been observed in Sargasso Sea samples (Paul. G. Davis, pers. com.; lorica height 14.2  $\mu$ m, diam. at orifice 4.2  $\mu$ m, max. diam. 7.3  $\mu$ m).

#### **Species diversity**

The total number of species encountered in our warm water samples is 80 (Table 2). This represents close to 50% of all loricate choanoflagellate species described so far. There is a fairly obvious link between the material available to us, in terms of the mere number of water samples



Fig. 16. (A) Map showing the approximate geographical positions of material that is part of the analysis presented in Fig. 17. Data sources: (1) NEW, N.E. Greenland; Thomsen and Østergaard 2017; (2) NOW, N.W. Greenland; Thomsen and Østergaard 2017; (3) Disko Bay, West Greenland; Thomsen and Østergaard 2017; (4) Igloolik, Canada; Daugbjerg et al. 1991; (5) Baltic Sea; Thomsen 1979, Vørs 1992, Ikävalko and Thomsen 1997, Ikävalko 1998; (6) Denmark; Thomsen et al. 2016; (7) Kilsfjorden, Norway; Espeland and Throndsen 1986; (8) Southampton, UK; Tong 1997a; (9) Newfoundland, Canada; McKenzie et al. 1997; (10) Gulf of St. Lawrence, Canada; Bérard-Therriault et al. 1999; (11) Rhode Island, USA; Menezes 2005; (12) Sargasso sea; Thomsen and Østergaard 2019a and subsequent papers; (13) Algiers; Leadbeater 1974; (14) Croatia and Montenegro; Leadbeater 1973, 1974; (15) Mediterranean Sea, Alexandria; Thomsen and Østergaard 2019a and subsequent papers; (16) Caribbean Sea; Thomsen and Østergaard 2019a and subsequent papers; (17) South Brazil; Bergesh et al. 2008; (18) Beagle Channel, Patagonia; Thomsen, unpublished; (19) Weddell Sea, Antarctica; Thomsen et al. 1990, Thomsen et al. 1997; (20) Prydz Bay, Antarctica; Marchant 1985, Marchant and Perrin 1990; (21) Andaman Sea, Thailand; Thomsen and Østergaard 2019a and subsequent papers; (22) Darwin, Australia; Lee et al. 2003; (23) West Australia; Thomsen and Østergaard 2019a and subsequent papers; (24) Shark Bay, Australia; Tong 1997b; (25) Sydney, Australia; Tong et al. 1998; (26) New Zealand; Moestrup 1979; Moestrup, unpublished; (27) Taiwan; Hara et al. 1997; Hara unpublished; (28) Japan; Hara et al. 1997; Hara, unpublished; (29) Subarctic North Pacific; Booth 1990; (30) California, USA; Thomsen et al. 1991; (31) Gulf of California, Mexico; Thomsen and Østergaard 2019a and subsequent papers; (32) North Pacific Central Gyre; Hoepffner and Haas 1990; (33) Equatorial Pacific Ocean; Thomsen and Østergaard 2019a and subsequent papers. (B) Number of species observed at each locality.

and LM/TEM preparations, and the total number of species listed from each individual site. The most thoroughly screened locations (West Australia and the Andaman Sea) are each represented by more than 60 species, whereas for the least intensively sampled regions (the Caribbean Sea, and the Eastern Mediterranean Sea) we have only listed approximately 30 species. For comparison it can be emphasized that similarly intensely sampled regions, i.e. Danish coastal waters and Disko Bay, West Greenland, are each represented by a maximum of 45 species. It thus appears that the tropical Indian Ocean sites sampled (the Andaman Sea and West Australia) do represent unique biodiversity hot spots.

To put the regions here analysed into a proper perspective, we have assembled biodiversity data worldwide (Fig. 16A, B) and conducted a matrix analysis (Fig. 17) in the shape of a two-way clustering of localities (35) versus species (presence-absence data; the diagram includes the 50 most decisive species).



**Fig. 17.** Shade matrix plot (Primer 7) showing relationships among clusters of samples (35) and species (50 most important taxa only/ presence-absence data). Both samples and species are hierarchically clustered independently. Clusters are further identified by letters (species: A-D; localities: E-H). See text for further explanation.

It is immediately obvious (Fig. 17), judged from the overall diagonal patterning of the matrix, that specific subsets of samples largely defined by temperature are matched by fairly clear-cut subgroups of species. The warm water sites reported on here cluster closely together (ca. 60% similarity) with maximum similarity (ca. 90%) between the Andaman Sea and the West Australian collections, and match up with a cluster of 9 species that have their maximum occurrences in such warm water habitats (Fig. 17D/ H; i.e. Campyloacantha imbricata, Campyloacantha spinifera, Cosmoeca ceratophora, Cosmoeca subulata, Pleurasiga echinocostata, Polyfibula elatensis, Stephanacantha Syndetophyllum pulchellum, dichotoma. Thomsenella infundibuliformis). It is similarly evident that there is a well-defined square within the matrix that defines associations between both Arctic and Antarctic cold-water habitats and cold-water adapted species (Fig. 17B/F; i.e. Acanthocorbis unguiculata, Bicosta antennigera, B. spinifera, Calliacantha longicaudata, C. natans, Diaphanoeca pedicellata, Parvicorbicula quadricostata, P. socialis). Notice that there is a North Atlantic cluster of sites (Baltic Sea, Denmark, Southampton (UK) and Rhode Island (USA)) that is overall very similar to the cold-water community, however, with a wider range of species represented. The matrix segment marked 'C' (Fig. 17) comprises species that are truly cosmopolitan forms (i.e. Bicosta minor, Cosmoeca norvegica, C. ventricosa, Calliacantha simplex, Diaphanoeca grandis, Parvicorbicula circularis, Pleurasiga min*ima*). The most divergent matrix element (Fig. 17A/E) is represented by Antarctic Weddell Sea icebiota samples which display a wide range of forms not encountered elsewhere. Samples from South Brazil and the North Pacific Central Gyre cluster separately very likely due to limitations in the number of taxa recorded from these sites (Fig. 16B). The inevitable conclusion to be extracted from the kind of analysis undertaken here, is that there are consistent and well-defined temperature dependent species associations with clear distributional constrictions, as well as groups of cosmopolitan species that have the potential to occur everywhere.

### Conclusions

While the genus *Polyfibula* is unambiguously defined based on overall lorica constructional details and especially the facets on transverse costal strips, it applies to all remaining species examined here, that the generic allocation is negotiable and likely to change once supplemental molecular data becomes available. In order to generally move loricate choanoflagellate taxonomy ahead, it is urgently required to seek supporting molecular evidence. A cultivation approach is one obvious way to go (see e.g. Nitsche and Arndt 2008; Schiwitza et al. 2019). However, it is likely that a vast range of species, and in particular those that prefer more oceanic habitats, will be very hard and time consuming to ever establish in culture. An alternative, or perhaps rather supplemental approach, is sequencing based on the extraction of single cells from a suspension of nanoflagellates concentrated from freshly collected water samples (see e.g. Nitsche et al. 2017; Schiwitza and Nitsche 2021). While this does work adequately when dealing with large forms, that can be recognized and unambiguously identified from mixed populations at low magnification in the inverted microscope (×40-60 objective), this approach will not suffice, when it comes to small loricate forms that share basic features in lorica design and overall construction. What is needed to move fast ahead, linking specific morphotypes with their unique sequences, is an approach that allows for an initial incontestable species identification, based on e.g. an examination of natural, wet and non-coated material in an environmental scanning electron microscope, and a subsequent extraction of the specific cell for sequencing. While the identification part and the position-fixing is fairly straightforward to accomplish, it is the subsequent manipulation (extraction) of the cell that is likely to be seriously challenging. No matter the approach taken it appears that an initial accessibility to flow-sorted cells, prepared as described by e.g. Kamennaya et al. (2018), might be essential in optimizing the success-rate in terms of securing quality approved data linking specific morphotypes to molecular profiles.

It is evident from Fig. 16A that the loricate choanoflagellate research community has by now achieved, based on classical light and electron microscopical descriptive techniques, a fairly comprehensive global coverage of species distributions. This has in turn contributed to e.g. a basic understanding of how temperature in particular impacts on large-scale species occurrence patterns. The approach that we have taken in the current series of papers on warm water loricate choanoflagellate biodiversity, is to reduce often complex regional sampling programmes (along multiple transects covering several depths) to a single species file. While this has proved adequate with reference to the taxonomic and morphometric data analysis, and also in the context of underpinning that temperature is a decisive factor in global biodiversity patterns, it conceals, however, species associations differentiating e.g. near coastal versus offshore habitats, surface water versus fluorescence maximum layers, and also where relevant (e.g. the Andaman Sea, Thailand) possible seasonal patterns in species occurrences. Once adequately progressed the molecular tools will be keystone elements in puzzling out any such small-scale patterns in species occurrences.

# Author contribution statement

Helge A. Thomsen (HAT) has conceived and designed the analysis, generated funding, and collected and processed samples. HAT is further responsible for compiling and writing the paper. Jette B. Østergaard (JBO) has been much involved in data collection and sample processing with reference to the Andaman Sea and the Pacific Ocean. Seiko Hara (SH) has contributed data from the Seto Inland Sea, Japan, and has been instrumental in finalizing the later versions of the manuscript.

#### Acknowledgements

Crew and scientists on board the research vessels ('Dana', 'Vædderen', 'Chakratong Tongyia', 'Malcolm Baldrige') are acknowledged for good collaborative spirit and for providing excellent facilities for collecting and processing our samples. Dr. Gunni Ærtebjerg and Steffen Mariager Pedersen are acknowledged for providing the samples from outside the Alexandria Harbour. We are grateful to Dr. Sergio Sanudo-Wilhelmy and Kurt Buck for organizing the trip to Gulf of California. Carol Kosman is warmly acknowledged for her meticulous contribution to this paper. Main funding sources, i.e. the Independent Research Fund Denmark, the Carlsberg Foundation, Danida (Denmark's development cooperation), the Højgaard Foundation, and the Danish Centre for Marine Research are acknowledged for their generous contributions to our research. The present work was carried out as part of the Galathea3 expedition under the auspices of the Danish Expedition Foundation (Galathea3 contribution no. 135), and as part of the Sargasso-Eel 2014 Expedition.

## References

- Bérard-Therriault, L., Poulin, M., Bossé, L., 1999. Guide d'identification du phytoplancton marin de l'estuaire et du Golfe du Saint-Laurent incluant également certains protozoaires. Publ. spec. sci. halieut. Aquat. 128, 1–387.
- Bergesch, M., Odebrecht, C., Moestrup, Ø., 2008. Loricate choanoflagellates from the South Atlantic coastal zone (~32 ° S) including the description of *Diplotheca tricyclica* sp. nov.. Biota Neotrop 8, 111–122.
- Booth, B.C., 1990. Choanoflagellates from the subarctic North Pacific Ocean, with description of two new species. Can. J. Zool. 68, 2393–2402.
- Buck, K., 1981. A study of choanoflagellates (Acanthoecidae) from the Weddell Sea, including a description of *Diaphanoeca multiannulata* n. sp. J. Protozool. 28, 47–54.
- Clarke, K.R., Gorley, R.N., 2015. PRIMER v7: User Manual/ Tutorial. PRIMER-E: Plymouth.
- Clarke, K.R., Gorley, R.N., Somerfierld, P.J., Warwick, R.M., 2014. Change in marine communities: an approach to statistical analysis and interpretation. PRIMER-E: Plymouth.
- Daugbjerg, N., Hansen, L.E., Skovgaard, K., Østergaard, J.B., 1991. An investigation on some protist plankton groups from the marine waters around Igloolik Island, NWT, Canada. In: Jørgensen, M. (ed.), Arctic Biology Course 1989, Igloolik Island, NWT Canada. Publ. by Zoological Museum, Univ. of Copenhagen, pp. 39-59.
- Espeland, G., Throndsen, J., 1986. Flagellates from Kilsfjorden, southern Norway, with description of two new species of Choanoflagellida. Sarsia 71, 209–226.

- Hallegraeff, G.M., 1983. Scale-bearing and loricate nanoplankton from the East Australian Current. Bot. Marina 26, 493–515.
- Hara, S., Sheu, J., Chen, Y.L., Takahashi, E., 1997. Choanoflagellates (Sarcomastigophora, Protozoa) from the coastal waters of Taiwan and Japan (II): Species composition and biogeography. Zool. Stud. 36, 98–110.
- Hoepffner, H., Haas, L.W., 1990. Electron microscopy of nanoplankton from the North Pacific central gyre. J. Phycol. 26, 421-439.
- ICZN 1999. International Code of Zoological Nomenclature, Fourth Edition: The International Trust for Zoological Nomenclature, London, UK, 306 pp.
- Ikävalko, J., 1998. Further observations on flagellates within sea ice in northern Bothnian Bay, the Baltic Sea. Polar Biol. 19, 323–329.
- Ikävalko, J., Thomsen, H.A., 1997. The Baltic Sea ice biota (March 1994): a study of the protistan community. Eur. J. Protistol. 33, 229–243.
- Kamennaya, N.A., Kennaway, G., Fuchs, B.M., Zubkov, M.V., 2018. "Pomacytosis" – Semi-extracellular phagocytosis of cyanobacteria by the smallest marine algae. PLoS Biol. 16,. https://doi.org/10.1371/journal.pbio.2003502 e2003502.
- Leadbeater, B.S.C., 1973. External morphology of some marine choanoflagellates from the coast of Jugoslavia. Arch. Protistenk. 115, 234–252.
- Leadbeater, B.S.C., 1974. Ultrastructural observations on nanoplankton collected from the coast of Jugoslavia and the Bay of Algiers. J. Mar. Biol. Ass. U.K. 54, 179–196.
- Leadbeater, B.S.C., 2015. The Choanoflagellates: Evolution, biology and ecology. Cambridge University Press.
- Lee, W.J., Brandt, S.M., Vørs, N., Patterson, D.J., 2003. Darwin's heterotrophic flagellates. Ophelia 57, 63–98.
- Manton, I., Bremer, G., 1981. Observations on lorica structure and aspects of replication in the *Pleurasiga sphyrelata* Thomsen complex (= *Polyfibula* spp., gen.n.) (Choanoflagellata). Zool. Scr. 10, 273–291.
- Marchant, H.J., 1985. Choanoflagellates in the Antarctic marine food chain. In: Siegfried, W.R., Condy, P.R., Laws, R.M. (Eds.), Antarctic Nutrient Cycles and Food Webs. Springer-Verlag, Berlin Heidelberg, pp. 271–276.
- Marchant, H.J., Perrin, R.A., 1990. Seasonal variation in abundance and species composition of choanoflagellates (Acanthoecidae) at Antarctic coastal sites. Polar Biol. 10, 499–505.
- McKenzie, C.H., Deibel, D., Thompson, R.J., MacDonald, B.A., Penney, R.W., 1997. Distribution and abundance of choanoflagellates (Acanthoecidae) in the coastal cold ocean of Newfoundland, Canada. Mar. Biol. 129, 407–416.
- Menezes, S., 2005. Nanoplankton Biodiversity in the Pettaquamscutt River Estuary, Rhode Island, U.S.A PhD thesis. Univ. Rhode Island, p. 319.
- Moestrup, Ø., 1979. Identification by electron microscopy of marine nanoplankton from New Zealand, including the description of four new species. New Zeal. J. Bot. 17, 61–95.
- Moestrup, Ø., Thomsen, H.A., 1980. Preparation of shadow-cast whole mounts. In: Gantt, E. (Ed.), Handbook of Phycological Methods. Vol. III. pp. 385-390 (Cambridge).
- Nitsche, F., Arndt, H., 2008. A new choanoflagellate species from Taiwan: Morphological and molecular biological studies

of *Diplotheca elongata* nov. spec., and *D. costata*. Eur. J. Protistol. 44, 220–226.

- Nitsche, F., Thomsen, H.A., Richter, D.J., 2017. Bridging the gap between morphological species and molecular barcodes exemplified by loricate choanoflagellates. Eur. J. Protistol. 57, 26–37.
- Schiwitza, S., Nitsche, F., 2021. A needle in the haystack mapping sequences to morphology exemplified by the loricate choanoflagellate *Enibas thessalia* sp. nov. (Acanthoecida, Acanthoecidae). Protist 172, (1). https://doi. org/10.1016/j.protis.2020.125782 125782.
- Schiwitza, S., Arndt, H., Nitsche, F., 2019. First description of an euryoecious acanthoecid choanoflagellate species, *Enibas tolerabilis* gen. et sp. nov. from a salar in the Chilean Andes based on morphological and transcriptomic data. Eur. J. Protistol. 67, 106–113.
- Soto-Liebe, K., Collantes, G., Kuznar, J., 2007. New records of marine choanoflagellates off the Chilean coast. Invest. Mar. Valparaíso 35, 113–120.
- Takahashi, E., 1981. Floristic study of ice algae in the sea ice of a lagoon, Lake Saroma, Hokkaido, Japan. Mem. National Inst. Polar Res. Series E: Biol. Med. Sci. 34, 49–56.
- Thomsen, H.A., 1973. Studies on marine choanoflagellates I. Silicified choanoflagellates of the Isefjord (Denmark). Ophelia 12, 1–26.
- Thomsen, H.A., 1976. Studies on marine choanoflagellates II. Fine-structural observations on some silicified choanoflagellates from the Isefjord (Denmark), including the description of two new species. Norw. J. Bot. 23, 33–51.
- Thomsen, H.A., 1978. Nanoplankton from the Gulf of Elat (= Gulf of Aquaba), with particular emphasis on the choanoflagellates. Isr. J. Zool. 27, 34–44.
- Thomsen, H.A., 1979. Electron microscopical observations on brackish-water nannoplankton from the Tvärminne area, S.W. Coast of Finland. Acta Bot. Fenn. 110, 11–37.
- Thomsen, H.A., 1982. Planktonic choanoflagellates from Disko Bugt, West Greenland, with a survey of the marine nanoplankton of the area. Meddr. Grønland. Bioscience 8, 1–35.
- Thomsen, H.A., Boonruang, P., 1983. A microscopical study of marine collared flagellates (Choanoflagellida) from the Andaman Sea, SW Thailand: Species of *Stephanacantha* gen. nov. and *Platypleura* gen. nov. Protistologica 19, 193– 214.
- Thomsen, H.A., Buck, K.R., 1991. Choanoflagellate diversity with particular emphasis on the Acanthoecidae. In: Patterson, D.J., Larsen, J. (Eds.), Free-Living Heterotrophic Flagellates. Clarendon Press, Oxford, pp. 259–284.
- Thomsen, H.A., Moestrup, Ø., 1983. Electron microscopical investigations on two loricate choanoflagellates (Choanoflagellida), *Calotheca alata* gen. et sp. nov. and *Syndetophyllum pulchellum* gen. et comb. nov., from Indo-Pacific localities. Proc. R. Soc. Lond. B Biol. Sci. 219, 41–52.
- Thomsen, H.A., Østergaard, J.B., 2017. Acanthoecid choanoflagellates from the Atlantic Arctic region a baseline study. Heliyon 3,. https://doi.org/10.1016/j. heliyon.2017.e00345 e00345.
- Thomsen, H.A., Østergaard, J.B., 2019a. Loricate choanoflagellates (Acanthoecida) from warm water seas. I.

*Conioeca* gen. nov. and *Nannoeca* Thomsen. Eur. J. Protistol. 67, 77–88.

- Thomsen, H.A., Østergaard, J.B., 2019b. Loricate choanoflagellates (Acanthoecida) from warm water seas. II. *Bicosta, Apheloecion, Campyloacantha* and *Saroeca*. Eur. J. Protistol. 67, 114–131.
- Thomsen, H.A., Østergaard, J.B., 2019c. Loricate choanoflagellates (Acanthoecida) from warm water seas. III. *Acanthocorbis* Hara and Takahashi and *Stephanoeca* Ellis. Eur. J. Protistol. 69, 52–69.
- Thomsen, H.A., Østergaard, J.B., 2019d. Loricate choanoflagellates (Acanthoecida) from warm water seas. IV. *Cosmoeca* Thomsen. Eur. J. Protistol. 71, https://doi.org/ 10.1016/j.ejop.2019.125632 125632.
- Thomsen, H.A., Østergaard, J.B., 2019e. Loricate choanoflagellates (Acanthoecida) from warm water seas. V. *Thomsenella* Özdikmen (= *Platypleura* Thomsen). Eur. J. Protistol. 71,. https://doi.org/10.1016/j.ejop.2019.125633 125632.
- Thomsen, H.A., Østergaard, J.B., 2021. Loricate choanoflagellates (Acanthoecidae) from warm water seas. VIII. *Crinolina* Thomsen and *Diaphanoeca* Ellis. Eur. J Protistol. (in press).
- Thomsen, H.A., Buck, K.R., Coale, S.L., Garrison, D.L., Gowing, M.M., 1990. Loricate choanoflagellates (Acanthoecidae, Choanoflagellida) from the Weddell Sea, Antarctica. Zool. Scr. 19, 367–387.
- Thomsen, H.A., Buck, K.R., Chavez, F.P., 1991. Choanoflagellates of the central California waters: Taxonomy, morphology and species assemblages. Ophelia 33, 131–164.
- Thomsen, H.A., Garrison, D.L., Kosman, C., 1997. Choanoflagellates (Acanthoecidae, Choanoflagellida) from the Weddell Sea, Antarctica, taxonomy and community structure with particular emphasis on the ice biota; with preliminary remarks on choanoflagellates from Arctic sea ice (Northeast Water Polynya, Greenland). Arch. Protistenk. 148, 77–114.
- Thomsen, H.A., Nitsche, F., Richter, D.J., 2016. Seasonal occurrence of loricate choanoflagellates in Danish inner waters. Protist 167, 622–638.
- Thomsen, H.A., Kamennaya, N., Zubkov, M.V., Østergaard, J. B., 2020a. Loricate choanoflagellates (Acanthoecida) from warm water seas. VI. *Pleurasiga* Schiller and *Parvicorbicula* Deflandre. Eur. J. Protistol. 75, https://doi.org/10.1016/j. ejop.2020.125717 125717.
- Thomsen, H.A., Kamennaya, N., Zubkov, M.V., Østergaard, J. B., 2020b. Loricate choanoflagellates (Acanthoecida) from warm water seas. VII. *Calotheca* Thomsen and Moestrup, *Stephanacantha* Thomsen and *Syndetophyllum* Thomsen and Moestrup. Eur. J. Protistol. 76, https://doi.org/10.1016/j. ejop.2020.125728 125728.
- Tong, S.M., 1997a. Choanoflagellates in Southampton Water including the description of three new species. J. Mar. Biol. Ass. U.K. 77, 929–958.
- Tong, S.M., 1997b. Heterotrophic flagellates from the water column in Shark Bay, Western Australia. Mar. Biol. 128, 517–536.

- Tong, S.M., Nygaard, K., Bernard, C., Vørs, N., Patterson, D.J., 1998. Heterotrophic flagellates from the water column in Port Jackson, Sydney, Australia. Eur. J. Protistol. 34, 162–194.
- Vørs, N., 1992. Heterotrophic amoebae, flagellates and heliozoa from the Tvärminne area, Gulf of Finland, in 1988–1990. Ophelia 36, 1–109.
- Vørs, N., Buck, K.R., Chavez, F.P., Eikrem, W., Hansen, L.E., Østergaard, J.B., Thomsen, H.A., 1995. Nanoplankton of the equatorial Pacific with emphasis on the heterotrophic protists. Deep-Sea Res. 42, 585–602.