



Mobile Optical Remote Sensing for quantification of Ammonia and Methane emissions from Dairy Farms in California

Vechi, Nathalia Thygesen; Mellqvist, Johan; Offerle, Brian; Samuelsson, Jerker; Scheutz, Charlotte

Link to article, DOI:

[10.5194/egusphere-egu21-10911](https://doi.org/10.5194/egusphere-egu21-10911)

Publication date:

2021

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Vechi, N. T., Mellqvist, J., Offerle, B., Samuelsson, J., & Scheutz, C. (2021). *Mobile Optical Remote Sensing for quantification of Ammonia and Methane emissions from Dairy Farms in California*. EGU21-10911. Abstract from EGU General Assembly 2021. <https://doi.org/10.5194/egusphere-egu21-10911>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Mobile Optical Remote Sensing for quantification of Ammonia and Methane emissions from Dairy Farms in California.

Nathalia Thygesen Vechi^{1,2}, Johan Mellqvist^{2,3}, Brian Offerle³, Jerker Samuelsson³, and Charlotte Scheutz¹

¹Technical University of Denmark, Department of Environmental Engineering, Kgs. Lyngby, Denmark (navei@env.dtu.dk)

²Department of Space, Earth and Environment, Chalmers University, Göteborg, Sweden.

³FluxSense AB, SE-41296 Göteborg, Sweden.

Solar occultation flux (SOF) and Mobile extractive FTIR (MeFTIR) are techniques used for over 20 years to quantify industrial emissions of VOCs, CH₄, and others, from refineries in the USA, Europe, and Asia. Here, they were combined to assess methane (CH₄) and ammonia (NH₃) from concentrated animal feeding operations (CAFOs) in the San Joaquin Valley (SJV), California. SOF and MeFTIR were used to measure NH₃ column, and ground concentrations of NH₃ and CH₄, respectively. SOF retrieves the gas column concentration from the solar spectra using a solar track, directing the light to a FTIR spectrometer, while crossing the gas plume. Subsequently, a direct flux approach combines the retrieved columns with wind information to obtain the mass fluxes of ammonia. In this survey, the wind information was acquired by a wind LIDAR, which measures wind speed and direction in the interval of 10 – 300 m. On the other hand, Methane emissions were quantified using a unique indirect flux approach by combining the estimated ammonia fluxes and the NH₃:CH₄ ratios measured from the ground concentration using MeFTIR.

Two field campaigns performed in spring and autumn studied emissions from 14 single dairy CAFOs. The daily emissions from the single farms averaged $96.4 \pm 38.4 \text{ kg}_{\text{NH}_3} \text{ h}^{-1}$ and $411 \pm 185.4 \text{ kg}_{\text{CH}_4} \text{ h}^{-1}$, respectively, for NH₃ and CH₄ with the corresponding emission factors (EF) per animal unit of $11.3 \pm 3.8 \text{ g}_{\text{NH}_3} \text{ h}^{-1} \text{ AU}^{-1}$ and $50.3 \pm 24.1 \text{ g}_{\text{CH}_4} \text{ h}^{-1} \text{ AU}^{-1}$. The uncertainty of ammonia measurements was 17 % in a standard confidence interval (CI) and 37 % in a 95 % CI, with the largest uncertainty associated with the wind measurements. Furthermore, the methane uncertainty estimations averaged 27 % in a standard CI, and 52 % in a 95 % CI, dominated by the ammonia fluxes uncertainty. Comparison between Annual or daily EFs obtained by SOF to other quantification approaches, have to take into consideration the SOF measurement conditions, day-time and sunny weather, due to their effects on the NH₃ emissions. The study contributed to develop the knowledge of dairy CAFOs emission, and to strengthen the role of optical remote sensing techniques, bridging the gap between satellites and stationary measurement approaches.