



## Using experimental animal data from ToxValDB to derive points of departure for application in LCIA and comparative risk screening

Aurisano, Nicolo; Huang, Lei; Jang, Suji; Chiu, Weihsueh A; Judson, Richard S. ; Jolliet, Olivier; Fantke, Peter

*Publication date:*  
2021

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Aurisano, N., Huang, L., Jang, S., Chiu, W. A., Judson, R. S., Jolliet, O., & Fantke, P. (2021). *Using experimental animal data from ToxValDB to derive points of departure for application in LCIA and comparative risk screening*. Abstract from International Society of Exposure Science 2021.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Using experimental animal data from ToxValDB to derive points of departure for application in LCIA and comparative risk screening

Nicolò Aurisano<sup>1</sup>, Lei Huang<sup>2</sup>, Suji Jang<sup>3</sup>, Weihsueh A. Chiu<sup>3</sup>, Richard Judson<sup>4</sup>, Olivier Jolliet<sup>2</sup>, Peter Fantke<sup>1</sup>

<sup>1</sup>Quantitative Sustainability Assessment, Department of Technology, Management and Economics, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark

<sup>2</sup>Department of Environmental Health Sciences, University of Michigan, Ann Arbor, Michigan, 48109, USA

<sup>3</sup>Department of Veterinary Integrative Biosciences, College of Veterinary Medicine and Biomedical Sciences, Texas A&M University, College Station, Texas, USA

<sup>4</sup>National Center for Computational Toxicology, U.S. Environmental Protection Agency, Research Triangle Park, NC 27711, USA

Life cycle impact assessment (LCIA) and comparative risk screening rely on chemical-specific points of departure (PODs) from regulatory toxicity data sources to evaluate toxicological impacts on human health from chemical exposures. However, regulatory PODs are not available for the majority of chemicals to which humans can be exposed. We thus aim to broaden the coverage of chemicals by using available *in vivo* data to estimate PODs that most closely mimic one that would be selected in a regulatory assessment context. As a starting point, we extracted and curated experimental animal data available from the US EPA ToxValDB, focusing on oral repeated-dose studies and three non-cancer effect-level types: lowest observed adverse effect level (LOAEL), no-observed adverse effect level (NOAEL) and benchmark dose lower bound (BMDL). The curation process included harmonization of units into mg/kg-d followed by extrapolations from subchronic or subacute studies to chronic, from different effect-level types to benchmark dose (BMD) and from tested animal species to humans. After curation, in the case of data-rich chemicals with at least 10 data points, we fit the resulting data to a lognormal distribution. For data-poor chemicals with less than 10 data points, we fit the distribution while applying a fixed standard deviation of  $\log_{10}=0.55$ , based on the average standard deviation across data-rich chemicals in our curated dataset. PODs were then derived for 9037 chemicals based on the 5<sup>th</sup>-ile of the fitted lognormal distribution. The resulting POD values ranged by orders of magnitude from  $1e-7$  mg/kg-d to  $1e4$  mg/kg-d across the considered substances, with a median POD=7 mg/kg-d. For the 746 chemicals with available regulatory PODs, we observed a good correlation with our derived 5<sup>th</sup>-ile PODs ( $R^2=0.70$  and  $RMSE=0.64$  of the  $\log_{10}$ -transformed values). These results suggest that the proposed method is able to derive PODs consistent with regulatory values in a high-throughput approach, thus substantially increasing the coverage of chemical substances for application in LCIA and risk screening. Next steps include using this curated dataset to train a machine-learning-based prediction model to estimate PODs for an even wider range of substances without any experimental animal data available. *This abstract does not necessarily reflect US EPA policy.*