



Hydrodynamic trade-offs in different flagellar arrangement

Asadzadeh, Seyed Saeed; Walther, Jens H.; Kiørboe, Thomas

Publication date:
2022

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Asadzadeh, S. S., Walther, J. H., & Kiørboe, T. (2022). *Hydrodynamic trade-offs in different flagellar arrangement*. Abstract from *Microscale Ocean Biophysics 6.0*, Port de Soller, Spain.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Hydrodynamic trade-offs in different flagellar arrangements

Seyed Saeed Asadzadeh¹, Jens H. Walther^{2,3}, Thomas Kiørboe¹,

¹Centre for Ocean Life, National Institute of Aquatic Resources, Technical University of Denmark

²Department of Mechanical Engineering, Technical University of Denmark

³Computational Science and Engineering Laboratory, ETH, Zrich, Switzerland

Technical
University of
Denmark



Microscale Ocean Biophysics 6.0, 2022

Flagellates are few-micron sized unicellular organisms equipped with one or a few flagella, and play a key role at the base of oceanic food webs as the main consumers of bacteria and phytoplankton. The beating flagellum propels the organism through the water and generates a feeding current that facilitates prey encounters. At the same time the stirring generated by the flagellum exposes the cells to their flow-sensing predators. While hydrodynamics of flagella is most often studied in the context of propulsion, efficient foraging is likely a much more important component of their fitness than propulsion per se, giving rise to a great diversity of flagellar arrangements, morphology, and beat kinematics. We hypothesize that the significance of this diversity represents different outcomes of the fundamental trade-offs between resource acquisition, predator avoidance, and propulsion. Using computational fluid dynamics, we study these hydrodynamic trade-offs and investigate how at low Reynolds number, where viscosity impedes predator-prey contact, flagellates yield the necessary feeding current structure and at the same time avoid flow sensing-predators to secure their success and key role in the microbial food webs.