

Blood-based gene expression as non-lethal tool for inferring salinity-habitat history of European eel (Anguilla anguilla)

Bertolini, Francesca; Rohtla, Mehis; Parzanini, Camilla; Tomkiewicz, Jonna; Durif, Caroline M.F.

Published in: Scientific Reports

Link to article, DOI: 10.1038/s41598-022-26302-y

Publication date: 2022

Document Version Publisher's PDF, also known as Version of record

Link back to DTU Orbit

Citation (APA): Bertolini, F., Rohtla, M., Parzanini, C., Tomkiewicz, J., & Durif, C. M. F. (2022). Blood-based gene expression as non-lethal tool for inferring salinity-habitat history of European eel (*Anguilla anguilla*). *Scientific Reports, 12*, Article 22142. https://doi.org/10.1038/s41598-022-26302-y

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- · You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

scientific reports

OPEN



Blood-based gene expression as non-lethal tool for inferring salinity-habitat history of European eel (Anguilla anguilla)

Francesca Bertolini^{1,2⊠}, Mehis Rohtla^{3,4}, Camilla Parzanini⁵, Jonna Tomkiewicz¹ & Caroline M. F. Durif³

The European eel is a facultative catadromous species, meaning that it can skip the freshwater phase or move between marine and freshwater habitats during its continental life stage. Otolith microchemistry, used to determine the habitat use of eel or its salinity history, requires the sacrifice of animals. In this context, blood-based gene expression may represent a non-lethal alternative. In this work, we tested the ability of blood transcriptional profiling to identify the different salinity-habitat histories of European eel. Eels collected from different locations in Norway were classified through otolith microchemistry as freshwater residents (FWR), seawater residents (SWR) or inter-habitat shifters (IHS). We detected 3451 differentially expressed genes from blood by comparing FWR and SWR groups, and then used that subset of genes in a machine learning approach (i.e., random forest) to the extended FWR, SWR, and IHS group. Random forest correctly classified 100% of FWR and SWR and 83% of the IHS using a minimum of 30 genes. The implementation of this non-lethal approach may replace otolith-based microchemistry analysis for the general assessment of life-history tactics in European eels. Overall, this approach is promising for the replacement or reduction of other lethal analyses in determining certain fish traits.

The European eel (*Anguilla anguilla*) is considered a facultative catadromous fish. While it spawns in the ocean, it displays several strategies during its continental growth phase by either growing in freshwater systems, skipping the freshwater phase or shifting between various salinity habitats (i.e., seawater, brackish water, freshwater)¹. Stock assessment is almost only based on the freshwater contingent of the species². Eels living their entire lifecycle in seawater (i.e., skipping the freshwater phase) are rarely included in assessments and their proportion relative to the whole population is unknown. Moreover, when eels are surveyed, it is unknown whether some have recently shifted between habitats. This distribution pattern complicates stock assessment and management of this species.

Otolith microchemistry is a method commonly applied to determine salinity-history in fish and thereby their use of habitats. Such analysis rely on the variation of ambient water chemistry, as chemical elements are incorporated into the otolith in a predictive manner throughout the life of the fish^{1,3}. Otolith microchemical analysis is reliable but requires the sacrifice of the animal. Nonetheless, it is so far the only method which provides migratory information throughout the entire life of a fish.

More recently, fatty acid analysis has also been considered to determine habitat salinity history of eel. The use of fatty acid biomarkers relies on the variation of dietary sources across habitats (in particular between marine and freshwater habitats), which is then reflected in the fatty acid composition of the consumer⁴. This analysis typically requires biopsy of the muscle among other tissues and/or organs (e.g., liver), providing relatively recent (i.e., weeks to several months) feeding information.

In this context, blood-based biomarkers may represent a non-lethal alternative to determine individual salinity-habitat history. Blood is a living tissue that transports molecules throughout the body. A blood sample can, therefore, reflect an individual's physiological state regarding its health, nutrition, reproductive development, stress and/or metabolism^{5–8}. Moreover, blood can be sampled without euthanasia in a high number of species.

¹National Institute of Aquatic Resources, Technical University of Denmark, Kongens Lyngby, Denmark. ²Division of Animal Sciences, Department of Agricultural and Food Sciences, University of Bologna, Bologna, Italy. ³Institute of Marine Research, Austevoll Research Station, Storebø, Norway. ⁴Estonian Marine Institute, University of Tartu, Tartu, Estonia. ⁵Department of Nutritional Sciences, University of Toronto, Canada. ^{\Box}email: franb@ aqua.dtu.dk; francesca.bertolini3@unibo.it

For identification of possible biomarkers, blood transcriptome is of particular interest, as gene expression is a crucial regulator of cell functions. Here, individual variability, physiological status and external factors, highly influence gene expression⁹. Through the improvement of Next Generation Sequencing (NGS) technologies, and the development of RNA-seq¹⁰ and bioinformatics tools, it is now possible to simultaneously profile the transcriptional activity of all genes in a desired tissue, from small starting amount of RNA. This has accelerated the discovery of informative genes that can represent biomarkers linked to specific physiological conditions and with a wide range of applications^{9,11,12}.

Whole blood transcriptome analysis has been used in several mammalian species, including humans, to identify biomarker indicators of pathological conditions, immune competences, and reproductive performance¹³⁻¹⁷. In mammalian blood, the only nucleated and hence transcriptionally active cells are the white cells (i.e., leucocytes). In contrast, red blood cells (i.e., erythrocytes) of non-mammalian vertebrates are also nucleated and hence transcriptionally active¹⁸. This makes non-mammalian blood potentially more informative than mammalian blood.

The recent and limited number of studies targeting blood transcriptomics in non-mammalian vertebrates, such as birds and reptiles, show that genetic activity between blood and liver have a high degree of similarity, especially in terms of biological processes^{19,20}. The liver is an organ with high functional gene expression diversity, and it is frequently used in transcriptomic investigations. Blood transcriptomic-based analyses show applications in ecology, ecophysiology, and toxicology studies where gene transcripts in the blood have been studied as candidate indicators of behavior, physiological condition, environmental impacts and other phenotypic differences, particularly for wild species ²⁰⁻²³.

Similar outcomes have been produced in the even more limited number of studies in fish, both in experimental and in field settings. A study on fathead minnow *Pimephales promelas*, investigating exposure to environmentally relevant concentrations of chemical substances, showed how blood responded with a greater number of altered genes compared to liver, and how they shared the same biological altered pathways²⁴. Another study detected 563 blood transcript biomarkers that can be used for non-lethal sex differentiation in the Asian swamp eel *Monopterus albus*²⁵.

The aim of the present study was to investigate whether whole blood transcriptional profile could be used to infer the salinity-habitat history from eels collected in the wild. To do this, we sampled blood, otoliths, and muscle tissue from European eels collected in different salinity habitats to compare transcriptomes with the otolith microchemistry (entire life) and fatty acid profiles (recent migratory history), which served to validate individual eel salinity histories. We identified candidate genes that could be used as transcriptomic-based biomarkers to discriminate salinity-habitat history and applied a machine learning approach to evaluate the ability of those biomarkers to match the classification derived by otolith microchemistry analysis. In recent years, machine learning algorithms have been used for the analysis of high-throughput deep sequencing data due to their computational efficiency in finding generalizable patterns from high-dimensional data obtained from a small number of samples²⁶. Here, we used random forest (RF), a supervised machine learning algorithm that is widely used in classification and regression problems²⁷.

Material and methods

Statement. Sampling and handling of eels in this study were approved by the Norwegian Animal Research Authority and all procedures followed local animal welfare regulations (FOTS id 15952) and are in accordance with ARRIVE guidelines.

Samples collection. Eels were collected in July and August 2020, in five sites in Norway which represented three different salinity habitats (Fig. 1): freshwater (FW; Arendal and Bergen), brackish water (BW; Arendal), and seawater (SW; Bergen and Haugesund). Eels were caught using fyke nets (mesh size at the cod end was ~8 mm, knot-to-knot, and 11 mm along the diagonal) and eel pots (mesh size was ~ 10 mm, knot-to-knot, and 15 mm diagonal). Brackish waters are often characterized by salinity stratification making it difficult to assign an exact salinity value to each location. Therefore, we considered SW sites to correspond to salinities >30 ppt, BW to comprise sites corresponding to salinities between 0.5–30 ppt, and FW sites to be <0.5 ppt.

A total of 151 individuals were captured and anaesthetized with clove oil and measured for total length (mm), wet mass (g), eye diameter (mm) and fin length (mm). These measurements were used to assign a silvering stage, where stages I and II represent eels in their growth phase (classic "yellow" phase), stage III indicates a pre-migrant phase, and stages IV and V are the two migratory "silver" phases²⁸.

From 60 anaesthetized animals, approximately 600 μ l of blood was collected in lithium heparine tubes, mixed with 2 volumes of RNA-later (Invitrogen) and stored at – 20 °C prior to RNA extraction. Anaesthetized eels were then euthanized and dissected. The sex of all captured eels was determined by macroscopic observation of gonads. Skinless, white muscle tissue close to the dorsal fin (~ 1.0 × 0.5 cm) was sampled for fatty acid analysis. Sagittal otoliths were dissected for microchemistry analysis (Table S1).

Otolith and fatty acid-based classification. Transversal otolith thin sections were prepared for chemical analysis of otoliths. Otolith thin sections were analysed for ²⁴ Mg, ⁴³Ca, ⁵⁵Mn, ⁸⁸Sr and ¹³⁷Ba using laser ablation inductively coupled plasma mass spectrometry at the University of Tartu (Department of Geology). A continuous line scan was traced from the core to the edge using a laser speed of 5 μ m/s and laser beam diameter of 40 μ m. Eels were classified as seawater residents (SWR), freshwater residents (FWR) or inter-habitat shifters (IHS) based on established otolith freshwater reference values of Sr:Ca and Ba:Ca (see Rohtla et al.²⁹ for additional details on chemical analysis and data interpretation).

Lipids were extracted using a modified version of the Folch et al.³⁰ method in a chloroform:methanol (2:1) solution, and fatty acids analyzed as methyl esters (FAME) through gas-chromatography and flame ionization





detection (GC-FID) at the Department of Chemistry and Biology of Ryerson University, Toronto, Canada. Individual fatty acids were identified using a series of standards, including a 39-component FAME mix (GLC-463, Nu-Check Prep Inc.), a marine PUFA mix inclusive of 22:1n-11 (11-docosenoic acid methyl ester; PUFA Mix No.1, Supelco Inc.), and 18:4n-3 (stearidonic acid; SMB00291, Sigma-Aldrich). The M/F ratio was used to determine eel diet³¹ This ratio provides information on a largely marine-(M; higher values) versus freshwater-based diet (F; lower values), and hence, indirectly, information on the feeding habitat (i.e., SW, FW), depending on the presence of a few characteristic fatty acids in the muscle tissue. Intermediate values were assumed to represent eels feeding and living in BW and/or eels regularly moving between SW and FW.

RNA extraction and sequencing and data mining. RNA extraction was performed following a modified Trizol (Invitrogen) protocol, starting from 200 µl of mixed whole blood + RNA-later. RNA integrity was assessed with Bioanalyzer, to estimate the RNA Integrity Number (RIN) for each of the samples. All samples had RIN > 8 and were therefore evaluated as intact and suitable for sequencing. Paired end mRNA-seq 150 bp was performed at Novogene co (China).

Read quality was assessed by Fastqc³². Reads were trimmed with Trimmomatic v0.38³³, removing the first 9 bases at the beginning of the reads, as well as part of reads that had lower quality and reads that were shorter than 36 bp after trimming (HEADCROP:9, SLIDINGWINDOW: 4:15, MINLEN:36). Mapping of trimmed reads was performed with Tophat2 v0.13³⁴ with default options using the latest version of the European eel reference genome and annotation fAngAng1.pri (NCBI; GCF_013347855.1) to guide the read mapping. A further filter was performed with Samtools v1.10³⁵, removing reads that mapped in multiple places and sorting the reads by read name, as condition to run the reads count. For every samples, reads count at each annotated gene was performed with htseq-count³⁶.

Differential expression and gene enrichment. Transcriptome analysis was performed at first with the overall dataset using the Deseq2 R package³⁷ considering the normalized log2-fold change. Differential expression was then done with the same package considering only FWR and SWR derived from the otolith analyses. Here, genes with adj p values < 0.05 were considered as significantly differentially expressed (DE) and used for subsequent gene enrichment and Random Forest (RF) analyses.

Gene enrichment was performed with upregulated and downregulated genes separately using Panther (http:// www.pantherdb.org/), considering Gene Ontology (GO) biological processes as annotation set and *Danio rerio* as reference gene set. Only GO terms with FDR p < 0.05 were considered. Redundant GO terms were removed with REVIGO³⁸, utilizing the *Danio rerio* database and applying SimRel as semantic similarity measure.

Random forest. In a classification context, RF allows to assign a unit for which the class is unknown to a pre-determined group, using the so-called majority rule to aggregate the 'B' predictions obtained from the different trees in the forest. The final predicted class is the most commonly occurring one. In RF, as trees are fitted to bootstrapped subsets of the data set, some observations are left aside each time a tree is built. This leads to a valid estimate of the prediction error of a random forest, which is the so-called out-of-bag (OOB) error²⁷. Predictors can then be ranked according to their relevance in the classification rule in two measures: the Mean Decrease in the Gini Index and the Mean Accuracy Decrease²⁷.

RF on the reduced gene set was performed over normalized gene expression data, obtained through a logtransformation of the whole gene set with Deseq2³⁷. Then, the log-transformed values of the DE genes were used for the RF analysis considering the whole sample set. Samples were classified according to the otolith analyses into three groups: FWR, SWR, and IHS. RF analysis was performed with the R package randomForest³⁹, with 700 trees. Each analysis was repeated 10 times and the average values were calculated. From the first round of analyses with all the DE genes, Mean Decrease in the Gini Index and the Mean Accuracy Decrease were also extracted. Then, genes with no contribution (=0) to Gini and Accuracy in at least one of the groups (i.e., FWR, IHS, SWR) were removed. The remaining genes were sorted according to their Mean Decrease in the Gini Index or Mean Accuracy Decrease. RF was then run again with the same parameters for ten times each, to assess the OOB and the classification error of panels using the top 150, 100, 50 and 30 genes based on their Mean Decrease in the Gini Index and the Mean Accuracy Decrease values.

Results

Comparison between transcriptional profiling, otolith microchemistry and fatty acid pro-files. Among the 60 eels included in this study, 26 individuals were captured in SW, 7 in BW, and 27 in FW. Otolith analysis classified 11 of these eels as FWR, 23 as IHS, and 25 eels as SWR, and 1 sample was unclear (SWR/IHS). The fatty acid-based classification reported 19 animals as from FW, 19 as BW and 13 as SW, while the habitat of 8 eels remained undecided, and 1 not classified (Table S1).

The RNA-seq data production and subsequent trimming yielded 43,008,662 \pm 3,559,138 high-quality reads used for mapping. Approximately 76.71% \pm 2.69 of the reads were successfully mapped to the reference genome (Table S1).

Figure 2 shows the principal component analysis (PCA) based on the blood transcriptomic analysis, where 2 outliers were omitted. Sample clusterization by silvering stage did not show any evident grouping (Fig. 2a), and this factor was hence not considered a main driver of variation in this analysis. In contrast, sample clusterization by salinity habitat provided a clearer picture, with SW and FW eels occupying two close but distinctive clusters on the y-axis, and BW eels encompassing these clusters (Fig. 2b). Labelling according to the results from the otolith analysis also highlighted differences between FWR and SWR eels, with IHS samples plotted between the two habitats (Fig. 2c). Labelling according to fatty acid profile was concordant with the results provided by the previous classifications (Fig. 2d). The uncertain individuals were closer to the "FW" area of the plot for the FW/ BW and closer to the SW area for the SW/BW. Altogether, habitat salinity was a major driver for the clusterization of the transcriptomic profile, and otolith-based classification produced the lowest number of uncertainties in the classification, as well as a clear division between SWR and FWR.

Differential gene expression between salinity habitats. DE was calculated comparing FWR and SWR eels. The analysis detected 3451 genes with adjpvalue < 0.05. Among these, 1496 genes were upregulated when eels were classified as SWR and 1955 genes were upregulated when eels were classified as FWR. Four genes were removed from the analysis, as they were outliers based on visual inspection of the plot distribution (Fig. 3, table s2).

Enrichment analyses reported 143 enriched Gene Ontology (GO) biological terms for the genes more expressed among FWR, and 250 for genes that were more expressed among SWR. After removing redundant



Figure 2. Principal component analysis (PCA) of the samples based on log2transformed transcriptomic values labelled based on their sex, that is silvering stage for female (I, II, III) and male (**a**), sampling site (**b**), otolith-based history (**c**) and fatty-acid based history (**d**).

terms 76 GO terms for downregulated and 143 Go terms for the upregulated were retained (Tables S3 and S4). A total of 19 GO terms overlapped between the two gene sets, and this may indicate common pathways between the two environments (i.e., different genes involved in the same pathways). These terms are mainly related to regulation and organization of basic cells components, particularly RNA expression (e.g., signal transduction, regulation of transcription by RNA polymerase II, organelle organization, chromosome organization). Most GO terms revealed distinct pathways that may be activated in different salinity environments. In FWR, GO terms related with developmental processes, morphogenesis and immune functions (e.g., cranial skeletal system development, cartilage development, hematopoiesis, Wnt signaling and immune system development; Fig. 4 and Table S3). In SWR, the most relevant processes were linked to ATP production and ion transport, response to stress and fatty-acid oxidation (e.g., energy-coupled proton transport, electrochemical gradient, ATP metabolic process, electron transport chain, mitochondrial transport, fatty acid beta-oxidation; Fig. 4 and Table S4).

Repeated PCA based plot considering only the DE genes, but expanded on the whole sample set, enhanced the separation between salinity profiles (Fig. 5). The division was particularly clear for the otolith-based analysis, where the DE genes were detected (Fig. 5a). There, the only sample that could not be classified in the otolith analysis (labelled as "ind") was classified as IHS after the blood transcriptome-based positioning in the PCA plot. Despite being less prominent, a gradient was also detected in samples classified according to their lipid profiles. There, it was more evident that the eels with intermediate values of the M/F ratio (i.e., FW/BW and SW/BW) tended to be closer to the FW and SW part of the plot respectively (Fig. 5b).

Classification of eel into salinity habitats using random forest. RF classification, using the complete set of 3451 DE genes, correctly classified 79% of the eels into their salinity habitat (average OOB = 21%). Mis-assignments concerned mostly FWR and IHS eel (Table 1). The subset of genes further selected considering both Mean Decrease in the Gini Index and Mean Accuracy ranking improved the overall results, especially in the correct assignment of FWR individuals. For both ranking categories, correct classification increased by retaining only the top ranked genes until reaching a maximum value of rate of 93.1% (OOB = 6.9%) for both rankings (Table 1). The results were quite similar, with 100% of correctly assigned SWR and FWR (CCP FWR = 1 and



Figure 3. Distribution of differentially expressed genes, with genes more expressed in freshwater resident eels (blue) and genes more expressed in seawater resident eels (green). Black dots represents non-significant differentially expressed genes. X-axis reports the intensity of gene expression change expressed in Log2fold change.

positive regulation of transcription, DNA-templated	regulation of cell mmunication	regulation of signaling	regulation of transcription by RNA polymerase II	calclass-mediated signaling	cell-cell signaling by wnt	Wnt signaling pathway	demethylation	glycarryboszbolgist metricik process	displicity-odd pater oddath parent in the collination leafy sering paterny	electron transport chain GPI anchor	RNA nodification	cytoplasmic translation aromatic compound	nucleicacid Chosphodester bond hydrolysis mesendateg mesend	protein acylation	proteasomal protein catabolic process	gene expression	RNA splicing	cellular cellul cellul tesponse to	ar res o stre	sponse ss
regulation of gene expression posit	regulation of obtainer-containing compound about process tive re	regulation of biosynthetic gulation	positive regulation of cellular	calciu s	im-mec ignalin	liated g	demeti cellular pro	phosphorus metabolic hylation	Execution elongation Execution Execution Execution Execution Execution Execution Execution Execution	metabolic process histone acetylation	hatena notification	process biosynthetic process biosynthetic process	calular network compound metholoic process	protein protein naturation by (Fe-ric) cluster transfer	cellular amide metabolic process	eterocycle catabolic process	process caluter nitrigen compound calutoric process	estimulus esti cal photorec	ystem ,	egulation of calibre antice webbilic processa egulation
transcrip of cellular hacromolecula biosynthetic process pr	etabolic rocess	biological process reg	biological process pulation of	intracellular signal transduction	cellular response to stimulus	·	modificati process macromole modificati	ion s cule ion	ubi vi snRNA metabolic	quitin a the	depe multiv	ndent /esicu constato constato	t prote	ein cat dy so proteolysis	abolic ting	athwa	ess ay	cell differenti nervous s	ation tr	positive
regulation reg of nitrogen compound rei metabolic propess st	gulation of re sponse f to F	rocess regation cellular p rocess regation	iological process ve regulation of r macromolecule rithetic process	chromatin organization chro	cytosk organiz mosor	eleton zation	egulation molecular function	developmental process	process snRNA processing	nucleoside riphosphate metabolic process	process protein maturation	metabolic u process celular nacronolecula bioxynthetic process	ipicetosphan biosyntosic process lipid biosynthetic process	primary metabolic process cellular protein metabolic	ATP metabolic process methylatic	c signific cyclic compound metabolic process collula metaboli	1111	regul of i trans	ation on port	regulation of TOR feignating of TOR signaling
endothelial to hematopoietic	c c organ	C ANADORICAL Alfundaria	animalorgan	orga chromosome organization		on or	ganismal	biological regulation	RNA 3-end processing	ncRNA metabolic process	pepide metabolic process	crypine cyclic compound biosynthetic process	nacronolecula metabolic procesa	organic substance metabolic	heterocycl biosynthe moores	le protein ic metabol		protein folding	process utilizing autophag mechanis	process
cartilage	developme			regulatio	organiz on regu	ation of	omponent ganization	localization	proton Exceptort, down electrochemics product	mRNA transpor	t transp	ort ocaliz	NA ribo zation biog	some nitoci organosis	hondition vization org	acuole panization	complex subunit	cell cycle	calkiar component organization or biogenesi	cellular process
developmer cranial	emato	poietic	animal organ development	regu	lation	of re	sponse to	ovement of cell or subcellular component	ene	rgy co ransp	upled	l proto	on tone	ribor	ucleo	prote	- organeille in	biological regulation		elgnaling
development	antomica			of catalytic activity		decular S	timulus s	ignaling	elect	roche	mical	gradi	ent	compl	ex bio	gene	sis	response to stimulus	Actin Sameré-base procesa	oation homeostasis
hemopoiesis	structure developme	developmen	developmentel process	GTPas	se media	ited p	ellular rocess _{co}	cell mmunication	endropbernik reficulien to Golgi resicie-mediae		nitrogen compound transport	organic sub transpo	ort horse	ingrosiin x kubunit vitation	nutin comp organi	ular onentactin c zation orga	sytoskeleton anization	novement of cell or subcellular	cal-cal attenion an	

Figure 4. Treemap of the downregulated GO term (left) and upregulated Goterm (right). The nested rectangles, representing the branches of the tree, are not scaled. See Tables S3 and S4 for *p*-values and fold changes, as well as details of the names and genes.

CCP SWR=1) in all the four subsets (150-100-50 and 30 genes) and only four misplaced animals belonging to IHS in the 50 and 30 genes panels (CCP IHS=0.84; Table 1). Misclassified samples were B121, B128, B173 and B181 which were classified as IHS by otolith analyses, but as SWR by RF, reflecting the recent salinity habitat, (i.e., capture location in SW). This overlap in the results derives also from the fact that most genes, at least in the 30 genes panels (93% for the 30 gene panels), overlapped between the two rankings (Table 2). Among the top 30 genes for mean decrease Gini and mean accuracy, 10 genes (*mctp2a, inppl1b, asap1b, itk, adra1ba, rerea, tead1b, nelfb, acin1a* and *rev1*) are included in one or more biological processes previously detected in the enrichment analysis. The processes were mainly related to general metabolic and regulation processes, or developmental processes, hematopoiesis and immune system (Table S5).





		Mean decr	ease in the	gini	Mean accuracy decrease					
	All DE	150	100	50	30	150	100	50	30	
OOB error rate (%)	18.97	8.62	8.62	6.9	6.9	8.62	8.62	6.9	6.9	
CCP FWR	0.80	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	
CCP SWR	0.96	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	
CCP IHS	0.67	0.80	0.80	0.84	0.84	0.80	0.80	0.84	0.84	

Table 1. OOB error rate (%) and the CPP of the different gene sets: all differentially expressed, and top 150, 100, 50 and 30 bases on Mean Decrease in the Gini and Mean Accuracy Decrease. CCP = (1-classification error).

Discussion

The salinity-habitat history of European eel can be determined with reasonable accuracy using whole blood transcriptomic analysis. This method can replace or reduce lethal-based assessments (e.g., using otolith microchemistry, fatty acid analysis). This is relevant for fish species in general, but more specifically for sensitive and/ or (critically) endangered species, such as the European eel.

In the present study, RNA-seq performed on small blood amounts of 60 adult eels (58 after removing of outliers), collected in different salinity habitats (i.e., FW, SW, and BW) was first assessed through PCA plots based on the normalized transcriptomic count in relation to sampling location, otolith-based classification, fatty acid-based analysis, and to other external drivers linked to reproductive traits that can influence transcriptomics (i.e., silvering stage and sex). Here, we observed a pattern driven by salinity, independently of the means of classification (i.e., collection site, otolith microchemistry, or fatty acid profiles) in contrast to silvering stage.

Gene symbol	Gene name	chromosome	MeanDecreaseAccuracy	MeanDecreaseGini
rasal1	rasGAP-activating-like protein 1	NC_049210.1	0.0083	0.4798
mctp2a	Multiple C2 domains, transmembrane 2a	NC_049216.1	0.0041	0.2491
LOC118211351	Zinc finger E-box-binding homeobox 2-like	NC_049213.1	0.0033	0.2325
rnf138	E3 ubiquitin-protein ligase RNF138	NC_049204.1	0.0034	0.2300
LOC118227625	paired box protein Pax-5-like	NC_049205.1	0.0037	0.2274
fgd6	FYVE, RhoGEF and PH domain containing 6	NC_049207.1	0.0028	0.2197
inppl1b	Nositol polyphosphate phosphatase-like 1b	NC_049203.1	0.0034	0.2165
hic2	Hypermethylated in cancer 2 protein-like	NC_049210.1	0.0024	0.2131
LOC118231111	Osteoclast stimulatory transmembrane protein-like	NC_049207.1	0.0027	0.2094
LOC118208373	filamin-A-like	NC_049202.1	0.0034	0.2013
LOC118219836	CCN family member 1-like	NC_049211.1	0.0026	0.1967
LOC118225135	cadherin-24-like	NC_049204.1	0.0026	0.1909
LOC118229298	C-C motif chemokine 20-like	NC_049206.1	0.0025	0.1892
LOC118235156	uncharacterized LOC118235156	NC_049209.1	0.0023	0.1702
asap1b	ArfGAP with SH3 domain, ankyrin repeat and PH domain 1b	NC_049204.1	0.0021	0.1660
LOC118226320	Nuclear factor interleukin-3-regulated protein-like	NC_049204.1	0.0014	0.1520
Itk	IL2 inducible T cell kinase	NC_049203.1	0.0018	0.1493
adra1ba	adrenoceptor alpha 1Ba	NC_049209.1	0.0018	0.1461
LOC118233432	Dynein regulatory complex protein 11-like	NC_049208.1	0.0018	0.1420
Rerea	Arginine-glutamic acid dipeptide (RE) repeats a	NC_049213.1	0.0016	0.1390
tead1b	TEA domain family member 1b	NC_049203.1	0.0015	0.1293
hivep3	Transcription factor HIVEP3	NC_049201.1	0.0013	0.1276
LOC118235830	myelin-associated glycoprotein-like	NC_049201.1	0.0017	0.1271
LOC118211187	forkhead box protein J3-like	NC_049213.1	0.0016	0.1265
nelfb	Negative elongation factor complex member B	NC_049210.1	0.0012	0.1190
acin1a	Apoptotic chromatin condensation inducer 1a	NC_049208.1	0.0013	0.1189
mob3b*	mob kinase activator 3b	NC_049214.1	0.0011	0.1158
zap70	tyrosine-protein kinase ZAP-70	NC_049204.1	0.0013	0.1150
rev1*	REV1 DNA directed polymerase	NC_049215.1	0.0012	0.1143
LOC118229170	isthmin-2-like	NC_049206.1	0.0013	0.1113
agpat4+	1-acylglycerol-3-phosphate O-acyltransferase 4 (lysophosphatidic acid acyltransferase, delta)	NC_049218.1	0.0013	0.1046
epb41l1+	Band 4.1-like protein 1	NC_049213.1	0.0012	0.0995

Table 2. Genes included in the two 30-gene panel for random forest classification. *Gene unique for the 30top gene panel based on gini values. *Gene unique for the 30 top gene panel based on accuracy values.

.....

This clusterization was observed when considering a reduced 3451 DE gene dataset, which was then used for machine learning analysis. The number of studies investigating blood transcriptomics in non-mammals is very limited, and hence there is no map of candidate genes that could be affected by salinity changes in whole blood.

Physiology related to salinity habitats. Investigations on the adaptation of teleost fish to different salinity conditions have focused on gill and gut epithelium. In these tissues, hypersalinity (i.e., >35 ppt) leads to an increase of Na⁺, K⁺-dependent ATPase activities^{40–42}. In plasma, Na⁺ and Cl⁻ ion levels increase in eury-haline species⁴¹. In our study, SWR eels showed a higher expression of genes related to ATP production ("mito-chondrial ATP synthesis coupled electron transport", "ATP synthesis coupled electron transport", "ATP synthesis coupled proton transport", "ATP biosynthetic process"). Therefore, at least in the blood, the activity seems to go in the opposite direction—expressing genes that are involved in ATP production, particularly ATP synthase genes (atp5f1e, atp5l, atp5mc3a, atp5mea, atp5mf, atp5pd, atp5pf) and NADH dehydrogenase (ndufa10, ndufa7, ndufa8, ndufa11, ndufb8, ndufc2, ndufs2, ndufs6, ndufv3). Ion transport and homeostasis is in line with the plasma of other tissues, with genes related to ion homeostasis, and regulation pathways that are significantly enriched in SWR.

In European eel, diet and forage habitat may represent a major driver of variation in the fatty acid composition of eel muscle tissue⁴. In fact, fatty acid-based analysis has a high discriminating power across salinity habitats. This is also valid regarding blood when incorporating transcriptomic analysis, even with a lower power than otolith-based clusterization. Although not yet investigated in fish, erythrocyte fatty acid composition is considered one of the most stable biomarker for assessing long-term dietary intake or endogenous biosynthesis and metabolism, from several weeks to months prior, in humans⁴³. In this study, genes related to fatty acid oxidation and fatty acid beta-oxidation biological processes were more largely represented in the SWR group. Fatty acid oxidation is an important process for ATP production and/or fatty acid storage in muscle/adipose tissue. Indeed,



Figure 6. Otolith Sr:Ca (blue) and Ba:Ca (green) profiles of the four inter-habitat shifting eels (*Anguilla anguilla*) misclassified by the random forest analysis on blood transcriptomes.

at the same age SW eels were larger than FW eels²⁹. Fatty acid oxidation that mainly occurs in the mitochondria, involves a repeated sequence of reactions that result in the conversion of fatty acids to acetyl-CoA^{44,45}. This is concordant with what was observed in our data, where ATP synthesis is also upregulated in the SWR. In mammals, erythrocytes do not oxidize fatty acids because they lack mitochondrial or aerobic metabolism, thus relying on cytosolic energy generation instead. As for fish, even if there are reductions in biosynthetic processes and in the ability to mount heat shock responses with cell aging (common in other vertebrates), erythrocytes of fish do not appear to lose functions like the capacity to perform ion and gas transport, as total protein concentration in young red blood cells is not affected by cell age^{46,47}.

Osmoregulation is one of the most energetically costly metabolic activities in teleost, as both sea- and freshwater deviate from the salinity of fish body fluids⁴⁸. Therefore, a large amount of energy is consumed by fish to maintain their osmotic homeostasis during acclimation to either freshwater or hyper-saline water^{49,50}. At a molecular level, this can have an impact on fish development, growth, reproduction, and other physiological and metabolic activities⁵⁰. This is in line with the high number of identified GO terms that are linked to development and morphogenesis and to phenotypical observation of European eels, where size was bigger in SWR eels compared to FWR and IHS at the same age^{4,29}. About 300 genes that are linked to morphogenesis and development were upregulated in FWR and not in SWR. Since development is a complex matter that requires activation and suppression of several pathways, at the present state, it is not possible to provide a more detailed description of these pathways. To be noted is the presence of biological processes related to the immune system development and regulation among the pathways, which were enhanced in FWR. Variations in water salinity may impact the immune function of fish^{51,52}. Here, the GO term "cell-cell signaling by wnt" was detected. The Wnt signaling pathway is critical for adult tissue maintenance, remodeling and regeneration, embryo development, and many cellular processes that include cell motility and cytoskeleton restructuring⁵³. The Wnt signaling pathway is associated with cellular remodeling in fishes, and it may play a role in salinity adaptation^{54,55}. The Wnt signaling also regulates the progenitor cell homeostasis, hereby controlling hematopoiesis that was also upregulated in the FWR group. Even if two of the major genes involved in this pathway (i.e. cdc42 and wnt11) were not significantly differentially expressed, other genes such as ccdc88c, tet2, tet3 and lgr4, which have a known function in this pathway, were differentially expressed in our study⁵⁶⁻⁵⁸.

Biomarkers for classification of salinity-life history. RF performed using the DE genes as first list, and then reduced using different rankings allowed the correct assignment of all eel samples (except four) into their salinity habitat. The misclassification of these four samples was detected in all gene panels and were classified as IHS by otolith analysis and as BW by fatty acid analysis. Particularly, otolith analysis detected that those misclassified samples had early FW experience (Fig. 6). Otolith analysis can further characterize IHS, by estimating when and how many times, shifts between FW and SW habitats have occurred²⁹. This indicates that transcriptomics analysis may fail to show very early FW experience. However, in order to clarify this, more subcategories among the IHS group to be run with RF would be needed. Despite RF tolerating low sample numbers, subcategories of IHS group were not performed in RF analysis due to our relatively limited sample size.

Machine learning is useful to infer phenotypes to samples whose phenotype is unknown or not given in the analysis (i.e., test population) based on information collected from samples with the available phenotype(s) of interest (i.e., training population). In order to maximize the usage of our samples in this work, we did not split our dataset into a training and test. In fact, RF uses bootstrap to create random permutation in the data that leads to the definition of an OOB population for each individual tree in the forest. The OOB population consists of all the samples that are not included in the bootstrap population used to build a given single tree and that can be used to obtain internal unbiased estimates of the prediction error and to evaluate variable importance³⁹. The ability of transcriptomic-based biomarkers to distinguish with a relatively high level of accuracy (75%) eels with habitat-shifting history (IHS) from residents (FWR or SWR) regardless of the salinity at the sampling site suggests that salinity habitat history may leave a fingerprint in blood transcriptomics. This could hypothetically be through e.g., epigenetic mechanisms; however, further studies would be needed to decipher this aspect. In this context, controlled laboratory studies might help to identify how long these transcriptional signals reflecting different environments last.

Conclusions

Our investigation showed that blood transcriptional profile is influenced by salinity habitat history. The combination of machine learning and transcriptomic profiling allowed the assessment of salinity-habitat history with high accuracy, including habitat shifting behaviors. As habitat shifting behaviors may involve several shifts throughout the eel's continental life, additional studies will be needed to assess the extent of the prediction power of blood transcriptomics. Still, for complete reconstructions of IHS chronological salinity history, otolith microchemistry seems inevitable. However, with an adequate increase and maintenance of the training population, using otolith microchemistry analysis as known phenotype, there may be the possibility to greatly reduce, and in the future substitute, otolith microchemistry analysis for general estimation of fish salinity history. This approach is promising for the replacement or reduction of other lethal analyses in fish research, especially for critically endangered species, such as the European eel. Collecting blood and determining salinity-habitat history during annual monitoring surveys would provide important information improving management of this species. Overall knowledge on eel habitat use, such as migratory season and proportion of individuals that shift habitat, should be considered in the species monitoring. This could highlight the importance of this contingent (in terms of numbers) and the need to improve migratory conditions and habitat availability.

Data availability

Raw RNA-seq data are available on EMBL-EBI under Accession Number PRJEB52485.

Received: 19 May 2022; Accepted: 13 December 2022 Published online: 22 December 2022

References

- 1. Daverat, F. et al. Phenotypic plasticity of habitat use by three temperate eel species, Anguilla anguilla, A. japonica and A. rostrata. Mar Ecol. Prog. Ser. 308, 231 (2006).
- International Council for the Exploration of the Sea. JOINT EIFAAC/ICES/GFCM WGEEL REPORT 2014 Report of the Joint EIFAAC/ICES/GFCM Working Group on Eel. 203 (2014).
- Tzeng, W. N., Severin, K. P. & Wickström, H. Use of otolith microchemistry to investigate the environmental history of European eel Anguilla anguilla. Mar. Ecol. Prog. Ser. 149, 73 (1997).
- 4. Parzanini, C. *et al.* Trophic ecology of the european eel (*Anguilla anguilla*) across different salinity habitats inferred from fatty acid and stable isotope analysis. *Can. J. Fish. Aquat. Sci.* **78**, 1721 (2021).
- Stoot, L. J. et al. Use of portable blood physiology point-of-care devices for basic and applied research on vertebrates: A review. Conserv. Physiol. https://doi.org/10.1093/conphys/cou011 (2014).
- Ribeiro, M. N., Ribeiro, N. L., Bozzi, R. & Costa, R. G. Physiological and biochemical blood variables of goats subjected to heat stress: A review. J. Appl. Anim. Res. 46, 1036 (2018).
- Ahmed, I., Reshi, Q. M. & Fazio, F. The influence of the endogenous and exogenous factors on hematological parameters in different fish species: A review. Aquac. Int. 28, 869. https://doi.org/10.1007/s10499-019-00501-3 (2020).
- Minias, P. The use of haemoglobin concentrations to assess physiological condition in birds: A review. Conserv. Physiol. https:// doi.org/10.1093/conphys/cov007 (2015).
- Connon, R. E., Jeffries, K. M., Komoroske, L. M., Todgham, A. E. & Fangue, N. A. The utility of transcriptomics in fish conservation. J. Exp. Biol. https://doi.org/10.1242/jeb.148833 (2018).
- Wang, Z., Gerstein, M. & Snyder, M. RNA-Seq: A revolutionary tool for transcriptomics. Nat. Rev. Genet. https://doi.org/10.1038/ nrg2484 (2009).
- 11. Chu, Y. & Corey, D. R. RNA sequencing: Platform selection, experimental design, and data interpretation. *Nucleic Acid Ther.* 22, 271 (2012).
- Sudhagar, A., Kumar, G. & El-Matbouli, M. Transcriptome analysis based on RNA-Seq in understanding pathogenic mechanisms of diseases and the immune system of fish: A comprehensive review. *Int. J. Mol. Sci.* https://doi.org/10.3390/ijms19010245 (2018).
- Chaussabel, D. Assessment of immune status using blood transcriptomics and potential implications for global health. Semin. Immunol. https://doi.org/10.1016/j.smim.2015.03.002 (2015).
- He, Y., Maltecca, C., Tiezzi, F., Soto, E. L. & Flowers, W. L. Transcriptome analysis identifies genes and co-expression networks underlying heat tolerance in pigs. *BMC Genet.* 21, 1 (2020).
- 15. Jaing, C. *et al.* Gene expression analysis of whole blood RNA from pigs infected with low and high pathogenic African swine fever viruses. *Sci. Rep.* **7**, 1 (2017).
- 16. Scott, M. A. *et al.* Whole blood transcriptomic analysis of beef cattle at arrival identifies potential predictive molecules and mechanisms that indicate animals that naturally resist bovine respiratory disease. *PLoS ONE* **15**, e0227507 (2020).
- 17. Surlis, C. *et al.* Blood immune transcriptome analysis of artificially fed dairy calves and naturally suckled beef calves from birth to 7 days of age. *Sci Rep* **8**, 1 (2018).
- Chiari, Y. & Galtier, N. RNA extraction from sauropsids blood: Evaluation and improvement of methods. *Amphib. Reptil.* 32, 136 (2011).

- Waits, D. S., Simpson, D. Y., Sparkman, A. M., Bronikowski, A. M. & Schwartz, T. S. The utility of reptile blood transcriptomes in molecular ecology. *Mol. Ecol. Resour.* 20, 308 (2020).
- 20. Watson, H., Videvall, E., Andersson, M. N. & Isaksson, C. Transcriptome analysis of a wild bird reveals physiological responses to the urban environment. *Sci. Rep.* **7**, 1 (2017).
- 21. Banerjee, S. M. *et al.* Species and population specific gene expression in blood transcriptomes of marine turtles. *BMC Genom.* 22, 1 (2021).
- 22. Franchini, P. *et al.* Animal tracking meets migration genomics: Transcriptomic analysis of a partially migratory bird species. *Mol. Ecol.* **26**, 3204 (2017).
- Wang, Y., Guo, J., Wang, L., Tian, H. & Sui, J. Transcriptome analysis revealed potential mechanisms of differences in physiological stress responses between caged male and female magpies. *BMC Genom.* 20, 1 (2019).
- Rodríguez-Jorquera, I. A. et al. Blood transcriptomics analysis of fish exposed to perfluoro alkyls substances: Assessment of a Non-lethal sampling technique for advancing aquatic toxicology research. Environ. Sci. Technol. 53, 1441 (2019).
- Ding, W., Cao, L., Cao, Z. & Bing, X. Transcriptome analysis of blood for the discovery of sex-related genes in ricefield eel Monopterus albus. Fish. Physiol. Biochem. 46, 1507 (2020).
- Piles, M. et al. Machine learning applied to transcriptomic data to identify genes associated with feed efficiency in pigs. Genet. Sel. Evol. 51, 1 (2019).
- 27. Hastie, T., Tibshirani, R. & Friedman, J. Random forests. In The Elements of Statistical Learning (Springer, 2009).
- Durif, C., Dufour, S. & Elie, P. The silvering process of Anguilla anguilla: A new classification from the yellow resident to the silver migrating stage. J. Fish. Biol. 66, 1025 (2005).
- Rohtla, M., Daverat, F., Arts, M.T., Browman, H.I., Parzanini, C., Skiftesvik, A.B., Thorstad, E.B., van der Meeren, T., Vøllestad, L.A., Durif, C.M. Habitat use and growth of yellow-stage European eel in coastal and freshwater ecosystems in Norway. *Can. J. Fish. Aquat. Sci.* (2022).
- Folch, J., Lees, M. & Sloane Stanley, G. H. A simple method for the isolation and purification of total lipides from animal tissues. J. Biol. Chem. 226, 497 (1957).
- Parzanini, C. et al. Discrimination between freshwater and marine fish using fatty acids: Ecological implications and future perspectives. Environ. Rev. https://doi.org/10.1139/er-2020-0031 (2020).
- Andrews, S. FastQC: A quality control tool for high throughput sequence data. http://www.bioinformatics.babraham.ac.uk/proje cts/fastqc/. Babraham Bioinform. (2010).
- Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics* 30(15), 2114–2120. https://doi.org/10.1093/bioinformatics/btu170matic (2014).
- 34. Kim, D. *et al.* TopHat2: Accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genom. Biol.* **14**, 1 (2013).
- 35. Li, H. et al. The sequence alignment/map format and SAMtools. Bioinformatics 25, 2078 (2009).
- 36. Anders, S., Pyl, P. T. & Huber, W. HTSeq-A Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**, 166 (2015).
- 37. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genom. Biol.* **15**, 1 (2014).
- Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. Revigo summarizes and visualizes long lists of gene ontology terms. PLoS ONE 6, e21800 (2011).
- 39. Liaw, A. & Wiener, M. Classification and regression by randomForest. R News 2, 18 (2002).
- Cao, Q. et al. Physiological mechanism of osmoregulatory adaptation in anguillid eels. Fish Physiol. Biochem. https://doi.org/10. 1007/s10695-018-0464-6 (2018).
- 41. Gonzalez, R. J. The physiology of hyper-salinity tolerance in teleost fish: A review. J. Comp. Physiol. B Biochem. Syst. Environ. Physiol. https://doi.org/10.1007/s00360-011-0624-9 (2012).
- Kültz, D. Physiological mechanisms used by fish to cope with salinity stress. J. Exp. Biol. https://doi.org/10.1242/jeb.118695 (2015).
 Brenna, J. T., Plourde, M., Stark, K. D., Jones, P. J. & Lin, Y. H. Best practices for the design, laboratory analysis, and reporting of
- trials involving fatty acids. Am. J. Clin. Nutr. https://doi.org/10.1093/ajcn/nq089 (2018).
 44. Tocher, D. R. Metabolism and functions of lipids and fatty acids in teleost fish. Rev. Fish. Sci. https://doi.org/10.1080/713610925
- (2003).
 45. Console, L. *et al.* The link between the mitochondrial fatty acid oxidation derangement and kidney injury. *Front. Physiol.* https://doi.org/10.3389/fphys.2020.00794 (2020).
- Lund, S. G., Phillips, M. C. L., Moyes, C. D. & Tufts, B. L. The effects of cell ageing on protein synthesis in rainbow trout (Oncorhynchus mykiss) red blood cells. J. Exp. Biol. 203, 2219 (2000).
- Phillips, M. C. L., Moyes, C. D. & Tufts, B. L. The effects of cell ageing on metabolism in rainbow trout (*Oncorhynchus mykiss*) red blood cells. J. Exp. Biol. 203, 1039 (2000).
- 48. Bœuf, G. & Payan, P. How should salinity influence fish growth? Comp. Biochem. Physiol. C Toxicol. Pharmacol. 130, 411 (2001).
- 49. Si, Y. *et al.* Liver transcriptome analysis reveals extensive transcriptional plasticity during acclimation to low salinity in *Cynoglossus semilaevis*. *BMC Genom.* **19**, 1 (2018).
- 50. Zhang, X. *et al.* RNA-Seq analysis of salinity stress-responsive transcriptome in the liver of spotted sea bass (*Lateolabrax maculatus*). *PLoS ONE* **12**, e0173238 (2017).
- 51. Birrer, S. C., Reusch, T. B. H. & Roth, O. Salinity change impairs pipefish immune defence. *Fish. Shellfish Immunol.* **33**, 1238 (2012). 52. Tian, L., Tan, P., Yang, L., Zhu, W. & Xu, D. Effects of salinity on the growth, plasma ion concentrations, osmoregulation, non-
- specific immunity, and intestinal microbiota of the yellow drum (*Nibea albiflora*). Aquaculture 528, 735470 (2020).
 53. Angers, S. & Moon, R. T. Proximal events in Wnt signal transduction. Nat. Rev. Mol. Cell Biol. https://doi.org/10.1038/nrm2717
- 53. Angers, S. & Moon, R. I. Proximal events in Writ signal transduction. *Nat. Rev. Mol. Cell Biol.* https://doi.org/10.1038/nrm2/1/ (2009).
- 54. Cui, J. et al. Transciptome analysis of the gill and swimbladder of Takifugu rubripes by RNA-Seq. PLoS ONE 9, e85505 (2014).
- Jeffries, K. M. *et al.* Divergent transcriptomic signatures in response to salinity exposure in two populations of an estuarine fish. *Evol. Appl.* 12, 1212 (2019).
- Ekici, A. B. et al. Disturbed Wnt signalling due to a mutation in CCDC88C causes an autosomal recessive non-syndromic hydrocephalus with medial diverticulum. *Mol. Syndromol.* 1, 99 (2010).
- Glinka, A. *et al.* LGR4 and LGR5 are R-spondin receptors mediating Wnt/β-catenin and Wnt/PCP signalling. *EMBO Rep.* 12, 1055 (2011).
- Xu, Q. et al. Loss of TET reprograms Wnt signaling through impaired demethylation to promote lung cancer development. Proc. Natl. Acad. Sci. USA 119, e2107599119 (2022).

Acknowledgements

We thank Per Henning Gjuvsland for eel fishing; Nataliia Tiutcheva Bang for technical assistance; Computerome 2,0 for computing support; Päärn Paiste for helping with chemical analyses of otoliths, and Michael T. Arts for further financial help and mentorship. This work was financially supported by the Nordic Council of Ministers, grant number 200005.

Author contributions

F.B., C.D. and J.T. conceived the ideas and designed methodology; C.D., C.P., M.R. and F.B. collected the data. F.B., M.R. and C.P. analyzed the data; F.B. led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at https://doi.org/ 10.1038/s41598-022-26302-y.

Correspondence and requests for materials should be addressed to F.B.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2022