



## Validation of a QTL associated with resistance to *Vibrio anguillarum* in rainbow trout (*Oncorhynchus mykiss*)

Karami, Asma Mohammad; Marana, Moonika Haahr; Mathiessen, Heidi; Dalsgaard, Inger; Nielsen, Torben Fejer; Kania, Per Walter; Buchmann, Kurt

*Published in:*  
Acta Veterinaria Scandinavica

*Link to article, DOI:*  
[10.1186/s13028-023-00692-z](https://doi.org/10.1186/s13028-023-00692-z)

*Publication date:*  
2023

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Karami, A. M., Marana, M. H., Mathiessen, H., Dalsgaard, I., Nielsen, T. F., Kania, P. W., & Buchmann, K. (2023). Validation of a QTL associated with resistance to *Vibrio anguillarum* in rainbow trout (*Oncorhynchus mykiss*). *Acta Veterinaria Scandinavica*, 65, Article 28. <https://doi.org/10.1186/s13028-023-00692-z>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

BRIEF COMMUNICATION

Open Access



# Validation of a QTL associated with resistance to *Vibrio anguillarum* in rainbow trout (*Oncorhynchus mykiss*)

Asma Mohammad Karami<sup>1</sup>, Moonika Haahr Marana<sup>1</sup>, Heidi Mathiessen<sup>1</sup>, Inger Dalsgaard<sup>2</sup>,  
Torben Fejer Nielsen<sup>3</sup>, Per Walter Kania<sup>1</sup> and Kurt Buchmann<sup>1\*</sup> 

## Abstract

Vibriosis is a bacterial disease in fish caused by the Gram negative bacterium *Vibrio anguillarum* with severe impact on rainbow trout (*Oncorhynchus mykiss*) farming. Sustainable control methods should be developed and we here show that marker assisted selective breeding of fish naturally resistant to the disease is feasible. We have validated the use of a single nucleotide polymorphism (SNP) marker SNP AX-89,945,921 (QTL on chromosome 21). The QTL was previously found associated with resistance to vibriosis and described following a genome wide association analysis (GWAS) of trout exposed to the bacterium. For this validation spawners were genotyped by use of the 57 K Axiom® Trout Microarray (Affymetrix) and homozygous male fish carrying the allele with the SNP AX-89,945,921 were then selected and used to fertilize eggs from outbred female trout resulting in fish all carrying the SNP (QTL-fish). Control fish (non-QTL fish) were produced by fertilizing the same batch of eggs by use of male parents negative for the SNP. The fish were exposed in freshwater to *V. anguillarum* (water bath infection) at 19 °C. A total of 900 fish were challenged in a common garden set-up in triplicate. A bacterial solution of *V. anguillarum* (serotype O1) was added to each of three freshwater fish tanks, each with 150 QTL and 150 non-QTL fish. Fish were tagged by tail fin cut (upper/lower) to discern the two groups, whereafter fish were monitored around the clock to detect disease signs and remove moribund fish. Clinical vibriosis developed within two days in non-QTL-fish (overall morbidity of 70%). QTL fish developed clinical signs later and the morbidity was significantly lower and did not reach 50%. Rainbow trout farming may benefit from using the QTL associated with higher resistance towards vibriosis. The effect may be optimized in the future by use of both male and female parents homozygous for the marker allele.

**Keywords** Fish, Marker associated selective breeding, Single nucleotide polymorphism, SNP, Vibriosis

## Findings

*Vibrio anguillarum* is a Gram negative bacterium causing vibriosis in a range of wild and cultured fish species including salmonids [1]. Maricultured rainbow trout in Danish waters suffer from the disease throughout the production period [2], and although a commercial tri-valent vaccine is available it is not fully protective under field conditions [3]. An alternative approach to reach control of disease in aquacultured fish is based on selective breeding of naturally resistant fish [4]. The application of genome wide association analysis (GWAS) and

\*Correspondence:  
Kurt Buchmann  
kub@sund.ku.dk

<sup>1</sup> Department of Veterinary and Animal Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, DK-1870 Frederiksberg C, Denmark

<sup>2</sup> National Institute of Aquatic Resources, Technical University of Denmark, DK-2800 Kgs. Lyngby, Denmark

<sup>3</sup> Aquasearch ova ApS, Hedegaardsvej 8, DK-7190 Billund, Denmark



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

search for genetic markers has lifted the breeding efficacy significantly [5]. This has proven successful for identification of genomic markers in fish associated with higher resistance towards viral diseases [6–10], various bacterial diseases [11–13] and parasitic diseases [14, 15]. A genetic background for vibriosis resistance has been reported for Japanese flounder (*Paralichthys olivaceus*) [16, 17], tongue sole (*Cynoglossus semilaevis*) [18] and turbot (*Scophthalmus maximus*) [19], and we recently showed that *V. anguillarum* resistance in rainbow trout (*Oncorhynchus mykiss*) is associated with SNPs on rainbow trout chromosome 21 (Omy21) [20]. In order to validate this finding and seeking a practical application of the knowledge we have performed a comparative challenge study exposing trout with high and low frequencies of the SNP. We produced QTL-fish by fertilizing rainbow trout eggs with sperm from three brood stock males, which were homozygous for the SNP AX-89,945,921 (associated with *V. anguillarum* resistance [20]). Non-QTL-fish were produced by fertilizing eggs from the same batch with sperm from three males negative for the SNP. The eggs were from 30 outbred female spawners carrying the SNP AX-89,945,921 at a low frequency, 7% homozygous and 44% heterozygous for the SNP associated with resistance. It was estimated that the frequency of heterozygotes in the non-QTL fish (fertilized by negative males) was 29% and that 71% of the non-QTL fish were fully negative for the SNP. QTL fish group comprises 29% homozygotes and 71% heterozygotes. The typing was achieved by use of the 57 k SNP chip (Axiom®Trout, Affymetrix, San Diego, United States) [21].

The eggs from the breeding station were hatched and reared at the disease-free recirculated hatchery (Bornholm Salmon Hatchery). The fish population delivering eggs for the validation trial differed from the population delivering eggs for the original QTL study [20]. Thus, no genetic exchange had occurred for decades between these farms, Fårup Mølle in Jelling and Refsgaard in Egtved (Jutland, Denmark), respectively. The fish were fed by pelleted feed (Inicio, Biomar, Denmark). When reaching a mean body weight of 6 g fish (1200 degree days) the fish were transported (3 h) in oxygenated cooled freshwater to the infection facility at the University of Copenhagen and acclimated at 19 °C. Full survival was confirmed after transport. Before experimentation fish were acclimatized and tagged in the infection facility. For acclimatization and tagging fish were placed in separate fish tanks with aerated and recirculated (internal filters, Eheim, Germany) freshwater at 19 °C for one week. In order to differentiate fish during their pathogen exposure in a common garden set-up, we tagged the fish during the acclimatization week by tail cut under anaesthesia (immersion into 50 mg/L Tricaine Methane Sulphonate

MS222, Sigma-Aldrich, Denmark). Fish carrying a high frequency of the SNP associated with resistance (QTL fish) had their upper part of the tail fin cut. Non-QTL fish, with a low frequency of the SNP, had their lower tail fin cut.

Fish (mean weight 6 g, mean length 8 cm) were exposed at 19 °C to *V. anguillarum*. We used the same bacterial isolate (serotype O1, isolate 130,829–1/3) as applied for the challenge of rainbow trout on which the original GWAS and QTL description was based [20]. The bacterium was cultured in KB-medium over 72 h at 20 °C, and the fish were exposed by adding a solution ( $2.4 \times 10^9$  colony forming units (cfu)/mL) of the bacterium to the fish tanks. A common garden set-up was applied, whereby both QTL-fish and non-QTL fish were kept in the same fish tank securing that the two types of fish were exposed to similar environmental conditions. Fish were exposed for 2 h in triplicate aerated 100 L fish tanks (with a reduced water volume of 30 L) with  $2 \times 150$  fish in each with a final *V. anguillarum* concentration of  $3.12 \times 10^7$  cfu/mL. Following exposure water was replenished to 100 L and two internal Eheim biofilters were placed in each tank. The fish were kept in constant light during the entire period of observation. Control fish in three fish tanks were kept in a separate but similar room under corresponding conditions but without being exposed to *V. anguillarum*.

The rainbow trout were monitored around the clock by checking the fish every second hour. In case moribund fish, showing severe clinical signs (equilibrium disturbance, exophthalmia, skin ulcers), were noted, the affected fish were removed from the trial and euthanized in an over dosage of MS222 (Tricaine Methane Sulphonate 300 mg/L) and recorded as mortality. Head kidney swabs of the diseased fish were inoculated on blood-agar plates for confirmation of the causative pathogen [2].


Mortality curves for the six different groups were prepared by the Graph Pad Prism version 9.50. Data were first tested for normality by the D'Agostino and Pearson test. A two tailed paired Student's t-test using a probability level of 1% (0.01) was applied to test for differences between groups. First step was to compare mortality rates between replicates within both the non-QTL group and within the QTL-group. Finally the non-QTL fish groups were tested against the QTL fish groups (both for the replicates and the combined groups).

The first disease signs were observed on day 1 post-exposure (1 dpe) in one specimen of the non-QTL fish and from 2 dpe the number of diseased fish in this group increased markedly in all triplicates. Morbidity in non-QTL fish accelerated at 2 dpe before noon, where only one QTL-fish but 34 non-QTL fish died. Morbidity decreased later at 4 dpe and the curve flattened on

5 dpe, whereafter the experiment was terminated on 7 dpe. Thus, the percentage of fish affected in the three non-QTL-groups reached 61, 70 and 72% (Fig. 1, left). This mortality level was for all non-QTL groups significantly higher compared to the QTL groups ( $P < 0.01$ ). The QTL-fish showed a significantly lower disease rate and in none of the groups the overall percentage of fish with clinical signs surpassed 50% (Fig. 1, left). Within the two groups (QTL and non-QTL fish) the subgroups did not differ significantly, and when combined within groups the non-QTL fish showed an over-all morbidity rate of 70%, whereas the QTL-fish remained at 49% (Fig. 1, right). Also, when combined the morbidity rates were significantly lower in QTL-fish compared to non-QTL fish ( $P = 0.0035$ ). Control fish which were kept without *V. anguillarum* exposure showed no disease signs and mortality at all.

Vibriosis is considered an important disease problem in rainbow trout farming but even with vaccines aquaculture enterprises may benefit from use of naturally resistant fish strains [4, 20]. Breeding of disease resistant fish leads to improved health of aquacultured fish, which again will increase survival, improve feed efficiency and reduce usage of medicines such as antibiotics. Convincing results have been presented for Atlantic salmon (*Salmo salar*) with regard to viral diseases, including infectious pancreatic necrosis virus (IPNV) and salmonid alpha virus (SAV) [6, 7, 9, 10]. Loci associated with disease resistance in rainbow trout have been presented against the bacterium *Flavobacterium psychrophilum* [11–13], the bacterium *V. anguillarum* [20] and the parasite *Ichthyophthirius multifiliis* [14, 15]. Other studies suggested loci for vibriosis resistance in flatfish, such as Japanese flounder [16, 17], tongue sole [18] and turbot

[19]. This study has validated the QTL for vibriosis resistance in trout. We performed controlled infection experiments and confirmed that the QTL for natural resistance against *V. anguillarum* described by Karami et al. [20] may find a practical application and be used for selective breeding of resistant fish strains. It is noteworthy that the rainbow trout populations used for description of the QTL [20] is genetically different from the population used for validation in this study. No exchange of genetic material between these farms (Fårup Mølle and Refsgaard trout farms) has taken place for decades, which suggests a generic application of the QTL. We showed in a triplicate and controlled study that morbidity and thereby expected mortality could be reduced from 70% to less than 50% simply using fish with male parents homozygous for the SNP AX-89,945,921. The QTL offspring produced comprised both homozygous (estimated to 29%) and heterozygous trout (estimated to 71%). It should therefore be investigated if production of a batch of fully homozygous offspring can increase the natural resistance to *V. anguillarum*. In the present study we merely used homozygous male parents and outbred female spawners. Selection of both female and male spawners homozygous for the SNP is likely to increase the resistance of the rainbow trout offspring. We therefore hypothesize that the natural resistance of those fish towards vibriosis would be even higher. It is recommended to evaluate this approach in a future study. In this context genetic variance and dominance effects should be considered as well. Future experiments should therefore produce fully homozygous offspring by using male and female parents homozygous for the allele and evaluate their susceptibility/resistance to the vibriosis.



**Fig. 1** Left: Mortality curves (triplicate trials) for QTL fish and non-QTL fish following bath exposure to *V. anguillarum*. Right: Mortality curves (trials combined) for QTL fish and non-QTL fish following bath exposure to *V. anguillarum*. Mean percentage and SD shown

## Abbreviations

GWAS	Genome wide association analysis
IPNV	Infectious pancreatic necrosis virus
Omy21	<i>Oncorhynchus mykiss</i> chromosome 21
QTL	Quantitative trait loci
SAV	Salmonid alpha virus
SNP	Single nucleotide polymorphism

## Acknowledgements

Aquaculturist Gert Jørgensen, Bornholm Salmon Hatchery, hatched and reared the fish under disease free conditions.

## Prior publication

Data have not been published previously.

## Author contributions

KB and TFN designed the investigation, selected spawners and supervised production of the experimental fish. ID produced the bacterial culture for the challenge procedure. KB, MHM, HM and AMK conducted the infection process and monitored the fish during the experiment. KB and AMK drafted the manuscript and PWK performed the statistics and graphics. All authors have read and approved the final version of the manuscript.

## Funding

Open access funding provided by Royal Library, Copenhagen University Library. This research received funding from Innovation Fund Denmark, grant 8090-00002B (TECHFISH).

## Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

## Declarations

### Ethics approval and consent to participate

The study was conducted under license 2019-15-0201-01614 issued by the Experimental Animal Inspectorate, Danish Ministry of Environment and Food. The animals were handled according to high ethical standards and national legislation.

### Consent for publication

Not applicable.

### Competing interests

The author director Torben Nielsen, Aquasearch ova ApS, is producing trout eggs. The other authors declare that they have no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

Received: 13 March 2023 Accepted: 22 June 2023

Published online: 26 June 2023

## References

1. Amaro C, Fouz B, Sanjuan E, Romalde JL. Vibriosis. In: Leong JA, Buchmann K, Woo PTK, editors. Climate change and infectious fish diseases. Wallingford UK: CAB International; 2020. p. 182–210.
2. Pedersen K, Skall HF, Lassen-Nielsen AM, Nielsen TF, Henriksen NH, Olesen NJ. Surveillance of health status on eight marine rainbow trout, *Oncorhynchus mykiss* (Walbaum), farms in Denmark in 2006. *J Fish Dis*. 2008;31:659–67. <https://doi.org/10.1111/j.1365-2761.2008.00941.x>.
3. Marana MH, Sepulveda D, Chen D, Al-Jubury A, Jaafar RM, Kania PW, et al. A pentavalent vaccine for rainbow trout in danish aquaculture. *Fish Shellfish Immunol*. 2019;88:344–51.
4. Gjedrem T, Baranski M. The Success of Selective Breeding in Aquaculture. In: Gjedrem T, Baranski M, editors. Selective Breeding in Aquaculture: An Introduction. London New York: Springer Dordrecht Heidelberg; 2009. p. 13–22.
5. Yáñez JM, Barria A, Lopez ME, Moen T, Garcia BF, Yoshida GM, Xu P. Genome-wide association and genomic selection in aquaculture. *Rev Aquac*. 2022. <https://doi.org/10.1111/raq.12750>.
6. Houston RD, Haley CS, Hamilton A, Guy DR, Tinch AE, Taggart JB, et al. Major quantitative trait loci affect resistance to infectious pancreatic necrosis in Atlantic salmon (*Salmo salar*). *Genetics*. 2008;178:1109–15. <https://doi.org/10.1534/genetics.107.082974>.
7. Moen T, Baranski M, Sonesson AK, Kjøglum S. Confirmation and fine-mapping of a major QTL for resistance to infectious pancreatic necrosis in Atlantic salmon (*Salmo salar*): population-level associations between markers and trait. *BMC Genomics*. 2009;10:368.
8. Verrier ER, Dorson M, Mauger S, Torhy C, Ciobotaru C, Hervet C, et al. Resistance to rhabdovirus (VHSV) in rainbow trout: identification of a major QTL related to innate mechanisms. *PLoS ONE*. 2013;8:e55302. <https://doi.org/10.1371/journal.pone.0055302>.
9. Hillestad B, Makvandi-Nejad S, Krasnov A, Moghadam HK. Identification of genetic loci associated with higher resistance to pancreas disease (PD) in Atlantic salmon (*Salmo salar* L). *BMC Genomics*. 2020;21:388. <https://doi.org/10.1186/s12864-020-06788-4>.
10. Pavelin J, Jin YH, Gratacap RL, Taggart JB, Hamilton A, Verner-Jeffreys DW, et al. The nedd-8 activating enzyme gene underlies genetic resistance to infectious pancreatic necrosis virus in Atlantic salmon. *Genomics*. 2021;113:3842–50.
11. Wiens G, Vallejo RL, Leeds TD, Palti Y, Hadidi SS, Liu S, et al. Genetic correlation between cold water disease resistance and spleen index in a domesticated population of rainbow trout: identification of QTL on chromosome Omy19. *PLoS ONE*. 2013;8:e75749. <https://doi.org/10.1371/journal.pone.0075749>.
12. Fraslin C, Dechamp N, Bernard M, Krieg F, Hervet C, Guyomard R, Quillet E. Quantitative trait loci for resistance to *Flavobacterium psychrophilum* in rainbow trout: effect of the mode of infection and evidence of epistatic interactions. *Genet Sel Evol*. 2018;50:1–16. <https://doi.org/10.1186/s12711-018-0431-9>.
13. Fraslin C, Brard-Fudulea S, D'Ambrosio J, Bestin A, Charles M, Haffrey P, et al. Rainbow trout resistance to bacterial cold water disease: two new quantitative trait loci identified after a natural disease outbreak on a french farm. *Anim Genet*. 2019;50:293–7. <https://doi.org/10.1111/age.12777>.
14. Jaafar R, Ødegård J, Mathiessen H, Karami AM, Marana MH, Jørgensen LVG, et al. Quantitative trait loci (QTL) associated with resistance of rainbow trout *Oncorhynchus mykiss* against the parasitic ciliate *Ichthyophthirius multifiliis*. *J Fish Dis*. 2020;43:1591–602. [10.1111/jfd.13264](https://doi.org/10.1111/jfd.13264).
15. Buchmann K, Nielsen T, Mathiessen H, Marana MH, Duan Y, Jørgensen LVG, et al. Validation of two QTL associated with lower *Ichthyophthirius multifiliis* infection and delayed time-to-death in rainbow trout. *Aquac Rep*. 2022. <https://doi.org/10.1016/j.aqrep.2022.101078>.
16. Wang L, Fan C, Liu Y, Zhang Y, Liu S, Sun D, et al. A genome scan for quantitative trait loci associated with *Vibrio anguillarum* infection resistance in japanese flounder (*Paralichthys olivaceus*) by bulked segregant analysis. *Mar Biotechnol*. 2014;16:513–21. <https://doi.org/10.1007/s10126-014-9569-9>.
17. Sakamoto T, Chen S. Genome-wide SNP identification for the construction of a high-resolution genetic map of japanese flounder (*Paralichthys olivaceus*): applications to QTL mapping of *Vibrio anguillarum* disease resistance and comparative genomic analysis. *DNA Res*. 2015;22(2):161–70. <https://doi.org/10.1093/dnares/dsv001>.
18. Tang Z, Guo L, Liu Y, Shao C, Chen S, Yang G. Location of *Vibrio anguillarum* resistance-associated trait loci in half-smooth tongue sole *Cynoglossus semilaevis* at its microsatellite linkage map. *Chin J Oceanol Limn*. 2016;34(6):1309–19. <https://doi.org/10.1007/s00343-016-5160-8>.
19. Zhang K, Han M, Liu Y, Lin X, Liu X, Zhu H, et al. Whole-genome resequencing from bulked-segregant analysis reveals gene set based association analyses for the *Vibrio anguillarum* resistance of turbot (*Scophthalmus maximus*). *Fish Shellfish Immunol*. 2019;88:76–83. <https://doi.org/10.1016/j.fsi.2019.02.041>.

20. Karami AM, Ødegård J, Marana MH, Zuo S, Jaafar R, Mathiessen H, Buchmann KA. Major QTL for resistance to *Vibrio anguillarum* in rainbow trout. *Front Genet.* 2020;11:607558. <https://doi.org/10.3389/fgene.2020.607558>.
21. Palti Y, Gao G, Liu S, Kent MP, Lien S, Miller MR, Rexroad CE III, Moen T. The development and characterization of a 57K single nucleotide polymorphism array for rainbow trout. *Mol Ecol Res.* 2015;15:662–72. <https://doi.org/10.1111/1755-0998.12337>.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

