



Species which may act as vectors or reservoirs of diseases covered by the Animal Health Law

Listed pathogens of crustaceans

EFSA Panel on Animal Health and Welfare (AHAW); Nielsen, Søren Saxmose; Alvarez, Julio; Bicout, Dominique; Calistri, Paolo; Canali, Elisabetta; Drewe, Julian Ashley; Garin-Bastuji, Bruno; Gonzales Rojas, José Louis; Smith, Christian Gortazar

Total number of authors:
36

Published in:
EFSA Journal

Link to article, DOI:
[10.2903/j.efsa.2023.8172](https://doi.org/10.2903/j.efsa.2023.8172)

Publication date:
2023

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
EFSA Panel on Animal Health and Welfare (AHAW), Nielsen, S. S., Alvarez, J., Bicout, D., Calistri, P., Canali, E., Drewe, J. A., Garin-Bastuji, B., Gonzales Rojas, J. L., Smith, C. G., Herskin, M., Michel, V., Miranda Chueca, M. A., Padalino, B., Spooler, H., Ståhl, K., Velarde, A., Viltrop, A., Winckler, C., ... Roberts, H. (2023). Species which may act as vectors or reservoirs of diseases covered by the Animal Health Law: Listed pathogens of crustaceans. *EFSA Journal*, 21(8), Article e08172. <https://doi.org/10.2903/j.efsa.2023.8172>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ADOPTED: 21 June 2023

doi: 10.2903/j.efsa.2023.8174

Species which may act as vectors or reservoirs of diseases covered by the Animal Health Law: Listed pathogens of fish

EFSA Panel on Animal Health and Welfare (AHAW),
 Søren Saxmose Nielsen, Julio Alvarez, Dominique Bicoût, Paolo Calistri, Elisabetta Canali,
 Julian Ashley Drewe, Bruno Garin-Bastuji, José Louis Gonzales Rojas,
 Christian Gortazar Smith, Mette Herskin, Virginie Michel, Miguel Angel Miranda Chueca,
 Barbara Padalino, Hans Spoolder, Karl Ståhl, Antonio Velarde, Arvo Viltrop, Christoph Winckler,
 Isabelle Arzul, Shetty Dharmaveer, Niels Jørgen Olesen, Morten Schiøtt, Hilde Sindre,
 David Stone, Niccolò Vendramin, Mariana Aires, Inmaculada Aznar Asensio, Sotiria-Eleni
 Antoniou, Fulvio Barizzzone, Sofie Dhollander, Marzia Gnocchi, Anna Eleonora Karagianni,
 Linnea Lindgren Kero, Irene Pilar Munoz Guajardo, Alessia Rusina and Helen Roberts

Abstract

Vector or reservoir species of five fish diseases listed in the Animal Health Law were identified, based on evidence generated through an extensive literature review (ELR), to support a possible updating of Regulation (EU) 2018/1882. Fish species on or in which highly polymorphic region-deleted infectious salmon anaemia virus (HPRA ISAV), Koi herpes virus (KHV), epizootic haematopoietic necrosis virus (EHNV), infectious haematopoietic necrosis virus (IHNV) or viral haemorrhagic septicaemia virus (VHSV) were detected, in the field or during experiments, were classified as reservoir species with different levels of certainty depending on the diagnostic tests used. Where experimental evidence indicated transmission of the pathogen from a studied species to another known susceptible species, the studied species was classified as a vector species. Although the quantification of the risk of spread of the pathogens by the vectors or reservoir species was not part of the terms of reference, such risks do exist for the vector species, since transmission from infected vector species to susceptible species was proven. Where evidence for transmission from infected fish was not found, these were defined as reservoirs. Nonetheless, the risk of the spread of the pathogens from infected reservoir species cannot be excluded. Evidence identifying conditions that may prevent transmission by vectors or reservoir fish species during transport was collected from scientific literature. For VHSV, IHNV or HPRA ISAV, it was concluded that under transport conditions at temperatures below 25°C, it is likely (66–90%) they will remain infective. Therefore, vector or reservoir species that may have been exposed to these pathogens in an affected area in the wild, aquaculture establishments or through water supply can possibly transmit VHSV, IHNV or HPRA ISAV into a non-affected area when transported at a temperature below 25°C. The conclusion was the same for EHN and KHV; however, they are likely to remain infective under all transport temperatures.

© 2023 European Food Safety Authority. *EFSA Journal* published by Wiley-VCH GmbH on behalf of European Food Safety Authority.

Keywords: Vector, Reservoir, highly polymorphic region-deleted infectious salmon anaemia virus, Koi herpes virus, cyprinid herpesvirus-3, epizootic haematopoietic necrosis virus, infectious haematopoietic necrosis virus

Requestor: European Commission

Question number: EFSA-Q-2022-00457

Correspondence: ahaw@efsa.europa.eu

Panel members: Søren Saxmose Nielsen, Julio Alvarez, Dominique Bicout, Paolo Calistri, Elisabetta Canali, Julian Ashley Drewe, Bruno Garin-Bastuji, José Luis Gonzales Rojas, Christian Gortázar, Mette S Herskin, Virginie Michel, Miguel Ángel Miranda, Barbara Padalino, Paolo Pasquali (active until 5 April 2023), Helen Clare Roberts, Hans Spoolder, Karl Ståhl, Antonio Velarde, Arvo Viltrop and Christoph Winckler.

Declaration of interest: If you wish to access the declaration of interests of any expert contributing to an EFSA scientific assessment, please contact interestmanagement@efsa.europa.eu.

Acknowledgements: The Panel wishes to thank Christos Palaikostas and Ilaria Carmosino for contributing to the extensive literature review that formed the basis of the assessment.

Amendment: An editorial correction was carried out that does not materially affect the contents or outcome of this scientific output. In the second paragraph in the conclusions before Section 4.2 three species were added (*Carrassius Carassius*, *Diplodus sargus* and *Pagrus pagrus*). To avoid confusion, the original version of the output has been removed from the EFSA Journal, but is available on request.

Suggested citation: EFSA AHAW Panel (EFSA Panel on Animal Health and Welfare), Nielsen, S. S., Alvarez, J., Bicout, D., Calistri, P., Canali, E., Drewe, J. A., Garin-Bastuji, B., Gonzales Rojas, J. L., Smith, C. G., Herskin, M., Michel, V., Miranda Chueca, M. A., Padalino, B., Spoolder, H., Ståhl, K., Velarde, A., Viltrop, A., Winckler, C., ... Roberts, H. (2023). Species which may act as vectors or reservoirs of diseases covered by the Animal Health Law: Listed pathogens of fish. *EFSA Journal*, 21 (8), 1–45. <http://doi.org/10.2903/j.efsa.2023.8174>

ISSN: 1831-4732

© 2023 European Food Safety Authority. *EFSA Journal* published by Wiley-VCH GmbH on behalf of European Food Safety Authority.

This is an open access article under the terms of the [Creative Commons Attribution-NoDerivs](https://creativecommons.org/licenses/by-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited and no modifications or adaptations are made.

EFSA may include images or other content for which it does not hold copyright. In such cases, EFSA indicates the copyright holder and users should seek permission to reproduce the content from the original source.



The EFSA Journal is a publication of the European Food Safety Authority, a European agency funded by the European Union.



Summary

Term of Reference 1 (ToR 1) requested the European Food Safety Authority (EFSA) to assess which species or groups of species of aquatic animals pose a considerable risk for spreading pathogens causing the diseases of aquatic species listed in COMMISSION IMPLEMENTING REGULATION (EU) 2018/1882 EU Regulation 2016/429. This Opinion specifically focuses on assessing vector or reservoir species of five diseases of fish, i.e. highly polymorphic region-deleted infectious salmon anaemia virus (Δ ISAV), Koi herpes virus (KHV) (CyHV3), epizootic haematopoietic necrosis virus (EHNV), infectious haematopoietic necrosis virus (IHNV) and viral haemorrhagic septicaemia virus (VHSV). The aim of the assessments is to indicate if the Annex to Implementing Regulation (EU) 2018/1882, listing those vectors or reservoir species, needs to be updated. EFSA was not requested to update the list of susceptible species, listed in the same Implementing Regulation, as this work is already being coordinated by the Reference laboratories of the EU and the World Organisation for Animal Health (WOAH). In addition, it was agreed that a species cannot be classified as both susceptible and vector or reservoir species.

The following working definitions were agreed for the assessment: a fish species can be considered a **vector** when the pathogen has been identified in or on the fish species and it has been demonstrated to transmit the pathogen to susceptible species. To be considered a **reservoir** species, the pathogen should have been identified in or on the fish species, but evidence of transmission of the pathogen to susceptible species could not be found. It should be cautioned, however, that these are working definitions to address the term of reference. A clear separation between reservoir, vectors and susceptible species is not always easily made in the field, especially for aquatic animal diseases.

Although the quantification of the risk of spread of the pathogens by the vectors or reservoir species was not part of the terms or reference, such risks do exist for the vector species, since transmission from infected vector species to susceptible species was proven. Where evidence for transmission from infected fish was not found, these were defined as reservoirs. Nonetheless, the risk of the spread of the pathogens from infected reservoir species cannot be excluded.

An ELR has been carried out to gather all published peer-reviewed scientific evidence available on parameters needed to assess the role of aquatic species (only fish species) as vectors or reservoirs. The detailed methods for searching the literature, study selection, data collection and quality assurance are described in detail in EFSA (2023). The data, extracted from the eligible literature, were assessed in two steps. In the first step, the working group experts individually identified those studies where the target pathogens were detected with reference tests (i.e. WOAH, EURL) in or on fish species, either in experimental or field settings, with a high (> 90%) certainty. This immediately led to the classification as reservoir or vector species (the latter only for experimental studies with proven transmission of the pathogen from the vector to the susceptible species). Also, those studies were identified that led to a clear exclusion of the species as vector or reservoir (> 90% certainty) due to negative test results.

In a second step, the studies with inconclusive (< 66% certainty) results were discussed in smaller groups and then consolidated by the whole working group. The cut-off level for classifying species as vectors or reservoirs was set at a minimum certainty of 66%.

The results of the assessment indicated that:

The following species is considered a reservoir for **EHNV** with > 90% certainty: *Tandanus tandanus* (dewfish).

The following species are considered reservoir species for **EHNV** with 66–90% certainty: *Maccullochella peelii* (Murray cod), *Macquaria novemaculeata* (Australian bass) and *Macquaria ambigua* (callop).

The following species are considered vector species for **KHV** with > 90% certainty: *Carassius auratus* (goldfish), *Carassius gibelio* (Prussian carp), *Ctenopharyngodon idella* (grass carp), *Gymnocephalus cernua* (Eurasian ruffe), *Hypophthalmichthys molitrix* (silver carp), *Rutilus rutilus* (common roach) and *Tinca tinca* (tench).

The following species are considered reservoir species for **KHV** with > 90% certainty: *Acipenser gueldenstaedtii* (Russian sturgeon), *Acipenser oxyrinchus* (Atlantic sturgeon), *Acipenser ruthenus* \times *Huso huso* (hybrid sterlet \times beluga), *Barbatula barbatula* (stone loach), *Gasterosteus aculeatus* (three-spine stickleback), *Perca fluviatilis* (European perch) and *Scardinius erythrophthalmus* (Pearl roach).

The following species are considered reservoir species for **KHV** with 66–90% certainty: *Oreochromis niloticus* (Nile tilapia) and *Pseudorasbora parva* (topmouth gudgeon).

The following species are considered reservoir species for **HPRA ISAV** with > 90% certainty: *Clupea harengus* (Atlantic herring), *Oncorhynchus masou* (masu salmon) and *Oncorhynchus kisutch* (coho salmon).

The following species are considered reservoir species for **HPRA ISAV** with 66–90% certainty: *Gadus morhua* (Atlantic cod), *Oncorhynchus keta* (chum salmon).

The following species are considered reservoir species for **IHN** with > 90% certainty: *Aulorhynchus flavidus* (tube-snout), *Acipenser transmontanus* (white sturgeon), *Clupea pallasii pallasii* (Pacific herring), *Cyprinus carpio* (common carp), *Danio rerio* (zebrafish), *Perca flavescens* (American yellow perch) and *Seriola quinqueradiata* (amberjack).

The following species are considered reservoir species for **IHN** with 66–90% certainty: *Cymatogaster aggregata* (shiner perch) and *Oncorhynchus gorbuscha* (pink salmon).

The following species are considered reservoir species for **VHSV** with > 90% certainty: *Anguilla anguilla* (European eel), *Argentina sphyraena* (lesser argentine), *Belone belone* (garfish), *Cottus pollux* (Japanese fluvial sculpin), *Cyprinus carpio* (common carp), *Enchelyopus cimbrius* (fourbeard rockling), *Epinephelus akaara* (Hong Kong grouper), *Esox lucius* × *Esox masquinongy* (tiger muskellunge), *Eutrigla gurnardus* (grey gurnard), *Gadiculus argenteus* (silvery pout), *Gadus chalcogrammus* (Alaska pollock), *Hippoglossus hippoglossus* (Atlantic halibut), *Ictalurus punctatus* (channel catfish), *Merluccius productus* (Pacific hake), *Moxostoma anisurum* (silver redhorse), *Moxostoma macrolepidotum* (shorthead redhorse sucker), *Oncorhynchus mykiss* (rainbow trout) × *Salvelinus fontinalis* (American brook charr), *Oryzias latipes* (Japanese rice fish), *Pagrus major* (red sea bream), *Percopsis omiscomaycus* (trout perch), *Petromyzon marinus* (sea lamprey), *Pomoxis annularis* (white crappie), *Rhinogobius* sp. (Yoshinobori complex; Japanese goby), *Reinhardtius hippoglossoides* (Greenland halibut), *Salvelinus alpinus* (Arctic char), *Scorpaena porcus* (black scorpionfish), *Sebastes inermis* (black rockfish) and *Seriola quinqueradiata* (amberjack).

The following species are considered reservoir species for **VHSV** with 66–90% certainty: *Alosa pseudoharengus* (alewife), *Anoplopoma fimbria* (sablefish), *Carassius auratus* (goldfish), *Fundulus diaphanous* (banded killifish), *Hypomesus pretiosus* (surf smelt), *Lota lota* (burbot), *Notemigonus crysoleucas* (golden shiners), *Oncorhynchus mykiss* (rainbow trout) × *Salmo trutta* (Amu-Darya trout), *Oncorhynchus mykiss* (rainbow trout) × *Salvelinus alpinus* (Alpine Char), *Oncorhynchus mykiss* (rainbow trout) × *Salvelinus namaycush* (American Lake Charr).

In addition, a list of vector or reservoir species for which no peer reviewed studies with evidence of its role as vector or reservoir was found, are suggested to be removed from the current list Commission Implementing Regulation 1882/2018.

Term of Reference 2 (ToR 2) requested EFSA to assess the suitability of the conditions under which fish species should be regarded as vectors or reservoirs for the purposes of movements. These conditions are set out in Annex I to Commission Delegated Regulation (EU) 2020/990 and in Annex XXX to Commission Delegated Regulation (EU) 2020/692. Alternative conditions had to be proposed, if the conditions in those Regulations were considered not to prevent the transmission of the targeted pathogen via movement of vectors or reservoirs.

To provide a concise answer within the time frame of the mandate, it was decided to focus the assessment on those conditions that would prevent transmission facilitated by the movement of vectors and reservoirs, for which scientific evidence was available. In a first step, the experts in the working group carried out a narrative literature review to collect any evidence from scientific literature identifying conditions that may prevent transmission by vectors. In addition, information on the duration of the experimental studies and the water temperature were compiled during the ELS carried out for ToR 1, collecting the ranges of the different durations and temperatures for which transmission has been proven for the different pathogens by the different vector species. Then, the experts concluded by consensus if the collected evidence was sufficient to support the need to alter the conditions stipulated in Annex I to Commission Delegated Regulation (EU) 2020/990 and in Annex XXX to Commission Delegated Regulation (EU) 2020/692.

For VHSV, IHNV or HPRA ISAV, it was concluded that under transport conditions at temperatures below 25°C, it is likely (66–90% certainty) that VHSV, IHNV and HPRA ISAV will remain infective.

Therefore, vector or reservoir species that may have been exposed to VHSV, IHNV or HPRA ISAV in an affected area can possibly transmit VHSV, IHNV or HPRA ISAV when transported at a temperature below 25°C into a non-affected area. Exposure in a VHSV, IHNV or HPRA ISAV affected area may have occurred if the vector or reservoir originate from (a) an aquaculture establishment or group of

aquaculture establishments, where susceptible species or reservoir or other vector species are kept, (b) the wild, where they may have been exposed to susceptible, reservoir or other vector species or (c) an aquaculture establishment supplied with water possibly contaminated with VHSV, IHNV or HPRΔ ISAV.

For EHN and KHV, it was concluded that both pathogens may remain infective under all transport temperatures. Therefore, vectors or reservoir species can transmit these viruses when they originate from the same sources as listed in points (a), (b) and (c) above, when transported at any temperature. It was suggested that these conclusions may be considered for amendments to Annex I of Reg 2020/990 and Annex XXX of Reg 2020/692.

Table of contents

Abstract.....	1
Summary.....	3
1. Introduction.....	7
1.1. Background and Terms of Reference as provided by the requestor.....	7
1.2. Interpretation of the Terms of Reference.....	8
1.2.1. Term of Reference 1: Assessment of potential vectors and reservoir species of diseases of fish, crustaceans and molluscs listed in Annex II to the AHL.....	8
1.2.2. Term of Reference 2: Conditions under which fish species shall be regarded as vectors or reservoirs of diseases of fish listed in Annex II to the AHL.....	9
2. Data and methodologies.....	9
2.1. Methodologies.....	9
2.1.1. Term of Reference 1: Assessment of potential vectors and reservoir species of pathogens of fish, listed in Annex II to the AHL.....	9
2.1.1.1. First step: Individual assessments by the fish experts of the data extracted from specific papers assigned to them.....	10
2.1.1.2. Second step: group discussion.....	11
2.1.2. Term of Reference 2: Conditions under which fish species shall be regarded as vectors or reservoirs of diseases of fish listed in Annex II to the AHL.....	11
2.2. Data.....	12
2.2.1. Term of Reference 1: Assessment of potential vectors and reservoir species of diseases of fish listed in Annex II to the AHL.....	12
2.2.2. Term of Reference 2: Conditions under which fish species shall be regarded as vectors or reservoirs of diseases of fish listed in Annex II to the AHL.....	12
3. Assessment.....	12
3.1. Term of Reference 1: Assessment of potential vectors and reservoir species of diseases of fish listed in Annex II to the AHL.....	12
3.2. Term of Reference 2: Conditions under which fish species shall be regarded as vectors or reservoirs of pathogens listed in Annex II to the AHL.....	27
4. Conclusions.....	28
4.1. Term of Reference 1: Assessment of potential vectors and reservoir species of diseases of fish listed in Annex II to the AHL.....	28
4.2. Term of Reference 2: Conditions under which fish species shall be regarded as vectors or reservoirs of diseases of fish listed in Annex II to the AHL.....	30
References.....	31
Abbreviations.....	35
Appendix A – Currently listed vector or reservoir species without sufficient evidence in peer-reviewed papers. ...	36
Appendix B – Vector or reservoir species mentioned in peer-reviewed papers that were excluded in the ELR	39
Annex A – Protocol.....	45

1. Introduction

1.1. Background and Terms of Reference as provided by the requestor

In accordance with Article 8 of Regulation (EU) 2016/429 (AHL), the disease-specific rules for listed diseases provided in the AHL, and the rules adopted pursuant to that Regulation, apply to listed species. In compliance with that Article, the Commission shall establish a list of animal species or groups of species, which pose a considerable risk for the spread of specific listed diseases based on the capability of those animals to carry those specific diseases. Animal species or groups of animal species shall only be added to the list if they pose a considerable risk for the spread of a specific listed disease because they are vectors or reservoirs for that disease, or scientific evidence indicates that such role is likely.

The list of vector species, which is set out in the fourth column of the table in the Annex to Implementing Regulation (EU) 2018/1882, was carried forward from the list, which was previously set out in Commission Regulation (EU) 1251/2008. The Commission now requires scientific advice to inform an amendment to that list, to ensure that only species, which comply with Article 8 of the AHL, are listed. This amendment may involve species, which are currently set out in the fourth column of the Annex to Implementing Regulation (EU) 2018/1882 being removed and/or new species being added to that list.

It should be noted that vector species of aquatic animals are not listed in the WOAHL Aquatic Code¹ or in the WOAHL Aquatic Manual.² In the disease specific chapters of the WOAHL Aquatic Manual however, as well as listing susceptible species, other species which have shown incomplete evidence of susceptibility are listed, as are species in which PCR positive results have been reported, but where an active infection has not been demonstrated. In 2020, the EU Reference Laboratories (EURLs) for fish, crustaceans and molluscs, with the assistance of experts, reviewed those non-susceptible species, which are listed in the WOAHL Manual, in an effort to determine whether or not, they could be considered to be vectors of specific listed diseases. The reports which have been prepared by the EURLs and which have been furnished to the Commission, may be of assistance to the risk assessor in providing the scientific advice which is currently sought. The three reports (concerning fish, molluscs and crustaceans) accompany this letter. It should, however, be noted that these reports also contain information concerning susceptible species to the listed diseases, which is not pertinent to this request for a scientific opinion.

In addition, for those species and groups of species referred to above, which should be listed in accordance with Article 8 of the AHL, scientific advice is also required concerning the conditions under which these species should be regarded as vectors or reservoirs for the purposes of movements.

The conditions under which these species should be regarded as vectors are set out in Annex I to Commission Delegated Regulation (EU) 2020/990³ and in Annex XXX to Commission Delegated Regulation (EU) 2020/692⁴. It should be noted that the conditions set out in Annex I to Commission Delegated Regulation (EU) 2020/990 are not identical to the conditions set out in Annex XXX to Commission Delegated Regulation (EU) 2020/692, and both sets of conditions are different to those which were previously set out in columns 3 and 4 of Annex I to Commission Regulation (EC) 1251/2008.

Terms of reference

In view of the above, the Commission asks EFSA for a scientific opinion on the listing of vector species of aquatic animals in accordance with Article 8 of Regulation (EU) 2016/429, as follows:

- 1) For each of the aquatic diseases listed in Annex II to the AHL, an assessment concerning which species or groups of species of aquatic animals pose a considerable risk for their spread based on the fact that:
 - i) they are vector species or reservoirs for that disease, or
 - ii) scientific evidence indicates that such role is likely.

¹ WOAHL Aquatic Animal Health Code, 2021, 23rd Edition.

² WOAHL Aquatic Manual, 2021, 8th Edition.

³ Commission Delegated Regulation (EU) 2020/990 of 28 April 2020 supplementing Regulation (EU) 2016/429 of the European Parliament and of the Council, as regards animal health and certification requirements for movements within the Union of aquatic animals and products of animal origin from aquatic animals (OJ L 221, 28 April 2020, p. 42).

⁴ Commission Delegated Regulation (EU) 2020/692 of 30 January 2020 supplementing Regulation (EU) 2016/429 of the European Parliament and of the Council as regards rules for entry into the Union, and the movement and handling after entry of consignments of certain animals, germinal products and products of animal origin (OJ L 174, 3.6.2020, p. 379).

For each of the species or groups of species, which are assessed to be vector species or reservoirs of the listed diseases, or where scientific evidence indicates that such role is likely, they should be aquatic animals, which are not already listed as susceptible to the listed disease.

- 2) For each of the species or groups of species, which are assessed to fulfil the requirements for listing by virtue of being a vector or reservoir of a listed disease, or where scientific evidence indicates such a role is likely, an assessment of the suitability of the conditions under which they should be regarded as vectors or reservoirs for the purposes of movements. These conditions are set out in Annex I to Commission Delegated Regulation (EU) 2020/990 and in Annex XXX to Commission Delegated Regulation (EU) 2020/692, however, alternative conditions should be proposed, if the conditions, which are set out in those Regulations, are assessed to be unsuitable.

1.2. Interpretation of the Terms of Reference

1.2.1. Term of Reference 1: Assessment of potential vectors and reservoir species of diseases of fish, crustaceans and molluscs listed in Annex II to the AHL

Term of Reference 1 (ToR 1) requests EFSA to provide a list of vector species or reservoir species of pathogens of fish, crustaceans and molluscs, listed in Annex II to the AHL, aiming to update the fourth column of the Annex to Implementing Regulation (EU) 2018/1882.

EFSA was not requested to update the list of susceptible species, already listed in the third column of the same Implementing Regulation. In addition, it was agreed that a species cannot be classified simultaneously as susceptible as well as vector or reservoir species.

This work is complementary to the work that was coordinated by the EURL and WOAHC concerning the identification of susceptible species.

This Scientific Opinion focuses on all life stages, including eggs, sperm and gametes of **fish** belonging to the superclass Agnatha and to the classes Chondrichthyes, Sarcopterygii and Actinopterygii. The pathogens listed by the AHL affecting fish are:

- Highly polymorphic region (HPR)-deleted infectious salmon anaemia virus (HPRΔ ISAV)
- Koi herpes virus (KHV) = Cyprinid herpesvirus-3 (CyHV3)
- Epizootic haematopoietic necrosis virus (EHNV)
- Infectious haematopoietic necrosis virus (IHNV)
- Viral haemorrhagic septicaemia virus (VHSV)

It was agreed that for this assessment, a fish species can be considered a **vector** when the pathogen has been identified in or on the species and it has been demonstrated to transmit the pathogen to susceptible species, or there is scientific evidence that indicates that this transmission is likely. In addition, the vector species must not already be listed as susceptible to the respective pathogen.

Vectors may transmit pathogenic agents to susceptible species in two ways: (i) the pathogenic agent can multiply within the vector's body and then be transmitted to other susceptible species; (ii) the pathogenic agent can remain in or on the vector without multiplying and be mechanically transmitted to other susceptible species.

To be considered a **reservoir** species, on the other hand, the pathogen has been identified in or on the fish species, but evidence of transmission of the pathogen to susceptible species is not available. In addition, it was agreed that the reservoir species must not already be listed as susceptible to the respective pathogen.

It should be cautioned, however, that these are working definitions to address the term of reference. A clear separation between reservoir, vectors and susceptible species is not always easily made on the basis of field observations alone, and for aquatic animal diseases in particular.

Although the quantification of the risk of spread of the pathogens by the vectors or reservoir species was not part of the terms of reference, such risks do exist for the vector species, since transmission from infected vector species to susceptible species was proven. Where evidence for transmission from infected fish was not found, these were defined as reservoirs. Nonetheless, the risk of the spread of the pathogens from infected reservoir species cannot be excluded.

1.2.2. Term of Reference 2: Conditions under which fish species shall be regarded as vectors or reservoirs of diseases of fish listed in Annex II to the AHL

The list of potential vectors and reservoir species developed in ToR 1 should be considered as vectors or reservoirs for movements in the EU, provided that certain conditions are fulfilled.

The conditions in the EC Delegated Reg 2020/990 Annex I specify that the species may be regarded as vectors when animals are present in: (a) an aquaculture establishment or group of **aquaculture establishments** where susceptible species listed in column 3 of that table in Annex 1, are kept; or (b) the **wild** where they can be **exposed to susceptible species** listed in column 3 of that table.

The conditions in EC Delegated Reg 2020/692 Annex XXX stipulate that Vectors may be regarded as the species that have been in contact with listed susceptible species listed in column 3 of the table in the Annex to Commission Implementing Regulation (EU) 2018/1882 through **co-habitation or through water supply**.

It should be noted that although these two delegated acts explicitly mention vectors, it is assumed that the same conditions apply for reservoirs. Thus, when vector species and reservoir species do not fulfil these conditions, they can be moved provided that the transport complies with the EU regulations and all the measures have been implemented which would prevent the contamination or infection of the transported species.

To address ToR 2, besides the conditions already laid down in EU Reg 2020/990 Annex I and Reg EC Delegated 2020/692 there are other conditions that need to be fulfilled by a species to be considered a vector. Evidence found in the scientific literature related to the above factors for the specific pathogens was scrutinised and summarised. If there was no proof that certain specific conditions can exclude that the fish species will act as potential vector or reservoir, there was no change in the conditions already laid down in the above-mentioned regulations.

2. Data and methodologies

2.1. Methodologies

2.1.1. Term of Reference 1: Assessment of potential vectors and reservoir species of pathogens of fish, listed in Annex II to the AHL


An extensive literature review (ELR) has been carried out to gather all scientific evidence available on parameters needed to assess the role of aquatic species as vectors or reservoir species of specific pathogens of fish, listed by the AHL. To assess the evidence the following review questions were posed:

Review questions:

- 1) For vector species: What is the evidence generated by experimental infection studies, demonstrating transmission of 'Pathogen A' from 'Vector species X' on- or in which Pathogen A was detected, to a 'Species Y'?



- 2) For reservoir species: What is the evidence generated by experimental infection studies or field studies, demonstrating the detection of Pathogen A on- or in Reservoir species X, without further evidence of transmission of pathogen A to a 'Species Y'?



As agreed in the interpretation of the ToRs, to define a vector species, proof of onwards transmission from species X to species Y is needed. This proof is usually not available from field detections. Field detections imply Pathogen A was detected in species X during outbreak investigations, prevalence studies or any other study where the pathogen was detected in fish in a specific area or farm. In these situations, it cannot be definitively proven that species Y has been infected through species X and not from any other source of infection.

The detailed methods for searching the literature, the study selection, data collection and quality assurance are described in detail in EFSA (2023).

The data set was generated with the relevant information extracted from the eligible literature needed to answer the review questions. Then, the assessment methodology for deciding if the information was sufficient to classify the fish species as a potential vector or reservoir species, according to the working definition provided in Section 1.2.1, was applied in two steps.

2.1.1.1. First step: Individual assessments by the fish experts of the data extracted from specific papers assigned to them

Questions:

How certain are you that species X is a RESERVOIR species based on the evidence generated through the ELR (for field and experimental infection studies that did not investigate species Y and field studies where infection of species Y could not be proven)?

How certain are you that species X is a VECTOR species based on the evidence generated through the ELR (for experimental infection studies that have also investigated infection of species Y)?

In the first step, the experts were asked to identify the species for which a clear 'yes' or 'no' could be answered on the above questions with a high certainty (**> 90% certainty**). The experts were asked to provide the reasoning for their choice and reminded to respect the working definition of vectors and reservoirs, and not to consider other information that was not collected or extracted from the eligible peer-reviewed literature that was outside the scope of the working definition, e.g. on observed clinical signs.

As a guidance to help the decision-making, the following criteria were agreed a priori among the experts:

• Positive results (> 90% certainty):

Experimental infections: there is higher certainty when evidence from experimental infections is available compared to field studies because the animals are infected under controlled conditions so there is no need for sequencing or confirmatory tests. The following tests are accredited by the EU reference laboratory (EURL) and the WOAHA as reference tests for the concerned pathogens:

- VHSV and IHNV (EURL for fish and crustacean diseases, 2021)
 - cell culture plus ELISA with Monoclonal antibodies or
 - qPCR assays based on Jonstrup et al. (2012) for VHSV; Purcell et al. (2013) for IHNV
- EHNIV (EURL for fish and crustacean diseases, 2021)
 - Cell culture plus IFAT, ELISA or PCR, Real time PCR. Sequencing is required
- KHV (WOAHA, 2023)
 - qPCR (Gilad, 2004; Engelsma et al., 2013)
- ISAV (EURL for fish and crustacean diseases, 2021)
 - HPR del- cell + IFAT; qPCR (Snow et al., 2004), plus sequencing

VECTOR: when at least one positive reference test for 'Species X' and 'Species Y' was reported.

RESERVOIR: where there is at least one positive reference test for 'detection of Pathogen A in Species X' (and negative or not tested for Species 'Y').

Field studies: as they are subject to more uncertainty, ideally two reference tests taken from the same animals or reported in the same paper are needed to conclude Pathogen A was truly present in an animal from species X:

RESERVOIR: when positive for two reference tests for 'Species X'.

- **Negative results (0–10% certainty) [this is equivalent to 90–100% certainty that species X is not a vector or reservoir].**

The review should have captured only papers where Pathogen A had been detected in Species X. However, there are some papers where negative results were recorded for Pathogen A detection in species X, e.g. when several diagnostic tests were used to detect pathogen A in species X, and not all results were positive. Depending on the specific situation (e.g. if other studies are available or not), negative results in Species X can provide more than 90% certainty that Species X is NOT a RESERVOIR species based on the evidence extracted from the literature.

In addition, in transmission experiments, where negative results for Pathogen A detection in Species Y were recorded (susceptible species), the assessment focussed on Species X as reservoir species and the same method as described above was followed.

Any positive test result that was not one of the above situations were considered as inconclusive. The inconclusive results were elaborated in the next step of the assessment (group discussion).

2.1.1.2. Second step: group discussion

- **Smaller expert working group discussion**
 - The individual judgements were presented and discussed to reach a consensus judgement between 3–4 experts on fish diseases.
 - Only inconclusive cases were discussed, and experts were asked to identify a more precise certainty range for the inconclusive assessments:
 - Likely 66–90%
 - As likely as not 33–66%
 - Unlikely 10–33%
- **Whole working group**
 - The results of the smaller expert group were presented, discussed and consolidated by the whole working group.
 - The cut-off level for classifying species as vectors or reservoirs was set at a minimum certainty of 66%.

Since some fish species could be the subject of different studies with different study design and methodological quality, their assessment could result in different classifications. In these situations, the classification as vector prevailed above the classification as reservoir species, as evidence of transmission was present. Nonetheless, all the outcomes of all the assessments of different studies with a certainty > 66% are provided in the assessment section (Tables 1–3), but only the classification with the highest risk for transmission was taken up in the conclusions. Studies of species for which the assessments had a certainty < 66% are provided in Appendix B, Table B.1.

Finally, it should be noted that one the limitation of the assessment based ELR is that it was exclusively based on peer-reviewed evidence. Current lack of qualitative evidence or published studies on specific species does not mean the species cannot play a role as vector or reservoir. Therefore, the assessment should be updated when new evidence becomes available.

2.1.2. Term of Reference 2: Conditions under which fish species shall be regarded as vectors or reservoirs of diseases of fish listed in Annex II to the AHL

Several conditions need to be fulfilled for a fish species to be able act as a vector or reservoir of a pathogenic agent for the purposes of movements.

The conditions laid down in EC Delegated Reg 2020/990 Annex I and EC Delegated Reg 2020/692, focus on **the exposure to a pathogenic agent**. The vectors or reservoirs should have been

exposed to the pathogenic agent at source. There are other conditions, however, that will influence if a potential vector species transmits the pathogenic agent to a susceptible species at the destination:

- **Contact with susceptible species:** The vectors or reservoirs should be in contact at the place of destination with uninfected susceptible/listed species.
- **Survival of the pathogen in or on the vector or reservoir:** the tenacity of the specific pathogenic agent will play a role in the probability of survival of the pathogen until the exposure and possible infection of a susceptible species.
- **Environmental conditions:** there are many different environmental conditions which could impact the persistence of a pathogen outside the vector or reservoir or within the vector or reservoir, at the source, during transport or at the destination. These conditions include temperature, pH, salinity, pollutants, turbidity, UV radiation and microbial water quality. However, it is presumed the water quality would not change significantly during the journey when vectors are moved to their destination.
- **Duration of the journey:** the shorter the duration of the journey between place of origin and destination, the more viable pathogenic organisms can be found, as decay for all pathogens is a function of time (Oidtmann et al., 2017).
- **Experimental infections:** temperature in combination with time are the most common factors, which affect persistence of aquatic animal pathogens. The conditions during experimental infection should be considered (such as the use of sterilised water, mud or suspended solids) and the effect of UV light and temperature as they can impact the time during which the pathogen can persist.
- **Testing at the origin:** Test sensitivity, test specificity and sampling protocol to determine the pathogen-free status of the consignment should be considered. Fallow periods between restocking farms following confirmed outbreaks should be considered (WOAH, 2022).

To deliver a concise and timely Scientific Opinion, it was agreed not to provide an exhaustive description of all those possible conditions. On the contrary, it was decided to focus only on those conditions that would PREVENT transmission facilitated by the movement of vectors and reservoirs, for which scientific evidence is available. In a first step, the experts in the working group carried out a narrative literature review to collect any evidence from scientific literature identifying conditions that may prevent transmission by vectors. In addition, information on the duration of the experimental studies and the water temperature were compiled during the ELS, carried out for ToR 1, collecting the ranges of the different durations and temperatures for which transmission has been proven for the different pathogens by the different vector species. Then, the experts concluded by consensus if the collected evidence was sufficient to support the need to alter the conditions stipulated in Annex I to Commission Delegated Regulation (EU) 2020/990 and in Annex XXX to Commission Delegated Regulation (EU) 2020/692.

2.2. Data

2.2.1. Term of Reference 1: Assessment of potential vectors and reservoir species of diseases of fish listed in Annex II to the AHL

The detailed data set extracted through the ELR is available in EFSA (2023) – see Annex.

2.2.2. Term of Reference 2: Conditions under which fish species shall be regarded as vectors or reservoirs of diseases of fish listed in Annex II to the AHL

The detailed data set extracted through the ELR is available in EFSA (2023) – see Annex.

3. Assessment

3.1. Term of Reference 1: Assessment of potential vectors and reservoir species of diseases of fish listed in Annex II to the AHL

Table 1 summarises the results of the assessment of potential vectors of AHL-listed pathogens of fish. The outcome of the assessment was that *Ctenopharyngodon idella* (grass carp), *Gymnocephalus cernua* (Eurasian ruffe), *Hypophthalmichthys molitrix* (silver carp), *Rutilus rutilus* (common roach) and

Tinca tinca (Tench) are considered to be vector species for **KHV** with more than 90% certainty, and *Carassius auratus* (goldfish) and *Carassius gibelio* (Prussian carp) with a certainty between 66 and 90%. It should be mentioned that all these species were also already suggested as vector species by the previous EURL report (EURL, 2022).

The assessment was based on the evidence from experimental infection studies that was generated by the ELR. The reasoning and level of certainty of the proposed classification is provided. More detailed data that were extracted from eligible studies, but that were considered as insufficient evidence, can be found in EFSA (2023).

Table 1: Proposed **vectors** of the AHL-listed fish pathogen Koi herpes virus based on evidence from experimental infection studies, with certainty and reasoning of classification

Vector Species X	Presence in EU*	Transmission route investigated	Species Y	Pathogen detection method with positive results in species Y	Certainty of classification as vector species	Reasoning for classification as vector species	Reference	Suggested classification by previous EURL report (2022)
AHL-listed fish pathogen: KHV								
Carassius auratus (goldfish)	Yes	Cohabitation	Cyprinus carpio (common carp)	PCR	66–90%	DNA demonstrated only in one individual of species Y	Radosavljevic et al. (2012)	Vector/reservoir
				Nested PCR	66–90%	DNA demonstrated in Species Y	Bergmann et al. (2010)	
				ISH				
				IFAT	66–90%	Two independent (RT)-PCRs detected both virus DNA and also RNA, indicating expression of genes and replication. No virus isolation.	El-Matbouli and Soliman (2011)	
				RT-PCR				
Seq								
Carassius gibelio (Carassius auratus gibelio in paper) (Prussian carp)	Yes	Cohabitation	Cyprinus carpio (common carp)	PCR	66–90%	DNA demonstrated only in one individual of species Y.	Radosavljevic et al. (2012)	Vector/reservoir
Ctenopharyngodon idella (grass carp)	Yes	Cohabitation	Cyprinus carpio (common carp)	PCR and nested PCR	90–100%	PCR product detected in species Y	Kempton et al. (2012)	Vector/reservoir
				PCR	66–90%	DNA demonstrated only in one individual of species Y	Radosavljevic et al. (2012)	
Gymnocephalus cernua (Eurasian ruffe)	Yes	Cohabitation	Cyprinus carpio (common carp)	PCR and nested PCR	90–100%	PCR product detected in species Y	Kempton et al. (2012)	Vector/reservoir
Hypophthalmichthys molitrix (silver carp)	Yes	Cohabitation	Cyprinus carpio (common carp)	PCR and nested PCR	90–100%	PCR product detected in species Y	Kempton et al. (2012)	Vector/reservoir
Rutilus rutilus (common roach)	Yes	Cohabitation	Cyprinus carpio (common carp)	PCR and nested PCR	90–100%	PCR product detected in species Y	Kempton et al. (2012)	Vector/reservoir

Vector Species X	Presence in EU*	Transmission route investigated	Species Y	Pathogen detection method with positive results in species Y	Certainty of classification as vector species	Reasoning for classification as vector species	Reference	Suggested classification by previous EURL report (2022)
<i>Tinca tinca</i> (tench)	Yes	Cohabitation	<i>Cyprinus carpio</i> (common carp)	PCR and nested PCR	90–100%	PCR product detected in species Y	Kempton et al. (2012)	Vector/reservoir
				PCR	66–90%	DNA demonstrated only in one individual of species Y	Radosavljevic et al. (2012)	

*Source: [Gbif.org](https://gbif.org); PCR: polymerase chain reaction; Seq: sequencing; RT-PCR: reverse transcription polymerase chain reaction; IFAT: immunofluorescence antibody test; ISH: in situ hybridisation; EURL: EU Reference Laboratory.

Table 2 summarises the results of the assessment of potential reservoir species of AHL-listed pathogens of fish, based on the evidence from experimental infection studies, generated by the ELR. However, in these studies, no evidence of transmission to species Y was found or studied. The level of certainty and reasoning of the proposed classification is provided. More detailed data from studies that did not provide sufficient evidence are in EFSA (2023).

The outcome of the assessment was that *Tandanus tandanus* (dewfish) is considered a reservoir species for **EHNV** with > 90% certainty and *Maccullochella peelii* (Murray cod), *Macquaria novemaculeata* (Australian bass) and *Macquaria ambigua* (callop) with 66–90% certainty.

Clupea harengus (Atlantic herring) and *Oncorhynchus masou* (masu salmon) are considered a reservoir species for **HPRΔ ISAV** with > 90% certainty; and *Gadus morhua* (atlantic cod); *Oncorhynchus keta* (chum salmon) and *Oncorhynchus kisutch* (coho salmon) with 66–90% certainty.

Acipenser transmontanus (white sturgeon), *Clupea pallasii pallasii* (Pacific herring), *Cyprinus carpio* (common carp), *Danio rerio* (zebrafish), *Perca flavescens* (American yellow perch) and *Seriola quinqueradiata* (Japanese amberjack) are considered a reservoir species for **IHN** with > 90% certainty.

Acipenser ruthenus × *Huso huso* (hybrid sterlet × beluga), *Barbatula barbatula* (stone loach) and *Pseudorasbora parva* (topmouth gudgeon) are considered a reservoir species for **KHV** with > 90% certainty; and *Gasterosteus aculeatus* (three-spine stickleback), *Oncorhynchus mykiss* (rainbow trout), *Perca fluviatilis* (European perch), *Rutilus rutilus* (roach), *Scardinius erythrophthalmus* (pearl roach) and *Tinca tinca* (tench) with 66–90% certainty.

Acanthopagrus schlegelii (black porgy), *Cottus pollux* (Japanese fluvial sculpin), *Cyprinus carpio* (common carp), *Epinephelus akaara* (Hong Kong grouper), *Esox lucius* × *Esox masquinongy* (tiger muskellunge), *Hippoglossus hippoglossus* (Atlantic halibut), *Ictalurus punctatus* (channel catfish), *Oryzias latipes* (Japanese rice fish), *Pagrus major* (red sea bream), *Petromyzon marinus* (sea lamprey), *Rhinogobius* sp. (Yoshinobori complex; Japanese goby), *Salvelinus fontinalis* (brook trout), *Sebastes inermis* (black rockfish) and *Seriola quinqueradiata* (Japanese amberjack) are considered reservoir species for **VHSV** with more than 90% certainty. *Carassius auratus* (goldfish), *Notemigonus crysoleucas* (golden shiners), *Oncorhynchus mykiss* (rainbow trout) × *Salvelinus alpinus* (Alpine charr), *Oncorhynchus mykiss* (rainbow trout) × *Salvelinus namaycush* (American Lake Charr) and *Oncorhynchus mykiss* (rainbow trout) × *Salmo trutta* (Amu-Darya trout) are considered a reservoir species for VHSV with 66–90% certainty.

The species that were already indicated by the previous report from EURL (EURL, 2022) as vector species are shown in the Table 2. As agreed during the interpretation of the terms of reference, in this review only pathogen detection in species Y was considered as evidence of transmission, whereas for the assessment for the previous EURL report may have included a broader range of evidence for transmission to species Y, including the demonstration of clinical signs.

Table 2: Proposed **reservoirs** of AHL-listed fish pathogens based on evidence from experimental infection studies, with certainty and reasoning of classification

Reservoir species X	Presence in EU*	Pathogen detection method in species X with positive results	Certainty of classification as reservoir species	Reasoning for classification as reservoir species	Reference	Suggested classification by previous EURL report (2022)
AHL-listed fish pathogen: EHN						
<i>Maccullochella peelii</i> (Murray cod)	No	CellCult	66–90%	Virus isolation after bath exposure	Langdon (1989)	Vector/reservoir
<i>Macquaria ambigua</i> (callop)	No	CellCult	66–90%	Virus isolation and mortality after experimental infection by IP	Langdon (1989)	Vector/reservoir
<i>Macquaria novemaculeata</i> (Australian bass)	No	CellCult	66–90%	Virus isolation and mortality after experimental infection by IP	Langdon (1989)	Not assessed
<i>Tandanus tandanus</i> (dewfish)	No	PCR CellCult	90–100%	Five of 15 positive for EHN (virus isolation) in IP injection challenge	Becker et al. (2013)	Vector/reservoir
AHL-listed fish pathogen: HPRΔ ISAV						
<i>Clupea harengus</i> (Atlantic herring)	Yes	RT-PCR	90–100%	One reference test was performed. No virus isolation.	Nylund et al. (2002)	Vector/reservoir
<i>Gadus morhua</i> (Atlantic cod)	Yes	PCR Sec	66–90%	Detected in the brain of cod by RT-PCR and qRT-PCR after 45 days IP	Grove et al. (2007)	Not assessed
<i>Oncorhynchus keta</i> (chum salmon)	Yes	CellCult	66–90%	Virus isolated from some individuals with low titres	Rolland and Winton (2003)	Not assessed
<i>Oncorhynchus kisutch</i> (coho salmon)	Yes	CellCult	66–90%	Virus isolated from some individuals with low titres	Rolland and Winton (2003)	Vector/reservoir
<i>Oncorhynchus masou</i> (masu salmon)	Yes	RT-PCR	90–100%	Virus detected by PCR in Masu salmon but no ISAV detected in Atlantic salmon after cohabitation with Masu salmon	Ito et al. (2014)	Vector/reservoir
AHL-listed fish pathogen: IHN						
<i>Acipenser transmontanus</i> (white sturgeon)	Yes	Plaque assay	66–90%	Experimental infection using immersion prove to cause mortality in larval stages. Virus is re-isolated from larval stages.	LaPatra et al. (1995)	Vector/reservoir
<i>Clupea pallasii pallasii</i> (Pacific herring)	No	Plaque assay	90–100%	IHN recovered from larvae tissue by RT-PCR after bath infection. No mortality or virus shedding.	Hart et al. (2011)	Vector/reservoir

Reservoir species X	Presence in EU*	Pathogen detection method in species X with positive results	Certainty of classification as reservoir species	Reasoning for classification as reservoir species	Reference	Suggested classification by previous EURL report (2022)
<i>Cyprinus carpio</i> (common carp)	Yes	PCR Plaque assay	90–100%	Cell culture and PCR positive but no active infection after immersion	Palmer and Emmenegger (2014)	Vector/reservoir
<i>Danio rerio</i> (zebrafish)	Yes	RT-qPCR	90–100%	One reference test was performed. Increased virus load	Briolat et al. (2014)	Not assessed
		CellCult	90–100%	Zebrafish experimentally infected- virus detected by isolation in cell culture, no shedding in water or mortality	LaPatra et al. (2000)	
<i>Perca flavescens</i> (American yellow perch)	Yes	PCR Plaque assay	90–100%	Cell culture and PCR positive and some clinic after immersion	Palmer and Emmenegger (2014)	Vector/reservoir
<i>Seriola quinqueradiata</i> (Japanese amberjack)	No	PCR CellCul	90–100%	Virus replication after IP injection only in amberjack	Ito and Kamaishi (2021)	Not assessed
AHL-listed fish pathogen: KHV						
<i>Acipenser ruthenus</i> × <i>Huso huso</i> (hybrid sterlet × beluga)	Yes	Nested-PCR	90–100%	Detected by PCR	Pospichal et al. (2016)	Vector/reservoir
<i>Barbatula barbatula</i> (stone loach)	No	Nested-PCR	90–100%	Detected by PCR	Pospichal et al. (2016)	Vector/reservoir
<i>Gasterosteus aculeatus</i> (three-spine stickleback)	Yes	PCR	66–90%	KHV detected by standard PCR method (relatively high number of copies)	Fabian et al. (2013)	Not assessed
<i>Oncorhynchus mykiss</i> (rainbow trout)	Yes	PCR	66–90%	Infection by cohabitation and immersion, PCR positive	Bergmann and Cieslak (2016)	Vector/reservoir
<i>Perca fluviatilis</i> (European-perch)	Yes	PCR	66–90%	KHV detected by standard PCR method (relatively high number of copies)	Fabian et al. (2013)	Vector/reservoir
<i>Pseudorasbora parva</i> (topmouth gudgeon)	Yes	qPCR	90–100%	One reference test was performed. Positive results in fish exposed to removal of skin mucus and in fish stressed by scaring by net.	Pospichal et al. (2018)	Not assessed
<i>Rutilus rutilus</i> (roach)	Yes	PCR	66–90%	KHV detected by standard PCR method (relatively high number of copies)	Fabian et al. (2013)	Vector/reservoir
<i>Scardinius erythrophthalmus</i> (pearl roach)	Yes	PCR	66–90%	KHV detected by standard PCR method (relatively high number of copies)	Fabian et al. (2013)	Not assessed

Reservoir species X	Presence in EU*	Pathogen detection method in species X with positive results	Certainty of classification as reservoir species	Reasoning for classification as reservoir species	Reference	Suggested classification by previous EURL report (2022)
<i>Tinca tinca</i> (tench)	Yes	PCR	66–90%	KHV detected by standard PCR method (relatively high number of copies)	Fabian et al. (2013)	Vector/reservoir
AHL-listed fish pathogen: VHSV						
<i>Acanthopagrus schlegeli</i> (black porgy)	No	IFAT	90–100%	Virus detected by isolation from dead fish	Isshiki et al. (2003)	Vector/reservoir
<i>Carassius auratus</i> (goldfish)	Yes	CellCult	66–90%	VHSV re-isolated after IP.	Getchell et al. (2015)	Vector/reservoir
		PCR				
		HIS				
		PCR	90–100%	PCR positive after co habitation	El-Matbouli et al. (2007)	
		Seq				
LAMP assay						
<i>Cottus pollux</i> (Japanese fluvial sculpin)	No	RT-PCR	90–100%	Bath challenge produced mortality of 80%, VHSV detected by PCR and virus isolation from dead fish	Ito and Olesen (2013)	Vector/reservoir
		CellCult				
<i>Cyprinus carpio</i> (common carp)	Yes	RT-qPCRR	90–100%	Virus detected virus isolation in cell culture from target organs up to 28 dpe and by PCR up to 90 dpe.	Cornwell et al. (2013b)	Vector/reservoir
		CellCult				
		RT-qPCRR	90–100%	Detected by PCR	Goodwin et al. (2012)	
		Plaque assay	90–100%	Reasoning?	Emmenegger et al. (2013)	
<i>Epinephelus akaara</i> (Hong Kong grouper)	No	IFAT	90–100%	Virus detected by isolation from both dead and surviving fish	Isshiki et al. (2003)	Vector/reservoir
<i>Esox lucius</i> × <i>Esox masquinongy</i> (tiger muskellunge)	No	RT-qPCRR	90–100%	Isolated virus, cross lesions and characteristic histology detected. No transmission	Groocock et al. (2012)	Vector/reservoir
		CellCult				
		ImmHis				
		RT-qPCRR	90–100%	Oral infection 6 of 16 positive by both PCR and isolation in cell culture	Getchell et al. (2013)	
		CellCult				
His						

Reservoir species X	Presence in EU*	Pathogen detection method in species X with positive results	Certainty of classification as reservoir species	Reasoning for classification as reservoir species	Reference	Suggested classification by previous EURL report (2022)
<i>Hippoglossus hippoglossus</i> (Atlantic halibut)	Yes	CellCult	90–100%	Virus re-isolated on cell culture from moribund fish and identified by ELISA.	Bowden (2003)	Vector/reservoir
		ELISA				
		ELISA	66–90%	Virus isolated only from 1 individual	Snow et al. (2005)	Not assessed
<i>Ictalurus punctatus</i> (channel catfish)	Yes	RT-qPCRR	90–100%	Detected by PCR and virus Isolated and characteristic histology could be detected in 1 fish, but 28 dpe no virus was detected. No shedding of virus of transmission to other fish shown.	Groocock et al. (2012)	Vector/reservoir
		CellCult				
		RT-qPCRR	90–100%	Detected by PCR in 20 of 20 at 15°C; in 18 of 20 at 20°C; in 20 of 20 at 25°C	Goodwin et al. (2012)	
<i>Notemigonus crysoleucas</i> (golden shiners)	No	CellCult	90–100%	Detected by RT-qPCR, no sequencing	Cornwell et al. (2013a,b)	Vector/reservoir
		RT-qPCR				
<i>Oncorhynchus mykiss</i> (rainbow trout) × <i>Salvelinus alpinus</i> (Alpine Charr)	Yes	CellCult	66–90%	Text mentions virus isolation after experimental infection by immersion, but the data are not displayed	Dorson et al. (1991)	Vector/reservoir
<i>Oncorhynchus mykiss</i> (rainbow trout) × <i>Salvelinus namaycush</i> (American Lake Charr)	Yes	CellCult	66–90%	Text mentions virus isolation after experimental infection by immersion, but the data are not displayed	Dorson et al. (1991)	Vector/reservoir
<i>Oncorhynchus mykiss</i> (rainbow trout) × <i>Salmo trutta</i> (Amu-Darya Trout)	Yes	CellCult	66–90%	Focus is on mortality and resistance – but scarce details regarding virus detection. No virus isolated	Dorson et al. (1991)	Vector/reservoir
<i>Oryzias latipes</i> (Japanese rice fish)	No	RT-PCR	90–100%	Bath challenge produced mortality of 100%, VHSV detected by PCR and virus isolation from dead fish.	Ito and Olesen (2013)	Vector/reservoir
		CellCult				
<i>Pagrus major</i> (red sea bream)	Yes	CellCult	90–100%	High mortality and virus isolation after IP injection	Ito et al. (2004)	Vector/reservoir
<i>Petromyzon marinus</i> (sea lamprey)	Yes	CellCult	90–100%	Virus detected by both PCR, cell culture and immunohistochemistry in target organs	Coffee et al. (2017)	Susceptible
		RT-PCR				
		His				
		ImmHis				

Reservoir species X	Presence in EU*	Pathogen detection method in species X with positive results	Certainty of classification as reservoir species	Reasoning for classification as reservoir species	Reference	Suggested classification by previous EURL report (2022)
<i>Rhinogobius</i> sp. (Yoshinobori complex; Japanese goby)	No	PCR CellCult	90–100%	Bath challenge produced mortality of 10%, VHSV detected by PCR and virus isolation from dead fish	Ito and Olesen (2013)	Vector/reservoir
<i>Sebastes inermis</i> (black rockfish)	No	CellCult	90–100%	High mortality and virus isolation after IP injection	Ito et al. (2004)	Vector/reservoir
<i>Seriola quinqueradiata</i> (Japanese amberjack)	No	CellCult	90–100%	High mortality and virus isolation after IP injection	Ito et al. (2004)	Vector/reservoir

*Source: [Gbif.org](https://gbif.org); PCR: polymerase chain reaction; His: histology; Seq: sequencing; RT-PCR: reverse transcription polymerase chain reaction; RT-qPCR: quantitative real-time reverse transcription PCR; IFAT: immunofluorescence antibody test; IP: intraperitoneal injection; ISH: in situ hybridisation; CellCult: cell culture; ImmHis: immune-histology; EURL: EU Reference Laboratory; dpe; days post exposure.

Table 3 summarises the results of the assessment of potential reservoir species of AHL-listed pathogens of fish, based on the evidence from field studies that was generated by the ELR. As from these studies potential transmission could not be evaluated, their role as potential reservoir species was assessed.

The outcome of the assessment was that *Oncorhynchus kisutch* (Coho salmon) is considered a reservoir species for **HPRΔ ISAV** with more than 90% certainty.

Aulorhynchus flavidus (tube-snout) is considered a reservoir species for **IHN** with more than 90% certainty. *Clupea pallasii pallasii* (Pacific herring), *Cymatogaster aggregata* (shiner perch) and *Oncorhynchus gorbuscha* (pink salmon) are considered a reservoir species for **IHN** with 66–90%.

Acipenser oxyrinchus (Russian sturgeon), *Acipenser gueldenstaedtii* (Atlantic sturgeon) and *Oreochromis niloticus* (Nile tilapia) are considered a reservoir species for **KHV** with more than 90% certainty.

The following species are considered reservoir species for **VHSV** with > 90% certainty: *Anguilla anguilla* (European eel), *Argentina sphyraena* (lesser argentine), *Belone belone* (garfish), *Cyprinus carpio* (common carp), *Enchelyopus cimbrius* (FOURBEARD rockling), *Eutrigla gurnardus* (grey gurnard), *Gadiculus argenteus* (Silvery pout), *Gadus chalcogrammus* (Alaska pollock), *Ictalurus punctatus* (channel catfish), *Merluccius productus* (Pacific hake), *Moxostoma anisurum* (silver redhorse), *Moxostoma macrolepidotum* (shorthead redhorse sucker), *Percopsis omiscomaycus* (trout perch), *Petromyzon marinus* (sea lamprey), *Pomoxis annularis* (white crappie), *Reinhardtius hippoglossoides* (Greenland halibut), *Salvelinus alpinus* (Arctic charr) and *Scorpaena porcus* (black scorpionfish).

Alosa pseudoharengus (alewife), *Anoplopoma fimbria* (sablefish), *Catostomus commersonii* (white sucker), *Fundulus diaphanous* (Banded killifish), *Hypomesus pretiosus* (surf smelt) and *Lota lota* (burbot) and *Trisopterus minutus* (poor cod) are considered reservoir species for **VHSV** with 66–90% certainty.

More details of field studies that provided insufficient evidence are included in EFSA (2023).

Finally, Table A.1 in Appendix A lists all the currently listed reservoir and/or vector species in Commission Implementing Regulation 1882/2018 for which no eligible papers were found during the ELR. It is therefore suggested to remove these species from the list.

Table 3: Proposed **Reservoir** of AHL-listed fish pathogens based on evidence from field studies, retrieved through extensive literature review, with certainty and reasoning of classification

Reservoir species X	Presence in the EU*	Pathogen detection method in species X with positive results	Certainty of classification as reservoir species	Reasoning for classification as reservoir species	Reference	Suggested classification by previous EURL report (2022)
AHL-listed fish pathogen: HPRΔ ISAV						
<i>Oncorhynchus kisutch</i> (coho salmon)	Yes	RT-PCR	90–100%	Detected both by PCR and ELISA	Kibenge et al. (2002)	Vector/reservoir
AHL-listed fish pathogen: IHN						
<i>Aulorhynchus flavidus</i> (tube-snout)	No	CellCult DNA probe	90–100%	Two out of 72 fish tested positive by DNA probe and neutralising antibody test	Kent et al. (1998)	Vector/reservoir
<i>Clupea pallasii pallasii</i> (Pacific herring)	No	Neutralising antibody test CellCult DNA probe	66–90%	One out of 162 fish tested positive by DNA probe and neutralising antibody test	Kent et al. (1998)	Vector/reservoir
<i>Cymatogaster aggregata</i> (shiner perch)	No	CellCult DNA probe	66–90%	One out of 307 fish tested positive by DNA probe and neutralising antibody test	Kent et al. (1998)	Vector/reservoir
<i>Oncorhynchus gorbuscha</i> (Pink salmon)	Yes	CellCult RT-PCR	66–90%	1 of 72 fish positive by PCR	Breyta et al. (2017)	Vector/reservoir
AHL-listed fish pathogen: KHV						
<i>Acipenser gueldenstaedtii</i> (Russian sturgeon)	Yes	PCR and nested PCR Seq IFAT InSiHy	90–100%	Positive on more than two reference tests	Kempter et al. (2009)	Vector/reservoir
<i>Acipenser oxyrinchus</i> (Atlantic sturgeon)	Yes	PCR and nested PCR Seq IFAT InSiHy	90–100%	Positive on more than two reference tests	Kempter et al. (2009)	Vector/reservoir
<i>Oreochromis niloticus</i> (Nile tilapia)	Yes	PCR His TEM	90–100%	Internal organs test positive with reference method (PCR) and TEM	Wahidi et al. (2019)	Not assessed

Reservoir species X	Presence in the EU*	Pathogen detection method in species X with positive results	Certainty of classification as reservoir species	Reasoning for classification as reservoir species	Reference	Suggested classification by previous EURL report (2022)
AHL-listed fish pathogen: VHSV						
<i>Alosa pseudoharengus</i> (alewife)	No	PCR Seq	66–90%	Detected with both RT-PCR and sequencing	Stepien and Niner (2020)	Vector/reservoir
<i>Anguilla anguilla</i> (European eel)	Yes	CellCult ImmHis Seroneutralization	90–100%	Pathogenicity assay in RT-PCR plus isolation on cell and identification with Mabs	Castric et al. (1992)	Vector/reservoir
<i>Anoplopoma fimbria</i> (sablefish)	No	PCR	66–90%	Detected by PCR.	Hedrick et al. (2003)	Vector/reservoir
<i>Argentina sphyraena</i> (lesser argentine)	Yes	CellCult ELISA	90–100%	Detected by isolation in cell culture and ELISA in 1–3 fish	Mortensen et al. (1999)	Vector/reservoir
<i>Belone belone</i> (garfish)	Yes	RT-mPCR CellCult Seq	90–100%	Positive by cell culture and detection by RT-PCR and sequencing	Ogut and Altuntas (2014)	Vector/reservoir
<i>Catostomus commersonii</i> (white sucker)	Yes	PCR CellCult	66–90%	Detected in only 2 of 101 fish, high Ct-values, no cultivation	Cornwell et al. (2011)	Vector/reservoir
<i>Cyprinus carpio</i> (common carp)	Yes	RT-PCR CellCult Seq	90–100%	Positive by RT-PCR, cell culture and sequencing	Thompson et al. (2011)	Vector/reservoir
<i>Enchelyopus cimbrius</i> (fourbeard rockling)	Yes	CellCult ELISA	90–100%	Two reference tests were performed with positive results	Mortensen et al. (1999)	Vector/reservoir
<i>Eutrigla gurnardus</i> (grey gurnard)	Yes	CellCult ELISA PCR Seq	90–100%	Internal organs positive on cell culture in 1 of 3 pools	Wallace et al. (2016)	Vector/reservoir
<i>Fundulus diaphanous</i> (banded killifish)	No	CellCult RT-qPCR	66–90%	Positive by PCR and no virus isolation by cell culture	Bain et al. (2010)	Vector/reservoir

Reservoir species X	Presence in the EU*	Pathogen detection method in species X with positive results	Certainty of classification as reservoir species	Reasoning for classification as reservoir species	Reference	Suggested classification by previous EURL report (2022)
<i>Gadiculus argenteus</i> (silvery pout)	Yes	rRT-PCR Seq	90–100%	Only one reference test was performed, but only PCR positive in 1 pool. However, confirmation by sequence analysis.	Sandlund et al. (2014)	Vector/reservoir
<i>Gadus chalcogrammus</i> (Alaska pollock)	No	PCR CellCult TEM	90–100%	Positive by cell culture and RT-PCR	Meyers et al. (1999)	Vector/reservoir
<i>Hypomesus pretiosus</i> (surf smelt)	No	PCR CellCult Plaque assay Seq	66–90%	Detected in 60 of 60 dead fish by virus isolation and PCR/sequencing	Hedrick et al. (2003)	Vector/reservoir
<i>Ictalurus punctatus</i> (channel catfish)	Yes	RT-PCR CellCult Seq	90–100%	Positive by RT-PCR, cell culture and sequencing	Thompson et al. (2011)	Not assessed
<i>Lota lota</i> (burbot)	Yes	nRT-PCR CellCult	66–90%	Virus isolation positive, confirmed by nRT-PCR. 1 fish positive.	Al-Hussiney et al. (2011)	Vector/reservoir
<i>Merluccius productus</i> (Pacific hake)	No	CellCult PCR TEM Nested PCR Seq	90–100%	Seventeen of 19 fish positive by cell culture and RT-PCR	Meyers et al. (1999)	Vector/reservoir
<i>Moxostoma anisurum</i> (silver redhorse)	No	RT-PCR CellCult Seq	90–100%	Detected by both isolation in cell culture and PCR, also pathology	Faisal et al. (2012)	Vector/reservoir
<i>Moxostoma macrolepidotum</i> (shorthead redhorse sucker)	No	RT-PCR CellCult Seq	90–100%	Detected by both isolation in cell culture and PCR, also pathology	Faisal et al. (2012)	Vector/reservoir
		RT-PCR CellCult Seq	90–100%	Positive by RT-PCR, cell culture and sequencing	Thompson et al. (2011)	Vector/reservoir
		RT-PCR CellCult Seq				
		RT-PCR CellCult Seq				

Reservoir species X	Presence in the EU*	Pathogen detection method in species X with positive results	Certainty of classification as reservoir species	Reasoning for classification as reservoir species	Reference	Suggested classification by previous EURL report (2022)
<i>Percopsis omiscomaycus</i> (trout perch)	No	RT-PCR CellCult Seq	90–100%	Positive by RT-PCR, cell culture and sequencing	Thompson et al. (2011)	Vector/reservoir
<i>Petromyzon marinus</i> (sea lamprey)	Yes	RT-PCR CellCult Seq	90–100%	Positive by RT-PCR, cell culture and sequencing	Thompson et al. (2011)	Susceptible
<i>Pomoxis annularis</i> (white crappie)	No	nRT-PCR Seq CellCult ImmHis	90–100%	Positive by cell culture, IHC and RT-PCR, further confirmed by sequencing	Al-Hussinee et al. (2011)	Vector/reservoir
<i>Reinhardtius hippoglossoides</i> (Greenland halibut)	Yes	RT-PCR Seq CellCult Immunodot Seroneutralisation TEM	90–100%	Positive by cell culture, IHC and RT-PCR	Lopez-Vazquez et al. (2006) and Dopazo et al. (2002)	Vector/reservoir
<i>Salvelinus alpinus</i> (Arctic char)	Yes	CellCult IFAT	90–100%	Detected in 5 fish by isolation in cell culture	Knuesel et al. (2003)	Vector/reservoir
<i>Scorpaena porcus</i> (black scorpionfish)	No	RT-mPCR CellCult Seq	90–100%	Both CPE and detection by RT-PCR and sequencing	Ogut and Altuntas (2014)	Vector/reservoir
<i>Trisopterus minutus</i> (poor cod)	Yes	CellCult ELISA	66–90%	Only one pool WAS positive but a reference method was used	King et al. (2001)	Vector/reservoir

*Source: [Gbif.org](https://gbif.org); PCR: polymerase chain reaction; His: histology; TEM: transmission electron microscope; Seq: sequencing; RT-PCR: reverse transcription polymerase chain reaction; RT-qPCR: quantitative real-time reverse transcription PCR; IFAT: immunofluorescence antibody test; ISH: In situ hybridisation; CellCult: cell culture; ImmHis: immune-histology; EURL: EU Reference Laboratory; nRT-PCR: nested reverse transcription PCR; RT-mPCR: reverse transcription multiplex PCR.

3.2. Term of Reference 2: Conditions under which fish species shall be regarded as vectors or reservoirs of pathogens listed in Annex II to the AHL

For a species to act as a vector there should be prior exposure to the pathogen of interest at the place of origin. That is, contact with susceptible species, other vector species or reservoir species or a pathogen-contaminated environment in the period before movement should have happened.

The type of aquaculture establishment from where the vector species is moved will influence the probability of exposure to the pathogen at the place of origin, going from low risk in closed systems to increasing risk in semi-closed and open water aquaculture systems. Nonetheless, it should be mentioned that even in very high biosecurity, closed, aquaculture systems in affected areas, introductions of listed pathogens can occur and high biosecurity conditions of the establishment, when in an affected area, cannot provide 100% assurance of pathogen freedom before movement of the vector species.

Potential survival of the pathogen during the journey will mainly depend on the duration of the journey, the tenacity of the pathogen and temperatures and water quality during transport. The duration between exposure to the potential source of infection and then exposure to naïve stocks of farmed aquatic animals should take account of the incubation period, any latent period and pre-movement testing. The temperature and water quality can reduce the persistence of pathogen that may be present in the carrying water/matrix. However, the impact of these parameters is specific to each pathogen.

- Epizootic haematopoietic necrosis
EHNV is extremely resistant to drying and can survive for months in water. For these reasons, it is presumed that EHNV would persist for months to years in a fish farm in water and sediment as well as on equipment. EHNV will only be inactivated after 24 h at 40°C and within 15 min at 60°C (Langdon, 1989). There is a lack of information on possible effect of salinity; however, only freshwater species were shown to be susceptible. **Considering the available evidence, it is very likely to almost certain (90–100% certainty) that EHNV will remain infective at any possible transport condition.**
- Viral haemorrhagic septicaemia
VHSV is sensitive to enzymatic degradation, environments with high bacterial load and high temperatures (above 28°C). VHSV can survive outside the host tissue in fresh water and sea water, but is affected by temperature, ultraviolet (UV) exposure, microbial community and suspended sediments. VHSV is tolerant of high salt concentrations such as in brine-treated fish (Skall et al., 2015). The viral glycoprotein of VHSV is misfolded at temperatures above 22–24°C (Lorenzen, 1997, Lorenzen and Olesen, 1995) and aquatic animals kept at temperatures above this temperature should not pose a risk as vectors. VHSV is inactivated at 30°C within 24 h (Bovo et al., 2005). **Under transport conditions at temperatures below 25°C, it is likely (66–90% certainty) that VHSV will remain infective.**
- Infectious haematopoietic necrosis
IHNV stability in host tissues during storage and processing is largely influenced by temperature. IHNV can survive outside the host tissue in fresh water and sea water, but is affected by temperature, UV exposure, microbial community and suspended sediments. When exposed to sunlight (UV-A and UV-B), IHNV at the water surface is rapidly inactivated with six orders of magnitude of virus rendered non-infectious within 3 h (Garver et al., 2013). In addition, infectious virus is inactivated by the microbial community within the water source and with increased amounts of suspended sediments (Garver et al., 2013; Kamei et al., 1987). IHNV is inactivated within 90 min at 32°C. (Bovo et al., 2005), and as VHSV G-protein is misfolded at temperatures above 22–24°C (Cain et al., 1999). **Under transport conditions at temperatures below 25°C, it is likely (66–90% certainty) that IHNV will remain infective.**
- Koi herpes virus disease
KHV can remain infectious in water up to 21 h at water temperatures between 23 and 25°C (Perelberg et al., 2003). Other studies in Japan have displayed a significant reduction in the infectious titre of KHV within 3 days in environmental water or sediment samples at 15°C, while KHV remained for more than 7 days infective after being added to sterilised water samples

(Shimizu et al., 2006). In Japan, KHV DNA was detected in river water samples at temperatures of between 9 and 11°C, 4 months prior to a KHV disease outbreak being observed (Haramoto et al., 2007). Using PCR, KHV DNA was detected in high quantities in water samples collected at eight sites along the Yura river system during 3 months after a KHV disease outbreak at water temperatures ranging from 28.4 to 14.5°C (Minamoto et al., 2009a). In Lake Biwa, Japan, KHV was found to be widely distributed at different sites in the lake, 5 years after the first observed KHV outbreak. Mean concentrations of KHV in the lake water showed annual variation, with a peak in the summer and a decline in winter, and the virus was most prevalent in turbid, eutrophic water found in the lake margins (Minamoto et al., 2009b).

Considering the available evidence, it is very likely to almost certain (90–100%) that KHV will remain infective at any possible transport condition.

- Infection with HPR-deleted infectious salmon anaemia virus
 ISAV has been shown to be readily inactivated by UV exposure and ozone treatment (Liltved et al., 2006). It is also sensitive to low-pH values (e.g. pH 4, 30 min) and to temperatures above 37°C. Exposure to 56°C for 30 min will inactivate the virus (Falk et al., 1997). In natural seawater at 10°C, declining virus titres were shown during a period of 72 h (Vike et al., 2014). Results from an intraperitoneal challenge experiment indicated a loss of infectivity between 3 and 6 h (Vike et al., 2014). However, the infectivity will also be affected by the presence of particles and organic material in the water, which may potentially aid or reduce the stability of the virus.
Under transport conditions at temperatures below 25°C, it is likely (66–90% certainty) that HPR-deleted infectious salmon anaemia virus will remain infective.

Table 4 summarises the conditions under which fish species should be regarded as vectors or reservoirs and should be considered to amend Annex I of Reg 2020/990 and Annex XXX of Reg 2020/692.

Table 4: Proposed conditions under which fish species listed shall be regarded as vectors or reservoirs

Name of listed pathogen	Conditions to be considered to amend Annex I of Reg 2020/990 and Annex XXX of Reg 2020/692
EHN _V , KHV	Vector or reservoir species that were exposed to EHN _V in an affected area can possibly transmit EHN _V when transported into a non-affected area. Exposure in the affected area may have occurred if they originate from: <ul style="list-style-type: none"> (a) an aquaculture establishment or group of aquaculture establishments, where susceptible species, reservoir species or other vector species are kept; or (b) the wild, where they may have been exposed to susceptible, reservoir or other vector species; (c) an aquaculture supplied with water possibly contaminated with EHN_V.
VHS, IHN _V and HPRΔ ISAV	Vector or reservoir species that were exposed to VHSV in an affected area can possibly transmit VHSV when transported at a temperate below 25°C into a non-affected area. Exposure in the affected area may have occurred if the vector/reservoir originate from: <ul style="list-style-type: none"> (a) an aquaculture establishment or group of aquaculture establishments, where susceptible species, reservoir species or other vector species are kept; (b) the wild in an affected area, where they may have been exposed to susceptible species, reservoir species or other vector species; (c) an aquaculture supplied with water possibly contaminated with VHSV.

4. Conclusions

4.1. Term of Reference 1: Assessment of potential vectors and reservoir species of diseases of fish listed in Annex II to the AHL

EHN_V

Reservoir

- Tandanus tandanus* (dewfish) is considered a reservoir species for EHN_V with > 90% certainty and *Maccullochella peelii* (Murray cod), *Macquaria novemaculeata* (Australian bass) and *Macquaria ambigua* (callop) with 66–90% certainty.

- No evidence or insufficient evidence was generated by the ELR for the following species currently listed Vectors/reservoirs in Commission Implementing Regulation 1882/2018: *Aristichthys nobilis*, *Carassius auratus*, *Carassius carassius*, *Cyprinus carpio*, *Hypophthalmichthys molitrix*, *Leuciscus* spp., *Rutilus rutilus*, *Scardinius erythrophthalmus* and *Tinca tinca*.

KHV

Vector

- The following species are considered vector species for KHV with > 90% certainty: *Carassius auratus* (goldfish), *Carassius gibelio* (Prussian carp), *Ctenopharyngodon idella* (grass carp), *Gymnocephalus cernua* (Eurasian ruffe), *Hypophthalmichthys molitrix* (silver carp), *Rutilus rutilus* (common roach) and *Tinca tinca* (tench).

Reservoir

The following species are considered reservoir species for KHV with > 90% certainty: *Acipenser gueldenstaedtii* (Russian sturgeon), *Acipenser oxyrinchus* (Atlantic sturgeon), *Acipenser ruthenus* × *Huso huso* (hybrid sterlet × beluga), *Barbatula barbatula* (stone loach), *Gasterosteus aculeatus* (three-spine stickleback), *Perca fluviatilis* (European perch) and *Scardinius erythrophthalmus* (pearl roach).

- The following species are considered reservoir species for KHV with 66–90% certainty: *Oreochromis niloticus* (Nile tilapia) and *Pseudorasbora parva* (topmouth gudgeon).

HPRA ISAV

Reservoir

- *Clupea harengus* (Atlantic herring), *Oncorhynchus masou* (masu salmon) and *Oncorhynchus kisutch* (coho salmon) are considered reservoir species for HPRΔ ISAV with > 90% certainty.
- *Gadus morhua* (Atlantic cod), *Oncorhynchus keta* (chum salmon), *Oncorhynchus kisutch* (coho salmon) are considered reservoir species for HPRΔ ISAV with 66–90% certainty.

IHN

Reservoir

- The following species are considered reservoir species for IHNV with > 90% certainty: *Aulorhynchus flavidus* (tube-snout), *Acipenser transmontanus* (white sturgeon), *Clupea pallasii* (Pacific herring), *Cyprinus carpio* (common carp), *Danio rerio* (zebrafish), *Perca flavescens* (American yellow perch) and *Seriola quinqueradiata* (Japanese amberjack).
- *Cymatogaster aggregata* (shiner perch) and *Oncorhynchus gorbuscha* (pink salmon) are considered reservoir species for IHNV with 66–90% certainty.
- No evidence or insufficient evidence was generated by the ELR for the following species (including animals other than fish) currently listed Vectors/reservoirs in Commission Implementing Regulation 1882/2018: *Acipenser baerii*, *Acipenser gueldenstaedtii*, *Acipenser ruthenus*, *Acipenser stellatus*, *Acipenser sturio*, *Ameiurus melas*, *Aristichthys nobilis*, *Astacus astacus*, *Carassius auratus*, *Carassius carassius*, *Clarias gariepinus*, *Gadus morhua*, *Hippoglossus hippoglossus*, *Hypophthalmichthys molitrix*, *Huso huso*, *Ictalurus punctatus*, *Ictalurus* spp., *Leuciscus* spp., *Melanogrammus aeglefinus*, *Platichthys flesus*, *Pacifastacus leniusculus*, *Procambarus clarkii*, *Pangasius pangasius*, *Rutilus rutilus*, *Sander lucioperca*, *Scardinius erythrophthalmus*, *Silurus glanis*, *Tinca tinca*.

VHSV

Reservoir

- The following species are considered reservoir species for VHSV with > 90% certainty: *Anguilla anguilla* (European eel), *Argentina sphyraena* (lesser argentine), *Belone belone* (garfish), *Cottus pollux* (Japanese fluvial sculpin), *Cyprinus carpio* (common carp), *Enchelyopus cimbrius* (fourbeard rockling), *Epinephelus akaara* (Hong Kong grouper), *Esox lucius* × *Esox masquinongy* (tiger muskellunge), *Eutrigla gurnardus* (grey gurnard); *Gadiculus argenteus* (silvery pout), *Gadus chalcogrammus* (Alaska pollock), *Hippoglossus hippoglossus* (Atlantic

halibut), *Ictalurus punctatus* (channel catfish), *Merluccius productus* (Pacific hake), *Moxostoma anisurum* (silver redhorse), *Moxostoma macrolepidotum* (shorthead redhorse sucker), *Oncorhynchus mykiss* (rainbow trout) × *Salvelinus fontinalis* (American Brook Charr), *Oryzias latipes* (Japanese rice fish), *Pagrus major* (red sea bream), *Percopsis omiscomaycus* (trout perch), *Petromyzon marinus* (sea lamprey), *Pomoxis annularis* (white crappie), *Rhinogobius* sp. (Yoshinobori complex; Japanese goby), *Reinhardtius hippoglossoides* (Greenland halibut), *Salvelinus alpinus* (Arctic charr), *Scorpaena porcus* (black scorpionfish), *Sebastes inermis* (black rockfish) and *Seriola quinqueradiata* (Japanese amberjack).

- The following species are considered reservoir species for **VHSV** with 66–90% certainty: *Alosa pseudoharengus* (alewife), *Anoplopoma fimbria* (sablefish), *Carassius auratus* (goldfish), *Fundulus diaphanous* (banded killifish), *Hypomesus pretiosus* (surf smelt), *Lota lota* (burbot), *Notemigonus crysoleucas* (golden shiners), *Oncorhynchus mykiss* (rainbow trout) × *Salmo trutta* (Amu-Darya trout), *Oncorhynchus mykiss* (rainbow trout) × *Salvelinus alpinus* (Alpine Char), *Oncorhynchus mykiss* (rainbow trout) × *Salvelinus namaycush* (American Lake Charr).
- No evidence or insufficient evidence was generated by the ELR for the following species currently listed Vectors/reservoirs in Commission Implementing Regulation 1882/2018: *Acipenser baerii*; *Acipenser gueldenstaedtii*; *Acipenser ruthenus*; *Acipenser stellatus*; *Acipenser sturio*, *Ameiurus melas*, *Argyrosomus regius*, *Aristichthys nobilis*, *Carassius Carassius*, *Clarias gariepinus*, *Dentex dentex*, *Dicentrarchus labrax*, *Diplodus puntazzo*, *Diplodus sargus*, *Diplodus vulgaris*, *Epinephelus aeneus*, *Epinephelus marginatus*, *Huso huso*, *Hypophthalmichthys molitrix*, *Ictalurus* spp., *Leuciscus* spp., *Morone chrysops* x, *Morone saxatilis*, *Mugil cephalus*, *Oreochromis* sp., *Pagellus bogaraveo*, *Pagellus erythrinus*, *Pangasius pangasius*, *Pagrus pagrus*, *Rutilus rutilus*, *Salvelinus fontinalis*, *Sander lucioperca*, *Scardinius erythrophthalmus*, *Sciaenops ocellatus*, *Silurus glanis*, *Solea senegalensis*, *Solea solea*, *Sparus aurata*, *Thunnus* spp., *Thunnus thynnus*, *Tinca tinca*, *Umbrina cirrose*.
- The assessment was exclusively based on peer reviewed evidence and should be updated when new evidence becomes available.

4.2. Term of Reference 2: Conditions under which fish species shall be regarded as vectors or reservoirs of diseases of fish listed in Annex II to the AHL

- **VHSV, IHNV or HPR-deleted infectious salmon anaemia virus**
 - Under transport conditions at temperatures below 25°C, it is likely (66–90% certainty) that VHSV, IHNV and HPRΔ ISAV will remain infective.
 - Vector or reservoir species can transmit VHSV, IHNV or HPRΔ ISAV when transported at a **temperature below 25°C** into a non-affected area. Exposure in an VHSV, IHNV or HPRΔ ISAV affected area may have occurred if the vector or reservoir originate from:
 - a) an aquaculture establishment where susceptible species or reservoir or other vector species of are kept;
 - b) the wild, where they may have been exposed to susceptible, reservoir or other vector species;
 - c) an aquaculture establishment supplied with water possibly contaminated with VHSV, IHNV or HPRΔ ISAV.
- **EHN and KHV**
 - It is very likely to almost certain (90–100% certainty) that EHN and KHV will remain infective at **any possible transport condition**.
 - Vector or reservoir species can transmit EHN when transported into a non-affected area. Exposure in the affected area may have occurred if they originate from:
 - a) an aquaculture establishment, where susceptible species, reservoir species or other vector species are kept; or
 - b) the wild, where they may have been exposed to susceptible, reservoir or other vector species;
 - c) an aquaculture supplied with water possibly contaminated with EHN.

References

- Al-Hussiney L, Lord S, Stevenson RMW, Casey RN, Groocock GH, Britt KL, Kohler KH, Wooster GA, Getchell RG, Bowser PR and Lumsden JS, 2011. Immunohistochemistry and pathology of multiple Great Lakes fish from mortality events associated with viral hemorrhagic septicemia virus type IVb. *Diseases of Aquatic Organisms*, 93, 117–127.
- Bain MB, Cornwell ER, Hope KM, Eckerlin GE, Casey RN, Groocock GH, Getchell RG, Bowser PR, Winton JR, Batts WN, Cangelosi A and Casey JW, 2010. Distribution of an invasive aquatic pathogen (viral hemorrhagic septicemia virus) in the Great Lakes and its relationship to shipping. *PLoS One*, 5, e10156.
- Becker JA, Tweedie A, Gilligan D, Asmus M and Whittington RJ, 2013. Experimental infection of Australian freshwater fish with epizootic haematopoietic necrosis virus (EHNV). *Journal of Aquatic Animal Health*, 25, 66–76. <https://doi.org/10.1080/08997659.2012.747451>
- Bergmann SM and Cieslak M, 2016. Is There any species specificity in infections with aquatic animal herpesviruses?—the koi herpesvirus (KHV): an alloherpesvirus model. *Fisheries and Aquaculture Journal*, 07. <http://dx.doi.org/10.4172/2150-3508.1000169>
- Bergmann SM, Fichtner D, Skall HF, Schlotfeldt HJ and Olesen NJ, 2003. Age- and weight-dependent susceptibility of rainbow trout *Oncorhynchus mykiss* to isolates of infectious haematopoietic necrosis virus (IHNV) of varying virulence. *Diseases of Aquatic Organisms*, 55, 205–210.
- Bergmann SM, Schütze H, Fischer U, Fichtner D, Riechardt M, Meyer K, Schrudde D and Kempter J, 2009. Detection of koi herpes virus (KHV) genome in apparently healthy fish. *Bulletin of the European Association of Fish Pathologists*, 29, 145–152.
- Bergmann SM, Lutze P, Schütze H, Fischer U, Dauber M, Fichtner D and Kempter J, 2010. Goldfish (*Carassius auratus auratus*) is a susceptible species for koi herpesvirus (KHV) but not for KHV disease (KHVD). *Bulletin of the European Association of Fish Pathologists*, 30.
- Bergmann SM, Dabels J, Klafack S, Jin Y, Lee P, Hofmann AC, Wang Y, Wang Q, Li Y, Zeng W, Lusastuti A, Zheng S, Jin Y, Kielpinska J and Monaghan S, 2021. Serological responses to Koi herpesvirus (KHV) in a non-cyprinid reservoir host. *Journal of Fish Diseases*, 44, 1229–1236.
- Bovo G, Hill B, Husby A, Håstein T, Michel C, Olesen NJ, Storset A and Midtlyng PJ, 2005. Work package 3 report: Pathogen survival outside the host, and susceptibility to disinfection, Fish Egg Trade, ISBN: 82-91743-36-3. Available online: <file:///C:/Users/linne/Downloads/Fisheggtrade-wp3.pdf> and <file:///C:/Users/linne/Downloads/VHSV-and-IHNV-diagnostic-manual-v2021-2.pdf>.
- Bowden TJ, 2003. A study of the susceptibility of Atlantic halibut, *Hippoglossus hippoglossus* (L.), to viral haemorrhagic septicaemia virus isolated from turbot, *Scophthalmus maximus* (L.). *Journal of Fish Diseases*, 26, 207–212. <https://doi.org/10.1046/j.1365-2761.2003.00445.x>
- Breyta R, Brito I, Kurath G and Ladeau S, 2017. Infectious hematopoietic necrosis virus virological and genetic surveillance 2000–2012. *Ecology*, 98, 283.
- Briolat V, Jouneau L, Carvalho R, Palha N, Langevin C, Herbolme P, Schwartz O, Spaink HP, Levraud JP and Boudinot P, 2014. Contrasted innate responses to two viruses in zebrafish: Insights into the ancestral repertoire of vertebrate IFN-stimulated genes. *Journal of Immunology*, 192, 4328–4341.
- Cain KD, Byrne KM, Brassfield AL, LaPatra SE and Ristow SS, 1999. Temperature dependent characteristics of a recombinant infectious hematopoietic necrosis virus glycoprotein produced in insect cells DAO 36:1-10|Full text in pdf format.
- Castric J, Jeffroy J, Bearzotti M and Kinkelin P, 1992. Isolation of viral haemorrhagic septicaemia virus (VHSV) from wild elvers *Anguilla Anguilla*. *Bulletin of the European Association of Fish Pathologists*, 12, 21–23.
- Coffee LL, Getchell RG, Groocock GH, Cornwell ER, Wooster GA, Lumsden JS and Bowser PR, 2017. Pathogenesis of experimental viral hemorrhagic septicemia virus IVb infection in adult sea lamprey (*Petromyzon marinus*). *Journal of Great Lakes Research*, 43, 119–126.
- Cornwell ER, Eckerlin GE, Getchell RG, Groocock GH, Thompson TM, Batts WN, Casey RN, Kurath G, Winton JR, Bowser PR, Bain MB and Casey JW, 2011. Detection of viral hemorrhagic septicemia virus by quantitative reverse transcription polymerase chain reaction from two fish species at two sites in lake superior. *Journal of Aquatic Animal Health*, 23, 207–217. <https://doi.org/10.1080/08997659.2011.644411>
- Cornwell ER, Bellmund CA, Groocock GH, Wong PT, Hambury KL, Getchell RG and Bowser PR, 2013a. Fin and gill biopsies are effective nonlethal samples for detection of viral hemorrhagic septicemia virus genotype IVb. *Journal of Veterinary Diagnostic Investigation*, 25, 203–209.
- Cornwell ER, LaBuda SL, Groocock GH, Getchell RG and Bowser PR, 2013b. Experimental infection of koi carp with viral hemorrhagic septicemia virus type IVb. *Journal of Aquatic Animal Health*, 25, 36–41. <https://doi.org/10.1080/08997659.2012.732653>
- Cornwell ER, Anderson GB, Coleman D, Getchell RG, Groocock GH, Warg JV, Cruz AM, Casey JW, Bain MB and Bowser PR, 2015. Applying multi-scale occupancy models to infer host and site occupancy of an emerging viral fish pathogen in the Great Lakes. *Journal of Great Lakes Research*, 41, 520–529.

- Dopazo CP, Bandín I, López-Vazquez C, Lamas J, Noya M and Barja JL, 2002. Isolation of viral hemorrhagic septicemia virus from Greenland halibut *Reinhardtius hippoglossoides* caught at the Flemish Cap. *Diseases of Aquatic Organisms*, 50, 171–179. <https://doi.org/10.3354/dao050171>
- Dorson M, Chevassus B and Torhy C, 1991. Comparal susceptibility of three species of char and of rainbow trout × char triploid hybrids to several pathogenic salmonid viruses. *Diseases of Aquatic Organisms*, 11, 217–224.
- Dorson M, Torhy C and Kinkelin P, 1994. Viral haemorrhagic septicaemia virus multiplication and interferon production in rainbow trout and in rainbow trout × brook trout hybrids. *Fish & Shellfish Immunology*, 4, 369–381.
- EFSA (European Food Safety Authority), Algers B, Blokhuis HJ, Broom DM, Costa P, Domingo M, Greiner M, Guemene D, Hartung J, Koenen F, Müller-Graf C, Morton DB, Osterhaus A, Pfeiffer DU, Roberts R, Sanaa M, Salman M, Sharp JM, Vannier P, Wierup M, 2008. Scientific Opinion of the Panel on AHAW on a request from the European Commission on aquatic animal species. *EFSA Journal* 2008;6(11):808, 144 pp. <https://doi.org/10.2903/j.efsa.2008.808>
- EFSA (European Food Safety Authority), Gnocchi M, Aires M, Alvarez J, Arzul I, Asensio Aznar I, Bicot JD, Carmosino I, Ashley Drewe J, Dharmaveer S, Garin Bastuji B, Karagianni AE, Miranda Chueca MA, Olesen NJ, Palaikostas C, Roberts H, Saxmose Nielsen S, Schiøtt M, Sindre H, Stone D, Rusina A, Vendramin N and Dholland S, 2023. Extensive literature review on vectors and reservoirs of AHL-listed pathogens of fish. EFSA supporting publication 2023:EN-8123, 23 pp. <https://doi.org/10.2903/sp.efsa.2023.EN-8123>
- El-Matbouli M and Soliman H, 2011. Transmission of Cyprinid herpesvirus-3 (CyHV-3) from goldfish to naïve common carp by cohabitation. *Research in Veterinary Science*, 90, 536–539. <https://doi.org/10.1016/j.rvsc.2010.07.008>
- El-Matbouli M, Saleh M and Soliman H, 2007. Detection of cyprinid herpesvirus type 3 in goldfish cohabiting with CyHV-3-infected koi carp (*Cyprinus carpio koi*). *The Veterinary Record*, 161, 792–793.
- Emmenegger EJ, Moon C, Hershberger PK and Kurath G, 2013. Virulence of viral hemorrhagic septicemia virus (VHSV) genotypes Ia, IVa, IVb, and IVc in five fish species. *Diseases of Aquatic Organisms*, 107, 99–111.
- EURL for Fish and Crustacean Diseases, 2021. DIAGNOSTIC METHODS AND PROCEDURES FOR THE SURVEILLANCE AND CONFIRMATION OF INFECTION WITH VHSV AND IHN. Available online: <file:///C:/Users/dhollso/Downloads/VHSV-and-IHN-diagnostic-manual-v2021-2.pdf>.
- EURL, 2022. European Union Reference Laboratory for Fish and Crustacean Diseases. National Institute of Aquatic Resources. Technical University of Denmark. Scientific opinions and recommendations for listing fish species susceptible to infection with list A, C and E diseases according to EU/2018/1882. Ref. Ares(2022)4996074, 7 August 2022. pp 40.
- Fabian M, Baumer A and Steinhagen D, 2013. Do wild fish species contribute to the transmission of koi herpesvirus to carp in hatchery ponds? *Journal of Fish Diseases*, 36, 505–514.
- Faisal M, Shavali M, Kim RK, Millard EV, Gunn MR, Winters AD, Schulz CA, Eissa A, Thomas MV, Wolgamood M, Whelan GE and Winton J, 2012. Spread of the emerging viral hemorrhagic septicemia virus strain, genotype IVb, in Michigan, USA. *Viruses*, 4, 734–760.
- Falk K, Namork E, Rimstad E and Mjaaland S, 1997. Dannevig BH Characterization of infectious salmon anemia virus, an orthomyxo-like virus isolated from Atlantic salmon (*Salmo salar* L.). *Journal of Virology*, 71, 9016–9023. <https://doi.org/10.1128/JVI.71.12.9016-9023.1997>. Available online: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC230202/>
- Follett JE, Meyers TR, Burton TO and Geesin JL, 1997. Comparative susceptibilities of Salmonid Species in Alaska to Infectious Hematopoietic Necrosis Virus (IHN) and North American Viral Hemorrhagic Septicemia Virus (VHSV). *Journal of Aquatic Animal Health*, 9, 34–40.
- Gaede L, Steinbrück J, Bergmann SM, Jäger K, Grafe H, Schoon HA, Speck S and Truyen U, 2017. Koi herpesvirus infection in experimentally infected common carp *Cyprinus carpio* (Linnaeus, 1758) and three potential carrier fish species *Carassius carassius* (Linnaeus, 1758); *Rutilus rutilus* (Linnaeus, 1758); and *Tinca tinca* (Linnaeus, 1758) by quantitative real-time PCR and in-situ hybridization. *Journal of Applied Ichthyology*, 33, 776–784.
- Garver KA, Mahony AAM, Stucchi D, Richard J, Van Woensel C and Foreman MG, 2013. Estimation of parameters influencing waterborne transmission of infectious hematopoietic necrosis virus (IHN) in Atlantic Salmon (*Salmo salar*). *PLoS One*, 8, e82296. <https://doi.org/10.1371/journal.pone.0082296>
- Getchell RG, Cornwell ER, Grocock GH, Wong PT, Coffee LL, Wooster GA and Bowser PR, 2013. Experimental Transmission of VHSV Genotype IVb by Predation. *Journal of Aquatic Animal Health*, 25, 221–229. <https://doi.org/10.1080/08997659.2013.811126>
- Getchell RG, Erkinharju T, Johnson AO, Davis BW, Hatch EE, Cornwell ER and Bowser PR, 2015. Goldfish *Carassius auratus* susceptibility to viral hemorrhagic septicemia virus genotype IVb depends on exposure route. *Diseases of Aquatic Organisms*, 115, 25–36. <https://doi.org/10.3354/dao02872>
- Goodwin AE, Merry GE and Noyes AD, 2012. Persistence of viral RNA in fish infected with VHSV-IVb at 15°C and then moved to warmer temperatures after the onset of disease. *Journal of Fish Diseases*, 35, 523–528.
- Grocock GH, Frattini SA, Cornwell ER, Coffee LL, Wooster GA, Getchell RG and Bowser PR, 2012. Experimental infection of four aquacultured species with viral hemorrhagic septicemia virus type IVb. *Journal of the World Aquaculture Society*, 43, 459–476.

- Grove S, Hjortaas MJ, Reitan L and Dannevig BH, 2007. Infectious salmon anaemia virus (ISAV) in experimentally challenged Atlantic cod (*Gadus morhua*). *Archives of Virology*, 10, 1829–1837.
- Hart LM, Traxler GS, Garver KA, Richard J, Gregg JL, Grady CA, Kurath G and Hershberger PK, 2011. Larval and juvenile Pacific herring *Clupea pallasii* are not susceptible to infectious hematopoietic necrosis under laboratory conditions. *Diseases of Aquatic Organisms*, 93, 105–110. <https://doi.org/10.3354/dao02294>
- Haramoto E, Kitajima M, Katayama H and Ohgaki S, 2007. Detection of koi herpesvirus DNA in river water in Japan. *Journal of Fish Diseases*, 30, 59–61.
- Hedrick RP, Batts WN, Yun S, Traxler GS, Kaufman J and Winton JR, 2003. Host and geographic range extensions of the North American strain of viral hemorrhagic septicemia virus. *Diseases of Aquatic Organisms*, 55, 211–220. <https://doi.org/10.3354/dao055211>
- Isshiki T, Nagano T and Miyazaki T, 2003. Susceptibility of various marine fish species to viral hemorrhagic septicemia virus isolated from Japanese flounder. *Fish Pathology*, 38, 113–115.
- Ito T and Kamaishi T, 2021. Japanese amberjack *Seriola quinqueradiata* and Red Sea bream *Pagrus major* susceptibility to Infectious hematopoietic necrosis virus (IHN) isolate. *Diseases of Aquatic Organisms*, 146, 1–8.
- Ito T and Olesen NJ, 2007. Viral haemorrhagic septicaemia virus (VHSV) remains viable for several days but at low levels in the water flea *Moina macrocopa*. *Diseases of Aquatic Organisms*, 127, 11–18.
- Ito T and Olesen NJ, 2013. Susceptibility of various Japanese freshwater fish species to an isolate of viral haemorrhagic septicaemia virus (VHSV) genotype IVb. *Diseases of Aquatic Organisms*, 107, 1–8. <https://doi.org/10.3354/dao02667>
- Ito T, Mori K-i, Arimoto M and Nakajima K, 2004. Virulence of Viral Hemorrhagic Septicemia Virus (VHSV) Isolates from Japanese Flounder *Paralichthys olivaceus* in Rainbow Trout and Several Species of Marine Fish. *Fish Pathology*, 39, 103–104.
- Ito T, Oseko N and Otodate M, 2014. Susceptibility of Amago trout, *Oncorhynchus masou macrostomus* (Günther) to an isolate of infectious salmon anaemia virus. *Journal of Fish Diseases*, 38, 237–240. <https://doi.org/10.1111/jfd.12226>
- Jensen B, Reschova S, Cinkova K, Ariel E and Vesely T, 2011. Common carp (*Cyprinus carpio*) and goldfish (*Carassius auratus*) were not susceptible to challenge with ranavirus under certain challenge conditions. *Bulletin of the European Association of Fish Pathologists*, 31, 112–118.
- Jin Y, Adamkowska N, Kielpinska J and Bergmann SM, 2020. Detection of koi herpesvirus (KHV) and carp oedema virus (CEV) in invasive round goby, *Neogobius melanostomus* Pallas, 1814, from Poland and Germany. *Journal of Veterinary Research*, 64, 247–251.
- Jørgensen PEV, Castric J, Hill B, Ijungberg O and Kinkelin Pde, 1994. The occurrence of virus infections in elvers and eel (*Anguilla Anguilla*) in Europa with particular reference to VHSV and IHN. *Aquaculture*, 123, 11–19.
- Kamei Y, Yoshimizu M, Ezura Y and Kimura T, 1987. Effects of estuarine and marine waters on the infectivities of infectious hematopoietic necrosis virus (IHN) and infectious pancreatic necrosis virus (IPNV). *Bulletin of Fisheries Sciences, Hokkaido University*, 38, 271–285. Available online: [https://eprints.lib.hokudai.ac.jp/dspace/bitstream/2115/23960/1/38\(3\)_P271-285.pdf](https://eprints.lib.hokudai.ac.jp/dspace/bitstream/2115/23960/1/38(3)_P271-285.pdf)
- Kempton J, Sadowski J, Schütze H, Fischer U, Dauber M, Fichtner D, Panicz R and Bergmann SM, 2009. Koi herpes virus: do acipenserid restitution programs pose a threat to carp farms in the disease-free zones? *Acta Ichthyologica et Piscatoria*, 39, 119–126.
- Kempton J, Kielpinski M, Panicz R, Sadowski J, Myslowski B and Bergmann SM, 2012. Horizontal transmission of koi herpes virus (KHV) from potential vector species to common carp. *Bulletin of the European Association of Fish Pathologists*, 32, 212–219.
- Kent ML, Traxler GS, Kieser D, Richard J, Dawe SC, Shaw RW, Prosperi-Porta G, Ketcheson J and Evelyn TPT, 1998. Survey of Salmonid pathogens in Ocean-caught fishes in British Columbia, Canada. *Journal of Aquatic Animal Health*, 10, 211–219.
- Kibenge MT, Opazo B, Rojas AH and Kibenge FSB, 2002. Serological evidence of infectious salmon anaemia virus (ISAV) infection in farmed fishes, using an indirect enzyme-linked immunosorbent assay (ELISA). *Diseases of Aquatic Organisms*, 51, 1–11. <https://doi.org/10.3354/dao051001>
- King JA, Snow M, Smail DA and Raynard RS, 2001. Distribution of viral haemorrhagic septicaemia virus in wild fish species of the North Sea, north east Atlantic Ocean and Irish Sea. *Diseases of Aquatic Organisms*, 47, 81–86. <https://doi.org/10.3354/dao047081>
- Kim W-S, Oh S-Y and Oh M-J, 2013. Susceptibility of marine medaka *Oryzias dancena* to fish pathogenic viruses. *Journal of Fish Pathology*, 26, 283–287.
- Knuesel R, Segner H and Wahli T, 2003. A survey of viral diseases in farmed and feral salmonids in Switzerland. *Journal of Fish Diseases*, 26, 167–182. <https://doi.org/10.1046/j.1365-2761.2003.00447.x>
- Langdon JS, 1989. Experimental transmission and pathogenicity of epizootic hematopoietic necrosis virus (EHNV) in redfin perch, *Perca fluviatilis* L., and 11 other teleosts. *Journal of Fish Diseases*, 12, 295–310. <https://doi.org/10.1111/j.1365-2761.1989.tb00318.x>
- Lapatra SE, Jones GR, Lauda KA, McDowell TS, Schneider R and Hedrick RP, 1995. White Sturgeon as a Potential Vector of Infectious Hematopoietic Necrosis Virus. *Journal of Aquatic Animal Health*, 7, 225–230.

- LaPatra SE, Barone L, Jones GR and Zon LI, 2000. Effects of infectious hematopoietic necrosis virus and infectious pancreatic necrosis virus infection on hematopoietic precursors of the zebrafish. *Blood Cells, Molecules, and Diseases*, 26, 445–452.
- Lee W-L, Yun H-M, Kim S-R, Jung S-J and Oh M-J, 2007. Detection of viral hemorrhagic septicemia virus (VHSV) from marine fish in the South Western Coastal Area and East China Sea. *Journal of Fish Pathology*, 20, 201–209.
- Lievens B, Frans I, Heusdens C, Juste A, Jonstrup SP, Lieffrig F and Willems KA, 2011. Rapid detection and identification of viral and bacterial fish pathogens using a DNA array-based multiplex assay. *Journal of Fish Diseases*, 34, 861–875.
- Lilthved H, Vogelsang C, Modahl I and Dannevig BH, 2006. Dannevig high resistance of fish pathogenic viruses to UV irradiation and ozonated seawater. *Aquacultural Engineering*, 34, 72–82. Available online: <https://www.sciencedirect.com/science/article/abs/pii/S0144860905000683>
- Liu Z, Teng Y, Liu H, Jiang Y, Xie XY, Li H, Lv J, Gao L, He J, Shi X, Tian F, Yang J and Xie C, 2008. Simultaneous detection of three fish rhabdoviruses using multiplex real-time quantitative RT-PCR assay. *Journal of Virological Methods*, 149, 103–109.
- Lopez-Vazquez C, Raynard RS, Bain N, Snow M, Bandin I and Dopazo CP, 2006. Genotyping of marine viral haemorrhagic septicaemia virus isolated from the Flemish Cap by nucleotide sequence analysis and restriction fragment length polymorphism patterns. *Diseases of Aquatic Organisms*, 73, 23–31.
- Lorenzen N and Olesen NJ, 1995. Multiplication of VHS virus in insect cells. *Veterinary Research*, 26, 428–432. Available online: <https://hal.science/hal-00902369/document>
- Maganha SRL, Cardoso PHM, Balian SC, Almeida-Queiroz SR, Fernandes AM and Sousa RLM, 2022. Molecular detection and phylogenetic analysis of Cyprinid herpesvirus 3 in Brazilian ornamental fish. *Brazilian Journal of Microbiology*, 53, 1807–1815.
- Matras M, Stachnik M, Borzym E, Maj-Paluch J and Reichert M, 2019. Potential role of different fish species as vectors of koi herpesvirus (CyHV-3) infection. *Journal of Veterinary Research*, 63, 507–511.
- Meyer K, Bergmann SM, Mvd M and Steinhagen D, 2012. Detection of Koi herpesvirus: impact of extraction method, primer set and DNA polymerase on the sensitivity of polymerase chain reaction examinations. *Aquaculture Research*, 43, 835–842.
- Meyers TR, Short S and Lipson K, 1999. Isolation of the North American strain of viral hemorrhagic septicemia virus (VHSV) associated with epizootic mortality in two new host species of Alaskan marine fish. *Diseases of Aquatic Organisms*, 38, 81–86. <https://doi.org/10.3354/dao038081>
- Minamoto T, Honjo MN, Uchii K, Yamanaka H, Suzuki AA, Kohmatsu Y, Iida T and Kawabata Z, 2009a. Detection of cyprinid herpesvirus 3 DNA in river water during and after an outbreak. *Veterinary Microbiology*, 135, 261–266.
- Minamoto T, Honjo MN and Kawabata Z, 2009b. Seasonal distribution of Cyprinid Herpesvirus 3 in Lake Biwa, Japan. *Applied and Environmental Microbiology*, 75, 6900–6904.
- Mortensen HF, Heuer OE, Lorenzen N, Otte L and Olesen NJ, 1999. Isolation of viral haemorrhagic septicaemia virus (VHSV) from wild marine fish species in the Baltic Sea, Kattegat, Skagerrak and the North Sea. *Virus Research*, 63, 95–106. [https://doi.org/10.1016/s0168-1702\(99\)00062-3](https://doi.org/10.1016/s0168-1702(99)00062-3)
- Nishizawa T, Kinoshita S, Kim WS, Higashi S and Yoshimizu M, 2006. Nucleotide diversity of Japanese isolates of infectious hematopoietic necrosis virus (IHN) based on the glycoprotein gene. *Diseases of Aquatic Organisms*, 71, 267–272.
- Nylund A, Devold M, Mullins J and Plarre H, 2002. Herring (*Clupea harengus*): A host for Infectious Salmon Anemia Virus (ISAV). *Bulletin of the European Association of Fish Pathologists*, 22, 311–318.
- Ogut H and Altuntas C, 2014. Survey of viral haemorrhagic septicaemia virus in wild fishes in the southeastern Black Sea. *Diseases of Aquatic Organisms*, 109, 99–106. <https://doi.org/10.3354/dao02728>
- Oidtmann B, Dixon P, Way K, Joiner C and Bayley AE, 2017. Risk of waterborne virus spread – review of survival of relevant fish and crustacean viruses in the aquatic environment and implications for control measures. *Reviews in aquaculture*, 10, 641–669.
- Palmer AD and Emmenegger EJ, 2014. Susceptibility of koi and yellow perch to infectious hematopoietic necrosis virus by experimental exposure. *Journal of Aquatic Animal Health*, 26, 78–83. <https://doi.org/10.1080/08997659.2014.886634>
- Polinski MP, Fehringer TR, Johnson KA, Snekvik KR, LaPatra SE, LaFrentz BR, Ireland SC and Cain KD, 2010. Characterization of susceptibility and carrier status of burbot, *Lota lota* (L.), to IHN, IPNV, *Flavobacterium psychrophilum*, *Aeromonas salmonicida* and *Renibacterium salmoninarum*. *Journal of Fish Diseases*, 33, 559–570. <https://doi.org/10.1111/j.1365-2761.2010.01152.x>
- Pospichal A, Piackova V, Pokorova D and Vesely T, 2016. Susceptibility of stone loach (*Barbatula barbatula*) and hybrids between sterlet (*Acipenser ruthenus*) and beluga (*Huso huso*) to cyprinid herpesvirus 3. *Veterinárni Medicina*, 61, 249–255.
- Pospichal A, Pokorova D, Vesely T and Piackova V, 2018. Susceptibility of the topmouth gudgeon (*Pseudorasbora parva*) to CyHV-3 under no-stress and stress conditions. *Veterinárni Medicina*, 63, 229–239.
- Radosavljevic V, Jeremic S, Cirkovic M, Lako B, Milicevic V, Potkonjak A and Nikolin V, 2012. Common fish species in polyculture with carp as cyprinid herpes virus 3 carriers. *Acta Veterinaria*, 62, 675–681.

- Rianto WB, Uun Y, Mohamad F and Sri A, 2019. Expression of HSP70 in koi herpesvirus-infected tilapia (*Oreochromis niloticus*). *Russian Journal of Agricultural and Socio-Economic Sciences*, 3, 39–44.
- Rolland JB and Winton JR, 2003. Relative resistance of Pacific salmon to infectious salmon anaemia virus. *Journal of Fish Diseases*, 9, 511–520.
- Sandlund N, Gjerset B, Bergh Ø, Modahl I, Olesen NJ and Johansen R, 2014. Screening for viral hemorrhagic septicemia virus in marine fish along the Norwegian coastal line. *PLoS One*, 9, e108529.
- Shimizu T, Yoshida N, Kasai H and Yoshimizu M, 2006. Survival of koi herpesvirus (KHV) in environmental water. *Fish Pathology*, 41, 153–157.
- Skall HF, Olesen NJ and Møllergaard S, 2005. Prevalence of viral haemorrhagic septicaemia virus in Danish marine fishes and its occurrence in new host species. *Diseases of Aquatic Organisms*, 66, 145–151.
- Skall HF, Jørgensen C and Olesen NJ, 2015. Evaluation of the effect of percolation and NaCl solutions on viral haemorrhagic septicaemia virus (VHSV) under experimental conditions. *Aquaculture*, 448, 507–511. Available online: <https://reader.elsevier.com/reader/sd/pii/S0044848615300727?token=69EBA8A7D2AEFF7031E04A2E4D00C945C3A846A287B1B76F4F36E335EB63CA9571E666758BE3EACB86A59A8ADDEEB4F3&originRegion=eu-west-1&originCreation=20230307112820>
- Snow M, Raynard RS and Bruno DW, 2001. Comparative susceptibility of Arctic char (*Salvelinus alpinus*), rainbow trout (*Oncorhynchus mykiss*) and brown trout (*Salmo trutta*) to the Scottish isolate of infectious salmon anaemia virus. *Aquaculture*, 196, 47–54.
- Snow M, King JA, Garden A and Raynard RS, 2005. Experimental susceptibility of Atlantic cod, *Gadus morhua* (L.), and Atlantic halibut, *Hippoglossus hippoglossus* (L.), to different genotypes of viral haemorrhagic septicaemia virus. *Journal of Fish Diseases*, 28, 737–742.
- Stepien CA and Niner MD, 2020. Evolutionary trajectory of fishPiscine novirhabdovirus (=Viral Hemorrhagic Septicemia Virus) across its Laurentian Great Lakes history: Spatial and temporal diversification. *Ecology and Evolution*, 10, 9740–9775.
- Thompson TM, Batts WN, Faisal M, Bowser P, Casey JW, Phillips K, Garver KA, Winton J and Kurath G, 2011. Emergence of Viral hemorrhagic septicemia virus in the North American Great Lakes region is associated with low viral genetic diversity. *Diseases of Aquatic Organisms*, 96, 29–43. <https://doi.org/10.3354/dao02362>
- Vike S, Oelckers K, Duesund H, Erga SR, Gonzalez J, Hamre B, Frette O and Nylund AJ, 2014. Infectious salmon anemia (ISA) virus: infectivity in seawater under different physical conditions. *Aquatic Animal Health*, 26, 33–42. <https://doi.org/10.1080/08997659.2013.864720>
- Wallace IS, Donald K, Munro LA, Murray W, Pert CC, Stagg H, Hall M and Bain N, 2016. A survey of wild marine fish identifies a potential origin of an outbreak of viral haemorrhagic septicaemia in wrasse, Labridae, used as cleaner fish on marine Atlantic salmon, *Salmo salar* L., farms. *Journal of Fish Diseases*, 38, 515–521.
- Wolf K, 1988. *Fish viruses and fish viral diseases*. Ithaca.
- WOAH, 2022. *Aquatic Animal Health Code*. Chapter 4.7 Following in aquaculture. Available online: https://www.woah.org/fileadmin/Home/eng/Health_standards/aahc/current/chapitre_following_aquaculture.pdf
- WOAH, 2023. *Manual of Diagnostic Tests for Aquatic Animals 2022*. Chapter 2.3.6. Infection with Koi herpes virus. Available online: <https://www.woah.org/en/what-we-do/standards/codes-and-manuals/aquatic-manual-online-access/>

Abbreviations

AHL	Animal Health Law
EHNV	epizootic haematopoietic necrosis virus
ELR	extensive literature review
EURL	EU Reference Laboratory
His	histology
HPRA	ISAV highly polymorphic region-deleted infectious salmon anaemia virus
IHN	infectious haematopoietic necrosis virus
ISH	<i>in situ</i> hybridisation
ITS	internal transcribed spacer
KHV	Koi herpes virus
PCR	polymerase chain reaction
RFLP	restriction fragment length polymorphism RFTM Ray's fluid thioglycollate medium
ToR	Term of Reference
Seq	sequencing
VHSV	viral haemorrhagic septicaemia virus
WOAH	World Organisation for Animal Health

Appendix A – Currently listed vector or reservoir species without sufficient evidence in peer-reviewed papers.

Table A.1: Currently listed Vectors/reservoirs in Commission Implementing Regulation 1882/2018 for which no evidence or insufficient evidence was generated by the extensive literature review

Infection with EHN				
VECTOR/ RESERVOIRS* SPECIES Scientific name	VECTOR/ RESERVOIRS* SPECIES Common name	Reference	Reasoning	Certainty
<i>Aristichthys nobilis</i>	Bighead carp	No eligible paper found	NA	NA
<i>Carassius auratus</i>	Goldfish	Jensen et al. (2011)	In the study it was not possible to re-isolate virus from any of the samples from goldfish experimentally challenged with EHN.	0–10%
<i>Carassius carassius</i>	Crucian carp	No eligible paper found	NA	NA
<i>Cyprinus carpio</i>	Common carp	Jensen et al. (2011)	In the study, it was not possible to re-isolate virus from any of the samples from carp experimentally challenged with EHN.	NA
<i>Hypophthalmichthys molitrix</i>	Silver carp	No eligible paper found	NA	NA
<i>Leuciscus</i> spp.	Chub spp.	No eligible paper found	NA	NA
<i>Rutilus rutilus</i>	Roach	No eligible paper found	NA	NA
<i>Scardinius erythrophthalmus</i>	Rudd	No eligible paper found	NA	NA
<i>Tinca tinca</i>	Tench	No eligible paper found	NA	NA
Infection with VHS				
VECTOR/ RESERVOIRS* Scientific name	VECTOR/ RESERVOIRS* Common name	Reference	Reasoning	Certainty
<i>Acipenser baerii</i>	Siberian sturgeon	No eligible paper found	NA	NA
<i>Acipenser gueldenstaedtii</i>	Danube sturgeon	No eligible paper found	NA	NA
<i>Acipenser ruthenus</i>	Sterlet sturgeon	No eligible paper found	NA	NA
<i>Acipenser stellatus</i>	Starry sturgeon	No eligible paper found	NA	NA
<i>Acipenser sturio</i>	Sturgeon	No eligible paper found	NA	NA
<i>Ameiurus melas</i>	Black bullhead	No eligible paper found	NA	NA
<i>Argyrosomus regius</i>	Meagre	No eligible paper found	NA	NA
<i>Aristichthys nobilis</i>	Bighead carp	No eligible paper found	NA	NA
<i>Clarias gariepinus</i>	North African catfish	No eligible paper found	NA	NA
<i>Dentex dentex</i>	Common dentex	No eligible paper found	NA	NA
<i>Dicentrarchus labrax</i>	European seabass	No eligible paper found	NA	NA
<i>Diplodus puntazzo</i>	Sharpsnout seabream	No eligible paper found	NA	NA
<i>Diplodus vulgaris</i>	Common two-banded seabream	No eligible paper found	NA	NA
<i>Epinephelus aeneus</i>	White grouper	No eligible paper found	NA	NA
<i>Epinephelus marginatus</i>	Dusky grouper	No eligible paper found	NA	NA
<i>Huso huso</i>	Beluga	No eligible paper found	NA	NA
<i>Hypophthalmichthys molitrix</i>	Silver carp	No eligible paper found	NA	NA

<i>Ictalurus</i> spp.	Catfish spp.	No eligible paper found	NA	NA
<i>Leuciscus</i> spp.	Chub spp.	No eligible paper found	NA	NA
<i>Morone chrysops</i> ×	White bass ×	No eligible paper found	NA	NA
<i>Morone saxatilis</i>	Striped bass hybrids	No eligible paper found	NA	NA
<i>Mugil cephalus</i>	Flathead grey mullet	No eligible paper found	NA	NA
<i>Oreochromis</i>	Tilapia	No eligible paper found	NA	NA
<i>Pagellus bogaraveo</i>	Blackspot seabream	No eligible paper found	NA	NA
<i>Pagellus erythrinus</i>	Common pandora	No eligible paper found	NA	NA
<i>Pangasius pangasius</i>	Pangas catfish	No eligible paper found	NA	NA
<i>Rutilus rutilus</i>	Roach	No eligible paper found	NA	NA
<i>Salvelinus fontinalis</i>	Brook trout	No eligible paper found	NA	NA
<i>Sander lucioperca</i>	pikeperch	No eligible paper found	NA	NA
<i>Scardinius erythrophthalmus</i>	Rudd	No eligible paper found	NA	NA
<i>Sciaenops ocellatus</i>	Red drum	No eligible paper found	NA	NA
<i>Silurus glanis</i>	Wels catfish	No eligible paper found	NA	NA
<i>Solea senegalensis</i>	Senegalese sole	No eligible paper found	NA	NA
<i>Solea solea</i>	Common sole	No eligible paper found	NA	NA
<i>Sparus aurata</i>	Gilthead seabream	No eligible paper found	NA	NA
<i>Thunnus</i> spp.	True tuna spp.	No eligible paper found	NA	NA
<i>Thunnus thynnus</i>	Atlantic bluefin tuna	No eligible paper found	NA	NA
<i>Tinca tinca</i>	Tench	No eligible paper found	NA	NA
<i>Umbrina cirrosa</i>	Shi drum	No eligible paper found	NA	NA

Infection with IHNV

VECTOR/ RESERVOIRS* Scientific name	VECTOR/ RESERVOIRS* Common name	Reference	Reasoning	Certainty
<i>Acipenser baerii</i>	Siberian sturgeon	No eligible paper found	NA	NA
<i>Acipenser gueldenstaedtii</i>	Danube sturgeon	No eligible paper found	NA	NA
<i>Acipenser ruthenus</i>	Sterlet sturgeon	No eligible paper found	NA	NA
<i>Acipenser stellatus</i>	Starry sturgeon	No eligible paper found	NA	NA
<i>Acipenser sturio</i>	Sturgeon	No eligible paper found	NA	NA
<i>Ameiurus melas</i>	Black bullhead	No eligible paper found	NA	NA
<i>Aristichthys nobilis</i>	Bighead carp	No eligible paper found	NA	NA
<i>Astacus astacus</i>	Noble Crayfish	No eligible paper found and not a fish species	NA	NA
<i>Carassius auratus</i>	Goldfish	No eligible paper found	NA	NA
<i>Carassius carassius</i>	Crucian carp	No eligible paper found	NA	NA
<i>Clarias gariepinus</i>	North African catfish	No eligible paper found	NA	NA
<i>Gadus morhua</i>	Atlantic cod	No eligible paper found	NA	NA
<i>Hippoglossus hippoglossus</i>	Atlantic halibut	No eligible paper found	NA	NA
<i>Hypophthalmichthys molitrix</i>	Silver carp	No eligible paper found	NA	NA
<i>Huso huso</i>	Beluga	No eligible paper found	NA	NA
<i>Ictalurus punctatus</i>	Channel catfish	No eligible paper found	NA	NA
<i>Ictalurus</i> spp.	Catfish spp.	No eligible paper found	NA	NA
<i>Leuciscus</i> spp.	Chub spp.	No eligible paper found	NA	NA
<i>Melanogrammus aeglefinus</i>	Haddock	No eligible paper found	NA	NA
<i>Pacifastacus leniusculus</i>	Signal Crayfish	No eligible paper found	NA	NA

<i>Procambarus clarkii</i>	Red Swamp crawfish	No eligible paper found and not a fish species	NA	NA
<i>Pangasius pangasius</i>	Pangas catfish	No eligible paper found	NA	NA
<i>Platichthys flesus</i>	European flounder	No eligible paper found	NA	NA
<i>Rutilus rutilus</i>	Roach	No eligible paper found	NA	NA
<i>Sander lucioperca</i>	Pikeperch	No eligible paper found	NA	NA
<i>Scardinius erythrophthalmus</i>	Rudd	No eligible paper found	NA	NA
<i>Silurus glanis</i>	Welsh catfish	No eligible paper found	NA	NA
<i>Tinca tinca</i>	Tench	No eligible paper found	NA	NA

*Regulation 1882/2018 does not differentiate between vectors or reservoir species.

Appendix B – Vector or reservoir species mentioned in peer-reviewed papers that were excluded in the ELR

Table B.1 lists species for which studies were retrieved through searching the electronic databases, but that were excluded because they did not pass the eligibility criteria; or species for which evidence was extracted from relevant studies, but the certainty of the assessment was too low for classification as vector or reservoir.

Table B.1: Species that were excluded during the eligibility screening and assessment

Infection with EHN _V					
Scientific name	Reference	Certainty	Reasoning	Conclusion WG and AHAW Panel	Suggested classification in previous EURL report (2022)
<i>Craterocephalus fulvus</i>	Becker et al. (2013)	0–10%	Bath challenge and IP challenge. Same or slightly increased mortality as control fish depending on the species. No virus detected.	Not classified	Not assessed
<i>Hypseleotris klunzingeri</i>					
<i>Maccullochella macquariensis</i>					
<i>Maccullochella peelii</i>					
<i>Macquaria ambigua</i>					Vector/Reservoir
<i>Mogurnda adspersa</i>					Vector/Reservoir
<i>Nannoperca australis</i>					Not assessed
<i>Salmo salar</i>	Langdon (1989)	NA	The study was excluded in the first level of screening because it investigates the presence of the pathogen in rainbow trout, <i>Salmo gairdneri</i> , which is an already known susceptible species.	Not assessed	Vector/Reservoir
Infection with HPR _Δ ISAV					
Scientific name	Reference	Certainty	Reasoning	Conclusion WG and AHAW Panel	Suggested classification in previous EURL report (2022)
<i>Salvelinus alpinus</i>	Snow et al. (2001)	33–66%	According to the paper, ISAV could only be detected by RT-PCR following IP-injection of ISAV. No positives by cell cultivation. 9/10 fish positive at 28 dpi and no fish positive at 40 dpi.	Not classified	Not assessed

Infection with VHSV					
Scientific name	Reference	Certainty	Reasoning	Conclusion WG and AHAW Panel	Suggested classification in previous EURL report (2022)
<i>Acanthopagrus schlegelii</i>	EFSA. Aquatic species susceptible to diseases listed in Directive 2006/88/EC 1 Scientific Opinion of the Panel on Animal Health and Welfare (AHAW) Adopted on the 11th of September 2008. The EFSA Journal. 2008; September: 1–144.	NA	Excluded because it's not a primary research study	Not assessed	Vector/Reservoir
<i>Ammodytes personatus</i>	Skall et al. (2005)	NA	Excluded because it's not a primary research study	Not assessed	Vector/Reservoir
<i>Gasterosteus mustelus</i>	Skall et al. (2005)	NA	Excluded because it's not a primary research study	Not assessed	Vector/Reservoir
<i>Glyptocephalus stelleri</i>	Lee et al. (2007)	NA	Excluded because the full text is not written in any EU language	Not assessed	Vector/Reservoir
<i>Hoplobrotula armata</i>	Lee et al. (2007)	NA	Excluded because the full text is not written in any EU language	Not assessed	Vector/Reservoir
<i>Larimichthys polyactis</i>	Lee et al. (2007)	NA	Excluded because the full text is not written in any EU language	Not assessed	Vector/Reservoir
<i>Liparis tessellatus</i>	Lee et al. (2007)	NA	Excluded because the full text is not written in any EU language	Not assessed	Vector/Reservoir
<i>Moina macrocopa</i>	Ito and Olesen (2007)	NA	Excluded at the first level of screening because the study does not investigate any not known susceptible species	Not assessed	Vector/Reservoir
<i>Oncorhynchus aguabonita</i>	Wolf (1988)	NA	Excluded because the full text is not written in any EU language	Not assessed	Vector/Reservoir
<i>Oryzias dancena</i>	Kim et al. (2013)	NA	Excluded because the full text is not written in any EU language	Not assessed	Vector/Reservoir

<i>Pampus argenteus</i>	Lee et al. (2007)	NA	Excluded because the full text is not written in any EU language	Not assessed	Vector/Reservoir
<i>Oryzias dancena</i>	Kim et al. (2013)	NA	Excluded because the full text is not written in any EU language	Not assessed	Vector/Reservoir
<i>Sebastes schlegeli</i>	Isshiki et al. (2003)	10–33%	Unclear %, antibodies detected by VNT.	Not classified	Not assessed
<i>Scorpaena izensis</i>	Lee et al. (2007)	NA	Excluded because the full text is not written in any EU language	Not assessed	Vector/Reservoir
<i>Scylliorhinus torazame</i>	Lee et al. (2007)	NA	Excluded because the full text is not written in any EU language	Not assessed	Vector/Reservoir
<i>Semotilus corporalis</i>	Cornwell et al. (2015)	NA	Excluded because the full text wasn't available	Not assessed	Vector/Reservoir
<i>Seriola dumerili</i>	OIE. Viral Haemorrhagic Septicaemia. OIE Aquatic Animal Disease Cards. 2017	NA	Excluded because it's not a primary research study	Not assessed	Vector/Reservoir
<i>Trichiurus lepturus</i>	Lee et al. (2007)	NA	Excluded because the full text is not written in any EU language	Not assessed	Vector/Reservoir

Infection with IHN					
Scientific name	Reference	Certainty	Reasoning		Suggested classification in previous EURL report (2022)
<i>Anguilla anguilla</i>	Bergmann et al. (2003)	NA	Excluded at the second level of screening because the study investigates the susceptibility of rainbow trout, which is an already known susceptible species (the origin of the isolates was not taken into account)	Not assessed	Vector/Reservoir
	Jørgensen et al. (1994)	NA	Excluded at the second level of screening because IHN was not identified in/on species X.	Not assessed	Vector/Reservoir

<i>Lota maxima</i>	Polinski et al. (2010)	33–66%	Reduced survival and virus re-isolation. Mortality in negative control not explained.	Not classified	Vector/Reservoir
<i>Plecoglossus altivelis</i>	Nishizawa et al. (2006)	NA	Excluded at the first level of screening	Not assessed	Vector/Reservoir
<i>Schopthalmus maximus</i>	EFSA (2008) Scientific Opinion of the Panel on AHAW on a request from the European Commission on aquatic animal species	NA	Excluded because it's not a primary research study	Not assessed	Vector/Reservoir
	Liu et al. (2008)	10–33%	Lack of details on diagnostic methods	Not classified	Vector/Reservoir
<i>Thymallus thymallus</i>	Follett et al. (1997)	NA	According to the study, Arctic grayling was refractory to experimental infection with IHNV	Not assessed	Vector/Reservoir

Infection with KHV					
Scientific name	Reference	Certainty	Reasoning	Conclusion WG and AHAW Panel	Suggested classification in previous EURL report (2022)
<i>Ameiurus nebulosus</i>	Matras et al. (2019)	10–33%	Detection of KHV DNA up to 4 weeks.	Not classified	Not assessed
<i>Ancistrus</i> spp.	Bergmann et al. (2009)	33–66%	Non-validated method used to detect KHV and a few positive found at 2nd round of nested PCR in species X. Conclusion is not supported by data.	Not classified	Not assessed
<i>Ameiurus nebulosus</i>	Fabian et al. (2013)	0–10%	KHV detected by standard PCR method, at VERY low copy number (< 3 in mean for all positive species). Only DNA is detected in exposed carp (which were KHV free) no transfer of disease. Possibly water contamination with DNA.	0–10%	Not assessed

<i>Carassius auratus</i>	Lievens et al. (2011)	0–10%	PCR positive, but only gills were tested. Paper does not focus on evidence collection on vectors or reservoirs	Not classified	Not assessed
	Meyer et al. (2012)	NA	Not relevant: pathogen injection to study immune response	Not classified	Not assessed
<i>Esox lucius</i>	Fabian et al. (2013)	0–10%	KHV detected by standard PCR method, at VERY low copy number (< 3 in mean for all positive species). Only DNA is detected in exposed carp (which were KHV free) no transfer of disease. Possibly water contamination with DNA.	0–10%	Not assessed
<i>Gasterosteus aculeatus</i>	Matras et al. (2019)	10–33%	Detection of KHV DNA only 2 weeks after infection.	Not classified	Not assessed
<i>Gobio gobio</i>	Fabian et al. (2013)	0–10%	KHV detected by standard PCR method, at VERY low copy number (< 3 in mean for all positive species). Only DNA is detected in exposed carp (which were KHV free) no transfer of disease. Possibly water contamination with DNA.	0–10%	Not assessed
<i>Hyphessobrycon eques</i>	Maganha et al. (2022)	10–33%	Only one detection by PCR and sequencing	Not classified	Not assessed
<i>Leuciscus idus</i>	Bergmann et al. (2009)	33–66%	Non-validated method used to detect KHV and a few positive found at 2nd round of nested PCR in species X. Conclusion is not supported by data.	Not classified	Vector/Reservoir

<i>Leuciscus leuciscus</i>	Fabian et al. (2013)	0–10%	KHV detected by standard PCR method, at VERY low copy number (< 3 in mean for all positive species). Only DNA is detected in exposed carp (which were KHV free) no transfer of disease. Possibly water contamination with DNA.	0–10%	Not assessed
<i>Misgurnus anguillicaudatus</i>	Maganha et al. (2022)	10–33%	Only one detection by PCR and sequencing	Not classified	Not assessed
<i>Neogobius melanostomus</i>	Jin et al. (2020)	0–10%	Only states tissue, not clear if it is gills included. Could be environmental contamination.	Not classified	Not assessed
<i>Oncorhynchus mykiss</i>	Bergmann et al. (2021)	33–66%	According to the findings in the paper, a portion of rainbow trout can be experimentally (immersion/ cohabitant) infected with KHV-E (no disease), produce antibodies against the virus and transfer the virus to naive carp. No isolation of virus in cell culture from either species.	Not classified	Not assessed
<i>Oreochromis niloticus</i>	Rianto et al. (2019)	10–33%	Only gills were tested and lack of details of pathogen confirmation by reference methods	Not classified	
<i>Pygocentrus nattereri</i>	Maganha et al. (2022)	10–33%	Only one detection by PCR and sequencing	Not classified	Not assessed
<i>Xiphophorus maculatus</i>	Maganha et al. (2022)	10–33%	Only one detection by PCR and no sequencing	Not classified	Not assessed

PCR: polymerase chain reaction; IP: intraperitoneal injection; EURL: EU Reference Laboratory; IP: intraperitoneal challenge; dpi: days post infection; VNT: virus neutralisation test.

Annex A – Protocol

Annex A is available under the Supporting Information section on the online version of the scientific output.