



## Properties of Excitons Bound to Ionized Donors

**Skettrup, Torben; Suffczynski, M.; Gorzkowski, W.**

*Published in:*  
Physical Review B Condensed Matter

*Link to article, DOI:*  
[10.1103/PhysRevB.4.512](https://doi.org/10.1103/PhysRevB.4.512)

*Publication date:*  
1971

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Skettrup, T., Suffczynski, M., & Gorzkowski, W. (1971). Properties of Excitons Bound to Ionized Donors. *Physical Review B Condensed Matter*, 4(2), 512-517. <https://doi.org/10.1103/PhysRevB.4.512>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Properties of Excitons Bound to Ionized Donors

T. Skettrup

*Physics Laboratory III, Technical University of Denmark, Lyngby, Denmark*

and

M. Suffczynski and W. Gorzkowski

*Institute of Physics, Polish Academy of Sciences, Warsaw, Poland*

(Received 11 December 1970)

Binding energies, interparticle distances, oscillator strengths, and exchange corrections are calculated for the three-particle complex corresponding to an exciton bound to an ionized donor. The results are given as functions of the mass ratio of the electron and hole. Binding of the complex is obtained for mass ratios up to 0.426. The interparticle distances are up to 50 times larger than the corresponding exciton radius. The oscillator strengths are about  $10^4$  times greater than those of free excitons, while the exchange corrections for the complex are comparable to those of free excitons. The results are applied to CdS and ZnO and compared with experimental results on these.

## I. INTRODUCTION

The three-particle complex consisting of an electron and a hole bound to an ionized donor is the simplest possible bound-exciton complex. Hopfield<sup>1</sup> has estimated the binding energies of bound-exciton complexes by extrapolations of molecular binding energies. More accurate calculations of binding energies for the three-particle complex were carried out by means of variational methods by Sharma and Rodriguez<sup>2</sup> and Suffczynski *et al.*<sup>3</sup>

Here we compute binding energies of the three-particle complex for a practically important interval of electron-hole mass ratios  $m_e/m_h$  by adding a long-range tail to the variational function. The wave functions obtained in this variational procedure are then, furthermore, used for computing other characteristic quantities of the center, such as interparticle distances, oscillator strengths, and exchange corrections as a function of  $m_e/m_h$ .

## II. GROUND-STATE ENERGY

The ground-state energy of the three-particle complex was calculated by variational means in the effective-mass approximation. The effective-mass Hamiltonian  $H$  for the complex is conveniently scaled as follows:

$$H^1 = H/2E_a$$

$$= -\frac{1}{2} (m_h/m_e) \Delta'_e - \frac{1}{2} \Delta'_h + 1/R'_h + 1/R'_e - 1/R'_{eh}. \quad (1)$$

Here  $R_h$  and  $R_e$  are the distances from the ionized donor to the hole and the electron, respectively, while  $R_{eh}$  is the distance between the electron and hole. Further,

$$E_a = \frac{e^4 m_h}{2\hbar^2 (4\pi\epsilon_0\epsilon)^2} = \frac{\hbar^2}{2m_h a_h^2} = \frac{e^2}{2(4\pi\epsilon_0\epsilon) a_h} \quad (2)$$

and

$$a_h = 4\pi\epsilon_0\hbar^2/e^2 m_h \quad (3)$$

are energy and radius for the hydrogenlike acceptor in the semiconductor. The  $e$  is the electronic charge,  $m_e$  and  $m_h$  are electron and hole masses, respectively,  $\epsilon_0$  is the permittivity vacuum, while  $\epsilon$  is the low-frequency value of the dielectric constant of the material. The primed coordinates are then measured in units of  $a_h$ . They are thus scaled with the dielectric constant and the hole mass. It is seen that the scaled Hamiltonian depends on only one parameter, the mass ratio  $m_e/m_h$ . In case of anisotropic masses or dielectric constants a transformation<sup>4</sup> to the isotropic case can be applied. Hence, only the isotropic solutions will be considered here.

The trial function was of the form,

$$f(\vec{R}_e, \vec{R}_h) = R_h^p \sum_{i=1}^N X_i \exp[-(A_i R'_e + B_i R'_{eh} + C_i R'_h)]$$

$$+ X_{N+1} e^{-A_0 R'_e} \frac{e^{-C_0 R'_h} - e^{-C R'_h}}{R'_h}, \quad (4)$$

where the variational parameters are  $X_i$ ,  $A_i$ ,  $B_i$ ,  $C_i$ ,  $C_0$ , and  $C$ , whereas  $1/A_0$  is the radius of the donor.  $p$  and  $N$  were usually 2 and 4, respectively. Previous calculations<sup>3</sup> of the ground-state energy were performed only with the first part of this trial function. The inclusion of the last term which is similar to that used by Rotenberg and Stein<sup>5</sup> is particularly useful when  $m_e/m_h$  is close to the value where this complex ceases to be bound. Near this limit the center is characterized, approximately, by an electron in the donor level and a loosely bound hole as described by the last term in the trial function.

By means of (4) the variational estimate of the ground-state energy was obtained as a function of




FIG. 1. Binding energy  $E$  for the three-particle center relative to the donor energy  $E_d$  as a function of the mass ratio.  $m_e/m_h = 0$  corresponds to the  $H_2^+$  molecule. The binding energy of an exciton to an ionized donor may be found from Eq. (5).

$m_e/m_h$ . The results are given in Fig. 1. The ratio of the ground-state energy  $E$  to the donor energy  $E_d$  varies monotonically with the mass ratio.<sup>6</sup> We obtain binding of the complex up to  $m_e/m_h = 0.426$  where we found  $(E - E_d)/E_d = (1.8 \times 10^{-4})\%$ . The binding energy given in Fig. 1 is that of a hole bound to a neutral donor. The complex may equally well be visualized as an ionized donor binding an exciton with a binding energy

$$E - E_{ex} = \left[ \left( 1 + \frac{m_e}{m_h} \right) \frac{E - E_d}{E_d} + \frac{m_e}{m_h} \right] E_{ex}, \quad (5)$$




FIG. 2. Expectation values of the interparticle distances in the three-particle complex in units of the effective Bohr radius of the hole  $a_h$  ( $a_h = 4\pi\epsilon_0\epsilon\hbar^2/e^2m_h$ ). The case of  $m_e/m_h = 0.20$  is explicitly shown (in scale 1:2).

where  $E_{ex}$  is the binding energy of the free exciton.

### III. INTERPARTICLE DISTANCES

It is important to know approximately the distances between the particles in the complex since the validity of the effective-mass equation (1) breaks down when these distances are too small. It is, furthermore, of interest to calculate these distances because they yield the extension of the center and hence the largest possible concentration of bound excitons before their envelopes substantially overlap. The expectation values of the distances are computed by means of the trial functions which gave lowest ground-state energy. The results are shown in Figs. 2 and 3. It is seen that all the distances are greater than the donor radius. The use of the effective-mass equation is therefore justified. In the limit  $m_e/m_h \rightarrow 0$ ,  $\langle R_e \rangle \rightarrow \langle R_{eh} \rangle \rightarrow 0.69 \times \langle R_h \rangle$  as for the  $H_2^+$  complex. In the limit where the complex becomes ionized  $\langle R_{eh} \rangle$  and  $\langle R_h \rangle$  are large, and  $\langle R_e \rangle$  is slightly larger than the donor radius  $a_d$ .

We have also computed the average interaction energies between the particles which we define as  $\langle e^2/8\pi\epsilon_0\epsilon R \rangle$ , where  $R$  is the distance between the particles in question. These energies are two times smaller than the expectation values of the corresponding potential energies. The results are given in Fig. 4. It is seen that the binding energy of the complex appears as the sum of the interaction energies between the particles. It is interesting to note that there exists an upper limit of  $R_h$  (and  $R_{eh}$ ) since curves (a) and (b) cross at  $m_e/m_h = 0.426$ . Here, the repulsive potential energy of the hole equals the attractive potential energy, and the hole ceases to be bound. At this point  $R_h$  and  $R_{eh}$  are about 50 times greater than  $a_d$ , the donor radius, corresponding to the largest possible extension of the complex. This relatively large extension of the wave function in the direct space is consistent with the observation of Reynolds *et al.*<sup>7</sup> that the phonon side bands for three-particle complexes are sharper than for four-particle complexes. This indicates a more localized wave function in  $k$  space for three-particle complexes.




FIG. 3. Ratios of the expectation values of the interparticle distances in the three-particle complex.




FIG. 4. Expectation values of the interaction energies, defined as half the potential energies. It is seen that above the limit mass ratio the repulsive energy for the hole is larger than the attractive energy, and the complex cannot exist.

#### IV. OSCILLATOR STRENGTHS

The oscillator strength  $f_c$  of the complex may be obtained by computing the momentum matrix element  $\langle p \rangle$  between the crystal ground state and the excited state corresponding to the three-particle complex. From the definition of the oscillator strength, one then finds<sup>8</sup>

$$f_c = \frac{2|\langle p \rangle|^2}{3m\hbar\omega_0} = \left( \frac{2|p_{cv}|^2}{3m} \right) \frac{1}{\hbar\omega_0} \left| \sum_{\vec{R}} f(\vec{R}, \vec{R}) \right|^2$$

$$= \frac{P^2}{\hbar\omega_0} \left| \sum_{\vec{R}} f(\vec{R}, \vec{R}) \right|^2, \quad (6)$$

where  $m$  is the free-electron mass,  $\omega_0$  is the frequency of the transition, and  $f(\vec{R}_e, \vec{R}_h)$  is the envelope function for the three-particle complex in Eq. (4). The  $f(\vec{R}_e, \vec{R}_h)$  should be normalized such that

$$\sum_{\vec{R}_e, \vec{R}_h} |f(\vec{R}_e, \vec{R}_h)|^2 = 1. \quad (7)$$

The  $p_{cv}$  is the usual one-electron  $p$ -matrix element between valence and conduction band, and  $P^2$  is the value of the terms in the brackets. Cardona<sup>9</sup> reports that the value of  $P^2$  is remarkably constant within the different groups of materials. For II-VI compounds he finds  $P^2 \approx 21$  eV, and for III-V and IV-IV materials,  $P^2 \approx 23$  eV. The modifying factor  $|\sum f(R, R)|^2$ , which multiplies the pure band-to-band strength in (6), is not just the probability that the electron and hole are in the same unit cell, as is usually assumed in optical transitions. The factor is the square of the sum of the amplitude probabilities that the electron and hole are in same unit cell or, if Fourier transformed, that the electron and hole have opposite  $k$  vectors.

Introducing the oscillator strength  $f_{ex}$  per unit cell  $\Omega$  of the free excitons, one obtains<sup>8,10</sup>

$$f_c = f_{ex} \frac{\omega_{ex}}{\omega_0} \frac{\pi a_{ex}^3}{\Omega} \left| \sum_{\vec{R}} f(\vec{R}, \vec{R}) \right|^2, \quad (8)$$

where  $\omega_{ex}$  and  $a_{ex}$ , respectively, are the resonance frequency and the radius of the free excitons. By computing  $|\sum f(R, R)|^2$ , the ratio  $f_c/f_{ex}$  may be obtained. Rashba and Gugenishvili<sup>11</sup> and Henry and Nassau<sup>8</sup> have done this with a wave function for a short-range potential representing very loosely bound-exciton complexes. Here we compute  $|\sum f(R, R)|^2$  for excitons bound to ionized centers by means of the wave function Eq. (4) obtained in the variational procedure.

The computed values of  $|\sum f(\vec{R}, \vec{R})|^2$  for the three-particle complex are given in Fig. 5 vs  $m_e/m_h$ . It is seen that these values are of the order of 10 which means that the strengths of these complexes are of the order  $10^4$  times greater than the strengths of free excitons.

It is interesting to know the limiting value of the oscillator strength when the center ceases to be bound. To calculate this we use Eq. (6). Now, the complex consists of an electron in the donor level and a free hole in the valence band. For this complex we have

$$f_{\vec{k}}(R_e, \vec{R}_h) = \left( \frac{\Omega}{N\pi a_d^3} \right)^{1/2} e^{-R_e/a_d} e^{i\vec{k} \cdot \vec{R}_h}, \quad (9)$$

where  $N$  is the number of unit cells of volume  $\Omega$  in the crystal,  $a_d$  is the donor radius, and  $\vec{k}$  is the wave vector of the hole. Then,

$$\left| \sum_{\vec{R}} f(R, R) \right|^2 = \sum_{\vec{k}} \left| \sum_{\vec{R}} f_{\vec{k}}(R, R) \right|^2$$

$$= \frac{N\Omega}{(2\pi)^3} \int_{\vec{k}} \frac{1}{N\Omega\pi a_d^3} \frac{64\pi^2 a_d^6}{(1 + a_d^2 |\vec{k}|^2)^4} d^3 k$$

$$= 1, \quad (10)$$

where the contribution from each  $k$  vector has been




FIG. 5. Enhancement factor for the oscillator strength of the three-particle complex. The strength may be obtained from Eq. (6) or (8). Above the limit mass ratio, the enhancement factor is 1, corresponding to the integrated strength of the recombination of a free hole and a bound electron.




FIG. 6. Factor determining the exchange correction for the three-particle complex.  $\Omega$  is the volume of one molecule in the solid in units of the effective Bohr radius of the hole  $a_h$ . Since  $\sum |f(R, R)|^2$  is proportional to  $\Omega$ , the factor shown in this figure is the proportionality constant  $\sum |f(R, R)|^2/\Omega$ . The exchange splitting may be obtained from Eq. (13).

taken into account. The oscillator strength for the recombination of a free hole and a bound electron therefore is

$$f = P^2/\hbar\omega_0. \quad (11)$$

Note that this strength is independent of the mass ratio and the donor radius.

#### V. EXCHANGE CORRECTION

The levels corresponding to parallel and antiparallel orientations of the electron and hole spins are split by the exchange interaction. Only the state with antiparallel spins is optically allowed. Thomas and Hopfield<sup>12</sup> mentioned that the exchange splitting is proportional to the probability that the electron and hole are in the same unit cell, i. e.,  $\sum_{\mathbf{R}} |f(\mathbf{R}, \mathbf{R})|^2$ . Henry *et al.*<sup>13</sup> applied this result to weakly bound excitons. Here, we compute the exchange splitting of the three-particle complex by means of the wave functions of Eq. (4).

The exchange splitting may be computed from first-order perturbation theory. This has been done for free excitons.<sup>14</sup> We modify these results to apply to the case of the three-particle complex. Eq. (19) in Ref. 14 then becomes

$$\Delta_0 = J_0 \sum_{\mathbf{R}} |f(\vec{\mathbf{R}}, \vec{\mathbf{R}})|^2, \quad (12)$$

where  $J_0$  is the exchange integral of the conduction- and valence-band Wannier functions for the same unit cell. Equation (12) represents only the short-range part of the exchange interaction. The long-range part [Eq. (11) in Ref. 14] vanishes for a single isotropic center. If, however, all three-particle complexes in the crystal are taken into account, a long-range exchange splitting is found. The optically active excitations of this system of centers have different energies, depending on its transverse or longitudinal character. This splitting between longitudinal and transverse states of the

whole system of centers is due to the long-range exchange interaction. It may be computed in the same way as for free excitons.<sup>14</sup> From Eqs. (14) and (15) in Ref. 14 one then obtains for the total exchange correction of the three-particle complex

$$\Delta_T = J_0 \sum_{\mathbf{R}} |f(\vec{\mathbf{R}}, \vec{\mathbf{R}})|^2 - \frac{1}{3} (\mu_x^2 N_c / \epsilon_0 \epsilon') |\sum_{\mathbf{R}} f(\vec{\mathbf{R}}, \vec{\mathbf{R}})|^2 \quad (\vec{\mathbf{k}} \perp \vec{\mu}), \quad (13)$$

$$\Delta_L = J_0 \sum_{\mathbf{R}} |f(\vec{\mathbf{R}}, \vec{\mathbf{R}})|^2 + \frac{2}{3} (\mu_x^2 N_c / \epsilon_0 \epsilon') |\sum_{\mathbf{R}} f(\vec{\mathbf{R}}, \vec{\mathbf{R}})|^2 \quad (\vec{\mathbf{k}} \parallel \vec{\mu}). \quad (14)$$

Here  $\Delta_T$  is the exchange energy of the transverse states (optically active),  $\Delta_L$  is that of the longitudinal states,  $\mu_x$  is the magnitude of the dipole moment associated with a valence-conduction-band transition,  $\epsilon_0$  is the permittivity of vacuum, whereas  $\epsilon'$  is the background dielectric constant at the transition frequency, and  $N_c$  is the concentration of centers. A detailed calculation, based on the assumption that the random distribution of centers may be replaced by an ordered distribution, shows that  $|\sum f(\mathbf{R}, \mathbf{R})|^2$  appears as the factor due to the centers in the long-range terms and not  $\sum |f(\mathbf{R}, \mathbf{R})|^2$  which only is present in the short-range terms.

Note that the long-range exchange splitting between transverse and longitudinal states depends on the concentration of complexes in the crystal, and that it is proportional to the oscillator strength (6) of the complex. It should be noticed that there are three relevant energy states for the complex: The state with parallel spin which is not affected by exchange, and the states with antiparallel spins for which the transverse and longitudinal states are split.

The  $\sum |f(\vec{\mathbf{R}}, \vec{\mathbf{R}})|^2$  is given in Fig. 6 for the three-particle complex. It is of the order of  $10^{-3}$  which is the same order as for free excitons. Hence, the short-range exchange correction for free and bound excitons should be comparable. The long-range exchange correction  $\Delta_{LR}$  for the bound excitons may be expressed in terms of the long-range exchange correction  $\Delta_{free}$  for free excitons and the oscillator strength ratio (8) as follows:

$$\Delta_{LR} = \Delta_{free} N_c \Omega f_c / f_{ex}. \quad (15)$$

Since  $f_c/f_{ex}$  is of the order  $10^4$ ,  $N_c$  should be of the order  $10^{18} \text{ cm}^{-3}$  to get  $\Delta_{LR}$  comparable with  $\Delta_{free}$  ( $\Delta_{free} = -\frac{1}{3} E_{LT}$ , where  $E_{LT}$  is the splitting between transverse and longitudinal free excitons<sup>14</sup>). This high content of impurities will, however, because of the large oscillator strengths of bound excitons, give rise to a bound-exciton absorption comparable to the free-exciton absorption. This results in a broad absorption edge below the band gap without any structure. For most measurements on bound-

TABLE I. Results computed for CdS and ZnO compared with experimental results. Only the three experimental values of  $m_e/m_h$ ,  $m_e$ , and  $E_{ex}$  are used in the calculations.

	CdS	ZnO
Parameters		
$\sigma = m_e/m_h$	0.20 <sup>a</sup>	0.21 <sup>b</sup>
$m_e$	0.20 <sup>a</sup>	0.24 <sup>c</sup>
$E_{ex}$ (meV)	27 <sup>a</sup>	59 <sup>d</sup>
Computed results		
$(E - E_d)/E_d$ (%)	3.1	2.7
$E - E_d$ (meV)	1.0	1.9
$E - E_{ex}$ [Eq. (5)] (meV)	6.4	13.4
$ \sum f(R, R) ^2$	31	29
$f_c$ [Eq. (6)]	255	180
$f_c/f_{exA}$ <sup>e</sup>	$1.0 \times 10^5$	$0.69 \times 10^5$
$\sum  f(R, R) ^2/\Omega$ [ $a_h^{-3}$ ]	$0.45 \times 10^{-3}$	$0.49 \times 10^{-3}$
$\sum  f(R, R) ^2$ <sup>f</sup>	$0.195 \times 10^{-3}$	$0.375 \times 10^{-3}$
$\Delta_0$ <sup>g</sup> [Eq. (12)] (meV)	0.49	0.83
$\Delta_{LR}$ [Eq. (15)] (meV)	-0.044	-0.061
$\Delta_{exch}$ (meV)	0.45	0.77
Experimental values		
$\Gamma_{6-free}$ (eV)	2.55266 <sup>h</sup>	3.3758 <sup>d</sup>
$(E - E_{ex})_{Expt}$ (meV)	4.66 - 7.10 <sup>h</sup>	11.3, 17.0, 20.9 <sup>i</sup>
$(\Delta_{exch})_{Expt}$ (meV)	0.31 <sup>j</sup>	0.9 <sup>k</sup>

<sup>a</sup>Reference 4.

<sup>b</sup>Reference 15.

<sup>c</sup>W. Baer, Phys. Rev. **154**, 785 (1967).

<sup>d</sup>D. G. Thomas, J. Phys. Chem. Solids **15**, 86 (1960).

<sup>e</sup>D. G. Thomas and J. J. Hopfield, Phys. Rev. **116**, 573 (1959); Ref. 14.

<sup>f</sup> $\Omega_{CdS} = 49.4 \text{ \AA}^3$ ,  $\Omega_{ZnO} = 23.7 \text{ \AA}^3$  (per molecule),  $a_h$  is found from  $a_h = 0.53 \text{ \AA}$  [ $13.6 \text{ eV} \times \sigma^2/E_{ex}(1+\sigma)m_e$ ]<sup>1/2</sup>.

<sup>g</sup> $J_0 = 2.2 \text{ eV}$  (ZnO, Ref. 14). For CdS,  $J_0$  is found from  $J_0 = \frac{1}{3}(\Delta_L + 2\Delta_T)\pi a_{ex}^3/\Omega$ , where  $\Delta_L = A_L - A_F$  and  $\Delta_T = A - A_F$

from Ref. 4, and  $a_{ex}$  is the exciton radius [see also Eq. (28) in Ref. 14]. Result:  $J_0 = 2.5 \text{ eV}$  for CdS.

<sup>h</sup>Reference 17.

<sup>i</sup>I. Filinski and T. Skettrup, *Proceedings of the Ninth International Conference on the Physics of Semiconductors*, edited by S. M. Ryvkin (Nauka, Leningrad, 1968), p. 216.

<sup>j</sup>Reference 12.

<sup>k</sup>Reference 16.

exciton levels one is therefore restricted to use samples with  $N_c < 10^{18} \text{ cm}^{-3}$  so that  $\Delta_{LR} < \Delta_{free} \approx \Delta_0$ . Hence,  $\Delta_{LR}$  is usually only a minor correction to  $\Delta_0$ .

## VI. COMPARISON WITH EXPERIMENT

The properties of the three-particle centers in CdS and ZnO have been computed by means of the results obtained in the previous sections. The values are given in Table I and compared with experimental results on these materials. The parameters used in the calculations are  $m_e/m_h$ ,  $m_e$ , and  $E_{ex}$ . The values for  $m_e$  and  $m_h$  determined from experiments vary somewhat. We have chosen  $m_e/m_h = 0.20$  for<sup>4</sup> CdS and  $m_e/m_h = 0.21$  for<sup>15</sup> ZnO both of these being obtained from measurements on excitons. Thus, there are no adjustable parameters in

the calculation. The experimental energy separations which should be compared with the computed ones are those between the optically inactive states, because only these are insensitive to exchange.

The optically inactive free exciton ( $\Gamma_{6,free}$ ) has been observed for both materials. The optically inactive bound exciton ( $\Gamma_{6,bound}$ ) is found from the observed  $\Gamma_{5,bound}$  line by subtracting the experimental exchange splittings.<sup>12,16</sup> The binding energies quoted in Table I are the energy separations between the  $\Gamma_{6,free}$  and the  $\Gamma_{6,bound}$  exciton lines.

The observed values vary somewhat because the binding energies depend on the type of donor which bind the exciton.<sup>17</sup> These effects of central-cell corrections for the different kinds of donors are not included in our calculations of the binding energies.

For CdS we compute a binding energy of 6.4 meV

whereas the experimental values vary from 4.6 to 7.1 meV. For ZnO we obtain 13.4 meV and experimental numbers range from 11.3 to 20.9 meV. Thus, for both materials we obtain a good agreement, since central-cell corrections are not considered here. Also the computed exchange splitting agree well with the observed values. The oscillator strengths are predicted to be 255 for CdS and 180 for ZnO. The oscillator strengths have not yet been measured. The computed values are 10–100 times greater than those reported for excitons bound to neutral centers.<sup>6</sup> However, the fact that the three-particle centers are actually observed at helium temperature where only a few donors are ionized indicates that the strengths must be quite large.

#### VII. SUMMARY

Properties of the three-particle complex, consisting of an electron and a hole bound to an ionized donor have been computed as a function of  $m_e/m_h$ .

The ground-state energy was calculated in the effective-mass approximation by the variational method. The energy depends only on the mass ratio  $m_e/m_h$  and varies monotonically with this. The wave functions derived from the variational procedure were used for computing interparticle distances and oscillator strengths of the complex. The exchange splitting between states with parallel and antiparallel electron and hole spins was also computed, and it was found that long-range effects may be important when the concentration of centers is large. Finally the properties of the three-particle complexes in CdS and ZnO were calculated. By comparing with experimental values, good agreement was obtained.

#### ACKNOWLEDGMENT

One of the authors (M. S) would like to thank Professor N. I. Meyer for the kind hospitality extended to him at the technical University of Denmark.

<sup>1</sup>J. J. Hopfield, *Proceedings of the Seventh International Conference on the Physics of Semiconductors, Paris, 1964* (Dunod, Paris, 1964), p. 725.

<sup>2</sup>R. R. Sharma and S. Rodriguez, *Phys. Rev.* **153**, 823 (1967).

<sup>3</sup>M. Suffczynski, W. Gorzkowski, and R. Kowalczyk, *Phys. Letters* **24A**, 453 (1967); M. Suffczynski and W. Gorzkowski, *II-VI Semiconducting Compounds, Providence, 1967*, edited by D. G. Thomas (Benjamin, New York, 1967), p. 384; W. Gorzkowski and M. Suffczynski, *Phys. Letters* **29A**, 550 (1969).

<sup>4</sup>J. J. Hopfield and D. G. Thomas, *Phys. Rev.* **122**, 35 (1961).

<sup>5</sup>M. Rotenberg and J. Stein, *Phys. Rev.* **182**, 1 (1969).

<sup>6</sup>J. M. Levy-Leblond, *Phys. Rev.* **178**, 1526 (1969).

<sup>7</sup>D. C. Reynolds, C. W. Litton, T. C. Collins, and E. N. Frank, *Proceedings of the Tenth International Conference on the Physics of Semiconductors, MIT, 1970*, p. 519 (unpublished).

<sup>8</sup>C. H. Henry and K. Nassau, *Phys. Rev. B* **1**, 1628

(1970).

<sup>9</sup>M. Cardona, *J. Phys. Chem. Solids* **24**, 1543 (1963).

<sup>10</sup>A. E. Glaubermann, A. V. Pundik, S. W. Kozitskii, and M. A. Ruvinskii, *Phys. Status Solidi* **32**, 61 (1969).

<sup>11</sup>E. I. Rashba and G. E. Gurgenshvili, *Fiz. Tverd. Tela* **4**, 1029 (1962) [*Sov. Phys. Solid State* **4**, 759 (1962)].

<sup>12</sup>D. G. Thomas and J. J. Hopfield, *Phys. Rev.* **128**, 2135 (1962).

<sup>13</sup>C. H. Henry, R. A. Faulkner, and K. Nassau, *Phys. Rev.* **183**, 798 (1969).

<sup>14</sup>T. Skettrup and I. Balslev, *Phys. Status Solidi* **40**, 93 (1970).

<sup>15</sup>R. E. Dietz, J. J. Hopfield, and D. G. Thomas, *J. Appl. Phys. Suppl.* **32**, 2282 (1961).

<sup>16</sup>D. C. Reynolds, C. W. Litton, and T. C. Collins, *Phys. Rev.* **140**, A1726 (1965).

<sup>17</sup>K. Nassau, C. H. Henry, and J. W. Shiever, *Proceedings of the Tenth International Conference on the Physics of Semiconductors, MIT, 1970*, p. 639 (unpublished).