Erratum: Structure of amorphous selenium studied by neutron diffraction

Hansen, Flemming Yssing; Knudsen, Torben Steen; Carneiro, K.

Published in:
Journal of Chemical Physics

Link to article, DOI:
[10.1063/1.431721](https://doi.org/10.1063/1.431721)

Publication date:
1975

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Hansen, F. Y., Knudsen, T. S., & Carneiro, K. (1975). Erratum: Structure of amorphous selenium studied by neutron diffraction. *Journal of Chemical Physics*, 63(7), 3182-3182. <https://doi.org/10.1063/1.431721>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

i. e., the $u \rightarrow 0$ approximation. It can be readily shown that with these rate constants the Master equation has an analytic solution. The population of the J level at time t , $N_J(t)$ is then of the form $N_J(t) = N_J(\infty) + [N_J(0) - N_J(\infty)] \exp(-\alpha t) = N_J(\infty)[1 - \exp(-\alpha t)] + N_J(0) \exp(-\alpha t)$. If $N_J(\infty) < N_J(0)$, $N_J(t)$ decays exponentially to its equilibrium value; if $N_J(\infty) > N_J(0)$, $N_J(t)$ rises exponentially to its equilibrium value; while if $N_J(\infty) = N_J(0)$, $N_J(t)$ stays constant. These features are evident in Fig. 1 and in the numerical simulations carried out by other workers.¹⁻³ Had we employed the $u \rightarrow 0$ approximation, the only changes in Fig. 1 would be an exact exponential relaxation of the moments and the absence of the weak "overshoot" in $N_J(t)$ for $J=6$ (and 7). For lower values of u (or for a less skewed initial distribution than that employed in Fig. 1) the results would be even better. The only realistic exception²¹ is H_2 .

The intermediate region, $u \sim 1$, is under active study. The preliminary evidence is that the maximal entropy procedure¹⁰ which is valid at both extremes will cover this region as well.

*Work supported by the Israel Commission for Basic Research, the Israel Academy of Sciences and Humanities.

¹J. C. Polanyi and K. B. Woodall, *J. Chem. Phys.* **57**, 1574 (1972).

²R. D. Levine and R. B. Bernstein, *Molecular Reaction Dynamics* (Clarendon, Oxford, 1974).

³H. W. Chang and D. W. Setser, *J. Chem. Phys.* **58**, 2298 (1973).

⁴W. H. Duerwer and D. W. Setser, *J. Chem. Phys.* **58**, 2310

(1973).

⁵R. B. Bernstein, *J. Chem. Phys.* **62**, 4570 (1975).

⁶I. Oppenheim, K. E. Shuler, and G. H. Weiss, *Adv. Mol. Relax. Process.* **1**, 13 (1967).

⁷J. I. Steinfeld, I. Burak, D. G. Sutton, and A. V. Nowak, *J. Chem. Phys.* **52**, 5421 (1970).

⁸W. D. Breshears, *Chem. Phys. Lett.* **20**, 429 (1973).

⁹R. M. Osgood, Jr., P. B. Sackett, and A. Javan, *J. Chem. Phys.* **60**, 1464 (1974).

¹⁰I. Procaccia and R. D. Levine, *J. Chem. Phys.* **62**, 3819 (1975).

¹¹R. B. Bernstein and R. D. Levine, *J. Chem. Phys.* **57**, 434 (1972).

¹²R. D. Levine and R. B. Bernstein, *Chem. Soc. Faraday Discuss.* **55**, 100 (1973).

¹³R. D. Levine and R. B. Bernstein, *Acc. Chem. Res.* **7**, 393 (1974).

¹⁴C. Rebick, R. D. Levine, and R. B. Bernstein, *J. Chem. Phys.* **60**, 4977 (1974).

¹⁵I. Procaccia and R. D. Levine (accepted for publication in *J. Chem. Phys.*).

¹⁶M. Robinson and J. I. Steinfeld, *Chem. Phys.* **4**, 467 (1974).

¹⁷I. Procaccia and R. D. Levine, *J. Chem. Phys.* **62**, 2496 (1975).

¹⁸A. M. G. Ding and J. C. Polanyi, *Chem. Phys.* (to be published).

¹⁹The validity of this approximation can be judged not only in terms of the predicted bulk behavior, Fig. 1, but also on the basis of individual rate constants. As long as $(E_J + E_{J'})/2|E_J - E_{J'}| = O(1)$, it is valid in the $u \rightarrow 0$ limit.

²⁰K. G. Anlauf, D. S. Horne, R. G. Macdonald, J. C. Polanyi, and K. B. Woodall, *J. Chem. Phys.* **57**, 1561 (1972).

²¹When we used artificially high values of B_e (or low values of T) so that $u > 1$, the populations determined by the numerical solution of the master equation show the typical $V-T$ behavior, where the relaxation time decreases as E_J increases.

ERRATA

Erratum: Structure of amorphous selenium studied by neutron diffraction [*J. Chem. Phys.* **62**, 1556 (1975)]

F. Yssing Hansen and T. Steen Knudsen

Fysisk-Kemisk Institut, The Technical University of Denmark, DK 2800 Lyngby, Denmark

K. Carneiro

Danish Atomic Energy Research Establishment, Risø, DK 4000 Roskilde, Denmark

Since the publication of this article it has been brought to the authors' attention that the method developed to extrapolate the structure factor, measured by neutron diffraction, is very similar to the procedure described previously by J. H. Konnert and J. Karle in *Acta Cryst. A* **29**, 702 (1973). We regret not

having made reference to their work in the original manuscript.

Further, if we had known about this work, we should probably not, in the abstract of our paper, have referred to our method as "new."