



Biomarkers in skin autoimmunity—An update on localised scleroderma

Hernández-Bustos, Adrián; Bolos, Begona; Astakhova, Kira

Published in:
Skin Health and Disease

Link to article, DOI:
[10.1002/ski2.335](https://doi.org/10.1002/ski2.335)

Publication date:
2024

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Hernández-Bustos, A., Bolos, B., & Astakhova, K. (2024). Biomarkers in skin autoimmunity—An update on localised scleroderma. *Skin Health and Disease*, 4(2), Article e335. <https://doi.org/10.1002/ski2.335>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

REVIEW ARTICLE

Biomarkers in skin autoimmunity—An update on localised scleroderma

Adrián Hernández-Bustos | Begona Bolos | Kira Astakhova 

Department of Chemistry, Technical University of Denmark, Kongens Lyngby, Denmark

CorrespondenceKira Astakhova.
Email: kiraas@kemi.dtu.dk**Funding information**

Danmarks Tekniske University, Grant/Award Number: Alliance 102140

Abstract

Human autoimmune diseases are complex and highly diverse conditions that can be of localised or systemic nature. Understanding the basic biology of autoimmune diseases goes hand in hand with providing the clinics with valuable biomarkers for managing these diseases. The focus of this review is paid to localised scleroderma, an autoimmune disease affecting skin and subcutaneous tissue. Localised scleroderma has very few serological biomarkers for clinical analyses distinguishing it from main differentials, and yet noneffective prognostic biomarkers. With this regard, the review covers well-established and new biomarkers such as cell surface proteins, auto-antibodies and cytokines. In recent few years, several new biomarkers have been suggested, many provided with modern genomic studies. This includes epigenetic regulation of DNA, RNA transcriptomics and regulatory RNA such as microRNA and long non-coding RNA. These findings can for the first time shed light on the genetic mechanisms behind the disease and contribute to improved diagnosis and treatment.

1 | INTRODUCTION

Mechanisms of autoimmune diseases are still not fully understood, inspiring the research community to study them with continuously advanced experimental and clinical approaches. Thanks to these efforts it is now confirmed that multiple autoimmune diseases are caused by a prolonged or chronic inflammatory response.^{1,2} These responses have origin in genetic predisposition and epigenetic factors.¹ Such precise biomarkers could be used for tracking diseases and potentially, guide treatment. Besides diagnostic and treatment follow up purposes, monitoring common biomarkers could reveal biological details on different autoimmune diseases and their interconnection.

Localized scleroderma (LS), or morphea, is an autoimmune disease typically leading to skin inflammation

and thickness distributed in various parts of the body, as well as affecting the extracutaneous tissues. Localised scleroderma can be classified based on the extent and degree of fibrosis into different groups. Traditionally, it has been classified into subtypes plaque, generalised, bullous, linear, and deep.³ However, according to the Padua consensus, it can be divided in the following groups: circumscribed (including subtypes superficial and deep), linear (including the subtypes trunk and head), generalised, panclerotic and mixed,⁴ being this the most accepted classification.⁵ While its pathogenesis is not fully known, factors such as genetic predisposition and epigenetics,^{6–8} vascular dysregulation or Th1/Th2/Th17 imbalance through the different stages⁹ lead to the activation of inflammatory and profibrotic pathways that ends into an upregulation on collagen deposition.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Authors. Skin Health and Disease published by John Wiley & Sons Ltd on behalf of British Association of Dermatologists.

With a reported incidence of 3.4–27 per 100,000,¹⁰ or 4–27 new cases per million children per year reported in,⁹ LS affects mainly women and one third of the patients are paediatric.¹⁰ The treatment is mainly based on methotrexate with efficacy 67% in paediatric patients.^{11–13} Corticosteroids are applied as well. However, most patients relapse after corticosteroids' discontinuation.¹⁴

Localised scleroderma and systemic sclerosis (Ssc) share skin histopathologic changes, reason why sometimes certain subtypes of LS as pansclerotic LS can be confounded with SSc.¹⁵ However, these two diseases differ in the distribution and pattern of skin involvement. The associated extracutaneous and internal organ manifestations are mild in LS,¹⁶ and are more severe and common in SSc.¹⁷ Remarkably, LS cannot evolve into SSc. Immunogenetically, human leukocytes antigens (HLA) type II genes are upregulated in LS, being type II specific upregulation, while in SSc it is type I upregulation that takes place.¹⁸ As it will be further discussed, both diseases share the presence of certain pathogenic autoantibodies, nevertheless, are genetically different.

Protein biomarkers for LS have been far less successful compared to other autoimmune diseases such as lupus or rheumatoid arthritis (RA). Since there is no serological marker specific for the disease or characteristic parameters in LS patients,¹⁹ diagnostics are based on clinical observations and, in cases where the LS type is not evident or the inflammatory state has to be defined, skin biopsies are used.^{20,21} Both serological and tissue samples are used to monitor LS. Other methods are based on imaging, using techniques as ultrasonography,^{22,23} infrared thermography,^{13,24} and optical coherence tomography.²⁵ Nevertheless, misdiagnose²⁶ and risk of extracutaneous involvement which commonly associates with morbidity in LS,²⁷ require fast and robust methods to diagnose and monitor LS and its subtypes.

As to the genetic signature of LS, specific HLA class I and class II alleles are associated with generalised and linear subtypes. Interestingly, LS is immunogenetically distinct from other scleroderma types.⁶ However, risk alleles in morphea are also associated with conditions such as RA and other autoimmune conditions.⁶ The role of HLA products in regulating interactions of immune cells is well known,²⁸ and therefore, the specific HLA profile of morphea could lead to B cells producing certain cytokines and autoantibodies contributing to disease progression.^{6,29}

The main objective of this review is to give an update on relevant biomarkers for diagnosing and studying LS. In addition to previously reviewed biomarkers,³⁰ we also include genomic studies, cell surface and endothelial biomarkers, and give a brief overview of upcoming clinical trials that aim at genetic studies of LS.

What is already known?

- Localised scleroderma (LS) is an autoimmune disease affecting skin and subcutaneous tissue
- LS has very few serological biomarkers for clinical analyses distinguishing it from main differentials, and yet noneffective prognostic biomarkers.

What does this study add?

- This review covers well-established and new biomarkers such as cell surface proteins, autoantibodies and cytokines
- New promising biomarkers include epigenetic markers, regulatory RNA and regulatory proteins

2 | BIOMARKERS IN LOCALISED SCLERODERMA

In the chapter below we sum up recent advances in biomarkers in scleroderma and classify them according to clinical relevance as follows: biomarkers of disease activity, biomarkers that enable distinguishing LS from the main differentials, and prognostic indicators including those that can monitor response to treatment. Both clinically established biomarkers and those in the research stage are included, to show the ongoing development in the field and to make the clinicians aware of the upcoming new tests.

2.1 | Biomarkers of disease activity and extracutaneous involvement

Cell surface and endothelial biomarkers can be evaluated using skin biopsy and reveal disease activity and complications (Table 1). Cluster of differentiation CD34 is a biomarker used to identify certain cell populations, particularly hematopoietic stem and progenitor cells, endothelial progenitor cells, and various types of mesenchymal cells. CD34 is a cell surface glycoprotein expressed on endothelial cells, including those present in blood vessels. Studies have shown decreased expression of CD34 on endothelial cells in LS lesions.^{31,32}

Factor XIIIa (FXIIIa) is a biomarker that is commonly used to identify and characterise certain cells within the skin, particularly dendritic cells known as dermal dendrocytes or FXIIIa + dendritic cells. These cells are a type of antigen-presenting cell found in the skin and play a role in wound healing, tissue repair, and immune

TABLE 1 Overview of biomarkers used in Localised scleroderma (LS) diagnosis and studies for assessing LS activity and complications (2000–2022).^a

Biomarker	Tissue (T) or serum (S) sample	Activity (A) and/or complications (C)	Cohort size (year)	Biological association	Details	(Refs.)
CD34	T	A	20 keloid, 20 lacerations, 20 LS patients (2022)	Cell surface glycoprotein expressed on endothelial cells.	Decreased expression of CD34 on endothelial cells in localised scleroderma lesions. CD34 can also be used to differentiate from other fibrotic diseases or to follow treatment response.	31,32
FXIIIa	T	A	30 LS patients (2013)	Transglutaminase related to blood clots stabilisation by fibrin cross-linking.	FXIIIa + dendritic cells can provide insights into the immune and inflammatory processes occurring in the affected skin.	32
CD1a	T	A	6 LS patients, 3 HC (2008)	Cell surface protein in immune Langerhans dendritic cells. These are related to immune responses and antigen presentation.	They can be used to identify and study Langerhans cells in the skin and are associated with disease onset and progression.	33
CD86	T	A	6 LS patients, 3 HC (2008)	Cell surface protein primarily associated with antigen-presenting cells. It plays a role on immune responses, T-cell activation and regulation.	Increased expression of CD86 can indicate immune activation and inflammation in the affected skin of individuals with LS.	34
sCD30	S	A	55 LS patients, 15 SSc patients, 20 HC (2000)	Circulating form of CD30, a cell surface protein primarily expressed on activated T cells.	Potential biomarker to assess immune system activity and inflammation, as well as disease progression or response to treatment.	35
RF	S	A/C	43 LS patients, 14 HC (2005)	RFs are antibodies that target the Fc portion of IgGs as immune response regulators.	RF was present in 30% of LS patients. Patients with generalised morphea can be differentiated by the dysregulated isotype of RF.	36
ANAs	S/T	A/C	750 LS patients ^a (2005)	In LS, autoreactive T and B cells produce pathogenic antibodies such ANAs, AHAs or anti-ssDNA autoantibodies that attack healthy tissues.	Present in patients with autoimmune connective tissue disease. ANAs were found in the serum of 36%–57.9% of LS patients (depends on the cohort). It is associated to risk of relapse and extracutaneous involvement.	37–42
anti-ssDNA autoantibodies	S	A/C	69 LS patients		AHAs were found in 32%–39% of the LS patients, and anti-ssDNA in 29%–30% of the LS patients, correlating to disease severity.	43
AHAs	S	A/C	71 HC (2008)		Both are commonly proposed as markers for higher risk of muscle and joint morbidity	

(Continues)

TABLE 1 (Continued)

Biomarker	Tissue (T) or serum (S) sample	Activity (A) and/or complications (C) association	Cohort size (year)	Biological association	Details	(Refs.)
Vimentin p16	S/T	A	20 keloid, 20 lacerations, 20 LS patients (2022)	Vimentin has been suggested to be the main myofibroblast general marker of the fibrotic process. p16 is a tumour suppressor protein.	Vimentin, p16 and FOXP3 were overexpressed in the affected skin for all three fibrotic pathologies.	31
FOXP3				FOXP3 suppresses expression of many genes including IL-2 and effector T-cell cytokines.		
IL-2	S	A	73 LS patients	Dysregulated production of cytokines has a critical role in autoimmune diseases and LS.	All the cytokines were increased and related to disease activity.	44
IL-2R	S		26 HC (2016)			
IL-12	S					
CXCL9	S					
CXCL10	S					
CCL18	S	A	74 LS patients		CCL18 allowed to differentiate between active and inactive disease.	45
sVCAM	S	A	22 HC (2019)	sVCAM is related to the transport of the leukocytes to the area where inflammation takes place.	These markers were found upregulated in the serum of LS patients	
Gal-9	S	A		Gal-9 is a β -galactoside binding lectin known for its immunomodulatory role.		
TIE-1	S	A		TIE-1 has a role in the regulation of lymphatic and cardiovascular development.		
SPARC	S	A	15 LS patients	Fibrosis mediator present only at sites of tissue remodelling and wound repair for different diseases.	SPARC was upregulated in patients with all types of sclerosis, especially elevated in those with LS.	46
			78 Sclerotic patients			
			15 HC (2017)			
MBP antibodies	S	A	70 LS patients	In LS, autoreactive T and B cells produce pathogenic antibodies that attack healthy tissues.	MBP was found in 71.4% of a cohort of 50 patients, differing significantly from HCs, SSc patients and correlating to higher disease activity.	47,48
			30 SSc patients			
			35 HC (2022)			

^aAHA, Antihistone antibodies; ANA, antinuclear antibodies; anti-ssDNA, anti-single strand DNA; CD, Cluster Differentiation; FOXP3, forkhead box P3; FXIIa, Coagulation factor XIIa; Gal-9, Galectin-9; HC, healthy control; IL, interleukin; LS, localized scleroderma; MBP, Myelin Basic Protein; RF, Rheumatoid Factor; sCD, soluble Cluster Differentiation; SPARC, Secreted Protein Acidic and Rich in Cysteine; SSc, Systemic Sclerosis; sVCAM, soluble vascular cell adhesion molecule; TIE-1, Tyrosine-protein kinase receptor.

regulation. In the context of LS, FXIIIa + dendritic cells can provide insights into the immune and inflammatory processes occurring in the affected skin.³²

CD1a is a cell surface protein that serves as a biomarker for a specific type of immune cell known as Langerhans cells. Langerhans cells are dendritic cells present in the skin and mucous membranes, and they play a significant role in immune responses and antigen presentation. In LS, CD1a is used as a biomarker to identify and study Langerhans cells in the skin associated with disease onset and progression.³³

CD86, also known as B7-2, is a cell surface protein primarily associated with antigen-presenting cells, particularly dendritic cells, macrophages, and B cells. CD86 plays a critical role in immune responses, including T-cell activation and regulation. In the context of LS, CD86 is relevant for monitoring the immune response and inflammatory processes. Increased expression of CD86 can indicate immune activation and inflammation in the affected skin of individuals with LS.³⁴

Soluble CD30 (sCD30) is a circulating form of CD30, a cell surface protein and member of the tumour necrosis factor receptor superfamily. CD30 is primarily expressed on activated T cells, including Th2 cells and T-regulatory cells, and is involved in immune responses. In the context of LS, serum levels of sCD30 has been investigated as a potential biomarker to assess immune system activity and inflammation.³⁵ sCD30 is associated with Th2-type immune responses, which are involved in allergic and inflammatory conditions. Changes in sCD30 levels could be indicative of disease progression or response to treatment.

All these biomarkers are studied in skin biopsies with immunohistochemistry methods. Alternatively, the cell types can be detected in blood with cell sorting techniques.

As to serological tests, they are not common for the diagnosis of LS, even not recommended in guidelines.^{20,49} However, serological biomarkers such as aldolase, creatinine phosphokinase, lactate dehydrogenase, C-reactive protein and rheumatoid factor (RF) are commonly measured (Table 1) and associate with the LS activity.^{20,37,50} RF was present in 30% of a cohort of 43 LS patients and 14 controls, especially for patients with generalised morphea which can be differentiated by the dysregulated isotype of RF.³⁶

The opinion on relevance of ANAs, extractable nuclear antigen antibodies and antibodies to single stranded DNA (anti-ssDNA) is conflicted since a big percentage of LS patients (50% or more, depending on the cohort and the subtype of LS) is negative on them.^{20,49} In different cohorts, the biggest of them being 671 LS patients,³⁷ ANAs were only found in the serum of 36%–57.9% of them.^{37–42} In addition, antinuclear antibodies (ANA) positivity combined with older LS onset age is thought to be a potential marker for risk of

relapse.⁴¹ Besides, ANA positivity has been associated with extracutaneous involvement.¹⁶ This is the case in the study by Li et al. who found extracutaneous involvement associated with more medication use, longer treatment durations, and greater disease burden.¹⁶

Anti-histone antibodies (AHAs) were found in 32%–39% of the LS patients, and anti-ssDNA in 29%–30% of the LS patients.^{42,43} A cohort of 187 LS patients found AHAs and anti-ssDNA antibodies in only 12% and 8% of the patients, respectively.⁵¹ Despite the dependence of the autoantibodies levels on the cohort and the low percentage of patients that are positive on them, some authors propose the use of them combined as a marker for higher risk of muscle and joint morbidity.⁵²

A recent study showed that myelin basic protein (MBP) antibodies measured in serum of the 27.3% of a cohort of 139 LS patients correlate to higher disease activity.⁴⁷ Smaller cohort of 50 LS patients found these antibodies in 71.4% of the patients.⁴⁸

Along with autoantibodies, cytokines have a unique signature in LS.^{8,42} Thus, serum levels of a cohort consisting of 73 LS patients and 26 healthy controls proved that TH1-related interleukins IL-2, IL-2R and IL-12, as well as chemokines CXCL9 and CXCL10 were increased and related to disease activity.⁴⁴ Similar sized cohorts reported upregulated levels of chemokines CXCL9, CXCL10 and CCL18.^{42,45} Serum levels of CCL18 chemokine were the most useful to differentiate from active and inactive disease, and its respective gene expression was increased at the inflammatory borders of the LS lesions, being overall a good marker to monitor the disease.⁴⁵ Interestingly, lectin Gal-9, tyrosine-protein kinase receptor, soluble vascular cell adhesion molecule was also found upregulated in serum samples from the same cohort, being also potential markers.

miRNAs serve as essential regulators of cell differentiation, proliferation and survival. The involvement of miRNAs in the functioning and regulation of the skin cells and skin diseases is a rapidly advancing area in dermatological research.⁵³ miRNAs have been identified to play a key role in the pathogenesis, diagnosis, and treatment of the skin diseases. It remains unknown how general miRNA regulators are, and if there are specific regulators for LS. Considering that miRNAs regulate specific pathways, specificity of miRNA to LS would be a result of having identified such a specific regulatory pathway.

To date, miRNAs have been identified to demonstrate significant effects in diverse skin inflammatory conditions such as wounds, cancer, psoriasis, scleroderma, dermatomyositis, for example, reviewed in Singhvi et al.⁵³ miRNA-29, miRNA-21 and miRNA-483-5p were investigated in several skin conditions, mainly in SS.^{54–57} These miRNAs were also found to be

upregulated in LS study. miRNA-7 and miRNA-196, both related to collagen expression, have also been studied and found downregulated in LS patients, being a potential marker for LS.^{7,58} Using 38 samples of LS patients and matched controls, it was shown that the serum levels of multiple miRNAs, that is, miRNA-181b-5p, miRNA-223-3p, miRNA-21-5p, let 7i-5p, miRNA-29a-3p and miRNA-210-3p were significantly increased in the LS patients compared to the healthy control (50). The level of let-7i in the female LS patients correlated negatively with disease activity scores body surface area and modified Localised Skin Severity Index (mLoSSI). Moreover, the female patients with inactive LS had significantly higher level of let-7i in comparison to those with active disease. The exact role of those miRNA molecules has not been revealed in LS and long-term longitudinal research is pivotal to confirm their prognostic value.

2.2 | Biomarkers enabling distinguishing localised scleroderma from the main differentials

It is important to differentiate LS from other conditions with similar skin manifestations to ensure accurate diagnosis and appropriate treatment. Some key skin conditions for differentiation from LS include SSc, Lichen Sclerosus, Eosinophilic Fasciitis, Morpheaform Basal Cell Carcinoma, Scleromyxedema and Localised Sclerodermaid Chronic Graft-versus-Host Disease (cGVHD).⁵⁹ Today, accurate diagnosis involves a thorough clinical evaluation, medical history review, skin biopsy, histopathological examination, and may sometimes include additional diagnostic tests or imaging studies.

The main differential for LS is with no doubt SSc. A recent study found MBP antibodies in the serum of 71.4% of a cohort of 70 LS patients, differing significantly from healthy controls and SSc patients, and showing relation to pain symptoms and higher disease activity.⁴⁸

Attachment of carbohydrate is among the most common post-translational modifications of proteins. For immunoglobulins, it affects recognition of antigens and interaction with immune cells. The recent study recruited 93 LS patients, 298 SSc patients, and 436 healthy controls, to conduct immunoglobulin proteomics assessment.⁶⁰ N-glycans of purified immunoglobulin G were obtained from plasma and detected by tandem mass spectrometry. The authors examined whether the IgG-Galactose (Gal) ratio differed between different subtypes of scleroderma. The IgG-Gal ratio was significantly higher in SSc patients (1.139 ± 0.870) than in LS patients (0.485 ± 0.280) and controls (0.395 ± 0.190). The IgG-Gal ratio successfully distinguished SSc patients from LS and controls (area under the curve = 0.88 and 0.81, respectively). IgG-Gal ratios

were abnormal in SSc patients and were associated with disease severity. The IgG-Gal ratio therefore shows potential as a biomarker for the diagnosis of SSc and the differentiation from LS.

Fibrosis is a common pathophysiological response of many tissues and organs subjected to chronic injury. Despite the diverse aetiology of keloid, lacaziosis and LS, the process of fibrosis is present in the pathogenesis of all these three entities beyond other individual clinical and histological distinct characteristics. Tafuri et al. report on fibrosis immunohistochemistry study in 20 skin paraffinized samples each of these three chronic cutaneous inflammatory diseases.³¹ The presence of α -smooth muscle actin (α -SMA) and vimentin cytoskeleton antigens, CD31, CD34, Ki67, p16; CD105, CD163, CD206 and FOXP3 antigens; and the central fibrotic cytokine Transforming growth factor- β (TGF- β) was determined by immunohistochemistry. Vimentin was overexpressed in comparison to α -SMA for all three pathologies. CD31 and CD34-positive blood vessel endothelial cells were present throughout the reticular dermis. Just in LS, Ki67 expression was almost absent, while. P16-positive levels were higher and observed in reticular dermis of keloidal collagen in keloids, in collagen bundles in scleroderma and in the external layers of the granulomas in lacaziosis. α -actin positive cells and rarely CD34 positive cells was found mainly in keloids, possibly related to higher p16 antigen expression, which accounts for cell senescence. CD105-positive cells were found in perivascular tissue in close contact with the adventitia in keloids and scleroderma, while, in lacaziosis, these cells were mostly observed in conjunction with collagen deposition in the external granuloma layer. Low FOXP3 expression was observed in all lesion types. Transforming growth factor- β was exclusive to keloid and lacaziosis lesions. For all the proposed lesion types, vimentin has been suggested to be the main myofibroblast general marker of the fibrotic process, while endothelial-to-mesenchymal transition (EndoMT), mesenchymal stem cells and M2 macrophages may not play a role. Similar to the case described by Tafuri et al., the also fibrosis mediator Secreted Protein Acidic and Rich in Cysteine (SPARC) is present only at sites of tissue remodelling and wound repair for different diseases. Tsuruta et al. observed that SPARC was upregulated in serum of patients with sclerosis of a small cohort of 15 patients with LS, 78 patients with other kinds of sclerosis and 15 healthy controls.⁴⁶ Even though serum levels of SPARC were upregulated in patients with all types of sclerosis, it was especially elevated in those with LS being this protein a promising and selective candidate for LS diagnostics.

Genetic aspects of LS can aid its effective differentiation as well. Mirizio et al. reported results from RNA sequencing (RNAseq) of skin biopsies from

juvenile-onset LS.⁶¹ LS gene signatures compared to healthy controls showed a distinct expression of an inflammatory response gene signature (IRGS) composed of interferon genes $IFN\gamma$ -, $IFN\alpha$ -, and $TNF\alpha$ -associated genes. Gene enrichment analysis showed that the IRGS, including interferon-inducible chemokines such as CXCL9, CXCL10, CXCL11, and $IFN\gamma$ itself, was more highly expressed in LS patients with more inflammatory lesions. The prevalence of the $IFN\gamma$ signature in the lesion biopsies of active LS patients indicates that these genes reflect clinical activity parameters and may be the promoters of early inflammatory disease. Importantly for sequencing experiment strategies and sampling, the use of paraffinized skin for sequencing was proven to be an effective substitute for fresh skin by comparing gene expression profiles.

Besides genomic markers, small regulatory RNA are exciting objects of study. Among others, microRNA are powerful short RNA molecules that are responsible for regulation of gene replication, transcription, and translation. Over decades, microRNA detection has been challenged by the fact that they are short living and present in the sample only at low (<1 fM) concentrations. As a result, before 2020, one of the few nucleic acid biomarkers described in LS was let-7a, related to several important cell pathways as DNA damage, Janus kinase/signal transducer and activator of transcription proteins (JAK/STAT) pathway, cell cycle or apoptosis. The expression of let-7a in dermal fibroblasts in a small cohort of 7 SSc patients, 7 LS patients and 7 healthy controls showed that the levels of the miRNA were under-regulated in LS patients, being a potential tool for differentiate LS from SSc patients.⁶²

2.3 | Prognostic biomarkers including biomarkers indicating response to treatment

When the diagnosis LS is established and the treatment has been initiated, prognostic biomarkers can offer precise follow up and personalised management plan. Research on specific biomarkers for monitoring treatment response in LS is still evolving, and there is no universally accepted biomarker for this purpose. Clinicians primarily assess treatment response through regular clinical examinations, monitoring changes in skin thickness, texture, and overall disease activity. They may use validated assessment tools such as the Localised Scleroderma Skin Severity Index to quantify skin involvement and monitor response to treatment.

Skin biopsies may be performed before and after treatment to evaluate histological changes in the skin, such as collagen deposition and inflammation, which can indicate treatment response. Among others, sCD30 has been reported as relevant to monitor treatment outcome.⁶³

Fibroblasts are the primary cell type involved in the excessive production of collagen, leading to skin fibrosis in LS. Common markers for fibroblasts include vimentin³¹, fibroblast-specific protein 1,⁶⁴ and platelet-derived growth factor receptor beta⁶⁵; all can be assessed using sera samples from patients and predict LS progression.

General blood biomarkers associated with inflammation, immune activation, and fibrosis (e.g., cytokines, growth factors, autoantibodies) can help assessing the disease activity and response to treatment in LS. Moreover, imaging with high-frequency ultrasound or magnetic resonance imaging can provide information on skin changes in LS. In a clinical trial completed in 2013, the CD34 and FXIIIa were used to monitor outcome of skin treatment with flashlamp pulsed dye laser.³² Thirty patients with plaque morphea were treated with the laser. Sessions were performed biweekly for 6 months and led to improvement in skin condition, also upon follow-up 12 months after the last laser treatment. An increased number of CD34-positive cells were found in both the upper and the lower dermis, accompanied by reduced FXIIIa-positive cells in the latter. Therefore, using the laser treatment with complementary use of CD34 and FXIIIa tests could be a potent new management plan for LS.

Several other cytokines have recently attracted attention in autoimmunity research and clinics. Interleukin-6 (IL-6) is a pro-inflammatory cytokine that plays a significant role in immune regulation and inflammation. It is involved in various autoimmune and inflammatory conditions, including LS. Soluble IL-6 (sIL-6) refers to the form of IL-6 that is found in the bloodstream and is measurable through blood tests. In LS, serum sIL-6 levels may be elevated, reflecting an active inflammatory process. However, research regarding sIL-6 specifically in LS is still evolving, and there may not yet be a standardized or widely accepted reference range for sIL-6 levels in this condition. Recent mechanistic studies supported a model whereby IL-6 trans-signalling driven by CD4 T cell-derived soluble IL-6 receptor complexed with fibroblast-derived IL-6 promoted excess extracellular matrix gene expression.⁶⁶ MISTRG6 mice transplanted with scleroderma skin demonstrated multiple fibrotic responses centred around human IL-6 signalling, which was improved by the presence of healthy bone marrow-derived immune cells. These results highlight the importance of IL-6 trans-signalling in pathogenesis of scleroderma and the ability of healthy bone marrow-derived immune cells to mitigate disease.⁶⁶ Hence, monitoring serum sIL-6 levels may have diagnostic and prognostic implications in LS, helping to assess disease activity and predict disease progression. Targeting IL-6 or its receptor with specific therapies (e.g., tocilizumab) is being explored in various autoimmune conditions, and it may have potential applications in LS as well.

Similarly, the specific role and levels of soluble IL-2 receptor (sIL-2) in LS have not been extensively studied, and there may not yet be established reference ranges or widely accepted biomarker status for sIL-2 in this condition. Nevertheless, IL-2 is a critical cytokine involved in the activation and proliferation of T cells. Elevated levels of sIL-2 could suggest increased T-cell activity in LS, potentially contributing to the inflammatory response.⁶⁷ In a 2018 study, the connection between soluble endothelial leukocyte adhesion molecule-1 (sE-selectin) and sIL-2R and the severity of skin lesions in various subtypes of LS was studied. Evaluation of disease severity, the location of skin lesions, the duration of symptoms and disease activity were assessed in relation to different LS types (generalised, plaque and linear) in patients with LS. The study included 42 patients with LS and 41 healthy subjects. Significantly higher serum levels of sE-selectin and sIL-2 were observed in the LS study group when compared with the control group ($p < 0.001$). The highest concentrations of sE-selectin and sIL-2 were observed in patients with the generalised subtype of LS. A positive, statistically significant, curvilinear relationship was shown amid the mLoSSI and sE-selectin and sIL-2 concentrations in the LS study group.⁶⁷

Tumour necrosis factor- α (TNF- α) is a pro-inflammatory cytokine that plays a crucial role in various immune and inflammatory responses. It is produced by a variety of immune cells and is known to be involved in the pathogenesis of several autoimmune and inflammatory conditions. In the context of LS (morphea), research has indicated the potential involvement of TNF- α in the disease process.⁴² Elevated levels of TNF- α in LS may indicate an active inflammatory process within the affected skin. Tumour necrosis factor- α can stimulate the production of collagen and other extracellular matrix components. In LS, increased TNF- α levels may contribute to the excessive collagen deposition and fibrosis seen in affected skin. Given its role in inflammation and fibrosis, TNF- α has been explored as a potential target for therapeutic interventions in LS. Tumour necrosis factor- α inhibitors (e.g., infliximab, etanercept) have been studied in some cases to assess their effectiveness in managing LS.^{68,69}

Together, recent data on cytokines sIL-6, sIL-2 and TNF- α aligns with previous recognition of pro-inflammatory state in LS. It is though exciting new aspect that these cytokines can become treatment targets and prognostic markers in LS.^{42,66–69}

Overall, assessment of treatment response in LS is often multifaceted, involving a combination of clinical, imaging, histopathological, and patient-reported measures. Future research is expected to shed light on specific biomarkers that can reliably predict and monitor

treatment response, leading to more personalised and effective treatment strategies for individuals with LS.

Genomic biomarkers are a large field that also holds a potential to provide with prognostic biomarkers for LS. Recently, genetic signature in skin biopsies from LS patients has been analysed with next-generation sequencing technologies. Saracino et al. performed a study on 16 LS patients with epidermal whole genome sequencing protocol.⁷⁰ No single affected gene or single nucleotide variant has been found. However, many potential disease-relevant pathogenic variants were present, including ADAMTSL1 and ADAMTS16. A highly proliferative, inflammatory and profibrotic epidermis profile was seen, with significantly overexpressed TNF α -via-NF κ B, TGF β , IL6/JAKSTAT and IFN-signalling, apoptosis, p53 and Kirsten rat sarcoma virus (KRAS)-responses. This is a highly diverse group of genes and pathways. Noteworthy, affected KRAS might link LS genetics to skin cancer.⁷¹ The authors also highlight that upregulated Interferon Alpha Inducible Protein 27 and downregulated (Laminin Subunit Alpha 4) LAMA4 potentially represent initiating epidermal 'damage' signals and enhanced epidermal-dermal communication. Localised scleroderma dermis exhibited significant profibrotic, B-cell and IFN-signatures, and upregulated morphogenic patterning pathways such as Wnt (being this one used to induce fibrosis in scleroderma models in mice⁷²). Overall, the study supports the absence of somatic epidermal mosaicism in LS, and identifies potential disease-driving epidermal mechanisms, epidermal-dermal interactions, and disease-specific dermal differential-gene-expression in LS.

Next, Schutt et al. analysed skin transcriptome in skin biopsy tissues from children with juvenile LS compared to paediatric healthy controls.¹⁸ In this study, differentially expressed genes (DEGs) were assessed for correlations with histopathologic and clinical features in children with juvenile LS and were used to group the children into distinct genetic clusters based on immunophenotype. RNA-Seq was performed on sections of paraffin-embedded skin tissue obtained from 28 children with juvenile LS and 10 paediatric healthy controls. As a result, 589 significant DEGs were identified in children with juvenile LS as compared to healthy controls. Hierarchical clustering was used to demonstrate 3 distinct juvenile LS immunophenotype clusters (Figure 1a). In one cluster, inflammation-related pathways were up-regulated, corresponding to the histologic skin inflammation score. In the second cluster, fibrosis-related pathways were up-regulated. In the third cluster, gene expression in the skin corresponded to the patterns seen in healthy controls. Up-regulation of HLA class II genes was observed within the first cluster (characterised by predominant inflammation) (Figure 1b), a feature that has also been observed in the peripheral blood of patients with




FIGURE 1 RNA transcriptome expression analyses of the skin of children with localised scleroderma (LS). (a) Dendrograms from hierarchical cluster mapping using complete linkage Euclidean distance show the groups of children with LS identified based on gene expression (genes listed on bottom) and skin histopathologic features (unique clusters designated as inflammatory, fibroproliferative, or healthy-like), stratified by clinical disease activity status. Numbers to the right of the dendrograms represent individual skin biopsy samples. Map clustering confirmed distinct differences in juvenile LS patients compared to healthy controls. B and C, Results of pathway analyses show the functional pathways for genes that were up-regulated (b) and those that were down-regulated (c) in the skin of children with juvenile LS relative to healthy controls. Horizontal blue bars show the *p* values on a logarithmic scale. Vertical orange lines represent the ratio of genes listed to the number of genes associated with each pathway. AR, androgen receptor; CSB, Cockayne syndrome complementation group B protein; GALNT12, N-acetylgalactosaminyltransferase 12; MHC, major histocompatibility complex; PD-1, programmed death 1; PKN1, serine/threonine protein kinase N1; PRC2, polycomb-repressive complex 2; rRNA, ribosomal RNA; SIRT1 sirtuin 1; TCA, tricarboxylic acid; TCR, T cell receptor; tRNA, transfer RNA.¹⁸

morphea and in the skin of patients with SSc. Moreover, the histologic scores of skin inflammation (based on numbers and categories of inflammatory cell infiltrates) were significantly correlated with the expression levels of HLA-DPB1, HLA-DQA2, HLA-DRA, and STAT1

genes. Collagen thickness correlated with the expression levels of collagen organization genes as well as with genes found to be correlated with the severity of inflammation, including genes for major histocompatibility complex (MHC) class I, MHC class II, and IFN γ

signalling. Identifying 3 distinct genetic signatures and associated genes is a significant step forward in terms of diagnosing and differentiating LS from SSc.

As it can already be seen in the work of Schutt et al. (Figure 1c) with the downregulation of DNA methylation pathway in LS patients,¹⁸ at the level of DNA genome, epigenetic regulation is another exciting field of study that also includes new findings for LS. Coit et al. studied DNA methylation differences and similarities between juvenile systemic sclerosis and juvenile LS compared to matched healthy controls.⁷³ Genome-wide DNA methylation changes in peripheral blood mononuclear cell samples were assessed using the MethylationEPIC array followed by bioinformatic analysis and limited functional assessment. A total of 105 and 144 differentially methylated sites were identified compared to healthy controls in juvenile systemic sclerosis and juvenile LS, respectively. Most differentially methylated sites and genes represented were unique to either juvenile systemic sclerosis or juvenile LS suggesting a different underlying epigenetic pattern in both diseases. Among shared differentially methylated genes, methylation levels in a CpG site in FGFR2 can distinguish between LS and healthy PBMCs with a high accuracy. Canonical pathway analysis revealed that inflammatory pathways were enriched in genes differentially methylated in SSc, including STAT3, NF- κ B, and IL-15 pathways. In contrast, the Hippo signalling pathway was enriched in LS. This study revealed important insights into juvenile-onset LS and suggested a potentially novel epigenetic diagnostic biomarker for LS.

Single-cell genomics is the study of the individuality of cells with a sequencing technique. Although young, the field has now entered a more mature stage and is beginning to provide with valuable new clues on healthy immune system and on autoimmune diseases. Especially for highly heterogeneous diseases, such as LS, single cell techniques can truly contribute to in-depth understanding of the disease pathology, underlying genetic signatures and their dynamic revelation in the disease phenotypes. This can become a path to discover new prognostic biomarkers including those indicating and even predicting response to treatment, as it already happened in oncology field. In 2020, Mirizio et al. conducted a pilot single-cell RNAseq on paired skin biopsy specimens from 3 patients with LS, exploring different sample preparation strategies for 10 \times Genomics sequencing.⁷⁴ Levels of cell viability and yield were comparable between frozen and freshly preserved cells. Furthermore, gene expression between preservation methods was collectively significantly correlated and conserved across all 18 identified cell cluster populations. The average expression of genes for major cell groups, such as keratinocytes, T/NK cells, DC/macrophages, fibroblasts, and pericytes, demonstrated a strong correlation between the average

counts for each gene across all cells in the respective group. This suggests that employing standardized cryopreservation protocols for the skin tissue will help facilitate multi-site collaborations looking to identify mechanisms of disease in disorders characterised by cutaneous pathology with single cell technology.

Another regulatory RNA that has been recently studied in LS is long non-coding RNA (lncRNA). lncRNAs are approx. Two hundred nucleotides in length and lack protein-coding potential. Increasing evidence indicates that lncRNAs exert an irreplaceable role in disease initiation, progression, and are novel molecular biomarkers for diagnosis and prognosis of most human diseases. Furthermore, lncRNAs and the pathways they influence might represent promising therapeutic targets. Profiling of inflammatory cells in skin samples from paediatric LS was performed, aiming at lncRNA profile analysis.⁷⁵ Among them, CD4+ T-cells were up-regulated. Co-culture dermal fibroblasts with CD4+ T-cells promoted fibrosis of fibroblasts. Candidate lncRNAs were further explored by lncRNAs-seq between the normal skin tissues and paediatric LS tissues, and the lncRNAs-seq between fibroblasts co-cultured with CD4+ T lymphocytes and control fibroblasts. By comparing the two datasets, the authors identified eight up-regulated (LINC01184, BAALC-AS1, AF165147.1, TRAM2-AS1, MIR100HG, CHROMR, LINC00665, ZEB1-AS1) and three down-regulated (LINC00662, CARMN, PAX8-AS1) lncRNAs, which were the potential lncRNAs for the phenotype of paediatric LS. The identified lncRNAs can become both valuable diagnostic markers and treatment candidates for paediatric LS.

2.4 | Biomarkers in trials

To date, there are seven completed trials focusing on LS, from which one published their results (2018 completed, phase II, NCT02915835).⁷⁶ As to the recruiting studies, morphea in adults and children clinical trial has been initiated (NCT01808937).⁷⁷ With five hundred participants and completion planned in 2027, the trial will be first prospective six-year-long study where both adults and children with LS will be analysed longitudinally for multiple disease parameters including disease activity, other autoimmune conditions, and excitingly, for DNA signature.

Nice Hospital, France, is currently recruiting for a trial of microRNA in LS (NCT04148716).⁷⁸ Prospective study with 18 participants will provide with a 2-year insight into miRNA signature in LS. The team is studying the involvement of pro-fibrotic “key” miRNAs called “FibromiRs”, including 3 miRNAs from the DNMT3a locus (miR-199a-3p, miR-199a-5p and miR-214 - characterised by the host laboratory) associated with monitoring the response to TGF- β in fibroblasts

and their potential interaction with pharmacological treatments such as nintedanib and/or PPAR γ agonists. The approach is part of a pilot study that can lead to a larger project after validation of the hypotheses.

3 | CONCLUSIONS

Research in biomarkers is an extensive field in autoimmunity that constantly introduces new biomarkers and biological insights into the field. Despite the lack of a single optimal biomarker, a combination of them could be efficient to diagnose and assess the state of the disease.

In LS, the diagnostic perspective with existing biomarkers is still not as good as in other autoimmune diseases. This is due to the low specificity and sensitivity of the common biomarkers, which are often shared with other autoimmune diseases such as lupus and RA. Differentiation among sub-types of sclerodermas and from other skin autoimmune conditions are also important tasks that require improved approaches. Lack of prognostic biomarkers including those reflecting response to treatment also attracts increasing attention in the research and clinical communities. Here recent genomic investigations are exciting contributions, with lncRNA, microRNA and epigenetic biomarkers being a potent new direction for research and clinics.

With three recruiting trials for new treatment strategies of LS (accessed November 2023), it becomes relevant to look up at genomic signature and responses to medication at critical checkpoints of the disease. Recent studies highlighted in this review suggest multiple proteomic and genetic biomarkers that could aid in-depth investigation of LS and improve treatment strategies.

ACKNOWLEDGEMENTS

We acknowledge Dr Kathryn S. Torok, Division of Rheumatology, Department of Pediatrics, Children's Hospital of Pittsburgh, University of Pittsburgh, Pittsburgh, PA, United States, for fruitful discussions related to this review. DTU Alliance fellowship (#102140) by the Technical University of Denmark is acknowledged for funding of this work. The financial source was not involved in the paper preparation process.

CONFLICT OF INTEREST STATEMENT

None to declare.

AUTHOR CONTRIBUTIONS

Adrian Hernandez-Bustos: Writing – original draft (lead); Writing – review & editing (equal). **Begona Bo-los:** Writing – original draft (supporting). **Kira Astakhova:** Conceptualisation (lead); Writing – original draft (equal); Writing – review & editing (lead).

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ETHICS STATEMENT

Not applicable.

ORCID

Kira Astakhova  <https://orcid.org/0000-0003-4878-0301>

REFERENCES

- Rao X, Sigdel KR. Regulation of inflammation in autoimmune diseases. *J Immunol Res*. 2019. <https://doi.org/10.1155/2019/7403796>
- Surace AEA, Hedrich CM. The role of epigenetics in autoimmune/inflammatory disease. *Front Immunol*. 2019;10:1–16. <https://doi.org/10.3389/fimmu.2019.01525>
- Peterson LS, Nelson AM, Su WPD. Classification of morphea (localized scleroderma). *Mayo Clin Proc*. 1995;70(11):1068–76. <https://doi.org/10.4065/70.11.1068>
- Laxer RM, Zulian F. Localized scleroderma. *Curr Opin Rheumatol*. 2006;18(6):606–13. <https://doi.org/10.1097/01.BOR.0000245727.40630.C3>
- Prasad S, Zhu JL, Schollaert-Fitch K, Torok KS, Jacobe HT. An evaluation of the performance of current morphea subtype classifications. *JAMA Dermatol*. 2021;157(4):399–405. <https://doi.org/10.1001/JAMADERMATOL.2020.5809>
- Jacobe H, Ahn C, Arnett FC, Reveille JD. Major histocompatibility complex class I and class II alleles may confer susceptibility to or protection against morphea: findings from the morphea in adults and children cohort. *Arthritis Rheumatol*. 2014;66(11):3170–7. <https://doi.org/10.1002/art.38814>
- Makino T, Jinnin M, Etoh M, Yamane K, Kajihara I, Makino K, et al. Down-regulation of microRNA-196a in the sera and involved skin of localized scleroderma patients. *Eur J Dermatol*. 2014;24(4):470–6. <https://doi.org/10.1684/EJD.2014.2384>
- Kurzinski K, Torok KS. Cytokine profiles in localized scleroderma and relationship to clinical features. *Cytokine*. 2011;55(2):157–64. <https://doi.org/10.1016/j.cyto.2011.04.001>
- Papara C, De Luca DA, Bieber K, Vorobyev A, Ludwig RJ. Morphea: the 2023 update. *Front Med*. 2023;10. <https://doi.org/10.3389/FMED.2023.1108623>
- Penmetsa GK, Sapra A. Morphea, StatPearls; 2023. <https://www.ncbi.nlm.nih.gov/books/NBK559010/>. Accessed 4 May 2023.
- Joly P, Bamberger N, Crickx B, Belaich S. Treatment of severe forms of localized scleroderma with oral corticosteroids: follow-up study on 17 patients. *Arch Dermatol*. 1994;130(5):663–4. <https://doi.org/10.1001/ARCHDERM.1994.01690050133027>
- Zulian F, Vallongo C, Patrizi A, Belloni-Fortina A, Cutrone M, Alessio M, et al. A long-term follow-up study of methotrexate in juvenile localized scleroderma (morphea). *J Am Acad Dermatol*. 2012;67(6):1151–6. <https://doi.org/10.1016/j.jaad.2012.03.036>
- Zulian F, Martini G, Vallongo C, Vittadello F, Falcini F, Patrizi A, et al. Methotrexate treatment in juvenile localized scleroderma: a randomized, double-blind, placebo-controlled trial. *Arthritis Rheum*. 2011;63(7):1998–2006. <https://doi.org/10.1002/art.30264>
- Zhao M, Wu J, Wu H, Sawalha AH, Lu Q. Clinical treatment options in scleroderma: recommendations and comprehensive review. *Clin Rev Allergy Immunol*. 2022;62(2):273–91. <https://doi.org/10.1007/S12016-020-08831-4>
- Glaser D, Torok KS. Evaluation and treatment of pediatric localized scleroderma: pearls and updates. *Curr Treat Options Rheumatol*. 2021;7:1–20. <https://doi.org/10.1007/S40674-021-00170-5>

16. Li SC, Higgins GC, Chen M, Torok KS, Rabinovich CE, Stewart K, et al. Extracutaneous involvement is common and associated with prolonged disease activity and greater impact in juvenile localized scleroderma. *Rheumatology*. 2021;60(12):5724–33. <https://doi.org/10.1093/RHEUMATOLOGY/KEAB238>
17. Bernatsky S, Joseph L, Pineau CA, Belisle P, Hudson M, Clarke AE. Scleroderma prevalence: demographic variations in a population-based sample. *Arthritis Care Res*. 2009;61(3):400–4. <https://doi.org/10.1002/ART.24339>
18. Schutt C, Mirizio E, Salgado C, Reyes-Mugica M, Wang X, Chen W, et al. Transcriptomic evaluation of juvenile localized scleroderma skin with histologic and clinical correlation. *Arthritis Rheumatol*. 2021;73(10):1921–30. <https://doi.org/10.1002/art.41758>
19. Kreuter A, Krieg T, Worm M, Wenzel J, Moinszadeh P, Kuhn A, et al. German guidelines for the diagnosis and therapy of localized scleroderma. *J Ger Soc Dermatol*. 2016;14(2):199–216. <https://doi.org/10.1111/ddg.12724>
20. Asano Y, Fujimoto M, Ishikawa O, Sato S, Jinnin M, Takehara K, et al. Diagnostic criteria, severity classification and guidelines of localized scleroderma. *J Dermatol*. 2018;45(7):755–80. <https://doi.org/10.1111/1346-8138.14161>
21. Furuhashi T, Torii K, Ikumi K, Kato H, Nishida E, Morita A. Ultraviolet A1 phototherapy for the treatment of localized scleroderma. *J Dermatol*. 2020;47(7):792–5. <https://doi.org/10.1111/1346-8138.15368>
22. Wang Y, ling Shan J, yan Chen H, feng Wu Z. Comparison of 2-D shear wave elastography with clinical score in localized scleroderma: a new method to increase the diagnostic accuracy. *J Dermatol*. 2019;46(2):131–8. <https://doi.org/10.1111/1346-8138.14713>
23. Pérez M, Zuccaro J, Mohanta A, Tijerin M, Laxer R, Pope E, et al. Feasibility of using elastography ultrasound in pediatric localized scleroderma (morphea). *Ultrasound Med Biol*. 2020;46(12):3218–27. <https://doi.org/10.1016/j.ultrasmedbio.2020.08.007>
24. Zulian F, Culp R, Sperotto F, Anton J, Avcin T, Baildam EM, et al. Consensus-based recommendations for the management of juvenile localised scleroderma. *Ann Rheum Dis*. 2019;78(8):1019–24. <https://doi.org/10.1136/annrheumdis-2018-214697>
25. Su P, Cao T, Tang MBY, Tey HL. In vivo high-definition optical coherence tomography: a bedside diagnostic aid for morphea. *JAMA Dermatol*. 2015;151(2):234–5. <https://doi.org/10.1001/jamadermatol.2014.2668>
26. Zulian F, Vallongo C, de Oliveira SKF, Punaro MG, Ros J, Mazur-Zielinska H, et al. Congenital localized scleroderma. *J Pediatr*. 2006;149(2):248–51. <https://doi.org/10.1016/j.jpeds.2006.04.052>
27. Li SC. Scleroderma in children and adolescents: localized scleroderma and systemic sclerosis. *Pediatr Clin North Am*. 2018;65(4):757–81. <https://doi.org/10.1016/j.pcl.2018.04.002>
28. Simmonds M, Gough S. The HLA region and autoimmune disease: associations and mechanisms of action. *Curr Genom*. 2007;8(7):453–65. <https://doi.org/10.2174/138920207783591690>
29. Hasegawa M, Fujimoto M, Kikuchi K, Takehara K. Elevated serum levels of interleukin 4 (IL-4), IL-10, and IL-13 in patients with systemic sclerosis - PubMed. *J Rheumatol*. 1997;24:328–32. <https://pubmed.ncbi.nlm.nih.gov/9034992/> Accessed 26 March 2021.
30. Snarskaya ES, Vasileva KD. Localized scleroderma: actual insights and new biomarkers. *Int J Dermatol*. 2022;61(6):667–74. <https://doi.org/10.1111/IJD.15811>
31. Tafuri WL, Tomokane TY, Silva AMG, Kanashiro-Galo L, Mosser DM, Quaresma JAS, et al. Skin fibrosis associated with keloid, scleroderma and Jorge Lobo's disease (Iacaziosis): an immunohistochemical study. *Int J Exp Pathol*. 2022;103(6):234–44. <https://doi.org/10.1111/IJEP.12456>
32. Tawfik AA, Shokir H, Soliman M, Salah L, Fathy S. Pulsed dye laser in the treatment of localized scleroderma and its effects on CD34+ and factor XIIIa+ cells: an immunohistochemical study. *Am J Clin Dermatol*. 2013;14(3):235–41. <https://doi.org/10.1007/S40257-013-0027-7>
33. Xie Y, Zhang X, Wakasugi S, Makino T, Inoue Y, Ihn H. Immunohistochemical characterization of the cellular infiltrate in localized scleroderma. *Int J Dermatol*. 2008;47(5):438–42. <https://doi.org/10.1111/J.1365-4632.2008.03615.X>
34. Xie Y, Zhang X, Inoue Y, Wakasugi S, Makino T, Ihn H. Expression of CD1a and CD86 on scleroderma Langerhans cells. *Eur J Dermatol*. 2008;18:50–4. <https://doi.org/10.1684/EJD.2008.0310>
35. Ihn H, Yazawa N, Kubo M, Yamane K, Sato S, Fujimoto M, et al. Circulating levels of soluble CD30 are increased in patients with localized scleroderma and correlated with serological and clinical features of the disease. *J Rheumatol*. 2000;27:698–702. <https://europepmc.org/article/med/10743811> Accessed 31 October 2023.
36. Mimura Y, Ihn H, Jinnin M, Asano Y, Yamane K, Tamaki K. Rheumatoid factor isotypes in localized scleroderma. *Clin Exp Dermatol*. 2005;30:405–8. <https://doi.org/10.1111/j.1365-2230.2005.01776.x>
37. Zulian F, Athreya BH, Laxer R, Nelson AM, Feitosa de Oliveira SK, Punaro MG, et al. Juvenile localized scleroderma: clinical and epidemiological features in 750 children. An international study. *Rheumatology*. 2006;45:614–20. <https://doi.org/10.1093/rheumatology/kei251>
38. Zulian F, Vallongo C, Woo P, Russo R, Ruperto N, Harper J, et al. Localized scleroderma in childhood is not just a skin disease. *Arthritis Rheum*. 2005;52:2873–81. <https://doi.org/10.1002/art.21264>
39. Wu EY, Li SC, Torok KS, Virkud YV, Fuhlbrigge RC, Rabinovich CE. Baseline description of the juvenile localized scleroderma subgroup from the childhood arthritis and Rheumatology research alliance legacy registry. *ACR Open Rheumatol*. 2019;1:119–24. <https://doi.org/10.1002/acr2.1019>
40. Leitenberger JJ, Cayce RL, Haley RW, Adams-Huet B, Bergstresser PR, Jacobe HT. Distinct autoimmune syndromes in morphea. *Arch Dermatol*. 2009;145:545–50. <https://doi.org/10.1001/archdermatol.2009.79>
41. Kurzinski KL, Zigler CK, Torok KS. Prediction of disease relapse in a cohort of paediatric patients with localized scleroderma. *Br J Dermatol*. 2019;180:1183–9. <https://doi.org/10.1111/bjd.17312>
42. Torok KS, Kurzinski K, Kelsey C, Yabes J, Magee K, Vallejo AN, et al. Peripheral blood cytokine and chemokine profiles in juvenile localized scleroderma: T-helper cell associated cytokine profiles. *Semin Arthritis Rheum*. 2015;45:284–93. <https://doi.org/10.1016/j.semarthrit.2015.06.006> Peripheral
43. Arkachaisri T, Fertig N, Pino S, Medsger TA. Serum autoantibodies and their clinical associations in patients with childhood- and adult-onset linear scleroderma. A single-center study. *J Rheumatol*. 2008;35:2439–44. <https://doi.org/10.3899/jrheum.080098>
44. O'Brien JC, Cyrus N, Jacobe H. Identification of cytokine biomarkers in morphea. *J Invest Dermatol*. 2016;136:S2. <https://doi.org/10.1016/j.jid.2016.02.033>
45. Mertens JS, de Jong EMGJ, van den Hoogen LL, Wienke J, Thurlings RM, Seyger MMB, et al. The identification of CCL18 as biomarker of disease activity in localized scleroderma. *J Autoimmun*. 2019;101:86–93. <https://doi.org/10.1016/j.jaut.2019.04.008>
46. Tsuruta M, Fukushima S, Hayano S, Yoshikawa T, Kawabe K, Makino T, et al. Secreted protein acidic and Rich in cysteine, a new biomarker of localized scleroderma. *J Invest Dermatol*. 2017;137:S270. <https://doi.org/10.1016/j.jid.2017.07.652>
47. Burger E, Paniagua R, Monson N, Jacobe H. Myelin basic protein antibodies as a biomarker in morphea. *J Invest Dermatol*. 2018;138:S5. <https://doi.org/10.1016/j.jid.2018.03.031>
48. Zhu JL, Paniagua RT, Chen HW, Florez-Pollack S, Kunzler E, Teske N, et al. Autoantigen microarrays reveal myelin basic protein autoantibodies in morphea. *J Transl Med*. 2022;20. <https://doi.org/10.1186/S12967-022-03246-5>
49. Knobler R, Moinszadeh P, Hunzelmann N, Kreuter A, Cozzio A, Mouthon L, et al. European Dermatology Forum S1-guideline on the diagnosis and treatment of sclerosing diseases of the skin,

- Part 1: localized scleroderma, systemic sclerosis and overlap syndromes. *J Eur Acad Dermatol Venereol.* 2017;31:1581–94. <https://doi.org/10.1111/jdv.14458>
50. Florez-Pollack S, Kunzler E, Jacobe HT. Morphea: current concepts. *Clin Dermatol.* 2018;36:475–86. <https://doi.org/10.1016/j.clindermatol.2018.04.005>
 51. Warner Dharamsi J, Victor S, Aguwa N, Ahn C, Arnett F, Mayes MD, et al. Morphea in adults and children cohort III: nested case-control study - the clinical significance of autoantibodies in morphea. *JAMA Dermatol.* 2013;149(10):1159–65. <https://doi.org/10.1001/jamadermatol.2013.4207>
 52. Khatri S, Torok KS, Mirizio E, Liu C, Astakhova K. Autoantibodies in morphea: an update. *Front Immunol.* 2019;10:1–12. <https://doi.org/10.3389/fimmu.2019.01487>
 53. Singhvi G, Manchanda P, Krishna Rapalli V, Kumar Dubey S, Gupta G, Dua K. MicroRNAs as biological regulators in skin disorders. *Biomed Pharmacother.* 2018;108:996–1004. <https://doi.org/10.1016/J.BIOPHA.2018.09.090>
 54. Li Y, Huang J, Guo M, Zuo X. MicroRNAs regulating signaling pathways: potential biomarkers in systemic sclerosis, genomics. *Proteom Bioinform.* 2015;13(4):234–41. <https://doi.org/10.1016/J.GPB.2015.07.001>
 55. Li Y, Zhang J, Lei Y, Lyu L, Zuo R, Chen T. MicroRNA-21 in skin fibrosis: potential for diagnosis and treatment. *Mol Diagn Ther.* 2017;21(6):633–42. <https://doi.org/10.1007/S40291-017-0294-8>
 56. Peng WJ, Tao JH, Mei B, Chen B, Li BZ, Yang GJ, et al. MicroRNA-29: a potential therapeutic target for systemic sclerosis. *Expert Opin Ther Targets.* 2012;16(9):875–9. <https://doi.org/10.1517/14728222.2012.708339>
 57. Chouri E, Servaas NH, Bekker CPJ, Affandi AJ, Cossu M, Hillen MR, et al. Serum microRNA screening and functional studies reveal miR-483-5p as a potential driver of fibrosis in systemic sclerosis. *J Autoimmun.* 2018;89:162–70. <https://doi.org/10.1016/J.JAUT.2017.12.015>
 58. Etoh M, Jinnin M, Makino K, Yamane K, Nakayama W, Aoi J, et al. microRNA-7 down-regulation mediates excessive collagen expression in localized scleroderma. *Arch Dermatol Res.* 2013;305(1):9–15. <https://doi.org/10.1007/S00403-012-1287-4>
 59. Rongioletti F, Ferrelli C, Atzori L, Bottoni U, Soda G. Scleroderma with an update about clinico-pathological correlation. *G Ital Dermatol Venereol.* 2018;153(2):208–15. <https://doi.org/10.23736/S0392-0488.18.05922-9>
 60. Liu Q, Lin J, Han J, Zhang Y, Lu J, Tu W, et al. Immunoglobulin G galactosylation levels are decreased in systemic sclerosis patients and differ according to disease subclassification. *Scand J Rheumatol.* 2020;49(2):146–53. <https://doi.org/10.1080/03009742.2019.1641615>
 61. Mirizio E, Liu C, Yan Q, Waltermire J, Mandel R, Schollaert KL, et al. Genetic signatures from RNA sequencing of pediatric localized scleroderma skin. *Front Pediatr.* 2021;9:669116. <https://doi.org/10.3389/FPED.2021.669116/FULL>
 62. Makino K, Jinnin M, Hirano A, Yamane K, Eto M, Kusano T, et al. The downregulation of microRNA let-7a contributes to the excessive expression of type I collagen in systemic and localized scleroderma. *J Immunol.* 2013;190(8):3905–15. <https://doi.org/10.4049/jimmunol.1200822>
 63. Kreuter A, Krieg T, Worm M, Wenzel J, Gambichler T, Kuhn A, et al. Diagnosis and therapy of localized scleroderma. *J Der Dtsch Dermatol Ges.* 2009;7(s6). <https://doi.org/10.1111/J.1610-0387.2009.07178.X>
 64. Reich A, Meurer M, Eckes B, Friedrichs J, Muller DJ. Surface morphology and mechanical properties of fibroblasts from scleroderma patients. *J Cell Mol Med.* 2009;13(8b):1644–52. <https://doi.org/10.1111/J.1582-4934.2008.00401.X>
 65. Iwayama T, Olson LE. Involvement of PDGF in fibrosis and scleroderma: recent insights from animal models and potential therapeutic opportunities. *Curr Rheumatol Rep.* 2013;15(2):304. <https://doi.org/10.1007/S11926-012-0304-0>
 66. Odell ID, Agrawal K, Sefik E, Odell AV, Caves E, Kirkiles-Smith NC, et al. IL-6 trans-signaling in a humanized mouse model of scleroderma. *Proc. Natl. Acad. Sci. U.S.A.* 2023;120(37). <https://doi.org/10.1073/PNAS.2306965120>
 67. Wodok-Wieczorek K, Salwowska N, Syguta E, Wodok A, Wcislo-Dziadecka D, Bebenek K, et al. The correlation between serum E-selectin levels and soluble interleukin-2 receptors with relation to disease activity in localized scleroderma. *Postep Dermatol Alergol.* 2018;35(6):614–9. <https://doi.org/10.5114/ADA.2018.77613>
 68. Ferguson ID, Weiser P, Torok KS. A case report of successful treatment of recalcitrant childhood localized scleroderma with infliximab and leflunomide. *Open Rheumatol J.* 2015;9(1):30–5. <https://doi.org/10.2174/18743129014090100030>
 69. Buka RL, Cunningham BB. Connective tissue disease in children. *Pediatr Ann.* 2005;34(3):225–38. <https://doi.org/10.3928/0090-4481-20050301-11>
 70. Saracino AM, Kelberman D, Otto GW, Gagunashvili A, Abraham DJ, Denton CP. Unravelling morphoea aetiopathogenesis by next-generation sequencing of paired skin biopsies. *Arch Dermatol Res.* 2023;315(7):2035–56. <https://doi.org/10.1007/S00403-023-02541-5>
 71. Yeh YW, Cheng CC, Yang ST, Tseng CF, Chang TY, Tsai SY, et al. Targeting the VEGF-C/VEGFR3 axis suppresses Slug-mediated cancer metastasis and stemness via inhibition of KRAS/YAP1 signaling. *Oncotarget.* 2017;8(3):5603–18. <https://doi.org/10.18632/ONCOTARGET.13629>
 72. Wei J, Melichian D, Komura K, Hinchcliff M, Lam AP, Lafyatis R, et al. Canonical Wnt signaling induces skin fibrosis and subcutaneous lipatrophy: a novel mouse model for scleroderma? *Arthritis Rheum.* 2011;63(6):1707–17. <https://doi.org/10.1002/art.30312>
 73. Coit P, Schollaert KL, Mirizio EM, Torok KS, Sawalha AH. DNA methylation patterns in juvenile systemic sclerosis and localized scleroderma. *Clin Immunol.* 2021;228:108756. <https://doi.org/10.1016/J.CLIM.2021.108756>
 74. Mirizio E, Tabib T, Wang X, Chen W, Liu C, Lafyatis R, et al. Single-cell transcriptome conservation in a comparative analysis of fresh and cryopreserved human skin tissue: pilot in localized scleroderma. *Arthritis Res Ther.* 2020;22(1):263. <https://doi.org/10.1186/S13075-020-02343-4>
 75. Gong Y, Liu H, Li G, Sun L. Identification of lncRNA expression profiles in pediatric localized scleroderma. *J Cosmet Dermatol.* 2022;21(11):6422–7. <https://doi.org/10.1111/JOCD.15318>
 76. Nct, riociguat in scleroderma associated digital ulcers. (2016). <https://clinicaltrials.gov/study/NCT02915835>. Accessed 15 November 2023.
 77. N.L. of M. of the U. government NIH, morphea in adults and children (MAC) cohort study: a morphea registry and DNA repository | ClinicalTrials.gov, (n.d.). <https://www.clinicaltrials.gov/study/NCT01808937>. Accessed 15 November 2023.
 78. N.L. of M. of the U. government NIH, microRNAs in Systemic Scleroderma | ClinicalTrials.gov, (n.d.). <https://www.clinicaltrials.gov/study/NCT04148716?distance=50&cond=NCT04148716&rank=1>. Accessed 15 November 2023.

How to cite this article: Hernández-Bustos A, Bolos B, Astakhova K. Biomarkers in skin autoimmunity—An update on localised scleroderma. *Skin Health Dis.* 2024;4(2):e335. <https://doi.org/10.1002/ski2.335>