Density functional theory calculations for the hydrogen evolution reaction in an electrochemical double layer on the Pt(111) electrode

Skulason, Egill; Karlberg, Gustav; Rossmeisl, Jan; Pedersen, Thomas Bligaard; Greeley, Jeffrey Philip; Jonsson, Hannes; Nørskov, Jens Kehlet

Published in:
Physical Chemistry Chemical Physics

Link to article, DOI:
[10.1039/b700099e](https://doi.org/10.1039/b700099e)

Publication date:
2007

[Link back to DTU Orbit](#)

Citation (APA):
Skulason, E., Karlberg, G., Rossmeisl, J., Pedersen, T. B., Greeley, J. P., Jonsson, H., & Nørskov, J. K. (2007). Density functional theory calculations for the hydrogen evolution reaction in an electrochemical double layer on the Pt(111) electrode. *Physical Chemistry Chemical Physics*, 9, 3241. <https://doi.org/10.1039/b700099e>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Electronic Supporting Information

Density functional theory calculations for the hydrogen evolution reaction in an electrochemical double layer on the Pt(111) electrode

E. Skúlason^{1,2}, G. S. Karlberg¹, J. Rossmeisl¹, T. Bligaard^{1,2},
J. Greeley¹, H. Jónsson^{3,*}, J. K. Nørskov^{1,*}

¹Center for Atomic-scale Materials Design, Department of Physics, Building 307, NanoDTU, Technical University of Denmark, DK-2800 Lyngby, Denmark

²Science Institute, University of Iceland, Dunhaga 3, IS-107 Reykjavik, Iceland

³Faculty of Science, VR-II, University of Iceland, IS-107 Reykjavik, Iceland

Figure S1: The variation of the activation energy (circles) and the reaction energy (squares) with number of Pt layers used for the Density Functional Theory calculations. These tests were made by calculating the IS, TS and FS for the Heyrovsky reaction ($H^+ + e^- + H_{ad} \rightarrow H_2$) with 1-8 Pt layers. The energy values have converged completely using 6 Pt layers in the model system. In the present study we have chosen 3 Pt layers to save computational time. The difference between the 3 Pt layers and the 8 Pt layers test cases is very small, 0.07 eV for the activation energy and 0.11 eV for the reaction energy.

* Corresponding authors, emails: hj@hi.is & norskov@fysik.dtu.dk

Figure S2: The variation of the work function with increasing vacuum distance between the repeated slabs. The test case is the IS for the Heyrovsky reaction ($H^+ + e^- + H_{ad} \rightarrow H_2$). The work function has completely converged with 12 Å of vacuum. In the present study we use 10 Å of vacuum for this system. The difference in the work function obtained by using 10 Å or 12 Å of vacuum is very small, or 0.035 eV, not affecting the trends observed in the study.