



## Return-map for semiconductor lasers with optical feedback

**Mørk, Jesper; Tromborg, Bjarne; Sabbatier, H.; Sørensen, Mads Peter**

*Published in:*  
1999 IEEE LEOS Annual Meeting Conference Proceedings

*Link to article, DOI:*  
[10.1109/LEOS.1999.811790](https://doi.org/10.1109/LEOS.1999.811790)

*Publication date:*  
1999

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Mørk, J., Tromborg, B., Sabbatier, H., & Sørensen, M. P. (1999). Return-map for semiconductor lasers with optical feedback. In *1999 IEEE LEOS Annual Meeting Conference Proceedings* (Vol. 2, pp. 445-446). IEEE. <https://doi.org/10.1109/LEOS.1999.811790>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Return-map for semiconductor lasers with optical feedback


*J. Mørk and B. Tromborg*

Research Center COM, Technical University of Denmark  
Build. 349, DK-2800 Lyngby, Denmark. E-mail: jm@com.dtu.dk

*H. Sabbatier, and M.P. Sørensen*

Dept. of Mathematical Modelling Technical University of Denmark  
Building 321, DK-2800 Lyngby, Denmark.

It is well known that a semiconductor laser exposed to moderate optical feedback and biased near threshold exhibits the phenomenon of low-frequency intensity fluctuations (LFF) [1-6]. While this behavior can be numerically simulated using the so-called Lang-Kobayashi model [2], the interpretation of the phenomenon has remained a controversy [1-6]. The LFF consist of a sudden drop in intensity followed by a build-up in steps of the external cavity roundtrip time,  $\tau$ , before a new drop-out occurs. The phenomenon has been attributed to a kind of chaotic itinerancy [3] and a bifurcation cascade has been identified very recently [6]. These results give insight into the behavior observed on a short time-scale, but do not explain some of the pronounced features of the LFF seen for moderate feedback levels; namely the stepwise build-up and its characteristic time of about 15 steps close to the solitary laser threshold. In this paper we present new results related to the slow time-scale behaviour of LFF which give a simple explanation of these general characteristics, as well as providing a new tool for studying the statistics of the LFF.




**Fig. 1.** Calculated return-map for low-frequency fluctuations. Feedback level  $\kappa=0.15$ , bias current  $J=J_{th}$  (threshold current without feedback). The inset shows a schematic of the return-map. In the actual calculation, two iterations of the  $\tau$  are

shown. **Fig. 2.** Time-dependence of photon number calculated from the return-map and a stochastic noise term.


Fig. 1 shows the calculated return-map for the system between the intensity at step  $n$  and the following step,  $n+1$ . Calculations are based on the iterative model derived in [2,4] which treats the entire spectrum of the laser emission within  $\tau$  as the basic variable and therefore is valid even in the case where fast and irregular, possibly chaotic, dynamics occur on a very short time scale. Starting out from the stationary state, the intensity drops to the lower branch of the map due to noise-induced switching over a potential barrier [1], the barrier maximum corresponding to the position of an unstable solution [2]. The intensity is now forced to build up until the point  $I_n = I_{bi}$  is reached. After this point, the system is bistable and a new switching to the low intensity branch is possible after a stochastic dwell time

determined by the noise of the system. This explains the overall slow time scale behaviour observed experimentally. In Fig. 1 we have made two cycles of iteration and the discrete points are seen to fall almost exactly on the same curve, demonstrating for the first time that a return-map can describe the dynamics. Fig. 2 shows an example of a calculated time trace using the map when a stochastic noise term is added, and closely resembles experimental observations.

In Fig. 3 we investigate the parameter dependence of the number of steps needed to reach



the point of bistability,  $n_{bi}$ , which gives a lower limit to the period of the LFF,  $T_{low} = n_{bi}\tau$ . For fixed feedback strength,  $\kappa$ , the period increases rapidly when the threshold current with feedback is reached and drops for increasing bias current. Conversely, when the laser is biased close to the threshold current without feedback,  $J_{th}$ , about 15 external roundtrips are required for the intensity to build-up, varying only weakly with  $\kappa$ . Both of these predictions of the return-map are in good agreement with experimental observations, and provides, to our knowledge, the first explanation of these generally observed dependencies. The return-map also provides a straightforward explanation of the “dead” or “forbidden” zone of LFF periods recently observed experimentally [5], and should be useful for studying the statistical properties of the LFF.



**Fig. 3.** Bias current dependency of the number of external cavity roundtrips to achieve bistability for feedback levels  $\kappa=0.15$  and  $0.30$ . The inset shows the  $\kappa$  dependency when the laser is biased at the threshold current of the laser without feedback.

- [1] C. H. Henry and R. F. Kazarinov, *IEEE J. Quantum Electron.*, **QE-22**, 294 (1986).
- [2] J. Mørk, B. Tromborg, P. L. Christiansen, *IEEE J. Quantum Electron.*, **24**, 123 (1988).
- [3] I. Fischer et al., *Phys. Rev. Lett.*, **76**, 220 (1996).
- [4] B. Tromborg, J. Mørk, and V. Velichansky, *Quantum Semiclass. Opt.*, **9**, 831 (1997).
- [5] D. Sukow, J. R. Gardner, and D. J. Gauthier, *Phys. Rev. A.*, **56**, R3370 (1997).
- [6] A. Hohl and A. Gavrielides, *Phys. Rev. Lett.*, **82**, 1148 (1999).