

Bioactivity and phylogeny of the marine bacterial genus Pseudoalteromonas

Vynne, Nikolaj Grønnegaard

Publication date: 2011

Document Version Publisher's PDF, also known as Version of record

Link back to DTU Orbit

Citation (APA): Vynne, N. G. (2011). *Bioactivity and phylogeny of the marine bacterial genus Pseudoalteromonas*. Technical University of Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.





Ph.D. thesis

by

Nikolaj Grønnegaard Vynne

2011

Technical University of Denmark

National Food Institute

Division of Industrial Food Research

Cover illustrations:

Map of the Galathea 3 route (www.galathea3.dk)

Structure of the antibiotic indolmycin

Excerpt of 16S rRNA phylogenetic tree

P. luteoviolacea S4054 grown in MMM

PREFACE

The present Ph.D. study has been conducted at the National Food Institute (previously the National Institute for Aquatic Resources) at the Technical University of Denmark. The study lasted from January 2009 to December 2011 under the supervision of Professor Lone Gram. The project included a 12 week research stay with Professor Debra Milton at the Department of Molecular Biology, Umeå University, Sweden.

The Ph.D. project was carried out as part of the collaborative research project "Discovery of novel bioactive bacteria and natural products and their use to improve human health and safety" funded by the Programme Commission on Health, Food and Welfare under the Danish Council for Strategic Research.

The work resulted in the preparation of four manuscripts which form the basis of this thesis. In addition, one genome announcement is in preparation for the Journal of Bacteriology, and collaborative efforts within the frame of the research project will likely result in a co-authorship (not included in this thesis).

Nikolaj G. Vynne

December 2011

SUMMARY

The purpose of this Ph.D. project was to evaluate a global collection of marine *Pseudoalteromonas* bacteria as a source of novel bioactive compounds, and to investigate the distribution and production of such compounds among different species within the *Pseudoalteromonas* genus. The strain collection was obtained during the research cruise "Galathea 3", which circumnavigated the Earth while screening marine bacteria for the ability to inhibit *Vibrio anguillarum* 90-11-287. *Pseudoalteromonas* strains were one of the most frequently isolated genera.

The *Pseudoalteromonas* strains were evaluated for their ability to repeatedly inhibit the fish pathogen *Vibrio anguillarum* 90-11-287 or *Staphylococcus aureus* 8325. Based on previous work, a hypothesis that antagonistic *Pseudoalteromonas* strains primarily were pigmented and surface associated was investigated. This Ph.D. work confirmed that surface-associated strains were significantly more likely to possess stable antibacterial activity and be pigmented. *Pseudoalteromonas* strains are known as prolific producers of bioactive secondary metabolites; hence screening the global strain collection for production of novel antibiotics was initiated. Novel quinolone-related compounds were described, but were not antibacterial. Several antibacterial compounds known from other sources were identified, for instance indolmycin which was hitherto only known from terrestrial *Streptomycetes*. Genome sequencing of *P. luteoviolacea* S4054 revealed up to 11 biosynthetic pathways with unknown products, confirming the potential for discovery of new secondary metabolites from *Pseudoalteromonas* strains.

The elaborate secondary metabolite production led me to speculate whether it was possible to use secondary metabolites to assist in species identification within this genus. This would also provide information on the use of 16S rRNA gene sequences to dereplicate strain collections in biodiscovery efforts. A phylogenetic study of 16S rRNA gene sequences of the *Pseudoalteromonas* strains confirmed the division into two clades; one consisted of bioactive pigmented strains and one predominantly of inactive non-pigmented strains. Correlating this to a dendrogram based on the secondary metabolites in each strain showed that some strains clustered together in a species-specific way, whereas other strains did not cluster near strains of the same species. Hence, secondary metabolite production was not unequivocally reflected in the secondary metabolite profile, possibly due to the limited resolving power of the 16S rRNA gene. The species *P. luteoviolacea* showed an interesting pattern indicative of phylotype specific antibiotic production strains. Detailed phylogenetic analysis of an expanded collection of *P. luteoviolacea* strains showed confirmed that production of antibiotics was related to phylogeny within this species, which indicates that

the underlying biosynthetic pathways are maintained under selective pressure and hence are important traits for the organism.

One strain stood out during work with the strain collection, in part because of its production of an intense black pigment in contrast to its phylogenetic placement within the non-pigmented clade. This strain was subsequently shown to represent a new bacterial species named *Pseudoalteromonas galatheae*.

Initial studies revealed the potential production of regulatory compounds involved in cell to cell signaling within some strains of the species *P. luteoviolacea*. Since such mechanisms are known to govern antibiotic production in some bacteria, this was investigated. A quorum sensing system controlling a putative novel biosynthetic pathway with high homology to the *lux* system of *Vibrio fischeri* was identified in *P. luteoviolacea* S4054. The signal molecule was potentially a new acylated homoserine lactone (AHL) like compound, and the AHL synthethase was phylogenetically distinct from related synthethases. This expands our knowledge of bacterial signaling and *lux* homologue system, and further work will resolve is this system has implications for antibiotic production.

In summary, this Ph.D. work explored the phylogeny and chemical diversity of the genus *Pseudoalteromonas*. Novel compounds were discovered but they possessed no antibiotic activity. However, analysis of the genome sequence of *P. luteoviolacea* S4054 revealed genetic potential for discovery of secondary metabolites not known within this species. Secondary metabolites were not unequivocally representative of species assignments, but on an intra-species level the use of detailed phylogenetic analysis showed phylotype specific production of antibiotics within the species *P. luteoviolacea*. These findings validate the genus *Pseudoalteromonas* as a potential source of novel secondary metabolites and may be useful when designing future biodiscovery strategies. The novel species *P. galatheae* was described which contributed to resolving the taxonomy of the genus. This thesis also provides evidence of a quorum sensing system related to the *lux* system of *Vibrio fischeri* but relying on putatively novel signaling molecule encoded by a distinct synthethase, which might be involved in the regulation of antibiotics production.

Resumé

Formålet med dette Ph.D. studie var at udforske det bioaktive potentiale af indsamlede marine *Pseudoalteormonas* bakterier, samt at kortlægge hvorledes produktionen af bioaktive stoffer er fordelt imellem de enkelte *Pseudoalteromonas* arter. Stammesamlingen blev grundlagt under en jordomsejling med forskningsekspeditionen "Galathea 3", hvor marine bakterier blev screenet for hæmmende effekt mod *Vibrio anguillarum* 90-11-287. *Pseudoalteromonas* stammer udgjorde en af de primære slægter, der blev isoleret.

Pseudoalteromonas stammerne blev testet for hæmmende effekt mod *Vibrio anguillarum* 90-11-287 og *Staphylococcus aureus* 8325. Baseret på tidligere arbejder udviklede jeg den hypotese at *Pseudoalteromonas* stammer der udviser en stabil hæmmende effekt oftere er pigmenterede eller isoleret fra et overfladeskrab. Det blev bekræftet via en statistisk analyse af de indsamlede bakterier. Det er kendt at *Pseudoalteromonas* stammer producerer bioaktive sekundære metabolitter, og på den baggrund igangsatte jeg en screening af de indsamlede *Pseudoalteromonas* stammer for at identificere nye stoffer med antibakteriel aktivitet. Denne screening resulterede i identifikation af ukendte quinolon-relatedere stoffer, men de var ikke antibakterielle. Derimod blev der isoleret allerede kendte antibiotika, f.eks. indolmycin som hidtil udelukkende var fundet i *Streptomyces* bakterie isoleret fra jord. Genomsekventering af *P. luteoviolacea* S4054 afslørede op til 11 biologiske pathways med ukendte slutprodukter der kunne være involverede i syntesen af sekundære metabolitter, hvilket bekræfter potentialet for opdagelse af nye stoffer fra *Pseudoalteromonas* bakterier.

Den omfattende produktion af sekundære metabolitter blandt *Pseudoalteromonas* stammerne førte til den ide at de sekundære metabolitter måske kunne bruges som et redskab til at adskille de forskellige arter inden for slægten. Dette ville også afklare om 16S rRNA gensekvenser kan bruges som et redskab til dereplikering af bakteriestammer i forbindelse med screening for sekundære metabolitter. En fylogeni baseret på 16S rRNA gensekvenser viste, som ventet baseret på tidligere studier, at stammerne opdeltes i to grupper. Den ene gruppe indeholdt hæmmende pigmenterede stammer, mens den anden gruppe primært indeholdt ikke-pigmenterede stammer uden hæmmende aktivitet. Ved sammenligning med et dendrogram basereret på produktion af sekundære metabolitter kunne det konstateres at nogle arter havde en ensartet sekundær metabolitprofil, mens andre arter producerede forskellige sekundære metabolitter alt efter hvilken stamme der blev undersøgt. Derfor kunne 16S rRNA gensekvenser ikke bruges som pålidelig metode til dereplikation af bioaktive *Pseudoalteromonas* stammer. Arten *P. luteoviolacea* udviste dog et interessant mønster der tydede på en overensstemmelse med fylogenetisk opdeling og produktion af sekundære metabolitter. En detaljeret fylogenetisk analyse baseret på recombinase A og gyrase B generne afslørede at produktionen af to forskellige antibiotika var begrænset til hver sin fylogenetiske gruppering. Dette tyder på at generne der er involveret i biosyntesen af disse antibiotika bliver bevaret under et selektivt pres, som betyder at antibiotikaproduktionen er et vigtigt træk for organismen.

Under arbejdet med stammesamlingen stod én stamme ud på grund af dens karakteristiske sorte pigment, som den producerede på trods af at fylogenetisk analyse placerede stammen i den ikke-pigmenterede gruppe. Denne stamme skulle vise sig at repræsentere en ny bakterieart som blev nanvgivet *Pseudoalteromonas galatheae*.

Indledende studier viste, at visse *P. luteoviolacea* stammer producerede stoffer, der potentielt var involveret i kommunikation imellem cellerne. Sådanne mekanismer regulerer produktionen af antibiotika i visse andre bakterier, og blev derfor undersøgt nærmere i *P. luteoviolacea* S4054. Et quorum sensing system med stor lighed til *lux* systemet fra *Vibrio fischeri* blev identificeret. Selve signalmolekylet tegner til at være en ukendt acyleret homoserin lakton (AHL) analog, og i en fylogenetisk undersøgelse adskilte AHL syntasen PluI sig tidligt fra de nærmest beslægtede proteiner fra *Vibrio* bakterier, hvilket tyder på at det er en hidtil ukendt AHL syntase. Dette styrker vor viden om bakterielle signalsystemer, og videre arbejde vil klarlægge hvorvidt dette system har betydning for antibiotikaproduktionen i *P. luteoviolacea* S4054.

For at opsummere har dette Ph.D. studie udforsket fylogeni og kemisk diversitet i slægten *Pseudoalteromonas*. Ingen nye antibiotika blev identificeret, men nye stoffer blev opdaget og det genetiske potentiale af *P. luteoviolacea* S4054 tyder Sekundære metabolitter var ikke universelt brugbare til artsidentifikation, men inden for arten *P. luteoviolacea* afspejledes produktionen af antibiotika i den fylogenetiske opdeling. Disse opdagelser tyder på at *Pseudoalteromonas* stammer er lovende mål for opdagelse af hidtil ukendte stoffer med biologisk aktivitet, og detaljeret fylogenetisk opklaring kan bruges som et værktøj i opdagelsen af sådanne nye stoffer. Den nye art *P. galatheae* blev beskrevet hvilket bidrog til en afklaring af taxonomien inden for slægten *Pseudoalteromonas*. Under dette Ph.D. arbejde blev det opdaget at *P. luteoviolacea* S4054 besad et quorum sensing system, der potentielt regulerer produktionen af antibiotika. Quorum sensing systemet havde ligheder med det *lux* system, der kendes fra *V. fischeri*, men de enkelte komponenter var ikke fuldt ud identiske og repræsenterer muligvis nye varianter af AHL syntase og AHL signal molekyler.

TABLE OF CONTENTS

Preface	I
Summary	II
Resumé	IV
Table of contents	VI
List of articles	

1.0 Introduction and background of the Ph.D. project1
1.0.1 Scope of the PhD project3
2.0 Bioactive secondary metabolites from marine bacteria5
2.0.1 The marine environment as a source of novel microbial and chemical diversity7
2.0.2 Biosynthesis of bacterial secondary metabolites8
2.1 Strategies for discovery of antibacterial compounds from marine bacteria12
2.1.1 Screening for <i>Pseudoalteromonas</i> strains in the Galathea 3 project12
2.1.2 Culturability and the influence of culture conditions on discovery of secondary metabolites from
marine bacteria14
2.1.3 Genomics and metagenomics as tools in biodiscovery: A marine perspective
2.2 Bacterial taxonomy in biodiscovery19
2.3 Conclusions from chapter 121
3.0 The genus <i>Pseudoalteromonas</i> – ecology and bioactivity
3.1 <i>Pseudoalteromonas</i> phylogeny and taxonomy23
3.2 Ecology of Pseudoalteromonas27
3.2.1 <i>Pseudoalteromonas</i> species interact with higher eukaryotes
3.3 <i>Pseudoalteromonas</i> as probiotics in aquaculture
3.4 Pathogenicity of Pseudoalteromonads31
3.5 Bioactive compounds produced by <i>Pseudoalteromonas</i>
3.5.1 Non-luteoviolacea Pseudoalteromonas33

3.5.2 Pseudoalteromonas luteoviolacea
3.6 Enzymatic activities of Pseudoalteromonads41
3.7 Anti-fouling potential of <i>Pseudoalteromonas</i> species43
3.8 Conclusions from chapter 244
4.0 Quorum sensing in regulation of bacterial production of antimicrobial compounds45
4.1 The Vibrio fischeri LuxI/LuxR quorum sensing model45
4.1.1 Acylated homoserine-lactones: Structure and analysis47
4.2 The role of quorum sensing in regulating bacterial production of antibiotics
4.3 Quorum sensing in <i>Pseudoalteromonas</i> 52
4.4 Conclusion from chapter 354
5.0 Conclusion
5.0.1 Bioactive secondary metabolites from marine bacteria
5.0.2 The genus <i>Pseudoalteromonas</i> – ecology and bioactivity55
5.0.3 Quorum sensing in regulation of bacterial production of antimicrobial compounds
5.1 Outlook
6.0 Acknowledgements
7.0 References

LIST OF ARTICLES

- Vynne N, Månsson M, Nielsen KF, Gram L (2011) Bioactivity, chemical profiling, and 16S rRNAbased phylogeny *of Pseudoalteromonas* strains collected on a global research cruise. Mar. Biotechnol. 13: 1062-1073.
- Vynne N, Månsson M, Nielsen KF, Gram L (2012) Antibiotic chemoprofile and RecA and GyrB sequence analyses reveal three phylotypes within the marine *Pseudoalteromonas luteoviolacea*.
 Manuscript submitted to Applied and Environmental Microbiology.
- 3 Vynne N, Christiansen G, Gram L (2012) *Pseudoalteromonas galatheae* S3431 sp. nov., a novel *Pseudoalteromonas* species isolated from a deep-sea polychaete. Manuscript in preparation.
- 4 Vynne N, Månsson M, Nielsen KF, Milton D, Gram L (2012) Production of a putatively novel acylated homoserine lactone-like quorum sensing activating molecule by *Pseudoalteromonas luteoviolacea* S4054. Manuscript in preparation.

Non peer reviewed publications:

Vynne N, Gram L (2012) Draft genome sequence of *Pseudoalteromonas luteoviolacea* S4054, an indolmycin producing marine bacterium. In preparation.

Bioactivity and phylogeny of the marine bacterial genus *Pseudoalteromonas*

Ph.D. thesis

by

Nikolaj Grønnegaard Vynne

2011

Technical University of Denmark

National Food Institute

Division of Industrial Food Research

1.0 INTRODUCTION AND BACKGROUND OF THE PH.D. PROJECT

The increase in hospital and community acquired infections caused by drug resistant bacteria is currently making an impact on societies worldwide. Developing new antibiotics is of the outmost importance, and searching for antibiotics that have a novel mechanism of action or are structurally distinct from those currently in use is one startegy to combat the spread of existing resistance mechanisms. Unfortunately, a combination of factors led to a decline in development of antibiotics, beginning in the late 1960'ies and lasting almost 40 years (Fischbach & Walsh, 2009). In this so called innovation gap, no new classes of antibiotics were introduced (figure 1). In fact, 73% of the approved antibiotics between 1981 and 2005 belonged to just three antibiotic classes: β -lactams, macrolides and quinolones (Hughes & Fenical, 2010;Newman & Cragg, 2007).



Figure 1: An illustration of the innovation gap from 1962 to 2000. No new classes of antibiotics were developed, leading to an almost empty drug pipeline when resistance against known antibiotic classes emerged. Adapted from Fischbach *et al.* (2009).

Part of the reason for this lack of new antibiotic classes is the failure of combinatorial chemistry to deliver a significant number of lead compounds to the pharmaceutical development pipeline (Newman & Cragg, 2007) and the low profitability of developing antibiotics compared to classic blockbuster drugs within e.g. cancer or metabolic conditions (Malik, 2008;Nathan & Goldberg, 2005).

The marine environment is considered an underexplored source of novel bioactive natural products (Fenical, 1993;Gulder & Moore, 2009;Imhoff *et al.*, 2011;Molinski *et al.*, 2008) and the structural diversity of marine natural products (Newman & Cragg, 2007) make them a promising target for discovery of novel antibiotics. The potential for discovery of structurally diverse novel antibiotics is enormous considering oceans cover 70% of Earth and even the marine Polar Regions are teeming with life. Antagonistic compounds are widely believed to play important roles in microbial ecology, for instance through chemical warfare among bacteria or as weapons in preypredator interactions. The dual role of antibiotics in human therapeutics and microbial interactions is fascinating, and we are only just beginning to understand the context of this within research areas such as development of antibiotic resistance, microbial signaling, bacteria-eukaryote interactions and microbial community dynamics.

During the research expedition 'Galathea 3', the project 'Stars of the ocean' isolated antagonistic marine bacteria. The majority of the isolated bacterial strains were identified as species of *Vibrio*, *Pseudoalteromonas* or the *Roseobacter* clade. (Gram *et al.*, 2010). A subsequent grant from the Strategic Research Council for Food, Health and Welfare allowed the establishment of the project "Discovery of novel bioactive bacteria and natural products and their use to improve human health and safety" involving two research groups from the Technical University of Denmark (DTU) and three from University of Copenhagen (KU). The main research areas were divided, with work at the National Food Institute focused on cultivation and ecology of marine bacteria and screening for antagonistic compounds, DTU Systems Biology focused on purification and structure elucidation of bioactive compounds, KU-LIFE focused on virulence inhibition in *Staphylococcus aureus* by marine natural products and interactions of marine bacterial metabolites with the human immune system, and KU-Health focused on marine natural products in interference with quorum sensing.

1.0.1 SCOPE OF THE PH.D. PROJECT

This Ph.D. work targeted the group of strains isolated during the 'Galathea 3'-expedition which belong to the marine Pseudoalteromonas genus. Pseudoalteromonads are known to frequently be bioactive (Bowman, 2007), and one of my primary aims was to screen the global collection of Pseudoalteromonas strains for antagonistic activity in order to isolate novel antibiotics. Bioactive Pseudoalteromonas strains are often found in association with higher eukaryotes or marine surfaces (Holmström & Kjelleberg, 1999), and we observed that production of antibacterial compounds often co-occurred with pigmentation. The nature of the strain collection allowed me to address the hypothesis that antagonistic *Pseudoalteromonas* are predominantly pigmented and surface associated (Vynne et al., 2011) using statistical tools. The bioactive potential of Pseudoalteromonads is well known (Bowman, 2007), hence screening a diverse global collection of *Pseudoalteromonas* strains for production of novel secondary metabolites might result in interesting compounds with e.g. antibiotic activity and provide an overview of the chemical diversity of the genus. Additionally, I hypothesized that secondary metabolites might be useful taxonomic markers to assist in species discrimination among Pseudoalteromonas strains since the majority of bioactive strains form a distinct clade when subjected to phylogentic analysis of 16S rRNA gene sequences (Gauthier et al., 1995). This might also facilitate biodiscovery strategies by providing an overview of the distribution of compounds among Pseudoalteromonas species. In working with the strain collection, one strain stood out in particular due to its production of an intense black pigment. This strain was identified as a new species; Pseudoalteromonas galatheae (Vynne et al., 2012a). During the screening for novel antibiotics I observed that four P. luteoviolacea strains based 16S rRNA phylogeny divided into two groups, each producing the common bioactive violacein and a specific antibiotic for the phylogenetic group in question. Based on these observations and studies of species specific production of secondary metabolites in the Actinomycete genus Salinispora (Jensen et al., 2007), I developed a hypothesis that phylogenetic reconstruction of the evolutionary relationship among Pseudoalteromonas strain could be used to guide biodiscovery efforts at the sub-species level and avoid characterization of overlapping secondary metabolite profiles. This I showed within the species Pseudoalteromonas luteoviolacea (Vynne et al., 2012b). The strain P. luteoviolacea S4054 produced molecules capable of activating several quorum sensing monitor strains. The presence of a putative cell-density dependant

signaling system could suggest that one or more interesting phenotypes such as antibiotic production were controlled in this manner, which I investigated experimentally. The work on cell signaling is not yet concluded, however I have shown that a signaling system is present in *P. luteoviolacea* and that it might regulate a novel biosynthetic pathway (Vynne *et al.*, 2012c). The signaling molecules are AHL-like and may also be novel. Finally, the potential for secondary metabolite production was assessed based on genome analysis, which indicated that strain S4054 has the potential to produce several so far unknown compounds (Vynne & Gram, 2012).

This Ph.D. thesis contains an overview of drug discovery strategies within marine bacteria and how efforts in microbial ecology and drug discovery may benefit from interacting (chapter 2). Chapter 3 describes the ecology, industrial applications and biodiscovery potential of the marine bacterial genus *Pseudoalteromonas*, and chapter 4 presents an overview of quorum sensing and the relevance of quorum sensing in drug discovery.

2.0 BIOACTIVE SECONDARY METABOLITES FROM MARINE BACTERIA

Bacterial secondary metabolites have historically played a central role in discovery of novel pharmaceutical lead compounds. In particular, terrestrial Actinobacteria have been a major source of biologically active secondary metabolites and most antibiotics in use today are either natural products or natural product derived (Newman & Cragg, 2007). Among the success stories are the discovery of streptomycin and tetracycline, with tetracycline and synthetic derivatives still being backbone drugs in treatment of bacterial infections when resistance is not present. Recent years have seen a dramatic increase in the discovery rate of marine natural products, many of which originate from marine microorganisms (figure 2). As we learn that marine eukaryotes may harbor distinct and species specific bacterial epibionts, it is becoming evident that some chemical compounds previously attributed to the macroorganism are in fact produced by epibiont bacteria. The discovery and origin of bryostatins, currently undergoing clinical trials as an anti-cancer agent, exemplifies this. Bryostatin 1, currently in phase I clinical trials against cancer, was originally isolated from samples of the marine bryozoan *Bugula neritina* and was immediately recognized for



Figure 2: Number of new marine natural compounds discovered broken down by phylum. The compounds discovered from 1965 to 2005 were summed up and averaged. Adapted from Blunt *et al.* (2009)

its potent anticancer activity (Pettit *et al.*, 1982). Initially thought to be a product of the bryozoan animal, it was later revealed that the biosynthetic genes were actually present in the bacterial symbiont "*Candidatus* Endobugula sertula" (Sudek *et al.*, 2006) but not in the bryozoan. Production of bryostatins was also reduced after an antibiotic treatment and accompanying reduction in the number of bacterial symbionts (Sudek *et al.*, 2006) offering strong evidence that the bryostatins are in fact of bacterial origin. The marked increase in discovery rates of microbially produced marine natural products likely in part due to an increased awareness of microbial epibionts as potential sources of bioactive metabolites (Mearns-Spragg *et al.*, 1998;Penesyan *et al.*, 2009;Sudek *et al.*, 2006), but it is also an acknowledgement of the importance of marine microorganisms as a promising source of biologically active molecules. In 2010 the marine drugs pharmaceutical pipeline contained twelve drugs in phase I to III clinical trials (table 1) mainly targeting cancer, and four approved drugs (Mayer *et al.*, 2010).

Clinical			Marine	
status	Compound name	Chemical class	organism	Disease area
Approved	Cytarabine, Ara-C	Nucleoside	Sponge	Cancer
	Viderabine, Ara-A	Nucleoside	Sponge	Antiviral
	Ziconotide	Peptide	Cone snail	Pain
	Trabectidin (ET-743) (EU registered only)	Alkaloid	Tunicate	Cancer
Phase III	Eribulin Mesylate (E7389)	Macrolide	Sponge	Cancer
	Soblidotin (TZT 1027)	Peptide	Bacterium	Cancer
Phase II	DMXBA (GTS-21)	Alkaloid	Worm	Cognition, Schizophrenia
	Plinabulin (NPI-2358)	Diketopiperazine	Fungus	Cancer
	Plitidepsin	Depsipeptide	Tunicate	Cancer
	Elisidepsin	Depsipeptide	Mollusc	Cancer
	PM1004	Alkaloid	Nudibranch	Cancer
	Tasidotin, Synthadotin (ILX-651)	Peptide	Bacterium	Cancer
	Pseudopterosins	Diterpene glycoside	Soft coral	Wound healing
Phase I	Bryostatin 1	Polyketide	Bryozoa*	Cancer
	Hemiasterline (E7974)	Tripeptide	Sponge	Cancer
	Marizomib (Salinosporamide A; NPI-0052)	β-lactone-γ-lactam	Bacterium	Cancer

Table 1: The status of the marine pharmaceutical pipeline, 2010. Modified from Mayer et al. (2010).

*: Evidence of bacterial origin, see text.

Among the advantages of marine drug leads is a hypothesis that compounds of marine origin are very potent due to the inevitable dilution that occurs in this environment, and the structural diversity of the molecules is stunning (Grabowski *et al.*, 2008). An additional distinct advantage of pursuing bacterial metabolites as drug leads is that, presumably, biosynthetic pathways of bacterial origin are more amenable to heterologous recombinant expression in for instance *E. coli*, which has the potential to eliminate the very real problem of a stable, cost-efficient supply of compound for tests and clinical trials (Molinski *et al.*, 2008). A recent success of biodiscovery within marine bacteria is the compound salinosporamide A. Salinosporamide A is a potent 20S proteasome inhibitor in clinical trials as anti-cancer agent (Feling *et al.*, 2003). It was discovered from a novel streptomycete and the development process included several valuable proof-ofconcept milestones for marine drugs, for instance the implementation of a seawater based fermentation process in a cGMP setting (Fenical *et al.*, 2009).

2.0.1 THE MARINE ENVIRONMENT AS A SOURCE OF NOVEL MICROBIAL AND CHEMICAL DIVERSITY

In the past, a common view was that the oceans were depleted in nutrients and presented harsh environmental parameters such as high salinity and a changing pH, which would not allow for microbial growth (MacLeod, 1965). This perception has changed, and today the marine environment is known to be teeming with microbial life considered a promising source of novel microbial and chemical diversity (Delong, 2007). A high diversity of marine bacteria are often found in marine surface associated microbial communities (Holmström et al., 2002;Webster et al., 2001). Increasing evidence suggests that marine microbial communities can be highly specific to particular niches. A comparison of the surface communities of co-occurring marine eukaryotes revealed distinct bacterial communities when examined at the species level (Longford et al., 2007), which is in agreement with later studies in which bacterial communities were found to be host-specific to several algae species (Lachnit et al., 2009). The bacterial community associated with marine sponges consists of specialist bacteria specific to a certain sponge species, sponge associated bacteria that are found on several sponge species but not in seawater and generalist marine bacteria found both on sponges and in seawater (Taylor et al., 2004). Even within one host individual, the composition of the epibiotic bacteria may be spatially distinct as reported for sponges (Meyer & Kuever, 2008; Thiel et al., 2007) and algae (Staufenberger et al., 2008). The niche specificity of bacterial communities is conceptually useful when trying to understand the vast diversity of marine bacteria and the implications on natural products discovery. Antagonistic

interactions among marine bacteria are frequently observed, and bacterially produced antibiotics are believed to be involved in microbial interactions and affect the structure of microbial communities (Long & Azam, 2001; Rypien et al., 2010; Slattery et al., 2001). In a recent study attempting to clarify interaction dynamics in soil microbial communities, a large number of pairwise interactions were observed among streptomycetes isolated from individual soil grains (Vetsigian et al., 2011). Based on these interactions, the authors suggest a model in which the original microbial community is highly likely to evolve and adapt rapidly through for instance acquisition or adaptation of antibiotic producing genetic pathways, subsequently countered by the emergence of antibiotic resistance (Vetsigian et al., 2011). It stands to reason that molecules involved in such interactions may evolve as microbial communities change, and thus microbial diversity and the assembly of diverse microbial communities may drive the evolution of chemical diversity. Hence, the understanding that many antibacterial compounds are involved in microbial interactions and shaping of microbial communities combined with the knowledge that microbial communities often are niche specific ties a link from microbial ecology to discovery of novel bioactive compounds (Imhoff et al., 2011). Understanding the implications of microbial diversity and the ecology of antibiotics may be of great assistance in maximizing the output of biodiscovery efforts. Developments in analytical techniques such as MALDI-TOF-MS (Esquenazi et al., 2008) have the potential to be of great assistance in research on the spatial location and identification of secondary metabolites within complex communities which has the potential to offer new and exciting insights at the interface of microbial ecology and natural products chemistry.

2.0.2 BIOSYNTHESIS OF BACTERIAL SECONDARY METABOLITES

Bioactive natural products of bacterial origin cover an enormous structural diversity and include multiple compound classes, such as terpenes, polyketides and nonribosomal peptides. These compounds are synthesized by cellular enzymes which are often encoded by genes organized on the same locus in so called biosynthetic pathways. Two of the classes with a well understood biosynthetic pathway are polyketides (PKs) and nonribosomal peptides (NRPs), which are often isolated from marine microorganisms (Hughes & Fenical, 2010). The synthesis machineries for these two classes of compounds are conceptually alike. The active domains are contained within one large protein in a modular structure where the synthesized molecule is transferred from one domain to the next as the catalytic reactions proceed (Staunton & Weissman, 2001). The canonic type I PK biosynthetic machinery depicted in figure 3. In PK synthesis, each module mediates one extension step of the PK chain, including any modification such as reduction of a ketone to a hydroxyl. Examples of antibacterial polyketids of marine origin are shown in figure 4 below.



Figure 3: Modular organization of erythromycin polyketide biosynthesis, a PKS type I pathway. Each module (except the loading and end modules) contains the essential ketosynthase, acyltransferase and acyl carrier protein domains required for elongation of the polyketide chain, whereas additional domains alter the substitutions on the chain. Adapted from Staunton and Weissman (2001).



Figure 4: Antibacterial polyketides of marine origin. BE-43472B (1) isolated from a marine *Streptomycete*; abyssomycin C (2) isolated from the marine actinomycete *Verrucosispora* sp; pestalone (3) isolated from a marine fungus; ariakemicin A (4) isolated from a marine gliding bacterium of the phylum *Bacteroides*. Modified from Huges & Fenical (2010).

A key difference among the two pathways is the use of different building blocks; polyketide synthetases (PKSs) rely on acetate (activated to form malonyl-CoA) and propionate (activated to form methyl-malonyl-CoA) (Cortes *et al.*, 1990) whereas nonribsomal synthetases (NRPSs) utilize amino acids (Schwarzer *et al.*, 2003). Variations of these archetypical PKS molecules exist, for instance a PKS gene cluster with no intrinsic AT domain (Piel, 2002). Instead, this pathway relies on discrete AT genes to synthesize the end product pederin. Antibacterial NRPS molecules are produced by marine bacteria (figure 5), and the generalized pathway shares the modular layout with PKS pathways, however the domains encode different functionalities. Marine NRPs include violacein and indolmycin, isolated from *P. luteoviolacea* strains in this Ph.D. work, which can be recognized as NRP compounds based on their tryptophan building blocks. Hybrid pathways further add to the potential structural diversity among natural products.





The modular nature of these biosynthetic machineries has led to interest in so called combinatorial biosynthesis as reviwed by (Weissman & Leadlay, 2005), where the enzymes catalyzing the synthesis reactions are modified based on the desired outcome (Cortes *et al.*, 1995). For instance, adding enoyl reductase domain to a module can eliminate a functional group and change the steric configuration of the molecule, thus potentially altering the biological activity. This metabolic engineering is gradually becoming 'plug-and-play' through the availability of distinct building blocks (Menzella *et al.*, 2005). This approach opens for a virtually unlimited cache of structural scaffolds, and provides researches with the ability to directly modify sites which are identified as important for pharmacological activity.

2.1 STRATEGIES FOR DISCOVERY OF ANTIBACTERIAL COMPOUNDS FROM MARINE BACTERIA

2.1.1 SCREENING PIPELINE FOR PSEUDOALTEROMONAS STRAINS IN THE GALATHEA 3 PROJECT

In this Ph.D. work, a screening pipeline was used to facilitate efficient screening for antibacterial compounds among the bacteria isolated during the research expedition Galathea 3. Bacterial strains were isolated from a range of diverse marine samples during the global expedition (Gram *et al.*, 2010) and pre-screened for antagonism towards *Vibrio anguillarum* 90-11-287 (Skov *et al.*, 1995), a potent fish pathogen. The bacterial strains were tentatively identified via BLAST queries of partial 16S rRNA gene sequences, and one of the dominant taxa was the genus *Pseudoalteromonas* which forms the topic of this thesis. The potentially antagonistic bacterial strains were screened for inhibitory activity following a principle similar to the outline in figure 6.



Figure 6: An overview of the principles in a screening strategy such as the one used in this Ph.D. work. Adapted from Molinari (2009).

The chemical analysis pipeline includes HPLC coupled with a detector, for instance an UV/VIS diode-array-detector. The HPLC is used to separate compounds in a given mixture by size, polarity or charge, resulting in a series of fractions which can be tested in the bioassay. Once a pure compound has been isolated, LC-MS can be used to obtain the accurate mass of the compound. The accurate mass is used to calculate the elemental composition and in combination with the formation of distinct adducts during LC-MS analysis, the compounds can often be dereplicated and identified with help from database searches (Nielsen *et al.*, 2011). However, to obtain the full structural information one must resort to nuclear magnetic resonance spectroscopy or x-ray crystallography.

The main bioassay used was a simple well diffusion agar assay (Hjelm *et al.*, 2004), where a target culture is cast into an agar substrate and test substances are added to wells cut in the agar (Figure 7).



Figure 7: Well diffusion agar assay of sterile filtered *Pseudoalteromonas* culture supernatants. The inhibition of *Vibrio angullarum* is seen as clearing zones of no bacterial growth, signifying the presence of an inhibitory compound in the tested supernatant.

This bioassay provides a very direct assessment of the antibacterial potential of a given sample, and repeated cycles of chemical fractionation followed by bioassays to verify activity is a proven method of obtaining relatively pure active compounds. This approach is often referred to as bioassay guided fractionation. This principle can be used for the identification of practically any compound where a specific bioassay exists, however, as part of the work in this thesis shows (Vynne *et al.*, 2012c), complications such as different detection limits among bioassays and chemical detectors or the inability to confine the active compound to one single fraction may occur. Also, as subsequent paragraphs reveal, the genome of *P. luteoviolaceae* potentially points to a range of bioactive compounds that are not expressed under standard laboratory growth conditions.

2.1.2 CULTURABILITY AND THE INFLUENCE OF CULTURE CONDITIONS ON DISCOVERY OF SECONDARY METABOLITES FROM MARINE BACTERIA

When determining if a bacterial strain produces for example antibiotics, growing the organism in pure cultures under controlled conditions is in most cases considered essential, and this approach has been greatly successful. It offers the immediate advantages of working with a pure culture growing under defined conditions for a high degree of reproducibility. Nevertheless, the fact that only a fraction of the cells in a given sample are culturable using standard laboratory approaches and growth substrates (an estimated 0.1 % for bacterioplankton (Gram et al., 2010)) suggests that a reliance on these classic techniques may also limit the potential for discovery of novel antibiotics. To compensate for this, several authors have taken alternative approaches to cultivation. A simple modification to growth conditions in liquid cultures was developed by (Yan et al., 2002) (figure 8). This presumably allows for increased biofilm formation and reduces oxygen limitations, and has the potential advantage of decreased exposure to autoinhibitory compounds. Culturing two marine Bacillus strains in this roller bottle setup induced production of two antimicrobial compounds which inhibited an MRSA test strain (Yan et al., 2002). Similarly, growth in biofilms may induce or enhance production of antibiotics as demonstrated for Phaeobacter 27-4 (Bruhn et al., 2007). This illustrates how important the physical conditions can be to antibiotic production.



Figure 8: Agar coated roller bottle design. The bottle rotates at 1 rpm, and bacterial biofilms form on the agar coated interior walls. The thickened portion of the agar allows for aeration of the liquid substrate. Adapted from Yan *et al.* (2002).

Laboratory growth substrates are often based on either a complex mix of carbohydrates and nitrogen designed to support fast bacterial growth, for instance yeast extract or peptone, or defined minimal substrates designed for selective enrichment of bacterial cultures or to support fastidious organisms. In the case of marine bacteria, neither substrate is likely to reflect the natural niche very and thus may not induce the production of key secondary metabolites. A recent study elegantly demonstrated how growth substrates may impact secondary metabolite production: A *Vibrio corallilyticus* strain shut down production of all secondary metabolites but the antibiotic andrimid when cultured on chitin or live chitinous *Artemia* instead of marine broth 2216 (Wietz *et al.*, 2011). This validates the use of growth substrates mimicking the natural nutrient regime, and is an excellent illustration of how microbial ecology and biodiscovery may go hand in hand. Preliminary results obtained during this Ph.D. work of *P. luteoviolacea* cultures grown on alga extracts suggests this strategy may indeed change the metabolic profile of the bacterium, and cultures grown in the complex growth substrate heart infusion broth supplemented with 2% sodium chloride caused a heavy production of extracellular polymers in the strain *P. luteoviolacea* S4054 (unpublished data).

Elicitation of antibiotic production may be achieved by co-culturing the producer organism with other microorganisms in an effort to simulate environmental parameters from the natural environment, such as signaling and stress factors. One of the earliest studies involving marine bacteria demonstrated how marine bacteria exposed to cells of *S. aureus* or *Pseudomonas aeruginosa* responded with increased antibacterial activity compared to non-exposed controls

(Mearns-Spragg et al., 1998). Similar results were obtained for marine sponge-associated bacteria (Kanagasabhapathy & Nagata, 2008) and in a marine streptomycete (Slattery et al., 2001). Three of four tested tropical marine epibiotic bacteria showed antifungal activity when exposed to live cells of the fungus for 24 h, whereas pure cultures of the epibiotic bacteria were not antifungal (Dusane *et al.*, 2011). This is a logical extension of the hypothesis that antibiotics have ecological significance, and conceivably are produced as chemical weapons in response to being challenged by foreign competitor cells. However, in some cases the problem itself is accessing the secondary metabolism of a so called unculturable microorganism. The use of high-throughput microtiter plate techniques, dilution methods and a growth substrate containing natural levels of nutrients led to the culturing of 14% of the cells in a marine water sample (Connon & Giovannoni, 2002), compared to the ca. 0.1% normally achieved (Gram et al., 2010). Among the cultures were several novel Proteobacteria clades. One of these clades was related to the SAR11 lineage, which is considered ubiquitous in the ocean surface layer and may contribute up to 50% of the total bacterioplankton cell numbers (Morris et al., 2002). SAR11 does not grow at standard laboratory techniques, but the use of the high-throughput, low nutrient technique allowed the successful culturing of SAR11 bacterial strains (Rappe et al., 2002) and subsequent improvements to the technique have allowed the culturing of more diverse novel SAR11 strains (Stingl et al., 2007). Other approaches include the use of various growth factors (D'Onofrio et al., 2010; Nichols et al., 2008), diffusion chambers (Bollmann et al., 2007) and membrane systems (Ferrari et al., 2008). Together, these innovative approaches to the fundamental microbiological task of culturing bacteria have opened new possibilities for culturing parts of the previously rarely isolated or unculturable majority of bacteria (Sogin et al., 2006), which allows access to practically unlimited diversity.

2.1.3 GENOMICS AND METAGENOMICS AS TOOLS IN BIODISCOVERY: A MARINE PERSPECTIVE

As the first bacterial genomes of prolific secondary metabolite producers were completed, it was observed that they contained gene clusters with homology to known biosynthetic pathways, but to which no known secondary metabolite could be associated. Although advanced culture techniques may activate some of these silent clusters one faces a daunting task if attempting to

elicit every biosynthetic pathway without further knowledge as for instance a genome sequence. With sequencing technologies rapidly advancing and sequencing costs dropping equally fast, genome and metagenome sequencing has become accessible to most labs in terms of both costs and the required expertise. Excellent software exists to aid in handling the large amounts of data generated by these approaches, and the modular nature of PKS and NRPS gene clusters makes in silico prediction of homologue loci an exciting opportunity. In fact, genome sequencing and analysis has already proven itself in discovery of previously non-described compounds from otherwise thoroughly studied bacteria. Prediction of novel secondary metabolites from such silent gene clusters are known from Streptomyces coelicolor (Challis & Ravel, 2000), where a silent NRPS pathway was shown to produce the novel iron chelator coelichelin under iron-limited growth conditions (Lautru et al., 2005). One of the first reports on secondary metabolites from thermophilic bacteria was based on genome analysis and identification of an orphan NRPS gene cluster in the thermophilic actinomycete *Thermobifida fusca*. The biosynthetic pathway showed unusual features including non-linear peptide assembly, and the end product was a siderophore molecule possessing a distinct novel iron chelation domain (Dimise et al., 2008). In the genome of the marine bacterium Pseudoalteromonas tunicata genome, a biosynthetic gene cluster was predicted based on the presence of an ATP-grasp domain (Blasiak & Clardy, 2009). The putative biosynthetic gene cluster, shown in figure 9, was then PCR amplified and cloned into a vector under control of an inducible promoter, and transformed into an E. coli host. Induction of the promoter resulted in the production and subsequent identification of two 3-formyl-tyrosine metabolites which were not previously identified in the otherwise well-studied P. tunicata strain D2 – possibly due to their lack of antibacterial activity.



Figure 9: Predicted biosynthetic gene cluster from the *P. tunicata* D2 genome. 1; structure of 3-formyl-Ltyrosine-L-threonine dipeptide, one of the discovered metabolites. Color codes indicate the proposed biosynthetic pathway. Adapted from Blasiak and Clardy (2010).

In this Ph.D. work, the genome sequence of *P. luteoviolacea* S4054 revealed between 7 (manually curated annotation of PKS and NRPS containing gene clusters in IMG/ER (Vynne & Gram, 2012)) and 14 (antiSMASH prediction (Medema *et al.*, 2011)) potential biosynthetic pathways, one of which was the violacein pathway. The violacein pathway is known (Antônio & Creczynski-Pasa, 2004;August *et al.*, 2000;Hoshino, 2011) and can be identified through homology searches, but none of the remaining potential biosynthetic pathways were homologue to pathways encoding known secondary metabolites. The antibiotics indolmycin and pentabromopseudilin were identified in *P. luteoviolacea* S4054 during this Ph.D. work, and their biosynthetic pathways are likely among the predicted gene clusters. However, since the genes involved in the biosynthesis are unknown, no pathway could be directly assigned to these compounds. The presence of up to 14 novel biosynthetic pathways opens the possibility that *P. luteoviolacea* S4054 may produce up to 11 as yet unknown secondary metabolites, highlighting the power of genomic analysis and the potential for genome assisted biodiscovery in marine *Pseudoalteromonas*.

Metagenomics have shown a potential as a biodiscovery related technique, as evidenced by the Global Ocean Sampling expedition, which based on the DNA sequence data obtained predicted a multitude of unknown proteins and protein families (Yooseph *et al.*, 2007). However, a general limitation of homology based studies is the reduced chance of finding something very different from what is already known, as genetic homology does imply a structural or conceptual similarity among the end products. Hence, *in silico* genomic and metagenomic studies offer a fantastic supplement to classical culture-based approaches but cannot be relied upon as the sole means of assessing the secondary metabolite diversity of a single strain or microbial community. A more

direct approach is using metagenomic DNA directly to create libraries of clones containing large inserts of metagenomic DNA and perform functional screens for expression of e.g. antibacterial activity. In a pioneering study on functional screening of uncultured marine symbionts, a functional screen of shotgun cloned 'metagenomic' DNA (mainly *Prochloron* sp.) led to the cloning and characterization of the biosynthetic pathway of the cytotoxic compound patellamide D DOI: 10.1002/cbic.200500210. A fosmid library was used to screen metagenomic DNA from microbial communities associated with a sponge (*Cymbastela concentrica*) and an alga (*Ulva australis*) for novel antibacterials, which resulted in the identification of three novel antibacterial proteins (Yung *et al.*, 2011).

2.2 BACTERIAL TAXONOMY IN BIODISCOVERY

Proper identification and taxonomical assignment is crucial to scientific communication, also within natural product discovery (Jensen, 2010). The production of secondary metabolites is often considered to be strain specific (Waksman & Schatz, 1943), in part due to the involvement of biosynthetic gene clusters in lateral gene transfer (LGT) (Egan et al., 2001b). This would seem to exclude bacterial taxonomy as a useful tool for dereplicating bacterial strains in a biodiscovery setting, however screening of a taxonomically dereplicated collection of actinomycetes led to a high number of discovered compounds compared to the strain througput (Goodfellow & Fiedler, 2010). Studies of the secondary metabolites produced by Salinispora species showed that a core set of metabolites are restricted to specific species (Jensen et al., 2007; Penn et al., 2009) and among species of the Roseobacter clade, production of a brown pigment and the antibacterial compound tropodithietic acid was shown to be related to growth conditions in a species-specific manner (Porsby et al., 2008). Hence, species specific chemoprofiles can exist. This is valid for some filamentous fungi as well, and has been used for guided discovery of secondary metabolites (Frisvad et al., 2008). Since the available resources often limit the amount of strains that can be included, not screening strains with identical or overlapping secondary metabolite profiles would be beneficial in biodiscovery efforts. In this Ph.D. work, a comparison between a statistical clustering based on secondary metabolites and 16S rRNA gene sequence phylogeny was made to reveal the extent, if any, of species specific secondary metabolite production in the

Pseudoalteromonas genus (Vynne *et al.*, 2011). Some strains, such as *P. ruthenica* or the *P. aurantia* – *P. citrea* complex were found to cluster together as expected based on 16S rRNA sequence identity. However, some species, e.g. *P. luteoviolacea*, were split in distinct clusters by chemoprofiling and strains of *P. rubra* were very diverse from a secondary metabolite point of view. This study showed that 16S rRNA gene sequences are insufficient for reliable dereplication of secondary metabolite profiles in *Pseudoalteromonas* strains. However, in a follow up study it was demonstrated that a more detailed phylogenetic analysis of GyrB and RecA sequences were capable of resolving phylogenetic groups matching the observed distribution of bioactive secondary metabolites (figure 10) (Vynne *et al.*, 2012b). In addition, the strains that produced the antibiotic indolmycin were exclusively isolated from algae, suggesting an ecology aspect is involved in production of this compound. The fixation of secondary metabolites within phylogenetic clusters is a strong indication that the biosynthetic gene clusters are under selective pressure and thus play an important functional role (Jain *et al.*, 1999), which further supports the importance of antagonistic compounds in microbial ecology. Furthermore, the detailed phylogeny could be used as a tool for dereplication of bacterial strains in biodiscovery.





0,002

2.3 CONCLUSIONS FROM CHAPTER 2

Marine bacteria are a proverbial 'treasure trove' of bioactive natural compounds, many of which are likely to possess interesting pharmacological activities. Advances in microbial ecology may be of great use within biodiscovery programs, as these bioactive compounds are highly likely to be involved in microbial ecology. For instance, sampling strategies could benefit from targeting specific niches if a certain biological activity was desired, or could simply aim at obtaining samples from as diverse microbial communities as possible. Once the biological material is collected, a dereplication of the bacterial strains may be carried out based on phylogeny and knowledge of microbial ecology to reduce the inclusion of overlapping secondary metabolomes. The biosynthetic potential of a *Pseudoalteromonas luteoviolacea* strain was assessed using bioinformatic approaches, and was found to contain up to 14 pathways involved in secondary metabolism. Accessing the biosynthetic potential via both culture dependent and culture independent techniques is likely to produce the best results, and modulation of growth conditions play an important role in this process.

3.0 THE GENUS *PSEUDOALTEROMONAS* – ECOLOGY AND BIOACTIVITY

The genus *Pseudoalteromonas* consists of obligate marine bacteria found ubiquitously in the marine environment. They are Gram-negative non-sporeforming straight or curved rods belonging to the *y-Proteobacteria*. The cells are motile by one or more polar flagella (Romanenko *et al.*, 2003) with some species reported to have lateral flagella in addition to polar. They are oxidase positive, vary in catalase reaction and are non-fermenting with a genomic DNA GC content of 37-50 mol% and several species are intensely pigmented (figure 11) (Gauthier *et al.*, 1995). Many *Pseudoalteromonas* show bioactivity such as antagonism towards other bacteria, hydrolytic enzymatic activities or anti-fouling activity. Pseudoalteromonads are considered obligate marine, but will grow on a range of nutrient regimes if marine salts are present. Some species, such as *P. ruthenica* can grow on NaCl as the sole salt source (Vynne, unpublished). All species grow at 20°C, with growth ranges within 4°C to 40°C. The bacteria are readily cultured on e.g. Marine Agar 2216 (Difco, USA) using routine culture techniques.



Figure 11: *Pseudoalteromonas* strains streaked on Marine Agar 2216.

Non-pigmented: *P. agarivorans*, yellow: *P. flavipulchra*, dark purple: *P. luteoviolacea* red: *P. rubra*

The first descriptions of *Pseudoalteromonas* species were published in the 1940s and 1950s when the isolates were assigned to the *Flavobacterium, Pseudomonas* or *Vibrio* genera (Bein, 1954;Berland *et al.*, 1969;ZoBell & Upham, 1944). In 1972, the genus *Alteromonas* was created to harbor marine aerobic bacteria with GC-mol% of 42.8 to 48 and a polar flagellum (Baumann *et al.*, 1972), however with the advent of 16S rRNA gene sequence analysis it was clear that the genus type strain *Alteromonas macleodii* formed a monophyletic cluster away from the remaining species in the genus (Gauthier *et al.*, 1995). To account for this discovery, the genus *Pseudoalteromonas* was created with *Pseudoalteromonas haloplanktis* as genus type strain and containing a total of 12 species. The genus now encompasses 36 species (Euzéby, 1997) including several that were recently described (Oh & Park, 2011;Xu *et al.*, 2010).

3.1 *Pseudoalteromonas* Phylogeny and Taxonomy

Understanding and navigating the taxonomy of Pseudoalteromonas species requires some attention to detail, as many species share a very high (>99%) 16S rRNA gene sequence identity. This can complicate the designation of species names to novel isolates, in particular since phenotypic traits will also often be similar and it is not practical to do DNA-DNA hybridization for each interesting novel strain. Thus, the reader is encouraged to study the phylogenetic relations among the Pseudoalteromonas type strains presented in figure 12, and keep in mind that there is likely a degree of overlap in what different authors identify as e.g. P. aurantia or P. citrea. The importance of proper species designation should not be understated. Accurate identification of a novel isolate is key in order to communicate science and allow other scientists to perform satisfactory literature searches as stressed in a recent review on marine natural products (Blunt et al., 2009). The use of 16S rRNA gene sequences to resolve bacterial taxonomy to the species level is widely accepted, cheap and fast to use. A 97% sequence identity is the accepted threshold for species delineation (Stackebrandt et al., 2002), however above 97% sequence identity does not confirm two strains as one species (Stackebrandt & Goebel, 1994). The gold standard remains DNA-DNA hybridization among species type strains where less than 70% DNA similarity is required to separate two strains into distinct species (Wayne et al., 1987), although the use of wholegenome sequence data may replace this method (Konstantinidis & Tiedje, 2005). The use of DNA-DNA hybridization, biochemical data and phylogenetic analysis of 16S rRNA genes is termed the polyphasic approach (Vandamme *et al.*, 1996).



Figure 12: Phylogenetic analysis of 16S rRNA sequences of *Pseudoalteromonas* type strains. Sequences were retrieved from GenBank, aligned in MEGA5 (Tamura *et al.*, 2011) and a phylogenetic tree inferred by maximum likelyhood analysis. Scale bar: 0.005 nucleotide substitutions per site.
In the course of this Ph.D. study, strain S3431 was identified as a potentially novel *Pseudoalteromonas* species. The strain was originally isolated from a polychate near Canal Conception in Chile and immediately caught our attention due to its deeply black pigment when cultured on marine agar 2216. Initial results of the S3431 16S rRNA gene sequence analysis showed that it had below 97% similarity to any described *Pseudoalteromonas* type strain. Further analyses of phenotypic traits and chemotaxonomic markers were conducted, and DNA-DNA hybridization was initiated. A later revision of the 16S rRNA sequence revealed sequencing errors and, after careful re-sequencing, the strain was found to have a 16S rRNA sequence identity of 99% to a number of non-pigmented *Pseudoalteromonas* type strains. However, it was decided that the preliminary results combined with the interesting pigmented phenotype was sufficient to warrant further investigations. DNA-DNA hybridizations (table 2), physiological data and biochemical tests revealed that S3431 represents a novel species given the name *Pseudoalteromonas galatheae* (Vynne *et al.*, 2012a).

Pseudoalteromonas species	Per cent DNA:DNA similarity to strain S3431
<i>P. agarivorans</i> DSM 14585^{T}	28.2
<i>P. aliena</i> LMG 22059 ^{T}	44.5
<i>P. atlantica</i> NCOMB 301^{T}	26.9
<i>P. carrageenovora</i> DSM 6820 ^T	14.8
<i>P. distincta</i> CIP 105340^{T}	44.5
<i>P. elyakovii</i> LMG 14908 [™]	44.2
<i>P. espejiana</i> DSM 9414 ^{T}	48.0
P. issachenkonii LMG19697 [™]	33.6
<i>P. nigrifaciens</i> LMG 2227 ^{T}	15.4
<i>P. paragorgicola</i> LMG 19696 ^T	58.4
<i>P. tetraodonis</i> DSM 9166 [™]	11.3
P.undina LMG 2880 ^T	13.8

Table 2: Per cent DNA-DNA similarity in 2X SSC at 66°C (Vynne *et al.*, 2012a). All related type strains have less than 70% DNA-DNA similarity to S3431, which is the accepted threshold for species delineation (Wayne *et al.*, 1987).

Unfortunately, the antibacterial effect that the strain was originally isolated for turned out to not be reproducible upon frozen storage and repeated culture in the lab. This is not uncommon, and may be due to several factors such as a need for specific nutrients or a lack of interaction with for instance co-cultured microorganisms. As described above, using culture conditions closer to the natural environment (C-source, associate microbiota) may re-evoke the production of antibacterial activity.

Figure 12 above reveals a characteristic feature of the genus, namely the phylogenetic division into two groups when 16S rRNA gene sequences of the type strains are analysed (Gauthier *et al.*, 1995;Ivanova *et al.*, 2004). One group predominantly consists of non-pigmented bacteria with little antibacterial activity while the majority of the other group is pigmented bacteria typically isolated from marine biotic surfaces. This latter group often produce compounds involved in microbial antagonism (Mikhailov *et al.*, 2002) and correlation between pigmentation and antifouling activity was reported for *P. tunicata* and *Pseudoalteromonas* sf57 (Huang *et al.*, 2011) where anti-fouling activity was lost in non-pigmented mutants. It is important to note that bioactivity is not exclusively linked to pigmentation as recently shown by Bernbom *et al.* (2011). A more comprehensive review is presented in the paragraph on bioactivity in *Pseudoalteromonas*.



Figure 13: Isolation sites of 101 *Pseudoalteromonas* strains with each color code representing one of the 8 clusters indicated in Table 1 in Vynne *et al.* (2011)

The present work investigated 101 novel environmental *Pseudoalteromonas* isolated globally from a range of marine sources (figure 13), and whilst all isolates in initial screenings displayed some

degree of antibacterial activity, reproducible inhibition of *Vibrio anguillarum* occurred significantly more frequent among pigmented strains (table 3). Analysis of 16S rRNA gene sequences supported the phylogenetic grouping of pigmented and non-pigmented strains respectively (Vynne *et al.*, 2011).

Table 3: Identity, antibacterial activity and pigmentation of *Pseudoalteromonas* strains froma global collection. Modified from Vynne *et al.* (2011)

ster	No o	f strains		No of fr	strains om	No of inhibitir	strains ng <i>Vibrio</i>	ct
16S rRNA clu	pigmented	non- pigmented	Related type strain	Surface	Water	pigmented	non- pigmented	<i>Vibrio</i> inhibit by EtAc extra
I	1	37	P.agarivorans	16	22	0	0	0
II	5	0	P.aurantia	2	3	5	0	0
	0	9	P.prydzensis	8	1	0	0	0
IV	3	3	P.phenolica	3	3	3	2	0
V	4	0	P.luteoviolacea	4	0	4	0	4
VI	9	0	P.rubra	9	0	9	0	0
VII	13	0	P.flavipulchra	13	0	13	0	0
VIII	15	0	P.ruthenica	14	1	15	0	15
S1712	0	1	P.mariniglutinosa	0	1	0	0	0
S3655	1	0	P.spongiae	1	0	0	0	0
Total	51	50		70	31	49	2	19

3.2 ECOLOGY OF PSEUDOALTEROMONAS

Pseudoalteromonas bacterial cells constitute 0.5-6% of the oceanic bacterioplancton (Wietz *et al.*, 2010) and 0.01%-2.1% of surface associated bacteria on a selection of marine surfaces (Skovhus *et al.*, 2004;Skovhus *et al.*, 2007). The strains included in this Ph.D. study originated from the open water column, from algae and seaweeds, fish skin or copepods as well as rocks and sediment, illustrating the wide range of environments to which *Pseudoalteromonas* species have adapted and successfully colonized.

The antagonistic *Pseudoalteromonas* species are frequently associated with marine microbial surface communities on biotic or abiotic surfaces (Gram *et al.*, 2010;Holmström & Kjelleberg, 1999). From an ecology point of view this makes sense, since a bacterium adapted to planktonic life will rarely want to find itself in a situation where production of antagonistic compounds would provide a competitive advantage. Intuitively one can imagine polar antagonistic compounds diffusing freely away resulting in concentrations much too low to exert their antagonistic effects, thus incurring a fitness (production) cost without any gains. Environmental biofilms consists of mixed microbial species and competition for nutrients and space is high (Egan *et al.*, 2008). Several strategies to overcome these challenges exist, e.g. production of antagonist compounds, production of EPS and modulation of bacterial and/or host signaling (Hibbing *et al.*, 2009). The high density of microbial cells in surface biofilms may also facilitate an increased rate of exchange of genetic material, allowing for exchange of e.g. biosynthetic pathways. Additionally the biofilm complex may physically limit the rate of diffusion (Stewart, 2003), making it easier for antagonistic species to maintain the local concentration of the bioactive compound.

Many *Pseudoalteromonas* species produce enzymes which degrade complex polysaccharides such as xylan, chitin or cellulose. In planktonic *Pseudoalteromonas* strains these enzymes likely allow rapid breakdown of particulate organic matter (POM) to support bacterial growth and survival, while surface associated *Pseudoalteromonas* could rely on such enzymes to break down host cell wall polysaccharides for nutrition. Cellulose induces assembly of type IV-like pili in *P. tunicata* which in turn promotes attachment to the algae *Ulva lactuca*, a known host of *P. tunicata* (Dalisay *et al.*, 2006) and thus could play a central role in initiating biofilm formation. It also suggests that *P. tunicata* may selectively colonize and feed on surfaces with specific available polysaccharides, providing a model for niche colonization within marine bacteria. It is hypothesized that some *Pseudoalteromonas* species may also colonize and interact with higher eukaryotes to the benefit of the host organism. In this scenario, the *Pseudoalteromonas* produce antagonistic compounds to e.g. protect the host from excessive fouling (Boyd *et al.*, 1999;Dobretsov & Qian, 2002;Holmström *et al.*, 2002) and may in turn acquire nutrients either by scavenging directly from the host via enzymatic degradation or from the increased concentration of organic matter found on any marine surface (Wahl, 1989). These are examples of how *Pseudoalteromonas* species have adapted to life on marine surfaces. In the present study, the strains with antagonistic effects towards *Vibrio anguillarum* and *Staphylococcus aureus* were significantly more likely to originate from a surface (Vynne *et al.*, 2011). This is consistent with bacterial antagonism as a selective advantage in the competitive environment of marine surfaces. However it was previously reported that the overall abundance of Pseudoalteromonads was lower in surface associated communities than in bacterioplankton (Skovhus *et al.*, 2007). This is likely a reflection of the multiple strategies that exist for microbial competition in surface communities.

3.2.1 *Pseudoalteromonas* species interact with higher eukaryotes

It is known that some marine host-microbe interactions are highly specific, to the point of a symbiosis. *Pseudoalteromonas* species are known to be associated with higher eukaryotes (Holmström & Kjelleberg, 1999), and the suite of bioactive compounds produced by some species may relate to such interactions. The macroalga *M. oxyspermum* will not develop its normal morphology unless it is colonized by certain marine bacteria (Tatewaki et al., 1983). Similarly, larvae of many bivalve species are more likely to settle where bacteria that secrete a chemical cue have already established (Hadfield & Paul, 2001). Indeed, biofilms of some Pseudoalteromonas species induce settlement of eukaryote larvae to actively promote macrofouling (Huggett et al., 2006; Tebben et al., 2011), while other species deter settlement (Holmström et al., 1992). Crustose corraline algae (CCA) are important in marine ecology as several organisms including sea urchin feed on these calcifying algae, and studies of CCA epiphytic bacteria provide excellent insight into the complexity of *Pseudoalteromonas* - eukaryote interactions. Hugget et al. (2006) reported that Pseudoalteromonas strains were among the bioactive bacteria isolated from a crustose corraline alga, and that some greatly induced settlement of larvae from the sea urchin Heliocidaris erythrogramma while other Pseudoalteromonads were poor inducers of larval settlement. Upon closer inspection, a BLAST search reveals that strains related to P. luteoviolacea were found in both the highly and poorly inducing group. It is tempting to speculate that some of these bacteria produce compounds with cytotoxic activity towards sea urchin to protect the CCA host although no direct evidence of this exists. Furthermore, it would be of interest to perform a detailed

phylogenetic study on these *P. luteoviolacea* strains, as this Ph.D. study has revealed a correlation between phylogeny and production of bioactive compounds (Vynne *et al.*, 2012b) and extending this approach could reveal any phylogenetic patterns in larval interactions which may be related to production of small-molecule compounds (Tebben *et al.*, 2011). Furthermore, a bachelor student in our group recently demonstrated that some strains of *P. luteoviolacea* are highly toxic to live *Artemia nauplii* whilst others have no adverse effect. This division reflected the antibiotic profile of the strains, since two strains producing pentabromopseudilin killed *Artemia* whereas two strains producing indolmycin had no effect (Anna Neu, unpublished data).

The *Pseudoalteromonas* eukaryote interactions could represent a strategy to attract potential host organisms and repel undesired settlement of macroscopic species, providing an indication of a microbe-host relationship with some symbiotic characteristics.

3.3 PSEUDOALTEROMONADS AS PROBIOTICS IN AQUACULTURE

The increase in fish rearing for human consumption has been dramatic and comes at a cost to the environment in part due to the accompanying use of antibiotics(Cabello, 2006). Probiotics have been defined by FAO/WHO as "live microorganisms which when administered in adequate amounts, confer a health benefit on the host." (FAO and WHO, 2001) Adding live antagonistic marine bacteria to an aquaculture system has reduced mortality in certain models and could reduce the need for antibiotics (Gatesoupe, 1997;Moriarty, 1998;Planas *et al.*, 2006;Rengpipat *et al.*, 1998). This provides a basic 'proof-of-concept' that antagonistic bacteria can act as probiotics. Based on their bacterial antagonist properties, *Pseudoalteromonas* strains are candidates for application as probiotics. Longeon *et al.* describes the probiotic effect of *Pseudoalteromonas* sp. X153 which produces an antibacterial protein of 87 kDa(Longeon *et al.*, 2004). Rearing of scallop in co-culture with *Pseudoalteromonas* sp. X153 significantly reduced scallop mortality but also, due to reasons unknown, slightly reduced the scallop growth rate. Interestingly, 16S rRNA gene sequence analysis showed a 99.9% identity to *P. piscicida*, a known fish pathogen. A co-culture of three commensal strains, one of them identified via 16S rRNA gene sequence similarity as *P. elyakovii*, of larvae of Atlantic halibut was shown to improve survival of the larvae when grazing on

bacteria inoculated prey (Bjornsdottir et al., 2010). Probiotic functions are not restricted to promoting animal life, as illustrated by the ability of a marine bacterium to enhance the growth rate of a range of plant species due to production of an enzyme with catalase activity (Dimitrieva et al., 2006). The bacterium was assigned to the Pseudoalteromonads based on biochemical and physiological tests, and identified by the authors as Pseudoalteromonas porphyrae based on 16S rRNA gene sequence similarity. However, the bacterium should likely be referred to as a strain of P. atlantica or P. elyakovii (both >97% 16S rRNA gene sequence similarity) as P. porphyrae is currently not an accepted species designation (Euzéby, 1997). These studies illustrate the potential of Pseudoalteromonads to function as probiotics within diverse aquaculture applications, and provide a glimpse into the possible roles played by assorted *Pseudoalteromonas* strains in the marine environment. In this thesis work, the fish pathogenic bacterium Vibrio anguillarum 90-11-287(Skov et al., 1995) was frequently used as test organism in agar based assays to evaluate production and activity of antagonistic compounds by *Pseudoalteromonas* strains. The ability of 51 strains to consistently inhibit V. anguillarum in a live cell assay (Vynne et al., 2011) suggests that these antagonistic *Pseudoalteromonas* strains might present interesting opportunities for research and industrial applications within aquaculture probiotics. In spite of the pathogenic traits of some Pseudoalteromonads, they are not considered a problem within aquaculture where e.g. Vibrio species will typically pose a greater problem (Toranzo et al., 2005). This opens the possibility for evaluating *Pseudoalteromonas* as fish probiotics on a case-by-case basis. Ideally non-pathogenic strains would be pursued. However, pathogenic traits may be specific towards certain species of fish or mollusks which may allow for selective use of for instance an algal pathogen in turbot aquaculture.

3.4 PATHOGENICITY OF PSEUDOALTEROMONADS

Whether discussing biotechnological applications or the ecology of *Pseudoalteromonas* strains, it is prudent to consider their potential for pathogenic interactions. The majority of *Pseudoalteromonas* strains are unlikely to be human pathogens as they do not grow well at 37°C, and there are no literature reports of infectious disease in humans caused by *Pseudoalteromonas* bacteria. While not strictly a result of pathogenicity, the poisoning that can occur when eating

certain fish of the order Tetradontiformes is attributed to an accumulation of diverse marine bacteria that produce the neurotoxin tetrodotoxin (Chau et al., 2011; Noguchi et al., 2006). Among these, Pseudoalteromonas tetraodonis was isolated from the skin slime of a pufferfish and produces tetrodotoxin (Simidu et al., 1990), hence it is likely involved in human poisoning following consumption of pufferfish. Pseudoalteromonas species also colonize and grow on icestored fish (Broekaert et al., 2011) but are unlikely to be a cause of infectious disease since they do not grow at human body temperature (37C). Going beyond the human perspective and into the natural marine environment of Pseudoalteromonas bacteria, a number of Pseudoalteromonas species are associated with infectious disease in fish, molluscs or seaweed and alga. As the name suggests, P. piscicida is a fish pathogen. Rapid killing was observed in challenge tests involving live bacterial culture (Bein, 1954). No attempts were made to identify the cause of killing, but no fish death was observed when using a different unidentified marine bacterium in the same experimental setup. A different strain of *P. piscicida* infects damselfish eggs causing significantly higher mortality than untreated controls or common fish pathogen Vibrio anguillarum (Nelson & Ghiorse, 1999). P. piscicida is known to produce several bioactive substances and one might speculate that the cytotoxic compound norharman could be involved in pathogenicity (Zheng et al., 2006). Pseudoalteromonas sp. LT-13 may be an opportunistic pathogen, taking advantage of the weakening that occurs upon infection in scallop larvae (Sandaa et al., 2008). An earlier study suggested *Pseudoalteromonas* sp. LT-13 acted as a true pathogen (Torkildsen *et al.*, 2005), however this may have been due to a general weakening of the scallop larvae due to the high bacterial load encountered in the experimental setup. A follow-up study using lower bacterial concentrations indicated that LT-13 is not a primary pathogen, but may be involved in infections as an opportunistic secondary pathogen (Sandaa et al., 2008). Pseudoalteromonas species have been isolated in numerical abundance from diseased pacific oysters (Garnier et al., 2007), but no direct evidence of infection was obtained through re-inoculation of a healthy population with a pure bacterial culture. P. elyakovii was isolated from diseased spots on Laminaria japonica (Sawabe et al., 2000) and was suggested to cause the disease.

The limited number of literature reports on pathogenic *Pseudoalteromonas* species suggests that these bacteria do not represent a danger to human health, and only rarely are involved in eukaryote disease.

32

3.5 BIOACTIVE COMPOUNDS PRODUCED BY PSEUDOALTEROMONAS

The wide occurrence of *Pseudoalteromonas* shows that members of this genus are successful in colonizing and persisting in a broad range of habitats. Bioactive substances such as enzymes, antimicrobials and antifouling cues are likely part of the reason for this success, enabling the producer strains access to nutrients or advantages through inhibition of competing microbes and modulation of the behavior of macroscopic animals. The bioactive molecules can be roughly divided into two groups: Small-molecule (molecular weight of less than ca. 3 kDa) non-proteinaceous compounds that are products of secondary metabolism and proteinacious compounds including both large enzyme complexes and small peptides. The chemical diversity of the compounds is staggering when considering compounds with antibacterial or cytotoxic action, making *Pseudoalteromonas* strains interesting targets for discovery efforts focused on bioactive compounds produced by *Pseudoalteromonas* strains is presented (table 4, see page 36). Enzyme activity and modulation of eukaryote behavior are covered in separate paragraphs. *P. luteoviolacea* will have a dedicated bioactivity paragraph, as this species has been a focal point of this Ph.D. work.

3.5.1 Non-luteoviolacea Pseudoalteromonas

Antibacterial activity is observed in multiple species of Pseudoalteromonads. The inhibitory spectrum varies however, as some species inhibit Gram-negative or Gram-positive bacteria exclusively while others have very broad or very narrow ranges of inhibition. An example of this is P. phenolica O-BC30 which produces novel structurally related anti-MRSA compounds. The compound MC21-A was bacteriocidal against MRSA Staphylococci and exhibited broad-spectrum antibacterial activity (Isnansetyo & Kamei, 2003) while MC21-B inhibited Gram-positive strains but not Gram-negative bacteria or fungi (Isnansetyo & Kamei, 2009). Similarly Pseudoalteromonas sp. F-420 produces the antibiotic korormicin, which is active against Gram-negative marine bacteria (Yoshikawa *et al.*, 1997). The compound targets a unique NADH:quinone reductase (Yoshikawa *et al.*, 1999) found in many marine bacteria which enables the bacteria to maintain a sodium-motive force under the conditions present in the marine environment (Unemoto & Hayashi, 1993). This

makes korormicin a highly specialized substance which targets marine bacteria, and more knowledge of its in situ role in ecology would be of great interest. Another interesting case are the structurally distinct class of small molecule antibiotic compounds called thiomarinols, produced by Pseudoalteromonas sp. SANK 73390 (Shiozawa *et al.*, 1993;Shiozawa *et al.*, 1995;Shiozawa *et al.*, 1997). Thiomarinols are active against both Gram-negative and Gram-positive bacteria, and are synthesized from the products of two biosynthetic pathways located on a plasmid (Fukuda *et al.*, 2011). Synthetic analogues have been engineered by mutagenesis of the producer bacterium (Murphy *et al.*, 2011). Pseudoalteromonas sp. CMMED 290 produces pentabromopseudilin (PBP), bromophene and two novel anti MRSA compounds structurally related to PBP and bromophene respectively (Fehér *et al.*, 2010). This is a good example of how biosynthetic intermediates may be isolated as interesting bioactive lead compounds based on bioassays which likely do not reflect the ecological role of the active compound. Prodigiosin and related compounds feature prominently in several bacteria, notably *Serratia marcescens*. Among the Pseudoalteromonads, *P. denitrificanas* (Kawauchi *et al.*, 1997;Kim *et al.*, 1999) and *P. rubra* (Fehér *et al.*, 2008;Gerber, 1971;Vynne *et al.*, 2011) produce the red pigments prodigiosin and cycloprodigiosin HCl (figure 14).



Figure 14: Fractionated ethyl acetate extracts of *P. rubra* strains (Maria Månsson)

This class of compounds has been investigated for several pharmacological effects as reviewed by (Bennett & Bentley, 2000), for example cycloprodigiosin has potential applications as an antimalarial (Kawauchi *et al.*, 1997) and immunosuppressive agent (Kim *et al.*, 1999). Recent research has shown that prodigiosin is a potential insecticide with activity against mosquito larvae

(Patil *et al.*, 2011), while others have provided evidence that prodigiosin functions as a UVprotective pigment which may be the actual reason for its wide-spread occurrence in marine bacteria (Boric *et al.*, 2011).

Some *Pseudoalteromonas* species produce antifungal compounds which are speculated to have a role in surface colonization. *P. issachenkonii* produces the antifungal compound isatin (Kalinovskaya *et al.*, 2004), which was suggested to confer a competitive advantage in colonizing algal thalli which is the natural niche of the producer species. Isatin-producing *Alteromonas* bacteria have also been isolated from shrimp embryos and it was shown that the *Alteromonas* bacteria protected the embryos from infection by a pathogenic fungus (Gil-Turnes *et al.*, 1989). *P. tunicata* produces an antifungal tambjamine compound structurally related to prodigiosin (Burke *et al.*, 2007). This is likely the compound that was responsible for the antifungal activity observed in co-culture biofilms of *P. tunicata* and a marine fungus, where *P. tunicata* was able to exclude the fungus from the biofilm (Franks *et al.*, 2006).

P. maricaloris KMM 636 produces the chromopeptides bromoalterochromide A and A' which are cytotoxic towards sea urchin eggs (Speitling *et al.*, 2007). We identified bromoalterochromide A in 12 strains of *P. flavipulchra* which are closely related to *P. maricaloris* and *P. pisicida*, and in one *P. phenolica* strain and one *P. rubra* strain. From these strains we also identified what is likely two novel bromoalterochromides, that were also produced in 2 strains of *P. aurantia / P. citrea* (Vynne *et al.*, 2011). Synthesis of bromoalterochromides is shared among several *Pseudoalteromonas* species, and the cytotoxic effect of bromoalterochromides A and A' provides evidence for the involvement of bromoalterochromides in bacterial-eukaryote interactions. Unfortunately, bromoalterochromides degrade rapidly upon exposure to UV light which makes them hard to purify and detect.

The antibacterial compound 2-*n*-pentyl-4-quinolinol was identified from a *Pseudoalteromonas* strain related to *P. citrea* and *P. aurantia* (Long *et al.*, 2003), and from *P. piscicida* A1-J11 (del Castillo *et al.*, 2008). During my Ph.D. work, the compound was identified in 5 of 5 strains related to *P. citrea* and *P. aurantia*, and in 1 of 13 strains related to *P. flavipulchra* and *P. piscicida* (Vynne *et al.*, 2011). An additional 3 quinolinone compounds were also identified in some of these strains, with no apparent species-specific pattern. 2-*n*-pentyl-4-quinolinol strongly inhibited *Vibrio harveyi*

and *Vibrio fischeri* but not other *Vibrio* sp. or *Pseudomonas* sp. (Long *et al.*, 2003). Interestingly, it was previously described as a primary antibiotic in a marine Pseudomonad (Wratten *et al.*, 1977) and as seen in figure 15 resembles the structure of the quinolone molecules utilized for signalling by *Pseudomonas aeruginosa* (Pesci *et al.*, 1999) which opens for discussion of the role of small-molecule antibiotics in the environment.



Figure 15: Chemical structure of 2-*n*-pentyl-4-quinolinol **1** and the *Pseudomonas* quinolone signal 2-heptyl-3-hydroxy-4-quinolone **2**.

Recent publications have raised questions regarding the role of antimicrobial compounds *in situ*. Many antibiotics have profound influence on gene expression levels in microorganisms exposed to sub-lethal concentrations of the antibiotics (Davies *et al.*, 2006), for instance subinhibitory concentrations of β -lactam or aminoglycoside antibiotics induce biofilm formation in *Pseudomonas aeruginosa* (Bagge *et al.*, 2004;Hoffman *et al.*, 2005). Biofilm formation is known to increase bacterial resistance towards antibiotics (Mah & O'Toole, 2001), and thus may be considered a defense mechanism. Indeed, laboratory studies indicate that antibiotic production may be of particular benefit in a structured environment (Chao & Levin, 1981) which may also be applicable for marine surface colonizers. Little is known of the effect of an antibiotic on the producer organism itself and the microbial community as a whole in a marine natural environment, and exploration of the *in situ* activities of naturally occurring antibiotics has the potential to shed a new light on marine microbial ecology.

Species	Source	Pigment	Bioactive compounds	Bioactivity
P. aliena	Seawater	+ (melanin)	Unknown compounds	Anti-tumorogenic
P. agarivorans	Seawater, ascidians	-	Unknown	Degrades algal polysaccharides
P. antarctica	Seawater, sea-ice, muddy soils	-	-	Cold-active enzymes
P. arctica	Seawater	-	-	-
P. atlantica	Seawater, marine alga	-	-	Degrades algal polysaccharides
P. aurantia	Seawater, alga	Yellow	2-pentyl-4-quinolinol	Antibacterial
			Macromolecule	Antibacterial
			Unknown	Antidiatom
P. byunsanensis	Tidal flat sediment	-	-	-
P. carrageenovora	Seawater, marine alga	-	-	Degrades algal polysaccharides
P. citrea	Seawater, mussels,	Yellow,	2-pentyl-4-quinolinol	Antibacterial
	ascidians, sponges	melanin	Unknown	Antidiatom
P. denitrificans	Seawater	Red	Cycloprodigiosin HCl	Antimalarial, immunosuppressant, anti-tumerogenic
			Unknown substance	Induces settlement of sea urchin Heliocidaris erythrogramma
P. distincta	Sponge	(melanin)	-	-
P. donghaensis	Seawater	-	-	-
P. elyakovii	Mussels, alga	-	-	-
P. espejiana	Seawater	-	-	-
P. flavipulchra	Seawater, copepod	Orange/yellow	2-pentyl-4-quinolinol	Antibacterial
P. haloplanktis	Seawater, scallop	-	Isovaleric acid	Antibacterial
			Unknown	Antidiatom
				Cold-active enzymes
P. issachenkonii	Alga	-	Isatin	Antifungal, hemolytic
P. lipolytica	Estuarine water	-	-	-
P. luteoviolacea	Seaweed, algae, rocks, seawater	Purple	Violacein	Antibacterial, cytotoxic
			Pentabromopseudilin	Antibacterial, inhibits myosin motor activity
			Indolmycin	Antibacterial
			L-amino acid oxidase	Antibacterial
			2,4-Dibromo-6-chlorophenol	Antibacterial, antifungal
P. maricaloris	Sponges	Yellow	Bromoalterochromides A and B	Antibacterial, cytotoxic towards sea urchins
P. marina	Tidal flat sediment	-	-	-

 Table 4: Bioactive compounds produced by Pseudoalteromonas species. Modified from Bowman (2007)

P. mariniglutinosa	Diatoms	-	-	-
P. nigrifaciens	Seawater, raw fish, salted foods, mussels	(melanin)	-	-
P. paragorgicola	Sponge	-	•	-
P. peptidolytica	Seawater	Yellow	Unknown compounds	Antimicrobial, hemolytic
P. phenolica	Seawater	Brown	2,2',3-tribromobiphenyl-4-4'- dicarboxylic acid 3,3',5,5'-tetrabromo-2,2'-	Antibacterial against Gram-positive bacteria Antibacterial
			biphenyldiol	
P. piscicida	Estuarine water, fish	Yellow	Protein	Antibacterial
	samples, copepous		2- <i>n</i> -pentyl-4-quinolinol	Antibacterial
	-		Unknown	Antidiatom
P. prydzensis	Seawater	-	-	-
P. rubra	Seawater	Red	Prodigiosin, cycloprodigiosin, 2-(p- hydroxybenzyl)prodigiosin	Antimalarial, immunosuppressant, insecticide, UV- protection
			Rubrenoic acids	Bronchodilatory
D. mathemism	Chall@ab.aaaaaataa	Dala anana	Unknown	Antidiatom
P. ruthenica	Sneimsn, seawater	Pale orange	Unknown compounds	Antimicropial
P. spongiae	Sponge	Pale orange	Unknown compounds	Strong induction of Hydroides elegans settlement
P. tetraodonis	Puffer fish	-	Tetrodotoxin	Neurotoxic
P translucida	Seawater	-	-	-
1. (14/15/46/44	Scawater			
P. tunicata	Tunicate	Dark green	Tambjamine Violacein AlpP L-amino acid oxidase	Kills <i>C. elegans</i> Antibacterial, cytotoxic Antibacterial, autoinhibitory
P. tunicata	Tunicate	Dark green	Tambjamine Violacein AlpP L-amino acid oxidase Unknown	Kills <i>C. elegans</i> Antibacterial, cytotoxic Antibacterial, autoinhibitory Antidiatom
P. tunicata	Tunicate	Dark green	Tambjamine Violacein AlpP L-amino acid oxidase Unknown Unknown protein	Kills <i>C. elegans</i> Antibacterial, cytotoxic Antibacterial, autoinhibitory Antidiatom Rapidly kills <i>C.</i> <i>elegans</i>
P. tunicata P. ulvae	Tunicate Marine alga	Dark green Purple	Tambjamine Violacein AlpP L-amino acid oxidase Unknown Unknown protein Unknown	Kills C. elegans Antibacterial, cytotoxic Antibacterial, autoinhibitory Antidiatom Rapidly kills C. elegans Inhibits settlement of invertebrate larvae and algal spores
P. tunicata P. ulvae	Tunicate Marine alga	Dark green	Tambjamine Violacein AlpP L-amino acid oxidase Unknown Unknown protein Unknown	Kills C. elegans Antibacterial, cytotoxic Antibacterial, autoinhibitory Antidiatom Rapidly kills C. elegans Inhibits settlement of invertebrate larvae and algal spores Antidiatom
P. tunicata P. ulvae P. ulvae	Tunicate Marine alga Seawater, fish	Dark green	Tambjamine Violacein AlpP L-amino acid oxidase Unknown Unknown protein Unknown Unknown	Kills C. elegans Antibacterial, cytotoxic Antibacterial, autoinhibitory Antidiatom Rapidly kills C. elegans Inhibits settlement of invertebrate larvae and algal spores Antidiatom, hemolytic
P. tunicata P. tunicata P. ulvae P. undina Pseudoalteromona	Tunicate Marine alga Seawater, fish	Dark green	Tambjamine Violacein AlpP L-amino acid oxidase Unknown Unknown protein Unknown Unknown Unknown Unknown Unknown Unknown	Kills C. elegans Antibacterial, cytotoxic Antibacterial, autoinhibitory Antidiatom Rapidly kills C. elegans Inhibits settlement of invertebrate larvae and algal spores Antidiatom, hemolytic
P. ulvae P. undina Pseudoalteromona SANK 73390	Tunicate Marine alga Seawater, fish	Dark green	Tambjamine Violacein AlpP L-amino acid oxidase Unknown Unknown protein Unknown Unknown	Kills C. elegans Antibacterial, cytotoxic Antibacterial, autoinhibitory Antidiatom Rapidly kills C. elegans Inhibits settlement of invertebrate larvae and algal spores Antidiatom, hemolytic
P. tunicata P. tunicata P. ulvae P. undina Pseudoalteromona SANK 73390 F-420	Tunicate Marine alga Seawater, fish Is sp. : Alga	Dark green	Tambjamine Violacein AlpP L-amino acid oxidase Unknown Unknown protein Unknown Unknown	Kills C. elegans Antibacterial, cytotoxic Antibacterial, autoinhibitory Antidiatom Rapidly kills C. elegans Inhibits settlement of invertebrate larvae and algal spores Antidiatom, hemolytic Antibacterial against Gram-negative bacteria
P. ulvae P. undina P. undina Pseudoalteromona SANK 73390 F-420 SWAT5	Tunicate Marine alga Seawater, fish ss sp. : Alga Marine particle	Dark green	Tambjamine ViolaceinAlpP L-amino acid oxidaseUnknown Unknown proteinUnknownUnknownHindersteinUnknownChiomarinols A, B, C, D, E, F, G Korormicin2-n-pentyl-4-quinolinol	Kills C. elegans Antibacterial, cytotoxic Antibacterial, autoinhibitory Antidiatom Rapidly kills C. elegans Inhibits settlement of invertebrate larvae and algal spores Antidiatom, hemolytic Antibacterial against Gram-negative bacteria Antibacterial

Unknown (a)	Seaweed	Unknown	Halophenol	Antibacterial, antifungal
CMMED 290	Nudibranch	Unknown	2,3,5,7-tetrabromobenzofuro- [3,2- <i>b</i>]pyrrole	Anti MRSA
			4,4',6-tribromo-2,2'-biphenol	Anti MRSA
			Pentabromopseudilin	Antibacterial, inhibits myosin motor activity
			Bromophene	Antibacterial
J010	Corraline algae	Unknown	Tetrabromopyrrole	Induces settlement and metamorphosis of coral larvae

(a) Jiang et al. 2006

3.5.2 PSEUDOALTEROMONAS LUTEOVIOLACEA

One of the first antibacterial compounds that were described from marine bacteria was pentabromopseudilin (PBP), identified as a novel compound produced by a bacterium later assigned to the *Pseudoalteromonas* (Burkholder *et al.*, 1966;Fenical, 1993). This compound was identified in a subset of *P. luteoviolacea* strains in this Ph.D. work (figure 16), and although it was claimed to also be produced by a *Chromobacterium* sp. (Andersen *et al.*, 1974) the bacterial strain in question shows biochemical characteristics resembling *P. luteoviolacea* over *Chromobacterium*, in particular a negative catalase reaction, survival for 3-4 days in plate cultures and optimum growth around room temperature (Gauthier, 1982;Gauthier & Flatau, 1976).



Figure 16: Small-molecule bioactive compounds produced by *P. luteoviolacea.* Violacein **1**, pentabromopseudilin **2** and indolmycin **3**.

PBP shows excellent *in vitro* activity against Gram-positive bacteria but is not active against Gramnegative bacteria or *Candida albicans* (Burkholder *et al.*, 1966). PBP is a potent inhibitor of myosin motor activity with clinical potential in treatment of malaria, cancer and heart disease (Fedorov *et* *al.*, 2009). In addition to PBP, *P. luteoviolacea* produces the purple pigment violacein. Violacein was reported to have antibacterial activity (Lichstein & Vandesand, 1945), though this was not observed in this Ph.D. work in tests against *P. luteoviolacea* strains, *Vibrio anguillarum* and *Staphylococcus aureus*. Violacein has potent cytotoxic activity (da Silva Melo *et al.*, 2000), and is likely used as a chemical defense molecule by the producing organisms (Matz *et al.*, 2004;Matz *et al.*, 2008). Violacein production is also described in *P. tunicata* (Matz *et al.*, 2008), *P. denitrificans* (Yada *et al.*, 2008) and in several other genera (Agematu *et al.*, 2011;Becker *et al.*, 2009;Hakvåg *et al.*, 2009;Lichstein & Vandesand, 1945;Wang *et al.*, 2009). Phylogenetic analysis of amino acid sequences of biosynthetic proteins involved in violacein synthesis indicate that the violacein biosynthesis in Pseudoalteromonads diverged at an early point (Hakvåg *et al.*, 2009). This may be the result of a genetic transfer event involving ancestral bacteria which since differentiated into *Pseudoalteromonas* species and violacein producing β-Proteobacteria, respectively, or it could indicate that violacein biosynthesis is an ancient pathway which originated in a common ancestor.

A third small-molecule antibiotic substance produced by *P. luteoviolacea* is indolmycin (Månsson *et al.*, 2010;Vynne *et al.*, 2011), as described in this Ph.D. work. Hitherto known only from *Streptomyces* species (Kanamaru *et al.*, 2001;von Wittenau & Els, 1961) indolmycin is a potent antibacterial agent with selective antibacterial activity against both Gram-negative and Gram-positive bacteria and particularly effective against *S. aureus* with a reported minimum inhibitory concentration of 0.25 µg/ml (Hurdle *et al.*, 2004) and *Helicobacter pylori* (Kanamaru *et al.*, 2001). The tryptophan-derived indolmycin inhibits the bacterial tryptophanyl-tRNA synthetase (Werner *et al.*, 1976) by competitive binding, and mutations leading to changes of specific amino acids in the tryptophanyl-tRNA synthetase or changes in the promotor sequence of the tryptophanyl-tRNA synthetase det *al.*, 2009). A *P. luteoviolacea* with no strain name given was reported to produce the antibacterial and antifungal compound 2,4-Dibromo-6-chlorophenol (Jiang *et al.*, 2000).

In the course of this Ph.D. study a link between phylogeny based on housekeeping genes and production of the bioactive compounds indolmycin and PBP was discovered (Vynne *et al.,* 2012b).

Indolmycin or PBP were produced by distinct phylogenetic groups, providing evidence for the importance of these compounds in the producer organisms as the biosynthetic pathways are maintained over evolutionary time.

In addition to the small-molecule compounds mentioned above, *P. luteoviolacea* is capable of synthesizing a macromolecular antibacterial substance. An L-amino acid oxidase (LAO) was characterized from two *P. luteoviolacea* strains by (Gomez *et al.*, 2008) and is likely synonymous with the macromolecule previously described in the type strain (Gauthier, 1976). Other LAOs are known from marine bacteria (Lucas-Elio *et al.*, 2005) and includes the AlpP autolysis protein produced by *P. tunicata* (Mai-Prochnow *et al.*, 2008) and an anti-MRSA agent produced by *P. flavipulchra* (Chen *et al.*, 2010). It is tempting to speculate that autolysis in *P. luteoviolacea* is due to LAO activity (Gauthier & Flatau, 1976), as suggested for *P. tunicata* (Mai-Prochnow *et al.*, 2004). The physiological role of LAOs in marine bacteria is unknown, though the antimicrobial effect of LAOs is due to the release of hydrogen peroxide when the LAO substrate is oxidized (Gomez *et al.*, 2008;Mai-Prochnow *et al.*, 2008). In addition to possible *in situ* antimicrobial effects they may increase available nitrogen by oxidizing amino acids in the local milieu as suggested for fungal LAOs (Davis *et al.*, 2005).

3.6 ENZYMATIC ACTIVITIES OF PSEUDOALTEROMONADS

Microbial life is dependent on enzyme activities. Extracellular enzymes play crucial roles in providing accessible nutrient sources and are often highly specialized for the niches of the producer organism. This ecology-driven specialization has gone on for the entirety of microbial evolution, and thus offers a wealth of highly specialized enzymes which may have uses within industries as diverse as dairy production and biofuels. The ubiquitous nature of the Pseudoalteromonads is an indication that they thrive in a wide range of environments, and arctic strains in particular have been investigated for enzymatic activities in the expectation that novel enzymes active at low temperature would be discovered. An overview of some of the currently described enzymatic activities is presented in table 5.

Many of the characterized enzymes from *Pseudoalteromonas* strains are from psychrophilic strains and exhibit cold-adapted properties. This opens for different applications such as treatment of dairy products to remove lactose at low temperature (Hoyoux *et al.*, 2001), however the cold adapted enzymes are often also less stabile and will denature or undergo conformational changes at a lower temperature than mesophilic enzymes (Georlette *et al.*, 2004).

Strain	Product	Reference
P. haloplanktis	Cold-adapted β-galactosidase	Hoyoux et al. 2001
Pseudoalteromonas sp.	Cold-adapted β-galactosidase	Fernandes <i>et al.</i> 2002
P. haloplanktis TAC125	Lipase PhTAC125 Lip1	de Pascale <i>et al.</i> 2008
Pseudoalteromonas sp. DY3	Cold-active cellulase	Zeng <i>et al.</i> 2006
Pseudoalteromonas sp. NJ276	Cold-active protease	Wang <i>et al.</i> 2008
Pseudoalteromonas sp. SM9913	Cold-adapted subtilase	Yan <i>et al.</i> 2009
P. haloplanktis TAB23	Cold-adapted α-amylase	Tutino <i>et al.</i> 2002
P. haloplanktis ANT/505	Pectate lyase	Van Truong <i>et al.</i> 2001
P. haloplanktis	Xylanase	Van Petegem <i>et al.</i> 2002
Pseudoalteromonas sp. MB-1	Endoglucanase	You <i>et al.</i> 2005
Pseudoalteromonas sp. 005NJ	α-l-rhamnosidase	Gaston Orrillo <i>et al.</i> 2007
Pseudoalteromonas arctica	Cold-adapted esterase	Al Khudary <i>et al.</i> 2010
P. haloplanktis	Cold-adapted cellulase	Violot <i>et al.</i> 2005
P. haloplanktis	DNA ligase	Georlette <i>et al.</i> 2000
P. carrageenovora	Carrageenase	Michel <i>et al.</i> 1999
Pseudoalteromonas sp. S9	Chitinase	Techkarnjanaruk <i>et al.</i> 1997
P. arctica GS230	Cold-adapted α-amylase	Lu <i>et al.</i> 2010
P. antarctica N-1	Agarase	Vera <i>et al.</i> 1998
Pseudoalteromonas sp. CKT1	Agarase	Xavier Chiura <i>et al.</i> 2000

Table 5: Enzyme activity from *Pseudoalteromonas* strains

Representative strains from the collection used in the present work were screened for production of industrially relevant enzymes in an agar based assay (figure 17 below) containing azurine cross-linked polysaccharides (Megazyme, Ireland). Six of 12 strains tested amylase positive and 5 of 12 tested curdlanase positive, with four strains testing positive in both assays (Vynne, unpublished).

Figure 17: Amylase assay on *Pseudoalteromonas* strains. The AZCL-substrate is seen as blue particles in the agar and dark blue halos indicate enzyme activity.



The positive strains were both pigmented and non-pigmented. A reason for the low number of observed enzymatic activities within the *Pseudoalteromonas* strains from this Ph.D. study could be that pigmented strains were overrepresented in the selection for screening, since the pigmented bacteria were the primary research interest due to their antagonist activities while the majority of the producer organisms listed in table 5 are of the non-pigmented clade. Also, since the screening took place at 25°C, some strains may have cold-adapted enzyme activities which were not detected due to the incubation temperature.

3.7 ANTI-FOULING POTENTIAL OF *PSEUDOALTEROMONAS* **SPECIES**

While some *Pseudoalteromonas* promote settlement of eukaryotes as previously described (Huggett *et al.*, 2006), many species have been investigated for their anti-fouling and anti-biofilm activities (Bernbom *et al.*, 2011;Boyd *et al.*, 1999;Dobretsov & Qian, 2002;Egan *et al.*, 2001a;Holmström *et al.*, 2002). Fouling is a build-up of biological material on a surface, which in the marine environment develops rapidly unless preventive measures are taken. Briefly, submerging a clean surface into the marine environment will cover the surface in a thin film of organic substances within minutes. Bacteria and diatoms will follow within hours, forming the basis for increased colonization by larvae of macrofouling organisms(Abarzua & Jakubowski, 1995;Colwell, 1983;Wahl, 1989). Fouling is a problem in the marine industries, particularly within shipping as a dense layer of fouling rapidly develops on a submerged hull reducing speed and fuel efficiency unless preventive measures are taken (Champ, 2000).

Pseudoalteromonas tunicata D2 produces anti-fouling compounds (Holmström *et al.*, 1998) shown to correlate with pigment production (Egan *et al.*, 2002). Other *Pseudoalteromonas* strains inhibit larvae and diatom settlement in direct cell interaction assays (Lee & Qian, 2003) and in tests of aqueous bacterial cell extracts (Dobretsov & Qian, 2002). Some antifouling effects may be due to bacterial antagonism towards other microorganisms which promote larval settlement, but there is no consistent link between antifouling and antibacterial compounds (Bernbom *et al.*, 2011).

3.8 CONCLUSION FROM CHAPTER 3

The genus *Pseudoalteromonas* has a complex and intricate taxonomy and underlying phylogeny. Care should be taken if 16S rRNA gene sequences are used as the only means of species identification when a novel Pseudoalteromonas isolate is being described, as this Ph.D. thesis demonstrate that even at 99% 16S rRNA sequence identity, it remains possible to describe new species. Correct taxonomical assignments are important for scientific communication and understanding evolutionary dynamics with a bacterial genus. Many *Pseudoalteromonas* species produce bioactive compounds and the present study expanded on the knowledge of bioactive small-molecule compounds and their distribution within the Pseudoalteromonas species. The potent antibacterial compound indolmycin was discovered for the first time in marine bacteria, and several other known bioactives were identified. It was discovered that, within a single species, three antibiotics were produced by distinct phylogenetic groups. The selective maintenance of the biosynthetic pathways suggests the antibiotics play an important role for the producer organism, likely in community interactions and protection given the antagonist nature of the compounds. The role of antibiotics in the natural environment is increasingly up for debate, as sublethal concentrations of antibiotics may in fact have dramatic effects on bacterial populations and hence what we perceive as antibiotics may in some cases serve a different purpose under natural conditions. As novel *Pseudoalteromonas* species and strains with interesting bioactivities are continuously described, it stands to reason that there is a rich untapped potential for biodiscovery of novel bioactive substances within the Pseudoalteromonads.

4.0 QUORUM SENSING IN REGULATION OF BACTERIAL PRODUCTION OF

ANTIMICROBIAL COMPOUNDS

Traditionally, bacteria were thought of as individual cells with no means to actively coordinate a change in the population of gene regulation and phenotype. As our understanding of bacterial behavior in both pure cultures and mixed communities has increased, it has become clear that intra- and inter-species cell-cell communication play a key role in coordinating gene expression and subsequently phenotype in both pure cultures and mixed communities. There are many types of bacterial communication (Blango & Mulvey, 2009; Dubey & Ben-Yehuda, 2011), however, one of the currently most studied is often referred to as quorum sensing (QS). The name quorum sensing reflects its reliance on cell numbers, and it is now known to regulate advanced phenotypes such as bioluminescence (Nealson & Hastings, 1979), virulence (Latifi et al., 1995), motility (Huber et al., 2001;Kohler et al., 2000), biofilm formation (Labbate et al., 2004) and antibiotic production (Berger et al., 2011; Duerkop et al., 2009; Hothersall et al., 2011). QS is mediated by small-molecule chemical compounds called auto-inducers, which diffuse freely into the extracellular environment and act as signal carriers. As part of the work with antibiotic producing Pseudoalteromonas, we discovered compounds that possibly could be autoinducers and hypothesized that these were involved in regulation/production of a particular antibiotic. This chapter outlines the general principles of QS and focuses on examples in which bacterial antibiotic production is QS regulated.

4.1 THE VIBRIO FISCHERI LUXI/LUXR QUORUM SENSING MODEL

The process of QS has been described in many bacterial species and regulates diverse physiological and chemical processes (see e.g. (Dobretsov *et al.*, 2009;Van Houdt *et al.*, 2007). Quorum sensing was first described in the marine bacteria *Vibrio fischeri* where it regulates bioluminescence (Nealson & Hastings, 1979). This cell density dependent QS system is studied in depth and forms the canonic Lux-based QS system (figure 18). Similar LuxIR-like QS systems have been widely described in Gram-negative bacteria (Whitehead *et al.*, 2001) and are characterized by containing homologues to the LuxI and LuxR proteins. Four main components are involved in LuxIR QS: the LuxI AHL synthetase, the acylated homoserine-lactone (AHL) signaling molecules, the LuxR transcriptional regulator and the genes encoding the QS regulated phenotype (Miller & Bassler, 2001). The LuxI protein is responsible for synthesis of the AHL molecules and even at low cell densities individual cells produce a small amount of the cognate AHL (Engebrecht & Silverman, 1984). As cell density increases, the local concentration of AHL rises and eventually reach a threshold concentration which activates the LuxIR regulatory system. The system is activated by LuxR proteins binding the cognate AHL, which induces a conformational change that allows the DNA binding domain of the LuxR protein to recognize and bind to a specific DNA sequence called a *lux*-box (Choi & Greenberg, 1992;Welch *et al.*, 2000;Zhu & Winans, 2001). Upon LuxR binding, transcription of the genes downstream of the *lux*-box is activated. The *luxI* gene may also be situated downstream of the LuxR binding site, which upon LuxR binding leads to an increased production of LuxI and subsequently increased AHL concentration creating a positive feedback loop. Several variations on this central concept exist, for instance LuxR can focus as a repressor protein (Fuqua *et al.*, 1996).



Figure 18: The canonical LuxI/LuxR QS system in *V. fischeri*. IM, inner membrane; OM, outer membrane. Adapted from Waters & Bassler (2005).

4.1.1 ACYLATED HOMOSERINE-LACTONES: STRUCTURE AND ANALYSIS

The AHL molecules are the active signal carrying molecules in LuxIR quorum sensing. Most diffuse freely in and out of the individual bacterial cells, enabling the AHL-producing bacteria – and possible "eavesdroppers" – to probe the concentration of signal molecules and monitor the microbial environment to respond in a coordinated, social way. Figure 19 shows the typical AHL compositions consisting of a homoserine lactone ring with a $C_4 - C_{18}$ N-acyl side chain (Fuqua *et al.*, 2001).

Figure 19: Acylated homoserine lactones produced by different bacterial species and recognized by corresponding LuxR homologues. The R side chain may also contain a number of double bonds. Adapted from Ng & Bassler (2009).



The lactone ring will open at alkaline pH, eliminating the signaling properties of the AHL. Susceptibility to this lactonolysis decrease as the N-acyl chain increases in length (Yates *et al.*, 2002). AHLs commonly have 3-oxo or 3-hydroxy substitutions, which influences signal specificity, and may have a number of unsaturated double bonds in the side chain. The biosynthesis of AHLs rely on cellular pools of acylated acyl carrier protein and S-adenosyl-L-methionine which are the substrates used by the LuxI AHL synthetase (Parsek *et al.*, 1999;Watson *et al.*, 2002). A very interesting recent study identified a new class of homoserine lactone QS signal molecules which are based on *p*-coumaric acid rather than cytoplasmic fatty acids and function in a LuxIR analogue QS system (Schaefer *et al.*, 2008).

AHLs are commonly identified via bioassays that rely on well characterized QS systems that detect AHL signals with varying sensitivity and specificity (Cha *et al.*, 1998;McClean *et al.*, 1997;Ravn *et al.*, 2001;Winson *et al.*, 1998). Often, these systems are engineered to function as biosensors, e.g. through a *lacZ* or luminescence reporter fusion (figure 20). Since the assays are based on naturally occurring pathways, a given biosensor will in most cases produce a stronger response when its cognate AHL is added, and the range of AHLs recognized varies among biosensors (Fekete *et al.*, 2007). The biosensors are also used in conjunction with methods from analytical chemistry in order to purify and identify the AHL compounds following bioassay guided fractionation strategies as described in chapter 1 but using AHL biosensors as indicator strains (Fekete *et al.*, 2007;Wang *et al.*, 2011).



Figure 20: Bioassay for detection of AHLs. The monitor strain is Agrobacterium *tumefaciens* pZLR4. pZLR4 contains the Lux homologue Tra-system with a *lacZ* fusion. The blue zones indicate that the Tra-system was activated and lacZ transcribed, leading to production of a blue pigment due to specific cleavage of 5bromo-4-chloro-indolyl-β-Dgalactopyranoside (Cha et al., 1998). The samples tested were C-18 fractions of an ethyl acetate extract of P. luteoviolacea S4054.

4.2 THE ROLE OF QUORUM SENSING IN REGULATING BACTERIAL PRODUCTION OF ANTIBIOTICS

One of the many phenotypes regulated by QS is, in some bacteria, the production of antibiotic compounds. The ability to regulate and coordinate production of an antibiotic compound is an obvious advantage to the producer organisms, e.g. in surface colonization where QS could restrict production of antibiotics to situations where a microbial community is established, cell density is high and action against competitors is needed.

The genus *Burkholderia* contains several species that utilize signaling via QS mechanisms. Among these, production of antibiotic compounds by *Burkholderia cepacia* and *Burkholderia thailandensis* is under QS regulation. The antibiotic bactobolin which is effective against both Gram-positive and Gram-negative bacteria was recently identified in *B. thailandensis* E264 (*Seyedsayamdost et al., 2010*). A biosynthetic pathway was proposed (figure 21) (Seyedsayamdost *et al., 2010*) and shown to be under LuxIR-homologue QS control with AHL production being essential for expression of antibiacterial activity (Duerkop *et al., 2009*). Bactobolin inhibited *Bacillus subtilis* which, like *B. thailandensis*, is a soil bacterium and the authors speculate that the QS regulated production may allow *B. thailandensis* to chemically defend itself during growth as microcolonies in the soil environment. Other LuxIR-homologue QS systems with high homology to QS systems in the closely related *Burkholderia mallei* and *Burkholderia pseudomallei* exist in *B. thailandensis* and are speculated to regulate virulence factors (Duerkop *et al., 2007*;Duerkop *et al., 2008*).



Figure 21: The QS regulated bactobolin pathway. I2 and R2 are LuxI and LuxR homologues, respectively. The red arrows are annotated as NRPS genes, green arrows indicate PKS genes, pink arrows are potential accessory antibiotic synthesis genes, orange arrows are putative transport genes, the brown arrow is a metallopeptidase, and gray indicates genes of unknown function. The numbers indicate genes targeted for molecular analyses. Modified from Duerkop *et al.* (2009)

The polyketide antibiotic mupirocin used in topical treatment of Staphylococci is produced by *Pseudomonas fluorescens* NCIMB 10586 and is under QS control (Thomas *et al.*, 2010). The genes *mupl* and *mupR* are regulatory genes with their products showing high amino acid homology to LuxI and LuxR, respectively (El-Sayed *et al.*, 2001). These genes are essential for biosynthesis of mupirocin and the MupI product was recently identified as 3-oxo-C10-HSL (Hothersall *et al.*, 2011). Interestingly, production of mupirocin was not increased by addition of exogenous 3-oxo-C10-HSL or *in trans* expression of *mupI*, suggesting that an additional control mechanism possibly related to the amidase homologue MupX influences production of mupirocin (Hothersall *et al.*, 2011). The involvement of the putative AHL-degrading enzyme MupX adds an extra level of complexity to this QS network and provides a glimpse of how tight regulatory control is employed by bacteria to ensure correctly timed production of, in this case, secondary metabolites with bioactivity.

The Gram-negative γ -Proteobacteria *Erwinia carotovora* and *Serratia* sp. ATTC 39006 produce the broad-spectrum β -lactam antibiotic carbapenem (Parker *et al.*, 1982). The production of carbapenem is under QS control in both bacteria (Bainton *et al.*, 1992;Thomson *et al.*, 2000) although interestingly the QS mechanisms are different with respect to the AHL binding molecules (figure 22 below). In *Erwinia*, the LuxR homologue CarR binds the AHL directly and subsequently activates transcription of the *car* biosynthetic pathway, in similar fashion as the canonical LuxIR system. In *Serratia* sp. ATTC 39006, two LuxR homologues exist: the AHL binding transcriptional repressor SmaR and the AHL independent transcriptional activator CarR (Cox *et al.*, 1998;Thomson *et al.*, 2000). C4-HSL or C6-HSL binds to the repressor SmaR which is then released and transcription is activated by CarR. The different approaches to regulation of the *car* biosynthetic pathway illustrate how LuxR homologues can act in different ways, possibly as a result of adaptation to different environments and variation in the role within regulatory cascades. In *Serratia* spp., QS is also involved in regulating production of the bioactive pigment prodigiosin (Thomson *et al.*, 2000) and several other interesting phenotypes such as swarming and biofilm formation (reviewed by Van Houdt *et al.* (2007)).

Members of the marine bacterial *Roseobacter* clade are widely isolated from the marine environment, often in association with marine alga (Buchan *et al.*, 2005). Some species of the Roseobacter clade are antibacterial (Gram *et al.*, 2010;Hjelm *et al.*, 2004;Martens *et al.*, 2007) and

50

this is frequently associated to production of the antibacterial compound tropodithietic acid (TDA) (Brinkhoff *et al.*, 2004;Bruhn *et al.*, 2007;Geng *et al.*, 2008;Porsby *et al.*, 2008). AHL production is widespread among *Roseobacter* strains (Bruhn *et al.*, 2007;Gram *et al.*, 2002;Wagner-Döbler *et al.*, 2005) and production of QS signals has been hypothesized to play an important role in *Roseobacter* ecology (Bruhn *et al.*, 2005;Slightom & Buchan, 2009).



Figure 22: QS systems regulating the carbapenem biosynthesis pathway in *Erwinia* and *Serratia* sp. ATCC 39006. In *Erwinia* carbapenem synthesis is induced at high cell density due to the AHL signal binding the LuxR homologue CarR and subsequent induction of the car biosynthetic pathway. In contrast, carbapenem biosynthesis in *Serratia* is activated by AHL binding to and inactivating the SmaR repressor, allowing the AHL independent *Serratia* CarR to induce the *car* pathway. Modified from Barnard *et al.* (2007).

The synthesis of TDA is cell-density dependent in some strains (Bruhn *et al.*, 2007), and recent work concluded that production of TDA is under QS control by a 3-OH-C10-HSL induced LuxIR homologue system in *Phaeobacter gallaciensis* DSM 17395 (Berger *et al.*, 2011). The *tdaA* gene regulates expression of the TDA biosynthetic pathway and is induced by the LuxR-homologue PgaR. As expected PgaR induces *tdaA* expression when stimulated with the Pgal AHL-product, but curiously TDA can also activate PgaR and induce *tdaA*. Although there is no distinct structural similarity between TDA and the cognate AHL (figure 23), TDA effectively functions as an

autoinducer acting on parts of a LuxIR homologue system, although much higher concentrations of TDA are needed to induce similar expression levels (Berger *et al.*, 2011).



4.3 QUORUM SENSING IN PSEUDOALTEROMONAS

Only few studies are published on QS in *Pseudoalteromonas*. Two strains have been investigated for QS aspects in their physiology (Guo *et al.*, 2011;Wang *et al.*, 2008), and *Pseudoalteromonas* spp. have been identified as AHL producers in subtidal biofilms (Huang *et al.*, 2008a;Huang *et al.*, 2008b). *Pseudoalteromonas* are also likely included in more general studies of marine microbiota, for instance a survey of coral-associated bacteria revealed *Pseudoalteromonas* strains that activated the biosensors *Agrobacterium tumefaciens* KYC55 or *Escherichia coli* K802NR-pSB1075 (Golberg *et al.*, 2011). *Pseudoalteromonas* sp. 520P1 produces two AHLs, with one inducing production of violacein in the wild type strain (Wang *et al.*, 2008). Recombinant expression of the violacein gene cluster under control of the WT promotor in *E. coli* did, however, not result in violacein production, whereas placing the gene cluster under control of a T7 promoter did (Zhang & Enomoto, 2011). This indicates that a regulatory system native to the WT strain may be essential for expression of the violacein gene cluster from *Pseudoalteromonas* 520P1. However despite attempts to identify *lux* related elements, neither *luxl, luxR* or a *lux*-box were identified, thus no conclusive evidence was presented.

AHL synthesis and quorum sensing regulated antibiotic production was reported for *Pseudoalteromonas* sp. NJ6-3-1, related to *P. denitrificans* (BLAST search of reported 16S rRNA nucleotide sequence) (Guo *et al.*, 2011). However, the authors claim that diketopiperazines are the inducer compounds. It is known that diketopiperazines can induce Lux homologue systems,

but it would be nice to see compelling evidence that the native AHLs do not induce a QS response, for instance AHL negative knockouts.

During this Ph.D. work it was discovered that the bioactive strain *P. luteoviolacea* S4054 produces one or more molecules that induce the biosensor strains *Agrobacterium tumefaciens* pZLR4 (Cha *et al.*, 1998) and *Chromobacterium violaceum* CV026 (McClean *et al.*, 1997) (Vynne *et al.*, 2012c). The *luxI* gene homologue *pluI* was identified through shotgun cloning and expression in *E. coli* JM109-pSB401 (Winson *et al.*, 1998) followed by sequencing by 'primer walking'. Later, whole genome sequencing revealed a putatively QS regulated operon structure downstream of the *pluI* gene, followed by a *luxR* homologue *pluR* (Vynne & Gram, 2012). The genomic organization is shown in figure 24.



Figure 24: Genomic organization of QS related elements in *P. luteoviolacea* S4054 and a putatively QS regulated operon. Predicted functions: Dark blue: *plul*, red: aminotransferase, yellow: dehydrogenase, green: methyltransferase, grey: hypothetical protein, purple: coenzyme F390 synthethase, pink: methylase, brown: lyase, olive: tryptophanyl t-RNA synthethase, light blue: *pluR*. Upstream of *pluI* a palindromic sequence reminiscent of *lux*-boxes was identified (not shown).

The function of the genes downstream of *luxl* is currently unknown, but several of the genes are annotated as enzymes that are known to be active in formation of the antibiotic indolmycin (Hornemann *et al.*, 1971), which correspondingly was identified in strain S4054. In spite of focused efforts it has not been possible to identify the chemical structure of the compound causing activation of the monitor strains. Fractionation using standard approaches in analysis of AHLs (Fekete *et al.*, 2007;Wang *et al.*, 2011) did not result in clear separation of the active compounds into distinct fractions. Raw extracts of the molecules lost the ability to induce a QS response in the biosensors after alkaline treatment but was stable upon acid treatment, as expected for AHLs (Vynne *et al.*, 2012c).

In my studies of a collection of *P. luteoviolacea* strains for phylogenetic and biodiscovery purposes, I discovered that only the indolmycin producing strains synthesized these molecules now denoted

AHL-like (Vynne *et al.*, 2012c). However, all strains produced the purple bioactive pigment violacein which is under QS control in *C. violaceum* (McClean *et al.*, 1997) and possibly in *Pseudoalteromonas* sp. 520P1 (Wang *et al.*, 2008). In *Serratia*, horizontal transfer of QS related elements have revealed interesting aspects of the impact of QS on microbial physiology. Transfer of a complete prodigiosin biosynthetic cluster to a nonpigmented *Serratia* strain placed prodigiosin biosynthesis under control of the native QS system, and correspondingly, transfer of the *luxIR* homologues *smalR* to a QS negative prodigiosin producing strain placed the native prodigiosin biosynthesis under QS control (Coulthurst *et al.*, 2006). These fascinating experiments show how some biosynthetic elements can be inherently disposed to QS regulation, and may offer an explanation to the variability in QS regulation of violacein synthesis.

4.4 CONCLUSION FROM CHAPTER 4

Quorum sensing plays an important role in microbial ecology as it enables bacteria to act as coordinated units rather than individual cells. This confers benefits such as biofilm formation, production of defensive antibiotics and expression of virulence factors (Miller & Bassler, 2001;Milton, 2006). Some *Pseudoalteromonas* strains, like many proteobacteria, produce AHLs and possess LuxIR homologue QS machineries which may be involved in regulating production of bioactive compounds. Clearly, the production of antibiotics is quorum sensing regulated in some bacteria and thus QS has a role in microbial interactions both as a direct signal and indirectly through regulating biosynthesis of antagonistic molecules that can affect the microbial community. It is well known that some bacteria have orphan LuxR genes in their genome (Patankar & González, 2009). These may be remnants from previous evolutionary events, but perhaps more interestingly, they may also allow non-AHL producing bacterial cells to "eavesdrop" on conversations among other AHL-producing bacteria. This could allow preemptive production of e.g. antibiotic resistance mechanisms to counter QS-induced antibiotic production or activation of otherwise silenced biosynthetic pathways. Consequently, understanding the role of QS in microbial ecology has the potential to benefit biodiscovery efforts focused on novel antibiotics.

5.0.1 BIOACTIVE SECONDARY METABOLITES FROM MARINE BACTERIA

This part of the thesis addressed marine bacteria as a source of bioactive secondary metabolites and strategies for biodiscovery efforts. In this study it was confirmed that Pseudoalteromonas strains exhibiting stable antibacterial activity are significantly more likely to be isolated from marine surfaces. The potential of marine bacteria to deliver novel bioactive compounds was confirmed. Although no strictly novel antibiotics were discovered, several antibiotics were identified and non-antibiotic novel compounds were discovered. The marine environment has great diversity and, as shown for P. luteoviolacea, application of high resolution phylogenetic reconstruction may provide a guiding tool to determine which bacterial strains should be included in further screening efforts. Genome sequencing of the bioactive strain P. luteoviolacea S4054 revealed a large unharnessed genetic potential with as many as 11 predicted biosynthetic pathways without a known product. As genome sequencing becomes ever more accessible, whole genome sequencing may become the preferred technique for direct comparison and dereplication of the biosynthetic potential of a collection of marine bacteria. Knowledge of the ecology of the wild type organism is very useful in such a case, as culturing the organism under specific environmentally relevant nutrient levels or physical conditions may result in the production of otherwise absent secondary metabolites, which may not be easily expressed in heterologous hosts.

5.0.2 THE GENUS *PSEUDOALTEROMONAS* – ECOLOGY AND BIOACTIVITY

The genus *Pseudoalteromonas* has broad biotechnological applications. This thesis addressed the phylogeny of the genus and its production of antibacterial compounds. The use of secondary metabolites in species identifications was not wholly successful as phylogenetic analysis revealed varying levels of species specificity in secondary metabolite production, with *P. ruthenica* as an example of a narrow species specific secondary metabolite profile and *P. rubra* as an example of a species with a very diverse secondary metabolite profile. The potent antibiotic indolmycin was

discovered in marine bacteria for the first time, which may be evidence of genetic exchange among the terrestrial and marine ecosystems. During the course of the study, a new species was described and given the name *Pseudoalteromonas galatheae* after the expedition on which it was first isolated. Within *Pseudoalteromonas luteoviolacea*, three antibiotics were produced in a phylotype specific manner. This provided evidence that these antibiotics are maintained under selective pressure and thus likely are important in microbial ecology.

5.0.3 QUORUM SENSING IN REGULATION OF BACTERIAL PRODUCTION OF

ANTIMICROBIAL COMPOUNDS

Quorum sensing regulates a range of potential phenotypes. In the context of this thesis, the interest has primarily been regulation of antibiotic production. A potentially novel AHL-like molecule was described from *P. luteoviolacea* S4054, and this may take part in regulation of a novel biosynthetic pathway. However, such molecules may also be involved in countless other interactions when released into a microbial community. Understanding novel quorum sensing systems and their influence on microbial interactions could help us not only discover novel natural products, but also to better understand the regulatory events in complex microbial communities both in nature and in medicine.

5.1 OUTLOOK

Discovery of novel antibiotics and other strategies to combat the spread of resistant human pathogenic bacteria will remain a priority. Efficient screening strategies combined with sampling of novel microbial diversity will likely remain one of the most promising avenues to achieve this, although revisiting known antibiotic producing strains with modern techniques may be necessary.

In the near future, whole genome sequencing and metagenome approaches appear poised to dominate within natural product discovery. This will offer nearly unlimited access to genetic and chemical diversity, and allow dereplication of biosynthetic gene clusters as one of the first steps in a biodiscovery program. As a result, the discovery of molecules with coveted pharmaceutical properties will accelerate. In addition, selective targeting of specific microbial communities for sequencing, e.g. sponge symbiotic bacteria or bacteria from dense biofilms, may lead to a higher rate of discovery than other wise obtained. The use of heterologous expression systems will provide access to some of these compounds, yet some gene clusters may not be properly expressed, or the end product may be toxic to the producer cell. Alternatively, combining genomic approaches with knowledge of the ecology and physiology of a bioactive bacterium may lead to production of novel compounds by the wild type strain. Genome sequencing would reveal the presence of any known signaling systems, which could then be activated to potentially induce novel biosynthetic pathways, or growth conditions may be adjusted based on *in silico* reconstructions of the metabolic network of the organism in order to induce expression of biosynthetic clusters, for instance by culturing under simulated natural conditions.

Knowledge of *in situ* microbial interactions and the role antibiotics play in the microbial community will naturally be of interest to microbial ecologists, but it may also help us understand why important phenomena such as resistance to antibiotics evolve, adapt and persist in the environment, and potentially make us better equipped to counter the emergence of antibiotic resistance in pathogenic bacteria.

6.0 ACKNOWLEDGEMENTS

I would like to thank my supervisor, Lone Gram, for her endless support and enthusiasm, and her ability to keep me focused whenever I was astray. Lone always had an open door when advice was needed, experimental results needed discussion, or frustrations abounded.

I also would like to thank Debra Milton, whose lab I visited at the Department of Molecular Biology, Umeå University, Sweden for 12 weeks in the spring of 2011. The stay was very inspirational, and Debra's knowledge of the molecular microbiology of marine bacteria was instrumental in beginning to unravel the quorum sensing system in *P. luteoviolacea* S4054. Thank you Debra for your time, helpfulness, and practical advice.

I would like to thank my past and present colleagues at the National Food Institute for the nice times, the good working atmosphere and stimulating discussions, scientific and non-scientific alike.

A big thank you is due for Maria Månsson and Kristian Fog Nielsen from DTU Systems Biology. Their chemical expertise and tireless efforts has been instrumental in the creation of the work behind this thesis, and I look forward to hopefully continuing our collaboration.

Finally, I would like to thank my family and my loving girlfriend Signe. Without your support and encouragement, this would not have been possible. Tak.

- Abarzua, S. & Jakubowski, S. (1995). Biotechnological investigation for the prevention of biofouling. 1. Biological and biochemical principles for the prevention of biofouling. *Mar Ecol-Prog Ser* 123, 301-312.
- Agematu, H., Suzuki, K. & Tsuya, H. (2011).*Massilia* sp. BS-1, a novel violacein-producing bacterium isolated from soil. *Biosci Biotech Bioch* 75, 2008-2010.
- Andersen, R. J., Wolfe, M. S. & Faulkner, D. J. (1974). Autotoxic antibiotic production by a marine *Chromobacterium. Mar Biol* 27, 281-285.
- Antônio, R. V. & Creczynski-Pasa, T. B. (2004). Genetic analysis of violacein biosynthesis by Chromobacterium violaceum. *Genet Mol Res* **3**, 85-91.
- August, P. R., Grossman, T. H., Minor, C., Draper, M. P., MacNeil, I. A., Pemberton, J. M., Call, K.
 M., Holt, D. & Osburne, M. S. (2000). Sequence analysis and functional characterization of the violacein biosynthetic pathway from *Chromobacterium violaceum*. J Mol Microb Biotech 2, 513-519.
- Bagge, N., Schuster, M., Hentzer, M., Ciofu, O., Givskov, M., Greenberg, E. P. & Høiby, N.
 (2004).*Pseudomonas aeruginosa* biofilms exposed to imipenem exhibit changes in global gene expression and β-lactamase and alginate production. *Antimicrob Agents Chemother* 48, 1175-1187.
- Bainton, N. J., Stead, P., Chhabra, S. R., Bycroft, B. W., Salmond, G. P., Stewart, G. S. & Williams,
 P. (1992).N-(3-oxohexanoyl)-L-homoserine lactone regulates carbapenem antibiotic production in *Erwinia carotovora*. *Biochem J* 288, 997.
- Baumann, L., Baumann, P., Mandel, M. & Allen, R. D. (1972). Taxonomy of aerobic marine eubacteria. J Bacteriol 110, 402.
- Becker, M. H., Brucker, R. M., Schwantes, C. R., Harris, R. N. & Minbiole, K. P. C. (2009). The bacterially produced metabolite violacein is associated with survival of amphibians infected with a lethal fungus. *Appl Environ Microb* **75**, 6635-6638.
- Bein, S. J. (1954). A study of certain chromogenic bacteria isolated from red tide water with a description of a new species. *B Mar Sci* **4**, 110-119.
- Bennett, J. W. & Bentley, R. (2000). Seeing red: The story of prodigiosin. Adv Appl Microbiol 47, 1-32.

- Berger, M., Neumann, A., Schulz, S., Simon, M. & Brinkhoff, T. (2011). Tropodithietic acid production in *Phaeobacter gallaeciensis* is regulated by N-Acyl homoserine lactone-mediated quorum sensing. *J Bacteriol* **193**, 6576-6585.
- Berland, B. R., Bianchi, M. G. & Maestrin, S. Y. (1969). Study of bacteria associated with marine algae in culture .I. Preliminary determination of species. *Mar Biol* 2, 350-&.
- Bernbom, N., Ng, Y. Y., Kjelleberg, S., Harder, T. & Gram, L. (2011). Marine bacteria from Danish coastal waters show antifouling activity against the marine fouling bacterium *Pseudoalteromonas* S91 and zoospores of the green alga *Ulva australis* independent of bacteriocidal activity. *Appl Environ Microb*, AEM.
- Bjornsdottir, R., Karadottir, E. G., Johannsdottir, J., Thorarinsdottir, E. E., Smaradottir, H., Sigurgisladottir, S. & Gudmundsdottir, B. K. (2010). Selection of bacteria and the effects of bacterial treatment of Atlantic halibut (*Hippoglossus hippoglossus* L.) eggs and larvae. *Aquaculture* 302, 219-227.
- Blango, M. G. & Mulvey, M. A. (2009). Bacterial landlines: contact-dependent signaling in bacterial populations. *Curr Opin Microbiol* **12**, 177-181.
- Blasiak, L. C. & Clardy, J. (2009). Discovery of 3-formyl-tyrosine metabolites from Pseudoalteromonas tunicata through heterologous expression. J Am Chem Soc 132, 926-927.
- Blunt, J. W., Copp, B. R., Hu, W. P., Munro, M. H. G., Northcote, P. T. & Prinsep, M. R. (2009). Marine natural products. *Nat Prod Rep* 26, 170-244.
- Bollmann, A., Lewis, K. & Epstein, S. S. (2007). Incubation of environmental samples in a diffusion chamber increases the diversity of recovered isolates. *Appl Environ Microb* **73**, 6386-6390.
- Boric, M., Danevcic, T. & Stopar, D. (2011). Prodigiosin from *Vibrio* sp. DSM 14379; A New UV-Protective Pigment. *Microb Ecol* 62, 528-536.
- Bowman, J. P. (2007). Bioactive compound synthetic capacity and ecological significance of marine bacterial genus *Pseudoalteromonas*. *Mar Drugs* 5, 220-241.
- Boyd, K. G., Adams, D. R. & Burgess, J. G. (1999). Antibacterial and repellent activities of marine bacteria associated with algal surfaces. *Biofouling* 14, 227-236.
- Brinkhoff, T., Bach, G., Heidorn, T., Liang, L., Schlingloff, A. & Simon, M. (2004). Antibiotic production by a *Roseobacter* clade-affiliated species from the German Wadden Sea and its antagonistic effects on indigenous isolates. *Appl Environ Microb* 70, 2560.
- Broekaert, K., Heyndrickx, M., Herman, L., Devlieghere, F. & Vlaemynck, G. (2011). Seafood quality analysis: Molecular identification of dominant microbiota after ice storage on several general growth media. *Food Microbiol* 28, 1162-1169.
- Bruhn, J. B., Gram, L. & Belas, R. (2007). Production of antibacterial compounds and biofilm formation by *Roseobacter* species are influenced by culture conditions. *Appl Environ Microb* 73, 442-450.
- Bruhn, J. B., Nielsen, K. F., Hjelm, M., Hansen, M., Bresciani, J., Schulz, S. & Gram, L.
 (2005). Ecology, inhibitory activity, and morphogenesis of a marine antagonistic bacterium belonging to the *Roseobacter* clade. *Appl Environ Microb* 71, 7263-7270.
- Buchan, A., Gonz+ílez, J. M. & Moran, M. A. (2005). Overview of the marine Roseobacter lineage. *Appl Environ Microb* **71**, 5665-5677.
- Burke, C., Thomas, T., Egan, S. & Kjelleberg, S. (2007). The use of functional genomics for the identification of a gene cluster encoding for the biosynthesis of an antifungal tambjamine in the marine bacterium *Pseudoalteromonas tunicata*. *Environ Microbiol* **9**, 814-818.
- Burkholder, P. R., Pfister, R. M. & Leitz, F. H. (1966). Production of a pyrrole antibiotic by a marine bacterium. *Appl Environ Microb* 14, 649-653.
- **Cabello, F. C. (2006).**Heavy use of prophylactic antibiotics in aquaculture: a growing problem for human and animal health and for the environment. *Environ Microbiol* **8**, 1137-1144.
- Cha, C., Gao, P., Chen, Y. C., Shaw, P. D. & Farrand, S. K. (1998). Production of acyl-homoserine lactone quorum-sensing signals by gram-negative plant-associated bacteria. *Mol Plant Microbe In* **11**, 1119-1129.
- **Challis, G. L. & Ravel, J. (2000).**Coelichelin, a new peptide siderophore encoded by the *Streptomyces coelicolor* genome: structure prediction from the sequence of its non-ribosomal peptide synthetase. *FEMS Microbiol Lett* **187**, 111-114.
- Champ, M. A. (2000). A review of organotin regulatory strategies, pending actions, related costs and benefits. *Sci Total Environ* 258, 21-71.
- Chao, L. & Levin, B. R. (1981). Structured habitats and the evolution of anticompetitor toxins in bacteria. *Proc Nat Acad Sci U S A* 78, 6324.
- Chau, R., Kalaitzis, J. A. & Neilan, B. A. (2011). On the origins and biosynthesis of tetrodotoxin. Aquat Toxicol 104, 61-72.
- Chen, W. M., Lin, C. Y., Chen, C. A., Wang, J. T. & Sheu, S. Y. (2010). Involvement of an L-amino acid oxidase in the activity of the marine bacterium *Pseudoalteromonas flavipulchra* against methicillin-resistant *Staphylococcus aureus*. *Enzyme Microb Tech* **47**, 52-58.
- **Choi, S. H. & Greenberg, E. P. (1992).**Genetic evidence for multimerization of LuxR, the transcriptional activator of *Vibrio fischeri* luminescence. *Mol Mar Biol Biotech* **1**, 408-413.
- Colwell, R. R. (1983). Biotechnology in the marine sciences. Science 222, 19.

- **Connon, S. A. & Giovannoni, S. J. (2002).** High-throughput methods for culturing microorganisms in very-low-nutrient media yield diverse new marine isolates. *Appl Environ Microb* **68**, 3878-3885.
- Cortes, J., Wiesmann, K. E., Roberts, G. A., Brown, M. J., Staunton, J. & Leadlay, P. F. (1995). Repositioning of a domain in a modular polyketide synthase to promote specific chain cleavage. *Science* 268, 1487.
- Cortes, J., Haydock, S. F., Roberts, G. A., Bevitt, D. J. & Leadlay, P. F. (1990). An unusually large multifunctional polypeptide in the erythromycin-producing polyketide synthase of Saccharopolyspora erythraea. *Nature* 348, 176-178.
- Coulthurst, S. J., Williamson, N. R., Harris, A. K. P., Spring, D. R. & Salmond, G. P. C. (2006). Metabolic and regulatory engineering of *Serratia marcescens*: mimicking phagemediated horizontal acquisition of antibiotic biosynthesis and quorum-sensing capacities. *Microbiol-SGM* 152, 1899.
- Cox, A. R. J., Thomson, N. R., Bycroft, B., Stewart, G., Williams, P. & Salmond, G. P. C. (1998). pheromone-independent CarR protein controls carbapenem antibiotic synthesis in the opportunistic human pathogen *Serratia marcescens*. *Microbiol-SGM* **144**, 201.
- D'Onofrio, A., Crawford, J. M., Stewart, E. J., Witt, K., Gavrish, E., Epstein, S., Clardy, J. & Lewis,
 K. (2010).Siderophores from neighboring organisms promote the growth of uncultured bacteria. *Chem Biol* 17, 254-264.
- da Silva Melo, P., Maria, S. S., Vidal, B. C., Haun, M. & Durán, N. (2000). Violacein cytotoxicity and induction of apoptosis in V79 cells. *In Vitro Cell Dev Biol Anim* **36**, 539-543.
- Dalisay, D. S., Webb, J. S., Scheffel, A., Svenson, C., James, S., Holmstr+Âm, C., Egan, S. & Kjelleberg, S. (2006). A mannose-sensitive haemagglutinin (MSHA)-like pilus promotes attachment of *Pseudoalteromonas tunicata* cells to the surface of the green alga Ulva australis. *Microbiol-SGM* 152, 2875.
- Davies, J., Spiegelman, G. B. & Yim, G. (2006). The world of subinhibitory antibiotic concentrations. *Curr Opin Microbiol* 9, 445-453.
- Davis, M. A., Askin, M. C. & Hynes, M. J. (2005). Amino acid catabolism by an areA-regulated gene encoding an L-amino acid oxidase with broad substrate specificity in *Aspergillus nidulans*. *Appl Environ Microb* 71, 3551.
- del Castillo, C. S., Wahid, M. I., Takeshi, Y. & Taizo, S. (2008). Isolation and inhibitory effect of anti-Vibrio substances from *Pseudoalteromonas* sp. A1-J11 isolated from the coastal sea water of Kagoshima Bay. *Fisheries Sci* **74**, 174-179.

Delong, E. F. (2007). Modern microbial seascapes. Nat Rev Microbiol 5, 755-757.

- Dimise, E. J., Widboom, P. F. & Bruner, S. D. (2008). Structure elucidation and biosynthesis of fuscachelins, peptide siderophores from the moderate thermophile *Thermobifida fusca*. *Proc Natl Acad Sci U S A* **105**, 15311-15316.
- Dimitrieva, G. Y., Crawford, R. L. & Y³ksel, G. (2006). The nature of plant growth promoting effects of a Pseudoalteromonad associated with the marine algae *Laminaria japonica* and linked to catalase excretion. *J Appl Microbiol* **100**, 1159-1169.
- **Dobretsov, S., Teplitski, M. & Paul, V. (2009).** Mini-review: quorum sensing in the marine environment and its relationship to biofouling. *Biofouling* **25**, 413-427.
- **Dobretsov, S. V. & Qian, P. Y. (2002).** Effect of bacteria associated with the green alga *Ulva reticulata* on marine micro-and macrofouling. *Biofouling* **18**, 217-228.
- Dubey, G. P. & Ben-Yehuda, S. (2011). Intercellular nanotubes mediate bacterial communication. *Cell* 144, 590-600.
- Duerkop, B. A., Herman, J. P., Ulrich, R. L., Churchill, M. E. A. & Greenberg, E. P. (2008). The Burkholderia mallei BmaR3-BmaI3 quorum-sensing system produces and responds to N-3hydroxy-octanoyl homoserine lactone. J Bacteriol **190**, 5137.
- Duerkop, B. A., Ulrich, R. L. & Greenberg, E. P. (2007). Octanoyl-homoserine lactone is the cognate signal for *Burkholderia mallei* BmaR1-Bmal1 quorum sensing. *J Bacteriol* **189**, 5034.
- Duerkop, B. A., Varga, J., Chandler, J. R., Peterson, S. B., Herman, J. P., Churchill, M. E. A., Parsek, M. R., Nierman, W. C. & Greenberg, E. P. (2009). Quorum-sensing control of antibiotic synthesis in *Burkholderia thailandensis*. *J Bacteriol* **191**, 3909.
- Dusane, D. H., Matkar, P., Venugopalan, V. P., Kumar, A. R. & Zinjarde, S. S. (2011). Cross-species induction of antimicrobial compounds, biosurfactants and quorum-sensing inhibitors in tropical marine epibiotic bacteria by pathogens and biofouling microorganisms. *Curr Microbiol*, 1-7.
- Egan, S., Holmström, C. & Kjelleberg, S. (2001a). *Pseudoalteromonas ulvae* sp nov., a bacterium with antifouling activities isolated from the surface of a marine alga. *Int J Syst Evol Microbiol* 51, 1499-1504.
- Egan, S., James, S., Holmström, C. & Kjelleberg, S. (2002). Correlation between pigmentation and antifouling compounds produced by *Pseudoalteromonas tunicata*. *Environ Microbiol* **4**, 433-442.
- Egan, S., Thomas, T. & Kjelleberg, S. (2008). Unlocking the diversity and biotechnological potential of marine surface associated microbial communities. *Curr Opin Microbiol* **11**, 219-225.
- Egan, S., Wiener, P., Kallifidas, D. & Wellington, E. M. H. (2001b). Phylogeny of *Streptomyces* species and evidence for horizontal transfer of entire and partial antibiotic gene clusters. *A van Leeuw J Microb* **79**, 127-133.

- El-Sayed, A. K., Hothersall, J. & Thomas, C. M. (2001). Quorum-sensing-dependent regulation of biosynthesis of the polyketide antibiotic mupirocin in *Pseudomonas fluorescens* NCIMB 10586. *Microbiol-SGM* 147, 2127.
- Engebrecht, J. A. & Silverman, M. (1984). Identification of genes and gene products necessary for bacterial bioluminescence. *Proc Natl Acad Sci U S A* 81, 4154.
- Esquenazi, E., Coates, C., Simmons, L., Gonzalez, D., Gerwick, W. H. & Dorrestein, P. C.
 (2008). Visualizing the spatial distribution of secondary metabolites produced by marine cyanobacteria and sponges via MALDI-TOF imaging. *Mol BioSyst* 4, 562-570.
- **Euzéby, J. P. (1997).**List of bacterial names with standing in nomenclature: a folder available on the Internet. *Int J Syst Evol Microbiol* **47**, 590-592.
- **FAO and WHO (2001).** Health and nutritional properties of probiotics in food including powder milk with live lactic acid bacteria. edn, vol. 85. Córdoba, Argentina: FAO.
- Fedorov, R., Böhl, M., Tsiavaliaris, G., Hartmann, F. K., Taft, M. H., Baruch, P., Brenner, B., Martin, R., Knölker, H. J. & other authors (2009). The mechanism of pentabromopseudilin inhibition of myosin motor activity. *Nat Struct Mol Biol* 16, 80-88.
- Fehér, D., Barlow, R., McAtee, J. & Hemscheidt, T. K. (2010). Highly brominated antimicrobial metabolites from a Marine *Pseudoalteromonas* sp. J Nat Prod.
- Fehér, D., Barlow, R. S., Lorenzo, P. S. & Hemscheidt, T. K. (2008). A 2-substituted prodiginine, 2-(p-hydroxybenzyl) prodigiosin, from *Pseudoalteromonas rubra*. *J Nat Prod* **71**, 1970-1972.
- Fekete, A., Frommberger, M., Rothballer, M., Li, X., Englmann, M., Fekete, J., Hartmann, A., Eberl, L. & Schmitt-Kopplin, P. (2007). Identification of bacterial *N*-acylhomoserine lactones (AHLs) with a combination of ultra-performance liquid chromatography (UPLC), ultra-highresolution mass spectrometry, and in-situ biosensors. *Anal Bioanal Chem* 387, 455-467.
- Feling, R. H., Buchanan, G. O., Mincer, T. J., Kauffman, C. A., Jensen, P. R. & Fenical, W.
 (2003).Salinosporamide A: a highly cytotoxic proteasome inhibitor from a novel microbial source, a marine bacterium of the new genus *Salinospora*. *Angew Chem Int Ed* 42, 355-357.
- Fenical, W. (1993). Chemical studies of marine bacteria developing a new resource. *Chem Rev* 93, 1673-1683.
- Fenical, W., Jensen, P. R., Palladino, M. A., Lam, K. S., Lloyd, G. K. & Potts, B. C. (2009). Discovery and development of the anticancer agent salinosporamide A (NPI-0052). *Bioorgan Med Chem* 17, 2175-2180.
- Ferrari, B. C., Winsley, T., Gillings, M. & Binnerup, S. (2008). Cultivating previously uncultured soil bacteria using a soil substrate membrane system. *Nat Protocols* **3**, 1261-1269.

Fischbach, M. A. & Walsh, C. T. (2009). Antibiotics for emerging pathogens. Science 325, 1089.

- Franks, A., Egan, S., Holmstrom, C., James, S., Lappin-Scott, H. & Kjelleberg, S. (2006). Inhibition of fungal colonization by *Pseudoalteromonas tunicata* provides a competitive advantage during surface colonization. *Appl Environ Microb* **72**, 6079-6087.
- Frisvad, J. C., Andersen, B. & Thrane, U. (2008). The use of secondary metabolite profiling in chemotaxonomy of filamentous fungi. *Mycol Res* **112**, 231-240.
- Fukuda, D., Haines, A. S., Song, Z., Murphy, A. C., Hothersall, J., Stephens, E. R., Gurney, R., Cox,
 R. J., Crosby, J. & other authors (2011). A natural plasmid uniquely encodes two
 biosynthetic pathways creating a potent anti-MRSA antibiotic. *PLoS ONE* 6, e18031.
- Fuqua, C., Parsek, M. R. & Greenberg, E. P. (2001). Regulation of gene expression by cell-to-cell communication: acyl-homoserine lactone quorum sensing. *Annu Rev Genet* **35**, 439-468.
- Fuqua, C., Winans, S. C. & Greenberg, E. P. (1996). Census and consensus in bacterial ecosystems: the LuxR-LuxI family of quorum-sensing transcriptional regulators. *Annu Rev Microbiol* 50, 727-751.
- Garnier, M., Labreuche, Y., Garcia, C., Robert, M. & Nicolas, J. L. (2007). Evidence for the involvement of pathogenic bacteria in summer mortalities of the pacific oyster *Crassostrea gigas*. *Microb Ecol* 53, 187-196.
- Gatesoupe, F. J. (1997). Siderophore production and probiotic effect of *Vibrio* sp. associated with turbot larvae, *Scophthalmus maximus*. *Aquat Living Resour* **10**, 239-246.
- Gauthier, G., Gauthier, M. & Christen, R. (1995). Phylogenetic analysis of the genera Alteromonas, Shewanella, and Moritella using genes coding for small-subunit ribosomal-RNA sequences and division of the genus Alteromonas into 2 genera, Alteromonas (emended) and Pseudoalteromonas gen nov, and proposal of 12 new species combinations. Int J Syst Bacteriol 45, 755-761.
- **Gauthier, M. J. (1976).** Modification of bacterial respiration by a macromolecular polyanionic antibiotic produced by a marine *Alteromonas*. *Antimicrob Agents Chemother* **9**, 361.
- Gauthier, M. J. (1982). Validation of the name Alteromonas Iuteoviolacea. Int J Syst Bacteriol 32, 82.
- Gauthier, M. J. & Flatau, G. N. (1976). Antibacterial activity of marine violet-pigmented *Alteromonas* with special reference to the production of brominated compounds. *Can J Microbiol* 22, 1612-1619.
- Geng, H., Bruhn, J. B., Nielsen, K. F., Gram, L. & Belas, R. (2008). Genetic dissection of tropodithietic acid biosynthesis by marine roseobacters. *Appl Environ Microb* 74, 1535-1545.
- Georlette, D., Blaise, V., Collins, T., D'Amico, S., Gratia, E., Hoyoux, A., Marx, J. C., Sonan, G., Feller, G. & other authors (2004). Some like it cold: biocatalysis at low temperatures. *FEMS Microbiol Rev* 28, 25-42.

- **Gerber, N. N. (1971).** Prodigiosin-like pigments from *Actinomadura (Nocardia) pelletieri*. J Antibiot **24**, 636-640.
- Gil-Turnes, M. S., Hay, M. E. & Fenical, W. (1989). Symbiotic marine bacteria chemically defend crustacean embryos from a pathogenic fungus. *Science* 246, 116.
- Golberg, K., Eltzov, E., Shnit-Orland, M., Marks, R. & Kushmaro, A. (2011). Characterization of quorum sensing signals in coral-associated bacteria. *Microb Ecol* 61, 783-792.
- Gomez, D., Espinosa, E., Bertazzo, M., Lucas-Elio, P., Solano, F. & Sanchez-Amat, A. (2008). The macromolecule with antimicrobial activity synthesized by *Pseudoalteromonas luteoviolacea* strains is an L-amino acid oxidase. *Appl Environ Microb* **79**, 925-930.
- **Goodfellow, M. & Fiedler, H. P. (2010).** A guide to successful bioprospecting: informed by actinobacterial systematics. *A van Leeuw J Microb*, 1-24.
- Grabowski, K., Baringhaus, K. H. & Schneider, G. (2008). Scaffold diversity of natural products: inspiration for combinatorial library design. *Nat Prod Rep* **25**, 892-904.
- Gram, L., Grossart, H. P., Schlingloff, A. & Kiørboe, T. (2002). Possible quorum sensing in marine snow bacteria: production of acylated homoserine lactones by Roseobacter strains isolated from marine snow. *Appl Environ Microb* 68, 4111-4116.
- **Gram, L., Melchiorsen, J. & Bruhn, J. B. (2010).** Antibacterial activity of marine culturable bacteria collected from a global sampling of ocean surface waters and surface swabs of marine organisms. *Mar Biotechnol* **12**, 439-451.
- Gulder, T. A. M. & Moore, B. S. (2009). Chasing the treasures of the sea bacterial marine natural products. *Curr Opin Microbiol* 12, 252-260.
- Guo, X. C., Zheng, L., Zhou, W. H., Cui, Z. S., Han, P., Tian, L. & Wang, X. R. (2011). A case study on chemical defense based on quorum sensing: antibacterial activity of sponge-associated bacterium Pseudoalteromonas sp NJ6-3-1 induced by quorum sensing mechanisms. Ann Microbiol 61, 247-255.
- Hadfield, M. G. & Paul, V. J. (2001). Natural chemical cues for settlement and metamorphosis of marine-invertebrate larvae. *Marine chemical ecology*, 431-461.
- Hakvåg, S., Fjærvik, E., Klinkenberg, G., Borgos, S. E. F., Josefsen, K. D., Ellingsen, T. E. & Zotchev,
 S. B. (2009). Violacein-producing *Collimonas* sp. from the sea surface microlayer of costal waters in Trøndelag, Norway. *Mar Drugs* 7, 576-588.
- Hibbing, M. E., Fuqua, C., Parsek, M. R. & Peterson, S. B. (2009). Bacterial competition: surviving and thriving in the microbial jungle. *Nat Rev Microbiol* **8**, 15-25.
- Hjelm, M., Bergh, Ø., Riaza, A., Nielsen, J., Melchiorsen, J., Jensen, S., Duncan, H., Ahrens, P., Birkbeck, H. & other authors (2004). Selection and identification of autochthonous

potential probiotic bacteria from turbot larvae (*Scophthalmus maximus*) rearing units. *Syst Appl Microbiol* **27**, 360-371.

- Hoffman, L. R., D'Argenio, D. A., MacCoss, M. J., Zhang, Z., Jones, R. A. & Miller, S. I. (2005). Aminoglycoside antibiotics induce bacterial biofilm formation. *Nature* 436, 1171-1175.
- Holmström, C., Egan, S., Franks, A., McCloy, S. & Kjelleberg, S. (2002). Antifouling activities expressed by marine surface associated *Pseudoalteromonas* species. *FEMS Microbiol Ecol* 41, 47-58.
- Holmström, C., James, S., Neilan, B. A., White, D. C. & Kjelleberg, S. (1998). *Pseudoalteromonas tunicata* sp. nov., a bacterium that produces antifouling agents. *Int J Syst Bacteriol* **48**, 1205-1212.
- Holmström, C. & Kjelleberg, S. (1999). Marine *Pseudoalteromonas* species are associated with higher organisms and produce biologically active extracellular agents. *FEMS Microbiol Ecol* 30, 285-293.
- Holmström, C., Rittschof, D. & Kjelleberg, S. (1992). Inhibition of settlement by larvae of *Balanus amphitrite* and *Ciona intestinalis* by a surface-colonizing marine bacterium. *Appl Environ Microb* 58, 2111-2115.
- Hornemann, U., Hurley, L. H., Speedie, M. K. & Floss, H. G. (1971). Biosynthesis of indolmycin. J Am Chem Soc 93, 3028-3035.
- **Hoshino, T. (2011).** Violacein and related tryptophan metabolites produced by *Chromobacterium violaceum*: biosynthetic mechanism and pathway for construction of violacein core. *Appl Microbiol Biotechnol* **91**, 1463-1475.
- Hothersall, J., Murphy, A. C., Iqbal, Z., Campbell, G., Stephens, E. R., Wu, J., Cooper, H., Atkinson,
 S., Williams, P. & other authors (2011). Manipulation of quorum sensing regulation in
 Pseudomonas fluorescens NCIMB 10586 to increase mupirocin production. *Appl Microbiol Biotechnol*, 1-10.
- **Hoyoux, A., Jennes, I., Dubois, P., Genicot, S., Dubail, F., Francois, J. M., Baise, E., Feller, G. & Gerday, C. (2001).**Cold-adapted β-galactosidase from the Antarctic psychrophile *Pseudoalteromonas haloplanktis. Appl Environ Microb* **67**, 1529-1535.
- Huang, Y. L., Ki, J. S., Case, R. J. & Qian, P. Y. (2008a). Diversity and acyl-homoserine lactone production among subtidal biofilm-forming bacteria. *Aquat Microb Ecol* 52, 185-193.
- Huang, Y. L., Ki, J. S., Lee, O. O. & Qian, P. Y. (2008b). Evidence for the dynamics of Acyl homoserine lactone and AHL-producing bacteria during subtidal biofilm formation. *The ISME journal* 3, 296-304.

- Huang, Y. L., Li, M., Yu, Z. & Qian, P. Y. (2011). Correlation between pigmentation and larval settlement deterrence by *Pseudoalteromonas* sp. sf57. *Biofouling* 27, 287-293.
- Huber, B., Riedel, K., Hentzer, M., Heydorn, A., Gotschlich, A., Givskov, M., Molin, S. +. & Eberl,
 L. (2001). The cep quorum-sensing system of *Burkholderia cepacia* H111 controls biofilm formation and swarming motility. *Microbiol-SGM* 147, 2517-2528.
- Huggett, M., Williamson, J., de Nys, R., Kjelleberg, S. & Steinberg, P. (2006). Larval settlement of the common Australian sea urchin *Heliocidaris erythrogramma* in response to bacteria from the surface of coralline algae. *Oecologia* **149**, 604-619.
- Hughes, C. C. & Fenical, W. (2010). Antibacterials from the sea. Chem-Eur J 16, 12512-12525.
- Hurdle, J. G., O'Neill, A. J. & Chopra, I. (2004). Anti-staphylococcal activity of indolmycin, a potential topical agent for control of staphylococcal infections. *J Antimicrob Chemothe* **54**, 549-552.
- Imhoff, J. F., Labes, A. & Wiese, J. (2011). Bio-mining the microbial treasures of the ocean: New natural products. *Biotechnol Adv*.
- **Isnansetyo, A. & Kamei, Y. (2003).**MC21-A, a bactericidal antibiotic produced by a new marine bacterium, *Pseudoalteromonas phenolica* sp. nov. O-BC30T, against methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* **47**, 480-488.
- **Isnansetyo, A. & Kamei, Y. (2009).** Anti-methicillin-resistant *Staphylococcus aureus* (MRSA) activity of MC21-B, an antibacterial compound produced by the marine bacterium *Pseudoalteromonas phenolica* O-BC30^T. *Int J Antimicrob Ag* **34**, 131-135.
- Ivanova, E. P., Flavier, S. & Christen, R. (2004). Phylogenetic relationships among marine Alteromonas-like proteobacteria: emended description of the family Alteromonadaceae and proposal of Pseudoalteromonadaceae fam. nov., Colwelliaceae fam. nov., Shewanellaceae fam. nov., Moritellaceae fam. nov., Ferrimonadaceae fam. nov., Idiomarinaceae fam. nov. and Psychromonadaceae fam. nov. Int J Syst Evol Microbiol 54, 1773-1788.
- Jain, R., Rivera, M. C. & Lake, J. A. (1999). Horizontal gene transfer among genomes: The complexity hypothesis. *Proc Natl Acad Sci U S A* 96, 3801.
- Jensen, P. R. (2010). Linking species concepts to natural product discovery in the post-genomic era. *J Ind Microbiol Biotechnol* **37**, 219-224.
- Jensen, P. R., Williams, P. G., Oh, D. C., Zeigler, L. & Fenical, W. (2007). Species-specific secondary metabolite production in marine actinomycetes of the genus *Salinispora*. *Appl Environ Microb* **73**, 1146-1152.
- Jiang, Z., Boyd, K. G., Mearns-Spragg, A., Adams, D. R., Wright, P. C. & Burgess, J. G. (2000). Two diketopiperazines and one halogenated phenol from cultures of the marine bacterium, *Pseudoalteromonas luteoviolacea*. *Nat Prod Let* **14**, 435-440.

- Kalinovskaya, N. I., Ivanova, E. P., Alexeeva, Y. V., Gorshkova, N. M., Kuznetsova, T. A., Dmitrenok, A. S. & Nicolau, D. V. (2004).Low-molecular-weight, biologically active compounds from marine *Pseudoalteromonas* species. *Curr Microbiol* 48, 441-446.
- **Kanagasabhapathy, M. & Nagata, S. (2008).** Cross-species induction of antibacterial activity produced by epibiotic bacteria isolated from Indian marine sponge *Pseudoceratina purpurea*. *World J Microb Biot* **24**, 687-691.
- Kanamaru, T., Nakano, Y., Toyoda, Y., Miyagawa, K. I., Tada, M., Kaisho, T. & Nakao, M. (2001). In vitro and in vivo antibacterial activities of TAK-083, an agent for treatment of *Helicobacter pylori* infection. *Antimicrob Agents Chemother* **45**, 2455-2459.
- Kawauchi, K., Shibutani, K., Yagisawa, H., Kamata, H., Nakatsuji, S., Anzai, H., Yokoyama, Y., Ikegami, Y., Moriyama, Y. & other authors (1997). A possible immunosuppressant, cycloprodigiosin hydrochloride, obtained from *Pseudoalteromonas denitrificans*. *Biochem Biophys Res Commun* 237, 543-547.
- Kim, H. S., Hayashi, M., SHIBATA, Y., Wataya, Y., Mitamura, T., Horii, T., Kawauchi, K., Hirata, H., Tsuboi, S. & other authors (1999). Cycloprodigiosin hydrochloride obtained from *Pseudoalteromonas denitrificans* is a potent antimalarial agent. *Biol Pharm Bull* 22, 532-534.
- **Kitabatake, M., Ali, K., Demain, A., Sakamoto, K., Yokoyama, S. & S+Âll, D. (2002).** Indolmycin resistance of *Streptomyces coelicolor* A3 (2) by induced expression of one of its two tryptophanyl-tRNA synthetases. *J Biol Chem* **277**, 23882.
- Kohler, T., Curty, L. K., Barja, F., Van Delden, C. & Pechere, J. C. (2000). Swarming of *Pseudomonas* aeruginosa is dependent on cell-to-cell signaling and requires flagella and pili. *J Bacteriol* 182, 5990.
- Konstantinidis, K. T. & Tiedje, J. M. (2005). Towards a genome-based taxonomy for prokaryotes. J Bacteriol 187, 6258.
- Labbate, M., Queck, S. Y., Koh, K. S., Rice, S. A., Givskov, M. & Kjelleberg, S. (2004). Quorum sensing-controlled biofilm development in *Serratia liquefaciens* MG1. *J Bacteriol* **186**, 692.
- Lachnit, T., Blumel, M., Imhoff, J. F. & Wahl, M. (2009). Specific epibacterial communities on macroalgae: phylogeny matters more than habitat. *Aquat Biol* 5, 181-186.
- Latifi, A., Winson, M. K., Foglino, M., Bycroft, B. W., Stewart, G. S. A. B., Lazdunski, A. & Williams, P. (1995). Multiple homologues of LuxR and LuxI control expression of virulence determinants and secondary metabolites through quorum sensing in *Pseudomonas aeruginosa* PAO1. *Mol Microbiol* 17, 333-343.
- Lautru, S., Deeth, R. J. & Bailey, L. M. (2005). Discovery of a new peptide natural product by *Streptomyces coelicolor* genome mining. *Nature* 1, 265-269.

- Lee, O. O. & Qian, P. Y. (2003). Chemical control of bacterial epibiosis and larval settlement of *Hydroides elegans* in the red sponge *Mycale adherens*. *Biofouling* **19**, 171-180.
- Lichstein, H. C. & Vandesand, V. F. (1945). Violacein, an antibiotic pigment produced by *Chromobacterium violaceum. J Infect Dis* 76, 47-51.
- Long, R. A., Qureshi, A., Faulkner, D. J. & Azam, F. (2003).2-n-Pentyl-4-quinolinol produced by a marine *Alteromonas* sp. and its potential ecological and biogeochemical roles. *Appl Environ Microb* 69, 568.
- Long, R. A. & Azam, F. (2001). Antagonistic interactions among marine pelagic bacteria. *Appl Environ Microb* 67, 4975-4983.
- Longeon, A., Peduzzi, J., Barthelemy, M., Corre, S., Nicolas, J. L. & Guyot, M. (2004). Purification and partial identification of novel antimicrobial protein from marine bacterium *Pseudoalteromonas* species strain X153. *Mar Biotechnol* **6**, 633-641.
- Longford, S. R., Tujula, N. A., Crocetti, G. R., Holmes, A. J., Holmstrom, C., Kjelleberg, S., Steinberg, P. D. & Taylor, M. W. (2007). Comparisons of diversity of bacterial communities associated with three sessile marine eukaryotes. *Aquat Microb Ecol* 48, 217-229.
- Lucas-Elio, P., Hernandez, P., Sanchez-Amat, A. & Solano, F. (2005). Purification and partial characterization of marinocine, a new broad-spectrum antibacterial protein produced by *Marinomonas mediterranea*. *BBA-Gen Subjects* **1721**, 193-203.
- MacLeod, R. A. (1965). The question of the existence of specific marine bacteria. *Microbiol Mol Biol R* 29, 9.
- Mah, T. F. & O'Toole, G. A. (2001). Mechanisms of biofilm resistance to antimicrobial agents. *Trends Microbiol* 9, 34-39.
- Mai-Prochnow, A., Lucas-Elio, P., Egan, S., Thomas, T., Webb, J. S., Sanchez-Amat, A. & Kjelleberg, S. (2008). Hydrogen peroxide linked to lysine oxidase activity facilitates biofilm differentiation and dispersal in several Gram-negative bacteria. *J Bacteriol* **190**, 5493.
- Mai-Prochnow, A., Evans, F., isay-Saludes, D., Stelzer, S., Egan, S., James, S., Webb, J. S. & Kjelleberg, S. (2004). Biofilm Development and Cell Death in the Marine Bacterium *Pseudoalteromonas tunicata*. *Appl Environ Microb* **70**, 3232-3238.
- Malik, N. N. (2008). Drug discovery: past, present and future. Drug Discov Today 13, 909-912.
- Månsson, M., Phipps, R. K., Gram, L., Munro, M. H., Larsen, T. O. & Nielsen, K. F. (2010). Explorative Solid-Phase Extraction (E-SPE) for Accelerated Microbial Natural Product Discovery. J Nat Prod 73, 1126-1132.
- Martens, T., Gram, L., Grossart, H. P., Kessler, D., Müller, R., Simon, M., Wenzel, S. C. & Brinkhoff, T. (2007). Bacteria of the Roseobacter clade show potential for secondary metabolite production. *Microb Ecol* 54, 31-42.

- Matz, C., Deines, P., Boenigk, J., Arndt, H., Eberl, L., Kjelleberg, S. & Jurgens, K. (2004). Impact of violacein-producing bacteria on survival and feeding of bacterivorous nanoflagellates. *Appl Environ Microb* 70, 1593.
- Matz, C., Webb, J. S., Schupp, P. J., Phang, S. Y., Penesyan, A., Egan, S., Steinberg, P. & Kjelleberg, S. (2008). Marine biofilm bacteria evade eukaryotic predation by targeted chemical defense. *PLoS ONE* **3**.
- Mayer, A., Glaser, K. B., Cuevas, C., Jacobs, R. S., Kem, W., Little, R. D., McIntosh, J. M., Newman,
 D. J., Potts, B. C. & other authors (2010). The odyssey of marine pharmaceuticals: a current pipeline perspective. *Trends Pharmacol Sci* 31, 255-265.
- McClean, K. H., Winson, M. K., Fish, L., Taylor, A., Chhabra, S. R., Camara, M., Daykin, M., Lamb, J. H., Swift, S. & other authors (1997). Quorum sensing and *Chromobacterium violaceum*: exploitation of violacein production and inhibition for the detection of N-acylhomoserine lactones. *Microbiol-SGM* 143, 3703.
- Mearns-Spragg, A., Bregu, M., Boyd, K. G. & Burgess, J. G. (1998). Cross-species induction and enhancement of antimicrobial activity produced by epibiotic bacteria from marine algae and invertebrates, after exposure to terrestrial bacteria. *Lett Appl Microbiol* 27, 142-146.
- Medema, M. H., Blin, K., Cimermancic, P., de Jager, V., Zakrzewski, P., Fischbach, M. A., Weber, T., Takano, E. & Breitling, R. (2011).antiSMASH: rapid identification, annotation and analysis of secondary metabolite biosynthesis gene clusters in bacterial and fungal genome sequences. Nucleic Acids Res 39, W339.
- Menzella, H. G., Reid, R., Carney, J. R., Chandran, S. S., Reisinger, S. J., Patel, K. G., Hopwood, D.
 A. & Santi, D. V. (2005). Combinatorial polyketide biosynthesis by de novo design and rearrangement of modular polyketide synthase genes. *Nat Biotechnol* 23, 1171-1176.
- Meyer, B. & Kuever, J. (2008). Phylogenetic diversity and spatial distribution of the microbial community associated with the Caribbean deep-water sponge Polymastia cf. corticata by 16S rRNA, aprA, and amoA gene analysis. *Microb Ecol* 56, 306-321.
- Mikhailov, V. V., Romanenko, L. A. & Ivanova, E. P. (2002). The genus *Alteromonas* and related Proteobacteria. In *The Prokaryotes*, 3rd ed., vol. 6. edn, vol. 6, pp. 597-645. Edited by In M.Dworkin, S.Falkow, E.Rosenberg, K.-H.Schleifer & E.Stackebrandt (ed.). New York, NY: Springer.
- Miller, M. B. & Bassler, B. L. (2001). Quorum sensing in bacteria. Annu Rev Microbiol 55, 165-199.
- Milton, D. L. (2006). Quorum sensing in vibrios: complexity for diversification. *Int J Med Microbiol* 296, 61-71.
- Molinski, T. F., Dalisay, D. S., Lievens, S. L. & Saludes, J. P. (2008). Drug development from marine natural products. *Nat Rev Drug Discov* 8, 69-85.

- Moriarty, D. J. W. (1998). Control of luminous Vibrio species in penaeid aquaculture ponds. Aquaculture 164, 351-358.
- Morris, R. M., Rappe, M. S., Connon, S. A., Vergin, K. L., Siebold, W. A., Carlson, C. A. & Giovannoni, S. J. (2002). SAR11 clade dominates ocean surface bacterioplankton communities. *Nature* **420**, 806-810.
- Murphy, A. C., Fukuda, D., Song, Z., Hothersall, J., Cox, R. J., Willis, C. L., Thomas, C. M. & Simpson, T. J. (2011). Engineered thiomarinol antibiotics active against MRSA are generated by mutagenesis and mutasynthesis of *Pseudoalteromonas* SANK73390. *Angew Chem Int Ed* 50, 3271-3274.
- Nathan, C. & Goldberg, F. M. (2005). The profit problem in antibiotic R&D. Nat Rev Drug Discov 4, 887-891.
- Nealson, K. H. & Hastings, J. W. (1979). Bacterial bioluminescence: its control and ecological significance. *Microbiol Mol Biol R* 43, 496.
- Nelson, E. J. & Ghiorse, W. C. (1999). Isolation and identification of Pseudoalteromonas piscicida strain Cura-d associated with diseased damselfish (Pomacentridae) eggs. *J Fish Dis* 22, 253-260.
- Newman, D. J. & Cragg, G. M. (2007). Natural products as sources of new drugs over the last 25 years. J Nat Prod 70, 461-477.
- Nichols, D., Lewis, K., Orjala, J., Mo, S., Ortenberg, R., O'Connor, P., Zhao, C., Vouros, P., Kaeberlein, T. & other authors (2008). Short peptide induces an "uncultivable" microorganism to grow *in vitro*. *Appl Environ Microb* 74, 4889-4897.
- Nielsen, K. F., Månsson, M., Rank, C., Frisvad, J. C. & Larsen, T. O. (2011). Dereplication of microbial natural products by LC-DAD-TOFMS. J Nat Prod.
- **Noguchi, T., Arakawa, O. & Takatani, T. (2006).** TTX accumulation in pufferfish. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics* **1**, 145-152.
- **Oh, Y. S. & Park, A. (2011).***Pseudoalteromonas donghaensis* sp. nov., isolated from seawater. *Int J Syst Evol Microbiol* **61**, 351.
- Parker, W. L., Rathnum, M. L., Wells Jr, J. S., Trejo, W. H., Principe, P. A. & Sykes, R. B. (1982).SQ 27,860, a simple carbapenem produced by species of *Serratia* and *Erwinia*. *J Antibiot* **35**, 653.
- Parsek, M. R., Val, D. L., Hanzelka, B. L., Cronan, J. E. & Greenberg, E. P. (1999). Acyl homoserinelactone quorum-sensing signal generation. *Proc Natl Acad Sci U S A* 96, 4360.
- Patankar, A. V. & González, J. E. (2009). Orphan LuxR regulators of quorum sensing. *FEMS Microbiol Rev* 33, 739-756.

- Patil, C. D., Patil, S. V., Salunke, B. K. & Salunkhe, R. B. (2011). Prodigiosin produced by Serratia marcescens NMCC46 as a mosquito larvicidal agent against Aedes aegypti and Anopheles stephensi. Parasitol Res, 1-9.
- Penesyan, A., Marshall-Jones, Z., Holmstrom, C., Kjelleberg, S. & Egan, S. (2009). Antimicrobial activity observed among cultured marine epiphytic bacteria reflects their potential as a source of new drugs. *FEMS Microbiol Ecol* **69**, 113-124.
- Penn, K., Jenkins, C., Nett, M., Udwary, D. W., Gontang, E. A., McGlinchey, R. P., Foster, B., Lapidus, A., Podell, S. & other authors (2009). Genomic islands link secondary metabolism to functional adaptation in marine Actinobacteria. *ISME J* 3, 1193-1203.
- Pesci, E. C., Milbank, J. B. J., Pearson, J. P., McKnight, S., Kende, A. S., Greenberg, E. P. & Iglewski, B. H. (1999). Quinolone signaling in the cell-to-cell communication system of *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* 96, 11229.
- Pettit, G. R., Herald, C. L., Doubek, D. L., Herald, D. L., Arnold, E. & Clardy, J. (1982). Isolation and structure of bryostatin 1. *J Am Chem Soc* 104, 6846-6848.
- Piel, J. (2002). A polyketide synthase-peptide synthetase gene cluster from an uncultured bacterial symbiont of Paederus beetles. *Proc Natl Acad Sci U S A* 99, 14002-14007.
- Planas, M., Pérez-Lorenzo, M., Hjelm, M., Gram, L., Uglenes Fiksdal, I., Bergh, Ø. & Pintado, J. (2006). Probiotic effect in vivo of *Roseobacter* strain 27-4 against *Vibrio* (*Listonella*) *anguillarum* infections in turbot (*Scophthalmus maximus L.*) larvae. *Aquaculture* 255, 323-333.
- Porsby, C. H., Nielsen, K. F. & Gram, L. (2008). Phaeobacter and Ruegeria species of the Roseobacter clade colonize separate niches in a Danish turbot (Scophthalmus maximus)rearing farm and antagonize Vibrio anguillarum under different growth conditions. Appl Environ Microb 74, 7356-7364.
- Rappe, M. S., Connon, S. A., Vergin, K. L. & Giovannoni, S. J. (2002). Cultivation of the ubiquitous SAR11 marine bacterioplankton clade. *Nature* **418**, 630-633.
- Ravn, L., Christensen, A. B., Molin, S., Givskov, M. & Gram, L. (2001). Methods for detecting acylated homoserine lactones produced by Gram-negative bacteria and their application in studies of AHL-production kinetics. *J Microbiol Meth* **44**, 239-251.
- Rengpipat, S., Phianphak, W., Piyatiratitivorakul, S. & Menasveta, P. (1998). Effects of a probiotic bacterium on black tiger shrimp *Penaeus monodon* survival and growth. *Aquaculture* 167, 301-313.
- Romanenko, L. A., Zhukova, N. V., Rohde, M., Lysenko, A. M., Mikhailov, V. V. & Stackebrandt, E. (2003). *Pseudoalteromonas agarivorans* sp. nov., a novel marine agarolytic bacterium. *Int J Syst Evol Microbiol* **53**, 125-131.

- Rypien, K. L., Ward, J. R. & Azam, F. (2010). Antagonistic interactions among coral-associated bacteria. *Environ Microbiol* 12, 28-39.
- Sandaa, R. A., Brunvold, L., Magnesen, T. & Bergh, Ø. (2008). Monitoring the opportunistic bacteria *Pseudoalteromonas sp. LT-13* in a great scallop, *Pecten maximus* hatchery. *Aquaculture* **276**, 14-21.
- Sawabe, T., Narita, M., Tanaka, R., Onji, M., Tajima, K. & Ezura, Y. (2000). Isolation of *Pseudoalteromonas elyakovii* strains from spot-wounded fronds of *Laminaria japonica*. *Nippon Suisan Gakkaishi* 66, 249-254.
- Schaefer, A. L., Greenberg, E. P., Oliver, C. M., Oda, Y., Huang, J. J., Bittan-Banin, G., Peres, C. M., Schmidt, S., Juhaszova, K. & other authors (2008). A new class of homoserine lactone quorum-sensing signals. *Nature* 454, 595-599.
- Schwarzer, D., Finking, R. & Marahiel, M. A. (2003). Nonribosomal peptides: from genes to products. *Nat Prod Rep* 20, 275-287.
- Seyedsayamdost, M. R., Chandler, J. R., Blodgett, J. A. V., Lima, P. S., Duerkop, B. A., Oinuma, K.
 I., Greenberg, E. P. & Clardy, J. (2010). Quorum-sensing-regulated bactobolin production by Burkholderia thailandensis E264. Org Lett 12, 716-719.
- Shen, B., Du, L., Sanchez, C., Edwards, D. J., Chen, M. & Murrell, J. M. (2001). The biosynthetic gene cluster for the anticancer drug bleomycin from Streptomyces verticillus ATCC15003 as a model for hybrid peptide-polyketide natural product biosynthesis. J Ind Microbiol Biotechnol 27, 378-385.
- Shiozawa, H., Kagasaki, T., Kinoshita, T., Haruyama, H., Domon, H., Utsui, Y., Kodama, K. & Takahashi, S. (1993). Thiomarinol, a new hybrid antimicrobial antibiotic produced by a murine bacterium: Fermentation, isolation, structure, and antimicrobial activity. J Antibiot (Tokyo) 46, 1834-1842.
- Shiozawa, H., Kagasaki, T., Torikata, A., Tanaka, N., Fujimoto, K., Hata, T., Furukawa, Y. & Takahashi, S. (1995). Thiomarinol-B and thiomarinol-C, new antimicrobial antibiotics produced by a marine bacterium. *J Antibiot (Tokyo)* **48**, 907-909.
- Shiozawa, H., Shimada, A. & Takahashi, S. (1997). Thiomarinols D, E, F and G, new hybrid antimicrobial antibiotics produced by a marine bacterium: Isolation, structure, and antimicrobial activity. J Antibiot (Tokyo) 50, 449-452.
- Simidu, U., Kita-Tsukamoto, K., Yasumoto, T. & Yotsu, M. (1990). Taxonomy of four marine bacterial strains that produce tetrodotoxin. *Int J Syst Evol Microbiol* **40**, 331.
- Skov, M. N., Pedersen, K. & Larsen, J. L. (1995). Comparison of pulsed-field gel electrophoresis, ribotyping, and plasmid profiling for typing of *Vibrio anguillarum* serovar O1. *Appl Environ Microb* 61, 1540-1545.

- Skovhus, T. L., Holmström, C., Kjelleberg, S. & Dahllöf, I. (2007). Molecular investigation of the distribution, abundance and diversity of the genus *Pseudoalteromonas* in marine samples. *FEMS Microbiol Ecol* **61**, 348-361.
- Skovhus, T. L., Ramsing, N. B., Holmström, C., Kjelleberg, S. & Dahllöf, I. (2004). Real-time quantitative PCR for assessment of abundance of *Pseudoalteromonas* species in marine samples. *Appl Environ Microb* 70, 2373-2382.
- Slattery, M., Rajbhandari, I. & Wesson, K. (2001). Competition-mediated antibiotic induction in the marine bacterium *Streptomyces tenjimariensis*. *Microb Ecol* **41**, 90-96.
- Slightom, R. N. & Buchan, A. (2009). Surface colonization by marine roseobacters: integrating genotype and phenotype. *Appl Environ Microb* **75**, 6027-6037.
- Sogin, M. L., Morrison, H. G., Huber, J. A., Welch, D. M., Huse, S. M., Neal, P. R., Arrieta, J. M. & Herndl, G. J. (2006). Microbial diversity in the deep sea and the underexplored "rare biosphere". *Proc Natl Acad Sci U S A* 103, 12115-12120.
- Speitling, M., Smetanina, O. E., Kuznetsova, T. A. & Laatsch, H. (2007). Marine bacteria. XXXV. Bromoalterochromides A and A ', unprecedented chromopeptides from a marine Pseudoalteromonas maricaloris strain KMM 636. *J Antibiot* **60**, 36-42.
- Stackebrandt, E., Frederiksen, W., Garrity, G. M., Grimont, P. A. D., Kampfer, P., Maiden, M. C. J., Nesme, X., Rossello-Mora, R., Swings, J. & other authors (2002). Report of the ad hoc committee for the re-evaluation of the species definition in bacteriology. Int J Syst Evol Microbiol 52, 1043-1047.
- Stackebrandt, E. & Goebel, B. M. (1994). Taxonomic note: A place for DNA-DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int J Syst Bacteriol* 44, 846-849.
- Staufenberger, T., Thiel, V., Wiese, J. & Imhoff, J. F. (2008). Phylogenetic analysis of bacteria associated with *Laminaria saccharina*. *FEMS Microbiol Ecol* 64, 65-77.
- Staunton, J. & Weissman, K. J. (2001). Polyketide biosynthesis: a millennium review. *Nat Prod Rep* 18, 380-416.
- Stewart, P. S. (2003). Diffusion in biofilms. J Bacteriol 185, 1485.
- **Stingl, U., Tripp, H. J. & Giovannoni, S. J. (2007).** Improvements of high-throughput culturing yielded novel SAR11 strains and other abundant marine bacteria from the Oregon coast and the Bermuda Atlantic Time Series study site. *ISME J* **1**, 361-371.
- Sudek, S., Lopanik, N. B., Waggoner, L. E., Hildebrand, M., Anderson, C., Liu, H., Patel, A., Sherman, D. H. & Haygood, M. G. (2006). Identification of the putative bryostatin polyketide synthase gene cluster from "*Candidatus* Endobugula sertula", the uncultivated microbial symbiont of the marine bryozoan *Bugula neritina*. J Nat Prod **70**, 67-74.

- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol*.
- Tatewaki, M., Provasoli, L. & Pintner, I. J. (1983). Morphogenesis of *Monostroma oxyspermum* (Kütz.) Doty (Chlorophyceae) in axenic culture, especially in bialgal culture. *J Phycol* **19**, 409-416.
- Taylor, M. W., Schupp, P. J., Dahllöf, I., Kjelleberg, S. & Steinberg, P. D. (2004). Host specificity in marine sponge-associated bacteria, and potential implications for marine microbial diversity. *Environ Microbiol* 6, 121-130.
- Tebben, J., Tapiolas, D. M., Motti, C. A., Abrego, D., Negri, A. P., Blackall, L. L., Steinberg, P. D. & Harder, T. (2011). Induction of larval metamorphosis of the coral *Acropora millepora* by tetrabromopyrrole isolated from a *Pseudoalteromonas* bacterium. *PLoS ONE* 6, e19082.
- Thiel, V., Neulinger, S. C., Staufenberger, T., Schmaljohann, R. & Imhoff, J. F. (2007). Spatial distribution of sponge–associated bacteria in the Mediterranean sponge *Tethya aurantium*. *FEMS Microbiol Ecol* **59**, 47-63.
- Thomas, C. M., Hothersall, J., Willis, C. L. & Simpson, T. J. (2010). Resistance to and synthesis of the antibiotic mupirocin. *Nat Rev Microbiol* 8, 281-289.
- Thomson, N. R., Crow, M. A., McGowan, S. J., Cox, A. & Salmond, G. P. C. (2000). Biosynthesis of carbapenem antibiotic and prodigiosin pigment in *Serratia* is under quorum sensing control. *Mol Microbiol* **36**, 539-556.
- Toranzo, A. E., Magariños, B. & Romalde, J. L. (2005). A review of the main bacterial fish diseases in mariculture systems. *Aquaculture* 246, 37-61.
- Torkildsen, L., Lambert, C., Nylund, A., Magnesen, T. & Bergh, +. (2005). Bacteria associated with early life stages of the great scallop, *Pecten maximus*: impact on larval survival. *Aquacult Int* **13**, 575-592.
- **Unemoto, T. & Hayashi, M. (1993).**Na+-translocating NADH-quinone reductase of marine and halophilic bacteria. *J Bioenerg Biomembr* **25**, 385-391.
- Van Houdt, R., Givskov, M. & Michiels, C. W. (2007). Quorum sensing in *Serratia*. *FEMS Microbiol Rev* **31**, 407-424.
- Vandamme, P., Pot, B., Gillis, M., De Vos, P., Kersters, K. & Swings, J. (1996). Polyphasic taxonomy, a consensus approach to bacterial systematics. *Microbiol Mol Biol R* 60, 407.
- Vecchione, J. J. & Sello, J. K. (2009). A novel tryptophanyl-tRNA synthetase gene confers high-level resistance to indolmycin. *Antimicrob Agents Chemother*, AAC-00723.
- Vetsigian, K., Jajoo, R. & Kishony, R. (2011). Structure and evolution of *Streptomyces* interaction networks in soil and *in silico*. *PLoS biology* 9, e1001184.

von Wittenau, M. S. & Els, H. (1961). The structure of indolmycin. J Am Chem Soc 83, 4678-4680.

- **Vynne, N. G., Christiansen, G. & Gram, L. (2012a).** *Pseudoalteromonas galatheae* S3431 sp. nov., a novel *Pseudoalteromonas* species isolated from a deep-sea polychaete. *In preparation for International Journal of Systematic and Evolutionary Microbiology*.
- Vynne, N. G. & Gram, L. (2012). Draft genome sequence of *Pseudoalteromonas luteoviolacea* S4054, an indolmycin producing marine bacterium. *In preparation for Journal of Bacteriology*.
- Vynne, N. G., Månsson, M., Nielsen, K. F. & Gram, L. (2012b). Antibiotic chemoprofile and RecA and GyrB sequence analyses reveal three phylotypes within the marine *Pseudoalteromonas luteoviolacea*. In preparation.
- Vynne, N. G., Månsson, M., Nielsen, K. F., Milton, D. L. & Gram, L. (2012c).*N*-acyl homoserine lactone-like compounds produced by a Luxl homologue in *Pseudoalteromonas luteoviolacea* strain S4054. *In preparation*.
- Vynne, N. G., Månsson, M., Nielsen, K. F. & Gram, L. (2011). Bioactivity, chemical profiling and 16S rRNA based phylogeny of *Pseudoalteromonas* strains collected on a global research cruise. *Mar Biotechnol*, doi:10.1007/s10126-011-9369-4.
- Wagner-Döbler, I., Thiel, V., Eberl, L., Allgaier, M., Bodor, A., Meyer, S., Ebner, S., Hennig, A., Pukall, R. & other authors (2005). Discovery of complex mixtures of novel long-chain quorum sensing signals in free–living and host-associated marine alphaproteobacteria. *Chembiochem* 6, 2195-2206.
- Wahl, M. (1989). Marine epibiosis. I. Fouling and antifouling: Some basic aspects. *Mar Ecol-Prog Ser* 58, 175-189.
- Waksman, S. A. & Schatz, A. (1943). Strain specificity and production of antibiotic substances. *Proc Natl Acad Sci U S A* 29, 74.
- Wang, H., Jiang, P., Lu, Y., Ruan, Z., Jiang, R., Xing, X. H., Lou, K. & Wei, D. (2009). Optimization of culture conditions for violacein production by a new strain of *Duganella* sp. B2. *Biochem Eng J* 44, 119-124.
- Wang, J., Quan, C., Wang, X., Zhao, P. & Fan, S. (2011). Extraction, purification and identification of bacterial signal molecules based on N–acyl homoserine lactones. *Microbial Biotechnology* **4**, 479-490.
- Wang, Y., Ikawa, A., Okaue, S., Taniguchi, S., Osaka, I., Yoshimoto, A., Kishida, Y., Arakawa, R. & Enomoto, K. (2008). Quorum sensing signaling molecules involved in the production of violacein by *Pseudoalteromonas*. *Biosci Biotechnol Biochem* 72, 1958-1961.

- Watson, W. T., Minogue, T. D., Val, D. L., von Bodman, S. B. & Churchill, M. E. A. (2002). Structural basis and specificity of acyl-homoserine lactone signal production in bacterial quorum sensing. *Mol Cell* 9, 685-694.
- Wayne, L. G., Brenner, D. J., Colwell, R. R., Grimont, P. A. D., Kandler, O., Krichevsky, M. I., Moore, L. H., Moore, W. E. C., Murray, R. G. E. & other authors (1987). Report of the ad hoc committee on reconciliation of approaches to bacterial systematics. *Int J Syst Evol Microbiol* 37, 463.
- Webster, N. S., Wilson, K. J., Blackall, L. L. & Hill, R. T. (2001). Phylogenetic diversity of bacteria associated with the marine sponge *Rhopaloeides odorabile*. *Appl Environ Microb* 67, 434.
- Weissman, K. J. & Leadlay, P. F. (2005). Combinatorial biosynthesis of reduced polyketides. *Nat Rev Microbiol* 3, 925-936.
- Welch, M., Todd, D. E., Whitehead, N. A., McGowan, S. J., Bycroft, B. W. & Salmond, G. P. C. (2000).N-acyl homoserine lactone binding to the CarR receptor determines quorumsensing specificity in *Erwinia*. *EMBO J* **19**, 631-641.
- Werner, R. G., Thorpe, L. F., Reuter, W. & Nierhaus, K. H. (1976). Indolmycin Inhibits prokaryotic tryptophanyl-tRNA ligase. *Eur J Biochem* 68, 1-3.
- Whitehead, N. A., Barnard, A. M. L., Slater, H., Simpson, N. J. L. & Salmond, G. P. C. (2001).Quorum–sensing in Gram-negative bacteria. *FEMS Microbiol Rev* 25, 365-404.
- Wietz, M., Gram, L., Jørgensen, B. & Schramm, A. (2010). Latitudinal patterns in the abundance of major marine bacterioplankton groups. *Aquat Microb Ecol* 61, 179-189.
- Wietz, M., Mansson, M. & Gram, L. (2011). Chitin stimulates production of the antibiotic andrimid in a *Vibrio coralliilyticus* strain. *Environ Microbiol Rep* **3**, 559-564.
- Winson, M. K., Swift, S., Fish, L., Throup, J. P., Jørgensen, F., Chhabra, S. R., Bycroft, B. W.,
 Williams, P. & Stewart, G. S. A. B. (1998). Construction and analysis of *luxCDABE*—based plasmid sensors for investigating N—acyl homoserine lactone-mediated quorum sensing.
 FEMS Microbiol Lett 163, 185-192.
- Wratten, S. J., Wolfe, M. S., Andersen, R. J. & Faulkner, D. J. (1977). Antibiotic metabolites from a marine Pseudomonad. *Antimicrob Agents Chemother* **11**, 411-414.
- Xu, X. W., Wu, Y. H., Wang, C. S., Gao, X. H., Wang, X. G. & Wu, M. (2010). *Pseudoalteromonas lipolytica* sp. nov., isolated from the Yangtze River estuary. *Int J Syst Evol Microbiol* 60, 2176.
- Yada, S., Wang, Y., Zou, Y., Nagasaki, K., Hosokawa, K., Osaka, I., Arakawa, R. & Enomoto, K. (2008). Isolation and characterization of two groups of novel marine bacteria producing violacein. *Mar Biotechnol* 10, 128-132.

- Yan, L., Boyd, K. G. & Grant Burgess, J. (2002). Surface attachment induced production of antimicrobial compounds by marine epiphytic bacteria using modified roller bottle cultivation. *Marine Biotechnology* 4, 356-366.
- Yates, E. A., Philipp, B., Buckley, C., Atkinson, S., Chhabra, S. R., Sockett, R. E., Goldner, M., Dessaux, Y., Camara, M. & other authors (2002).N-acylhomoserine lactones undergo lactonolysis in a pH-, temperature-, and acyl chain length-dependent manner during growth of Yersinia pseudotuberculosis and Pseudomonas aeruginosa. Infect Immun 70, 5635.
- Yooseph, S., Sutton, G., Rusch, D. B., Halpern, A. L., Williamson, S. J., Remington, K., Eisen, J. A., Heidelberg, K. B., Manning, G. & other authors (2007). The Sorcerer II Global Ocean Sampling expedition: expanding the universe of protein families. *PLoS biology* 5, e16.
- Yoshikawa, K., Nakayama, Y., Hayashi, M., Unemoto, T. & Mochida, K. (1999). Korormicin, an antibiotic specific for gram-negative marine bacteria, strongly inhibits the respiratory chain-linked Na+-translocating NADH: quinone reductase from the marine *Vibrio alginolyticus*. J Antibiot (Tokyo) **52**, 182.
- Yoshikawa, K., Takadera, T., Adachi, K., Nishijima, M. & Sanc, H. (1997). Korormicin, a novel antibiotic specifically active against marine Gram-negative bacteria, produced by a marine bacterium. *J Antibiot (Tokyo)* 50, 949-953.
- Yung, P. Y., Burke, C., Lewis, M., Kjelleberg, S. & Thomas, T. (2011). Novel antibacterial proteins from the microbial communities associated with the sponge *Cymbastela concentrica* and the green alga *Ulva australis*. *Appl Environ Microb* 77, 1512-1515.
- Zhang, X. & Enomoto, K. (2011). Characterization of a gene cluster and its putative promoter region for violacein biosynthesis in *Pseudoalteromonas* sp. 520P1. *Appl Microbiol Biotechnol* 90, 1963-1971.
- Zheng, L., Yan, X. J., Han, X. T., Chen, H. M., Lin, W., Lee, F. S. C. & Wang, X. R.
 (2006).Identification of norharman as the cytotoxic compound produced by the sponge (*Hymeniacidon perleve*)-associated marine bacterium *Pseudoalteromonas piscicida* and its apoptotic effect on cancer cells. *Biotechnol Appl Biochem* 44, 135-142.
- Zhu, J. & Winans, S. C. (2001). The quorum-sensing transcriptional regulator TraR requires its cognate signaling ligand for protein folding, protease resistance, and dimerization. Proc Natl Acad Sci U S A 98, 1507-1512.
- **ZoBell, C. E. & Upham, H. C. (1944).** A list of marine bacteria including descriptions of sixty new species. University of California Press.

Vynne N, Månsson M, Nielsen KF, Gram L (2011)

Bioactivity, chemical profiling, and 16S rRNA-based phylogeny *of Pseudoalteromonas* strains collected on a global research cruise.

Marine Biotechnology 13: 1062-1073.

ORIGINAL ARTICLE

Bioactivity, Chemical Profiling, and 16S rRNA-Based Phylogeny of *Pseudoalteromonas* Strains Collected on a Global Research Cruise

Nikolaj G. Vynne • Maria Månsson • Kristian F. Nielsen • Lone Gram

Received: 24 September 2010 / Accepted: 18 January 2011 / Published online: 9 February 2011 © Springer Science+Business Media, LLC 2011

Abstract One hundred one antibacterial Pseudoalteromonas strains that inhibited growth of a Vibrio anguillarum test strain were collected on a global research cruise (Galathea 3), and 51 of the strains repeatedly demonstrated antibacterial activity. Here, we profile secondary metabolites of these strains to determine if particular compounds serve as strain or species markers and to determine if the secondary metabolite profile of one strain represents the bioactivity of the entire species. 16S rRNA gene similarity divided the strains into two primary groups: One group (51 strains) consisted of bacteria which retained antibacterial activity, 48 of which were pigmented, and another group (50 strains) of bacteria which lost antibacterial activity upon sub-culturing, two of which were pigmented. The group that retained antibacterial activity consisted of six clusters in which strains were identified as Pseudoalteromonas luteoviolacea, Pseudoalteromonas aurantia, Pseudoalteromonas phenolica, Pseudoalteromonas ruthenica, Pseudoalteromonas rubra, and Pseudoalteromonas piscicida. HPLC-UV/VIS analyses identified key peaks, such as violacein in P. luteoviolacea. Some compounds, such as a novel bromoalterochromide,

Electronic supplementary material The online version of this article (doi:10.1007/s10126-011-9369-4) contains supplementary material, which is available to authorized users.

N. G. Vynne (⊠) · L. Gram
National Food Institute, Technical University of Denmark, Søltofts Plads, bldg. 221,
2800 Kgs. Lyngby, Denmark
e-mail: ngvy@food.dtu.dk

M. Månsson · K. F. Nielsen Department of Systems Biology, Technical University of Denmark, Søltofts Plads, bldg. 221, 2800 Kgs. Lyngby, Denmark were detected in several species. HPLC-UV/VIS detected systematic intra-species differences for some groups, and testing several strains of a species was required to determine these differences. The majority of non-antibacterial, non-pigmented strains were identified as *Pseudoalteromonas agarivorans*, and HPLC-UV/VIS did not further differentiate this group. *Pseudoalteromonas* retaining antibacterial were more likely to originate from biotic or abiotic surfaces in contrast to planktonic strains. Hence, the pigmented, antibacterial *Pseudoalteromonas* have a niche specificity, and sampling from marine biofilm environments is a strategy for isolating novel marine bacteria that produce antibacterial compounds.

Keywords *Pseudoalteromonas* · Antibacterial activity · Secondary metabolites · Bioprospecting · Galathea 3

Introduction

Compounds of relevance for the pharmaceutical and biotechnology industries are produced by marine microorganisms (Burgess et al. 1999), and it has been suggested that some compounds of pharmacological interest previously attributed to macroorganisms may in fact be of microbial origin (Bewley and Faulkner 1998; Simmons et al. 2008; Sudek et al. 2006). The emergence of multiresistant pathogenic bacterial strains and the failure of combinatorial and diversity-oriented chemistry to adequately supply the drug discovery pipeline (Newman 2008) have re-invigorated natural product chemistry as a path for discovery and development of new antibiotics. With this in mind, we isolated marine bacteria with antibacterial activity during the Danish Galathea 3 marine research expedition (Gram et al. 2010). The antibacterial strains were tentatively identified using 16S rRNA similarity, and one of the major groups of isolated bacteria was identified as *Pseudoalteromonas* (Gram et al. 2010).

The genus Pseudoalteromonas consists of Gramnegative marine bacteria belonging to the γ -proteobacteria and is present globally in marine waters where they constitute 0.5% to 6% of the total bacterioplankton (Wietz et al. 2010). They are heterotrophic aerobes and nonfermentative, and the cells are motile by one or more polar flagella. The genus divides into two groups: pigmented and non-pigmented species. The pigmented species are often producers of bioactive secondary metabolites (Bowman 2007) displaying cytotoxic (Zheng et al. 2006), antibacterial (Gauthier 1976b; Gauthier and Flatau 1976; Isnansetyo and Kamei 2003; Jiang et al. 2000; McCarthy et al. 1994), antifungal (Franks et al. 2006; Kalinovskava et al. 2004), or antifouling (Egan et al. 2001; Holmström et al. 2002) effects. It has been hypothesized that bioactive Pseudoalteromonas are primarily associated with higher organisms (Holmström and Kjelleberg 1999), suggesting an ecological role in which some bioactive species might play an active part in host defense against pathogens and fouling organisms (Holmström et al. 1996; Armstrong et al. 2001; Egan et al. 2008). A link between surface colonization and antibacterial activity has not been experimentally verified, although several studies have successfully isolated epiphytic bacteria with antibacterial activity from algae and other marine organisms (Armstrong et al. 2001; Boyd et al. 1999; James et al. 1996; Penesyan et al. 2009). The group of non-pigmented species has highly similar 16S rRNA gene sequences (Ivanova et al. 2004) and is rarely inhibitory against other microorganisms, although, Pseudoalteromonas haloplanktis strain INH produces isovaleric acid and 2-methylbutyric acid showing a broad spectrum of bacterial inhibition (Hayashida-Soiza et al. 2008).

Phylogeny and differentiation of bacterial species rely heavily on 16S rRNA gene similarity (Stackebrandt et al. 2002); however, 16S rRNA gene similarity does not provide sufficient differentiation below species level (Fox et al. 1992). Comparison of secondary metabolite production has supported species delineation within the actinomycete genus Salinispora, where strains with high 16S rRNA similarity were shown to belong to distinct species each with different specific metabolite profiles (Jensen et al. 2007). In mycology, comparison of chemical profiles (e.g., TLC, direct-infusion mass spectrometry, and HPLC with various detectors such as UV/VIS and/or mass spectrometry) of secondary metabolites has been widely used to identify and differentiate filamentous fungi (chemotaxonomy) also at sub-species level (Frisvad et al. 2008), and the chemophylogeny correlates well with phylogenetic analysis of sequences of specific housekeeping genes (e.g., chitin synthase, β -tubulin, and calmodulin; Geiser et al. 2007). Since several Pseudoalteromonas species produce a range of secondary metabolites, we hypothesized that chemical profiling and specific marker compounds could be indicative of bioactive potential and at the same time be useful in species identification or differentiation (Jensen et al. 2007).

The aim of the present study was to profile the secondary metabolites of these strains to determine if particular compounds serve as markers of strains or species with antibacterial activity and to determine if several strains of each species must be tested to assess the full bioactivity potential. As part of this, we provide accurate identification and phylogeny of these organisms by detailed 16S rRNA gene sequence comparative analysis. Since the bacteria were isolated from different sample types, our collection also allows us to address aspects of *Pseudoalteromonas* ecology such as the possible link between surface or planktonic lifestyle and antibacterial activity.

Materials and Methods

Strain Isolation Approximately 500 marine bacterial strains with antagonist activity against *Vibrio anguillarum* strain 90-11-287 (Skov et al. 1995) were isolated during the Danish Galathea 3 research expedition (Gram et al. 2010). One hundred one of these strains tentatively identified as *Pseudoalteromonas* species were included in the present study. *V. anguillarum* 90-11-287 was used as target strain since the expedition ship was not equipped to handle, e.g., potential human pathogens, and this *Vibrio* strain is in our experience very sensitive to antibacterial compounds from other marine bacteria (Hjelm et al. 2004).

Growth Media and Culture Conditions Pseudoalteromonas strains were grown in marine broth (MB) 2216 (Difco, Detroit, MI, USA) and on marine agar (MA) 2216 (Difco, Detroit, MI, USA) prepared in accordance with the manufacturer's instructions. Broth cultures were incubated under stagnant conditions at 25°C. Pigment production was determined by visual inspection of 48-h-old culture broths (MB) and colonies grown on MA for 24 to 48 h.

Antibacterial Activity Instant Ocean (IO) bioassay agar plates were prepared as described by Hjelm et al. (2004). Ten grams per liter agar, 3.3 g/l casamino acids (Difco 223050, Detroit, MI, USA), and 30 g/l Instant Ocean aquatic salts (Instant Ocean[®] Aquarium systems Inc., Sarrebourg, France) were added to distilled water and autoclaved. Glucose (0.4%) and 10 μ l/ml of *V. anguillarum* overnight culture were added to the cooled (44°C) IO and plates poured. The plates were allowed to dry for 15 min, and if used for well diffusion agar assays (WDAA), wells (diameter 6 mm) were punched. The inhibitory activity of live *Pseudoalteromonas* bacterial cells was tested by spotting 48-h-old MA grown colonies on freshly prepared IO agar plates containing *V. anguillarum*. Plates were incubated at 25°C and inspected for clearing zones in the growth of *V. anguillarum* after 24 h.

Cell-free supernatants were prepared to test for the presence of water-soluble antibacterial compounds secreted to the broth, and ethyl acetate extracts were prepared to test for production of non-polar antibacterial compounds. Each strain was grown in 20 ml of MB for 48 h. A 1.5-ml sample was withdrawn for 0.2 μ m filtering, and subsequently the remainder of the culture was extracted with an equal volume of ethyl acetate. The ethyl acetate fraction was transferred to a new vessel, evaporated to dryness, and redissolved in 2×0.5 ml ethyl acetate. The 1.5-ml cell-free sterile-filtered supernatant and the ethyl acetate extracts were stored at -20° C until tested in the WDAA (50 μ l sample per well) based on IO agar plates containing *V. anguillarum*. Controls (sterile MB and pure ethyl acetate) did not cause any inhibition zones.

The number of antibacterial *Pseudoalteromonas* strains in surface samples (e.g., algae, driftwood, fish, and sediment samples) was compared to their numbers in water samples by the Fisher's exact test (Fisher 1958). A 2×2 contingency table was used to test the hypothesis that presumed antibacterial strains with stable antibacterial activity were equally likely to be isolated from water samples and surface samples.

16S rRNA Gene Sequence Analyses A detailed phylogenetic analysis was performed on 16S rRNA sequences obtained in a previous study (Gram et al. 2010). For the analysis in this study, we conducted a BLAST (http://blast.ncbi.nlm.nih.gov) search against a compilation of Pseudoalteromonas type strain sequences retrieved from GenBank (list of type strains obtained from http://www.bacterio.cict.fr), and sequences of the type strains with a BLAST match in our strain collection were included in 16S rRNA sequence analysis to obtain a robust phylogenetic tree. Sequences from two additional Pseudoalteromonas strains were included: The genus type strain P. haloplanktis and the bioactive Pseudoalteromonas tunicata. Salinispora arenicola CNS-205 was used as outgroup. The sequences were aligned by the MAFFT online software (http://www.ebi.ac.uk/Tools/mafft/; Katoh et al. 2002) and curated with the Gblocks software on its least stringent settings (Castresana 2000; Talavera and Castresana 2007). The resulting alignment was processed using the MEGA4 software (Tamura et al. 2007) to create neighborjoining and minimum evolution trees. PhyML 3.0 was used to generate a maximum likelihood tree (Guindon and Gascuel 2003). Phylogenetic trees were generated under default parameters with 1,000 bootstrap replications for neighbor-joining and minimum evolution trees and 100 bootstrap replications for the maximum likelihood tree.

GenBank accession numbers for the *Pseudoalteromonas* strains used in this study are included in Supplementary Table 1.

HPLC-UV/VIS Analysis of Secondary Metabolites The strains were grown in static cultures in 10 ml MB for 3 days at 25°C, and for each species, one strain was cultured in triplicate to establish extraction and growth variation. Cultures were extracted with equal volumes of ethyl acetate, centrifuged, and the ethyl acetate was evaporated under N2 to dryness. Samples were redissolved in 300 μ l acetonitrile–water (1:1 v/v) and filtered through a 13-mm ID PFTE syringe filter. A subsample of 2 µl was then analyzed by reversed phase HPLC on an Agilent 1100 System equipped with a UV/VIS photo diode array detector (scanning 200-600 nm). Separation was done on a 100 mm×2 mm i.d., 3 μm Gemini C₆-phenyl column (Phenomenex, Torrance, CA, USA), running at 40°C using a binary linear solvent system of water (A) and acetonitrile (B; both buffered with 50 µl/l trifluoroacetic acid) at a flow of 300 μ l/min. The gradient profile was t=0 min, 5% B; t=22 min, 70% B; t=24.5, 100% B; t=27 min, 100% B; and t=29 min, B=5%, holding this for 8 min prior to the next injection. The chromatographic profiles were compared, subtracting peaks present in media blank extracts. Samples were analyzed in random order, and six of the first extracts were analyzed several times during the sequence to determine any retention time shifts. Cluster analysis was done on a matrix of detected / non-detected peaks (1/0) using NTSYSpc 2.20g (Exeter Software, Setauket, NY, USA). SAHN clustering was used by unweighted pair-group method (UPGMA) and simple distance measurement. Representative extracts were also analyzed by HPLC-UV/VIS-TOFMS in both positive and negative electrospray (Nielsen and Smedsgaard 2003). Peaks were tentatively identified by UV spectra and accurate mass data by matching in Antibase 2009 (35 930 microbial secondary metabolites; Wiley & Sons, Hoboken, NJ, USA; Nielsen et al. 2006).

Results

Pigmentation and Antibacterial Activity The one hundred one *Pseudoalteromonas* strains were originally isolated for their ability to inhibit *V. anguillarum* (Gram et al. 2010). However, on re-cultivation and re-testing for the ability to inhibit *V. anguillarum* after storage at -80° C for several months, only 51 strains retained inhibitory activity (Table 1). These 51 strains were all inhibitory when tested as live cultures in the "spot test" assay. Twenty of the strains produced water-soluble, diffusible, antibacterial substances as indicated by the ability of cell-free sterile-filtered supernatant to inhibit growth of *V. anguillarum* in the

16S rRNA cluster	No. of strains		Related type strain ^a	No. of strains from		No. of strains inhibiting <i>Vibrio</i>		Inhibition of <i>Vibrio</i>
	Pigmented	Non-pigmented		Surface samples	Water samples	Pigmented	Non-pigmented	extracts
I	1	37	P. agarivorans	16	22	0	0	0
II	5	0	P. aurantia	2	3	5	0	0
III	0	9	P. prydzensis	8	1	0	0	0
IV	3	3	P. phenolica	3	3	3	2	0
V	4	0	P. luteoviolacea	4	0	4	0	4
VI	9	0	P. rubra	9	0	9	0	0
VII	13	0	P. flavipulchra	13	0	13	0	0
VIII	15	0	P. ruthenica	14	1	15	0	15
S1727	0	1	P. mariniglutinosa	0	1	0	0	0
S3655	1	0	P. spongiae	1	0	0	0	0
Total	51	50		70	31	49	2	19

Table 1 Identity, antibacterial activity, and pigmentation of Pseudoalteromonas strains from a global collection

^a The type strain which the majority of the strains in the cluster were most closely related to

WDAA. These 20 strains were identified as Pseudoalteromonas phenolica (one strain), Pseudoalteromonas luteoviolacea (five strains), Pseudoalteromonas rubra (nine strains), Pseudoalteromonas citrea (one strain), and Pseudoalteromonas aurantia (four strains). Ethyl acetate extraction of culture broths resulted in 19 crude extracts which inhibited growth of V. anguillarum in the WDAA. Four of these were identified as P. luteoviolacea and were the only strains where both cellfree supernatant and crude ethyl acetate extracts inhibited Vibrio growth. The remaining 15 inhibitory crude extracts all originated from strains identified as Pseudoalteromonas ruthenica. The crude ethyl acetate extracts of 26 strains were intensely colored; however, only some of these extracts inhibited growth of Vibrio indicating that the pigments were not universally antibacterial. Cell-free culture supernatants and ethyl acetate crude extracts of strains with no growth inhibition of V. anguillarum in the "spot test" assay were also tested but showed no growth inhibition.

Forty-eight of the 51 antibacterial strains were pigmented, while two (S3431 and S3655) of the 50 non-active strains were pigmented (Table 1). Antibacterial activity was significantly more likely to be produced by pigmented strains as determined by Fisher's exact test (two-tailed pvalue of 0.0000). In total, 70 strains were isolated from surface swabs and 31 from water samples. Of the surfaceassociated strains, 45 remained active in the spot-assay in comparison to six of the water sample strains. The Fisher's exact test demonstrated a significant relation between surface association and stable antibacterial activity (two-tailed p value of 0.0000).

Pseudoalteromonas strains were isolated on all parts of the global cruise in both tropical and temperate waters (Fig. 1). Our strain collection is not large enough to allow for complete biogeographic analysis, but pigmented strains appeared to be more frequent in coastal areas whereas the non-pigmented strains appeared associated with open waters (Fig. 1).

16S rRNA Gene Sequence Analyses We initially performed a BLAST search querying the 16S rRNA gene sequence of each strain against the GenBank database. The results of this analysis were ambiguous, as some sequences returned more than 40 hits all with identical scores in the BLAST results (data not shown), frequently including the sequences of several different *Pseudoalteromonas* species. Therefore, a BLAST analysis was carried out querying the sequences against a complete set of *Pseudoalteromonas* type strains, and hence, each strain is matched with the best type strain BLAST match (Suppl Table 1).

The 16S rRNA gene sequences were used to cluster the strains by constructing a neighbor-joining tree, and branch support was verified by comparison to minimum evolution and maximum likelihood trees (Fig. 2). Nodes supported by an 80% bootstrap cutoff were collapsed when three or more strains were included in the cluster (Fig. 2). An exception was made for clusters VI and VII, which are shown as separate clusters due to obvious differences in phenotype (pigment, bioactivity, secondary metabolite profile). Ninety-nine of the strains fell into one of eight primary clusters.

Clusters I and III consisted of non-pigmented non-inhibitory strains (Table 1). Cluster I included 38 strains and the type strains of *P. haloplanktis, Pseudoalteromonas agarivorans, Pseudoalteromonas tetraodonis, Pseudoalteromonas paragorgicola, Pseudoalteromonas distincta, Pseudoalteromonas arctica, Pseudoalteromonas nigrifaciens, Pseudoalteromonas* Fig. 1 Isolation sites of 101 *Pseudoalteromonas* strains with each color code representing one of the eight clusters as indicated in Table 1. Each *circle* represents the isolation of one or more strains belonging to the given cluster. *White*=cluster I, *yellow*=cluster II, *gray*=cluster III, *green*= cluster IV, *violet*=cluster V, *red*=cluster VI, *orange*=cluster VII, *pink*=cluster VII, and *blue*=non-clustered strain



elyakovii, Pseudoalteromonas carrageenovora, Pseudoalteromonas marina, and Pseudoalteromonas aliena. Strain S3431—a black-pigmented strain in cluster I—did not show more than 97% similarity even when compared to the full GenBank database which suggests that S3431 could represent a novel *Pseudoalteromonas* species. Despite the low BLAST similarity score, phylogenetic analysis and tree construction placed strain S3431 in the diverse cluster I (the non-collapsed cluster I is shown in Supplementary Figure 1). Cluster III contained no type strains which supported the BLAST analysis where the strains in cluster III were 98% similar to the best type strain match (Supplementary Table 1).

The remaining six of the eight clusters contained pigmented strains. Pale yellow strains clustered with the

Fig. 2 Phylogenetic tree based on 16S rRNA gene sequences of the Galathea 3 Pseudoalteromonas strains. Sequences were aligned by MAFFT (default options), and the resulting alignment was used to generate a neighbor-joining tree in the MEGA4 software package (default settings, 1,000 bootstraps). S. arenicola CNS-205, GenBank accession number CP000850, GeneID: 5705939 was used as outgroup. Clusters containing two or more non-type strain sequences were collapsed. The scale bar represents 0.02 amino acid substitutions per nucleotide position. Circle nodes also occurred in minimum evolution and maximum likelihood trees



type strains of *P. citrea* and *P. aurantia* (cluster II) and four intensely purple strains grouped in cluster V with the type strain of P. luteoviolacea. Cluster VI contained nine redpigmented strains and the type strain of *P. rubra*, and cluster VII consisted of 12 intensely yellow strains, one pale yellow strain, and the Pseudoalteromonas flavipulchra, Pseudoalteromonas maricaloris, and Pseudoalteromonas piscicida type strains. Fifteen strains and their nearest BLAST match, P. ruthenica, formed cluster VII. These strains all produced a pale brown pigment. Cluster IV contained six strains and the type strain of P. phenolica. Four of the strains in this cluster had *P. phenolica* as their best type strain BLAST match; however, strain S1093 had P. luteoviolacea as its best match at 98% identity, while P. rubra and P. luteoviolacea type strains scored identically (97%) as the best matches for S2724. The strains in cluster IV were heterogeneous with respect to pigmentation, some were non-pigmented and others appeared brown.

Profiling of Secondary Metabolites The 101 strains and select type strains were separable in discrete groups by HPLC-UV/VIS (Fig. 3), and the triplicate profiles from an isolate of each species were very reproducible and could be superimposed on each other (data not shown). All of the 38 strains of the 16S rRNA cluster I fell into group A, in which no UV/VIS peaks were unique compared to the media blanks indicating that no secondary metabolites were produced. This large group also included all nine strains from 16S rRNA cluster III and strains of *P. phenolica* and *P. ruthenica* less proficient in secondary metabolite production. A summary of the detected compounds is shown in Table 2, and the producer organisms are shown by 16S rRNA cluster in Table 3.

Based on their production of specific metabolites, the majority of pigmented bacteria formed six main groups not including four *P. rubra* strains (Fig. 3). In the pigmented bacteria, a total of 26 distinct peaks were detected and included in the cluster analysis. We identified indolmycin, violacein, and prodigiosin among the significant peaks based on reference standards. Furthermore, nine peaks could be tentatively identified based on HPLC-UV/VIS-TOFMS results and data in Antibase2009. These nine included two likely novel bromoalterochromides and a brominated indole (Table 2).

Comparing the 16S rRNA gene sequence clusters with the chemical profiling revealed several patterns. Some compounds were exclusively produced by strains belonging to the same cluster whereas other compounds were produced across several strains from different clusters. All strains in cluster II (*P. aurantia*/*P. citrea*) shared production of compound B (retention time, RT 12.31 min) whereas the production of four other compounds F, Q, T, and U (RT 15.47, 16.60, 17.96, and 18.28 min) were scattered in the group. This included a novel bromoalterochromide (compound Q, RT 16.60 min) that was also found in cluster VII (*P. flavipulchra/P. piscicida*). Three of the other compounds were identified as quinolines based on distinct UV spectra and accurate mass.

P. luteoviolacea strains (cluster V) shared production of compound D (violacein, RT 14.29) in all four strains and the type strain but were sub-divided by compound A (indolmycin, RT 11.21) produced by two strains and compound Z (pentabromopseudilin, RT 22.65) produced by the two other strains and the type strain. This division is visible in Fig. 4, which shows chromatograms of the four strains in cluster V. Interestingly, also two *P. phenolica* strains produced compound Z.

Cluster VII (*P. flavipulchra* and *P. piscicida*) was chemically very homogeneous. All strains except one produced three bromoalterochromides P, Q, and R (RT 16.49, 16.60, and 17.10) of which Q and R were novel compounds. In contrast to cluster V and compound D, the production of P, Q, and R was not a unique marker for strains of this cluster as P, Q, and R were also detected in one strain from cluster II and one strain from cluster VI.

Thirteen of the 15 strains in cluster VIII, identified as *P. ruthenica*, shared a unique chemical profile and produced the compounds H, K, and O (RT 15.78, 15.98, 16.30) with characteristic UV spectra. None of these matched compounds in Antibase2009 and potentially constitute novel antibacterials. These compounds were not detected among strains from other clusters, yet they were not suitable as a distinct chemical marker for cluster VIII since no secondary metabolites were detected in the remaining two strains in this cluster or the type strain.

The strains in cluster VI were identified as *P. rubra* and were chemically very heterogeneous. Five of nine strains produced the red pigment prodigiosin (compound M, RT 16.00) which was not detected in strains of other clusters. Additionally, a multitude of known and non-identified compounds were detected in one or more strains in the cluster. In total, 16 compounds were detected within the cluster, and 12 of these were unique for this cluster. Only two strains shared an identical production of secondary metabolites, further stressing the chemical diversity among the strains in this cluster.

Discussion

We demonstrate in this study, in agreement with earlier findings (Bowman 2007), that species within the *Pseudoalteromonas* genus produce a range of secondary metabolites, some with antibacterial activity. Several species of the genus are intensely pigmented, and it is



Fig. 3 Dendrogram from cluster analysis of detected peaks from HPLC-UV/VIS detection of compounds in the ethyl acetate extracted broth and biomass. Data were processed in NTSYSpc 2.20q, with SAHN clustering by UPGMA and simple distance measurement

hypothesized that pigmentation co-occur with antibacterial activity (Egan et al. 2002). In our global collection of *Pseudoalteromonas* strains that demonstrated antibacterial activity on initial isolation, strains that were pigmented were significantly more likely to retain antibacterial activity on re-growth than non-pigmented strains. Several intensely colored organic extracts were not inhibitory against *V. anguillarum*, and hence, we do not believe that the pigments, in general, are the cause of the antibacterial activity although, e.g., the purple pigment violacein is a known antibiotic compound (Lichstein and Vandesand 1945).

Table 2 Ethyl acetate extractable secondary metabolites produced by pigmented Pseudoalteromonas strains

Compound	MI (Da) ^c	UV-max data	RT (min)	Code
Indolmycin ^a	257		11.21	А
2-Pentyl-4-quinolinol ^b	215		12.31	В
Novel mono-brominated indole	280		13.78	С
Violacein ^a	343		14.29	D
Unidentified	244	212 nm (100%), 250 nm (48%)	15.21	Е
2-n-Heptyl-(1H)-quinolin-4-one ^b	243		15.47	F
Unidentified	NI	228 nm (45%), 308 nm (100%)	15.70	G
Unidentified	386	<200 nm	15.78	Н
Unidentified	316	286 nm (100%)	15.80	Ι
Unidentified	NI	<200 nm	15.81	J
Unidentified	NI	228 nm (45%), 308 nm (100%)	15.98	Κ
Unidentified	676	<200 nm	15.99	L
Prodigiosin ^a	323		16.00	М
Unidentified	NI	<200 nm	16.26	Ν
Unidentified	NI	228 nm (45%), 308 nm (100%)	16.30	0
Bromoalterochromide A ^b	843		16.49	Р
Novel bromoalterochromide, 2 bromine	921		16.60	Q
Novel bromoalterochromide, 1 bromine	857		17.10	R
Unidentified	333	310 nm (100%)	17.20	S
2-n-Nonyl-(1H)-quinolin-4-one ^b	271		17.96	Т
Nonyl-quinolinone analog ^b	271		18.28	U
Unidentified	315	362 nm (100%)	18.90	V
Unidentified	244	218 nm (100%), 280 nm (82%)	19.42	W
Unidentified	NI	218 nm (100%), 288 nm (82%)	19.78	Х
Unidentified	NI	250 nm (100), 280 nm (86)	19.92	Y
Pentabromopseudilin ^b	549		22.65	Ζ

NI no ionization or MI could not be assigned using ESI⁺ and ESI⁻

^a Validated reference standard used for verification

^b Accurate mass and UV data fit the data from Antibase2009

^c Mono-isotopic mass

Nearly half of the isolated strains did not retain any antibacterial effect after frozen storage and sub-culturing despite being isolated on the original plates due to antibacterial activity. The observed loss of antibacterial activity may be due to a requirement for factors specific to local seawater, as initial tests for antibacterial activity were carried out using 50% local seawater (Gram et al. 2010). Furthermore, loss of antibacterial activity may be due to repression or inhibition of gene clusters encoding products that are required for secondary metabolite synthesis (e.g., by catabolite repression). A reduction in antibiotic production when the producer organism is grown in excess of a carbon source is a known phenomenon (Sanchez et al. 2010), and suppression of secondary metabolite production by excess concentrations of other substrate components is demonstrated in Streptomyces (Doull and Vining 1990). This could suggest that culturing the strains under nutrient limited conditions may reestablish production of antibacterial compounds. Also, during the original sampling and screening procedure, the agar plates may have harbored co-cultured microorganisms which potentially induce antibacterial activity as has been demonstrated by Mearns-Spragg et al. (1998). Hence, it may be possible to re-induce the antibacterial activity if the right conditions can be created.

Several bioactive *Pseudoalteromonas* have been isolated from higher organisms, and it has been hypothesized that antibacterial compounds may play a role in bacterial competition or as protective agents beneficial for the host organism (Holmström and Kjelleberg 1999). We provide statistical evidence that surface-associated presumed antibacterial pseudoalteromonads are significantly more likely to show stable production of antibacterial compounds than







×

×

×

×

×

16S cluster

×

×

LMG19699

P. ruthenica

S1727 S3655

×

Identification of peak by capital letter in Table 2

×

N



Fig. 4 HPLC-UV/VIS profiles of the ethyl acetate extracted broth and biomass of four *P. luteoviolacea* with the tailing violacein peak at 14–15 min and in two traces pentabromopseudilin at 22-23 min in contrast to indolmycin at 11 min in the two other traces

Pseudoalteromonas species isolated as planktonic cells. This suggests that production of antibacterial compounds may play an important role in the ability of *Pseudoalteromonas* strains to colonize and persist on surfaces submerged in the marine environment, as previously suggested for *P. tunicata* strain D2 (Rao et al. 2005).

The analysis of 16S rRNA gene sequences from our global collection of Pseudoalteromonas confirms that phylogenetic analysis results in a number of clusters encompassing predominantly pigmented species or non-pigmented species (Ivanova et al. 2004). Strain S3431 was the single pigmented strain in the so-called non-pigmented clusters. Novel diversity might be represented in cluster III which consisted of strains with 98% or less 16S rRNA gene sequence similarity to Pseudoalteromonas type strains and formed a separate cluster in the phylogenetic analysis. However, these strains showed no antibacterial activity and no small molecule metabolites were detected. Such novel diversity could still represent untapped biotechnological potential, producing, e.g., enzymes or peptides with biological activity, as known for other non-pigmented Pseudoalteromonas (Hoyoux et al. 2001; Violot et al. 2005).

Chemical profiling of the strains detected an array of secondary metabolites. In addition to complementing our analysis of 16S rRNA gene sequences, it also demonstrated that some compounds (e.g., violacein, prodigiosin) were characteristic of a species and other compounds were produced by several species, and we also detected intraspecies clusters of different secondary metabolite profiles. In a broad sense, the clustering based on 16S rRNA gene similarity agreed with the groups resulting from the chemophylogenetic analysis. However, some compounds were produced by organisms of different species that then clustered together using the secondary metabolites as basis. The chemical analysis separated the four isolated *P. luteoviolacea* strains into two distinct sub-groups, showing intra-species chemical diversity. The *P. luteoviolacea* strains produced violacein and

pentabromopseudilin which are active against gram-positive and gram-negative bacteria and the anti-staphylococcal agent indolmycin (Hornemann et al. 1971; Hurdle et al. 2004). Violacein and pentabromopseudilin have previously been detected in *P. luteoviolacea* (Gauthier 1976a; Laatsch and Pudleiner 1989), but to our knowledge, this is the first report of *Pseudoalteromonas* strains producing indolmycin (Månsson et al. 2010).

Within some species, all strains were consistently antibacterial. However, in others, such activity did not appear to be a consistent trait of the species. For instance, strains of the 16S cluster VI (P. phenolica) were heterogeneous in their ability to inhibit Vibrio in our assays, while all but one strain in the homogeneous cluster VII had identical metabolite profiles and all were inhibitory. Even more obvious was the heterogeneous chemical profiles within the P. rubra strains. All except one strain shared a chemical marker prodigiosin and/or RT 15.99 min but had major variations in 14 other compounds. This may in part be due to loss of ability to produce a compound. For instance, strain S2471 over time lost ability to produce the brominated indole (RT 13.78 min). Also, we note that the type strain DSM 6842 (ATCC 29570) did not in our culture produce prodigiosin which has been observed previously (Gauthier 1976b; Gauthier and Flatau 1976). The consistent bromoalterochromide production in the two species P. piscicida and P. flavipulchra/maricaloris (cluster VII) was expected (Speitling et al. 2007) and supported the high DNA sequence similarity between the two. This emphasizes the need to isolate and screen multiple strains from each species when bioprospecting within the genus Pseudoalteromonas, as even the homogeneous cluster VII contains one strain with a metabolite profile that does not share a single compound with the other strains in this cluster.

Several of the 26 detected peaks were known substances, with a majority known as antibacterials. These included violacein (Lichstein and Vandesand 1945), two bromopseudilins (Lovell 1966), two indolmycins (Werner 1980), four quinolines (Wratten et al. 1977), and prodigiosin (Kalesperis et al. 1975). Due to its very low aqueous solubility, violacein probably protects against predation rather than acts as a true antibiotic, and it has been shown to induce cell death in grazing organisms (Matz et al. 2008). Such compounds would be very beneficial for protection of a biofilm, which is likely how surface-associated Pseudoalteromonas would grow. The 14 compounds that could not be identified were mainly not identified due to poor ionization in ESI⁺ and ESI⁻ and/or several plausible candidates in Antibase2009. However, for chemotaxonomic studies, identity of the compounds is not necessary as long as they can be unambiguously identified between samples (Frisvad et al. 2008).

Within cluster VIII (*P. ruthenica*) and cluster II (*aurantia*/ *citrea*), we found examples where strains with highly similar 16S rRNA gene sequences (>99%) and with identical chemotaxonomy originated from geographically distinct locations. This latter observation is in agreement with studies on *Salinispora* biogeography and secondary metabolite production in which the authors show how strains of the marine bacterium *S. arenicola* isolated from worldwide locations are highly related and produce identical patterns of secondary metabolites (Jensen and Mafnas 2006; Jensen et al. 2007). In contrast, the *P. luteoviolacea* and *P. rubra* strains showed both local and global variations in their secondary metabolite profile, which one might speculate is due to adaptation to local specific niches.

In conclusion, we believe sampling from specific niches, e.g., biofilms on surfaces, to be of importance in discovery of novel secondary metabolites from the genus *Pseudoalteromonas*. While differences in metabolite patterns among species encourage isolation and screening of novel diversity, bioprospecting known *Pseudoalteromonas* species should not be ruled out. Investigation of multiple strains of one *Pseudoalteromonas* species can yield novel compounds due to intra-species variations within secondary metabolite profiles.

Acknowledgments We acknowledge Dr. Jesper B. Bruhn for valuable input during the early phase of this study. This study was supported by the Programme Commission on Health, Food and Welfare under the Danish Council for Strategic Research. The present work was carried out as part of the Galathea 3 expedition under the auspices of the Danish Expedition Foundation. This is Galathea 3 contribution no. p73.

References

- Armstrong E, Yan LM, Boyd KG, Wright PC, Burgess JG (2001) The symbiotic role of marine microbes on living surfaces. Hydrobiologia 461:37–40
- Bewley CA, Faulkner DJ (1998) Lithistid sponges: star performers or hosts to the stars. Angew Chem Int Ed 37:2162–2178
- Bowman JP (2007) Bioactive compound synthetic capacity and ecological significance of marine bacterial genus *Pseudoalteromonas*. Mar Drugs 5:220–241
- Boyd KG, Adams DR, Burgess JG (1999) Antibacterial and repellent activities of marine bacteria associated with algal surfaces. Biofouling 14:227–236
- Burgess JG, Jordan EM, Bregu M, Mearns-Spragg A, Boyd KG (1999) Microbial antagonism: a neglected avenue of natural products research. J Biotechnol 70:27–32
- Castresana J (2000) Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. Mol Biol Evol 17:540–552
- Doull JL, Vining LC (1990) Nutritional control of actinorhodin production by *Streptomyces coelicolor* A3(2)—suppressive effects of nitrogen and phosphate. Appl Microbiol Biotechnol 32:449–454
- Egan S, Holmström C, Kjelleberg S (2001) *Pseudoalteromonas ulvae* sp nov., a bacterium with antifouling activities isolated from the surface of a marine alga. Int J Syst Evol Microbiol 51:1499–1504

- Egan S, James S, Holmström C, Kjelleberg S (2002) Correlation between pigmentation and antifouling compounds produced by *Pseudoalteromonas tunicata*. Environ Microbiol 4:433–442
- Egan S, Thomas T, Kjelleberg S (2008) Unlocking the diversity and biotechnological potential of marine surface associated microbial communities. Curr Opin Microbiol 11:219–225
- Fisher RA (1958) Statistical methods for research workers. Hafner, New York
- Fox GE, Wisotzkey JD, Jurtshuk P (1992) How close is close: 16S rRNA sequence identity may not be sufficient to guarantee species identity. Int J Syst Evol Microbiol 42:166
- Franks A, Egan S, Holmstrom C, James S, Lappin-Scott H, Kjelleberg S (2006) Inhibition of fungal colonization by *Pseudoalteromonas tunicata* provides a competitive advantage during surface colonization. Appl Environ Microbiol 72:6079–6087
- Frisvad JC, Andersen B, Thrane U (2008) The use of secondary metabolite profiling in chemotaxonomy of filamentous fungi. Mycol Res 112:231–240
- Gauthier MJ (1976a) Morphological, physiological, and biochemical characteristics of some violet-pigmented bacteria isolated from seawater. Can J Microbiol 22:138–149
- Gauthier MJ (1976b) Alteromonas rubra sp. nov., a new marine antibiotic-producing bacterium. Int J Syst Bacteriol 26:459–466
- Gauthier MJ, Flatau GN (1976) Antibacterial activity of marine violetpigmented *Alteromonas* with special reference to the production of brominated compounds. Can J Microbiol 22:1612–1619
- Geiser DM, Klich MA, Frisvad JC, Peterson SW, Varga J, Samson RA (2007) The current status of species recognition and identification in *Aspergillus*. Stud Mycol 59:1
- Gram L, Melchiorsen J, Bruhn JB (2010) Antibacterial activity of marine culturable bacteria collected from a global sampling of ocean surface waters and surface swabs of marine organisms. Mar Biotechnol 12:439–451
- Guindon S, Gascuel O (2003) A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. Syst Biol 52:696
- Hayashida-Soiza G, Uchida A, Mori N, Kuwahara Y, Ishida Y (2008) Purification and characterization of antibacterial substances produced by a marine bacterium *Pseudoalteromonas haloplanktis* strain. J Appl Microbiol 105:1672–1677
- Hjelm M, Bergh Ø, Riaza A, Nielsen J, Melchiorsen J, Jensen S, Duncan H, Ahrens P, Birkbeck H, Gram L (2004) Selection and identification of autochthonous potential probiotic bacteria from turbot larvae (*Scophthalmus maximus*) rearing units. Syst Appl Microbiol 27:360–371
- Holmström C, Kjelleberg S (1999) Marine *Pseudoalteromonas* species are associated with higher organisms and produce biologically active extracellular agents. FEMS Microbiol Ecol 30:285–293
- Holmström C, James S, Egan S, Kjelleberg S (1996) Inhibition of common fouling organisms by marine bacterial isolates with special reference to the role of pigmented bacteria. Biofouling 10:251–259
- Holmström C, Egan S, Franks A, McCloy S, Kjelleberg S (2002) Antifouling activities expressed by marine surface associated *Pseudoalteromonas* species. FEMS Microbiol Ecol 41:47–58
- Hornemann U, Hurley LH, Speedie MK, Floss HG (1971) Biosynthesis of indolmycin. J Am Chem Soc 93:3028–3035
- Hoyoux A, Jennes I, Dubois P, Genicot S, Dubail F, Francois JM, Baise E, Feller G, Gerday C (2001) Cold-adapted betagalactosidase from the Antarctic psychrophile *Pseudoalteromo*nas haloplanktis. Appl Environ Microbiol 67:1529–1535
- Hurdle JG, O'Neill AJ, Chopra I (2004) Anti-staphylococcal activity of indolmycin, a potential topical agent for control of staphylococcal infections. J Antimicrob Chemothe 54:549–552
- Isnansetyo A, Kamei Y (2003) MC21-A, a bactericidal antibiotic produced by a new marine bacterium, *Pseudoalteromonas phenolica* sp. nov. O-BC30T, against methicillin-resistant *Staphylococcus aureus*. Antimicrob Agents Chemother 47:480–488

- Ivanova EP, Flavier S, Christen R (2004) Phylogenetic relationships among marine *Alteromonas*-like proteobacteria: emended description of the family *Alteromonadaceae* and proposal of *Pseudoalteromonadaceae* fam. nov., *Colwelliaceae* fam. nov., *Shewanellaceae* fam. nov., *Moritellaceae* fam. nov., *Ferrimonadaceae* fam. nov., *Idiomarinaceae* fam. nov. and *Psychromonadaceae* fam. nov. Int J Syst Evol Microbiol 54:1773–1788
- James SG, Holmström C, Kjelleberg S (1996) Purification and characterization of a novel antibacterial protein from the marine bacterium D2. Appl Environ Microbiol 62:2783–2788
- Jensen PR, Mafnas C (2006) Biogeography of the marine actinomycete Salinispora. Environ Microbiol 8:1881–1888
- Jensen PR, Williams PG, Oh DC, Zeigler L, Fenical W (2007) Species-specific secondary metabolite production in marine actinomycetes of the genus *Salinispora*. Appl Environ Microbiol 73:1146–1152
- Jiang Z, Boyd KG, Mearns-Spragg A, Adams DR, Wright PC, Burgess JG (2000) Two diketopiperazines and one halogenated phenol from cultures of the marine bacterium, *Pseudoalteromonas luteoviolacea*. Nat Prod Let 14:435–440
- Kalesperis GS, Prahlad KV, Lynch DL (1975) Toxigenic studies with the antibiotic pigments from *Serratia marcescens*. Can J Microbiol 21:213
- Kalinovskaya NI, Ivanova EP, Alexeeva YV, Gorshkova NM, Kuznetsova TA, Dmitrenok AS, Nicolau DV (2004) Lowmolecular-weight, biologically active compounds from marine *Pseudoalteromonas* species. Curr Microbiol 48:441–446
- Katoh K, Misawa K, Kuma K, Miyata T (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. Nucleic Acids Res 30:3059–3066
- Laatsch H, Pudleiner H (1989) Marine bakterien, I: synthese von pentabrompseudilin, einem phenylpyrrol aus Alteromonas luteoviolaceus. Liebigs Ann Chem 1989:863–881
- Lichstein HC, Vandesand VF (1945) Violacein, an antibiotic pigment produced by *Chromobacterium violaceum*. J Infect Dis 76:47–51
- Lovell FM (1966) Structure of a bromine-rich marine antibiotic. J Am Chem Soc 88:4510–4511
- Månsson M, Phipps RK, Gram L, Munro MH, Larsen TO, Nielsen KF (2010) Explorative solid-phase extraction (E-SPE) for accelerated microbial natural product discovery. J Nat Prod 73:1126–1132
- Matz C, Webb JS, Schupp PJ, Phang SY, Penesyan A, Egan S, Steinberg P, Kjelleberg S (2008) Marine biofilm bacteria evade eukaryotic predation by targeted chemical defense. PLoS ONE 3: e2744
- McCarthy SA, Johnson RM, Kakimoto D (1994) Characterization of an antibiotic produced by *Alteromonas luteoviolacea* Gauthier 1982, 85 isolated from Kinko Bay, Japan. J Appl Microbiol 77:426–432
- Mearns-Spragg A, Bregu M, Boyd KG, Burgess JG (1998) Crossspecies induction and enhancement of antimicrobial activity produced by epibiotic bacteria from marine algae and invertebrates, after exposure to terrestrial bacteria. Lett Appl Microbiol 27:142–146
- Newman DJ (2008) Natural products as leads to potential drugs: an old process or the new hope for drug discovery? J Med Chem 51:2589–2599
- Nielsen KF, Smedsgaard J (2003) Fungal metabolite screening: database of 474 mycotoxins and fungal metabolites for dereplication by standardised liquid chromatography-UV-mass spectrometry methodology. J Chromatogr A 1002:111–136

- Nielsen KF, Sumarah MW, Frisvad JC, Miller JD (2006) Production of metabolites from the *Penicillium roqueforti* complex. J Agric Food Chem 54:3756–3763
- Penesyan A, Marshall-Jones Z, Holmstrom C, Kjelleberg S, Egan S (2009) Antimicrobial activity observed among cultured marine epiphytic bacteria reflects their potential as a source of new drugs. FEMS Microbiol Ecol 69:113–124
- Rao D, Webb JS, Kjelleberg S (2005) Competitive interactions in mixed-species biofilms containing the marine bacterium *Pseudoalteromonas tunicata*. Appl Environ Microbiol 71:1729–1736
- Sanchez S, Chavez A, Forero A, Garcia-Huante Y, Romero A, Sanchez M, Rocha D, Sanchez B, Avalos M, Guzman-Trampe S, Rodriguez-Sanoja R, Langley E, Ruiz B (2010) Carbon source regulation of antibiotic production. J Antibiot 63:442–459
- Simmons TL, Coates RC, Clark BR, Engene N, Gonzalez D, Esquenazi E, Dorrestein PC, Gerwick WH (2008) Biosynthetic origin of natural products isolated from marine microorganisminvertebrate assemblages. P Natl A Sci USA 105:4587–4594
- Skov MN, Pedersen K, Larsen JL (1995) Comparison of pulsed-field gel electrophoresis, ribotyping, and plasmid profiling for typing of *Vibrio anguillarum* serovar O1. Appl Environ Microbiol 61:1540–1545
- Speitling M, Smetanina OE, Kuznetsova TA, Laatsch H (2007) Marine bacteria. XXXV. Bromoalterochromides A and A', unprecedented chromopeptides from a marine *Pseudoalteromonas maricaloris* strain KMM 636. J Antibiot 60:36–42
- Stackebrandt E, Frederiksen W, Garrity GM, Grimont PAD, Kampfer P, Maiden MCJ, Nesme X, Rossello-Mora R, Swings J, Truper HG, Vauterin L, Ward AC, Whitman WB (2002) Report of the ad hoc committee for the re-evaluation of the species definition in bacteriology. Int J Syst Evol Microbiol 52:1043–1047
- Sudek S, Lopanik NB, Waggoner LE, Hildebrand M, Anderson C, Liu H, Patel A, Sherman DH, Haygood MG (2006) Identification of the putative bryostatin polyketide synthase gene cluster from "Candidatus Endobugula sertula", the uncultivated microbial symbiont of the marine bryozoan Bugula neritina. J Nat Prod 70:67–74
- Talavera G, Castresana J (2007) Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. Syst Biol 56:564–577
- Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. Mol Biol Evol 24:1596
- Violot S, Aghajari N, Czjzek M, Feller G, Sonan GK, Gouet P, Gerday C, Haser R, Receveur-Brechot V (2005) Structure of a full length psychrophilic cellulase from *Pseudoalteromonas haloplanktis* revealed by X-ray diffraction and small angle X-ray scattering. J Mol Biol 348:1211–1224
- Werner RG (1980) Uptake of indolmycin in gram-positive bacteria. Antimicrob Agents Chemother 18:858
- Wietz M, Schramm A, Jørgensen B, Gram L (2010) Latitudinal patterns in the abundance of major marine bacterioplankton groups. Aquat Microb Ecol 61:179–189
- Wratten SJ, Wolfe MS, Andersen RJ, Faulkner DJ (1977) Antibiotic metabolites from a marine pseudomonad. Antimicrob Agents Chemother 11:411–414
- Zheng L, Yan XJ, Han XT, Chen HM, Lin W, Lee FSC, Wang XR (2006) Identification of norharman as the cytotoxic compound produced by the sponge (*Hymeniacidon perleve*)-associated marine bacterium *Pseudoalteromonas piscicida* and its apoptotic effect on cancer cells. Biotechnol Appl Biochem 44:135–142

Vynne N, Månsson M, Nielsen KF, Gram L (2012)

Antibiotic chemoprofile and RecA and GyrB sequence analyses reveal three phylotypes within the marine *Pseudoalteromonas luteoviolacea*.

Applied and Environmental Microbiology: Submitted.

1	Antibiotic chemoprofile and RecA and GyrB sequence analyses reveal three
2	phylotypes within the marine Pseudoalteromonas luteoviolacea
3	
4	Nikolaj G. Vynne ^{1*} , Maria Månsson ² , Kristian F. Nielsen ² and Lone Gram ¹
5	
6	¹ National Food Institute, Technical University of Denmark, Søltofts Plads 221, DK-2800 Kgs. Lyngby,
7	Denmark
8	² Department of Systems Biology, Technical University of Denmark, Søltofts Plads 221, DK-2800 Kgs. Lyngby,
9	Denmark
10	* corresponding author
11	ngvy@food.dtu.dk
12	+45 4525 2571
13	
14	
15	Running title: Chemoprofiles and phylotypes in <i>P. luteoviolacea</i>

1 Abstract

2 The bacterial species Pseudoalteromonas luteoviolacea produces antibiotic compounds and 3 Pseudoalteromonads are, in general, known for their bioactivity. This has made them an interesting topic of 4 research in the area of drug discovery. A major time sink in natural products discovery is the effort spent 5 rediscovering known compounds, and methods to avoid this are in high demand. We recently found that 6 four strains of P. Iuteoviolaceae indistinguishable by 16S rRNA gene sequence produced two different 7 antibiotic profiles. The purpose of the present study was to determine if this sub-division was true for a 8 larger group of *P. luteoviolaceae* isolates and to determine if it could be related to their phylogeny. These 9 analyses would shed light on inter-species evolution and serve to determine if simple phylogenetic tools 10 could assist in further discovery of bioactives in species already known to produce bioactive secondary metabolites. The 13 P. luteoviolacea strains divided into three chemoprofiles based on antibiotic 11 12 production (violacein, violacein + pentabromopseudilin, violacein + indolmycin). Detailed phylogenetic 13 analysis of GyrB and RecA translated amino acid sequences divided the strains into phylotypes related to antibacterial activity and production of the antibiotic compounds. Hence, we believe phylogeny within a 14 15 bioactive species has the potential to be a useful tool in biodiscovery efforts. Additionally, the correlation 16 between evolutionary groups and antibiotic production provides evidence that the antibiotic biosynthetic 17 pathways are maintained under selective pressure and hence likely confer important benefits for the 18 producer organism.

19

20 237 words.

2

1 Introduction

2

3 Many antibiotics used in treatment of infectious disease are of natural product origin, and despite high 4 hopes for new drug discovery strategies, alternative approaches to drug discovery such as combinatorial 5 chemistry has failed to adequately supply the drug pipeline (48). Therefore, we must revert to discovery of 6 novel natural products capable of inhibiting or killing pathogenic bacteria (6). Screening of microorganisms 7 (47) from extreme or underexplored environments(1) could be a promising approach as novel genetic 8 diversity ensuring adaptation to extreme conditions may encode production of novel compounds. The 9 marine environment is still considered an underexplored source of novel antimicrobial compounds and 10 especially marine microorganisms are viewed as a potential source of novel antibiotic compounds 11 (15,23,40). Indeed, marine bacteria belonging to the Cyanobacteria (60), Actinobacteria (49), 12 Pseudoalteromonas (2,62), and the Roseobacter clade (4,28) produce compounds with interesting 13 pharmacological properties. 14 15 Members of the genus *Pseudoalteromonas* are ubiquitous in the marine environment (65) and form two 16 phylogenetic groups supported by analysis of 16S rRNA gene sequences (46,62). One group predominantly 17 consists of non-pigmented pelagic bacteria with low or no antibacterial activity while the majority of 18 species in the other group is pigmented and antagonistic bacteria typically isolated from marine biotic 19 surfaces. The widespread occurrence of these bacteria is an indication of successful survival and 20 competition strategies in which a trait such as production of antagonistic compounds could play an 21 important ecological role (26). Consequently, the Pseudoalteromonas represent a promising target for

22 biodiscovery efforts.

23

One of the main challenges in natural products discovery is the effort squandered rediscovering known
compounds (9), hence so called 'dereplication' strategies to reduce the degree of rediscovery prior to

3
purification and structure elucidation steps are of outmost importance (29,50,55). One such strategy is 1 2 early stage dereplication informed by microbial taxonomy (36,37,58). However, this is complicated by the 3 inherent challenges of taxonomical classification (16) and more so, by the observation that closely related 4 strains of the same species may produce different antibiotics (62,63) while strains of different genera may 5 produce the same antibiotics as observed for e.g. the bioactive pigment violacein(18,39,64). Such cross-6 species production of compounds is likely due to lateral gene transfer of gene clusters responsible for 7 biosynthesis of these metabolites (14,51). Nonetheless, screening of a taxonomically dereplicated 8 collection of Actinobacteria led to discovery of a high number of novel compounds as compared to the 9 strain throughput (20). Hence, it may possible to apply knowledge of bacterial systematics and taxonomy as 10 a guide for efficient bioprospecting within other groups of bacteria. We previously reported (41,62) that 11 five Pseudoalteromonas luteoviolacea strains produced two combinations of the three antibacterial 12 compounds violacein (39), indolmycin (42), and pentabromopseudilin (PBP) (5) and accordingly may serve 13 as a model to explore this hypothesis.

14

15 The aim of the present study was investigate phylogeny as a tool for dereplication in biodiscovery efforts 16 and provide an evolutionary perspective on bacterial antagonism within the species Pseudoalteromonas 17 luteoviolacea. This was achieved through detailed analysis of GyrB and RecA sequences coupled with 18 chemical detection of three known antibacterial compounds; violacein, indolmycin, and 19 pentabromopseudilin. We found that the chemoprofile was related to phylogenetic affiliation which 20 validates phylogeny as an approach for dereplication of bacterial strains in a natural products perspective 21 and offers evidence that the underlying biosynthetic pathways are maintained through the course of 22 evolution.

23

24 Materials and methods

Bacterial strains and culture conditions. Thirteen *P. luteoviolacea* strains were used in this study (table 1).
They originated from distinct geographical areas and were primarily isolated from surface water or algae.
The strains were cultured in a marine minimal medium (MMM) (52) with 4.0 g/l mannose and 3.0 g/l
casamino acids. All strains were incubated at 25 °C and 200 rpm agitation.

5

6 Assays for inhibition of bacterial growth. Vibrio anguillarum 90-11-287 (57) and Staphyloccoccus aureus 7 8325 were cultured in tryptic soy broth (Difco, USA). Well diffusion agar assays were carried out as 8 previously described (28). The assay substrate contained 30 g/l Sea Salts (Sigma, USA) and 10 g/l agar. To 9 support growth of V. anguillarum 4 g/l glucose and 3 g/l casamino acids was added. An additional 5 g/l 10 peptone was added to S. aureus agar. The inter-species inhibition was tested by embedding each of the 13 11 strains in agar and testing inhibition by sterile filtered culture of the same strains. The agar-substrate was 12 modified for the inhibition assays targeting Pseudoalteromonas luteoviolacea: Glucose was substituted with 13 4 g/l mannose, and due to temperature sensitivity of the P. luteovioacea, standard bacteriological agar was replaced with low gelling point agarose (Sigma, USA). The growth substrate for the assays was cooled to 37 14 15 ^oC in a water bath before overnight bacterial culture was added, and the assays were subsequently 16 performed as for V. anguillarum and S. aureus.

17

18 Analytical detection of antibacterial compounds. Samples for LC-MS analyses were prepared from cultures 19 in MMM extracted with ethyl acetate (EtOAc). Extracts were dried under nitrogen and redissolved in 20 methanol (MeOH). LC-MS samples were analysed using an Agilent 1100 HPLC system with a diode array 21 detector (Waldbronn, Germany) coupled to an LCT TOF mass spectrometer (Micromass, Manchester, UK) 22 using a Z-spray electrospray (ESI) source. A Phenomenex Luna II C₁₈ column (50 mm ×2 mm, 3 µm) was used for separation, applying a linear acetonitrile (MeCN)-water (20 mM formic acid) 0.3 ml min⁻¹ gradient (15-23 24 100%) over 20 min at 40°C. For all LC-MS analyses, violacein and indolmycin were detected in positive 25 ionisation mode (ESI⁺), while pentabromopseudilin was detected in negative mode (ESI⁻).

1

2	PCR amplification and sequencing. DNA was purified from overnight P. luteoviolacea cultures using the
3	NucleoSpin Tissue kit (Machery-Nagel, Germany) or QIAGEN Genomic-Tip G/100 (QIAGEN, USA) following
4	the manufacturer's protocol. 16S rRNA genes from strains H33, H33S, NCIMB 1942, NCIMB 1944 and
5	NCIMB 2035 were amplified by PCR, and gyrB and recA genes were amplified from all strains. One reaction
6	consisted of 2.5 μ l 10X hot start PCR buffer (Fermentas, Canada), 2.5 μ l 2mM dNTP mix, 4 μ l 25 mM MgCl ₂ ,
7	0.8 μl 12.5 μM forward primer, 0.8 μl 12.5 μM reverse primer, 12.28 μl MilliQ H ₂ O, 0.2 μl Maxima Hot Start
8	<i>Taq</i> DNA polymerase (Fermentas, Canada) and 1 μ l DNA template at 50 ng/ μ l for a total volume of 25 μ l.
9	The reactions were performed on an Applied Biosystems Veriti 96 well cycler. 16S rRNA genes were
10	amplified according to [Porsby]. The primers and reaction conditions used for gyrB amplification were as
11	described in (66). Primers and conditions for amplification of <i>recA</i> fragments were according to (53).
12	Sequencing was done by Eurofins MWG Operon, Germany.
13	
14	Phylogenetic analysis. The 16S rRNA gene sequences were obtained from GenBank or by sequencing and
15	aligned in MEGA5 using MUSCLE (59). The evolutionary history was inferred using the maximum likelihood

16 method based on the Hasegawa-Kishino-Yano model (25), determined by MEGA5 as the best fitting model.

The phylogenetic tree was constructed using MEGA5 default settings. For analysis of *recA* and *gyrB*sequences, an alignment was created for each gene in MEGA5 using MUSCLE. The alignments were curated

19 manually and trimmed to be of equal length and in-frame for translation to amino acid sequence. The two

20 alignments were concatenated and phylogenetic analyses were performed in MEGA5. Evolutionary

21 relationships based on the encoded amino acid sequence were inferred using the maximum likelihood

22 method with Dayhoff matrix-based distance calculations (56). A phylogenetic tree was generated and

tested with 1000 bootstrap replications.

1	Minimum inhibitory concentrations (MIC) of violacein and indolmycin. MICs were determined for
2	violacein (Sigma, USA) and indolmycin (Bioaustralis, Australia) as these compounds were commercially
3	available. MIC assays were carried out in 96-well microtiter plates according to the guidelines by the clinical
4	and laboratory standards institute (7), with minor modifications. P. luteoviolacea strains, V. anguillarum 90-
5	11-287, and S. aureus 8325 were tested for sensitivity to indolmycin by determining the MIC. P.
6	luteoviolacea was cultured in MMM with 4 g/l mannose and 3 g/l casamino acids, V. anguillarum in MMM
7	with 4 g/l glucose and 3 g/l casamino acids, and S. aureus in MHB. Overnight bacterial cultures were diluted
8	to 10^5 CFU/ml and 90 μl added per well. 10 μl of antibiotic solution was added to each well. Indolmycin was
9	tested in serial two-fold dilution at final concentrations of 0.5 μ g/ml to 128 μ g/ml. Controls were included
10	for no antibiotic and for ethanol solvent. The well containing the lowest concentration of antibiotic that
11	had no visual bacterial growth after 48 hours corresponded to the MIC.
12	
13	Genbank nucleotide accession numbers. 16S rRNA nucleotide sequences JQ250820-JQ250824, gyrB
14	nucleotide sequences JQ280430 – JQ280442 and <i>recA</i> nucleotide sequences JQ280417-JQ280429.
15	
16	Results
17	
18	P. luteoviolacea production of antibacterial compounds and antibacterial activity. A collection of P.
19	luteoviolacea strains were tested for the ability to inhibit bacterial growth using either live cells or sterile
20	filtered culture supernatant. The strains were also tested for their ability to produce three known
21	antibacterial compounds previously identified within the species (table 2). All 13 strains produced the
22	purple antibacterialpigment violacein. Four strains did not produce other known antibiotics, three strains
23	produced indolmycin and five strains produced PBP (table 2). No strain produced both indolmycin and PBP.
24	Despite the violacein production by all strains, just 11 of the 13 strains inhibited V. anguillarum and S.
25	aureus in the live cell assay. In the test of sterile filtered culture supernatants, only the three indolmycin

producing strains inhibited both V. anguillarum and S. aureus. Supernatants from nine strains inhibited S. 1 2 aureus, and these strains produced indolmycin or PBP. The supernatant from the remaining four strains 3 were not inhibitory against any of the target organisms and these were the four strains that produced 4 neither indolmycin nor PBP. Sterile filtered culture supernatants of the P. luteoviolacea strains were also 5 tested for intra-species inhibition (table S1). Five sterile filtered supernatants inhibited the full spectrum of 6 P. luteoviolacea strains, including the producer strain itself: Two of the indolmycin producing strains 7 (S4047-1 and S4054) and three of the PBP producers (S2607, S4060-1 and 2ta16). Four supernatants 8 inhibited three to five *P. luteoviolacea* strains. Among these, interestingly the supernatants of strains H33 9 and H33S, both unable to inhibit V. anguillarum or S. aureus, did inhibit three (H33) and five (H33S) strains 10 respectively. Overall, however, , the intra-species inhibition patterns did not correlate with the antibiotic 11 production profile of the strains.

12

13 Phylogenetic analyses and assignment of phylotypes. A 16S rRNA gene sequence and a GyrB-RecA based phylogenetic tree were created. The 16S rRNA sequences were similar at a level of >99% and the 14 15 phylogenetic reconstruction showed no correlation to production of antibiotics (figure 1). In contrast, three 16 phylotypes were identified in the GyrB-RecA phylogeny (figure 2; I, II, and III) and corroborated the division 17 of P. luteoviolacea strains into groups with specific antibiotic production and bacterial inhibition. Phylotype 18 I was synonymous with the six PBP producing strains. Within phylotype II, the indolmycin producer \$4054 19 placed itself as a representative of an ancestral state to the two other indolmycin producers and, on a 20 separate branch, the strains H33 and H33S. Phylotype III was synonymous with the two strains NCIMB 1944 21 and NCIMB 2035 which were not antagonistic.

22

MIC of known antagonistic compounds. To investigate the potential role of indolmycin and violacein as
 antibacterial compounds, the MICs against *V. anguillarum, S. aureus,* and the 13 *P. luteovioloacea* strains
 were determined. Under these experimental conditions violacein did not inhibit any of the tested strains at

the concentrations included in the experiment; hence the MIC of violacein for all tested strains was > 128 μ g/ml. The MIC of indolmycin to *V. anguillarum* was 2 μ g/ml. *S. aureus* was inhibited by indolmycin at all tested concentrations, resulting in a MIC of $\leq 0.5 \mu$ g/ml. None of the *P. luteoviolaceae*, irrespective of antibiotic profile, were inhibited by indolmycin and MIC was > 128 μ g/ml

5

6 Discussion

7 Pseudoalteromonas luteoviolacea is a well-known producer of several bioactive secondary metabolites 8 which are believed to play important ecological roles for the producer organism, such as protection against 9 predation or bacterial antagonism. However, little is known of the phylogenetic distribution of antibiotic 10 compounds within this species. We recently (Mansson et al. 2010, Vynne et al 2011) found that within a 11 collection of just five strains two distinct antibiotic profiles were detected, prompting us to speculate if this 12 reflected phylogenetic diversity or was a mere phenotypic variation. In the present study, we expand our 13 investigation to a collection of 13 strains demonstrating three different antibiotic profiles within this species. We have attempted to further broaden our collection; however, were not able to obtain more 14 15 strains as P. luteoviolacea under certain conditions is autoinhibitory (18) and several laboratories no longer 16 had stock cultures. Our data demonstrate that biodiscovery processes can benefit from exploring several 17 strains of the same species.

18

All strains produced violacein, a hallmark compound of the species; however, the data presented here and currently in preparation (10) indicates that the antibacterial activity of violacein is limited. Alternatively, violacein likely acts as a cell-associated anti-predation compound (44) which is more in line with its low solubility in water and its clear cell-association. Violacein is not present in a simple sterile filtered supernatant and these were only antibacterial if harvested from strains producing either indolmycin or PBP. The indolmycin or PBP containing supernatants were equally inhibitory towards *S. aureus*, whereas only indolmycin containing supernatants inhibited the Gram-negative *V. anguillarum*. This is in agreement with

previous studies describing pentabromospeudilin as a compound targeting Gram-positive bacteria (5), 1 2 whereas indolmycin, previously only isolated from Streptomyces species (35,61), inhibits both Gram-3 positive and Gram-negative bacteria (35) and is very potent against staphylococci (31). The genetic basis of 4 the biosynthetic pathways of PBP and indolmycin is not known, but the available mechanistic insight 5 (24,30,54) suggests that it is two distinct biosynthetic pathways with no common precursor. However, as 6 indolmycin and violacein both are tryptophan derived it is not unlikely that their biosynthetic pathways are 7 interlinked at an early stage. Interestingly, the strains H33 and H33S did not produce indolmycin or PBP, yet 8 were active in the live cell assay and in intra-species inhibition. This could potentially be explained by some 9 degree of antibacterial activity of violacein or violacein analogues present in the growing colony, or it could 10 be related to macromolecular antibacterial substances as described in strains CPMOR-1 and CPMOR-2. 11 CPMOR-1 and CPMOR-2 produce a macromolecular L-amino acid oxidase with antibiotic activity (19), but in 12 contrast to H33 and H33S, CPMOR-1 and CPMOR-2 did not cause intra-species inhibition. Therefore it is

also possible that other as yet undiscovered antibiotics are produced by H33 and H33S.

14

15 Our analysis of 16S rRNA gene sequences (figure 1) resolved two clades with no apparent relation to 16 antibiotic production. However, the resolving power of the 16S rRNA gene is limited when performing 17 phylogenetic analyses below the species level and for many species, studies have shown the usefulness of 18 phylogenetic analyses of housekeeping genes such as recA and gyrB. Hence, analysis of antibacterial 19 secondary metabolite production and phylogenetic reconstruction based on the housekeeping genes recA 20 and gyrB divided the 13 strains into three phylotypes with specific production of antibiotics (figure 2). Such 21 a correlation among taxonomic units and secondary metabolite synthesis is well known from the world of 22 fungal natural products (17) and has also been reported for a marine actinobacteria (34). Analysis of GyrB 23 and RecA sequences provides a clearer resolution of taxonomy than 16S rRNA gene sequences when 24 comparing strains at or below the species level(8,27,67), which may explain why phylotype specific

- secondary metabolite production was only partially observed in our previous analysis of 16S rRNA gene
 based phylogeny and chemotypes in 101 *Pseudoalteromonas* strains (62).
- 3

4 The phylogenetic analysis placed the indolmycin producer S4054 near the divergence point of the non-5 indolmycin producers H33 and H33S. This indicates that indolmycin production is an ancestral trait, and we 6 speculate that H33 and H33S may have lost or silenced the indolmycin biosynthetic pathway to rely on a 7 different antagonistic compound in microbial competition since inhibition was observed in the live cell 8 assay, or that the divergence point was shortly before acquisition of the indolmycin producing trait. This 9 raises the question of which taxonomical level to consider when searching for novel chemical diversity, as 10 selection of biological material based on 16S rRNA gene sequences may exclude chemical diversity (33). It 11 is tempting to speculate that phylotype III employs a different strategy when colonizing and competing in 12 the marine environment as these strains showed no inhibition in live cell assays or tests of sterile 13 supernatants. However, antagonism may simply be silenced under the tested growth conditions, as it is known that environmental factors such as host components (3) or microbial interactions (11,45) may 14 15 facilitate the production of antagonistic compounds. Further insight could be gained via whole genome 16 sequencing in order to search for potential clusters of genes involved in e.g. polyketide synthesis or non-17 ribosomal peptide synthesis (22).

18

Often, genes involved in biosynthetic pathways occur grouped within the bacterial genome and are subjects of acquisition or reduction events (13,32,38). Production of antagonistic compounds that play a key ecological role for the producer organism is expected to be under strong selective pressure, as was observed for phylotype I and could be argued for in the indolmycin sub-population in phylotype II. This is consistent with the model of gene fixation in the complexity hypothesis (32) in which genes with a positive selection pressure are maintained at a high occurrence within a taxonomical unit as observed for three closely related *Salinispora* species (34). Species specific chemotypes are known in the marine environment

from e.g. the actinobacterial genus *Salinispora* (34) and were suggested to reflect adaptation to specific
environmental niches. Altogether, this suggests that the compounds indolmycin and PBP play important
roles in the ecology of the producer organism, as their production can be traced through a phylogenetic
reconstruction of closely related strains and thus are maintained on an evolutionary scale. Although the
present study has no data to support previous reports of the antibacterial activity of violacein, we believe
that the compound does play a key role in *P. luteoviolacea* ecology as for instance protection against
predatory microorganisms (43,44).

8

9 Based on the limit number of strains in this study, no apparent correlation among antibiotic production and 10 geographical location was observed within the P. luteoviolacea strains, although the majority of the 11 antagonistic strains were isolated from surfaces and, specifically, all indolmycin-producing strains were 12 isolated from macroalgae or seaweed. The apparent lack of correlation among antibiotic production and 13 geographic origin is in contrast to observations within the marine actinobacterium Salinispora arenicola, where biogeographic patterns including local endemism were observed among polyketide synthases likely 14 15 involved in synthesis of bioactive secondary metabolites (12). Two of the P. luteoviolaceae strains (4054 16 and 4060) had different antibiotic profile and were originally isolated from the same sample (21). No 17 systematic intra-species inhibition pattern was observed and all P. Iuteoviolacea strains, irrespective of 18 antibiotic production profile, were resistant to indolmycin. We speculate that the species gains an 19 ecological advantage by colonizing with strains producing different antibiotics, hence broadening the 20 competitive ability against other species.

21

In summary, we found that the marine bacterium *Pseudoalteromonas luteoviolacea* produces the
bioactives violacein, indolmycin, and PBP. Violacein as a pure compound did not inhibit any of the tested
bacteria, whereas indolmycin inhibited *V. anguillarum* and *S. aureus* but not the *P. luteoviolacea* strains.
We found that the antibiotic profile was correlated to – and hence could have been predicted by – the

phylogenetic analyses. This offered a glimpse of the evolutionary stability of the underlying biosynthetic
 pathways and provides a method for dereplicating bacterial strains as part of a natural product screening
 effort.

- 4
- 5
- 6 Acknowledgements. We would like to thank Dr. Farooq Azam and Dr. Krystal Rypien of Scripps Institution 7 of Oceanography, UCSD, for supplying strain P. luteoviolacea 2ta16; Dr. Antonio Sanchez-Amat of 8 University of Murcia for supplying strains P. luteoviolacea CPMOR-1 and CPMOR-2; and Dr. Tillman Harder 9 of University of New South Wales for supplying strains P. luteoviolacea H33 and H33S. This study was 10 supported by the Programme Commission on Health, Food and Welfare under the Danish Council for 11 Strategic Research. The present work was carried out as part of the Galathea 3 expedition under the 12 auspices of the Danish Expedition Foundation. This is Galathea 3 contribution no. pXXXXXX (to be added 13 if/when accepted)

1		
2		Reference List
3		
4 5	1.	Baker, D. D., M. Chu and U. Oza 2007. The value of natural products to future pharmaceutical discovery. Nat. Prod. Rep. 24 :1225-1244.
6 7	2.	Bowman, J. P. 2007. Bioactive compound synthetic capacity and ecological significance of marine bacterial genus <i>Pseudoalteromonas</i> . Mar. Drugs 5 :220-241.
8 9	3.	Boyd, K. G., D. R. Adams and J. G. Burgess 1999. Antibacterial and repellent activities of marine bacteria associated with algal surfaces. Biofouling 14 :227-236.
10 11 12	4.	Brinkhoff, T., G. Bach, T. Heidorn, L. Liang, A. Schlingloff and M. Simon 2004. Antibiotic production by a <i>Roseobacter</i> clade-affiliated species from the German Wadden Sea and its antagonistic effects on indigenous isolates. Appl. Environ. Microbiol. 70 :2560.
13 14	5.	Burkholder, P. R., R. M. Pfister and F. H. Leitz 1966. Production of a pyrrole antibiotic by a marine bacterium. Appl. Environ. Microbiol. 14 :649-653.
15 16	6.	Clardy, J., M. A. Fischbach and C. T. Walsh 2006. New antibiotics from bacterial natural products. Nat. Biotechnol. 24 :1541-1550.
17 18 19	7.	Clinical and laboratory standards institute . 2006. Guideline M7-A7: Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard. 7 . Clinical and laboratory standards institute. Pennsylvania.
20 21	8.	Coenye, T., D. Gevers, Y. V. Peer, P. Vandamme and J. Swings 2005. Towards a prokaryotic genomic taxonomy. FEMS Microbiol. Rev. 29 :147-167.
22 23	9.	Corley, D. G. and R. C. Durley 1994. Strategies for Database Dereplication of Natural Products. J. Nat. Prod. 57 :1484-1490.
24 25	10.	Delpin, M., J. Melchiorsen, M. Månsson, K. F. Nielsen, and L. Gram. 2012. Indolmycin is a major bioactive in bacterial antagonism by <i>Pseudolateromonas luteoviolacea</i> , strain S4054. In preparation .
26 27 28	11.	Dusane, D. H., P. Matkar, V. P. Venugopalan, A. R. Kumar and S. S. Zinjarde 2011. Cross-species induction of antimicrobial compounds, biosurfactants and quorum-sensing inhibitors in tropical marine epibiotic bacteria by pathogens and biofouling microorganisms. Curr. Microbiol. 1-7.
29 30 31	12.	Edlund, A., S. Loesgen, W. Fenical and P. R. Jensen 2011. Geographic distribution of secondary metabolite genes in the marine actinomycete <i>Salinispora arenicola</i> . Appl. Environ. Microbiol. 77 :5916-5925.
32 33	13.	Egan, S., P. Wiener, D. Kallifidas and E. M. H. Wellington 1998. Transfer of streptomycin biosynthesis gene clusters within streptomycetes isolated from soil. Appl. Environ. Microbiol. 64 :5061.

- Egan, S., P. Wiener, D. Kallifidas and E. M. H. Wellington 2001. Phylogeny of *Streptomyces* species
 and evidence for horizontal transfer of entire and partial antibiotic gene clusters. A. van Leeuw. J.
 Microb. **79**:127-133.
- Fenical, W. 1993. Chemical studies of marine bacteria developing a new resource. Chem. Rev.
 93:1673-1683.
- Fraser, C., E. J. Alm, M. F. Polz, B. G. Spratt and W. P. Hanage 2009. The bacterial species challenge:
 making sense of genetic and ecological diversity. Science 323:741.
- Frisvad, J. C., B. Andersen and U. Thrane 2008. The use of secondary metabolite profiling in
 chemotaxonomy of filamentous fungi. Mycol. Res. 112:231-240.
- Gauthier, M. J. and G. N. Flatau 1976. Antibacterial activity of marine violet-pigmented *Alteromonas* with special reference to the production of brominated compounds. Can. J. Microbiol. 22:1612-1619.
- 19. Gomez, D., E. Espinosa, M. Bertazzo, P. Lucas-Elio, F. Solano and A. Sanchez-Amat 2008. The
 macromolecule with antimicrobial activity synthesized by *Pseudoalteromonas luteoviolacea* strains is
 an L-amino acid oxidase. Appl. Microbiol. Biotechnol. **79**:925-930.
- Goodfellow, M. and H. P. Fiedler 2010. A guide to successful bioprospecting: informed by
 actinobacterial systematics. A. van Leeuw. J. Microb. 1-24.
- Gram, L., J. Melchiorsen and J. B. Bruhn 2010. Antibacterial activity of marine culturable bacteria
 collected from a global sampling of ocean surface waters and surface swabs of marine organisms.
 Mar Biotechnol 12:439-451.
- Gross, H. 2009. Genomic mining-a concept for the discovery of new bioactive natural products. Curr.
 Opin. Drug Discovery Dev. 12:207-219.
- 22 23. Gulder, T. A. M. and B. S. Moore 2009. Chasing the treasures of the sea bacterial marine natural
 products. Curr. Opin. Microbiol. 12:252-260.
- Hanefeld, U., H. G. Floss and H. Laatsch 1994. Biosynthesis of the marine antibiotic
 pentabromopseudilin. Part 1. The benzene ring. J. Org. Chem. 59:3604-3608.
- 26 25. Hasegawa, M., H. Kishino and T. Yano 1985. Dating of the human-ape splitting by a molecular clock
 27 of mitochondrial DNA. J. Mol. Evol. 22:160-174.
- Hibbing, M. E., C. Fuqua, M. R. Parsek and S. B. Peterson 2009. Bacterial competition: surviving and
 thriving in the microbial jungle. Nat. Rev. Microbiol. 8:15-25.
- Hilario, E., T. R. Buckley and J. M. Young 2004. Improved resolution on the phylogenetic
 relationships among *Pseudomonas* by the combined analysis of *atpD, carA, recA* and 16S rDNA. A. van
 Leeuw. J. Microb. 86:51-64.
- Hjelm, M., Ø. Bergh, A. Riaza, J. Nielsen, J. Melchiorsen, S. Jensen, H. Duncan, P. Ahrens, H.
 Birkbeck and L. Gram 2004. Selection and identification of autochthonous potential probiotic
 bacteria from turbot larvae (*Scophthalmus maximus*) rearing units. Syst. Appl. Microbiol. 27:360-371.

- Hook, D. J., E. J. Pack, J. J. Yacobucci and J. Guss 1997. Approaches to automating the dereplication
 of bioactive natural products the key step in high throughput screening of bioactive materials from
 natural sources. J. Biomol. Screen. 2:145.
- Hornemann, U., L. H. Hurley, M. K. Speedie and H. G. Floss 1971. Biosynthesis of indolmycin. J. Am.
 Chem. Soc. 93:3028-3035.
- Hurdle, J. G., A. J. O'Neill and I. Chopra 2004. Anti-staphylococcal activity of indolmycin, a potential
 topical agent for control of staphylococcal infections. J. Antimicrob. Chemothe. 54:549-552.
- Jain, R., M. C. Rivera and J. A. Lake 1999. Horizontal gene transfer among genomes: The complexity
 hypothesis. Proc. Natl. Acad. Sci. U. S. A. 96:3801.
- Jensen, P. R. 2010. Linking species concepts to natural product discovery in the post-genomic era. J.
 Ind. Microbiol. Biotechnol. 37:219-224.
- Jensen, P. R., P. G. Williams, D. C. Oh, L. Zeigler and W. Fenical 2007. Species-specific secondary metabolite production in marine actinomycetes of the genus *Salinispora*. Appl. Environ. Microbiol.
 73:1146-1152.
- Kanamaru, T., Y. Nakano, Y. Toyoda, K. I. Miyagawa, M. Tada, T. Kaisho and M. Nakao 2001. In vitro
 and in vivo antibacterial activities of TAK-083, an agent for treatment of *Helicobacter pylori* infection.
 Antimicrob. Agents Chemother. 45:2455-2459.
- Knight, V., J. J. Sanglier, D. DiTullio, S. Braccili, P. Bonner, J. Waters, D. Hughes and L. ZHANG 2003.
 Diversifying microbial natural products for drug discovery. Appl. Microbiol. Biotechnol. 62:446-458.
- 37. Larsen, T. O., J. Smedsgaard, K. F. Nielsen, M. E. Hansen and J. C. Frisvad 2005. Phenotypic
 taxonomy and metabolite profiling in microbial drug discovery. Nat. Prod. Rep. 22:672-695.
- 38. Lawrence, J. 1999. Selfish operons: the evolutionary impact of gene clustering in prokaryotes and
 eukaryotes. Curr. Opin. Genet. Dev. 9:642-648.
- 24 39. Lichstein, H. C. and V. F. Vandesand 1945. Violacein, an antibiotic pigment produced by
 25 *Chromobacterium violaceum*. J. Infect. Dis. 76:47-51.
- Long, R. A. and F. Azam 2001. Antagonistic interactions among marine pelagic bacteria. Appl.
 Environ. Microbiol. 67:4975-4983.
- Månsson, M., R. K. Phipps, L. Gram, M. H. Munro, T. O. Larsen and K. F. Nielsen 2010. Explorative
 Solid-Phase Extraction (E-SPE) for Accelerated Microbial Natural Product Discovery. J. Nat. Prod.
 73:1126-1132.
- 42. Marsh, W. S., A. L. Garretson and E. M. Wesel 1960. PA 155 A, B, and X. Antibiotics produced by a
 strain of *Streptomyces albus*. Antibiotics & Chemotherapy 10:316-320.
- Matz, C., P. Deines, J. Boenigk, H. Arndt, L. Eberl, S. Kjelleberg and K. Jurgens 2004. Impact of
 violacein-producing bacteria on survival and feeding of bacterivorous nanoflagellates. Appl. Environ.
 Microbiol. 70:1593.

- 44. Matz, C., J. S. Webb, P. J. Schupp, S. Y. Phang, A. Penesyan, S. Egan, P. Steinberg and S. Kjelleberg
 2008. Marine biofilm bacteria evade eukaryotic predation by targeted chemical defense. PLoS ONE 3
- 45. Mearns-Spragg, A., M. Bregu, K. G. Boyd and J. G. Burgess 1998. Cross-species induction and
 enhancement of antimicrobial activity produced by epibiotic bacteria from marine algae and
 invertebrates, after exposure to terrestrial bacteria. Lett. Appl. Microbiol. 27:142-146.
- 6 46. Mikhailov, V. V., L. A. Romanenko, and E. P. Ivanova. 2002. The genus *Alteromonas* and related
 7 Proteobacteria. In M.Dworkin, S.Falkow, E.Rosenberg, K.-H.Schleifer, and E.Stackebrandt (ed.). The
 8 Prokaryotes. 3rd ed., vol. 6.: 597-645. Springer. New York, NY.
- Molinari, G. 2009. Natural Products in Drug Discovery: Present Status and Perspectives. Guzmán,
 Carlos A. and Feuerstein, Giora Z. Pharmaceutical Biotechnology. 655: 13-27. Springer New York.
 Advances in Experimental Medicine and Biology.
- 48. Newman, D. J. 2008. Natural products as leads to potential drugs: An old process or the new hope for
 drug discovery? J. Med. Chem. 51:2589-2599.
- Nicolaou, K. C. and S. T. Harrison 2006. Total Synthesis of Abyssomicin C and atrop-Abyssomicin C.
 Angew. Chem. , Int. Ed. 45:3256-3260.
- Nielsen, K. F., M. Månsson, C. Rank, J. C. Frisvad and T. O. Larsen 2011. Dereplication of microbial
 natural products by LC-DAD-TOFMS. J. Nat. Prod.
- 51. Omura, S., H. Ikeda, J. Ishikawa, A. Hanamoto, C. Takahashi, M. Shinose, Y. Takahashi, H. Horikawa,
 H. Nakazawa and T. Osonoe 2001. Genome sequence of an industrial microorganism *Streptomyces avermitilis*: deducing the ability of producing secondary metabolites. Proc. Natl. Acad. Sci. U. S. A.
 98:12215.
- 52. Östling, J., A. E. Goodman and S. Kjelleberg 1991. Behaviour of IncP-1 plasmids and a miniMu
 transposon in a marine *Vibrio* sp. S14.: isolation of starvation inducible *lac* operon fusions. FEMS
 Microbiol. Ecol 86:83-94.
- 53. Pascual, J., M. C. Macián, D. R. Arahal, E. Garay and M. J. Pujalte 2010. Multilocus sequence analysis
 of the central clade of the genus *Vibrio* by using the 16S rRNA, *recA*, *pyrH*, *rpoD*, *gyrB*, *rctB* and *toxR*genes. Int J Syst Bacteriol 60:154-165.
- Peschke, J. D., U. Hanefeld and H. Laatsch 2005. Biosynthesis of the marine antibiotic
 pentabromopseudilin. 2. The pyrrole ring. Biosci. Biotechnol. Biochem. 69:628-630.
- Sashidhara, K. V. and J. N. Rosaiah 2007. Various dereplication strategies using LC-MS for rapid
 natural product lead identification and drug discovery. Nat. Prod. Commun. 2:193-202.
- Schwartz, R. M. and M. O. Dayhoff 1978. Matrices for detecting distant relationships. Atlas of protein
 sequence and structure 5:353-358.
- 57. Skov, M. N., K. Pedersen and J. L. Larsen 1995. Comparison of pulsed-field gel electrophoresis,
 ribotyping, and plasmid profiling for typing of *Vibrio anguillarum* serovar O1. Appl. Environ.
 Microbiol. 61:1540-1545.

- Smedsgaard, J. and J. Nielsen 2005. Metabolite profiling of fungi and yeast: from phenotype to
 metabolome by MS and informatics. J. Exp. Bot. 56:273-286.
- 59. Tamura, K., D. Peterson, N. Peterson, G. Stecher, M. Nei and S. Kumar 2011. MEGA5: molecular
 evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum
 parsimony methods. Mol. Biol. Evol.
- 60. Taori, K., V. J. Paul and H. Luesch 2008. Structure and activity of largazole, a potent antiproliferative
 agent from the Floridian marine cyanobacterium *Symploca* sp. J. Am. Chem. Soc. 130:1806-1807.
- 8 61. von Wittenau, M. S. and H. Els 1961. The structure of indolmycin. J. Am. Chem. Soc. 83:4678-4680.
- 9 62. Vynne, N. G., M. Månsson, K. F. Nielsen and L. Gram 2011. Bioactivity, chemical profiling and 16S
 rRNA based phylogeny of *Pseudoalteromonas* strains collected on a global research cruise. Mar.
 Biotechnol. doi:10.1007/s10126-011-9369-4.
- Waksman, S. A. and A. Schatz 1943. Strain specificity and production of antibiotic substances. Proc.
 Natl. Acad. Sci. U. S. A. 29:74.
- Wang, H., P. Jiang, Y. Lu, Z. Ruan, R. Jiang, X. H. Xing, K. Lou and D. Wei 2009. Optimization of
 culture conditions for violacein production by a new strain of *Duganella* sp. B2. Biochem. Eng. J.
 44:119-124.
- Wietz, M., A. Schramm, B. Jørgensen and L. Gram 2010. Latitudinal patterns in the abundance of
 major marine bacterioplankton groups. Aquat. Microb. Ecol. (Accepted)
- 46. Yamamoto, S. and S. Harayama 1995. PCR amplification and direct sequencing of *gyrB* genes with
 universal primers and their application to the detection and taxonomic analysis of *Pseudomonas putida* strains. Appl. Environ. Microbiol. 61:1104.
- Yamamoto, S., H. Kasai, D. L. Arnold, R. W. Jackson, A. Vivian and S. Harayama 2000. Phylogeny of
 the genus *Pseudomonas*: intrageneric structure reconstructed from the nucleotide sequences of *gyrB* and *rpoD* genes. Microbiology 146:2385.
- 25 26



Figure 1: Phylogenetic analysis of 16S rRNA gene sequences in MEGA5 (59) using *Pseudoalteromonas atlantica* (accession no. X82134) as outgroup (not shown in tree). The 16S rRNA gene sequences were aligned using MUSCLE, and the evolutionary history inferred using the maximum likelihood method based on the Hasegawa-Kishino-Yano model (25). A phylogenetic tree was constructed using the default MEGA5 settings. Scale bar: substitutions per site. Closed boxes: pentabromopseudilin producer, open circles: indolmycin producer. All strains produce violacein.



0,002

Figure 2: Phylogenetic analysis of concatenated RecA and GyrB amino acid sequences of 13 strains based on the maximum likelihood method. The evolutionary history was inferred by using the maximum likelihood method based on the Dayhoff matrix model(56). The tree with the highest log likelihood (-1597,5915) is shown. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. All positions containing gaps and missing data were eliminated. There were a total of 1000 positions in the final dataset. Evolutionary analyses were conducted in MEGA5 (59) using a discrete Gamma distribution to model evolutionary rate differences among sites (5 categories (+G, parameter = 0,1000)). Clades identified as phylotypes are indicated by roman numerals I-III. Scale bar: substitutions per site. Closed boxes: pentabromopseudilin producer, open circles: indolmycin producer. All strains produce violacein. **Table 1**: *P. luteoviolacea* strains used in this study. The phylogeneticposition of all strains within the *Pseudoalteromonas luteoviolacea* cladewas verified by analysis of partial 16S rRNA gene sequences (>1200 bp,data not shown).

Strain name	Origin	Source
DSM 6061 (T)	Mediterranian, Nice	Surface water
S2607	Pacific, Eastern Australia	Rock surface
S4060-1	Pacific, Costa Rica	Seaweed
2ta16	Florida Keys, USA	<i>M. annularis</i> coral
CPMOR-2	Mediterranian, Murcia	Surface water
NCIMB1944	Mediterranian, Nice	Surface water
S4047-1	Pacific, Costa Rica	Seaweed
S4054	Pacific, Costa Rica	Seaweed
CPMOR-1	Mediterranian, Murcia	Macroalgae
H33	Sydney, Australia	Unknown
H33S	Sydney, Australia	Unknown
NCIMB1942	Mediterranian, Nice	Surface water
NCIMB2035	Mediterranian, Nice	Surface water

Table 2: Inhibitory activity of *P. luteoviolacea* strains in the live cell and supernatant well diffusion agar assays and production of three known antibacterial compounds in marine minimal medium cultures. The production of antibacterial compounds was determined by LC-MS analysis. *V. anguil. Vibrio anguillarum.* x: Inhibition or compound production, respectively. -: No inhibition or compound production.

	Live cell assay inhibition		Inhibition super	by sterile natant	Antibacte	otype		
						Pentabromo-		hyl
Strain	V. anguil.	S. aureus	V. anguil.	S. aureus	Violacein	pseudilin	Indolmycin	4
DSM6061 ^T	х	х	-	х	х	х	-	I
S2607	х	х	-	х	х	x	-	I
S4060-1	х	х	-	х	х	x	-	I
2ta16	х	х	-	х	х	x	-	I
CPMOR-2	х	х	-	х	х	x	-	I
NCIMB 1944	х	x	-	х	х	х	-	I
S4047-1	х	x	х	х	х	-	x	П
S4054 WT	х	x	х	х	х	-	x	П
CPMOR-1	х	x	х	х	х	-	x	П
H33	х	х	-	-	х	-	-	П
H33s	х	х	-	-	х	-	-	П
NCIMB 1942	-	-	-	-	x	-	-	Ш
NCIMB 2035	-	-	-	-	x	-	-	III

Table S1: Intra-species inhibition by *P. luteoviolacea* sterile supernatant in well diffusion agar assays. Substrate blanks were included and did not

2 inhibit any organism. +: inhibition, (+): weak inhibition, -: no inhibition.

Producer						Inh	nibition (+) o	f target strai	า				
strain													
	DSM6061 (T)	S2607	S4047-1	S4054	S4060-1	2ta16	CPMOR-1	CPMOR-2	NCIMB 1942	NCIMB 1944	NCIMB 2035	H33	H33S
DSM6061 ^T	-	+	-	-	+	-	-	-	-	-	+	-	-
S2607	+	+	+	+	+	+	+	+	+	+	+	+	+
S4060-1	+	+	+	+	+	+	+	+	+	+	+	+	+
2ta16	+	+	+	+	+	+	+	+	+	+	+	+	+
CPMOR-2	-	-	-	-	-	-	-	-	-	-	-	-	-
NCIMB 1944	(+)	-	(+)	(+)	-	-	-	-	-	-	-	-	-
S4047-1	+	+	+	+	+	+	+	+	+	+	+	+	+
S4054	+	+	+	+	+	+	+	+	+	+	+	+	+
CPMOR-1	-	-	-	-	-	-	-	-	-	-	-	-	-
H33	-	-	-	-	+	-	-	-	-	-	+	+	-
H33S	-	+	-	-	+	+	-	-	-	-	+	+	-
NCIMB 1942	-	-	-	-	-	-	-	-	-	-	-	-	-
NCIMB 2035	-	-	-	-	-	-	-	-	-	-	-	-	-

Vynne N, Christiansen G, Gram L (2012)

Pseudoalteromonas galatheae S3431 sp. nov., a novel *Pseudoalteromonas* species isolated from a deep-sea polychaete

Manuscript in preparation.

1 Pseudoalteromonas galatheae S3431 sp. nov., a novel Pseudoalteromonas species

2 isolated from a deep-sea polychaete

- 3 Nikolaj G. Vynne¹*, Gunna Christiansen² and Lone Gram¹
- 4
- ⁵ ¹ Technical University of Denmark, National Food Institute, Søltofts Plads, bldg. 221, DK-2800 Kgs.
- 6 Lyngby, Denmark
- ⁷² University of Aarhus, Institute of Medical Microbiology and Immunology, Bartholin Building, DK-
- 8 8000 Aarhus C, Denmark
- 9
- 10 * corresponding author:
- 11 phone: +45 4525 2571
- 12 fax: +45 4588 4774
- 13 e-mail: ngvy@food.dtu.dk
- 14
- 15 December 26th 2011

1 Abstract

2 A novel aerobe marine bacterium was isolated from swab samples of an unidentified polychaete 3 near Canal Concepción, Chile. The strain was gram negative with a polar flagellum, and it grew 4 from 5 to 30 °C. Growth required a NaCl concentration of 1 to 6% (w/v) and pH 5.5-8.9, and a wide 5 range of carbohydrates were utilized as growth substrates. The main components of the total 6 cellular fatty acids were summed feature 3 ($C_{16:1} \omega 7c/C_{16:1} \omega 6c$, 35.82%), $C_{16:0}$ (20.43%) and $C_{17:1}$ 7 ω8c (10.85%). The DNA G+C content of the strain was 39.6 mol%. The most similar strain was *P.arctica* LMG 23753^T (99.7%), and 16S rRNA gene sequence identities ranged from 92.6% to 8 9 99.7% when compared to Pseudoalteromonas type strains. DNA-DNA reassociation values showed 10 less than 70% genomic DNA relatedness to Pseudoalteromonas type strains. Based on the 11 available phylogenetic and phenotypic data, this isolate represents a novel species of the genus Pseudoalteromonas. The name Pseudoalteromonas galatheae is proposed and the type strain is 12 $S3431^{T}$ = LMGXXXXX = DSM XXXX (to be added when/if accepted). 13

14

The genus *Pseudoalteromonas* contains several species that produce different bioactive compounds, and it has been found that for most species pigmentation co-occurs with production of bioactive metabolites (Bowman, 2007;Egan *et al.*, 2002;Vynne *et al.*, 2011). Among the known bioactives produced by pigmented members of the genus are antifouling, anti-bacterial and cytotoxic compounds, however, also non-pigmented species may be of biotechnological interest due to the production of novel enzyme activities (Bowman, 2007).

7 During a research cruise (Galathea 3), bacteria were collected from marine environments and 8 screened for antibacterial activity (Gram et al., 2010). Of the isolated strains, more than one hundred were identified by 16S rRNA gene sequence similarity as Pseudoalteromonas (Gram et al., 9 2010;Vynne et al., 2011). On February 3rd 2007, a swab sample was obtained from an unidentified 10 11 polychaete near Canal Concepción, Chile (-50.4498 ºN, -74.8912 ºE). From subcultures of this sample, strain S3431 was isolated due to production of an intense black pigment. Later analysis 12 would show a high level of 16S rRNA sequence similarity to strains of the so called non-pigmented 13 group of the genus Pseudoalteromonas (Gauthier et al., 1995), in clear contrast to the intense 14 black pigment produced by this strain. The most related type strain was *P. arctica* LMG 23753^T 15 16 (99.7% identical) as determined by a BLAST comparison of 16S rRNA gene sequence of S3431 and Pseudoalteromonas type strains. Several other type strains also had >99% identity and were 17 18 included in this study. Further taxonomic and physiological analysis of this new strain determined 19 that it represents a novel species within the *Pseudoalteromonas* genus.

Strain S3431 was routinely cultured for morphological and physiological characterisation in marine
broth 2216 (Difco, USA) and on marine agar 2216 (Difco, USA) at 25°C. A marine minimal medium
(MMM) containing no carbon source was used as substrate for BioLog assays (Östling *et al.*, 1991).
Temperature requirements for growth were tested in MB 2216 from 5°C to 30°C in 5° intervals,
and at 37°C. NaCl requirements were tested on ½YT agar medium containing 2 g/l yeast extract,
1.25 g/l tryptone, 10 g/l agar and NaCl in concentrations of 0% – 9% (w/v). The pH range
supporting growth was determined in MB, pH 4.2 – 10.3.

27

Cell shape and motility was investigated in wet mounts using an Olympus BX51 phase contrast
 microscope, while the presence of flagella was determined by transmission electron microscopy

1 (TEM). For TEM, colony mass from colonies grown on MA was resuspended in phosphate buffered 2 saline, pH 7.4. Seven μ l of this suspension were placed on a 300-mesh Ni-grid coated with carbon 3 film glow discharged for 30 seconds. The grid was washed using 2 drops of double distilled H₂O 4 and stained with 3 drops of PTA pH 6.9 for 15 seconds, after which the grid was dried by filter 5 paper. TEM was performed at 60 keV on a JEOL1010 transmission electron microscope fitted with 6 a digital camera.

7 Gram-testing was done using the 3% KOH method and aminopeptidase strips (BactiDent). Catalase 8 activity was tested using the 3% H₂O₂ method, and oxidase activity was tested using BD BBL DrySlide. The ability of the strain to ferment glucose was tested using Hugh & Leifsons media 9 10 modified for marine microorganisms by adding 2.0% Instant Ocean salts (Aquarium Systems, Inc., 11 Sarrebourg, France) (Hugh & Leifson, 1953). BioLog GN2 microtiterplates were used to test for 12 carbon assimilation: S3431 was grown on MA overnight and colony mass was resuspended to $OD_{600} = 1.0$ in MMM with no carbon source, supplemented with 5 mM thioglycolate. 150 μ l of this 13 suspension was added to each well of the GN2 microplate. The plate was manually read after 10 14 days of incubation at 25°C. 15

16 Hemolytic activity was tested on Blood Agar Base (Oxoid) with 5% defibrinated calf blood. Production of enzymes was tested by spotting colony mass of \$3431 on medium consisting of 17 28.75 ml/l buffer (85% phosphoric acid 0.08 M, boric acid 0.08 M, glacial acetic acid 0.08 M), 30 g/l 18 Instant Ocean salts and 10 g/l agar and destilled water to 1000 ml. The pH of the medium was 19 adjusted to 6.0 and enzyme substrate was added to a concentration of 0.1% (w/v). Azurine-20 crosslinked (AZCL)-amylose, AZCL-curdlan, AZCL-galactan, AZCL-rhamnogalacturon I, AZCL-xylose 21 and AZCL-dextran were tested. Azo-avicel was used as substrate to test production of endo-22 23 cellulases. Enzyme actitivity was detected by the presence of a colored zone in the agar surrounding the colony. AZCL-substrates and azo-avicel were purchased from Megazyme, Ireland. 24 25 Production of caseinase activity was tested on agar plates containing 100 g/l skim milk powder (Difco), 30 g/l Sea Salts (Sigma) and 15 g/l agar. Chitinase production was assayed on agar plates 26 containing 0.5 g/l peptone, 0.1 g/l yeast extract, 0.01 g/l FePo₄, 30 g/l Sea Salts (Sigma) and 0.2% 27 28 colloidal chitin (Weyland *et al.*, 1970).

1 The *Pseudoalteromonas* sp. S3431 16S-rRNA gene sequence was analysed using BLAST to

2 determine closely related species. A phylogenetic tree was created based on *Pseudoalteromonas*

3 type strain 16S-rRNA gene sequences. These were aligned using MUSCLE and a maximum

4 likelihood tree was created using the Kimura 2-parameter model and 1000 bootstrap replications.

5 Determination of GC-mol% of genomic DNA, cellular fatty acid profiling and DNA-DNA
6 reassociation analyses were carried out by the DSMZ.

7

8 Bacterial cells of the novel isolate were Gram-negative, motile rods with one polar flagella as 9 described for members of the genus *Pseudoalteromonas* (figure 1) (Mikhailov *et al.*, 2002). The cells were approximately $2.2 - 3.0 \mu m$ long and $0.7 - 0.8 \mu m$ wide. The isolate grew at 5 to $30^{\circ}C$, 10 but not at 35°C. No sporulation was observed. Growth was observed at pH 5.5 but not pH 4.3, and 11 at pH 8.9 but pH 9.7. The strain was capable of growth on ½YT agar containing 1 to 6% NaCl, but 12 not in 0.5 or 7% NaCl. The strain was oxidase and catalase positive, and capable of oxidative but 13 14 not fermentative glucose metabolism. Membrane vesicles appeared on several cells as evidenced 15 by TEM, which was previously observed in other *Pseudoalteromonas* species (Nevot *et al.*, 2006).

Carbon source assimilation in the BioLog GN2 microplate system showed that, in addition to the
 carbon sources listed in table 1, S3431 was able to utilize α-cyclodextrin, dextrin, glycogen, tween
 40, tween 80, D-Cellobiose, gentiobiose, lactulose, D-raffinose, acetic acid, D-galacturonic acid, α ketoglutaric acid, propionic acid, succinic acid, L-alaninamide, L-alanine, L-alanyl-glycine, L asparagine, L-glutamic acid, glycyl-L-glutamic acid, L-proline, L-serine, L-threonine, inosine, uridine,
 α-D-glucose-6-phosphate and D-glucose-6-phosphate as sole carbon sources.

22 The novel strain produced amylase but not curdlanase, galactanase, rhamnogalacturonase,

23 xylanase or dextranase. No endo-cellulases were produced. Chitinase production was not

24 detected, however agarolytic activity was observed on both chitin plates and agar medium

containing 0.5 g/l peptone, 0.1 g/l yeast extract, 0.01 g/l FePO₄·H₂O and 30 g/l Sea Salts (Sigma).

26 Caseinase activity was observed as clearing zones on skim milk agar.

The 16S-rRNA gene sequence was compared to 16S-rRNA sequences from *Pseudoalteromonas*type strains using the BLAST tool. The most similar strain was *P.arctica* LMG 23753^T (99.7%), and

identities ranged from 92.6% to 99.7%. In a minimum evolution phylogenetic tree, the strain
 placed itself firmly within the so called non-pigmented group (supplementary figure 1), in contrast

3 to the intense black pigment produced by this strain.

The DNA-DNA reassociation values of S3431 to type strains with 99% 16S rRNA identity are shown
in table 2, and were below the 70% recommended limit for delineating new species (Wayne *et al.*,
1987). G+C content in genomic DNA was 39.6 mol%, which is in range with other *Pseudoalteromonas* species (Gauthier *et al.*, 1995). The main cellular fatty acids were summed
feature 3 (C_{16:1} ω7c/C_{16:1} ω6c, 35.82%), C_{16:0} (20.43%) and C_{17:1} ω8c (10.85%) as shown in table 3
which confirms the position of strain S3431 within the genus *Pseudoalteromonas* (Ivanova *et al.*,

10 2000).

11

12 Based on the available phenotypic and phylogenetic information, strain S3431 should be

13 recognized as a new species within the genus *Pseudoalteromonas* and the name

14 *Pseudoalteromonas galatheae* sp. nov. is proposed.

15

16 Description of *Pseudoalteromonas galatheae* sp. nov.

Pseudoalteromonas galatheae (ga.la.the'ae. N.L. n. galathea, referring to the research expedition on
which the type strain was first isolated).

19 Cells are motile straight rods with polar flagella, $2.2 - 3.0 \mu m$ long and $0.7 - 0.8 \mu m$ wide. The cells 20 are Gram-negative, non-spore forming, strictly aerobe and mainly occur as single cells. When 21 grown on MA 2216 at 25 °C, the strain forms raised circular black to dark brown colonies, which appear smooth and shiny. A brown pigment diffusing into the agar is produced. Growth occurs at 4 22 - 30 °C, with no growth at 37 °C. Growth is observed within a pH range of 5.5-8.9. Substrate NaCl 23 24 content from 1-6% supports growth. The strain utilzes D-glucose, D-mannose, D-galactose, maltose, sucrose, melibiose, lactose, succinate, D-gluconate, D-mannitol, sorbitol, citrate, xylose, 25 trehalose, acetate, L-arginine, α -cyclodextrin, dextrin, glycogen, tween 40, tween 80, D-cellobiose, 26 27 gentiobiose, lactulose, D-raffinose, acetic acid, D-galacturonic acid, α -ketoglutaric acid, propionic

1 acid, succinic acid, L-alaninamide, L-alanine, L-alanyl-glycine, L-asparagine, L-glutamic acid, glycyl-

- 2 L-glutamic acid, L-proline, L-serine, L-threonine, inosine, uridine, α-D-glucose-6-phosphate and D-
- 3 glucose-6-phosphate as sole carbon sources. It does not grow on D-fructose, sorbitol, N-
- 4 acetylglucosamine or pyruvate. The strain produces amylase and hydrolyzes agar and casein, but is
- 5 not hemolytic. The primary cellular fatty acids are summed feature 3 ($C_{16:1} \omega 7c/C_{16:1} \omega 6c$, 35.82%),
- 6 $C_{16:0}$ (20.43%) and $C_{17:1}$ $\omega 8c$ (10.85%).
- 7 The type strain is S3431^T (= DSM XXXXX = LMG XXXXXX, to be added when accepted), isolated from
- 8 a marine polychaete in Canal Concepción, Chile. The DNA G+C mol content of the type strain is
- 9 39.6%.
- 10

11 Acknowledgements

- 12 We thank Dr. Bernhard Schink for ensuring a correct specific epithet. This study was supported by the
- 13 Programme Commission on Health, Food and Welfare under the Danish Council for Strategic Research. The
- 14 present work was carried out as part of the Galathea 3 expedition under the auspices of the Danish
- 15 Expedition Foundation. This is Galathea 3 contribution no. pXXXXXX (to be added if/when accepted)
- 16

1	
2	Reference List
3	
4	Bowman, J. P. (2007). Bioactive compound synthetic capacity and ecological significance of marine bacterial
5	genus Pseudoalteromonas. Mar Drugs 5, 220-241.
6	Egan, S., James, S., Holmström, C. & Kjelleberg, S. (2002). Correlation between pigmentation and
7	antifouling compounds produced by Pseudoalteromonas tunicata. Environ Microbiol 4, 433-442.
8	Gauthier, G., Gauthier, M. & Christen, R. (1995). Phylogenetic analysis of the genera Alteromonas,
9	Shewanella, and Moritella using genes coding for small-subunit ribosomal-RNA sequences and
10	division of the genus Alteromonas into 2 genera, Alteromonas (emended) and Pseudoalteromonas
11	gen nov, and proposal of 12 new species combinations. Int J Syst Bacteriol 45, 755-761.
12	Gram, L., Melchiorsen, J. & Bruhn, J. B. (2010). Antibacterial activity of marine culturable bacteria collected
13	from a global sampling of ocean surface waters and surface swabs of marine organisms. Mar
14	Biotechnol 12 , 439-451.
15	Hugh, R. & Leifson, E. (1953). The taxonomic significance of fermentative versus oxidative metabolism of
16	carbohydrates by various gram negative bacteria. J Bacteriol 66, 24-26.
17	Ivanova, E. P., Zhukova, N. V., Svetashev, V. I., Gorshkova, N. M., Kurilenko, V. V., Frolova, G. M. &
18	Mikhailov, V. V. (2000). Evaluation of phospholipid and fatty acid compositions as chemotaxonomic
19	markers of Alteromonas-like Proteobacteria. Current Microbiology 41, 341-345.
20	Mikhailov, V. V., Romanenko, L. A. & Ivanova, E. P. (2002). The genus Alteromonas and related
21	Proteobacteria. In <i>The Prokaryotes</i> , 3rd ed., vol. 6. edn, vol. 6, pp. 597-645. Edited by In M.Dworkin,
22	S.Falkow, E.Rosenberg, KH.Schleifer & E.Stackebrandt (ed.). New York, NY: Springer.
23	Nevot, M., Deroncelé, V., Messner, P., Guinea, J. & Mercadé, E. (2006). Characterization of outer
24	membrane vesicles released by the psychrotolerant bacterium <i>Pseudoalteromonas antarctica</i> NF3.
25	Environ Microbiol 8 , 1523-1533.

- Östling, J., Goodman, A. E. & Kjelleberg, S. (1991). Behaviour of IncP-1 plasmids and a miniMu transposon
 in a marine *Vibrio* sp. S14.: isolation of starvation inducible *lac* operon fusions. *FEMS Microbiol Ecol* 86, 83-94.
- Vynne, N. G., Månsson, M., Nielsen, K. F. & Gram, L. (2011). Bioactivity, chemical profiling and 16S rRNA
 based phylogeny of *Pseudoalteromonas* strains collected on a global research cruise. *Mar Biotechnol*, doi:10.1007/s10126-011-9369-4.
- Wayne, L. G., Brenner, D. J., Colwell, R. R., Grimont, P. A. D., Kandler, O., Krichevsky, M. I., Moore, L. H.,
 Moore, W. E. C., Murray, R. G. E. & other authors (1987). Report of the ad hoc committee on
 reconciliation of approaches to bacterial systematics. *Int J Syst Bacteriol* 37, 463.
- Weyland, H., Rüger, H.-J. & Schwarz, H. (1970). Zur Isolierung und Identifizierung mariner Bakterien. Ein
 Beitrag zur Standardisierung und Entwicklung adaequater Methoden. Veroeff Inst Meeresforsch
 Bremerhaven 12, 269-296.
- 13

- 1 **Figure 1:** TEM of cells of *Pseudoalteromonas galatheae* S3431 showing the polar flagellum.
- 2 Additionally, vesicle-like structures appear near the cell membrane.



Table 1: Characteristics of *Pseudoalteromonas* type strains. 1: *P. galatheae* S3431, 2: *P. arctica*^T, 3: *P. translucida*^T, 4: *P. haloplanktis*^T, 5: *P. agarivorans*^T, 6: *P. aliena*^T, 7: *P. atlantica*^T, 8: *P. carrageenovora*^T, 9: *P. distincta*^T, 10: *P. elyakovii*^T, 11: *P. espejiana*^T, 12: *P. issachenkonii*^T, 13: *P. nigrifaciens*^T, 14: *P. paragorgicola*^T, 15: *P. tetraodonis*^T, 16: *P. undina*^T.

Characteristic	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
DNA G-C content (mol%)	39.6	39	46.3	43.3	43.8	41- 43	41.2	39.5	43.8	38.5	43	43	39- 41	41.1	41.5	42.2
Growth at:																
4 ºC	+	+	+	-	-	+	-	(+)	-	v	-	+	+	+	+	+
30 ºC	+	-	+	v	+	+	+	+	+	+	+	+	+	+	+	ND
40ºC	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1% NaCl	+	+	+	+	+	v	+	+	+	+	+	+	+	+	+	+
3-6% NaCl	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
8% NaCl	-	+	-	+	+	-	-	+	+	+	+	+	+	-	+	+
9% NaCl	-	+	-	+	+	-	-	+	+	+	+	+	+	-	+	+
10% NaCl	-	-	-	+	-	-	-	+	+	+	+	+	+	-	+	+
Enzymatic activity:																
Catalase positive	+	ND	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Oxidase positive	+	ND	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Amylase	+	-	+	v	+	+	+	-	-	+	+	-	v	+	-	v
Caseinase	+	+	+	+	+	v	+	+	-	ND	ND	+	-	+	ND	ND
Agarase	+	-	-	-	+	-	+	-	-	-	-	-	-	-	-	-
Chitinase	-	ND	-	v	ND	-	-	-	-	-	-	+	-	-	-	+
Utilization of:																
D-Glucose	+	+	-	+	+	v	+	+	+	+	+	+	+	ND	+	+
L-Arabinose	-	ND	ND	-	-	ND	-	-	-	-	-	ND	ND	ND	ND	v
D-Mannose	+	-	ND	+	-	-	+	-	+	+	v	-	v	ND	-	-
D-Galactose	+	+	+	v	-	-	+	v	-	+	+	+	+	+	+	-
D-Fructose	-	-	ND	+	-	-	+	+	+	+	v	+	+	ND	-	-
Maltose	+	+	-	+	+	+	+	+	-	+	+	+	+	+	+	+
Sucrose	+	+	ND	v	-	ND	+	+	-	+	+	+	v	ND	+	+
Melibiose	+	+	-	-	-	-	+	+	+	v	+	+	+	-	-	-
Lactose	+	-	-	-	-	-	+	+	+	v	+	+	+	-	-	-
D-Gluconate	+	+	ND	-	-	-	-	-	-	-	-	-	+	ND	-	-
Glycerol	+	-	ND	-	-	-	+	+	+	-	v	-	+	ND	-	-
Succinate	+	-	ND	v	-	-	+	+	-	+	-	+	+	ND	v	+
D-Mannitol	+	+	+	v	+	ND	+	+	-	+	+	+	-	+	-	-
Sorbitol	-	ND	ND	-	ND	-	-	-	-	-	-	-	v	-	-	-
Citrate	+	-	-	+	-	-	v	+	+	v	+	+	+	-	+	-
N-Acetylglucosamine	-	-	ND	-	-	+	-	-	-	-	-	-	ND	ND	ND	+
Xylose	+	-	-	-	ND	-	-	-	-	+	+	-	v	ND	-	-
Trehalose	+	-	ND	-	ND	-	+	-	-	-	+	-	+	ND	-	v
Acetate	+	-	ND	+	-	-	+	+	+	v	+	+	+	ND	+	+
Pyruvate	-	+	ND	+	-	-	+	+	+	+	+	+	+	ND	+	+
L-Arginine	+	ND	-	-	ND	ND	-	-	ND	-	+	ND	v	-	-	v

Table 2: Per cent DNA-DNA similarity in 2X SSC at 66°C {Vynne, 2012 455 /id}. All related type strains have less than 70% DNA-DNA similarity to S3431, which is the accepted threshold for species delineation {Wayne, 1987 152 /id}. WIP: Work in progress.

Pseudoalteromonas species	Per cent DNA:DNA similarity to strain S3431
<i>P. agarivorans</i> DSM 14585^{T}	28.2
P. aliena LMG 22059 [™]	44.5
<i>P. atlantica</i> NCOMB 301^{T}	26.9
<i>P. carrageenovora</i> DSM 6820^{T}	14.8
<i>P. distincta</i> CIP 105340^{T}	44.5
<i>P. elyakovii</i> LMG 14908 [™]	44.2
P. espejiana DSM 9414 [™]	48.0
P. issachenkonii LMG19697 [™]	33.6
P. nigrifaciens LMG 2227 ^T	15.4
<i>P. paragorgicola</i> LMG 19696^{T}	58.4
<i>P. tetraodonis</i> DSM 9166^{T}	11.3
<i>P.undina</i> LMG 2880^{T}	13.8
<i>P. antarctica</i> CECT4664 ^T	WIP

significant amounts.	
Fatty acid	S3431 [™]
C _{12:0}	1.25
C _{14:0}	0.51
C _{15:0}	-
C _{16:0}	20.43
C _{17:0}	5.76
C _{18:0}	1.14
C _{15:1} ω8c	1.60
C _{17:1} ω8c	10.85
C _{17:1} ω6c	0.55
C _{18:1} ω7c	7.77
C _{16:0} iso	1.18
C _{17:0} iso	0.60
C _{17:0} anteiso	0.56
C _{10:0} 3OH	1.07
C _{11:0} 3OH	1.11
C _{12:0} 3OH	5.15

Summed feature 1 (C_{15:0} iso H/C_{13:0} 3OH)

Summed feature 3 ($C_{16:1} \omega 7c/C_{16:1} \omega 6c$)

Other

Table 3: Cellular fatty acid composition of *Pseudoalteromonas galatheae* S3431^T. Fatty acids
 constituting less than 0.5% of the total amount were not included. -, Not detected in

0.62

35.82

4.03

Supplementary figure 1: Phylogenetic tree based on 16S rRNA gene sequences of *Pseudoalteromonas* type strains. The novel species *P. galatheae* S3431 sp. nov. placed itself in the non-pigmented clade, near *P. translucida* DSM 14402^T, *P. antarctica* CECT 4664^T, *P. nigrifaciens* NCIMB 8614^T, and *P. haloplanktis* DSM6060^T. The 16S rRNA gene sequences were aligned in MEGA 5 and the phylogenetic reconstruction was based on the maximum likelihood method with the Kimura 2-parameter model, with a Gamma distribution model allowing invariant sites to model the evolutionary rate differences. Scale bar: 0,005 substitutions per nucleotide site.



⊢ 0,005
Vynne N, Månsson M, Nielsen KF, Milton D, Gram L (2012)

Production of a putatively novel acylated homoserine lactone-like quorum sensing activating molecule by *Pseudoalteromonas luteoviolacea* S4054

Manuscript in preparation.

- 1 Production of a putatively novel acylated homoserine lactone-like quorum sensing
- 2 activating molecule by *Pseudoalteromonas luteoviolacea* S4054
- 3
- 4 Vynne, N. G.¹*, Månsson, M.², Nielsen, K.F.², Milton, D.³ and Gram, L.¹
- ¹: National Food Institute, Technical University of Denmark, Søltofts Plads bld. 221, DK-2800 Kgs. Lyngby,
 Denmark
- ²: Department of Systems Biology, Technical University of Denmark, Søltofts Plads bld. 221, DK-2800 Kgs.
- 8 Lyngby, Denmark
- 9 ³: Department of Molecular Biology, Umeå University, S-901 87 Umeå, Sweden.
- 10
- 11 *: corresponding author.
- 12 e-mail: <u>ngvy@food.dtu.dk</u>
- 13 Keywords: Quorum sensing, *Pseudoalteromonas luteoviolacea*, antibiotics, lux,
- 14
- 15 Running title: AHL-like molecules in *Pseudoalteromonas luteoviolacea*

16 Abstract

17 The bioactive strain Pseudoalteromonas luteoviolacea S4054 produces antibiotics and compounds capable 18 of eliciting a signal response in the quorum sensing monitor strains Agrobacterium tumefaciens pZLR4, 19 Chromobacterium violaceum CV026, and Escherichia coli JM109 pSB401. Since quorum sensing in some 20 cases control the production of antibiotics, we hypothesized that quorum sensing might be involved in 21 control of antibiotic production by P. luteoviolacea S4054. Through shotgun cloning the plul gene was 22 identified as the gene coding for production of the signaling compounds. plul showed homology to luxl AHL 23 synthethases from Vibrionaceae and other Proteobacteria, but formed a monophyletic branch in 24 phylogenetic analysis of luxl homologues indicating that plul encodes a novel AHL synthethase. Genome 25 sequencing revealed that *plul* is located in an operon structure with downstream genes that may constitute 26 a biosynthetic pathway potentially involved in biosynthesis of the potent antibiotic indolmycin. 27 Immediately downsteam of the operon, the *luxR* homologue *pluR* was located on the opposite DNA strand. 28 Chemical analysis of the signaling molecules revealed AHL-like traits, however it was not possible to identify the compounds using standard methods for AHL identification indicating that the signaling molecules might 29 30 represent novel AHL-like compounds.

32 Introduction

33 Cell-density dependant signaling occurs widely among bacteria and is commonly termed quorum sensing 34 (QS). QS allows a population of cells the benefits of coordinated gene expression, sensing the presence of 35 other signal molecule producing microorganisms or possibly probing the local environmental conditions. A 36 common QS system in the y-Proteobacteria is the LuxI-LuxR system. This QS system was first described in 37 Vibrio fischeri (8), where it regulates the expression of bioluminescence genes in a cell-density dependent 38 manner. The signaling molecules of this system are N-acylated homoserine lactones (AHLs) and are 39 produced by LuxI AHL synthethases (9). AHLs are produced continuously at low cell-densities AHLs, but as 40 cell-density in the local environment increases, the concentration of AHLs reaches a threshold level and 41 activates the QS regulated system through binding to the transcriptional regulator LuxR. This activates gene 42 expression of the QS regulated phenotype, in the case of V. fischeri bioluminescence. In some bacteria, expression of the luxI gene is also increased leading to a strong autoinducing effect (25). Some non-AHL 43 molecules such as diketopiperazines are capable of modulating the Lux-system to activate or repress the 44 45 signaling response (11,15). Hence the QS regulated phenotypes may be influenced by AHLs or by other 46 compounds.

47 For some bacteria, QS no doubt holds an important role in microbial community interactions through 48 regulation of phenotypes such as biofilm formation, motility and production of antimicrobial compounds 49 (10). In the marine environment, multiple bacterial species will be present on most surfaces and 50 competition for space and nutrients is believed to play a major role in shaping the microbial community 51 (14). The production of antibacterial compounds is one way for a bacterium to modulate the surrounding 52 community and QS regulation of biosynthetic pathways for production of antibacterial compounds is well 53 known (6,32). Recently production of the potent antibacterial compound tropodithietic acid was demonstrated to be QS controlled in Phaeobacter gallaeciensis (1), a member of the ecologically significant 54 55 Roseobacter clade (2). Previous studies have indicated that AHLs are produced by Pseudoalteromonas

species associated with different microbial communities (17,18), however, few studies have addressed the actual regulatory role of the compounds. Violacein, which is a key metabolite of the *P. luteoviolacea* species and may protect against predation (22), is produced under QS regulation in both distant and closely related organisms such as *Chromobacterium violaceum* and *Pseudoalteromonas* sp. 520P1 (23,37). A LuxI-LuxR system in *P. luteoviolacea* could potentially regulate production of violacein, but also of other bioactive compounds such as pentabromopseudilin and indolmycin which we previously identified in a collection of *P. luteoviolacea* strains (20,35,35).

63 The purpose of the present study was to determine if QS regulatory systems could be involved in antibiotic production by P. luteoviolacea. We demonstrate that compounds inducing AHL-monitor bacteria are 64 65 present in some but not all violacein producing P. luteoviolacea strains. In one AHL-monitor-inducing strain, 66 P. luteoviolacea S4054, we identify a luxl gene homologue and we investigate the genomic context of this 67 gene to identify a gene homologue to known luxR genes. We identify a putative novel biosynthetic pathway in an operon structure with the *luxl* gene, which is likely quorum sensing regulated. Thorough chemical 68 69 analyses did, however, not lead to identification of any known AHL molecule and we conclude that the AHL-70 like compound(s) likely represent a novel AHL structure.

71

72 Materials and methods

Strains, plasmids and media. Thirteen *Pseudoalteromonas luteoviolacea* strains previously described (34)
were included in this study. *P. luteoviolacea* strains were routinely cultured at 25 °C in a marine minimal
medium (27) with 4 g/l mannose and 3 g/l casamino acids, unless otherwise mentioned. *Chromobacterium violaceum* CV026 was cultured in Luria broth (per litre: Bacto Tryptone, 10 g; Bacto yeast extract, 5 g; and
sodium chloride, 5 g) at 25 °C with 20 µg/ml kanamycin (23). *Agrobacterium tumefaciens* pZLR4 was
cultured in ABT medium (per litre: (NH₄)₂SO₄ 0.4 g, Na₂HPO₄ 0.6 g, KH₂PO₄ 0.3 g, NaCl 0.3 g, 1 mM MgCl₂,
0.1 mM CaCl₂, 0.01 mM FeCl₃, 2.5 mg thiamin supplemented with 0.5% glucose and 0.5% cas-amino acids)

(4) supplemented with 20 μg/ml gentamicin. The bacterial strains and plasmids used for cloning
procedures in this study are listed in table 1. *E. coli* strains were grown at 37 °C in Luria broth
supplemented with carbenicillin 100 μg/ml or tetracyclin 10 μg/ml. All broth cultures were incubated with
agitation, and supplemented with 1.2% agar as required.

84 Bioassays for detection of potential QS activating molecules. Well diffusion agar assays were used to 85 screen for the presence of QS activating molecules in sterile filtered culture supernatants and ethyl acetate 86 extracts. Ethyl acetate extracts were prepared as described for LC-MS analysis. To prepare the assays, 50 ml 87 overnight culture of C. violaceum CV026 was added to 100 ml molten LB-agar at 46 °C, supplemented with 88 20 μg/ml kanamycin (23), or 50 ml overnight culture of *A. tumefaciens* pZLR4 were added to 100 ml ABT-89 agar at 46 °C supplemented with 20 µg/ml gentamicin and 75 µg/ml X-gal (3). The solution was poured into 90 petri dishes to form a 5 mm thick layer. Wells were cut after the agar had solidified and 50 μ l sample was 91 loaded per well. The assays were incubated at 25 °C and read after 24 and 48 h. Simple T-streaks against C. 92 violaceum CV026 were used to screen E. coli clones for production of QS activating compounds (23).

The monitor strain *E. coli* JM109 pSB401 (40) was used as a heterologous host to express and detect
potential QS activating molecules. The *E. coli* JM109 clones containing both the pSB401 plasmid and a
pBluescript construct were plated on LB-agar with 50 µg/ml carbenicillin, 10 µg/ml tetracyclin, 20 µg/ml
IPTG and 50 µg/ml X-gal, incubated overnight at 37 °C and screened for bioluminescence using a LAS-4000
(Fujifilm Life Sciences, USA).

Thin layer chromatography (TLC). Ethyl acetate extracts or synthetic AHL standards were spotted onto a
C₁₈ TLC plate (TLC aluminum sheets 20×20 cm², RP-18 F_{254 S}, 1.05559. Merck 64271 Darmstadt, Germany)
and the plates were developed in 10 ml 60:40 (v/v) methanol/water until the front reached the top (28).
The TLC plate was dried and agar top-layers were prepared as for the well diffusion assay. The top layer of
bacterium-containing agar was poured onto the TLC plate immediately following preparation, and
incubated at 25 °C for 24h.

PCR conditions, DNA techniques, DNA sequencing and bioinformatics. PCR was performed as previously described (5,24). Unless otherwise stated, all conditions for the various DNA techniques were as described by (30). Reaction conditions for DNA-restriction enzymes were as suggested by the manufacturers. DNA sequencing of the unknown DNA regions obtained from shotgun cloning was performed by primer walking in two directions from known regions of DNA sequence using the primers listed in table 2. Sequence data were processed and assembled in CLC-bio DNA workbench, and MEGA 5 was used for sequence alignment and phylogenetic reconstruction.

111 Shotgun cloning to identify genes involved in QS. Genomic DNA was purified from P. luteoviolacea \$4054 using the QIAGEN Genomic-Tip/100 kit. The genomic DNA was digested with EcoRI or BamHI and ligated 112 113 into the corresponding pBluescript digest. The shotgun library obtained in pBluescript was transformed into 114 chemically competent E. coli JM109 pSB401, incubated overnight at 37 °C and screened for luminescence. 115 Luminescent clones were tested for induction of violacein production in C. violaceum CV026 by T-streaks, and the pBluescript vector from positive clones was cloned by transformation into chemically competent E. 116 117 *coli* DH5- α λpir , plated on LB-agar with 50 μ g/ml carbenicillin and incubated overnight at 37 °C. The 118 resulting clones were verified for activity in the C. violaceum CV026 bioassay and the pBluescript DNA insert 119 was sequenced using a primer walking method.

120 LC-MS and LC-MS/MS analyses. P. luteoviolacea S4054, E. coli NV5, and E. coli NV1 were grown in MMM 121 for 3 days at 25°C and the cultures were extracted with equal volumes of ethyl acetate, centrifuged, and 122 the ethyl acetate phase was evaporated under N2 to dryness. Samples were redissolved in 500 μ l MeOH 123 and analysed by LC-MS. LC-MS analyses were performed on an Agilent 1100 System equipped with a 124 UV/VIS photo diode array detector (scanning 200–600 nm). Separation was done on a 100 mm×2 mm i.d., 3 μm Gemini C6-phenyl column (Phenomenex, Torrance, CA, USA), running at 40°C using gradient (15-100% 125 over 20 minutes) of water (H2O) and acetonitrile (MeCN); both buffered with 50 µl/l trifluoroacetic acid) at 126 127 a flow of 300 μ l/min. LC-MS/MS analyses were performed on a maXis quadrupole time of flight mass

spectrometer (Bruker Daltonics, Bremen, Germany) equipped with an electrospray (ESI) ion source. The MS
was connected to an Ultimate 3000 UHPLC system (Dionex, Sunnyvale, CA) equipped with a diode-array
detector. Separation was performed at 40 °C on a 150 mm × 2.1 mm ID, 2.6 µm Kinetex C18 column
(Phenomenex) using a linear water/MeCN (both buffered with 20 mM formic acid) gradient starting from
15% MeCN and increased to 100% in 13 min at a flow of 0.4 mL min–1. Experiments were performed in ESI+
with a data acquisition range of *m/z* 100–1200 with collision energy of 40 V. The MS was calibrated using
sodium formate automatically infused prior to each analytical run.

Purification and structure elucidation of the QS activating molecule. Preparative TLC was performed on C18 plates (20 x 20 cm, Analtech). On each plate, approximately 20 mg of crude extract was loaded in 500 uL 50/50 (v/v) MeOH/H2O with a solvent streaker. Plates were developed in 60% MeOH in water. To visualize the active compounds, the plates were overlaid with agar containing *C. violaceum* CV026 or *A. tumefaciens* pZLR4 as described above but limited to a 3 cm wide band at the left and right edges.

140 Three different solid phase extraction columns were attempted on a small scale (2 mg crude extract):

141 Strata-X (30 mg/1 mL, Phenomenex), Strata-NH2 (100 mg/1 mL, Phenomenex), and diol (100 mg/1 mL,

142 Phenomenex). From all three columns, five fractions were collected and tested in the QS assays. From

143 Strata-X: 100% H2O, 25% MeOH in H2O, 40% MeOH, 60% MeOH, and 100% MeOH. From Strata-NH2: 95%

144 MeCN in H2O, 60% MeCN, 40% MeCN, 25% MeCN, and 100% H2O. From diol: 100% heptane, 100%

dichlormethane, 100% EtOAc, 50% EtOAc in MeOH, and 100% MeOH. Fractions were evaporated under N2

and redissolved in 100% MeOH immediately before biotesting.

147 An 8 L culture of *P. luteoviolacea* S4054 was grown in Marine Broth 2216 (Difco, USA). On day 3, a mixture

148 (50/50) of sterile XAD-7 (Sigma-Aldrich, St. Louis, MO) and Diaion HP20SS resin (Sigma-Aldrich) was added

to the broth (12 g of resin L–1). After 24 h the resin was filtered off and washed with water (2 x 2 L),

150 followed by extraction with methanol/water (50/50 v/v; 800 mL) (24h) and methanol (2 x 800 mL) (24 h).

151 All organic extracts were pooled and dried completely to give a crude extract (2.0712 g). Violacein and

indolmycin were quantitatively removed from the extract by cation-exchange as previously described (20).

153 The remaining extract was subjected to anion-exchange. The crude extract (640 mg) was re-dissolved in 154 methanol/water (50/50 v/v; 5 ml) and mixed with 3 g Strata-SAX (Phenomenex, Torrance, CA) and dried 155 before packing into a 100 g SNAP column (Biotage, Uppsala, Sweden). Using an Isolera flash purification 156 system (Biotage), the column was equilibrated (5 mL/min) upside-down with water + ammonium hydroxide 157 (2%). The column was then turned and washed with methanol with ammonium hydroxide (2%) (fraction 1). 158 The column was then stripped with methanol with formic acid (1%) and washed until no colour remained 159 (fraction 2). Fraction 1, enriched with QS activity, was redissolved in EtOAc and absorbed onto xx g Isolute 160 diol (Biotage) and added to a 100 g SNAP column with pure diol (10 g). A total of 12 fractions were collected 161 ranging from heptane, dichloromethane, EtOAc to pure MeOH (5 mL/min). The fractions with QS activity 162 (100 % dichloromethane to 20% dicholormethane) were pooled and further separated on a Luna II C18 163 column (250 × 10 mm, 5 μ m) (Phenomenex) using a 10-90% MeCN/H₂O gradient (buffered with 20 mM FA, 164 flow rate 5 mL/min) over 20 minutes on a Gilson 322 liquid chromatograph with a 215 liquid 165 handler/injector (BioLab, Risskov, Denmark). This yielded four fractions with QS activity. NMR spectra were 166 recorded on a Varian Unity Inova 500 MHz spectrometer equipped with a 5 mm probe using standard pulse 167 sequences. The signals of the solvent were used as internal references (δH 4.80 for D₂O). 168 Stability test of AHL-like compounds. The lactone ring of classic AHLs opens under alkaline conditions, 169 leading to a loss of QS inducing activity. To test the AHL-like nature of the compounds obtained from 170 chemical fractionation, the compounds and a mix of synthetic AHLs were suspended in solutions of pH 2, 171 pH 7, and pH 11 for two hours, after which samples were tested in well diffusion agar assays against the 172 monitor strain A. tumefaciens pZLR4. 173 Genome mining for luxI – luxR gene pairs and quorum sensing controlled gene clusters. A permanent 174 draft genome of *P. luteoviolacea* was obtained by sequencing and assembly as described elsewhere (33). 175 The genome sequence was submitted to the IMG ER pipeline for annotation and visualization (21). The 176 nucleotide sequence obtained from cloning experiments was identified and the neighboring gene clusters 177 investigated. *luxR* homologue genes were identified by a search for genes with the Pfam 03472 autoinducer binding domain or genes in COG2771. The resulting genes were submitted to BLASTX and LuxR to identify
LuxR homologues. The gene cluster downstream of the *luxI* homologue gene was investigated for its
putative biological function through analysis of the annotations and BLAST.

181

182 **Results and discussion**

Characterization and production of AHL-like molecules by Pseudoalteromonas luteoviolacea strains. 183 184 Thirteen Pseudoalteromonas luteoviolacea strains were cultured overnight in MMM and sterile filtered 185 supernatants were tested in well assays using C. violaceum CV026 or A. tumefaciens pZLR4 monitor strains. 186 Three strains (S4047-1, S4054, and CPMOR-1) activated both monitor strains indicating the presence of 187 AHL-like compounds in the growth supernatants. Ten strains did not activate either monitor strain, but 188 were previously shown to produce violacein. Hence, violacein production in P. luteoviolacea is not 189 universally regulated by molecules that elicit a response in traditional QS monitor strains. We previously 190 showed that the three strains are closely related produce indolmycin, an antibiotic compound not 191 produced by the remaining 10 P. luteoviolacea strains (34), hence it was tempting to speculate that 192 indolmycin production could be QS regulated. Organic extracts of overnight cultures of the QS-activating 193 strains were prepared and loaded on a TLC plate, subsequently overlaid with A. tumefaciens pZLR4 (figure 194 1). Figure 1 shows the presence of at least three putative AHL-like compounds in the culture supernatant of 195 strain S4054, which was chosen for further studies of the potential QS system.

196 Cloning and characterization of a *luxl* homologue from *P. luteoviolacea* S4054. To elucidate the genetics 197 of the putative QS system in *P. luteoviolacea* S4054 a shotgun cloning approach was applied, in which a 198 shotgun library was screened for its ability to induce luminescence in *E. coli* JM109 pSB401 when expressed 199 in the same host. Twenty three luminescent clones were isolated and tested against the *C. violaceum* 200 CV026 monitor strain. Clones NV-5 and NV-17 elicited the strongest response based on the intensity of 201 violacein production in CV026 and in TLC bioassays a pattern similar to the wild type was observed. Hence, 202 the DNA fragment cloned from S4054 encodes a gene which is expressed in *E. coli* DH5- α λ pir and enables 203 synthesis of AHL-like compounds. A restriction pattern analysis was carried out on the insert DNA 204 fragments from positive clones which indicated that NV-5 and NV-17 clones contained the same DNA 205 fragment, with an additional fragment in NV-17 (data not shown). Since NV-5 produced the strongest 206 response in CV026 and contained the shortest DNA insert, this construct was chosen for sequencing based 207 on a primer walking strategy. In the resulting nucleotide sequence the gene *plul* with homology to various 208 luxl genes was identified using BLAST. The 20 best BLASTX hits were used to produce the phylogenetic tree in figure 2, which placed Plul on a monophyletic branch adjacent to LuxI proteins from primarily 209 210 Vibrionaceae, which are known to synthetize AHL molecules (9). Hence, molecular and bioinformatics 211 analyses provided evidence that *plul* encodes a novel LuxI homologue AHL-synthethase involved in the production of AHL-like molecules by P. luteoviolacea S4054. 212

213 Analysis of the genomic context of *pluI*. To facilitate analysis of the genomic context of the *pluI* gene whole 214 genome sequencing was done (33). The genome confirmed the *plul* sequence obtained by cloning and 215 subsequent primer walking. Upstream of *plul* a palindromic sequence resembling a Lux-box was identified. 216 In canonic LuxIR systems the Lux-box is targeted by the LuxR protein for DNA binding suggesting that the 217 downstream genes are QS regulated. *plul* was situated in an operon structure (figure 3) and immediately 218 following the operon, a luxR homologue pluR was identified on the opposite DNA strand. The genes pluI 219 and *pluR* likely form a QS system for regulation of gene expression. The putative operon structure very 220 likely contained genes regulated by this PluIR system in a genomic organization parallel to the V. fischeri 221 *luxI-CDABEG* genes (8). Annotation of the genes contained in the operon structure showed functions such 222 as aminotransferases and a methyltransferase which are included in the biosynthetic pathway of the 223 antibiotic indolmycin (16), however as the genes involved in biosynthesis of indolmycin are not known, 224 further studies are required to determine the actual product of the putative *pluIR* regulated operon. The 225 presence of a tryptophanyl tRNA synthethase in the operon also suggests a link to indolmycin, which is 226 tryptophan derived and selectively targets tryptophanyl tRNA to exert its antibacterial activity (38).

Chemical analysis of AHL-like compounds from *P. luteoviolacea* S4054. The supernatant was analyzed for the presence of AHL-like molecules in the TLC assay with *Agrobacterium* and *Chromobacterium*. None of the signals were present in the blank medium. The strong response in the *A. tumefaciens* pZLR4 monitor strain coupled with a weaker response in *C. violaceum* CV026 suggests that the QS activating molecules share characteristics with 3-oxo substituted AHLs (3). When subjected to alkaline conditions, the synthetic AHL standards and the QS activating compounds isolated from strain S4054 lost the ability to elicit a response in the QS monitor strain *A. tumefaciens* pZLR4, as would be expected from AHLs (42).

234 Ethyl acetate extracts and supernatants of cultures of s4054 were investigated for the presence of AHLs 235 using validated LC-HRMS methods (36). No peaks matched the retention time or accurate mass of 236 commonly found AHL compounds or any analogues reported in literature (19). LC-MS/MS experiments (26) 237 with in-source fragmentation (40 V) were performed in order to confirm the presence of potential novel 238 AHL analogues. Blank medium spiked with AHL standards was used as positive control to ensure that there 239 was no ion suppression. However, the characteristic fragment of 102 Da corresponding to the lactone 240 moiety of the AHLs could not be observed in any of the *P. luteoviolacea* extracts. With the potential of a 241 completely novel QS compound, the metabolite profiles of s4054 and the E. coli clone NV5 were compared 242 as common peaks are likely candidates for the QS compound(s). The *E. coli* clone NV1 containing the 243 pBluescript vector without the *luxl* insert was used as negative control. However, no shared compounds 244 were detected between S4054 and clone NV5, suggesting that the AHL molecules were present in very low amounts or did not perform well in the chromatography. S4054 and NV5 were subjected to preparative 245 246 reverse phase C18 TLC in order to investigate more pure fractions of the QS compound, yet it was not 247 possible to obtain sufficient resolution. Likewise, it was not possible to obtain resolution of the QS activity 248 when subjecting the crude extracts to different solid phase extraction columns (C18, diol, amino) with a 249 finer particle size. As the QS compound might be present in concentrations below the detection limits of 250 our instruments, an 8 L culture was prepared and subjected to cation- and anion-exchange to remove 251 charged compounds. As expected for AHL-like compounds, the activity was recovered from the neutral

252 fraction. Subsequent purification was done on diol and C18 to obtain orthogonal separation of the 253 compounds. This led to four fractions with QS activity. However, it was not possible to deduce an accurate 254 mass from LC-MS analyses. The active fractions were analysed by NMR, which revealed the presence of an 255 NH-proton and a large aliphatic moiety consistent with an AHL; however, the amounts obtained were 256 insufficient for detailed structural analysis. Overall, NMR and chromatographic behavior suggest that we 257 are dealing with an AHL-like molecule, potentially with a modification in the homoserine lactone or with 258 multiple oxo/hydroxyl-groups causing tailing which makes it difficult to detect on columns traditionally 259 used for AHL purification (36).

260

262 Conclusion. P. luteoviolacea S4054 produced molecules that elicited QS responses in the monitor strains 263 C.violaceum CV026, A. tumefaciens pZLR4, and E. coli JM109 pSB401. This, in addition to genomic evidence 264 and indications from chemical analyses, provided strong evidence for the AHL-like nature of these 265 molecules, however we were unable to elucidate the complete structure of the active compounds. The *plul* 266 gene forms a monophyletic cluster in a phylogenetic analysis of *luxI* homologues which indicates that it 267 might represent a novel family of LuxI synthethase homologues. The *pluI* gene was located in an operon 268 structure with a potential biosynthetic pathway downstream of *plul*. This pathway was not related to the 269 violacein pathway, which appears to not be quorum sensing regulated in P. luteoviolacea, but some gene 270 functions corresponded to known biosynthetic steps in the biosynthesis of the potent antibiotic indolmycin. 271 Quorum sensing regulation of antibiotic production is potentially an important modulating factor in 272 microbial communities, and further work is warranted to elucidate the mechanisms of the P. luteoviolacea 273 S4054 QS system and its in situ function.

274

Acknowledgements. This study was supported by the Programme Commision on Health, Food and Welfare
 under the Danish Council for Strategic Research. The present work was carried out as part of the Galathea 3
 expedition under the auspices of the Danish Expedition Foundation. This is Galathea 3 contribution no.
 PXXXXXXX (to be added if/when accepted).



Figure 1: TLC of extract from *P. luteoviolacea* S4054. A: 10 µl ethyl acetate extract from *P. luteoviolacea* overnight culture in MMM. At least three spots are visible in lane A, indicative of three or more AHL-like compounds in the sample. The amounts of synthetic AHLs loaded as standards were: OHHL 10 pM, HHL 3 nM, and OHL 3.8 pM.



Figure 2: Phylogenetic tree of the top 20 BLASTX hits for the *plul* gene. The tree is based on a multiple sequence alignment of translated nucleotide sequences calculated in MEGA5 (31), which was also used to infer the phylogeny using the maximum likelihood method based on the Whelan and Goldman model (39). The rigidity of the tree was tested using 100 bootstrap replications as indicated for nodes supported 70% or higher. Plul forms a monophyletic branch adjacent to LuxI proteins from mostly *Vibrionaceae* and far removed from the *P. atlantica* protein. This supports the hypothesis that PluI represents a novel AHL synthethase. Scale bar: substitutions per site.



Figure 3: Genomic organization of QS related elements in *P. luteoviolacea* S4054 and a putatively QS regulated operon. Predicted functions: Dark blue: *plul*, red: aminotransferase, yellow: dehydrogenase, green: methyltransferase, grey: hypothetical protein, purple: coenzyme F390 synthethase, pink: methylase, brown: lyase, olive: tryptophanyl t-RNA synthethase, light blue: *pluR*. Upstream of *plul* a palindromic sequence reminiscent of a *lux*-box was identified (not shown).

Table 1: Bacterial strains and plasmids used in this study.

Strain or plasmids	Genotype or relevant markers	Source
Pseudoalteromonas luteoviolacea		
DSM 6061 (T)	Type strain, wild type	DSMZ
S2607	Wild type	(13)
S4047-1	Wild type	(13)
S4054	Wild type	(13)
S4060-1	Wild type	(13)
CPMOR-1	Wild type	(12)
CPMOR-2	Wild type	(12)
2ta16	Wild type	(29)
H33	Wild type	T. Harder, unpublished
H33S	Wild type	T. Harder, unpublished
NCIMB 1942	Wild type	NCIMB
NCIMB 1944	Wild type	NCIMB
NCIMB 2035	Wild type	NCIMB
Chromobacterium violaceum CV026 Agrobacterium tumefaciens pZLR4		
Escherichia coli		
DH5α-λ <i>pir</i> JM109	F ⁻ Ф80dlacZ∆M15 ∆(lacZYA-argF U169 deoR supE44 hsdR17 recA1 endA1 gyrA96 thi-1 relA1 recA1 endA1 gyrA96 thi hsdR17 supE44 relA1∆(lac- proAB) mcrA [F'traD36 proAB lacl ⁹ lacZ∆M	(7)
		(·-/
Plasmids		
pBluescript	Ap ^r ; ColE1 origin	Stratagene
pSB401	Tc ^r ; p15A origin	(40)

	Prime	er Sequence 5'-3'		
	Т3	AAT TAA CCC TCA CTA AAG GG		
	T7	GTA ATA CGA CTC ACT ATA GGG C		
	NV5-1	1 TCG GGT GAG AAT ATT		
	NV5-2	2 ACG TAA TCA ATG CCT		
	NV5-3	3 AAA ACA TGG CCA ACC		
	NV5-4	4 TTG AAG AGT TTG CGT		
	NV5-5	5 TAT CGC GGC AAA TCG		
	NV5-6	6 TGT TAC TTA ATG AAG		
	NV5-7	7 CTT CTT GCC AAT AGC		
	NV5-8	8 AGC TCT ATC CCT GAT		
	NV5-9	9 GTA TTG GCG CGT TCA		
	NV5-1	10 CTA CGT TAC GTT TAC		
	NV5-1	11 GCT TCA TTG TAA TAC		
	NV5-1	12 TGC AGT AGA TAA GCA		
	NV5-1	13 CGA GTT TAA AGC CGT		
	NV5-1	14 GTG AGC TGT TCG CAG		
286				
287				
200		D	oforonco Lict	
200		r.		
289				
290	1.	Berger, M., A. Neumann, S. Schulz, M. Sin	non and T. Brinkhoff 2011. Tropodithietic acid production	
291		in Phaeobacter gallaeciensis is regulated by N-Acyl homoserine lactone-mediated quorum sensing. J		
292		Bacteriol 193 :6576-6585.		
202	2	Buchan A. I. M. Gonz±ílez and M. A. Mo	ran 2005. Overview of the marine Recephacter lineage	
293	۷.	Buchan, A., J. M. Gonz-filez and M. A. Moran 2005. Overview of the marine Roseobacter lineage.		
234				
295	3.	Cha, C., P. Gao, Y. C. Chen, P. D. Shaw and	S. K. Farrand 1998. Production of acyl-homoserine lactone	
296		quorum-sensing signals by gram-negative plant-associated bacteria. Mol. Plant Microbe In. 11 :1119-		
297		1129.		
200			territe and the state of the state of the medicate back to be a long to be	
298	4.	Clark, D. J. and O. Maaløe 1967. DNA repl	ication and the division cycle in <i>Escherichia coli</i> . J. Mol. Biol.	
299		23 :99-112.		
300	5.	Croxatto, A., V. J. Chalker, J. Lauritz, J. Jas	s, A. Hardman, P. Williams, M. Cámara and D. L. Milton	
301		2002. VanT, a homologue of Vibrio harvey	<i>i</i> LuxR, regulates serine, metalloprotease, pigment, and	
302		biofilm production in Vibrio anguillarum. J	. Bacteriol. 184 :1617-1629.	
	-			
303	6.	Duerkop, B. A., J. Varga, J. R. Chandler, S.	B. Peterson, J. P. Herman, M. E. A. Churchill, M. R. Parsek,	
304		w. c. Nierman and E. P. Greenberg 2009.	Quorum-sensing control of antibiotic synthesis in	
305		Burknolaeria thallandensis. J. Bacteriol. 19	1.3909.	

- 306
 7. Dunn, A. K., M. O. Martin and E. V. Stabb 2005. Characterization of pES213, a small mobilizable
 307 plasmid from *Vibrio fischeri*. Plasmid 54:114-134.
- Engebrecht, J., K. Nealson and M. Silverman 1983. Bacterial bioluminescence: isolation and genetic
 analysis of functions from *Vibrio fischeri*. Cell **32**:773-781.
- Engebrecht, J. A. and M. Silverman 1987. Nucleotide sequence of the regulatory locus controlling
 expression of bacterial genes for bioluminescence. Nucleic Acids Research 15:10455-10467.
- Fuqua, C., M. R. Parsek and E. P. Greenberg 2001. Regulation of gene expression by cell-to-cell
 communication: acyl-homoserine lactone quorum sensing. Annu. Rev. Genet. 35:439-468.
- Galloway, W. R., J. T. Hodgkinson, S. D. Bowden, M. Welch and D. R. Spring 2011. Quorum sensing in Gram-negative bacteria: small-molecule modulation of AHL and Al-2 quorum sensing pathways.
 Chem Rev 111:28-67.
- Gomez, D., E. Espinosa, M. Bertazzo, P. Lucas-Elio, F. Solano and A. Sanchez-Amat 2008. The
 macromolecule with antimicrobial activity synthesized by *Pseudoalteromonas luteoviolacea* strains is
 an L-amino acid oxidase. Appl. Environ. Microb. **79**:925-930.
- Gram, L., J. Melchiorsen and J. B. Bruhn 2010. Antibacterial activity of marine culturable bacteria
 collected from a global sampling of ocean surface waters and surface swabs of marine organisms.
 Mar Biotechnol 12:439-451.
- Hibbing, M. E., C. Fuqua, M. R. Parsek and S. B. Peterson 2009. Bacterial competition: surviving and
 thriving in the microbial jungle. Nat. Rev. Microbiol. 8:15-25.
- Holden, M. T. G., S. Ram Chhabra, R. De Nys, P. Stead, N. J. Bainton, P. J. Hill, M. Manefield, N.
 Kumar, M. Labatte and D. England 1999. Quorum–sensing cross talk: isolation and chemical
 characterization of cyclic dipeptides from *Pseudomonas aeruginosa* and other Gram-negative
 bacteria. Molecular Microbiology 33:1254-1266.
- Hornemann, U., L. H. Hurley, M. K. Speedie and H. G. Floss 1971. Biosynthesis of indolmycin. J. Am.
 Chem. Soc. 93:3028-3035.
- Huang, Y. L., J. S. Ki, R. J. Case and P. Y. Qian 2008. Diversity and acyl-homoserine lactone production
 among subtidal biofilm-forming bacteria. Aquat. Microb. Ecol. 52:185-193.
- Huang, Y. L., J. S. Ki, O. O. Lee and P. Y. Qian 2008. Evidence for the dynamics of Acyl homoserine
 lactone and AHL-producing bacteria during subtidal biofilm formation. The ISME journal 3:296-304.
- Laatsch, H. 2011. Antibase 2011. <u>http://www</u>. wiley-vch. de/stmdata/antibase2011. php . Wiley-VCH:
 Weinheim, Germany.
- Månsson, M., R. K. Phipps, L. Gram, M. H. Munro, T. O. Larsen and K. F. Nielsen 2010. Explorative
 Solid-Phase Extraction (E-SPE) for Accelerated Microbial Natural Product Discovery. J. Nat. Prod.
 73:1126-1132.

```
    Markowitz, V. M., K. Mavromatis, N. N. Ivanova, I. Chen, A. Min, K. Chu and N. C. Kyrpides 2009.
    IMG ER: a system for microbial genome annotation expert review and curation. Bioinformatics
    25:2271.
```

- Matz, C., P. Deines, J. Boenigk, H. Arndt, L. Eberl, S. Kjelleberg and K. Jurgens 2004. Impact of
 violacein-producing bacteria on survival and feeding of bacterivorous nanoflagellates. Appl. Environ.
 Microb. 70:1593.
- McClean, K. H., M. K. Winson, L. Fish, A. Taylor, S. R. Chhabra, M. Camara, M. Daykin, J. H. Lamb, S.
 Swift and B. W. Bycroft 1997. Quorum sensing and *Chromobacterium violaceum*: exploitation of
 violacein production and inhibition for the detection of N-acylhomoserine lactones. Microbiol-SGM
 143:3703.
- 350 24. McGee, K., P. Hörstedt and D. L. Milton 1996. Identification and characterization of additional
 351 flagellin genes from *Vibrio anguillarum*. Journal of Bacteriology **178**:5188-5198.
- 352 25. Miller, M. B. and B. L. Bassler 2001. Quorum sensing in bacteria. Annu. Rev. Microbiol. 55:165-199.
- 353 26. Ortori, C. A., J. F. Dubern, S. R. Chhabra, M. C+ímara, K. Hardie, P. Williams and D. A. Barrett 2011.
 354 Simultaneous quantitative profiling of N-acyl-l-homoserine lactone and 2-alkyl-4 (1H)-quinolone
 355 families of quorum-sensing signaling molecules using LC-MS/MS. Analytical and Bioanalytical
 356 Chemistry :1-12.
- 357 27. Östling, J., A. E. Goodman and S. Kjelleberg 1991. Behaviour of IncP-1 plasmids and a miniMu
 358 transposon in a marine *Vibrio* sp. S14.: isolation of starvation inducible *lac* operon fusions. FEMS
 359 Microbiol. Ecol 86:83-94.
- Ravn, L., A. B. Christensen, S. Molin, M. Givskov and L. Gram 2001. Methods for detecting acylated
 homoserine lactones produced by Gram-negative bacteria and their application in studies of AHL production kinetics. J. Microbiol. Meth. 44:239-251.
- Rypien, K. L., J. R. Ward and F. Azam 2010. Antagonistic interactions among coral-associated
 bacteria. Environ. Microbiol. 12:28-39.
- 365 30. Sambrook, J. and D. W. Russell. 2001. Molecular cloning: a laboratory manual. 2. CSHL press.
- 366 31. Tamura, K., D. Peterson, N. Peterson, G. Stecher, M. Nei and S. Kumar 2011. MEGA5: molecular
 acount of the second state of the secon
- 369 32. Thomson, N. R., M. A. Crow, S. J. McGowan, A. Cox and G. P. C. Salmond 2000. Biosynthesis of
 370 carbapenem antibiotic and prodigiosin pigment in *Serratia* is under quorum sensing control. Mol.
 371 Microbiol. 36:539-556.
- 372 33. Vynne, N. G. and L. Gram 2012. Draft genome sequence of *Pseudoalteromonas luteoviolacea* S4054,
 373 an indolmycin producing marine bacterium. In preparation for Journal of Bacteriology
- 374 34. Vynne, N. G., M. Månsson, K. F. Nielsen and L. Gram 2012. Antibiotic chemoprofile and RecA and
 375 GyrB sequence analyses reveal three phylotypes within the marine *Pseudoalteromonas luteoviolacea*.
 376 In preparation
- 35. Vynne, N. G., M. Månsson, K. F. Nielsen and L. Gram 2011. Bioactivity, chemical profiling and 16S
 rRNA based phylogeny of *Pseudoalteromonas* strains collected on a global research cruise. Mar.
 Biotechnol. doi:10.1007/s10126-011-9369-4.

- Wang, J., C. Quan, X. Wang, P. Zhao and S. Fan 2011. Extraction, purification and identification of
 bacterial signal molecules based on N–acyl homoserine lactones. Microbial Biotechnology 4:479-490.
- Wang, Y., A. Ikawa, S. Okaue, S. Taniguchi, I. Osaka, A. Yoshimoto, Y. Kishida, R. Arakawa and K.
 Enomoto 2008. Quorum sensing signaling molecules involved in the production of violacein by
 Pseudoalteromonas. Biosci. Biotechnol. Biochem. **72**:1958-1961.
- 38. Werner, R. G., L. F. Thorpe, W. Reuter and K. H. Nierhaus 1976. Indolmycin Inhibits prokaryotic
 tryptophanyl-tRNA ligase. Eur. J. Biochem. 68:1-3.
- 387 39. Whelan, S. and N. Goldman 2001. A general empirical model of protein evolution derived from
 388 multiple protein families using a maximum-likelihood approach. Molecular Biology and Evolution
 389 18:691-699.
- Winson, M. K., S. Swift, L. Fish, J. P. Throup, F. Jørgensen, S. R. Chhabra, B. W. Bycroft, P. Williams
 and G. S. A. B. Stewart 1998. Construction and analysis of *luxCDABE*—based plasmid sensors for
 investigating N—acyl homoserine lactone-mediated quorum sensing. FEMS Microbiol. Lett. 163:185 192.
- Yanisch-Perron, C., J. Vieira and J. Messing 1985. Improved M13 phage cloning vectors and host
 strains: nucleotide sequences of the M13mpl8 and pUC19 vectors. Gene 33:103-119.
- Yates, E. A., B. Philipp, C. Buckley, S. Atkinson, S. R. Chhabra, R. E. Sockett, M. Goldner, Y. Dessaux,
 M. Camara and H. Smith 2002. N-acylhomoserine lactones undergo lactonolysis in a pH-,
 temperature-, and acyl chain length-dependent manner during growth of Yersinia pseudotuberculosis
 and Pseudomonas aeruginosa. Infect. Immun. **70**:5635.

Vynne N, Gram L (2012)

Draft genome sequence of *Pseudoalteromonas luteoviolacea* S4054, an indolmycin producing marine bacterium

Manuscript in preparation.

- 1 Draft genome sequence of *Pseudoalteromonas luteoviolacea* S4054, an indolmycin producing
- 2 marine bacterium
- 3

4 Nikolaj G. Vynne¹* and Lone Gram¹

⁵ ¹ Technical University of Denmark, National Food Institute, Søltofts Plads, bldg. 221, DK-2800 Kgs.

- 6 Lyngby, Denmark
- 7
- 8 * corresponding author:
- 9 phone: +45 4525 2571
- 10 fax: +45 4588 4774
- 11 e-mail: ngvy@food.dtu.dk
- 12
- 13 **Abstract:** The ability to produce antibacterial compounds is wide spread among members of the
- 14 genus *Pseudoalteromonas*. Here we present the 6.08-Mb draft genome sequence of *P. luteoviolacea*
- 15 S4054, which produces the antibiotics violacein and indolmycin, and investigate the potential for
- 16 further discoveries of bioactive compounds.
- 17
- 18 Marine bacteria are widely investigated as a potential source of novel molecules with
- 19 pharmacologically relevant activites. Several species of the genus *Pseudoalteromonas* produce
- 20 antibacterial, antifouling or cytotoxic compounds (9) (1), however the ecological role of these
- compounds is not yet fully understood. Often, the bioactive *Pseudoalteromonas* have been isolated
- 22 from the surfaces of marine eukaryotes and their bioactivity may be part of a symbiotic relationship
- 23 (2). *P. luteoviolacea* S4054 was isolated from a seaweed sample in the Pacific Ocean and it produces
- the antibacterial compounds violacein and indolmycin (9) (7). Violacein is produced by different
- 25 bacterial species, while to our knowledge indolmycin has not previously been isolated from a marine
- 26 bacterium. The biosynthetic pathways responsible for production of these compounds are candidates
- 27 for horizontal gene transfer events in S4054, however, only the violacein pathway is described on a
- 28 genetic level, enabling identification through homology searches.
- 29

Genomic DNA was obtained by successive phenol-chloroform purification steps. Mated paired-end
library preparation, Illumina Hi Seq 2000 sequencing runs and initial data processing were done by
BGI (Beijing, China). After quality filtering and trimming, the short reads were assembled and gaps
filled using SOAPdenovo (6). This resulted in 126 contigs at a depth of 30.8-fold genome coverage
and a total length of 6,076,381 bp. The draft genome was uploaded to the IMG/ER pipeline for gene

- annotation. A total of 5,268 genes were annotated, 5,121 of which were protein encoding. One
- 36 hundred and twenty six genes fell into COG categories in secondary metabolism, with 30 genes in
- 37 COG1020 (non-ribosomal peptide synthase modules and related proteins) and 20 genes in COG1028
- 38 (dehydrogenases). Ten genes encoded proteins involved in PKS biosynthesis. BLASTP homology
- 39 searches identified 5 genes encoding proteins VioA, B. C, D and E which form the violacein
- 40 biosynthetic pathway (5). Upstream of the VioA-E genes, a gene encoding a multi-antimicrobial and

1	toxic	compound extrusion pump was located, similar to the organization found in the	
2	Pseu	doalteromonas tunicata D2 genome (8). In addition to the violacein pathway, six biosynthetic	
3	pathv	ways were identified containing clusters of potential polyketide synthase or nonribosomal peptide	
4	synth	esis coding genes. Genes encoding surface attachment related functions such as curli, type IV	
5	pili ar	nd P pili were identified, suggesting a surface associated lifestyle. The genome contained a	
6	necro	osis inducing protein NPP1 homologue indicating a potential for plant pathogenicity (3). In	
7	additi	ion, the genome had potential for quorum sensing regulated processes based on the presence of	
8	one p	pair of <i>luxI - luxR</i> genes (4).	
9			
10	The o	data presented here provides further insight on the genetic potential for production of bioactive	
11	secondary metabolites within the pseudoalteromonads, and we believe it may advance studies on the		
12	ecolo	ogy of these organisms.	
13			
14	Nucleotide accession number The draft genome sequence for P. luteoviolacea S4054 is deposited		
15	in GenBank under the accession number XXXXXXXXXXXXXXXX.		
16			
17	Ackn	owledgements: This work was funded by the Programme Commission on Health, Food and	
18	Welfare under the Danish Council for Strategic Research. The present work was carried out as part of		
19	the G	alathea 3 expedition under the auspices of the Danish Expedition Foundation.	
20			
21		Reference List	
22			
23	1.	Bowman, J. P. 2007. Bioactive compound synthetic capacity and ecological significance of	
24		marine bacterial genus <i>Pseudoalteromonas</i> . Mar. Drugs 5 :220-241.	
25	2.	Egan, S., T. Thomas, and S. Kjelleberg. 2008. Unlocking the diversity and biotechnological	
26		potential of marine surface associated microbial communities. Curr. Opin. Microbiol. 11 :219-	
27		225.	
28	3.	Fellbrich, G., A. Romanski, A. Varet, B. Blume, F. Brunner, S. Engelhardt, G. Felix, B.	
29		Kemmerling, M. Krzymowska, and T. Nürnberger. 2002. NPP1, a Phytophthora-associated	
30		trigger of plant defense in parsley and <i>Arabidopsis</i> . Plant J. 32 :375-390.	
31	4.	Fuqua, C., S. C. Winans, and E. P. Greenberg. 1996. Census and consensus in bacterial	
32		ecosystems: the LuxR-LuxI family of quorum-sensing transcriptional regulators. Annu. Rev.	
33		Microbiol. 50 :727-751.	
3/			
54	5.	Hoshino, T. 2011. Violacein and related tryptophan metabolites produced by	
35	5.	Hoshino, T. 2011. Violacein and related tryptophan metabolites produced by <i>Chromobacterium violaceum</i> : biosynthetic mechanism and pathway for construction of	
35 36	5.	Hoshino, T. 2011. Violacein and related tryptophan metabolites produced by <i>Chromobacterium violaceum</i> : biosynthetic mechanism and pathway for construction of violacein core. Appl. Microbiol. Biotechnol. 91 :1463-1475.	
35 36 37	5. 6.	 Hoshino, T. 2011. Violacein and related tryptophan metabolites produced by <i>Chromobacterium violaceum</i>: biosynthetic mechanism and pathway for construction of violacein core. Appl. Microbiol. Biotechnol. 91:1463-1475. Li, R., Y. Li, K. Kristiansen, and J. Wang. 2008. SOAP: short oligonucleotide alignment program. 	

- Månsson, M., R. K. Phipps, L. Gram, M. H. Munro, T. O. Larsen, and K. F. Nielsen. 2010.
 Explorative Solid-Phase Extraction (E-SPE) for Accelerated Microbial Natural Product Discovery.
 J. Nat. Prod. 73:1126-1132.
- Thomas, T., F. F. Evans, D. Schleheck, A. Mai-Prochnow, C. Burke, A. Penesyan, D. S. Dalisay,
 S. Stelzer-Braid, N. Saunders, J. Johnson, S. Ferriera, S. Kjelleberg, and S. Egan. 2008. Analysis
 of the Pseudoalteromonas tunicata Genome Reveals Properties of a Surface-Associated Life
 Style in the Marine Environment. PLoS ONE 3.
- 9. Vynne, N. G., M. Månsson, K. F. Nielsen, and L. Gram. 2011. Bioactivity, chemical profiling and
 16S rRNA based phylogeny of *Pseudoalteromonas* strains collected on a global research cruise.
 Mar. Biotechnol. doi:10.1007/s10126-011-9369-4.
- 11
- 12