

Mucosal immune response in common carp (Cyprinus carpio L.)

Host pathogen interactions in relation to beta-glucan stimulation

Przybylska, Dominika Alicja

Publication date: 2012

Document Version Publisher's PDF, also known as Version of record

Link back to DTU Orbit

Citation (APA): Przybylska, D. A. (2012). Mucosal immune response in common carp (Cyprinus carpio L.): Host pathogen interactions in relation to beta-glucan stimulation. Technical University of Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- · You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Mucosal immune response in common carp (Cyprinus carpio L.) Host pathogen interactions in relation to β -glucan stimulation

Dominika A. Przybylska PhD Thesis 2012 Mucosal immune response in common carp (*Cyprinus carpio* L.)host pathogen interactions in relation to β-glucan stimulation

Dominika Alicja Przybylska

PhD Thesis

Mucosal immune response in common carp (*Cyprinus carpio* L.) host pathogen interactions in relation to β -glucan stimulation

Dominika Alicja Przybylska

National Food Institute

Biological Quality Research Group

Technical University of Denmark

2012

The thesis submitted for the degree of Doctor of Philosophy is based on a literature review and 3 scientific publications

Contents

Summary

Sammendrag

1.	General introduction	10		
	1.1. Immunity in fish	11		
	1.2. Mucosal immune system	14		
	1.2.1. Mucosal immunity in fish	14		
	1.2.2.1. Mucus	15		
	1.2.2.2. Skin	16		
	1.2.2.3. Gill	16		
	1.2.2.4. Gut	17		
	1.3. Wound healing process			
	1.3.1. The inflammatory response following the tissue damage	19		
	1.3.2. Cells involved in the inflammatory response	20		
	1.3.2.1 Neutrophils	21		
	1.3.2.2 Macrophages			
	1.3.2.3 Fibroblasts	23		
	1.3.2.4 Mucous cells	24		
	1.3.3. Molecules involved in the inflammatory response	24		
	1.3.3.1. Interleukin 1β (IL-1β)	25		
	1.3.3.2. Interleukin 6 (IL-6)			
	1.3.3.3. Interleukin 8 (IL-8)			
	1.3.3.4. Heat shock protein 70 (Hsp70)			
	1.3.4. Pathogen associated molecular patterns (PAMPs) and damage associated molecular patterns (DAMPs)	27		
	1.3.4.1 PAMPs stimulation	27		
	1.3.4.2 DAMPs stimulation			
	1.4 Stress reaction	31		
	1.4.1 Teleost hypothalamo-pituitary-interrenal (HPI) axis			
	1.4.2 Glucocorticoids in fish			
	1.5 Antibody response			
	1.5.1 IgM			
	1.5.2 IgZ1	34		

	1.6 β-glucans as immune modulators		
	1.6.1	Diversity and structure	35
	1.6.2	Administration routes and its effect on teleost immunity	37
	1.7 Methods applied in the study		
	1.7.1	The damage model	
	1.7.2	Intravenous injection	40
	1.7.3	Aeromonas hydrophila	41
	1.8 Objec	43	
2	General discussion		44
	2.1 The best use of immune modulators: bath, injection or fed?		44
	2.2 The u	45	
	2.3 Inflammatory response during wound healing process: DAMPs and PAMPs stimulation		46
	2.4 β-glu	48	
	2.5 Skin	– perfect tissue to follow β-glucan actions?	49
3.	Conclusions Perspective for future research Acknowledgements		
4.			
5.			
6.	Reference		
7.	Accompanying papers		

"Jeśli nie możesz ich przekonać, zamąć im w głowach"

"If you can't convince them, confuse them"

Harry Truman

Summary

Control of fish diseases is a great concern in aquaculture because of losses in the production. Drug choices for the treatment of common infectious diseases are becoming increasingly limited and expensive and, in some cases, unavailable due to the emergence of drug resistance in bacteria and fungi. This is why number of biological compounds, as an alternative to the drugs, has been used to reduce the risk of diseases and improve fish welfare by enhancement of non-specific defence system. Among them, β -glucans, naturally occurring polysaccharides found in the cell wall of plants, bacteria and fungi, are some of the most powerful and extensively investigated immune modulants. β -glucans have been proven to enhance the immune system and pharmacologically they are classified as biological response modifiers (BRM).

The focus of the present thesis was on: 1. creation of a model for the examination of the biological impact of two commercially available β -glucan enriched products on the wound healing process in common carp (*Cyprinus carpio* L.) in sterile, controlled conditions; 2. investigation of potential impact of intravenously injected β -glucan on mucosal immune response and immunoglobulin switch-like process in common carp. In order to reach these objectives, different methods were used such as real-time quantitative PCR (RT-PCR) in order to measure the expression of immune-related genes involved in wound healing process, ELISA for specific antibody detection, cortisol assay for measurement of stress level, respiratory burst assay for radical production measurement and image analysis.

The results of this study showed that previous infections gave rise to changes regarding texture quality parameters in fresh fish meat, and were a starting point for use of immune modulators such as β -glucans. Further work showed that bath in two commercially available β -glucan enriched products, specifically MacroGard and 6.3 kDa oat fiber, had a direct positive effects on the wound closure in common carp and promoted faster wound healing compared to non-treated fish. We showed the immunological and regenerative response following stimulation with PAMPs and DAMPs in controlled conditions, without the exposure to pathogens, which resulted in an inflammatory response by activating IL-1b, IL-6 and IL-8. Local differences in expression pattern dependent on stimulation by DAMPs alone or DAMPs/PAMPs combination. In addition, the absence of marked differences on the respiratory burst activity in head-kidney cells supports the idea of a localized immune response to the site of injury. Due to direct and constant contact between skin and β -glucan, bath treatment was an obvious choice to investigate. However, intravenous injection of β -glucan showed it has a biological effect on skin as well. Once again we observed immunological changes taking place in skin of common carp, with IgM/IgZ

immunoglobulin switch-like process. Finally, β -glucan seems to work in dose-dependent manner, with elevating cortisol level when injected at high doses.

Sammendrag

For at sikre en sund og rentabel i akvakultur produktion er kontrol af fiskesygdomme af stor betydning. Muligheden for brug af medicin til behandling af almindelige infektiøse sygdomme bliver mere og mere begrænset og kostbare, og er i nogle tilfælde ikke mulig på grund af resistens i bakterier og svampe. Derfor er der stor fokus på alternativer til medicinering, som kan reducere risikoen for sygdomme og forbedre fiskevelfærden gennem stimulering af det ikke-specifikke forsvarssystem. β-glucaner er naturligt forekommende polysaccharider, der findes i cellevæggen hos planter, bakterier og svampe. β-glucaner er nogle af de mest effektive og bedst undersøgte stoffer med immunmodulerende egenskaber i fisk. β-glucaner har vist sig at forbedre immunsystemet og farmakologisk klassificeres de som biologiske respons-modifikatiorer. Fokus i denne afhandling er: 1. Udvikling af en model til undersøgelse af den biologiske virkning af to kommercielt tilgængelige β-glucan berigede produkter på sårhelingsprocessen i almindelig karpe (Cyprinus carpio L.) under sterile, kontrollerede betingelser; 2. Undersøgelse af betydningen af intravenøst injiceret β-glucan på det mukosale-immunrespons og en antistof switch-lignende proces i karper. For at nå disse mål, blev der anvendt forskellige metoder, såsom kvantitativ realtime-PCR (RT-PCR) til måling af ekspressionen af immun-relaterede gener involveret i sårhelingsprocessen, ELISA for bestemmelse af specifikke antistoftitere samt cortisol-måling af stressniveau, real-time respiratorisk burst assay til bestemmelse af produktionen af frie-radikaler og billedanalyse af sårhelingsprocessen. Det er blevet vist at behandlede sygdomsudbrud giver anledning til ændringer i kvalitetsparametre som tekstur i frisk fiskekød. Hvilket har været udgangspunktet for yderligere at undersøge anvendelses af immunmodulatoren β-glucaner. Den videre undersøgelse af effekter af β -glucan har vist, at to kommercielt tilgængelige β -glucan berigede produkter, MacroGard og 6,3 kDa havrefibre, har en direkte positiv effekt på sårlukning i karper ved badbehandling med disse. Således var sårheling hurtigere i β-glucan behandlede fisk i sammenligning med ikke-behandlede fisk. Det immunologiske og regenerative respons efter stimulering med PAMPs (Pathogenassociated molecular patterns) og DAMPs (danger-associated molecular patterns molecules) under kontrollerede forhold, uden at eksponeringen for patogener er ligeledes blevet undersøgt, og en inflammatorisk reaktion ved aktivering af IL-1b, IL-6 og IL-8 blev fundet. Herudover blev der set lokale forskelle i ekspressionsmønsteret afhængige af om stimuleringen var med DAMPS alene eller DAMPS / PAMPs kombination. Desuden understøtter fraværet af væsentlige forskelle i den respiratorisk burst aktivitet i hoved-nyreceller ideen om at det lokale respons på et vævsbeskadigelsen er det primære innate respons. Yderligere blev der fundet immunologiske effekter af intravenøs injektion af β-glucan i huden. Således blev der observeret immunologiske forandringer i huden som tyder på en immunoglobulin switch-lignende proces med IgM / IgZ.

Endvidere blev det fundet at iv. injektion med β -glucan har en dosisafhængig effekt på stress niveauet i karper udtrykt som forøgelse af plasma cortisol niveau.

1. General introduction

The common carp (*Cyprinus carpio* L.) is probably the first fish species whose distribution was widely extended by human introduction, since the Romans introduced it from the River Danube throughout Europe (Balon, 1995). It is the third most frequently introduced species worldwide (Welcomme, 1992). The common carp also accounts for the world's second highest farmed fish production, mainly from polyculture in Asia (Milstein, 1992), and the production of ornamental varieties is even more important in monetary value (Balon, 1995). A change in the main objective of common carp production can be observed in Europe (FAO report 2004-2012; FAO. © 2004-2012. Cultured Aquatic Species Information Programme. Cyprinus carpio. Cultured Aquatic Species Information Programme). Since this species has outstanding importance in freshwater aquaculture, many aspects of its physiology, nutrition, genetics, and diseases have been studied during past decades. Adverse changes in the natural environment, the increasing intensity of carp production in many areas, extensive inter-regional transport of common carp and other cyprinids, and the ban on using several traditional medicaments (fungicides, antibiotics and insecticides) call for the intensification of research on carp diseases. The development of vaccines and immune modulators, to increase the natural resistance of fish to pathogens, seems to be the most promising solution for avoiding the application of antibiotics (Rodgers and Furones, 2009). Nowadays, number of biological compounds has been used to reduce the risk of diseases and improve fish welfare by enhancement of non-specific defence system. Among them, β -glucans are some of the most powerful and extensively investigated immune modulants. They are naturally occurring polysaccharides from cereals (e.g. barley, oat), algae, mushrooms and yeast cell walls. Glucan exerts a beneficial effect on a variety of experimental disease states of bacterial, viral, fungal and parasitic origin (DiLuzio and Williams, 1978; Williams et al., 1983, Williams and DiLuzio, 1980, 1985, Williams et al., 1978, Cook et al., 1980; Yano et al., 1991, Raa et al., 1992, Duncan and Klesius 1996, Rodriguez et al. 2009, Zhang et al. 2009). Many studies investigated the effect of beta glucans on fish immunity (Dalmo et al., 1997; Palic et al., 2006; 2007; Dalmo and Bøgwald, 2008; Yano et al., 1991; Raa et al., 1992; Duncan and Klesius 1996; Rodríguez et al., 2009; Zhang et al., 2009).

The overall theme of the PhD study is to investigate and understand the impact of the β -glucan as immune modulator in relation to the mucosal immune response in common carp (*Cyprinus carpio* L.). By introducing immune modulator from "outside" and "inside" we investigated parameters of innate immune response of common carp, such as pro-inflammatory cytokines expression, respiratory burst activity, antibody ratio and cortisol level. Two models were applied: bath model and examination of biological effect of β -glucan on wound healing process in carp, and intravenous

injection model, to investigate natural and specific antibody ratio and dose-related side effects of β -glucan.

1.1 Immunity in fish

Immunity system exists in animals (vertebrates and invertebrates) and plants (Ausubel 2005) to protect themselves against variety of microorganisms (Janeway and Medzitov 2002). Innate immunity is non-specific and fast-reacting defense mechanism, which includes physical parameters, cellular and humoral factors (Magnadóttir, 2004). Pathogens can be recognized by the adaptive immune system, a highly specific and diverse system characterized by memory occurrence.

Fish are the earliest vertebrates that have developed both arms of the immune system, reflected by the innate and the adaptive immune response (Figure 1). Mucosal immune system in skin, gill and gut forms the first line of defense and the connective tissues associated are densely populated with immune cells. Fish lack bone marrow as a primary immune organ, and their hematopoietic cells are produced in so-called head kidney (HK) where myeloid and lymphoid immune cells raise (van Muiswinkel, 1995). The capacity of the head-kidney to trap antigens and produce antibodies has been demonstrated in a few species (Ellis et al., 1989; Zapata and Cooper, 1990). Moreover head kidney combines immune functions with corticosteroid production and is an intriguing link between the endocrine and immune system (Manning, 1994). The HK together with the thymus, centre of T-lymphocytes maturation, are sites for the interaction of immune system with antigens (Tort et al., 2003). Fish lack lymph nodes as secondary immune organs (Tort et al., 2003). Blood filtration and erythrocytic destruction is performed by the melanomacrophagic centres, formed by accumulation of macrophages associated to elipsoid capillaries. These centers may retain antigens as immune complexes for long periods (Espenes et al., 1995).

Although immune system of teleost fish is different from that of mammals, there are some common features. Immune cells communicate via cytokines and associated receptors, which is important for a fast and effective attack on pathogens (Bird et al., 2006). Many genes encoding molecules associated with defense mechanism of fish have counter parts in mammals, suggesting a common ancestor of key mediators of immune system (Plouffe et al., 2005). In vertebrates, cytokines are the key regulators of both, innate and adaptive immunity and elicit broad range of actions on cell growth and differentiation (Bird et al, 2006). In general, the cytokines repertoire in fish is alike in mammals, however function of each fish cytokine is not as good described as in mammals. However, in fish many genes are duplicated and it results in the creation of numerous novel or

semi-novel genes and functions in fish, known as "more genes in fish than mammals" concept (Ohno 1970). It is therefore not surprising that there are already 5 genomic databases sequenced thus far for this taxon; the zebrafish, medaka, stickleback, tiger pufferfish and the green spotted pufferfish (www.ensembl.org). Large scale of teleost fish genome analysis has revealed numerous gene duplications that are thought to originate from the WGD (Stein 2007).

Tremendous progress has been made in isolating these molecules from fish in recent years, (ILs) such as IL-1 β (Zou et al., 1999; Bird et al., 2002; Seppola et al., 2007), IL-6 (Bird et al., 2005b; Costa et al., 2011), IL-8 (Seppola et al., 2007; Oehlers et al., 2009; Qiu et al., 2009.); tumor necrosis factor alpha (TNF- α) (Hirono et al., 2000; Secombs et al., 2001) and interferons e.g. interferon gamma (IFN- γ) (Zou et al., 2004), and many more are predicted to be found. Genetic diversity translates to protein diversity and as such it is therefore very possible that in teleost fish, there will be a lot of unique and differing functionalities amidst the background of conserved functions. In fact, many of these fish-specific features are now starting to be unraveled (Aoki et al., 2008).

The adaptive immune response provides the vertebrate immune system with the ability to recognize and remember specific pathogens, and to mount stronger and faster response to pathogens. To mount such an immune response, a specific pathogen or antigen needs to be recognized (Aoki et al., 2008). In teleostean fishes, as in mammals, a first antigen encounter will induce some longlived memory cells that retain the capacity to be stimulated by the antigen, which results in faster and more vigorous secondary response (dos Santos et al., 2001; Kaattari et al., 2002). The essential features of adaptive immunity are immunoglobulins and the major histocompatibility complex (MHC). The first evidence of for MHC genes in teleost fish was published by Hashimoto and coworkers in 1990. Today, MHC genes have been isolated and described for all major vertebrates, including cartilaginous fish (Barl, 1998; Ohta et al., 2000) and bony fish (Kruiswijk et al., 2002; Stet et al., 1998). The teleost fish MHC is equivalent to its mammalian counterpart, but the class I and class II genes appear in different linkage group and are not clustered (Stet et al., 2003). Mammals have five types of immunoglobulin (IgA, IgD, IgG, IgM and IgE; Elgert KD 1998; from Immunology: Understanding the immune system, Chapter 4). The main immunoglobulin studied in in teleost is tetrameric IgM (Håvarstein et al., 1998; Kaattari et al., 1998) and additionally, IgD and IgT/IgZ have been recently described, but functional implications have not yet been established (Flajnik, 2002; Danilova et al., 2005; Savan et al., 2005a; Savan et al., 2005b; Hansen et al., 2005; Zhang et al. 2010). Even so, IgZ or IgT were not sequenced in all teleost species (Bengten et al., 2006). IgG is not found in fish and no class switch seems to be possible (Magor et al., 1999). However, Przybylska and Nielsen (PAPER 3) have showed, that dose-related injection with β-

glucan cause immunoglobulin switch-like process in skin of common carp, where presence of main fish immunoglobulin IgM is overtaken by recently described, mucosal immunoglobulin IgZ1.

Common carp (Cyprinus carpio L.) is a close relative to zebrafish (*Danio rerio*), and they are representatives of the most abundant and widespread primary freshwater fish family, Cyprinidae (FishBase, <u>http://www.fishbase.org/</u>). In the current PhD thesis, some references will be made to zebrafish due to ample genomic resources including a nearly fully sequenced genome and over a million expressed sequence tags, while genomic data for the rest of the cyprinids are quite scarce (Orbam and Wu, 2006).



Figure 1. Mammalian model: illustration of cells involved in innate and adaptive immune response (Dranoff, 2004). The innate immune response functions as the first line of defense against infection. It consists of soluble factors, such as complement proteins, and diverse cellular components including granulocytes (basophils, eosinophils and neutrophils), mast cells, macrophages, dendritic cells and natural killer cells. The adaptive immune response is slower to develop, but manifests as increased antigenic specificity and memory. It consists of antibodies, B cells, and CD4+ and CD8+ T lymphocytes. Natural killer T cells and T cells are cytotoxic lymphocytes that straddle the interface of innate and adaptive immunity. Most of the cells involved in mammalian immunity are present in fish, and they are myeloid cells, neutrophilic granulocytes and macrophages. Pathogens are killed by natural cytotoxic cells, the fish equivalent of mammalian natural cells. Further, fish possess mast cells, however they do not have histamine granules (Mulero et al., 2007). In the present PhD thesis, both, innate (**PAPER 2**) and adaptive immunity (**PAPER 3**) have been examinated.

1.2 Mucosal immune system

The mucosal immune system in vertebrates is a part of the immune system that responds to and protects against pathogens that enter the body through mucosal surfaces. The mucosal membranes are endowed with powerful mechanical and chemical cleansing mechanisms that degrade and repel most foreign matter and, in addition, a large and highly specialised innate and adaptive mucosal immune system protects these surfaces (Holmgren and Czerkinsky, 2005). The mammalian mucosal immune system is composed of inductive sites, where antigens sampled from mucosal surfaces stimulate cognate naïve T and B lymphocytes, and effector sites, where the effector cells after extravasation and differentiation get to work (Brandtzaeg and Pabst, 2004). Inductive sites for mucosal immunity consists of organized mucosa-associated lymphoid tissue (MALT) as well as local and regional draining lymph nodes (LN), whereas the effector sites consists of distinctly different histological compartments, including the lamina propria (LP) of various mucosae, the stroma of exocrine glands and the surface epithelia (Brandtzaeg and Pabst, 2004). Depends on the anatomical regions, MALT is subdivided into: bronchus-associated lymphoid tissue (BALT), gutassociated lymphoid tissue (GALT) and skin-associated lymphoid tissue (SALT; Brandtzaeg and Pabst, 2004). In vertebrates, mucosal immune system provides three main functions: (i) mucus membrane protection against infections, (ii) prevention of the uptake of antigens, microorganisms and foreign materials, (iii) and avoidance of the of harmful immune response development against these pathogens (Holmgren and Czerkinsky, 2005).

1.2.1. Mucosal immunity in fish

The mucosal surface of skin, gill and intestine are the first barrier between the fish and surrounded environment. MALT is present in fish, however it is less organised than described in mammals: in contrast to mammals, fish do not possess organized lymphoid tissue such as Peyer's patches and M cells or secretory IgA (SIgA) have not been described in fish yet (Rombout et al., 2011). In fish, skin, gills and gut are considered as mucosal lymphoid organs and especially part of the hindgut (2nd segment) seems to have an inducing function (Rombout et al., 1993). Uptake of macromolecules by intestinal epithelial cells and many diffusely distributed leucocytes including intraperitoneal lymphocytes (IELs) are described in numerous fish species, such as common carp (Rombout et al., 1993), goldfish (Temkin and McMillan, 1986), Atlantic salmon (Uran et al., 2008). Moreover, as recently showed by Przybylska et al. (**PAPER 2**), carp skin responses to β-glucan stimulation when bathing, resulting in an inflammatory response by activating IL-1b, IL-6 and IL-8 and local differences in expression pattern dependent on stimulation by DAMPs alone or DAMPs/PAMPs combination.

1.2.1.1. Mucus

Fish and other aquatic vertebrates are covered with mucus blanket. The mucus on the epidermal layer of fish skin is synthesized and secreted by epidermal goblet cells. Mucus present in vertebrates consists of many components, two of which are predominantly important in gel formation. These are water and large filamentous, highly glycosylated glycoproteins called mucins (Verdugo 1990; Strous and Dekker, 1992; Bansil et al., 1995; Kesimer et al., 2010).

Fish mucus has wide range of functions: (i) protects underlying epithelium against various types of pathogens, (ii) respiration, (iii) ionic and osmotic regulation (Shephard, 1994). In vertebrates, before imping epithelial surface and cause damage, foreign particles and microorganisms are entrapped in mucus; mucus blanket is also a reservoir of numerous innate immune factors such as lysozyme, immunoglobulins, complement proteins, lectins, C-reactive proteins, proteolytic enzymes and various antibacterial proteins and peptides, e.g. defensins (Shepherd, 1994; Cole et al., 1997; Subramanian et al., 2007; Narvaez et al., 2010). Vertebrates' mucins and mucus have rheological properties (viscosity and elasticity), which are important for their physiological function (Roussel and Delmotte, 2004). Based on biochemical characterisation, 19 mucin genes have been identified in humans (Dekker et al., 2002), and their gene structure and role in the infection process has been proved (McAuley et al., 2007; Thornton et al., 2008). Up to date, only two mucin genes, Muc2c and Muc5b were cloned and sequenced in fish (van der Marel et al., 2012) cause accurate assembly of the mucin genes is difficult due to the large size of the centraltandem repeats (Cox et al., 2010). However, the mucin genes were highly conserved and showed a high similarity to mammalian Muc2 and Muc5B. Analysis of tissues and organs for mucin expression in carp revealed a clear separation in the site of expression between the two mucins. Muc5B was expressed with low copy numbers in brain and liver and with high copy numbers in skin and gills. Muc5B appears to be homologous to mammalian Muc5B. Muc5B has been found to be mainly expressed in the mucous glands of the respiratory mucosa and salivatory glands, as well as in the gall bladder, pancreas and cervix (Roussel and Delmotte, 2004). Carp Muc2 was only, but highly, expressed in the first and second intestinal segment (van der Marel et al., 2012). Moreover, differences Muc5B and Muc2C expression were observed in carp with different b-glucan feeding regimens. Consistent, but not significant, down-regulation of Muc2C in the intestine and gills was seen in the glucan fed fish, while Muc5B was significantly increased in skin, with slight up-regulation in gills. Additionally, Muc5B expression is affected by β -glucan enriched bath, as it was recently reported (Przybylska et al., PAPER 2) and correlate with work presented by van der Marel co-workers (2010) where water with increased bacterial load did not induce clinical symptoms in carp, however rapid skin mucosa respond have been seen, even if the

bacteria involved were considered to be non-pathogenic (van der Marel et al., 2010). In vertebrates, beta glucans and lipopolysaccharide (LPS) work through the same receptors, hence, the mechanism by which LPS, MacroGard and 6.3 kDa oat fiber stimulate Muc5b expression might be alike. Moreover, in humans, Muc5b is one of the largest gel-forming glycoprotein in the body, however cells may differ in the density of mucin granules, suggesting a diversity of the mucins synthesized even by an individual cell (Roussel and Delmotte, 2004; Kesimer et al., 2010; Sheehan et al., 2000).

1.2.1.2. Skin

Fish skin is a complex structure composed of two layers, the outer stratified epithelium (epidermis) and the inner layer (dermis), which are separated by a basal membrane that is composed of undifferentiated cells (Buchmann, 1999). In contrast to terrestrial vertebrates, fish skin comprises living cells and hence is metabolically active (Martin, 1997; Press CM and Evensen, 1999). Pluripotent cells in fish skin, actively participate in wound healing, tissue remodelling, and are able to differentiate into epithelial cells, mucous cells, club cells or sensory cells (Rakers et al., 2010). The thickness and cells composition of epidermis varies among fish species. In common carp, epithelial cells, goblet cells, club cells and keratynocytes are mainly located in the epidermis, while chromatophores, melanophores and xanthopores are found deeper in the tissue (Iger and Bonga, 1994). In fish, the epidermis forms not only the first barrier with against environment, but represents the initial site for complex immune responses against microorganisms (Kearn, 1999). Immunological components, such as lymphocytes, macrophages and granulocytes were found in skin of teleost species (Iger et al., 1988; Peleteiro and Richards, 1990; Herbomel et al., 2001). Presence and role of specific antibodies in skin and mucus against e.g. infection of skin by I. multifiliis was previously showed (Clark et al., 1996; Dickerson and Clark, 1998). Moreover, the presence of IgM was detected in the goblet cells of trout skin (Peleteiro and Richards, 1988). Recent work presented by Przybylska et al. (PAPER 2) and Przybylska and Nielsen (PAPER 3) showed that skin together with mucus is responding to stimulation with β-glucan immune.

1.2.1.3. Gill

Gills consist of lamellae, which comprise the main respiratory surfaces of the fish (Wilson and Laurent, 2002). Recent morphological studies on nile tilapia (*Oreochromis niloticus*) have described the branchial filament as a multilayered epithelium, such as in other teleostean species (Monteiro et al., 2010). Lymphocytes (Grove et al., 2006a; Lin et al., 1998), macrophages (Lin et al., 1998; Mulero et al., 2008), neutrophils (Lin et al., 1998), eosinophilic granulocytes (Barnett et

al., 1996; Lin et al., 1998; Mulero et al., 2007) and antibody-secreting cells (ASC) (Davidson et al., 1997; dos Santos et al., 2001a) have been observed in the gill associated lymphoid tissue (GIALT) of different fish species. Presence of the B cells and T preponderance cells was shown as well (Lin et al., 1999). Mucus production has been proven to be higher in the area surrounding the gill cover than in any other skin sites (Shephard 1994).

1.2.1.4. Gut

Fish do not possess organized GALT, and thus, have no lymphoid tissue such as (i) Peyer's patches, an organized lymphoid nodules and mesenteric lymph nodes, (ii) M cells, which sample antigen directly from the lumen of gut and deliver it to antigen presenting cells, (iii) secretory IgA, which is produced by activated B-cells in the mucosa where it forms immune complexes with pathogens and allergens, thereby preventing them from binding to and penetrating the intestinal mucosa (SIgA; Rombout et al., 2005; Rombout et al., 2010). Lamina propria of teleost gut harbors a variety of immune cells including, but not limited to macrophages, granulocytes, lymphocytes and plasma cells, whereas the surface epithelium compartment is mainly composed of T cells and few B cells. Similar to mammals, immunological differences are recognized along the different segments of the fish gastrointestinal tract. Most of our knowledge in that area relates to the differential uptake of particles in the anterior gut (foregut or first segment). In cyprinids like carp (Cyprinus carpio L.) (Rombout et al., 1985) and goldfish (Carassius auratus auratus) (Temkin and McMillan, 1986), large intraepithelial macrophages containing phagocytosed material have been observed and therefore these are thought to be the main antigen presenting cells. However, no macrophages were found in the gut of seabream (Sparus aurata); (Mulero et al., 2008). Presence of IgM positive cells in gut of sea bass, turbot, carp and cod was shown (Abelli et al., 1997; Fournier-Betz et al., 2000; Rombout et al., 1998; Rombout et al., 1993; Schrøder et al., 1998).

1.3 Wound healing process

Wounds in fish can be caused by pathogens, such ectoparasites as *Ichthyophthirius multifillis* and lice, Gram-negative bacteria as *Moritella viscosus*, as well as mechanical trauma (Gonzalez et al., 2007; Ingerslev et al., 2010; Krkosek et al., 2011). In vertebrates reduced skin integrity or mucus production facilitates entry of pathogens into the underlying tissue and vascular system, and therefore rapid recruitment of immune cells to the wound site is essential (Singer and Clark, 1999). In all animals, wound healing is a complex, well-organized process which can be roughly divided into three overlapping phases: inflammation, tissue formation and remodelling, in which blood cells, soluble mediators, resident cells (fibroblasts, endothelial cells, goblet cells) and extracellular matrix components are involved (Figure 2; Singer and Clark, 1999; Gillitzer and Goebeler, 2001; Kanter and Akpolat, 2008). In fish, so far, the inflammatory phase is the best described.



Figure 2. Time line for three phases of wound healing process in mammals. The same three phases are distinguished in wounds occurred in fish. Up to date, the inflammatory (inflammation) phase is the best described. Source Enoch and Price, 2004 (modified). In present work, wound closere in skin and muscle of common carp was investigated, and similar time-wise pattern was obserwed: inflammation phase with bleeding step (day 1), epithelium cells overgrowth (day 3) and complete wound closure at day 14 (**PAPER 2**).

1.3.1 The inflammatory response following the tissue damage

Wound healing process in vertebrates is accompanied by inflammatory reactions (Singer and Clark, 1999). In mammalian model, during tissue injury, blood vessels break and disrupt homeostasis, therefore, red blood cells together with platelets form a clot, to re-establish the homeostasis and provide provisional matrix for cell migration (Singer and Clark, 1999). At the same time, inflammatory reaction starts in order to clean up the site of injury and fight pathogens: numerous vasoactive mediators and chemokines are released to recruit neutrophils and macrophages to the site of injury (Singer and Clark, 1999; Gillitzer and Goebeler, 2001). They include interleukins, lymphokines and several related signalling molecules such as tumor necrosis factor alpha (TNF- α) and interferons. All mention above have important role to play: recruitment of inflammatory cells to the wound site as well as contribution to the regulation of re-epithelization, tissue remodelling and angiogenesis (Gillitzer and Goebeler, 2001). In mammals, in response to injury and release of danger associated molecular patterns (DAMPs), which are a danger signal for surrounding cells, numerous cytokines are secreted, and selectively activate and guide leukocyte subsets during wound healing and tissue repair (Gillitzer and Goebeler, 2001). In fish, immediately after injury, erythrocytes and thrombocytes become abundant in the affected area (Jiang and Doolittle, 2003). As in mammalian mmodel, fish neutrophiles and macrophages are of particular importance due to their phagocytic activity and cytokines and chemokines production (Gonzalez et al., 2007a, b; Martin and Leibovich, 2005). Fish neutrophils appear at the injured site very fast and have been observed in skin and musculature tissue two hours after mechanical damage (Gonzalez et al., 2007a).

Immune cells appearance at the injured site is possible due to locally secreted cytokines and chemokines, as well as DAMPs by the damaged cells. In vertebrates, chemokines poses unique potential for selective activation of various leukocyte subsets (Belperio et al., 2000). Due to neutrophils infiltrations, injured area is been cleaned of foreign particles and bacteria, and macrophages are taking over (Singer and Clark, 1999). Mammalian macrophages circulate in blood as monocytes although they mature and differentiate in specific tissue types (Geissmann et al., 2010). In response to specific chemoattractants (e.g interleukin-8, IL-8), monocytes are directed to the wound area and become active macrophages. They are considered a key cells in wound repair due to their capacity to produce inflammatory cytokines as well as growth factors (Riches, 1996). Recently, process of wound closure in skin of common carp was investigated (Przybylska et al., **PAPER 2**) where bathing in β-glucans, specifically MacroGard and 6.3 kDa oat fiber, have a direct positive effects on the wound closure in common carp. Immunological and regenerative response following stimulation with pathogen associated molecular pattern (PAMPs) and DAMPs in

controlled conditions, without the exposure to pathogens was showed. PAMPs/DAMPs stimulation resulted in an inflammatory response by activating IL-1b, IL-6 and IL-8, and local differences in expression pattern dependent on stimulation by DAMPs alone or DAMPs/PAMPs combination. Expression of main pro-inflammatory cytokines in skin confirms presence, of immune related cells, such as neutrophils and macrophages and their active response to stimulation. In addition, the absence of marked differences on the respiratory burst activity in head-kidney cells supports the notion of a localized immune response to the site of injury (Przybylska et al., **PAPER 2**).

Wound healing is a complex process, and it has been proven, that not only "typical" immune cells of haematopoietic origin, such as neutrophils or macrophages are involved in wound closure and cytokines production. Fibroblasts have been described in humans as the non-conventional cell type having immune regulation capabilities (Larsen et al., 1989; Chen et al., 2005). Recent studies in human have shown that fibroblasts can not only respond to cytokines, but also generate these signals (IL-1 β , IL-4, IL-6, IL-8, IL-10, IL-12, TNF- α and IFN- γ) and influence the behaviour of immune competent cells trafficking to the injury site (Larsen et al., 1989; Chen et al., 2005; Ruiz et al., 2007). Injury could potentiate immune-like performance of fibroblasts, as well as having a direct effect on the resident cells present in skin, such as epithelial cells, resident macrophages and lymphocytes. That hypothesis has been strengthen by results obtained from research on mammalian fibroblasts. Wei et al. (2002) have presented first evidences that glucan can directly stimulate human fibroblasts. Moreover, in vitro studies in mammals presented by Iocono et al. (2000) showed that IL-8/CXCL8 had an inflammatory effect on keratinocyte proliferation and collagen lattice contraction. Rennekampff et al. (2000) demonstrated results where IL-8/CXCL8 was shown to have stimulatory effect on murine keratinocytes proliferation in vitro. We believe, wound closure process in fish is alike the one described in mammals. It is supported by results presented by Ingerslev et al. (2010) and recently by Przybylska et al. (PAPER 2). Several fibroblast cell lines have been established and characterised in fish (Collet and Secombes, 2001; Ossum et al., 2004). Ingerslev *et al.*, (2010) reported that fish fibroblasts are susceptible to DAMPs.

1.3.2 Cells involved in the inflammatory response

Cells of vertebrates' immune system are active participants in the repair of injured skin (Singer and Clark, 1999). Immune cells can greatly impact the repair process at each of these stages: hemostasis and inflammation, proliferation, and scar formation/remodeling phases (Enoch and Price, 2004). Immediately after a skin injury platelets enter the wounded area, where they aggregate and release a host of mediators that initiate the coagulation cascade, as well as growth factors and cytokines involved in the recruitment of inflammatory and immune cells (Martin, 1997; Singerand and Clark,

1999). In vertebrates, neutrophils are responsible for destroying any invading microorganisms, whereas overtaking macrophages clear the wound of debris and foreign particles (Smith and Lumsden, 1983). Both neutrophils and macrophages release an assortment of growth factors and cytokines (Riches, 1996). Eventually, T cells and other members of the adaptive immune system may become involved in wound closure in mammals (Havran and Jameson, 2010). During re-epithelialization, angiogenesis, the scar formation and remodeling phase, fibroblasts increase in number and produce a permanent scar in the repaired skin (Chen et al., 2005; Ruiz et al., 2007).

1.3.2.1 Neutrophils

In all animals, neutrophils are the most numerous and important components of host defence against bacteria, viruses and fungal infections (Smith and Lumsden, 1983). In fish, the anterior kidney is the central immune organ where neutrophils are being produced (Bols et al., 2001), however they can be found in spleen and blood, and are commonly increased in inflammatory lesions (Afonso et al., 1998). In response to inflammation, fish neutrophils migrate from blood stream to site of infection, where they bind, engulf and kill pathogens by proteolytic enzymes, antimicrobial peptides and reactive oxygen radicals production (Lamas and Ellis, 1994; Dalmo et al., 1997; Palic et al., 2005). In mammals, by degranulation, neutrophils release antimicrobial factors as well as neutrophil extracellular traps (NETs) into the extracellular medium (Brinkmann et al., 2004; Cho et al., 2005). NETs have been recently described in mammalian and fish neutrophils, but not in other cell types. In vertebrates, NETs can be produced by stimulated neutrophils or in response to cytokine stimulation, and immobilize and kill pathogens (Brinkmann et al., 2004; Cho et al., 2005; Palic et al., 2007). In fish, immune and endocrine systems are linked, therefore neutrophils are affected by high cortisol level (Weyts et al., 1998a, b; Flik et al., 2006; Palic et al., 2007). It was showed that fish exposed to handling and crowding stress for 48 hours showed the decrease in neutrophil activity measured as NETs release (Palic et al., 2007). Fish and mammalian neutrophils share morphological and functional similarities in respect to phagocytosis, chemotaxis, respiratory burst intensity, degranulation and NETs release (Lamas and Ellis, 1994; Berman et al., 2003; Hermann et al., 2004; Rodrigues et al., 2004; Palic et al., 2005a, b). Degranulation of neutrophils and release of myeloperoxidase (MPO) is an important step in oxygen mediated killing of pathogens, and it appears to be essential step in exocytosis in neutrophil extracellular traps (Brinkmann et al., 2004). Recently, particulate β -1,3-1,6-glucan from baker's yeast was found to be a potent stimulator of fish neutrophil degranulation and an optimal neutrophil immune modulator during stress conditions in fathead minnow (Ainsworth, 1994; Palic et al., 2006; 2007). Previous reports have showed strong respiratory burst response of fish phagocytes to glucans in doses from 50 to 1000 µg/ml (Castro et al., 1999; Jørgensen and Robertsen, 1995). In

contrary to that, Przybylska et al, (submitted) have reported that bath supplemented with beta glucan enriched products had no effect on respiratory burst activity in neutrophils collected from head kidney of common carp. Nevertheless, during injury, neutrophils are the first leukocytes that migrate to into inflamed tissue (Chadzinska et al., 1999).

1.3.2.2 Macrophages

Macrophages in vertebrates arise from hematopoietic progenitors, which differentiate directly, or via circulating monocytes, into subpopulation of tissue macrophages (Geissmann et al., 2010). In higher vertebrates, macrophages are multifunctional: they can ingeste microbes, secrete wide range of biologically active molecules (hydrolases, complement components, enzyme inhibitors, cytokines, eikosanoides) and reactive oxygen and nitrogen species (Auger et al., 1992). Depending on the organ, resident macrophages of higher vertebrates posses morphological and phenotypic differences (Gordon and Taylor, 2005). Fish macrophages were shown to have a long life compared to neutrophils and play a crucial role in both, the innate and the adaptive immune system (Joerink et al., 2006). In the innate immune system, they act as phagocytic cells, and producing oxygen and nitrogen radicals. In the adaptive immune system, macrophage act as professional antigen presenting cells (APCs). Fish macrophages are migratory and actively phagocytic, they generate radical species and can produce a number of immunologically important molecules, e.g. cytokines (Secombs and Fletcher, 1992; Neumann et al., 1995; Jang et al., 1995). Hanington et al., (2009) have summarized the recent findings of macrophage development in teleost. Results presented by Joerink et al., (2006) have showed that carp head-kidney macrophages are heterogeneous and, upon *in vitro* stimulation, can give rise to functionally different polarization states. First evidences for innate and alternative macrophage polarization in vivo in common carp were presented by Chadzinska et al. (2008). Moreover, teleost macrophages become activated following stimulation with bacterial products (e.g. lipopolysaccharide) and/or soluble host-derived factors (Graham and Secombes, 1988; Jang et al., 1995; Neumann et al., 1995). β-glucans have been proved to increase fish host resistance to certain pathogens (Brattgjerd et al., 1994; Dalmo and Seljelid, 1995; Jørgensen and Robertsen, 1995). Moreover, macrophages are one of the key cell types involved in wound healing process in vertebrates. Although there is not direct link between β -glucans stimulation and fish macrophages activation, Przybylska et al., (PAPER 2) have showed that during wound healing process, PAMPs/DAMPs stimulation resulted in an inflammatory response by activating IL-1b, IL-6 and IL-8 which might be indirect evidence of macrophage activation.

1.3.2.3 Fibroblasts

In mammals, the fibroblast is the last specialised member of the connective tissue, whose main function is to maintain the structural integrity of the connective tissue by secreting precursors of the extracellular matrix rich of type I and type III collagen (Alberts, 2008). The fibroblasts are responsible for the synthesis, deposition, and remodeling of the extracellular matrix. Conversely, the extracellular matrix can have a positive or negative effect on the ability of fibroblasts to synthesize, deposit, remodel, and generally interact with the extracellular matrix in mammals (Xu J, Clark, 1996; Clark et al., 1995). In mammalian model, after migrating into wounds, fibroblasts commence the synthesis of extracellular matrix, and subsequently the provisional extracellular matrix is gradually replaced with a collagenous matrix (Clark et al., 1995). It has been proven, not only "typical" immune cells of haematopoietic origin, such as neutrophils or macrophages are involved in wound closure and cytokines production in wound healing process. Fibroblasts have been described in humans as the non-conventional cell type having immune regulation capabilities (Larsen et al., 1989; Chen et al., 2005). Recent studies in human have shown that fibroblasts can not only respond to cytokines, but also generate these signals (IL-1β, IL-4, IL-6, IL-8, IL-10, IL-12, TNF- α and IFN- γ) and influence the behaviour of immune competent cells trafficking to the injury site (Larsen et al., 1989; Nirodi et al., 2000; Chen et al., 2005; Ruiz et al., 2007). Once an abundant collagen matrix has been deposited in the wound, the fibroblasts stop producing collagen, and the fibroblast-rich granulation tissue is replaced by a relatively acellular scar.

In contrast to higher vertebrates, fish skin is not keratinised and hence, metabolically active (Martin, 1997; Press CM and Evensen, 1999). Several fibroblast cell lines have been established and characterised in fish (Collet and Secombes, 2001; Ossum et al., 2004). Ingerslev et al., (2010) reported that fish fibroblasts are susceptible to DAMPs. Moreover, Przybylska et al., (**PAPER 2**) have showed that fish bath in water with β -glucan enriched products were healing faster when compared to non-treated fish. High up-regulation of IL-1 β and IL-6 in skin samples from treated carp three days post injury could be explained by the ability of "moving" fibroblasts to express genes of immune functions before the more potent immune cells have been recruited. Presence of DAMPs (wounding) and PAMPs (β -glucan) could potentiate immune-like performance of fibroblasts, as well as having a direct effect on the resident cells present in skin, such as epithelial cells, resident macrophages and lymphocytes.

1.3.2.4 Mucous cells

Fish and other aquatic vertebrates are covered with mucus layer over their entire body surface. Skin, together with mucosa forms the first line of defence against pathogens and is an essential protective barrier in aquatic organisms. In skin of mammalian epithelia, goblet cells, cells responsible for active mucus production are situated (Kanter and Akpolat, 2008). In mucus, foreign particles and microorganisms are entrapped, before imping epithelial surface and cause damage (Rogers, 1994). Few studies have emphasized the role of goblet cells and mucus in wound healing in vertebrates (Geggel et al., 1984, Kanter and Akpolat, 2008). Goblet cells differentiation and functioning is affected by mucosal immunity in mammals (Ciacci et al., 2002), and gel-forming mucins can be regulated by inflammatory cytokines such as interleukins (IL-1 β , -6, -17) or exogenous factors (Roussel and Delmotte, 2004). In fact, Jin-Jang et al., (2003) in rat model, that have found goblet cells are always lost in the presence of intense inflammation and regulation by mucogenesis in goblet cells was associated with the cessation of inflammation. In fish, mucous cells usually originate in the middle layer of epidermis, where they develop and mature (Bullock and Roberts, 1974). Mucous cells of different types are found in fish skin (Pinky et al., 2008). It was reported by Mittal et al., (2000), that the major constituents of the surface secretions in fish, consists mainly of glycoproteins of heterogeneous molecular structure. The density of mucous cells varies greatly among fish species, body regions and growth stage (Bullock and Roberts, 1974; Yamamoto et al., 2011).

1.3.3 Molecules involved in the inflammatory response

In general, the extent of the inflammatory response in vertebrates can vary depends on the triggering stimuli (Enoch and Price, 2004). It is important that occurred immune response is powerful enough to clean up debris and combat pathogens subsequently. During inflammation phase in mammals, numerous cytokines are secreted, and selectively activate and guide leukocyte subsets during wound healing and tissue repair (Gillitzer and Goebeler, 2001). They work in autocrine and paracrine manner and more seldom in an endocrine way. In general, fish posses a similar repertoire of cytokines as mammals, however the function of each fish cytokine is not so well established as for mammalian cytokines (Secombes et al., 1996). Some of the cytokines are in high importance during wound healing process and interleukin 1 β (IL-1 β), interleukin 6 (IL-6), interleukin 8 (IL-8), the best described cytokines in fish (Scapigliati et al., 2000, Secombes et al., 1996) are among them.

1.3.3.1 Interleukin 1 β (IL-1 β)

In mammals two isoforms are present: IL-1 α and IL-1 β , but only IL-1 β was characterized in fish. IL-1 β is probably the the best described cytokine in fish. It is a main pro-inflammatory cytokine with multiple functions such as stimulating cell division, cells activation, expression of other cytokines and induction of fever. Main IL-1 β source in vertebrates are polymorphonuclear laukocytes (constitute nearly 50% of all cells at the wound site; Gillitzer and Goebeler, 2001) and macrophages, as well as some resident cells (Hübner et al., 1996, Feiken et al., 1995). In injured sites, IL-1ß activates the lining endothelial cells and cause enhancement of leukocyte migration. In vertebrates, IL-1β was shown to be strongly up-regulated during the inflammation (Hübner et al., 1996, Grellner et al., 2000; Chadzinska et al., 2007). Cortisol together with LPS synergistically stimulates the expression of IL-1ß in fish HK phagocytes (Engelsma et al., 2003), however in vitro studies in common carp showed inhibition of inflammatory cytokine due to acute stress (Saeij et al., 2003). Interleukin 1 is one of the cytokines produced by macrophages in response to an immunogenic stimulus in fish. Previous studies revealed that glucan treatment in fish enhanced the expression of interleukin 1 [Fujiki et al., 2000]. Injection with Astragalus polysaccharide, the IL-1b mRNA level increased in a dose-dependent manner in the head kidney, while no significant changes were found in the gill and spleen (Galina et al., 2009). Chadzinska et al., (2008) reported migration of inflammatory leukocytes from HK into peritoneum during intraperitoneal injection of zymosan particles. Constitutive expression of IL-1ß in peritoneal leukocytes (PTL) was two-fold higher than in HK, however during zymosan-induced peritonitis, IL-1ß expression showed twice higher expression in peritoneal leukocytes (PTL) than in HK. In addition, Przybylska et al., (**PAPER 2**) have showed changes in IL-1 β expression following wound healing process and stimulation with β-glucans. Furthermore, Przybylska and Nielsen (PAPER 3) reported changes in IL-1 β expression followed intravenous injection with β -glucan, suggesting response to PAMPs stimulation.

1.3.3.2 Interleukin 8 (IL-8)

Interleukin 8 belongs to CXC chemokine family and its secretion activates and tracks neutrophil granulocytes to site of inflammation in vertebrates (Jimenez et al., 2006). As is well established in mammals, CXCL8 is among the first chemokines expressed during inflammation. The proinflammatory cytokines like IL-1 b and TNF-a induce CXCL8 expression in endothelial cells, fibroblasts and local immune cells in mammals (Larsen et al., 1989; Strieter et al., 1989). IL-8 regulates is known regulator of neutrophil trafficking (Rennekampff et al., 2000; Werner and Grose, 2003, Grellner et al., 2000; Gillitzer and Goebeler, 2001) and augments angiogenesis in

mammals (Li et al., 2003; Yao et al. 2002). It was recently reported that in two fish species that belong to the cyprinid lineage, common carp and zebrafish, two CXCL8-like lineages evolved: CXCa_L1 (lineage 1) and CXCL8_L2 (lineage 2) (Abdelkhalek et al., 2009; van der Aa et al., 2010). Nevertheless, presence of two CXCL8 lineages in carp appears to be functional homologs to mammalian IL-8, and has crucial biological role in chemotaxis (van der Aa et al., 2010). Results presented by Sangrador-Vegas et al., (2002) reported that stimulation rainbow trout fibroblasts induce IL-8 expression. Recently, Przybylska et al., (**PAPER 2**) have showed that IL-8 is inducible in skin and muscle of common carp, by introduction of PAMPs and DAMPs.

1.3.3.3 Interleukin 6 (IL-6)

Expression of IL-8 in mammals is correlated with interleukin 6 (IL-6); (Nielsen et al., 2007). Interleukin 6 is a multifunctional cytokine that is produced by a variety of hematopoietic and nonhematopoietic cell types. Vertebrates IL-6 is produced and secreted by macrophage, neutrophils, T cells, as well as resident cells (20 Hübner et al., 1996, 14 Gillitzer and Goebeler, 2001; Lin et al., 2003). IL-6 seemd to be one of the key cytokined involved in wound healing process in mammals: experiments on diabetic or IL-6 knockout mice have shown drop in neutrophil and macrophage number, as well as a reduction in collagen production and deposition (Lin et al., 2003, Berdal et al., 2007). The administration of IL-6 at the wound site induces massive leukocyte infiltration in mice (Saba et al., 1996) and induces collagen production (Duncan and Berman, 1991; Greenwel et al., 1993). Orthologous of the mammalian IL-6 members have been identified in fish (Bird et al., 2005, Iliev et al., 2007, Castellana et al., 2008) and recently, Varela et al., 2011 have confirmed that zebrafish IL-6 shares structural characteristic with IL-6 from other vertebrates and its expression follows immune stimulation. IL-6 was found to follow β -glucans stimulation during bath treatment in skin and muscle of common carp (Przybylska et al., **PAPER 2**).

1.3.3.4 Heat shock protein 70 (Hsp70)

Heat shock proteins (Hsps) are highly conserved stress proteins, found in all prokaryotic and eukaryotic cells and are classified according to their molecular weight or to the way they are expressed. 70kDa Hsps are the most conserved and the most studied class of Hsps. In mammals, Hsp70 is highly inducible during inflammatory reaction caused by stress (Li et al., 1992; Trautinger, 2001; Jacquier-Sarlin et al., 1995; Guo et al., 2007; Novoselova et al., 2006). Hsps function as "danger signals" – damage associated molecular patterns (DAMPs), released endogenous structures from damaged or dead cells in all vertebrates (Matzinger 2002, Seong and Matzinger, 2004).

Hsp families have been reported in fish cell lines, primary cell cultures, as well as in various tissues from whole animals (Iwama et al., 2004, Toa et al., 2004). Up-regulation of the Hsps in fish was observed in various stress conditions e.g. exposure to bacterial pathogens (Jia et al., 2000), environmental contamination such as heavy metals (Duffy et al., 1999), pesticides (Hassanein et al., 1999). The Hsp response can vary according to tissue (Rabergh et al., 2000), Hsp families (Wagstaff et al., 1998), stressor (Airaksinen et al., 2003), species sensitivity (Basu et al., 2002) and season (Fader et al., 1994). In fish, Hsp70 responds to a wide range of substances and treatments such as hormones, chemicals, anoxia and viral infection (Deane et al., 2007; Erdogan et al., 2007; Feng et al., 2003; Purcell et al., 2004). Hsps are chaperone proteins, but are involved in immune response as well. They are "danger signals", released during e.g injury and alert surrounded cells about ongoing inflammation or pathogen presence. Stolte et al., (2009) showed that an immune stimulus modulates expression of Hsp70 *in vitro* and *in vivo*. As showed by Przybylska and Nielsen (**PAPER 3**), increase in Hsp70 expression during common carp intravenous injection with high dose of β -glucan was observed.

1.3.4 Pathogen associated molecular patterns (PAMPs) and damage associated molecular patterns (DAMPs)

1.3.4.1 PAMP recognition

The immune system has to be effecting in distinguishing "self" from "non-self" and combating pathogens (Figure 3). They are constrained, however, by relying on a limited and invariant repertoire of receptors to recognize microorganisms. The pathogen-recognition receptors of macrophages, neutrophils, and dendritic cells recognize simple molecules and regular patterns of molecular structure known as pathogen-associated molecular patterns (PAMPs), exogenous molecules (antigens) produced only by potential pathogens (Medzhitov and Janeway, 2000). Janeway Medzhitov (2002) proposed that infection would be detected by innate receptors for microbiological biomolecules that are either absent or different in mammals. These receptors are known as pattern recognition receptors (PRRs) and they recognize structures such as mannose-rich oligosaccharides, peptidoglycans and lipopolysaccharides in the bacterial cell wall, and unmethylated CpG DNA, which are common to pathogens and have been conserved during evolution (Roach et al., 2005). PRRs have several functions: stimulate ingestion of the pathogens they recognized, guide cells to the site of infection and induce production of effector molecules. Important receptors recognising the pathogens are the Toll-like receptors (TLRs) and Nod-like receptors (NLRs) (Medzhitov and Janeway, 2000). However, microorganisms evolve more rapidly than their hosts, and this may explain why the cells and molecules of the innate immune system

recognize only molecular structures that have reminded unchanged during evolution. Hostpathogen coevolution promotes that random selection promotes recognition sites with a broad specificity for essential non-variable molecular structures on the pathogen (e.g. lipopolysaccharide or other pathogen-associated molecular patterns, PAMPs). In all vertebrates, the primary function of PRRs and the innate immune response is to provide immediate protection from pathogens (Takeda et al., 2003; Thoma-Uszynski, et al., 2001). This initial recognition of pathogen leads to the production of a variety of inflammatory mediators, including chemokines, cytokines, vasoactive amines and eicosanoids (Medzhitov, 2008). The main and most immediate effect of these mediators is to elicit an inflammatory exudate locally: plasma proteins and leukocytes (mainly neutrophils) that are normally restricted to the blood vessels, access the extravascular tissues at the site of infection or injury (Medzhitov, 2008). In fish, polymorphonuclear granulocytes and mononuclear phagocytes stimulated by PAMPs produce inflammatory cytokines or reactive oxygen and nitrogen species (ROS); (Jault et al., 2004; Purcell et al., 2006).

Distinct pathogens express different PAMPs, hence a combination of specific sets of PRRs is essential for integrating an immune response against a specific pathogen. Four main types of PRRs have been described in fish: Toll-like receptors (TLRs; Jault et al., 2004; Rodriguez et al., 2005), NOD-like receptors (NLR; Laing, et al., 2008), C-type lectin receptors (CLRs; Soanes et al., 2008) and peptidoglycan recognition proteins (PGRPs; Li et al., 2007).

In vertebrates, in addition to their role in mammalian host defense from deleterious microbial infection, receptors for PAMPs recognition are also involved in various aspects of mammalian homeostasis such as development, the recognition of cellular and tissue injury, and tissue repair and regeneration. The repair and regeneration of tissue after injurious insult is a complex process. However, the first generation of information from this nascent field suggests that TLRs may be important in many stages of this process. In providing pro-survival signals and in preventing apoptosis, TLRs may dictate the threshold of injury and cellular death, thereby limiting the extent of damage to initial injury. In addition, signaling induced after injury, such as the TLR-dependent production of prostaglandins in tissue stroma (Fukata et al., 2006, Brown et al., 2007) or the up-regulation of anti-apoptotic factors may provide signals to keep both differentiated and progenitor cells within the tissue alive.

1.3.4.2 DAMPs recognition

In recent years it has been discovered that "self" cells and molecules are capable or triggering inflammatory response. In 1994, Metzinger proposed a Danger model, which suggests that the immune system is more concerned with damage than with foreignness, and is called into action by alarm signal from injured tissues, rather than by the recognition of non-self (Matzinger, 1994). In Danger model, "danger" defines if the immune response must take place or not (Matzinger, 2002; Matzinger 2002). According to this hypothesis, some kind of message should "tell" dendritic cells (DCs) that tissue cells are injured or dying in a violent, stressful or unexpected fashion (Matzinger, 2002; Matzinger, 2002). During damage, infection, necrosis and apoptosis, injured cells can release intra-cellular components such as mitochondrias, DNA, RNA, heat shock proteins, hyaluron or chromatin-associated proteins, and trigger immune response in the surrounding cells (Metzinger, 1994). In that case, any internal molecule that is normally not secreted, be it cytoplasmic, nuclear or membrane, could act as activator of dendritic cells. Thus any cell damage would be noticed, antigen-presenting cells would be activated and initiate an inflammation. If the damages were noninfectious, a small autoreactive response might be initiated, followed by inflammatory mechanism to clean up. Recent evidence points to various molecules capable of sounding the alarm, however the search for molecular messages used by the immune system to decipher danger is far from being over. So far, the following molecules are classified as "danger signals":

- Heat shock proteins (Hsps) scaffold proteins that bind, fold and protect other proteins and peptides inside cell compartments. Many of them are overexpressed under stress. Hsps can induce maturation of DCs, and on the other hand, they may efficiently transfer antigenic peptides from the cell that is succumbing to the dendritic cells (Palazon et al., 2008).
- RNA and DNA from apoptotic cells under normal circumstances, nucleic acids are confined within the limit of membranes and are not accessible, hence membrane damage abruptly expose them. Moreover, it was shown that upon activation/ death, neutrophils project nets of DNA that trap microbes and may also be considered as endogenous alarm (Wartha and Henriques-Normark, 2008).
- Nucleotides recent evidence also points to a role of nucleotides in danger signalling, including short-lived extracellular ATP (Piccini et al., 2008).
- Extra domain A (EDA) of fibronectin plays an active role in inflammation and remodelling, and recently was found to be expressed in inflamed tissues (Lasarte et al., 2007).

- Uric acid its content increases in injured cells and seems to be a principal endogenous danger signal that stimulates DCs (Shi et al., 2003).
- HMGB-1 a non-histone complex of the nuclear proteasome, ought to be confined into compartments that are not exposed (Rovere-Querini et al., 2004). It was shown to have stimulatory effect on macrophages, enhancing TNF-α expression and secretion (Bianchi and Manfredi, 2007).

Vertebrates' antigen presenting cells (APCs) can be activated from their immediate environment and respond to both, endogenous and exogenous signals. Vertebrates and bacteria have shared eons of evolutionary time, and thus receptors for endogenous and exogenous signals may have evolved simultaneously. Indeed, there is evidence that these receptors are often the same molecules. For example TLR4 is a receptor for the bacterial product lipopolysaccharide (LPS), the Hsp70 and the extracellular breakdown products of hyaluron; TLR2 binds bacterial lipoproteins and Hsp60; and TLR9 binds to DNA CpG sequence in all living organisms (Metzinger, 2002). Hence, it appears TLRs actions bind PAMPs and DAMPs.



Figure 3. Recognition of DAMPs and PAMPs by immune cells. DAMPs are released from the host cells during injury. PAMPs are recognized by PRR on the surface of immune cells. DAMPs and PAMPs bind to the same receptors on effective cells and trigger an inflammatory response. In current work, wound healing process in relation to DAMPs/PAPMs stimulation was examinated in skin and muscle of common carp (**PAPER 2**). In **PAPER 3**, immunoglobulins switch-like process was examinated following PAMPs intravenous injection into carps' system.

1.4 Stress reaction

1.4.1 Teleost hypothalamo-pituitary-interrenal (HPI) axis

When internal equilibria and proper physiological functioning are disturbed or threatened to become disturbed as result of external or internal stimuli (stressors), the animals physiological state is defined as stress (Chrousos and Gold, 1992). Animals respond to stress at different levels: body, tissue and cell (Iwama et al., 2004). On the body level the stress response is typically characterized by activation of the hypothalamus – pituitary – adrenal axis (HPA) and release of glucocorticoids and catecholamines.

Teleost fish have a fully functional stress axis. Their stress response is characterized by the activation of hypothalamo-pituitary-interrenal axis (HPI; Figure 4). Neurons in the nucleus preopticus of the hypothalamus (NPO) release corticotropin-releasing hormone (CRH). CRH stimulates corticotrope cells in the pituitary pars distalis (PD) to secrete adrenocorticotropic hormone (ACTH). ACTH stimulates cortisol release into the blood from interrenal cells in the head kidney (Flik et al., 2006). Interestingly, fish head kidney combines endocrine and immune functions, as it is major hematopoietic and lymphopoietic organ (Weyts et al., 1998, Flik et al, 2006). This organization suggests that the endproducts of the stress axis have direct paracrine access to the cells of the immune system. Vice versa, chemical signals of the immune system e.g. cytokines could exert direct paracrine action on the interrenal (steroid) and chromaffin (catecholaminergic) cells (Weyts et al., 1998).



Figure 4. Schematic representation of the hypothalamo-pituitary-interrenal (HPI) axis in teleostean fish. The hypothalamus secretes corticotropin-releasing factor (CRF), which stimulates the pituitary to release adrenocorticotropic hormone (ACTH). The interrenal gland stimulated with ACTH, releases cortisol, the primary stress hormone in fish. A negative feedback system acts on the hypothalamus to ensure homeostatic regulation. In **PAPER 3** of the present work, cortisol level in blood of common carp was investigated following PAMPs intravenous injection.

1.4.2 Glucocorticoids in fish

In vertebrates, the most commonly measured circulating hormone levels, as indicators of stress, are the catecholamines and corticosteroids. In teleost, cortisol is the principal glucocorticoid secreted by the interrenal tissue (steroidogenic cells) located in the head kidney (Iwama et al., 1999). In the circulation of fish, the bulk (80%) of cortisol is bound to serum steroid globulins, which leaves only 20% of the cortisol in its unbound and bioactive form (Flik and Perry, 1989). Basal plasma cortisol levels species-specific and stressor-specific, with range from 5-50 ng/ml (Martinez-Porchas et al., 2009). In teleost, this hormone is released after HPI axis activation when undergoes stress conditions (Mommsen et al., 1999). Plasma cortisol can rapidly increase and reach the peak within 30 minutes (Mommsen et al., 1999; Tanck et al., 2000). However, after cessation of the stressor,

plasma cortisol quickly drops again to basal levels. Moreover, once released to the circulation, cortisol is converted and inactivated to cortisone, which lacks the immune regulatory properties of cortisol (Weyts et al., 1997). Similar to mammals, glucocorticoids (GRs) are present in fish tissue and take part in the glucose metabolism, cognition and osmoregulation what is connected with the fact, that teleost fish do not possess the mineralocorticoid aldosterone, and use cortisol instead for osmoregulation (Dean et al., 2003). Although details of the formation of the GR heterocomplex are yet to be established in teleost fish, it has been assumed, based on the conservative sequence of Hsp70, Hsp90 and the GR ligand-binding region, that comparable processes takes place in all fish (Stolte et al., 2006).

Interestingly, *in vitro* studies showed that, in common carp, elevated cortisol level that occur during stress rescues neutrophilic granulocytes from apoptosis (Weyts et al., 1998b). Activated B-lymphocytes, however, show increased apoptosis when exposed to high levels of cortisol (Weyts et al., 1998a). Thus, in stressed fish, the innate immune system becomes activated. This is further illustrated by the redistribution of neutrophilic granulocytes from the head kidney into circulation (Huising et al., 2003). This stress-induced neutrophilia is considered to improve peripheral surveillance and is reported in mammals as well (Dhabhar et al., 1996).

1.5 Antibody response

Immunoglobulins (Igs) are the central component of the adaptive arm of the immune system. In mammals, five classes of Igs, including IgG, IgM, IgD, IgA and IgE, have been well identified, whereas in teleosts, IgM and IgD genes have been cloned and characterized (Wilson et al., 1990, 1997; Stenvik et al., 2000; Hordvik et al., 2002; Hirono et al., 2003; Saha et al., 2004; Savan et al., 2005). Although fish equivalents of IgG, IgA and IgE have not been identified, a third teleost Ig heavy chain isotype IgZ (IgZ - zebrafish; IgT – trout; IgH in fugu) has been recently described from fish (Danilova et al., 2005; Hansen et al., 2005; Gambon-Deza et al., 2010; Zhang et al., 2010). Although, the basic gene structure of the new class of Ig is conserved, not every teleost species, e.g. channel catfish seem to have the sequence of IgZ or IgT (Bengten et al., 2006).

1.5.1 IgM

The major immunoglobulin of fish blood serum is tetrameric IgM-like molecule (Koumansvandiepen et al., 1995). Due to the fact that teleost fish IgM shares some structural and functional characteristics with mammalian IgM, the ability of various teleost fish to produce heterogenous mixtures of IgM polymers, monomr (Rombout et al., 1993) and halfmer subunits (Pucci et al., 2003) is noteworthy. Due to usage of different methods for IgM detection, comparison of published data is complicated. In general, the first appearance of B-lymphocyte and immunoglobulins is late in marine species compared to fresh water species (Chantanachookhin et al., 1991). However, during larval and organs development IgM, is a pre-dominant immunoglobulin (Ryo et al., 2010). In adult fish, IgM is constitutively expressed in renal hematopoietic tissue, spleen and thymus, but expressed at low levels in the skin and gill (Saha et al., 2004). Recent gene expression studies in zebrafish have shown that IgM is more predominant in both lymphoid and non-lymphoid organs compared to IgZ (Danilova et al., 2005; Ryo et al., 2010). More importantly, the higher expression observed in the kidney compared to the spleen and gut might be attributed to the first appearance of IgM positive B-cells in the kidney, which may be transported to spleen, gut and other secondary lymphoid organs (Huttenhuis et al., 2006). Recently, it was showed that intravenous injection with β -glucan does not affect IgM expression in HK and skin of common carp (Przybylska and Nielsen, **PAPER 3**).

1.5.2 IgZ

IgZ and IgT were named differently because they were initially discovered in zebrafish and trout, respectively. IgZ and IgT share similar characteristics, e.g. heavy chains are composed of four Cregions and are highly homologous, indicating both of which represent the same immunoglobulin isotype (Danilova et al., 2005; Hansen et al., 2005), nevertheless IgZ and IgT genes differ in genome organization and location (Danilova et al., 2005). Functionally, both IgZ and IgT are expressed during embryonic development, indicating it could play a protective role during initial stages of larval development (Danilova et al., 2005; Hansen et al., 2005). However, in adult fish, IgZ is limited to the primary lymphoid tissue, whereas IgT is expressed in a variety of tissues, suggesting functional differences (Hansen et al., 2005). Ryo et al. (2010) have recently shown differential expression of the IgZ variants (IgZ1 and IgZ2 chimera) in response to pathogens and suggested that the IgZ subtypes in carps may have mutually exclusive humoral functions. It was shown that IgZ1 is expressed systematically and IgZ2 chimera is preferentially expressed at mucosal sites. IgZ1 and IgZ2 were equally expressed in head kidney, trunk kidney and spleen. In peripheral blood leukocytes, the expression of IgZ1 is significantly higher than in other tissues compared to IgZ2 and IgM, indicating that this antibody might be restricted to pathogens in blood. In contrast, the IgZ2 gene is expressed at relatively higher level in gut and gills, which in fish are the major routes of infection, thus this antibody might protect the host from epithelial pathogens. IgZ1 mRNA increase was seen during the blood parasite infection, T. borreli, as well as during vaccination with V. anguillarum antigens, which are rich in lipopolysaccharides (Ryo et al., 2010). In vertebrates, beta glucans and lipopolysaccharide (LPS) work through the same receptors, hence, we believe, the mechanism by which IgZ1 expression was triggered in T. borreli infection could be
alike while injecting beta glucans. Moreover, Ryo et al. (104. 2010) reported that IgZ1 functions as antibody that neutralizes foreign antigens in the fish blood. It could explain high expression of IgZ1 reported by Przybylska and Nielsen (**PAPER 3**) – if mechanisms triggered by LPS and β -glucan are alike, IgZ1 level should increase to clear circulation from injected β -glucan.

1.6 β-glucans as immune modulators

 β -glucans are naturally occurring polysaccharides from cereals (e.g. barley, oat), algae, mushrooms and yeast cell walls. They have various effects in all organisms from plants to vertebrates (Rosenberger, 1976). β -glucans easily form gel containing high-order structures of single spirals or triplet spirals due to its unique primary structure; therefore, its purification is extremely difficult, and consequently crude β -glucan fractions have been used in studies in mammals, rather than purified β -glucan (Kimura et al., 2007). In vertebrates, glucan exerts a beneficial effect on a variety of experimental disease states of bacterial, viral, fungal and parasitic origin (DiLuzio and Williams, 1978; Williams et al., 1983, Williams and DiLuzio, 1985, Yano et al., 1991, Raa et al., 1992, Duncan and Klesius 1996, Rodriguez et al. 2009, Zhang et al. 2009).

1.6.1 Diversity and structure

Glucans are polysaccharides that only contain glucose as structural components. Beta 1,3-D glucans are chains of D-glucose rings connected at the 1 and 3 positions. Smaller side chains branch off the 1,3 polysaccharide "backbone". The most active form of beta 1,3-D glucans are those that contain 1,6 side-chains branching off from the longer beta 1,3 glucan backbone (Akramiene et al., 2007). These are referred to as β -1,3/1,6 glucan, whereas 1,3/1,4 β -glucans are linear polysaccharides consisting of β -D-glucosyl residues linked through 1,3- β and 1,4- β -glycosidic linkages, and are localized in the cell wall of cereals, rice and wheat (Cui, 2001; Figure 5).

 β -glucans derived from different sources vary in structure and bioactivity (Akramiene et al., 2007). Differences in length of polysaccharide chain, extent of branching, and the length of those branches can result in the difference between material extracted by hot water, insoluble cell wall component and in different molecular weight (Akramiene et al., 2007). Research has shown that insoluble (1,3/1,6) β -glucan, has greater biological activity than that of its soluble (1,3/1,4) β -glucan counterparts (Ooi and Liu, 2000). Another variable is the fact that some of beta glucans exist as single strand chain, while backbones of the beta-1,3 glucans exist as double or triple stranded helix chain. Bohn and BeMiller (1995) have described β -glucans as biological response modifiers, in which activity correlates with branching degree and size. They have reported that less branched

glucans with molecular weight of 5 kDa-10 kDa were non-active. Conversely, Tanioka et al. (2011) have shown that barley-derived β -glucan (~2 kDa) stimulates maturation of mouse dendritic cells. In addition, up-regulation (~ x30) of IL-6 expression in bone marrow cells has been seen when compared to treatment with 40-70 kDa barely β -glucan (Tanioka et al., 2011). Przybylska et al., (**PAPER 2**) have reported that both β -glucans, MacroGard and 6.3 kDa oat fiber, promote wound healing process in common carp, however with higher wound closure ratio in group treated with MacroGard. Authors concluded it could be related to high branching level.

In general, *in vitro* studies on mammalian model have suggested that large molecular weight β glucans (e.g. zymosan) can directly activate leukocytes and stimulating their phagocytic, cytotoxic (Brown and Gordon, 2003). Williams et al. (1996) have shown that these carbohydrates can stimulate production of proinflammatory cytokines (IL-1 β , IL-8, IL-6 and TNF- α). Low or intermediate molecular weight β -glucans (e.g. glucan phosphate) posses biological activity *in vivo*, but their cellular effects are not clear (Williams et al., 1996). Data showed by Nakagawa et al. (2003), suggest that these β -glucans can stimulate production of IL-8 and IL-6, but no Il-1 β or TNF- α . Glucans at the size in a range of 5-10 kDa (e.g. laminarin) are considered as inactive (Bohn and BeMiller, 1995).

Final effect of β -glucans immune modulation depends on preparation and used protocol. In present thesis, two different β -glucan enriched products were examinated: MacroGard, provided by Biorigin (Sao Paulo, Brasil) and 6.3 kDa fiber oat provided by Scan Oat (Sweden). In order to dissolve soluble β -glucan fractions, solutions of both products were stirred for 1 h at 90 °C and autoclaved (121 °C, 15 min. 1atm.).



Figure 5. Diagram showing orientation and location of different β-glucan linkages.

1.6.2 Administration routes and its effect on teleost immunity

The immune modulatory effect of β -glucans has been shown in a number of studies (Yano et al., 1991, Raa et al., 1992, Duncan and Klesius 1996, Rodriguez et al. 2009, Zhang et al. 2009). These naturally occurring polysaccharides differ in length, extent of branching and bioactivity (Akramiene et al., 2007). In vertebrates, β -glucans of similar structure and molecular weight (m.w.) trigger different biological effects in in vivo and in vitro (William et al., 1996, Brown and Gordon; William et al., 1996, Nakagawa et al., 2003, Bohn and BeMiller, 1995). In vitro studies have shown that large molecular weight β -glucans (e.g. zymosan) can directly activate leukocytes and stimulating their phagocytic, cytotoxic (Brown and Gordon, 2003). Williams and co-workers have shown that these carbohydrates can stimulate production of pro-inflammatory cytokines (IL-1β, IL-8, IL-6 and TNF- α) (Williams et al., 1996). Low or intermediate molecular weight β -glucans (e.g. glucan phosphate) posses biological activity in vivo, but their cellular effects are not clear (Williams et al., 1996). Data showed by Nakagawa et al. (2003) suggest that these β -glucans can stimulate production of IL-8 and IL-6, but no Il-1β or TNF-α. Glucans at the size in a range of 5-10 kDa (e.g. laminarin) are considered as inactive in general (Bohn and BeMiller, 1995). On the other hand, even β-glucans of similar structure and molecular weight can show different biological activity in in vivo and in vitro studies (Akramiene et al., 2007). β-glucans have been shown to accelerate the wound healing process in mammals, however not yet in fish (Wolk and Danon, 1985, Hon Kwon et al., 2009).

Three different way of administration are available: injection, immersion or feed. Depends on purpose, the mode of immune stimulant application is very important and should be chosen with regard to, for example, labour input or stress factor. Immune stimulants are widely used in aquaculture. β -glucans injections is the most common and efficient route. Yano et al. (1991) injected carp (*Cyprinus carpio* L.) with glucan extracts from yeast prior to *Edwardsiella tarda* challenge. All untreated fish died within 3 days, whereas fish injected with β -glucan extract had survival rates of 60-80%. Data showed that the glucan-injected carps had higher activity of the alternative complement pathway. Chen and Ainsworth (1992) injected catfish (*Ictalurus punctatus*) with baker's yeast-derived glucan alone or with *E. tarda*. Data showed increase in non-specific defence parameters of phagocytic cell activity. Selvaraj et al. (2005) have demonstrated that intraperitoneal injection of yeast glucan significantly improved the survival rate of *C. carpio* against challenged with *A.hydrophila*. Administration of 500 and 1000 µg glucan/fish elicited 100% survival in carp. Robertson et al. (1990) reported that intraperitoneal injection of yeast glucan survival resulted in increase resistance in Atlantic salmon against challenged with *Vibrio anguillanum*, *Vibrio salmonicida* and *Yersinia ruckeri*. Catfish injected with yeast glucan showed increase

resistance against challenge with *Edwardsiella ictaluri* (Chen and Ainsworth, 1992). Taken all together, these data highlight the biggest advantage of injection: direct glucan- organ contact, which induces macrophages to enhance the non-specific cellular immune response consequently.

Feeding trials might be difficult to give full picture of immune stimulant effects due to chemical action of the enzymes and digestive fluids, which would change or reduce immune stimulants effectiveness, or because they would be removed from the body too quickly to have any effect. Moreover, only particles less than one micron are capable of passing through the walls of intestine and blood capillaries, these data indicate that oral administration as highly efficient (Selvaraj et al., 2005). Nevertheless, Roerstad et al. (1992) demonstrated that fish fed with MacroGard supplemented feed, showed increased resistance to infection caused by *Aeromonas salmonicida*. van der Marel et al. (2012) have shown that carp fed with β -glucan supplemented diet had higher mucosal immune response when compared to control group. Moreover, significantly higher expression levels of β -defensin 2 in gills and both β -defensin genes in skin were found.

Immersion method is an alternative, easy and fast route for β -glucans administration to fish in all sizes (Raa et al., 1992). However it is extremely expensive to introduce bathing trials to farmed fish species and in consequences very little information is available on bathing protocols and dosage. The study of Selvaraj et al. (2005) where bathing trials were carried out as fish immersion in the aerated glucan solution for about 90 min., showed that bathing did not increase survival percentage in common carp challenged with *Aeromonas hydrophila*, and no increase in antibody response was observed either. Fish probably cannot absorb glucan particles through the bathing route, as they are insoluble in nature. Results presented by Selvaraj et al. (2005) revealed that glucan administration through bathing and oral routes did not improve the survival percentage of carp. Chinook salmon immersed in glucan solution did not show protection against bath challenge with *Aeromonas salmonicida* (Nikl et al., 1993).

Many authors have also reported enhanced bacterial killing activity in glucan treated teleost fish (Yoshida et al., 1995; Jorgensen and Robertsen, 1995). Yano et al. (1991) injected carp with glucan extracts from yeast prior to *Edwardsiella tarda* challenge and observed 60-80% survival when compared to non-injected group. Glucan-injected carps had higher activity of the alternative complement pathway as well. Chen and Ainsworth, (1992) injected catfish (*Ictalurus punctatus*) with baker's yeast-derived glucan alone or with *E. tarda* and showed an increases in non-specific defence parameters of phagocytic cell activity, the specific immune response and protection against disease.

1.7 Methods applied in the study

1.7.1 The damage model

Few studies with damage model involved were performed in order to investigate the immune response in fish. Gonzalez et al. (2007) performed mechanical tissue damage in common carp in order to mimic the inflammatory responses to ectoparasite *I. multifiliis*. Ingerslev et al. (2010) investigated local tissue response in the musculature of rainbow trout (*Oncorhynchus mykiss*) following mechanical tissue damage and natural infection with *Moritella viscosus*. According to study of Selvaraj et al. (2005) immersion in beta glucan solution did not increase survival percentage in common carps challenged with *Aeromonas hydrophila*, and no increase in antibody response was observed. However, biological effect of bath supplemented with immune stimulants on innate immunity in fish was not addressed. In order to study the biological impact of two commercially available β -glucan enriched products on the wound healing process, fish were used to create wounds on the left side, above lateral line. Using an 8 mm biopsy punch, the skin and muscle tissues samples were collected. Apart sampling, an internal controls from non-injured side were taken, which allowed us to study local versus systemic response (Figure 6).



Figure 6. Illustration of wounding (A) and sampling (B-E) procedures (PAPER 2). Fish were wounded on the left side above lateral line (red circle) with 5 mm biopsy punch. During sampling, skin and muscle tissue were collected from the damaged site (red circle), as well as from non-damaged site (internal control, green circle) using 8 mm biopsy punch.

1.7.2 Intravenous injection

Three different way of administration are available: injection, immersion or feed. Depends on purpose, the mode of immune stimulant application is very important and should be chosen with regard to, for example, labour input or stress factor. Although, injection is the most time consuming and rough administration route, it is the most common and efficient route. Yano et al. (1991) injected carp with glucan extracts from yeast prior to *Edwardsiella tarda* challenge and observed 60-80% survival when compared to non-injected group. Glucan-injected carps had higher activity of the alternative complement pathway as well. Chen and Ainsworth, (1992) injected catfish (*Ictalurus punctatus*) with baker's yeast-derived glucan alone or with *E. tarda* and showed an increases in non-specific defence parameters of phagocytic cell activity, the specific immune

response and protection against disease. Taken all together, these data underlie the biggest advantage of injection, direct glucan – organs contact, which induces macrophages to enhance the non-specific cellular immune response.

Our bath model showed, that wound healing process occurs faster in carp kept in water supplemented with β -glucan enriched products. Immunizing substances, or inoculations, are generally given by injection. Intravenous injection guarantees that injected substance (medication, immune stimulant, etc.) will act more quickly and reach the subcutaneous tissues, the vascular tree, or an organ. We investigated if intravenous (i.v.) injection of *Aeromonas hydrophila* bacterin (BSK10; Nielsen et al., 2001) with two dosages of β -glucan derived from Baker's yeast has a biological effect on immune parameters in common carp. Fish were divided randomly into three groups of 23 individuals and all fish were injected with 100 µl of: 23 fish with 100 µl of bacterin in sterile PBS (B); 23 fish with 100 µl of bacterin + low dose of β -glucan (B-L β g) and, 23 fish with 100 µl of bacterin + high dose of β -glucan (B-H β g).

1.7.3 Aeromonas hydrophila

Aeromonas hydrophila, a fermentative rod-shaped, with a single polar flagella, is the most common bacteria present in all freshwater environments (Austin, 1999). This Gram-negative bacterium is an opportunistic pathogen that causes a wide range of diseases in fish (Nielsen et al., 2001). There are more than 120 serotypes of A. hydrophila described and each of them may show different growth in mucus and can produce different kinds of toxins. It does not cause problems in fish populations under normal conditions. Infection with A. hydrophila appears as a secondary infection, when fish are under environmental or physiological stress, injured or infected by other pathogens (Plumb et al., 1976; Pai et al., 1995 Fang et al., 2000). The disease caused by A. hydrophila is called motile aeromonad septicaemia (MAS) and its clinical signs in fish vary from tissue swelling, necrosis, ulceration and haemorrhagic septicaemia (Hazen et al., 1978a; Karunasagar et al., 1986; Aguilar et al., 1997; Azad et al., 2001; Figure 7). When it enters the body of its victim, it travels through the bloodstream to the first available organ. This bacterium can digest materials such as gelatin and haemoglobin. It produces aerolysin cytotoxic enterotoxin (ACT), toxin which posses multiple biological activities like ability to lyse red blood cells, destroy tissue and evoke a fluid secretory response in ligated intestinal loop models (Sha et al., 2002). Aerolisyn is an extracellular, soluble, hydrophilic protein that exhibits both hemolytic and cytolytic properties. Aerolysin binds to specific glycoprotein receptors on the surface of eucaryotic cells before inserting into the lipid bilayer and forms holes. It was shown that ACT caused levels of pro-inflammatory cytokines, such as TNF- α , IL - 1 β and IL - 6 but also that many bacterial toxins possess cytokine – inducing

activities and therefore the host pro-inflammatory cytokine response to ACT may contribute to the pathogenesis of *Aeromonas* infections (Chopra et al., 2000).



Figure 7. Koi carp infected with Aeromonas hydrophila with ulcer in the skin and muscle, and bleeding.

1.8 Objectives of the PhD study

This PhD study aimed to investigate the biological effect of the β -glucan as immune modulator of the mucosal surfaces in common carp (*Cyprinus carpio* L.) in relation to the wound healing process.

Thus, the aims of the current PhD thesis were:

- To examine, if previous infections had an impact on texture quality, and if these changes were detectable after recovery from bacterial infection.
- To create a model in controlled condition, where, without introducing the pathogen, we could mimic naturally occurring wounds, and examine the biological impact of two commercially available β-glucan enriched products on the wound healing process in common carp.
- Examine, if intravenous injection of β-glucan affects the skin mucosal immunity and can introduce an antibody change in *Cyprinus carpio* L.

Some references will be made to the mammalian system for explanation.

2 General discussion

Fast wound healing process and tissue regeneration is extremely important for all animals. In their natural environment fish are in intimate contact with a high amount of microorganisms and pathogens. Skin, together with mucosa forms the first line of defence against pathogens and is an essential protective barrier in aquatic organisms. Skin damage in fish caused e.g. by pathogens, such ectoparasites as Ichthyophthirius multifillis (Gonzalez et al., 2007) and lice (Krkosek et al., 2011), Gram-negative bacteria as Moritella viscosus (Ingerslev et al., 2010), as well as mechanical trauma reduce skin integrity and/or change mucus production, and can facilitates entry of pathogens into the underlying tissue and vascular system. Once the protective barrier is broken, the normal (physiologic) process of wound healing is immediately set in motion, and involves a complex series of interactions between different cell types, cytokine mediators, and the extracellular matrix (Singer and Clark, 1999; Kanter and Akpolat, 2008; Gillitzer and Goebeler 2001). The study of the interaction between the parasite and the fish host has been described by Gonzalez et al. (2007), however process of wound closure in a molecular perspective have so far not been described in details in common carp. Moreover, nowadays, number of biological compounds has been used to reduce the risk of diseases and improve fish welfare by enhancement the non-specific defence system. Although, among them, β -glucans are some of the most powerful and extensively investigated modulators of the immune system, their effect on skin mucosal immunity has not yet been addressed.

2.1 The best use of immune modulators: bath, injection or fed?

Immune modulators are widely used in aquaculture. One of the most frequent doubts is which way of administration is the best and how long the protection will last. The potential use of β -glucans as immune and growth stimulants in the large aquaculture sector has not been fully explored. Research on the use of immune modulators for the improving non-cellular immune response becomes more promising. The limited studies carried out, have been primarily restricted to the production cold water fish e.g. salmonids whilst the application of β -glucan to the large warm water fish and ornamental fish sectors has not been ascertained (Robertsen et al., 1990). Three different way of administration are available: injection, bath or oral. Depend on the purpose, the mode of application of immune modulator is very important and should be chosen with regard to labour input and stress factor. Second segment of hindgut in fish is a place where antigens up-take takes place (Rombout et al., 1985; Selvaraj et al., 2005) and, to a large extent, antibiotics and immune modulators are orally introduced to fish by incorporation into fed. In the present PhD thesis, we were interested in examinating mucosal immune response following direct stimulation with β -

glucan, and due to the fact that it remains unclear, under what changes β -glucan goes and subsequently triggers an immune reaction that can be detected in e.g. skin of fish (van der Marel et al., 2012), oral administration not addressed.

One of the aims of the present PhD study (**PAPER 2**) was on biological effect of β -glucans on wound healing process in common carp. Bath trial was an obvious choice for investigation if β -glucans can affect wound closure by "outside" stimulation, even though very limited information is available on bathing protocols and dosage, and all up-to date investigations were carried as short immersion in e.g. aerated glucan solution (Selvaraj et al., 2005). The idea was to keep wounded fish in constant and direct contact with β -glucans for two experimental weeks and monitor process of wound closure. By using bath treatment, it was also possible to investigate local and systemic immune response due to DAMPs/PAMPs stimulation. The results we obtained clearly showed that beta glucan enriched bath directly affected wound contraction dynamics and faster closure was observed.

Due to fact that injected substance can reach all organs, injection was showed to be the most effective route in immune modulants introduction (Yano et al., 1991; Chen and Ainsworth 1992). In the current thesis, intravenous injection of β -glucan was introduced (**PAPER 3**) to examine if by being present in the circulation system, glucan can cause any biological effect. Our aim was on if β -glucan injection into vein affects antibody response in the circulation and skin as well. Moreover, two different doses of β -glucan were used to investigate any dose-dependent manner or toxic effect. Since injection itself is very stressful for fish, stress response, including cortisol level and expression of Hsp70, was examined. Obtained results showed that effect of direct β -glucan stimulation by intravenous injection in common carp is rigorously correlated with dosage. Moreover, high amount of particulate β -glucan introduced to circulation system of carp elevates cortisol level, and at high dose can be toxic for fish. Furthermore, intravenous administration of β -glucan affects immunoglobulin expression in carp skin with immunoglobulin class change process, where presence of main fish immunoglobulin IgM is overtaken by recently described, mucosal immunoglobulin IgZ1. IgZ1 is being considered a mucosal immunoglobulin, and its high expression in skin of fish could increase efficiency in fighting pathogens invading skin layer.

2.2 The use of internal control samples: local vs. systemic reaction

Many years ago, Wolk and Danon (1985) have emphasized that potential application of glucans to promote wound healing in humans and other heterogenic populations of animals, would require an experimental model in which a wound on one side of the animal could be compared to a symmetrical wound on the other side of the same animal. Recently, Ingerslev et al., (2010) used a

model where during sterile damage in rainbow trout, internal control samples were collected as well. It turned out to be a powerful model to examine local versus systemic response in the fish. Presented results clearly showed that there were no significant differences in expression level between the internal controls from damaged fish in comparison to non-injured fish, and damaging a restricted area in the fish muscle resulted in response in that specific area. In our study, internal controls of non-damaged muscle tissue and skin have been taken. This allowed us to obtain more details on changes taking place in wounded and non-wounded areas due to DAMPs/PAMPs receptor(s) stimulation, and subsequently, information about the local versus systemic response in carp. In addition, no significant changes have been observed in non-wounded groups, and the absence of marked differences on the respiratory burst activity in head-kidney cells supports the idea of a localized immune response to the site of injury.

2.3 Inflammatory response during wound healing process: DAMPs and PAMPs stimulation

Wound healing can be defined as the physiology by which the body replaces and restores function to damaged tissues (Tortora and Grabowski, 1996). The classic model of wound healing is divided into three sequential, yet overlapping, phases: (1) inflammatory, (2) proliferative and (3) remodelling, and a set of complex biochemical events takes place in a closely orchestrated cascade to repair the damage (Singer and Clark, 1999; Kanter and Akpolat, 2008; Gillitzer and Goebeler 2001). By wounding procedure that was performed to mimic naturally occurring wounds, danger molecules were released from injured cells of skin and muscle of common carp (PAPER 2). Moreover, bath in water supplemented with β-glucan was PAMPs source. DAMPs activate immune response, but danger signals such RNA/DNA or heat shock proteins are as well involved in myogenesis and have high constitutive expression in the heart and muscles in fish (Sass et al., 1996). Immediate increase in IL-1β, IL-6 and IL-8 cytokines day post-damage in muscle collected from wounded-control group is likely to be the inflammatory response caused by the release of DAMPs e.g. heat shock proteins. Interestingly, in skin samples from wounded-control fish, different pattern of pro-inflammatory cytokine expression has been noted. No or low IL-1B, IL-6 and IL-8 expression has been detected one day after damage, however no gene expression was investigated at earlier time points. It is possible that stored IL-1 β , IL-6 and IL-8 were quickly secreted (hours) in response to injury and subsequently, new amount of these cytokines needed to be produce. The inversed pattern of gene expression in skin/muscle tissue has been seen in the βglucan supplemented group as well, with change in expression three days post wounding. In mammals, DAMPs and PAMPs can bind to the same receptors and therefore they may simultaneously effect when combined (Holtick et al., 2011). Receptors stimulation with DAMPs/PAMPs resulted in local expression of IL-1β, IL-6 and IL-8 genes, and that local

inflammatory response differences in expression pattern dependent on stimulation by DAMPs alone or DAMPs/PAMPs combination (Figure 8). In the injection experiment (PAPER 3) only IL- 1β expression was investigated. In fact, present results show no significant changes in IL-1 β expression at early phase of experiment. It can be explained by the inhibitory role of cortisol, seing that glucocorticoids control cytokine expression in several ways. One of them is interaction with NF-kB transcription factor, preventing it to attach to kB gene responsive elements (McKay and Cidlowski, 1999) and subsequently down-regulates expression of IL-1 β , which expression is regulated via the NF-kB pathway (Engelsma et al., 2001; McKay and Cidlowski, 1999). Additionally, not only the transcription of IL-1 is down-regulated by GC, also the stability of IL-1b mRNA is decreased by these steroids (Lee et al., 1988). On the other hand, we cannot rule out fact that, IL-1 β is main pro-inflammatory cytokine, which reaches peak within minutes after stimulation, so measurement 96 hours after stimulation does not give clear picture. But then again, we observe high and significant up-regulation 12 days post i.v. injection. At that time, cortisol level reaches basal line, and expression of IL-1ß starts. Moreover, fish from B-Hßg injected group appeared to have skin and muscle damage (personal observation) which could enhance IL-1ß expression in response to DAMPs. In mammals, corticosteroids have inhibitory effect on monocytes and macrophages (Baybutt and Holsboer, 1990), and that could explain no changes in IL-1 β expression, as macrophages are main source of IL-1 β .



Figure 8. Summary of the expression results in skin and muscle from bath and injection experiment. Sets of one, two and three arrows indicate a low (1-5 fold), intermediate (5-25 fold) and high (> 25 fold) differences in gene expression respectively. NS indicates no samples were tested. Results compiled from **PAPER 2** and partly from **PAPER 3**.

2.4 β-glucan immune modulator effect on non-immune cells

Skin injury (DAMPs) activates mucosal immune cells as well as epithelial resident cells. It has been proven, not only "typical" immune cells of haematopoietic origin, such as neutrophils or macrophages are involved in wound closure and cytokines production (Ingerslev et al., 2010). Chen and co-workers (2005) have described human fibroblasts as the non-conventional cell type having immune regulation capabilities. Recent studies in human have shown that fibroblasts can not only respond to cytokines, but also generate these signals (IL-1β, IL-4, IL-6, IL-8, IL-10, IL-12, TNF-α and IFN- γ) and influence the behaviour of immunocompetent cells trafficking to the injury site (Chen et al., 2005; Ruiz et al., 2007). In fish, several fibroblast cell lines have been established and characterised, and Ingerslev et al., (2010) reported that fish fibroblasts are susceptible to DAMPs (Collet and Secombes, 2001; Ossum et al., 2004). In the wounding experiment (PAPER 2), high up-regulation of IL-1ß and IL-6 in skin samples from wounded-MacroGard fish three days post injury could be explained by the ability of "moving" fibroblasts to express genes of immune functions before the more potent immune cells have been recruited. Presence of DAMPs/PAMPs could potentiate immune-like performance of fibroblasts, as well as having a direct effect on the resident cells present in skin, such as epithelial cells, resident macrophages and lymphocytes. Glucans' ability for direct stimulation of human fibroblasts has been shown by Wei et al., (2002).

Another cell type that was shown to be involved in wound healing process is goblet cells (Geggel et al., 1984; Kanter and Akpolat, 2008). Goblet cells, cells responsible for the production and maintenance of the protective mucus blanket, are exclusively lodged in epithelia. Due to goblet location, we expected that bath supplemented with beta glucans would have an effect on their function in common carp, and subsequently changes in Muc5b expression should be detected. High Muc5b up-regulation three and 14 days post-wounding has, however not been restricted to the wound area, but was a general response of the skin mucosa to the injury. Goblet cells differentiation and functioning in mammalian system is affected by mucosal immunity and gelforming mucins can be regulated by inflammatory cytokines such as interleukins (IL-1 β , -6, -17), tumour necrosis factor α (TNF- α) or exogenous factors (Ciacci et al., 2002; Roussel and Delmotte, 2004; Chakraborty et al., 2010; van der Marel et al., 2012) and similar mechanism could be present in fish. Therefore, during wound closure in common carp, high local inflammation and IL-1ß expression could hamper Muc5b expression at the site of injury. Our results can be compared to work presented by van der Marel co-workers (2010) where carp kept in water with increased bacterial load did not cause clinical symptoms, however rapid skin mucosa respond have been seen, even if the bacteria involved were considered to be non-pathogenic. In vertebrates, β -glucans and lipopolysaccharide (LPS) work through the same receptors, hence, the mechanism by which LPS,

MacroGard and 6.3 kDa oat fiber stimulate Muc5b expression might be alike. However, in humans, cells may differ in the density of mucin granules, suggesting a diversity of the mucins synthesized even by an individual cell (Roussel and Delmotte, 2004; Kesimer et al., 2010; Sheehan and Thornton 2000). Considering that based on mucins content, several different goblet cell types have been described, we cannot assure, that beta glucans bath did not have any effect on expression of other mucin granules present in mucous or goblet cells (Fletcher et al., 1976).

2.5 Skin – perfect tissue to follow β -glucan actions?

Skin together with mucus blanket is the first line of defence. Carps skin is a complex structure composed of the outer stratified, relatively thin epithelium (epidermis) and the inner layer (dermis) composed of dense fibrous connective tissue, which is attached to the underlying muscles, by a layer of loose connective tissue (Figure 9). However, in contrast to terrestrial vertebrates, cells from fish skin are non-keratinised and consequently metabolically active (Bullock and Roberts, 1974). During two conducted experiments, bath and injection trial, effects of used immune modulant were detected in skin. In the bath treatment (**PAPER 2**), direct and constant contact of β -glucans caused positive effect on wound closure. In the injection trial (**PAPER 3**), indirect effect was observed. Injected into vein β -glucan reached all internal organs, as well as skin.

Skin and mucues blanket are the first layers between fish and an environment, and beside mechanical barrier function, immune system was developed as well in mucosal compartment. Powerful mucosal immune system increases efficiency in fighting pathogens invading the host. One of the arms that could be used for that purpose is antibody. The output evolution, immunoglobulins diversified into several isotypes with specialized roles in innate and adaptive responses, in the systemic and mucosal compartments. In teleost, the most primitive vertebrates with immunoglobulins, due to lack of evidences of immunoglobulin specialization, IgM was regarded as the only functional antibody in systemic and mucosal compartments. However, in 2005, a new immunoglobulin isotype, called IgZ was found in zebrafish (IgZ - in zebrafish; also called IgT - teleostean; Solem and Stenvik, 2006; Danilova et al., 2005). The existence of two distinct B cell lineages, with mutuall expression either IgM or IgZ, was suggested. Moreover, the possible class-switch recombination process between IgM and IgT was rouled out (Danilova et al., 2005; Hansen et al., 2005). However, in 2010, Zhang and co-workers provided evidence for the existence of a previously unrecognized B cell lineage that expressed only IgT. In the same study, IgT⁺B cells have been showed to be potent phagocytic cells with ability for killing rainbow trouts' gut parasite Ceratomyxa shasta. In fish that survived infection, authors reported the substantial accumulation of IgT⁺ cells in the gut concomitantly with considerably increased (~51-fold) gut

mucus IgT. In contrast, the number of IgM⁺ B cells and concentration of IgM did not change in surviving fish. Further, substantial upregulation (~733-fold) of IgT gene expression was detected in gut of these fish, with almost negligible upregulation of IgM expression. Such a high expression of IgT in gut and protective role against gut parasite indicates, that it is the first nontetrapod immunoglobulin specialized in mucosal immunity (Zhang et al., 2010). In fish, skin, mucus, gills and gut together, form mucosal immune system. In present PhD thesis, IgM and IgZ1 antibody response following intravenous injection with different doses of β-glucan was examinated in skin and head kidney of common carp (PAPER 3). We observed high expression of IgZ1 in skin 12 and 23 days post injection (injection with B-L β g; ~390- and ~310-fold respectively) with no significant changes in IgM expression (Figure 10). Taking under consideration mucosal functions of IgT(Z) reported by Zhang el al. (2010) and further by Ryo et al., (2011), high IgZ1 expression detected in the injection experiment (PAPER 3) leads to the conclusion that $IgT(Z)^+$ B cells are major B cells in skin of common carp, with low cell count for IgM⁺ B cells. Zhang et al., (2010) found the $IgT(Z)^+$ B cell population constituted 16-28% of all trout B cells in the blood, spleen, head kidney and peritoneal cavity, while in gut they represented the main B cell subset. Results obtained from the injection experiment show that IgZ1 expression was significantly higher in skin than in common carp head kidney, suggesting that either $IgT(Z)^+$ B cells have the majority among cells in carp skin, or β -glucan injection enhanced proliferation of these cells (Figure 9). IgM⁺B cells are highly phagocytic and have intracellular killing abilities (Li et al., 2006). However Zhang et al., (2010) compared the phagocytic ability of $IgT(Z)^+$ and IgM^+ B cells, and they seemed to be alike, as well as capacities to proliferate and secrete immunoglobulin in response to microbial stimulation. Even so, in our injection experiment, only IgT(Z) gene expression was upreguated, with no changes in IgM expression (Figure 10). That draws the conclusion that only $IgT(Z)^+$ cells are present in the skin, as mention earlier, or that phagocytic activity of two distinct B cell lineages differs. Moreover, $IgT(Z)^+$ B cells seem to respond to β -glucan stimulation, which might confirm earlier findings that $IgT(Z)^+$ B cells are able to ingested and kill bacteria intracellularly (Zhang et al., 2010). Furthermore, $IgZ(T)^+$ B cells response to β -glucan might suggests presence of the receptor(s) through which, any actions with β -glucan will be more pronounced in fish skin. Hence, to obtain more data, studies on the fish cell line should be carry on.

Throughout the bath and intravenous injection, in spite of administration route, effect of β -glucan stimulation was always observed in skin of common carp. That raises a new hypothesis about β -glucan actions as immune modulant in fish immunity. β -glucans as polysaccharides that contain only glucose as structural components are "part" of fungi, and fungal infection in fish always occurs on skin surfaces. To clarify this hypothesis further studies on fish mucosal immunity following immune modulation are needed.



Figure 9. Schematic model of common carp skin. Fish skin is divided in to layers, the epidermis and dermis, which are separated by basal membrane. The outer surface of fish skin is protected by the mucosal blanket, formed by goblet cells (known also as mucous cells). In a healthy skin, many leucocytes can be found. Scales are originated from the dermal layer. Based on recent studies, $IgZ(T)^+$ B cells might be the major B cells present in skin.



Figure 10. Summary of the antibody expression results in skin and head kidney from injection experiment (paper 3). Sets of one, two and three arrows indicate a low (1-5 fold), intermediate (5-25 fold) and high (> 25 fold) differences in gene expression respectively. NS indicates no samples were tested.

3 Conclusions

- 1. Fish diseases have an impact on product quality-associated parameters, as the structural changes can be associated with the development of scars as a consequence of the regeneration process.
- 2. MacroGard and 6.3 kDa oat fiber, promote wound healing process in common carp, however with higher wound closure ratio in group treated with MacroGard.
- 3. The immunological and regenerative response following stimulation with DAMPs (wounding procedure) and PAMPs (β-glucan) in controlled conditions, without the exposure to pathogens results in an inflammatory response by activating main pro-inflammatory cytokines such as IL-1β, IL-6 and IL-8. Moreover, these changes have a local character and depend on receptor(s) mediated stimulation either with DAMPs alone or DAMPs/PAMPs combination.
- 4. DAMPs/PAMPs stimulation affected the inflammatory phase in wound healing process in common carp, and therefore bathing in β-glucans, specifically MacroGard and 6.3 kDa oat fiber, enhanced the wound closure in common carp.
- 5. Fish skin injury introduces DAMPs and activates cells present in mucosal layers, these of the immune function, as well as resident cells such as fibroblasts and goblet/mucous cells. High local inflammation and IL-1β expression could hamper Muc5b expression at the site of injury, which might explain why Muc5b expression was not higher in MacroGard treated group as would expect. Moreover, stimulation with DAMPs/PAMPs possibly increased the immune-like performance of fibroblasts.
- 6. Effect of direct β-glucan stimulation by intravenous injection in common carp is rigorously correlated with dosage: high amount of particulate β-glucan introduced to circulation system of carp elevates cortisol level, and at high dose can be toxic for fish. Strong stress reaction caused by handling and injection could be additionally boosted with high dose of injected β-glucan, which is recognized as pathogen associated molecular pattern (PAMP) and overwhelmed the immune system of carp.
- 7. Intravenous administration of β -glucan affects immunoglobulin expression in carp skin with immunoglobulin class change, where presence of main fish immunoglobulin IgM is overtaken by recently described, mucosal immunoglobulin IgZ1. High expression of IgZ(T) in skin might indicate that IgZ(T)⁺ B cells lineage posses receptor, through which stimulation with β -glucan occures.

8. Intravenous injection of *Aeromonas hydrophila* bacterin alone or together with β-glucan does not elevates production of antibodies against *A.hydrophila* if fish were earlier primed to *A. hydrophila*. Slight elevation in specific antibody level could be a short-term alert in response to injected PAMPs (bacterin and beta glucan) and temporarily affect antibody production.

4 Perspective for future research

The work conducted in my thesis has provided new insight into quality related issue in relation to previous diseases. Further, biological impact of β -glucan was investigated in relation to wound healing process common carp. To continue with investigation of immune modulators effect on fish immune system, intravenous injection experiment was performed. Nowadays, fish mucosal immunity is dynamically investigated area and new captivating opportunities arise. In continuation of these studies, several interesting subjects are worth examining:

(1) Fluorescent-labelled antibodies against β -glucans – a perfect tool for β -glucans localization?

Although the mechanism, by which β -glucan triggers immune response in teleost fish remains unknown, all three ways of introducing immune modulators, bath, injection and oral, are proved to stimulate immune system in common carp. Since mucosal immunity is dynamically developing area, new tools are opportunities arise. Monoclonal antibodies were raised against (1->3,1->4)beta-glucan and could be useful tool in the quantitation and immunocytochemical location of (1->3,1->4)-beta-glucans. Fluorescent-labelled antibodies against β -glucans could show how deep β glucan-enriched products such as MacroGard, penetrate wounded tissue. We have showed that bath in MacroGard and 6.3 kDa oat fiber enhanced the wound healing process when compared to nontreated group, but immune response differs in skin and muscle tissue. Usage of fluorescent-labelled antibodies against β -glucans could contribute to the discussion if there is any correlation between size of immune modulants and their variable immune potency. Similar issue is addressed in fed trials. It remains unclear, under what changes β -glucan goes in the gut of fish, and subsequently triggers an immune reaction that can be detected in e.g. skin of fish.

(2) Further investigation of IgM/IgZ1 class change in skin of common carp.

Among circulating lymphocytes in fish, B cells count for approximately 70%, however some resident B cells can be found e.g. in skin of common carp. Immunoglobulin M (IgM) and Z1 (IgZ1) are mutually exclusive on B cells. We showed high expression of IgZ1 in skin accompanying by no detection for IgM expression. However, it was showed that IgM is present in goblet cells in rainbow trout skin. It was also showed that no immunoglobulin class switching takes place in fish. Goblet cells staining against IgM and IgZ1 could provide more information about content of goblet cells before, during and after immune stimulation. Moreover, new assay in which separate examination of IgM/IgZ1 expression in skin resident B cells and goblet cells should be established, as recent studies have proven the existence of two distinct B cell lineages. Furthermore, seems like IgZ(T)⁺B cells are the main B cell lineage present in carp skin and responds to immune modulation.

Hence, further research should focuse on work with cell lines, looking for receptor(s) of interest through which β -glucan triggers the immune modulation in fish skin.

(3) Does β -glucan work in dose dependent manner?

Immune stimulants are valuable alternative for the control of fish diseases and may be useful in fish culture, as drug choices for the treatment of common infectious diseases are becoming increasingly limited and expensive and, in some cases, unavailable due to the emergence of drug resistance in bacteria and fungi. As mention in the introduction of this PhD thesis, three routes of administration of the immune modulants are possible. The most effective method of administration of immune modulants to fish is by injection. Many studies have reported so-called "optimal dose" for fish injection, however, the dosage range is wide. Oral and immersion methods have also been successfully used in fish, however, their efficacy decreases with long-term administration. Moreover, still little information about dosage and protocols is available, especially for bath treatments. Since dose-dependent side effects of immune modulants have not been well studied in fish, it is interesting to obtain more data on potential dose-related effects of injected β -glucan, including the timing, dosages, method of administration and the physiological condition of fish.

5 Acknowledgements

The work of this thesis was carried out at Technical University of Denmark during 2009-2012. Financial support was provided by European Community's Seventh Framework Programme (FP7/2007-2013) under Grant agreement number PITN-GA-2008-214505.

First of all I would like to express my gratitude to my supervisor Associate Professor Michael E. Nielsen for providing me the exciting opportunity to study in Denmark. It was not always an easy task, but I would like you to know I have appreciated your help, going way behind science, support, guidance and patience. Thank you also for your good company at the conferences and giving me opportunity to present my work. Thank you for reminding that this is a holistic PhD. These years have been worth it all©

Ogromne podziekowania dla mojej rodziny: Mamy, Taty, Dziadka i Brata za niekonczaca sie cierpliwosc, godzinne rozmowy i wsparcie. Za bycie zawsze po mojej stronie i pozwolenie abym uczyla sie na wlasnych bledach. Za pozwolenie mi wybrania wlasnej sciezki, mimo ze ta zaprowadzila mnie daleko od Nich. DZIEKUJE!

To Axel, "...word can't say what love can do..." for being with me during my ups and downs and taking care of me, for giving my courage to "just say it".

I would also like to thank my co-workers, the NEMO people, for the company during these fruitful years. Thank you for many discussion, challenges, brainstorms, and time together in the office, conferences, bars[©] For all fun we had during NEMO meetings.

Further, I would like to thank my friends "behind the scene" here in Denmark but also in Poland and England. For many, many, many chats and "drinking" on skype. Your presence and endless energy means a lot! Thank you for being there for me!

And last but not least: my huge gratitude to Geert Wiegertjes. For telling me about NEMO and for not willing to keep me in The Netherlands. You said "You will like it" and you were right[®]

Dominika A.Przybylska

June 2012

6 References

Abdelkhalek NK, Komiya A, Kato-Unoki, Somamoto T and Nakao M. Molecular evidence for the existence of two distinct IL-8 lineages of teleost CXC-chemokines. Fish and Shellfish Immunology 2009; 27:763–767.

Abelli L, Picchietti S, Romano N, Mastrolla L and Scapigliati G. Immunohistochemistry of gut-associated lymphoid tissue of the sea bass *Dicentrarchus labrax* (L.). Fish and Shellfish Immunology 1997; 7: 235–245.

Afonso A, Silva J, Lousadai S, Ellis AE and Silva MT. Uptake of neutrophils and neutrophilic components by macrophages in the inflamed peritoneal cavity of rainbow trout (*Oncorhynchus mykiss*). Fish and Shellfish Immunology 1998; 8: 319–338.

Aguilar A, Merino S, Rubires X and Tomas JM. Influence of osmolarity on lipopolysaccharides and virulence of *Aeromonas hydrophila* serotype O:34 strains grown at 37°C. Infection and Immunity 1997; 65: 1245-1250.

Ainsworth AJ. A β-glucan inhibitable zymosan receptor on channel catfish neutrophils. Veterinary Immunology and Immunopathology 1994; 41: 141-152.

Airaksinen S, Råbergh CMI, Lahti A, Kaatrasalo A, Sistonen L and Nikinmaa M. Stressor-dependent regulation of the heat shock response in zebrafish, *Danio rerio*. Comparative Biochemistry and Physiology 2003; 134: 839–846.

Alberts B. Molecular biology of the cell. New York: Garland Science, 2008.

Akramiene D, Kodrotas A, Didziapetriene J and Kevelaitis E. Effects of beta-glucans on the immune system. Medicina Lithuania 2007; 43: 597-606.

Aoki T, Takano T, Santos MS, Kondo H and Hirono I. Molecular Innate Immunity in Teleost Fish: Review and Future Perspectives. Fisheries for Global Welfare and Environment, 5th World Fisheries Congress 2008, 263–276.

Austin B. Emerging bacterial fish pathogens. Biulletin of the European Association of Fish Pathologists 1999; 19: 231-234.

Azad IS, Rajendran KV, Rajan JJS, Vijayan KK and Santiago TC. Virulence and histopathology of Aeromonas hydrophila (SAH 93) in experimentally infected tilapia, *Oreochromis mossambicus* (L.). Journal of Aquaculture in the Tropics 2001; 16: 265-275.

Balon EK. Origin and domestication of the wild carp, *Cyprinus carpio*: from Roman gourmets to the swimming flowers. Aquaculture 1995; 129: 3-48.

Bansil R, Stanley HE and Lamont JT. Mucin Biosynthesis. Annual Review of Physiology 1995; 57: 635-657.

Barton BA. Stress in Fishes: A Diversity of Responses with Particular Reference to Changes in Circulating Corticosteroids. Integrative and Comparative Biology 2002; 42:517–525.

Basu N, Kennedy CJ and Iwama GK. The effects of stress on the association between hsp70 and the glucocorticoid receptor in rainbow trout. Comparative Biochemistry and Physiology 2003; 134: 655–663.

Baybutt HN and Holsboer F. Inhibition of macrophage differentiation and function of cortisol. Endocrinology 1990; 127: 476-480.

Belperio JA, Keane MP, Arenberg DA, Addison CL, Ehlert JE, Burdick MD and Strieter RM. CXC chemokines in angiogenesis. Journal of Leukocyte Biology 2000; 68: 1-8.

Bengten E, Clem LW, Miller NW, Warr GW and Wilson M. Channel catfish immunoglobulins: Repertoire and expression. Developmental and Comparative Immunology 2006; 30: 77-92.

Berdal M, Appelbom HI, Eikrem JH, Lund Å, Zykova S, Busund L-T, Seljelid R and Jenssen T. Aminated – β -1,3-D-glucan improves wound healing in diabetic db/db mice. Wound Repair and Regeneration 2007; 15: 825–832.

Berman J, Hsu K and Look AT. Review: Zebrafish as a model organism for blood diseases. British Journal of Haematology, 2003; 123: 568–576.

Bianchi ME and Manfredi AA. High-mobility group box 1 (HMGB1) protein at the crossroads between innate and adaptive immunity. Immunological Reviews 2007; 220: 35–46.

Bird S, Zou J, Savan R, Kono T, Sakai M, Woo J and Secombes CJ. Characterisation and expression analysis of an interleukin 6 homologue in the Japanese pufferfish, *Fugu rubripes*. Developmental and Comparative Immunology 2005; 29: 775–789.

Bird S, Zou J and Secombes CJ. Advances in fish cytokine biology give clues to the evolution of a complex network. Current Pharmaceutical Design 2006; 12: 3051-3069.

Bird S, Zou J, Wang T, Munday B, Cunningham C and Secombes CJ. Evolution of interleukin-1β. Cytokine and Growth Factor Reviews 2002; 13: 483–50.

Bohn JA and BeMiller JN. $(1\rightarrow 3)$ - β -d-Glucans as biological response modifiers: a review of structure-functional activity relationships. Carbohydrate Polymers 1995; 28: 3-14.

Bols NC, Brubacher JL, Ganassin RC and Lee LEJ. Ecotoxicology and innate immunity in fish. Developmental and Comparative Immunology 2001; 25: 853-873.

Brandtzaeg P and Pabst R. Let's go mucosal: communication on slippery ground. Trends in Immunology 2004; 25: 570-577.

Brattgjerd S, Evensen Ø and Lauve A. Effect of injected yeast glucan on the activity of macrophages in Atlantic salmon, *Salmo salar* L., as evaluated by in vitro hydrogen peroxide production and phagocytic capacity. Immunology 1994; 83: 288-294.

Brinkmann V, Reichard U, Goosmann C, Fauler B, Uhlemann Y, Weiss DS, Weinrauch Y and Zychlinsky A. Neutrophil Extracellular Traps Kill Bacteria. Science 2004; 303: 1532-1535.

Brown GD and Gordon S. Fungal beta-glucans and mammalian immunity. Immunity 2003; 19: 311-315.

Brown SL, Riehl TE, Walker MR, Geske MJ, Doherty JM, Stenson WF and Stappenbeck TS. Myd88dependent positioning of Ptgs2-expressing stromal cells maintains colonic epithelial proliferation during injury. The Journal of Clinical Investigation 2007; 117: 258-269.

Buchmann K. Immune mechanisms in fish skin against monogeneans – a model. Folia Parasitologica 1999; 46: 1-9.

Bullock AM and Roberts RJ. The dermatology of marine teleost fish. The normal integument. Oceanography Marine Biology and Animal Review 1974; 13: 383-411.

Castellana B, Iliev DB, Sepulcre MP, MacKenzie S, Goetz FW, Mulero V and Planas JV. Molecular characterization of interleukin-6 in the gilthead seabream (*Sparus aurata*). Molecular Immunology 2008; 45: 3363-3370.

Castro R, Couso N, Aobach A and Lamas J. Effect of different β-glucans on the respiratory burst of turbot (*Psetta maxima*) and gilthead seabream (*Sparus aurata*) phagocytes. Fish and Shellfish Immunology 1999;

9: 529–541.

Chadzinska M, Kolaczkowska E, Seljelid R and Plytycz B. Morphine modulation of peritoneal inflammation in Atlantic salmon and CB6 mice. Journal of Leukocyte Biology 1999; 65: 590-596.

Chadzinska M, Leon-Kloosterziel KM, Plytycz B, Verburg-van Kemenade BML. In vivo kinetics of cytokine expression during peritonitis in carp: Evidence for innate and alternative macrophage polarization. Developmental and Comparative Immunology 2008; 32: 509–518.

Cbantanachookhin C, Seikaiband T and Tanaka M. Comparative study of the ontogenyof the lymphoidorgans in three species of marine fish. Aquaculture 1991; 99: 143-155.

Chen D and Ainsworth AJ. Glucan administration potentiates immune defence mechanisms of channel catfish *Ictalurus punctatus* Rafinesque. Journal of Fish Diseases 1992; 15: 295-304.

Chen B, Tsui S and Smith TJ. IL-1 β Induces IL-6 Expression in Human Orbital Fibroblasts: Identification of an Anatomic-Site Specific Phenotypic Attribute Relevant to Thyroid-Associated Ophthalmopathy. Journal of Immunology 2005; 175: 1310-1319.

Cho JH, Fraser IP, Fukase K, Kusumoto S, Fujimoto Y, Stahl GL, and Ezekowitz RAB. Human peptidoglycan recognition protein S is an effector of neutrophil-mediated innate immunity. Blood 2005; 106: 2551-2558.

Chopra AK, Xu X-J, Ribardo D, Gonzalez M, Kuhl K, Peterson JW and Houston CW. The cytotoxic enterotoxin of Aeromonas hydrophila induces proinflammatory cytokine production and activates arachidonic acid metabolism in macrophages. Infection and Immunity 2000; 68: 2808-2818.

Ciacci C, DiVizio D, Seth, R Insabato G, Mazzacca G, Podolsky DK and Mahida YR. Selective reduction of intestinal trefoil factor in untreated coeliac disease. Clinical and Experimental Immunology 2002; 130: 526–531.

Chakraborty S, Bonthu N, Swanson BJ and Batra SK. Role of mucins in the skin during benign and malignant conditions. Cancer Letters 2010; 301: 127-141.

Chrousos GPI and Gold PW. The concepts of stress and stress system disorders. JAMA: Journal of the American Medical Association 1992; 267: 1244-1252.

Clark TG, Lint T-L, AND Dickerson HW. Surface antigen cross-linking triggers forced exit of a protozoan parasite from its host. Immunology 1996; 93: 6825-6829.

Cohan VL, Scott AL, Dinarello CA and Prendergast RA. Interleukin-1 Is a Mucus Secretagogue. Cellular Immunology 1991; 136: 425-434.

Cole P. The damaging role of bacteria in chronic lung infection. Journal of Antimicrobial Chemotherapy - Supplement 1997; 40: 5–10.

Collet B and Secombes CJ. The rainbow trout (*Oncorhynchus mykiss*) Mx1 promoter: Structural and functional characterization. European Journal of Biochemistry 2001; 268: 1577-1584.

Costa MM, Maehr T, Diaz-Rosales P, Secombes CJ and Wang T. Bioactivity studies of rainbow trout (*Oncorhynchus mykiss*) interleukin-6: Effects on macrophage growth and antimicrobial peptide gene expression. Molecular Immunology 2011; 48: 1903–1916.

Cox CM, Sumners, LH Kim S, McElroy AP, Bedford MR and Dalloul RA. Immune responses to dietary βglucan in broiler chicks during an *Eimeria* challenge. Poultry Science 2010; 89:2597–2607. Cui X, Shin H, Song C, Laosinchai W, Amano Y, Brown RM. A putative plant homolog of the yeast β -1,3-glucan synthase subunitFKS1 from cotton (*Gossypium hirsutum* L.) fibers. Planta 2001; 213: 223-230.

Dalmo RA, Ingebrigtsen K and Bøgwald J. Non-specific defence mechanisms in fish, with particular reference to the reticuloendothelial system (RES). Journal of Fish Diseases 1997; 20: 241–273.

Dalmo RA and Seljelid R.The immunomodulatory effect of LPS, laminaran and sulphated laminaran [β (l,3)-D-glucan] on Atlantic salmon, *Salmo salar* L., macrophages *in vitro*. Journal of Fish Diseases 1995; 18: 175-185.

Danilova N, Bussmann J, Jekosch K and Steiner LA. The immunoglobulin heavy-chain locus in zebrafish: identification and expression of a previously unknown isotype, immunoglobulin Z. Nature Immunology 2005; 6: 295-302.

Davidson GA, Lin S-H, Secombes CJ and Ellis AE. Detection of specific and "constitutive" antibody secreting cells in the gills, head kidney and peripheral blood leukocytes of dab (*Limanda limanda*). Veterinary Immunology and Immunopathology 1997; 58: 363-374.

Dean DB, Whitlow ZW and Borski RJ. Glucocorticoid receptor upregulation during seawater adaptation in a euryhaline teleost, the tilapia (*Oreochromis mossambicus*). General and Comparative Endocrinology 2003; 132: 112–118.

Deane ED, Zhou L and Woo NYS. Effects of mitogenic hormones on HSP70 expression in a silver sea bream fibroblast cell line and a primary macrophage preparation. General and Comparative Endocrinology 2007; 152: 183–188.

Deivasigamani B. Structure of immune organ in edible catfish, *Mystus gulio*. Journal of Environmental Biology 2007; 28: 757-764.

Dekker J, Rossen JWA, Buller HA and Einerhand, AWC. The MUC family: An obituary 2002; 27: 126-131.

DiLuzio NR and Williams DL. Protective effect of glucan against systemic *Staphylococcus aureus* septicemia in mormal and leukemic mice. Infection and Immunity 1978; 20: 804-810.

Dhabhar FS and McEwen BS. Stress-Induced Enhancement of Antigen-Specific Cell-Mediated Immunity. The Journal of Immunology 1996; 156: 2608-2615.

dos Santos NMS, Taverne-Thiele JJ, Barnes AC, Van Muiswinkel WB,. Ellisae and Rombout JHWM. The gill is a major organ for antibody secreting cell production following direct immersion of sea bass (*Dicentrarchus labrax*, L.) in a *Photobacterium damselae* ssp. piscicida bacterin: an ontogenetic study. Fish and Shellfish Immunology 2001; 11: 65-74.

Dranoff G. Review: Cytokines in cancer pathogenesis and cancer therapy. Nature Reviews 2004; 4: 11-22.

Duffy LK, Scofield E, Rodgers T, Patton M and Bowyer RT. Comparative baseline levels of mercury, Hsp 70 and Hsp 60 in subsistence fish from the Yukon-Kuskokwim delta region of Alaska. Comparative Biochemistry and Physiology 1999; 124: 181-186.

Duncan PL and Klesius PH. Dietary immunostimulants enhance nonspecific immune responses in channel catfish but not resistance to Edwardsiella ictaluri. Journal of Aquatic Animal Health 1996; 8: 241-248.

Ellis, A.E., R.J. Roberts and P. Tytler: The anatomy and physiology of teleosts. In: Fish pathology (Ed.: RJ Roberts). Bailliere Tindall, London, UK 1989; 13-55.

Engelsma MY, Hougee S, Nap D, Hofenk M, Rombout JHWM, van Muiswinkel WB, Verburg-van Kemenade BML. Multiple acute temperature stress affects leucocyte populations and antibody responses in common carp, *Cyprinus carpio* L. Fish and Shellfish Immunology 2003; 15: 397-410.

Engelsma MY, Huising MO, van Muiswinkel WB, Flik G, Kwang J, Savelkoul HFJ and Verburg-van Kemenade BML. Neuroendocrine–immune interactions in fish: a role for interleukin-1. Veterinary Immunology and Immunopathology 2002; 87: 467-479.

Enoch S and Price P. Cellular, molecular and biochemical differences in the pathophysiology of healing between acute wounds, chronic wounds and wounds in the aged. Worlds wide wounds 2004.

Erdogan O, Atamanalp M, Sisman T, Aksakal E and Alak G. Effects of 2,2-dichlorovinyl dimethyl phosphate (DDVP) on Hsp70 gene expression in rainbow trout. Isreali Journal of Aquaculture Bamidgeh 2007; 59: 230-234.

Espenes A, Press CML, Dannevig BH and Landsverk T. Immune-complex trapping in the splenic ellipsoids of rainbow trout *(Oncorhynchusmykiss)*. Cell Tissue Research 1995; 282: 41-48.

Fader SC, Zhongmo YU and Spotila JR. Seasonal variation in heat shock proteins (hsp70) in stream fish under natural conditions. Journal of Thermal Biology 1994; 19: 335-341.

Fang HM, Ling KC, Ge R and Sin YM. Enhancement of protective immunity in blue gourami, *Trichogaster trichopterus* (Pallas), against *Aeromonas hydrophila* and *Vibrio anguillarum* by *A. hydrophila* major adhesin. Journal of Fish Diseases 2000; 23: 137-146.

FAO report 2004-2012; FAO. © 2004-2012. Cultured Aquatic Species Information Programme. Cyprinus carpio. Cultured Aquatic Species Information Programme.

Feiken E, Rømer J, Eriksen J and Lund LR. Neutrophils express tumor necrosis factor- α during mouse skin wound healing. The Journal of Invegastive Dermatology 1995; 105: 120-123.

Feng Q, Boone AN and Vijayan MM. Copper impact on heat shock protein 70 expression and apoptosis in rainbow trout hepatocytes. Comparative Biochemistry and Physiology 2003; 135: 345–355.

Fletcher TC, Jones R and Reid L. Identification of glycoproteins in goblet cells of epidermis and gill of plaice (*Pleuronectes platessa* L.), flounder (*Platichthys flesus* (L.)) and rainbow trout (*Salmo gairdneri* Richardson). The Histochemical Journal 1976; 8: 597-608.

Flik G and Perry SF. Cortisol stimulates whole body calcium uptake and the branchial calcium pump in freshwater rainbow trout. Journal of Endocrinology 1989; 120: 75-82.

Fournier-Betz V, Quentel C, Lamour F AND LeVenE A. Immunocytochemical detection of Ig-positive cells in blood, lymphoid organs and the gut associated lymphoid tissue of the turbot (*Scophthalmus maximus*). Fish and Shellfish Immunology 2000; 10: 187-202.

Fujiki K, Shin D-H, Nakao M and Yano T. Molecular cloning and expression analysis of carp (*Cyprinus carpio*) interleukin-1 β , high affinity immunoglobulin E Fc receptor γ subunit and serum amyloid A. Fish and Shellfish Immunology 2000; 10: 229-242.

Fukata M, Chen A, Klepper A, Krishnareddy S, Vamadevan AS, Thomas LS, Xu R, Inoue H, Arditi M, Dannenberg AJ and Abreu MT. Cox-2 Is Regulated by Toll-Like Receptor-4 (TLR4) Signaling: Role in Proliferation and Apoptosis in the Intestine. Gastroenterology 2006; 131: 862-877.

Galina J, Yin G, Ardo L and Jeney Z. The use of immunostimulating herbs in fish. An overview of research. Fish Physiology and Biochemistry 2009; 35: 669-676.

Gambo-Deza F, Sanchez-Espinel C and Magada-Mompo S. Presence of an unique IgT on the IGH locus in three-spined stickleback fish (*Gasterosteus aculeatus*) and the very recent generation of a repertoire of VH genes. Developmental and Comparative Immunology 2010; 34: 114-122.

Ganassin RC and Bols NC. Development of a monocyte/macrophage-like cell line, RTS11, from rainbow trout spleen. Fisha and Shellfish Immunology1998; 8: 457-476.

Geggel HS, Friend J, and Thofr RA. Conjunctival Epithelial Wound Healing. Investigative Ophthalmology and Visual Science 1984; 25: 860-863.

Geissmann F, Manz MG, Jung S, Sieweke MH, Merad M and Ley K. Development of Monocytes, Macrophages, and Dendritic Cells. Science 2010; 327: 656-661.

Gillitzer R and Goebeler M. Chemokines in cutaneous wound Healing. Journal of Leukocyte Biology 2001; 69: 513-521.

Gonzalez SF, Buchmann K and Nielsen ME. Real-time gene expression analysis in carp (*Cyprinus carpio* L.) skin: inflammatory responses caused by the ectoparasite *Ichthyophthirius multifiliis*. Fish and Shellfish Immunology 2007a; 22: 641-650.

Gonzalez SF, Huising MO, Stakauskas R, Forlenza M, Verburg-van Kemenade LBM, Buchmann K. Realtime gene expression analysis in carp (*Cyprinus carpio* L.) skin: inflammatory responses to injury mimicking infection with ectoparasites. Developmental and Comparative Immunology 2007b; 31: 244-254.

Gordon S and Taylor PR. Review: Monocyte and macrophage heterogeneity. Nature 2005; 5: 953-964.

Graham S and Secombes CJ. Theproductionofa macrophage- activating factor from rainbowtrout *Salmo gairdneri* leucocytes. Immunology 1988; 65: 293-297.

Greenwel P, Rubin J, Schwartz M, Hertzberg EL and Rojkind M. Liver fat-storing cell clones obtained from a CCl4-cirrhotic rat are heterogeneous with regard to proliferation, expression of extracellular matrix components, interleukin-6, and connexin 43. Laboratory Investigation; A journal of Technical Methods and Pathology 1993; 69: 210-216.

Grellner W, Georg T and Wilske J. Quantitative analysis of proinflammatory cytokines (IL-1 β , IL-6, TNF- α) in human skin wounds. Forensic Science International 2000; 113: 251-264.

Grove S, Johansen R, Reitan LJ, Press CM. Immune- and enzyme histochemical characterisation of leukocyte populations within lymphoid and mucosal tissues of Atlantic halibut (*Hippoglossus hippoglossus*). Fish and Shellfish Immunology 2006; 20: 693-708.

Guo S, Wharton W, Moseley P, and Shi H. Heat shock protein 70 regulates cellular redox status by modulating glutathione- related enzyme activities. Cell Stress and Chaperones 2007; 12: 245-254.

Hanington PC, Tam J, Katzenback BA, Hitchen SJ, Barreda DR, Belosevic M. Review: Development of macrophages of cyprinid fish. Developmental and Comparative Immunology 2009; 33: 411-429.

Hansen JD, Landis ED, and Phillips RB. Discovery of a unique Ig heavy-chain isotype (IgT) in rainbow trout: Implications for a distinctive B cell developmental pathway in teleost fish. Proceedings of The National Academy of Science of The United States of America 2005; 102: 6919-6924.

Hassanein HM, Banhawy MA, SolimanFM, Abdel-Rehim SA, Muller WEG and Schroder HC. Induction of Hsp70 by the Herbicide Oxyfluorfen (Goal) in the Egyptian Nile Fish, *Oreochromis niloticus*. Archives of Environmental Contamination and Toxicology 1999; 37: 78-84.

Hazen TC, Raker ML, Esch G and Fliermans CB. Ultrastructure of red sore lesions on largemouth bass micropterus – Salmonides association of the ciliate epistylis sp and the bacterium *Aeromonas hydrophila*. Journal of Protozoology 1978; 25: 351-355.

Havran WL and Jameson JM. Epidermal T Cells and Wound Healing. Journal of Immunology 2010; 15:

5423-5428.

Håvarstein LS, Aasjord PM, Ness S and Endresen C. Purification and partial characterization of an IgM-like serum immunoglobulin from Atlantic salmon (*Salmo salar*). Developmental and Comparative Immunology 1988; 12: 773-785.

Herbomel P, Thisse B and Thisse C. Zebrafish Early Macrophages Colonize Cephalic Mesenchyme and Developing Brain, Retina, and Epidermis through a M-CSF Receptor-Dependent Invasive Process. Developmental Biology 2001; 38: 274-288.

Hermann AC, Millard PJ, Blake SL, Kim CH. Development of a respiratory burst assay using zebrafish kidneys and embryos. Journal of Immunological Methods 2004; 292: 119-129.

Hirono I, Nam B-H, Enomoto J, Uchino K, Aoki T. Cloning and characterisation of a cDNA encoding Japanese flounder *Paralichthys olivaceus* IgD. Fish and Shellfish Immunology 2003; 15: 63-70.

Holmgren J and Czerkinsky C. Review: Mucosal immunity and vaccines. Nature Medicine 2005; 11: S45-S53.

Holtick U, Klein-Gonzalez N and von Bergwelt-Baildon MS. Potential of Toll-like receptor 9 agonists in combined anticancer immunotherapy strategies: synergy of PAMPs and DAMPs? Immunotherapy 2011; 3: 301-304.

Hordvik I, Berven FS, Solem ST, Hatten F and Endresen C. Analysis of two IgM isotypes in Atlantic salmon and brown trout. Molecular Immunology 2002; 39: 313-321.

Hübner G, Brauchle M, Smola H, Madlener M, Fässler R and Werner S. Differential regulation of proinflammatory cytokines during wound healing in normal and glucocorticoid-treated mice. Cytokine 1996; 8: 548-556.

Huising MO, Stolte E, Flik G, Savelkoul HFJ and Verburg-van Kemenade BML. CXC chemokines and leukocyte chemotaxis in common carp (*Cyprinus carpio* L.). Developmental and Comparative Immunology 2003; 27: 875-888.

Huttenhuis HBT, Ribeiro ASP, Bowden TJ, Van Bavel C, Taverne-Thiele AJ and Rombout JHWM. The effect of oral immuno-stimulation in juvenile carp (*Cyprinus carpio* L.). Fish and Shellfish Immunology 2006; 21: 261-271.

Iger Y, Abraham M, Dotan A, Fattal B and E. Rahamim E. Cellular responses in the skin of carp maintained in organically fertilized water. Journal of Fish Biology 1988; 33: 711-720.

Iger Y and Bonga SEW. Cellular responses of the skin of carp (*Cyprinus carpio*) exposed to acidified water. Cell and Tissue Research 1994; 275: 481-492.

Iliev DB, Castellana B, MacKenzie S, Planas JV and Goetz FW. Cloning and expression analysis of an IL-6 homolog in rainbow trout (*Oncorhynchus mykiss*). Molecular Immunology 2007; 44: 1803-1807.

Ingerslev HC, Lunder T, Nielsen ME. Inflammatory and regenerative responses in salmonids following mechanical tissue damage and natural infection. Fish and Shellfish Immunology 2010; 29: 440-450.

Iwama GK, Afonso LOB, Todgham A, Ackerman P and Nakano K. Are hsps suitable for indicating stressed states in fish? The Journal of Experimental Biology 2004; 207: 15-19.

Iwama GK, Vijayan MM, Forsyth RB and Ackerman PA. Heat Shock Proteins and Physiological Stress in Fish. American Zoologist 1999; 39: 901-909.

Jacquier-Sarlin MR, Jarnot L and Polla BS. Dirrefential expression and regulation of hsp70 and hsp90 by

phorbol esters and heat shock. The Journal of Biological Chemistry 1995; 270: 14094-14099.

Janeway CA and Medzhitov R. Innate immune recognition. Annual Review of Immunology 2002; 20:197-216.

Jang SI, Hardie LJ, and Secombes CJ. Elevation of rainbow trout *Oncorhynchus mykiss* macrophage respiratory burst activity with macrophage-derived supernatants. Journal of Leukocyte Biology 1995; 57: 943-947.

Jault C, Pichon L and Chluba J. Toll-like receptor gene family and TIR-domain adapters in *Danio rerio*. Molecular Immunology 2005; 40: 759-771.

Jia X, Patrzykat A, Devlin RH, Ackerman PA, Iwama GK and Hancock REW. Antimicrobial Peptides Protect Coho Salmon from *Vibrio anguillarum* Infections. Applied and Environmental Microbiology 2000; 66: 1928-1932.

Jimenez N, Coll J, Salguero FJ, Tafalla C. Co-injection of interleukin 8 with the glycoprotein gene from viral haemorrhagic septicemia virus (VHSV) modulates the cytokine response in rainbow trout (*Oncorhynchus mykiss*). Vaccine 2006; 24: 5615-5626.

Jin-Jang G, Di-Seng W and Hung-Tu H. Spontaneous remission of edema and regranulation of goblet cells in rat tracheae after capsaicin-induced acute inflammation. Anatomy and Embriology 2003; 206: 301-309.

Joerink M, Ribeiro CMS, Stet RJM, Hermsen T, Savelkoul HFJ and Wiegertjes GF. Head Kidney-Derived Macrophages of Common Carp (*Cyprinus carpio* L.) Show Plasticity and Functional Polarization upon Differential Stimulation. The Journal of Immunology 2006; 177: 61-69.

Jørgensen JB and Robertsen B. Yeast β -glucan stimulates respiratory burst activity of Atlantic salmon (*Salmo salar* L.) macrophages. Developmental and Comparative Immunology1995; 19: 43-57.

Kaattari S, Evans D and Klemer J. Varied redox forms of teleost IgM: an alternative to isotypic diversity? Immunological Reviews 1996; 166: 133-142.

Kaattari SL, Zhang HL, Khor IW, Kaattari IM and Shapiro DA. Affnity maturation in trout: clonal dominance of high affnity antibodies late in the immune response. Developmental and Comparative Immunology 2002; 26: 191-200.

Kanter M and Akpolat M. Vitamin C protects against ionizing radiation damage to goblet cells of the ileum in rats. Acta Histochemica 2008; 110: 481-490.

Karunasagar I, Ali PKMM, Jeyasekaran G and Karunasagar I. Ulcerative form of Aeromonas hydrophila infection of Catla catla. Current Science (Bangalore) 1986; 55: 1194-1195.

Kearn GC. The survival of monogenean (platyhelminth) parasites on fish skin. Parasitology 1999; 119: S57-S8.

Kesimer M, Makhov AM, Griffith JD, Verdugo P and Sheehan JK. Unpacking a gel-forming mucin: a view of MUC5B organization after granular release. American Journal of Physiology - Lung Cellular and Molecular Physiology 2010; 298: L15-L22.

Kimura Y, Sumiyoshi M, Suzuki T, Suzuki T and Sakanaka M. Effects of water-soluble low-molecular weight β -1,3-D-glucan (branch β -1,6) isolated from ureobasidium pullulans 1A1 strain black yeast on resistant stress in mice. Journal of Pharmacy and Pharmacology 2007; 59: 1137-1144.

Koumans-van Diepen JCE, Egberts E, Peixoto BR, Taverne N, and Rombout JHWM. B cell and immunoglobulin heterogeneity in carp *(Cyprinus* carpio L.); Immuno(cyto)chemical study. Developmentaland Comparative Immunology 1995; 19: 97-108.

Krkosek M, Connors BM, Ford H, Peacock S, Mages P, Ford JS, Morton A, Volpe JP, Hilborn R, Dill LM and Lewis MA. Fish farms, parasites, and predators: implications for salmon population dynamics. Ecological applications. 2011; 21: 897-914.

Kwon A-H, Qiu Z, Hashimoto M, Yamamoto K and Kimura T. Effects of medicinal mushroom (*Sparassis crispa*) on wound healing in streptozotocin-induced diabetic rats. The American Journal of Surgery 2009; 197: 503-509.

Laing KJ, Purcell MK, Winton JR and Hansen JD. A genomic view of the NOD-like receptor family in teleost fish: identification of a novel NLR subfamily in zebrafish. BMC Evolutionary Biology 2008; 8: 42-56.

Lamas J and Ellis AE. Atlantic salmon (*Salmo salar*) neutrophil response to *Aeromonas hydrophila*. Fish and Shellfish Immunology 1994; 4: 201-219.

Larsen CG, Anderson AO, Oppenheim JJ and Matsushima K. Production of interleukin-8 by human dermal fibroblasts and keratinocytes in response to interleukin-1 or tumour necrosis factor. Immunology 1989; 68:31-36.

Lasarte JJ, Casares N, Gorraiz M, Hervás-Stubbs S, Arribillaga L, Mansilla C, Durantez M, Llopiz D, Sarobe P, Borrás-Cuesta F, Prieto J and Leclerc C. The Extra Domain A from Fibronectin Targets Antigens to TLR4-Expressing Cells and Induces Cytotoxic T Cell Responses In Vivo. The Journal of Immunology 2007; 178:748-756.

Lee SW, Tsou A, Chan H, Thomas J, Petrie K, Eugui EM and Allison AC. Glucocorticoids selectively inhibit the transcription of the interleukin 1b gene and decrease the stability of interleukin 1b mRNA. Proceeding of the National Academy of Sciences of the United States of America 1988; 85: 1204-1208.

Li J, Barreda DR, Zhang Y-A, Boshra H, Gelman AE, Lapatra S, Tort L and Oriol SJ. B lymphocytes from early vertebrates have potent phagocytic and microbicidal abilities. Nature Immunology 2006; 7: 1116-1124.

Li GC and Ching-Hsien L. Heat shock protein HSP70 protects cells from oxidative stress. International Journal of Radiation Oncology Biology Physics – Supplement 1992; 24: 180.

Li A, Dubey S, Varney ML, Dave BJ and Singh RK. IL-8 Directly Enhanced Endothelial Cell Survival, Proliferation, and Matrix Metalloproteinases Production and Regulated Angiogenesis. The Journal of Immunology 2003; 170: 3369-3376.

Li X, Wang S, Qi J, Echtenkamp SF, Chatterjee R, Wang M, Boons G-J, Dziarski R and Gupta D. Zebrafish Peptidoglycan Recognition Proteins Are Bactericidal Amidases Essential for Defense against Bacterial Infections. Immunity 2007; 27: 518-529.

Lin S-H, Davidson GA, Secombes CJ and Ellis AE. A morphological study of cells isolated from the perfused gill of dab and Atlantic salmon. Journal of Fish Biology 1998; 53: 560-568.

Lin S-H, Ellis AE, Davidson GA and Secombes CJ. Migratory, respiratory burst and mitogenic responses of leucocytes isolated from the gills of rainbow trout (*Oncorhynchus mykiss*). Fish and Shellfish Immunology 1999; 9: 211-226.

Martin P. Wound Healing-Aiming for Perfect Skin Regeneration. Science 1997; 276: 75-81.

Martin P and Leibovich SJ. Inflammatory cells during wound repair: the good, the bad and the ugly. Trends in Cell Biology 2005; 15: 599-607.

Martínez-Porchas, Martínez -Córdova LR and Ramos-Eeriquez R. Cortisol and Glucose: Reliable indicators of fish stress? Pan-American Journal of Aquatic Sciences 2009; 4: 158-178.

Matzinger P. Memories are made of this? Nature 1994; 369: 605-606.

Matzinger P. An Innate Sense of Danger. Annals of the New York Acadademy of Sciences 2002; 961: 341-342.

Matzinger P. The Danger Model: A Renewed Sense of Self. Science 2002; 296: 301-305.

McAuley JL, Linden SK, Png CW, King RM, Pennington HL, Gendler SJ, Florin TH, Hill GR, Korolik V and McGuckin MA. MUC1 cell surface mucin is a critical element of the mucosal barrier to infection. Journal of Clinical Investigation 2007, 117: 2313-2324.

McKay LI and Cidlowski JA. Molecular control of immune/inflammatory responses: interactions between nuclear factorkB and steroid receptor-signaling pathways. Endocrine Reviews 1999; 20: 435-459.

Medzhitov R and Janeway CA. How does the immune system distinguish self from nonself? Immunology 2000; 12: 185-188.

Medzhitov R. Cell Death, Tissue Damage, and Inflammation. The Third International Damage Associated Molecular Pattern Molecules (DAMPs) and Alarmins Symposium iii-iDEAs in Pittsburgh, PA; USA 2008.

Mittal PS and Mittal AK. Characterisation of glycoproteins in the secretory cells in the operculum of an Indian hill stream fish *Garra lamta* (Hamilton) (Cyprinidae, Cypriniformes). Fish Physiology and Biochemistry 2002; 26: 275-288.

Mittal PS and Mittal AK. Glycoproteins in the Epithelium of Lips and Associated Structures of a Hill Stream Fish *Garra lamta* (Cyprinidae, Cypriniformes): A Histochemical Investigation. Anatomy, Histology, Embriology 2008; 37: 101-113.

Mommsen TP, Vijayan MM and Moon TW. Cortisol in teleosts: dynamics, mechanisms of action, and metabolic regulation. Reviews in Fish Biology and Fisheries 1999; 9: 211-268.

Monteiro SM, Oliveira E, Fontaínhas-Fernandes A and Sousa M. Fine Structure of the Branchial Epithelium in the Teleost *Oreochromis niloticus*. Journal of Morphology 2010; 271:621-633.

Mulero I, Sepulcre MP, Meseguer J, García-Ayala A and Mulero V. Histamine is stored in mast cells of most evolutionarily advanced fish and regulates the fish inflammatory response. PNAS: Proceedings of the National Academy of Sciences 2007; 104: 19434-19439.

Mulero I, Sepulcre MP, Roca FJ, Meseguer J, García-Ayala and Mulero v. Characterization of macrophages from the bony fish gilthead seabream using an antibody against the macrophage colony-stimulating factor receptor. Developmental and Comparative Immunology 2008; 32: 1151-1159.

Nakagawa Y, Ohno N and Murai T. Suppression by Candida albicans beta-glucan of cytokine release from activated human monocytes and from T cells in the presence of monocytes. Journal of Infectious Diseases 2003; 187: 710-713.

Narvaez E, Berendsen J, Guzmán, Gallardo JA and Mercado L. An immunological method for quantifying antibacterial activity in *Salmo salar* (Linnaeus, 1758) skin mucus. Fish and Shellfish Immunology 2010; 28: 235-239.

Neumann NF, Fagan D and Belosevic M. Macrophage activating factor(s) secreted by mitogen stimulated goldfish kideney leukocytes synergize with bacterial lipopolysaccharide to induce nitric oxide production in teleost macrophages. Developmental and Comparative Immunology 1995; 19: 473-482.

Nielsen ME, Høi L, Schmidt AS, Qian D, Shimada T, ShenJY and Larsen JL. Is *Aeromonas hydrophila* the dominant motile Aeromonas species that causes disease outbreaks in aquaculture production in the Zhejiang Province of China? Diesease of Aquatic Organisms 2001; 46: 23-29.

Nielsen AR, Mounier R, Plomgaard P, Mortensen OH, Penkowa M, Speerschneider T, Pilegaard H and Pedersen BK. Expression of interleukin-15 in human skeletal muscle – effect of exercise and muscle fibre type composition. The Journal of Physiology 2007; 584: 305-31.

Nielsen AR and Pedersen BK. The biological roles of exercise-induced cytokines: IL-6, IL-8, and IL-15. Applied Physiology, Nutrition and Metabolism 2007; 32: 833-839.

Nikl L, Evelyn TPT and Albright LJ. Trials with an orally and immersion-administered beta-1,3 glucan as an immunoprophylactic against Aeromonas salmonicida in juvenile chinook salmon *Oncorhynchus tshawytscha*. Diseases of Aquatic Organisms 1993: 17: 191-196.

Novoselova EG, Glushkova OV, Cherenkov DA, Parfenyuk SB, Novoselova TV, Lunin SM, Khrenov MO, Guzhova IV, Margulis BA, and Fesenko EE. Production of Heat Shock Proteins, Cytokines, and Nitric Oxide in Toxic Stress. Biochemistry 2006; 71: 376-383.

Oehlers SHB, Flores MV, Hall CJ, O'Toole R, Swift S, Crosier KE and Crosier PS. Expression of zebrafish cxcl8 (interleukin-8) and its receptors during development and in response to immune stimulation. Developmental and Comparative Immunology 2010; 34: 352-359.

Ohno S. Evolution by Gene Duplication. Springer, New York. 1970. 160.

Ooi VEC and Liu F. Immunomodulation and anti-cancer activity of polysaccharide-protein complexes. Current Medicinal Chemistry 2000; 7: 715-729.

Orban L and Wu Q. Cyprinds. In: Kocher T and Kole C. Genome Mapping and Genomics in Fishes and Aquatic Animals 20008; Chapter 2: 45-83.

Ossum CG, Hoffmann EK, Vijaya MM, Holte SE and Bols NC. Characterization of a novel fibroblast-like cell line from rainbow trout and responses to sublethal anoxia. Journal of Fish Biology 2004; 64: 1103-1116.

Pai R, Karunasagar I, Shetty HPC and Karunasagar I. The effect of some stress factors on infection of fish by *Aeromonas hydrophila*. Journal of Aquaculture in The Tropics 1995; 10: 29-35.

Palazon A, Dubrot J, Martinez-Ferero I, Rouzaunt-Subira A, Ochoa C, Perez-Garcia J, Hervas-Stubbs S and Melero I. Polly Matzinger's "danger model" finds its predicted danger-denoting self moieties. Immunologia 2008; 27: 205-211.

Palić D, Andreasen CB, Frank DF, Menzel BM and Roth JA. Gradient separation and cytochemical characterisation of neutrophils from kidney of fathead minnow (*Pimephales promelas* Rafinesque, 1820). Fish and Shellfish Immunology 2005a; 18: 263-267.

Palić D, Andreasen CB, Menzel BW and Roth JA. A rapid, direct assay to measure degranulation of primary granules in neutrophils from kidney of fathead minnow (*Pimephales promelas* Rafinesque, 1820). Fish and Shellfish Immunology 2005b; 19: 217-227.

Palić D, Andreasen CB, Ostojić J, Tell RM and Roth JA. Zebrafish (*Danio rerio*) whole kidney assays to measure neutrophil extracellular trap release and degranulation of primary granules. Journal of Immunological Methods 2007; 319: 87-97.

Peleteiro MC and Richards RH. Immunocytochemical studies on immunoglobulin-containing cells in the epidermis of rainbow trout *Salmo gaivdneri* Richardson: influence of bath vaccination. Journal of Fish BIology 1988; 32:845-858.

Peleteiro MC and Richards RH. Phagocytic cells in the epidermis of rainbow trout, *Salmo gairdneri* Richardson. Jotirmil of Fish Diseases 1990; 13: 225-232.

Piccini A, Carta S, Tassi S, Lasiglié D, Fossati G and Anna Rubartelli A. ATP is released by monocytes

stimulated with pathogen-sensing receptor ligands and induces IL-1 β and IL-18 secretion in an autocrine way. PNAS: Proceedings of the National Academy of Sciences 2008; 105: 8067-8072.

Pickering AD. Endocrine-induced pathology in stressed salmonid fish. Fisheries Research 1993; 17: 35-50.

Plumb JA, Grizzle JM and Defigueiredo J. Necrosis and bacterial infection in channel catfish *Ictalurus punctatus* following hypoxia. Journal of Wildlife Diseases 1976; 12: 247-253.

Pratt W. and Toft DO. Steroid Receptor Interactions with Heat Shock Protein and Immunophilin Chaperones. Endocrine Reviews 1997; 18: 306-360.

Press CM and Evensen Ø. The morphology of the immune system in teleost fishes. Fish and Shellfish Immunology 1999; 9: 309-318.

Pucci B, Coscia MR, Oreste U. Characterization of serum immunoglobulin M of the Antarctic teleost Trematomus bernacchii. Comparative Biochemistry and Physiology 2003; 135: 349-357.

Purcell MK, Kurath G, Garver KA, Herwig RP and Winton JR. Quantitative expression profiling of immune response genes in rainbow trout following infectious haematopoietic necrosis virus (IHNV) infection or DNA vaccination. Fish and Shellfish Immunology 2004; 17: 447-462.

Purcell MK, Smith KD, Aderem A, Hood L, Winton JR and Roach JC. Conservation of Toll-like receptor signaling pathways in teleost fish. Comparative Biochemistry and Physiology 2006; 77-88.

Qiu L, Zhang H, Yang K and S. Molecular cloning and mRNA expression analysis of interleukin-8 gene in Japanese sea perch (*Lateolabrax japonicus*). Molecular Biology Reports 2009; 36: 1099-1105.

Raa J, Roerstad G, Engstad R and Robertsen B. The use of immunostimulants to increase the resistance of aquatic organisms to microbial infections. In: Shariff IM, Subasinghe RP, Arthur JR, editors. Diseases in Asian Aquaculture II. Manila, Philipines: Fish Health Section, Asian Fisheries Society 1992: 39-50.

Rakers S, Gebert M, Uppalapati S, Meyer W, Maderson P, Sell AF, Kruse C and Paus R. Fish matters': the relevance of fish skin biology to investigative dermatology. Experimental Dermatology 2010; 19: 313-324.

Råbergh CMI, Airaksinen S, Soitamo A, Björklund HV, Johansson T, Nikinmaa M AND SistonenL. Tissuespecific expression of zebrafish (*Danio rerio*) heat shock factor 1 mRNAs in response to heat stress. The Journal of Experimental Biology 2000; 203: 1817-1824.

Rennekampff, H-O Hansbrough JH, Kiessig V, Doré C, Sticherling M and Schröder J-M. Bioactive Interleukin-8 Is Expressed in Wounds and Enhances Wound Healing. Journal of Surgical Research 2000; 93: 41-54.

Riches D. Cytokine and Macrophage Control of Lung Fibrosis. Pediatric Pulmonology – Supplement 1996; 13: S15.4

Roach JC, Glusman G, Rowen L, Kaur A, Purcell MK, Smith KD, Hood LE and Aderem A. The evolution of vertebrate Toll-like receptors. PNAS: Proceedings of the National Academy of Sciences 2005; 102: 9577-9582.

Robertson B, Rorstad G, Engstad R and Raa J. Enhancement of non-specific disease resistance in Atlantic salmon *Salmo salar* L. by a glucan from Saccharomyces ceravisiae cell walls. Journal of Fish Direases 1990; 13: 391-400.

Rodgers CJ and Furones MD. Antimicrobial agents in aquaculture: Practice, needs and issues. Options Méditerranéennes 2009; 86: 41-59.

Rodríguez I, Chamorro R, Novoa B and Figueras A. β-Glucan administration enhances disease resistance and some innate immune responses in zebrafish (*Danio rerio*). Fish and Shellfish Immunology 2009; 27: 369-373.

Rodríguez A, Esteban MÁ AND Meseguer J. Phagocytosis and Peroxidase Release by Seabream (*Sparus aurata* L.) Leucocytes in Response to Yeast Cells. The Anatomical Record 2003; 272: 415-423.

Rodriguez MF, Wiens GD, Purcell MK and Palti Y. Characterization of Toll-like receptor 3 gene in rainbow trout (*Oncorhynchus mykiss*). Immunogenetics 2005; 57: 510-519.

Romano N, Picchietti S, Taverne-Thiele AJ, Taverne N, Abelli L, Mastrolia L, Verburg-van Kemenade BML and Rombout JHWM. Distribution of macrophages during fish development: an immunohistochemical study in carp (*Cyprinus carpio*, L.). Anatomy and Embryology 1998; 198:31-41.

Rombout JHWM, Abelli L, Picchietti S, Scapigliati G and Kiron V. Review: Teleost intestinal immunology. Fish and Shellfish Immunology 2011; 31: 616-626.

Rombout JHWM, Huttenhuis HBT, Picchietti S and Scapigliati G. Phylogeny and ontogeny of fish leucocytes. Fish and Shellfish Immunology 2005; 19: 441-455.

Rombout JHWM, Lamers CHJ, Helfrich MH, Dekker A and Taverne-Thiele JJ. Uptake and transport of intact macromolecules in the intestinal epithelium of carp (*Cyprinus carpio* L.) and the possible immunological implications. Cell and Tissue Research 1985; 239: 519-530.

Rombout JHWM, Taverne-Thiele AJ and Villena MI. The gut-associated lymphoid tissue (GALT) of carp (*Cyprinus carpio* L.): an immunocytochemical analysis. Developmentaland ComparativeImmunology 1993; 17: 55-66.

Rosenberger R. The cell wall. In: Smith J, Berry D (eds) The Filamentous fungi, Vol 2 Biosynthesis and metabolism. Edward Arnold, London 1976; 328-334.

Roussel P and Delmotte P. The diversity of epithelial secreted mucins. Current Organic Chemistry 2004; 8: 413-437.

Rovere-Querini P, Capobianco A, Scaffidi P, Valentinis B, Catalanotti F, Giazzon M, Dumitriu IE, Müller S, Iannacone M, Traversari C, Bianchi ME and Manfredi AA. HMGB1 is an endogenous immune adjuvant released by necrotic cells. EMBO: European Molecular Biology Organization 2004; 5: 825-830.

Ruiz C, Pérez E, García-Martínez O, Díaz-Rodríguez L, Arroyo-Morales M and Reyes-Botella C. Expression of cytokines IL-4, IL-12, IL-15, IL-18, and IFNg and modulation by different growth factors in cultured human osteoblast-like cells. Journal of Bone and Mineral Metabolism 2007; 25: 286-292.

Ryo S, Wijdeven RHM, Tyagi A, Hermsen T, Kono T, Karunasagar I, Rombout JHWM, Sakai M, Verburgvan Kemenade BML and Savan R. Common carp have two subclasses of bonyfish specific antibody IgZ showing differential expression in response to infection. Developmental and Comparative Immunology 2010; 34:1183-1190.

Rørstad G, Aasjord PM and Robertsen B. Adjuvant effect of a yeast glucan in vaccines against furunculosis in Atlantic salmon (*Salmo salar* L.). Fish Shellfish Immunology1993; 3: 179-190.

Saba AA, Kaidi AA, Godziachvili V, Dombi GW, Dawe EJ, Libcke JH and Silva YJ. Effects of interleukin-6 and its neutralizing antibodies on peritoneal adhesion formation and wound healing. American Surgeon 1996; 62: 569-572.

Saeij JPJ, Verburg-van Kemenade LBM, van Muiswinkel WB and Wiegertjes GF. Daily handling stress reduces resistance of carp to *Trypanoplasma borreli*: in vitro modulatory effects of cortisol on leukocyte

function and apoptosis. Developmental and Comparative Immunology 2003; 27: 233-245.

Saha NR, Suetake H, Kikuchi K and Suzuki Y.Fugu immunoglobulin D: a highly unusual gene with unprecedented duplications in its constant region. Immunogenetics 2004; 56: 438–447.

Sangrador-Vegas A, Lennington JB and Smith TJ. Molecular cloning of and IL-8-like CXC chemokine and Tissue factor in rainbow trout (*Oncorhynchus mykiss*) by use of suppression substrative hybridization. Cytokine 2002; 17: 66-70.

Sass JB, Weinberg ES and Krone PH. Specific localization of zebrafish hsp90-alpha mRNA to myoD-expressing cells suggests a role for hsp90-alpha during normal muscle development. Mechanisms of development 1996; 54: 195-204.

Savan R, Aman A, Nakao M, Watanuki H and Sakai M. Discovery of a novel immunoglobulin heavy chain gene chimera from common carp (*Cyprinus carpio* L.). Immunogenetics 2005; 57: 458-463.

Scapigliati G, Bird S, Secombes CJ. Invertebrate and fish cytokines. European Cytokine Network 2000; 11: 354-361.

Schrøder MB, Flańio E, Pilström L and Jørgensen TØ. Localisation of Ig heavy chain mRNA positive cells in Atlantic cod (*Gadus morhua* L.) tissues; identified by in situ hybridization. Fish and Shellfish Immunology 1998; 8: 565-576.

Secombes CJ and Fletcher TC. The role of phagocytes protective mechanism in the fish. Annual Reviews of Fish Diseases 1991; 53-71.

Secombes CJ, Hardie LJ AND Daniels G. Cytokines in fish: an update. Fish and Shellfish Immunology 1996; 6: 291-304.

Selvaraj V, Sampath K and Sekar V. Administration of yeast glucan enhances survival and some nonspecific and specific immune parameters in carp (*Cyprinus carpio*) infected with *Aeromonas hydrophila*. Fish and Shellfish Immunology 2005; 19: 293-306.

Seong S-Y and Matzinger P. Hydrophobicity: an ancient damage- associated molecular pattern that initiates innate immune responses. Nature Reviews 2004; 4: 469-478.

Seppola M, Larsen AN, Steiro K, Robertsen B and Jensen I. Characterisation and expression analysis of the interleukin genes, IL-1β, IL-8 and IL-10, in Atlantic cod (*Gadus morhua* L.). Molecular Immunology 2008; 45: 887-897.

Sha J, Kozlova EV and Chopra AK. Role of various enterotoxins in *Aeromonas hydrophila*-induced gastroenteritis: generation of enterotoxin gene-deficient mutants and evaluation of their enterotoxic activity. Infection and Immunity 2002; 70: 1924-1935.

Shephard KL. Functions for fish mucus. Reviews in fish biology and fisheries 1994; 4: 401-429.

Sheehan JK, Brazezu C, Kutay S, Pigeon H, Kirkham S, Howard M and Thornton DJ. Physical characterization of the MUC5AC mucin: A highly oligomeric glycoprotein whether isolated from cell culture or in vivo from respiratory mucous secretions. Biochemical Journal 2000; 347: 37-44.

Shi Y, Evans JE and Rock KL. Molecular identification of a danger signal that alerts the immune system to dying cells. Nature 2003; 425: 516-521.

Singer AJ and Clark RAF. Cutaneous wound healing. The New England Journal of Medicine 1999; 341: 738-746.
Smith GS and Lumsden JH. Review of neutrophil adherence, chemotaxis, phagocytosis and killing. Veterinary Immunology and Immunopathology 1983; 4: 177-236.

Soanes KH, Ewart KVand Mattatall NR. Recombinant production and characterization of the carbohydrate recognition domain from *Atlantic salmon* C-type lectin receptor C (SCLRC). Protein Expression and Purification 2008; 59: 38-46.

Solem ST and Jørgen S. Antibody repertoire development in teleosts – a review with emphasis on salmonids and Gadus morhua L. Developmental and Comparative Immunology 2006; 30: 57-76.

Stein C, Caccamo M, Laird G, Leptin M. Conservation and divergence of gene families encoding components of innate immune response systems in the zebrafish. Genome Biol. 2007; 8: R251.

Stenvik J, Olsen K, Schroder M and Jorgensen TO. The expression of immunoglobulin D (IgD) in Atlantic cod (*Gadus morhua* L.). Developmental and Comparative Immunology – Supplement 2000; 24: S36-S237.

Stolte EH, Chadzinska M, Przybylska D, Flik G, Savelkoul HFJ and Verburg-van Kemenade BML. The immune response differentially regulates Hsp70 and glucocorticoid receptor expression in vitro and in vivo in common carp (*Cyprinus carpio* L.). Fish and Shellfish Immunology 2009; 27: 9-16.

Stolte EH, Verburg van Kemenade BML, Savelkoul HFJ and Flik G. Review: Evolution of glucocorticoid receptors with different glucocorticoid sensitivity. Journal of Endocrinology 2006; 190: 17-28.

Strieter RM, Kunkel SL, Showell HJ, Remick DG, Phan SH, Ward PA and Marks RM. Endothelial Cell Gene Expression of a Neutrophil Chemotactic Factor by TNF-α, LPS, and IL-1β. Science 1989; 243: 1467-1469.

Strous GJ and Dekker J. Mucin-Type Glycoproteins. Critical Reviews in Biochemistry and Molecular Biology 1992; 27: 57-92.

Subramanian S, MacKinnon S and Ross NW. A comparative study on innate immune parameters in the epidermal mucus of various fish species. Comparative Biochemistry and Physiology 2007; 148: 256-263.

Takeda K, Kaisho T and Akira S. Toll-like receptors. Annual Review of Immunology 2003; 21: 335-376.

Tanck MWT, Booms GHR, Eding EH, Bonga SEW AND Komen J. Cold shocks: a stressor for common carp. Journal of Fish Biology 2000; 57: 881-894.

Tanioka A, An W-W, Kuge T, Tsubaki K and Nakaya K. Barley Low Molecular Weight beta-Glucan Potently Induces Maturation of Mouse Dendritic Cells. Anticancer Research 2011; 31: 1647-1651.

Temkin RJ and McMillan DB. Gut-associated lymphoid tissue of the goldfish *Carassius aurata*. Journal of Morphology 1986; 190: 9-26.

Thoma-Uszynski S, Stenger S, Takeuchi O, Ochoa MT, Engele M, Sieling PA, Barnes PF, Röllinghoff M, Bölcskei PL, Wagner M, Akira S, Norgard MV, Belisle JT, Godowski PJ, Bloom BR and Modlin RL. Induction of Direct Antimicrobial Activity Through Mammalian Toll-Like Receptors. Science 2001, 291: 1544-1547.

Thornton DJ, Rousseau K and McGuckin MA. Structure and Function of the Polymeric Mucins in Airways Mucus Annual Review of Physiology 2008; 70: 459-486.

Toa DG, Afonso LOB and Iwama GK. Stress response of juvenile rainbow trout (*Oncorhynchus mykiss*) to chemical cues released from stressed conspecifics. Fish Physiology and Biochemistry 2004; 30: 103-108.

Tort L, Balasch JC and Mackenzie S. Fish immune system. A crossroads between innate and adaptive responses.

Tortora GJ and Grabowski SR. Human physiology; Human anatomy; Anatomy; Physiology. In Principles of anatomy and physiology. Harper Collins College, New York 1996.

Trautinger F. Heat shock proteins in the photobiology of human skin. Journal of Photochemistry and Photobiology 2001; 63: 70-77.

Uran PA, Aydin RJ, Schrama W, Verreth JAJ and Rombout JHWM. Soybean meal-induced uptake block in Atlantic salmon *Salmo salar* distal enterocytes. Journal of Fish Biology 2008; 73: 2571-2579.

van der Aa LM, Chadzinska M, Tijhaar E, Boudinot P and Verburg-van Kemenade BML. CXCL8 Chemokines in Teleost Fish: Two Lineages with Distinct Expression Profiles during Early Phases of Inflammation. Plos One 2010; 5: 1-13.

van der Marel M, Adamek M, Gonzalez SF, Frost P, Rombout JHWM, Wiegertjes GF, Savelkoul HFJ, Steinhagen D. Molecular cloning and expression of two β -defensin and two mucin genes in common carp (*Cyprinus carpio* L.) and their up-regulation after β -glucan feeding. Fish and Shellfish Immunology 2012; 32: 494-501.

van der Marel M, Caspari N, Neuhaus H, Meyer W, Enss M-L and Steinhagen D. Changes in skin mucus of common carp, *Cyprinus carpio* L., after exposure to water with a high bacterial load. Journal of Fish Diseases 2010; 33: 431-439.

Varela M, Dios S, Novoa B and Figueras A. Characterisation, expression and ontogeny of interleukin-6 and its receptors in zebrafish (*Danio rerio*). Developmental and Comparative Immunology 2012; 37: 97-106.

Wagstaff MJD, Smith J, Collaco-Moraes Y, de Belleroche JS, Voellmy R, Coffin RS and Latchman DS. Delivery of constitutively active form of heat shock factor using a virus vectorprotects neuronal cells from thermal or iachaemic stress but not from apoptosis. European Journal of Neuroscience 1998; 10: 3343-3350.

Wartha F and Henriques-Normark B. ETosis: A Novel Cell Death Pathway. Science Signalling 2008; 27: pe25.

Werner S AND Grose R. Regulation of Wound Healing by Growth Factors and Cytokines. Ohysiological Reviews 2003; 83: 835-870.

Wei D, Zhang L, Williams DL and Browder IW. Glucan stimulates human dermal fibroblast collagen biosynthesis through a nuclear factor-1 dependent mechanism. Wound Repair and Regeneration 2002; 10: 161-168.

Weyts FAA, Flik G, Rombout JHWM and Verburg-van Kemenade BML. Cortisol induces apoptosis in activated B cells, not in other lymphoid cells of the common carp, *Cyprinus carpio* L. Developmental and Comparative Immunology 1998a; 22: 551-562.

Weyts FAA, Flik G and Verburg-van Kemenade BML. Cortisol inhibits apoptosis in carp neutrophilic granulocytes. Developmental and Comparative Immunology1998b; 22: 563-572.

Weyts FAA, Verburg-van Kemenade BML and Flik G. Characterisation of Glucocorticoid Receptors in Peripheral Blood Leukocytes of Carp, *Cyprinus carpio* L. General and Comparative Endocrinology 1998; 111: 1–8.

Weyts FAA, Verburg-van Kemenade BML, Flik G, Lambert JGD and Bong SEW. Conservation of Apoptosis as an Immune Regulatory Mechanism: Effects of Cortisol and Cortisone on Carp Lymphocytes. Brain Behaviour and Immunity 1997; 11: 95-105.

Williams DL and DiLuzio NR. Immunopharmacologic modification of experimental viral diseases by glucan. Eos-Rivista Di Immunologia ed. Immunofarmacologia 1985 5: 78-82.

Williams DL, Mueller A and Browder W. Glucan-based macrophage stimulators – Areview of their antiinfective potential. Clinical Immunotherapeutics 1996; 5: 392-399.

Wilson M, Bengtén E, Miller NW, Clem LW, Pasquier LD AND Warr GW. A novel chimeric Ig heavy chain from a teleost fish shares similarities to IgD. Proceedings of the National Academy of Science 1997; 94: 4593-4597.

Wilson JM AND Laurent P. Fish Gill Morphology: Inside Out. Journal of Experimental Zoology 2002; 293: 192-213.

Wilson MR, Marcuz A, van Ginkel F, Milerl NW, Clem LW, Middleton D and Warr GW. The immunoglobulin M heavy chain constant region gene of the channel catfish, *Ictalurus punctatus*: an unusual mRNA splice pattern produces the membrane form of the molecule. Nucleic Acids Research 1990; 8: 5227-5233.

Wolk D and Danon D. Promotion of wound healing by yeast glucan evaluated on single animals. Medical Biology 1985; 63: 73-80.

Yamamoto T, Kawai K and Oshima S. Distribution of mucous cells on the body surface of Japanese flounder *Paralichthys olivaceus*. Journal of Fish Biology 2011; 78: 848-859.

Yano T, Matsuyama H and Mangindaan REP. Polysaccharide-induced protection of carp *Cyprinus carpio* L. against bacterial infection. Journal of Fish Diseases 1991; 14: 577-582.

Yao M, Li Q and Martins-Green M. Functions of the chemokine IL-8 in inflammation-induced angiogenesis. Molecular Biology of the cell - Supplement 2002; 13: 26a.

Yoshida T, Kruger R and Inglis V. Augmentation of non-specific protection in African catfish, Clarias gariepinus (Burchell), by the long-term oral administration of immunostimulants. Journalof Fish Diseases 1995: 18: 195-198.

Verdugi P. Goblet cells secretion and mucogenesis. Annual Review of Physiology 1990; 52: 157-176.

Zapata, A. and E.L.Cooper: The immune system: Comparative Histophysiology. John Wiley Sons, Chichester, U.K. (1990).

Zhang Y-A, Salinas I, Li J, Parra D, Bjork S, Xu Z, LaPatra SE, Bartholomew J and Sunyer JO. IgT, a primitive immunoglobulin class specialized in mucosal immunity. Nature 2010; 11: 827-836.

Zhang Z, Swain T, Bøgwald J, Dalmo RA and Kumari J. Bath immunostimulation of rainbow trout (*Oncorhynchus mykiss*) fry induces enhancement of inflammatory cytokine transcripts, while repeated bath induce no changes. Fish and Shellfish Immunology 2009; 26: 677-684.

PAPER I

Previous bacterial infection affects textural quality parameters of heattreated fillets from rainbow trout (*Oncorhynchus mykiss*)

Hans-Christian Ingerslev, Grethe Hyldig, **Dominika Przybylska**, Stina Frosch and Michael Engelbrecht Nielsen

(published)

PREVIOUS BACTERIAL INFECTION AFFECTS TEXTURAL QUALITY PARAMETERS OF HEAT-TREATED FILLETS FROM RAINBOW TROUT (ONCORHYNCHUS MYKISS)

H.-C. INGERSLEV, G. HYLDIG, D. PRZYBYLSKA, S. FROSCH and M.E. NIELSEN¹

Technical University of Denmark, National Food Institute, Division of Industrial Food Research, Biological Quality Group, Kgs. Lyngby, Denmark

¹Corresponding author. TEL: +45-23688290; FAX: +45-45884774; EMAIL: mice@food.dtu.dk

Received for Publication April 27, 2011 Accepted for Publication February 27, 2012

doi:10.1111/j.1745-4557.2012.00443.x

ABSTRACT

Sensory quality of fish meat is influenced by many parameters prior to slaughter. In the present study, it was examined if previous infections or damages in the muscle tissue influence product quality parameters in fish. Fillets from rainbow trout (*Oncorhynchus mykiss*) reared in seawater at a commercial fish farm were sensory evaluated for more than a year after recovery following physical tissue damage or infection by the bacterial pathogens *Yersinia ruckeri* and *Vibrio anguillarum*. The effect of vaccination was also included as some fish were vaccinated before bacterial challenge. The fish fillets were sensory examined as heat-treated and cold-smoked. Heat-treated fillets from nonvaccinated fish previously infected by *V. anguillarum* had changed textural characteristics and were less flaky, had a lower oiliness and a higher toughness and fibrousness in comparison with control fish. This article was the first to describe a correlation between previous infections in fish and changes in sensory-quality parameters.

PRACTICAL APPLICATIONS

This work contributes with knowledge about sensory-quality parameters of fish meat after recovery from infections and physical-tissue damage. Because the results demonstrate an influence on the texture from previous disease, the practical potentials of the results are valuable for the aquaculture industry. In order to minimize the effects of previous diseases on the sensory quality regarding the texture, these fishes should be processed as cold-smoked instead of being sold as raw meat. The established correlation between disease history stresses the importance of disease prevention in aquaculture production, e.g., vaccination of the fish.

INTRODUCTION

Fish farming is an expanding worldwide industry, and the amount of produced fish and the number of species is increasing every year (Fao 2008). Many biological and nonbiological parameters such as welfare, rearing, stress level at slaughter and the amount of fat in the fillet have been shown to influence subsequent quality parameters of farmed fish postslaughter (Sigurgisladottir *et al.* 1997; Poli *et al.* 2005; Ashley 2007; Morkore *et al.* 2007; Pettersson *et al.* 2009; Merkin *et al.* 2010). Some of the quality parameters influenced by these factors are the texture of the fillet (meat) as well as the content of connective tissue and coloration of the fillet (Sigholt *et al.* 1997; Sigurgisladottir *et al.* 1997). Disease outbreaks occur in fish farms, and the pathogens isolated can be a wide range of bacteria, viruses, parasites and fungi (Dalsgaard and Madsen 2000; Bondad-Reantaso *et al.* 2005; Pedersen *et al.* 2008). In general, infections are seen in both freshwater and seawater, and in all developmental stages of the fish. Survival and recovery of infected fish depends, however, on the type of infection, vaccination and veterinary treatment (Coyne *et al.* 2006; Raida and Buchmann 2008a).

Previous studies from commercial fish farms have reported changes in the tissue structure of the fillet of Atlantic salmon (*Salmo salar* L.) following bacterial and viral infections (Lunder *et al.* 1995; Lerfall *et al.* 2012). These changes develop as a result of heavy damages and bleedings in the musculature caused by pathogens (Moran *et al.* 1999). The structural

changes can be associated with the development of scars as a consequence of the regeneration process (Pedersen et al. 1994). Occasionally, examples of infections leading to tissue damage and bleedings are seen in Norwegian-salmon farming, when Atlantic salmon (S. salar) becomes infected by the bacterium Moritella viscosa or by the virus causing pancreas disease (PD) (Mcloughlin et al. 1996; Benediktsdottir et al. 1998). Recently, Lerfall et al. (2012) showed using a texture analyzer that cold-smoked fillets from infected or previous infected fish had a higher firmness relative to healthy fish. Usually, fish recovered from a M. viscosa or PD infection are downgraded from "superior quality" to "production fish," causing a lowered sales price on the market (Salte et al. 1994; Aunsmo et al. 2010). In addition to infections, physical tissue damage of the skin may also occur under fish farming conditions in connection to handling, transporting and crowding. In a number of studies, the structure and texture of meat have been directly linked to quality-related aspects (Trigo et al. 2008; Wolowiny 2008; Lonergan et al. 2010). However, until now it has not been shown whether previous infection or physical tissue damage may influence the sensory quality of fish meat regarding the texture of the fillet from recovered fish.

In Denmark, bacterial pathogens causing the most serious economical problems in the aquaculture industry are the gram-negative bacteria Yersinia ruckeri and Vibrio anguillarum (Larsen and Pedersen 1999; Tobback et al. 2007). Y. ruckeri usually causes infections in fry in freshwater, while V. anguillarum is mostly associated with larger fish in brackish- and saltwater. Both bacteria can lead to high mortality and the clinical signs include ulcerations and punctual bleedings in the musculature of the fish. Earlier findings have reported development of scars following infection by V. anguillarum (Pedersen et al. 1994; Stone et al. 1997). Usually, high ratio of fillets from slaughtered rainbow trout are sold as either fresh or smoked, making it relevant to examine those products. The aim of this study was to examine if physical tissue damage or previous infections of rainbow trout by Y. ruckeri, V. anguillarum or a combination of both influence sensory parameters of the fillet after recovery from disease. The studies were performed under regular farming conditions using both vaccinated and nonvaccinated fish.

MATERIALS AND METHODS

Overall Experimental Setup

The experimental setup was designed to mimic a natural farming situation implying an initial freshwater period and a later seawater-rearing period. The freshwater phase in the current experiment was subdivided into two parts, A and B. Facility A was located at the Technical University of Denmark

and was used from fry and until a fish size of about 50 g. At this location, the fish were experimentally infected by *Y. ruck-eri* and *V. anguillarum* and further medically treated. Another group of fish underwent a local, mechanical damage of the musculature. After recovery, all fish were transferred to a second freshwater facility, facility B, for further growth before being transferred to seawater cages (facility C).

Rearing of Fish at Facility A and Experimental Groups of Fish

Initially, unvaccinated rainbow trout (0+) fry (n = 4,300)reared at Horns Herred Ørredopdræt (Skibby, Zealand) were brought to the rearing facilities at facility A, the National Food Institute, Technical University of Denmark (DTU Food) in the Autumn of 2007. The experimental facilities at DTU Food consisted of two separate units: A1 and A2. Unit A1 was used for control fish and consisted of two 1,200-L tanks that were connected to a bio-filter with recirculated water and temperature control. Unit A2 was used for infection of fish and was composed of four 400-L tanks that were also connected to a bio-filter with recirculated water, temperature control and exposure of circulating water by UV light. Both rearing facilities were supplied daily with 10% freshwater and the content of ammonium and nitrite were measured in order to ensure good water quality. The water temperature was kept at 15C in both units and the fish were exposed to 16 h of light and 8 h of darkness. Ten different experimental groups of fish were obtained at facility A after vaccination and challenge with Y. ruckeri and V. anguillarum: (1) control fish; (2) Y. ruckeri challenged; (3) Y. ruckeri vaccinated + challenged; (4) V. anguillarum challenged; (5) *V. anguillarum* vaccinated + challenged; (6) *Y. ruckeri* challenged + *V. anguillarum* vaccinated; (7) Y. ruckeri challenged + V. anguillarum challenged; (8) Y. ruckeri vaccinated + challenged + V. anguillarum challenged; and (9) Y. ruckeri vaccinated + challenged + V. anguillarum vaccinated + challenged. Another group (10) was obtained after mechanical damage of the skin and musculature. Figure 1 shows how the different experimental groups were obtained. The procedures for vaccination, challenge and mechanical tissue damage are described later.

Each of the surviving fish from groups other than control were in August 2008 individually tagged by injection of electronic passive integrated transponder (PIT) tags into the peritoneal cavity (Product no. AB10400, Loligo Systems, Tjele, Denmark). Hence, all groups of fish could later be reared in the same cage. The individual tag numbers were read using an electronic PIT reader (Product no. AB10625, Loligo Systems).

Rearing of Fish at Facility B and C

In September 2008, the fish had reach a size of approximately 50 g and all experimental groups of fish (groups 1–10;



FIG. 1. OVERVIEW OF THE DIFFERENT EXPERIMENTAL GROUPS OF FISH OBTAINED AFTER VACCINATION, CHALLENGE AND MECHANICAL TISSUE DAMAGE AT FACILITY A

Control fish were not challenged. YRV, Yersinia ruckeri vaccinated; YRC, Yersinia ruckeri challenged; VAV, V. anguillarum vaccinated; VAC, Vibrio anguillarum challenged.

n = 1,500) were moved from facility A to the second freshwater facility (facility B) in Bisserup, Denmark. Facility B was composed of four 2,000-L circular tanks containing freshwater and connected to a bio-filter. The water quality parameters were equal to the conditions at facility A. The fish were equally distributed in the tanks upon arrival. Here, the fish were reared until transferred to one seawater cage with a diameter of 40 m and a depth of 4 m in the Spring of 2009. The size of the fish when transferred to seawater was approximately 150 g. The fish were reared in the cages until slaughter in November 2009.

Procedures for Vaccination against *Y. ruckeri* and Challenge

At a size of 3 g (n = 30, subsampling), 2,150 näive fish (group 1) were bath-vaccinated against *Y. ruckeri*. The bath-vaccination was performed using the Ermogen Vet. vaccine (Aqua Health, Novartis, Charlottetown, Canada) in October

2007. The procedures were in accordance with the manufacturer's instructions by diluting the vaccine 1:10 into water and further 30 s of bathing of the fish.

At a mean size of 14 g (n = 30, subsampling), 733 previously Y. ruckeri-vaccinated and 754 nonvaccinated, näive fish (group 1) were challenged with Y. ruckeri in March 2008 using the 99.8.60 strain (serotype O1). The strain was kindly provided by Dr. Inger Dalsgaard, National Veterinary Institute in Denmark and had previously been isolated from infected rainbow trout in a Danish fish farm. Bacteria were grown for 18 h at 25C in Luria Bertani (LB) medium with shaking until a final concentration of 109 bacteria/mL. Prior to this, a challenge test using two different amounts of bacteria was set up (106 and 107 bacteria per fish) in order to determine the dose; maximally, 50% of the fish died. The bacteria were diluted in 1× phosphate buffered saline (PBS). Based on these results, 106 bacteria were chosen for challenge of experimental fish. The bacteria were intraperitoneally (i.p.) injected $(100 \,\mu\text{L})$ in the fish using a syringe. When mortality was registered, the surviving fish were treated with antibiotics.

230

Procedures for Vaccination against *V. anguillarum* and Challenge

At a mean size of 30 g (n = 30, subsampling), approximately 1,200 naïve fish (group 1) and 150 previously *Y. ruckeri* challenged fish (group 3) were vaccinated against *V. anguillarum* in May 2008 using the AquaVacTM Vibrio vet. vaccine (Schering-Plough Animal Health). The fish were i.p. injected using 100 µL of vaccine per fish according to the manufacturer's instructions.

At a mean size of 40 g (n = 30, subsampling), 119 previously *Y. ruckeri*-vaccinated and challenged (group 3) plus 149 *V. anguillarum*-vaccinated fish were challenged with *V. anguillarum* in July 2008 using the strain 070602-1/2A (serotype 01). The strain was kindly provided by Dr. Inger Dalsgaard, National Veterinary Institute in Denmark and had previously been isolated from infected rainbow trout in a Danish fish farm. Bacteria were grown in LB medium containing 1% NaCl₂ for 24 h at 25C with shaking until a final concentration of 10⁹ bacteria/mL. A challenge test was also set up for *V. anguillarum*. Based on this, a dose of 10⁵ bacteria per fish was chosen for challenge of experimental fish. The bacteria were i.p. injected (100 μ L) into the fish using a syringe. Treatment with antibiotics was initiated 2 days postchallenge.

Treatment of *Y. ruckeri*- and *V. anguillarum*-Infected Fish by Antibiotics

Fish infected by either *Y. ruckeri* or *V. anguillarum* were fed with up to 1–1.5% of body weight per day with Aquavet feed (Biomar, Brande, Denmark, product no. 84894) containing sulfadiazine and trimethoprim antibiotics. Feeding was started a few days after mortality was observed and continued until it had ceased according to veterinary regulations. For *Y. ruckeri*, feeding was initiated 6 days postchallenge and for the following 8 days. For *V. anguillarum*, feeding was started 2 days postchallenge and continued for 4 days.

Bacteriology of Challenged Fish

After challenge by *Y. ruckeri* and *V. anguillarum* plating out from the head kidney of dead fish was performed in order to confirm presence of bacteria in the blood. Plating from *Y. ruckeri* infected fish was performed on standard LB plates, while LB plates containing 5% calf blood were used for *V. anguillarum*. Both species were incubated at 20C.

Plating out was also performed from noninfected control fish in order to verify that they were free of infection. Later in the experiment, after antibiotic treatment and recovery and prior to movement of the fish from the facilities at facility A, the fish were again screened for the presence of bacteria. This was performed in order to state the fish free of pathogens and not to bring bacterial pathogens from the laboratory facilities to the sea cages.

Mechanical Damage of the Musculature

Mechanical punctual damage of the skin and musculature was performed on the left side of 50 fishes 2 weeks prior to transfer to seawater using a specially constructed device. This has previously been shown to be a good control of muscle damage, which can be compared with following a bacterial infection (Ingerslev et al. 2010). The fish had not been vaccinated or challenged prior to the damage. Within minutes before the damage, the fishes were anesthetized using a dose of 50 mg/L of MS-222 (Sigma-Aldrich, St. Louis, MO). The device contained 8×4 (32) needles (25 g, Becton Dickinson, Brøndby, Denmark) equally distributed on an area of 2.7×6.3 cm. The length of the needles, which penetrated the fish skin and musculature, was 0.6 cm. The device was penetrated once through the skin below the dorsal fin with one half of the needles above the lateral line and the other half below. The procedures were conducted in accordance with the regulations set forward by the Danish Ministry of Justice and animal protection committees by Danish Animal Experiments Inspectorate permit 2007/561-1302 and in compliance with European Community Directive 86/609.

Feeding

The fish were fed with commercial trout feed of approximately 1–1.5% of body weight per day throughout the entire experiment. The sizes of the pellets were in accordance with the manufacturer's recommendations in relation to size of the fish. Until at a size of approximately 100 g the fish were fed with a diet of DAN-EX trout feed (Danafeed, Horsens, Denmark). Once transferred to the freshwater facilities at Bisserup and for the remaining experiment the fish were fed with Aqualife/EFICO trout feed and Ecolife Pearl (Biomar).

Enzyme-Linked Immunosorbant Assays for Detection of *Y. ruckeri* and *V. anguillarum*

An enzyme-linked immunosorbant assay (ELISA) was used to detect the presence of the *Y. ruckeri* O1 and *V. anguillarum* O1 in the plasma from slaughtered fish. The ELISAs were run separate for each antigen using the same protocol (Mikkelsen *et al.* 2006).

In the following, all steps in the protocol were carried out at room temperature.

In order to prepare the antigens (*Y. ruckeri* O1 and *V. anguillarum* O1) bacteria were grown for 12 and 24 h at room temperature in LB media, respectively. The bacteria were then centrifuged for 13 min at 1,800 rpm and resuspended in coating buffer (pH 9.6).

Initially, the wells of microtiter plates (Immunosorb, Nunc, Roksilde, Denmark, product no. 269620) were coated using 200 µL of antigen solution diluted in coating buffer (106 bacteria/well) and further incubated at 4C over night. Unbound bacteria were removed by three washes in PBS-Tween, pH 7.2. The following day, the plates were blocked using 200 µL of 0.5% bovine serum albumin (BSA) in PBS and shaken for 15 min. The wells were then further washed three times in PBS-Tween. To each well, 100 µL of rainbow trout serum diluted 1:10 in PBS with 0.1% BSA was added and the plates were gently shaken for 30 min. The wells were then washed in order to remove unbound plasma. Furthermore, 100 µL of rabbit-anti trout Ig diluted 1:3,000 in PBS with 0.1% BSA (Sigma Aldrich) was then added to the wells and the plates were gently shaken for 10 min and the wells were thereafter washed three times. Subsequently, 100 µL of horseradish peroxidase conjugated goat-anti rabbit antibody (Bio-Rad, Hercules, CA, product no. 172-1019) diluted 1:3,000 in PBS with 0.1% BSA was added to the wells followed by shaking for 10 min. The wells were then washed three times. Lastly, 100 µL of enzyme substrate (TMB; Sigma-Aldrich, product no. T5525) was added to the wells and the plates were incubated for 10 min. The reaction was stopped using 50 µL of stop solution (1 M H₂SO₄). It was added when a strong color in the wells was obtained or after maximally 15 min of incubation. The absorbance was when measured at 450 nm using a Synergy 2 multi-mode microplate reader (BioTek Instruments, Winooski, VT).

Fish Slaughtering and Fillet Smoking

Prior to slaughtering, fishes were killed by sharp blows to the head. They were further gill-cut and bled for 5 min in icewater. After slaughter and gutting, the fishes were stored on ice until the next day when they were filleted in a postrigor state. The fillets were further used for the applied sensory analysis. Fillets from the left side of the fish were used for smoking, while right side fillets were used for analysis of the heat-treated fish.

The fish fillets were smoked at Bisserup Havbrug according to standard procedures the same day as the fish were filleted. Initially, the fillets were dry-salted for 3 h followed by a short dripping of water. Smoke was generated using a mix of peat litter from beech and crowberry heather. The fillets were then smoked for 40 h at 28C and subsequently vacuum packed and stored at 2C prior to sensory evaluation.

Sensory Profiling

The trout fillets were evaluated for textural characteristics by sensory profiling either as heat-treated or cold-smoked. Because of loss of some of the experimental fish, only four groups in addition to the control fish were sensory evaluated. These were: (1) previously V. anguillarumchallenged fish (group 4); (2) previously mechanically damaged fish (group 10); (3) previously Y. ruckerivaccinated + challenged + V. anguillarum-challenged fish (group 8); and (4) previously Y. ruckeri-vaccinated + challenged + V. anguillarum-vaccinated + challenged (group 9). The fillets evaluated as heat treated and were stored on ice for 3 and 7 days before steaming. The cold-smoked fillets were evaluated 9 days after smoking. A sensory panel that consists of nine (for heat-treated) or eight (cold-smoked) assessors, who had been selected, tested, and specifically trained in descriptive analysis (Iso 1988, 1993) of rainbow trout, evaluated the fillets (Hyldig 2009a). The first three sessions were quantitative, i.e., the assessors were trained to evaluate the descriptors on a linear scale. Each attribute was evaluated using a 15-cm unstructured linear scale with two anchor points that were "little" and "much" of attribute intensity. The anchor points were placed 1.5 and 13.5 cm from zero on the scale (Meilgaard et al. 1999). The sensory analysis was performed in separated booths under normal daylight and at ambient temperature (Iso 1988). The assessors used water and flat bread to clean the palate between samples. Data were collected using a computer system (FIZZ Network Version 2.0, Biosystems, Couternon, France). One fillet from each fish was analyzed once by one single assessor. The samples were placed in individual porcelain bowls and covered with porcelain lids with three-digit codes. The amount of fish prepared for each assessor was 50 g. The fish samples for steaming were heated in a prewarmed convection oven (RATIONAL Combi-Dämpfer CCM, Neuruppin, Germany) with air circulation for 15 min at 100C. After heat treatment, the samples were immediately served for the panel. The sensory attributes for analyzing the textural characteristics of heat-treated fillets were: flakiness, juiciness, firmness, oiliness, toughness and fibrousness. For appearance, they were: bright, colored and discolored (Hyldig 2009b). For the cold-smoked fillets, they were: elastic (with a fork), elastic, firm, juicy, tough, crispy and oily. For appearance, they were: discolored, colored and iridescent (Hyldig 2009b). Each sample was cut in slices (3 mm wide) and three pieces were put into individual Petri dishes for each assessor. All evaluations were performed in separated booths under normal daylight and at ambient temperature.

Statistical Analysis

Statistical tests for difference in antibody titre levels were performed using *t*-test.

The statistical analysis of the sensory data was performed using one-tailed Fisher's exact test in order to compare the experimental groups with control fish. The specific test was used in order to give a statistically qualitative result because of the relatively small sample numbers. The survival curves of the challenged fish were plotted using a Kaplan–Meier plot and significance was tested using a log–rank test. All statistical analysis was performed in the Graphpad Prism 4.0c (Graph-Pad Software, San Diego, CA) for Macintosh software (GraphPad Software, La Jolla, CA).

RESULTS

Challenge by Y. ruckeri

Mortality following challenge by *Y. ruckeri* was initiated on day 3 postchallenge and continued until day 17 postchallenge (Fig. 2). Antibiotics was given from day 6 postchallenge and continued until the mortality had ceased according to veterinarian regulations. A final survival of 63.8 and 59.5% was obtained in vaccinated and naïve fish challenged by *Y. ruckeri*, respectively. No significant difference was seen between these two groups (P > 0.05). No mortality was observed for control fish.

Challenge by V. anguillarum

A number of 149 previously *Y. ruckeri* vaccinated + *Y. ruckeri* challenged + *V. anguillarum-vaccinated* fish were challenged by *V. anguillarum*. Furthermore, 123 previoulsy *Y. ruckeri* vaccinated + *Y. ruckeri* challenged and 124 nonvaccinated fish were challenged by *V. anguillarum*. Mortality was initiated on day 2 following challenge in naïve fish. The survival of this group following challenge by *V. anguillarum* was 82.4% (Fig. 3). This was significantly higher compared with nonvaccinated fish challenged against *Y. ruckeri* (59.5%, *P* < 0.05; see Figs. 2 and 3). In addition, vaccinated fish were protected against the bacterium because only 0.7% in total of vaccinated fish died following challenge (Fig. 3). For fish that were previ-



FIG. 2. CHALLENGE OF RAINBOW TROUT BY YERSINIA RUCKERI Control fish were not challenged. YRV, Yersinia ruckeri vaccinated; YRC, Yersinia ruckeri challenged.



FIG. 3. CHALLENGE OF RAINBOW TROUT BY VIBRIO ANGUILLARUM Control fish were not challenged. YRV, Yersinia ruckeri vaccinated; YRC, Yersinia ruckeri challenged; VAV, Vibrio anguillarum vaccinated; VAC, Vibrio anguillarum challenged.

ously vaccinated against and infected by *Y. ruckeri*, the final survival was 79.0%. Between this group and naïve fish challenged by *V. anguillarum*, there was no significant difference in survival (P > 0.05), but the survival of these two groups was significantly lower than for vaccinated fish (P < 0.05).

Clinical Signs of Infected Fish

Both bacteria gave rise to the classical clinical signs of disease as reported in earlier studies (Poppe and Bergh 1999). For *Y. ruckeri*, bleedings were observed in the region of the mouth and some fish showed bleedings in the fins. Internal signs were pale liver, bleedings and ascites. The clinical signs for *V. anguillarum*-infected fish were seen as external bleedings in the skin, musculature and at the base of the fins. For some fish, bleedings were visible in the musculature the whole way from the skin and into the peritoneal cavity. Internal signs included ascites, pale liver and bleedings in the peritoneal cavity.

Bacteriology following Challenge and Recovery

Following bacterial challenge, subsequent dead fish were immediately screened for the presence of bacteria. Plating of blood on agar plates from either *Y. ruckeri* or *V. anguillarum*infected fish appeared as white colonies. Furthermore, *V. anguillarum* were hemolytic on blood agar (results not shown).

After antibiotic treatment and subsequent recovery, a number (n = 10, subsampling) of previously infected fish were examined for the presence of pathogens before the fish were transferred to the sea cages. All examined fish were stated clear of both *Y. ruckeri* and *V. anguillarum* (results not shown).

Detection of *Y. ruckeri* and *V. anguillarum* by ELISA

The results of the ELISAs for detection of *Y. ruckeri* and *V. anguillarum* showed presence of antibodies against both bacte-





FIG. 4. ANTIBODIES AGAINST YERSINIA RUCKERI

Antibody levels were measured in the experimental groups 1, 3, 4, 6, 8, 9 and 10. The results are shown as absorbance at 450 nm + standard error of the mean. Control fish were not challenged. n = 9 fish in each group analyzed. YRV, *Yersinia ruckeri* vaccinated; YRC, *Yersinia ruckeri* challenged; VAV, *Vibrio anguillarum* vaccinated; VAC, *Vibrio anguillarum* challenged.

ria in all groups of fish (Figs. 4 and 5). For both bacteria, there was no statistical difference between any of the groups in antibody titre (*t*-test, P > 0.05).

Size of Fish at Termination of the Experiment

The fish were reared in the sea cages until slaughter in November 2009. The average weight (g \pm standard deviation [SD]) and length (cm \pm SD) of the fish at slaughter were 667.1 \pm 103.9 and 36.8 \pm 1.8, respectively (n = 111). Furthermore, there was no significant difference in size between fish from the different experimental groups (P > 0.05).

Sensory Analysis

A breakdown of the pump providing the water flow for the land-based fish tanks at Bisserup took place in April 2009. The technical failure resulted in a loss of 757 fish, thus, some of the experimental groups were lost. Because of this, the subsequent sensory analysis was reduced to the experimental groups described in Table 1 (groups 4, 8, 9 and 10). The numbers of fish from each group used for sensory analysis were 8 and 9 for cold-smoked fillets and heat-treated fillets, respectively. Significant differences were seen for fish stored

FIG. 5. ANTIBODIES AGAINST *VIBRIO ANGUILLARUM* Antibody levels were measured in the experimental groups 1, 3, 4, 6, 8, 9 and 10. The results are shown as absorbance at 450 nm + standard error of the mean. Control fish were not challenged. *n* = 9 fish in each group analyzed. YRV, *Yersinia ruckeri* vaccinated; YRC, *Yersinia ruckeri* challenged; VAV, *Vibrio anguillarum* vaccinated; VAC, *Vibrio anguillarum* challenged.

for 7 days on ice before heat-treating regarding texture (Table 1). For these, previously V. anguillarum infected fish (group 4) were significantly less flaky and had a lower oiliness in comparison with control fish (P < 0.001). Furthermore, they had a significantly higher toughness in comparison with control fish (P < 0.01). Fish previously vaccinated against and challenged with Y. ruckeri and later V. anguillarum challenged (group 8) did also have a significantly higher toughness, but also a higher fibrousness in comparison with control fish (P < 0.001). Fish vaccinated and challenged with Y. ruckeri and V. anguillarum (group 9) and mechanically damaged fish (group 10) did not show significant changes for any of parameters analyzed in comparison with control fish (P > 0.05). For heat-treated fish after 3 days of ice-storage and cold-smoked fillets, no difference was seen for any of the parameters tested (P > 0.05, results not shown).

DISCUSSION

The results presented herein establish a linkage between previous infections and sensory-quality parameters in rainbow trout recovered from disease. Fillets from previously infected and mechanically damaged rainbow trout were examined for sensory analysis as heat-treated and smoked in order to

TABLE 1.	SENSORY	ANALYSIS OF	HEAT-TREATED F	ILLETS STORED	ON ICE FOR 7 DAYS
----------	---------	-------------	----------------	---------------	-------------------

		Texture					Appearance			
Group number	Experimental treatment	Flakiness	Juiciness	Firmness	Oiliness	Toughness	Fibrousness	Brightness	Color	Discolor
4	VAC	_	n.s.	n.s	_	+	n.s.	n.s.	n.s.	n.s.
8	YRV + YRC + VAC	n.s.	n.s.	n.s.	n.s.	+	+	n.s.	n.s.	n.s.
9	YRV + YRC + VAV + VAC	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
10	Mechanical damage	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.

Experimental groups shown are 4, 8, 9 and 10. All experimental groups are compared with the control group using one-tailed Fisher's exact test. "+" indicates significantly higher level relatively to control fish (P < 0.001); "-" indicates significantly lower level relatively to control fish (P < 0.001). P > 0.001.

n = 9 fish evaluated in each of the groups.

n.s., nonsignificant; VAV, Vibrio anguillarum vaccinated; VAC, Vibrio anguillarum challenged; YRV, Yersinia ruckeri vaccinated; YRC, Yersinia ruckeri challenged.

examine the usual consumer products industrially processed from that species. The time points of 3 and 7 days of icestorage of heat-treated fillets prior to the sensory analysis were chosen in order to make the results comparable with what could be expected within a natural situation in fish farming. Furthermore, the texture of fish meat is known to undergo changes together with storage time, making it obvious to do measurements at different time points (Faergemand et al. 1995). Only fish that were previously challenged by V. anguillarum and processed as heat-treated were significantly different regarding the texture in comparison with control fish. Fillets from this group were less flaky, had a lower oiliness and a higher toughness in comparison with control fish. Furthermore, fish previously vaccinated against and challenged with Y. ruckeri and later V. anguillarumchallenged had a significantly higher toughness and fibrousness compared with control fish. These results show that previous infection has an impact on the resulting sensory quality of the fish meat from recovered fish. The findings are further supported by recent studies of Atlantic salmon infected or previous infected by PD (Lerfall et al. 2012). Here, the texture, although measured instrumentally, was directly influenced by the infection.

Usually, diseases are giving rise to bleedings and damages in the musculature, which further leads to an inflammatory response as the initial step of the tissue regeneration (Høie 1999; Larsen and Pedersen 1999). This is characterized by influx of immune cells in order to combat the pathogen and clear the infected tissue for dying and dead cells (Diegelmann and Evans 2004). Inflammation together with the later stages of tissue regeneration typically leads to establishment of scarring in affected regions, as seen for farmed Atlantic salmon following infection by *M. viscosa* (Salte *et al.* 1994). Scarring is a result of deposition of collagen between the live cells and this is believed to account for the textural differences between previously infected fish and control fish. The textural characteristic "hardness" has previously been correlated with the amount of collagen in the

meat from the shellfish Abelone (Haliotis discus) (Hatae et al. 1996). In beef meat, a correlation between meat tenderness and the collagen content has been shown, pointing out a significant influence of collagen on the textural characteristics of meat (Riley et al. 2005). The preparation of the heat-treated fillets was performed at a temperature of 100C. Previous experiments with similar treatments of fish meat of the same size have shown that the resulting core temperature thus reach approximately 70C (Green-Petersen and Hyldig 2010). Usually, at much lower temperatures about 35C, fish collagens denature and become soluble (Saito et al. 2001). Thus, it could be believed that the collagen content in meat would not have an influence on textural characteristics when meat is processed at those temperatures. However, in earlier studies it has been shown that also denatured (soluble) collagens in meat contribute to the textural characteristics (Lepetit 2008). The mechanical damage performed may have had limited impact on the fish. A less heavy inflammatory response could be expected following sterile physical damage in comparison with infected fish because of the absence of pathogens in the tissue (Ingerslev et al. 2010). Furthermore, nonvaccinated, V. anguillarumchallenged fish had the highest textural toughness indicating that vaccination protects against disease and thus tissue damage and scarring. The fish used in the study had been reared under conventional farming conditions and could then be compared with a natural situation. The infections of Y. ruckeri and V. anguillarum were performed in a way where challenged fish were treated by antibiotics a few days after mortality was observed in order to mimic the situation during a natural outbreak of disease in fish farming. The natural route of entry by bacterial pathogens in the host is through the outer surfaces such as the skin and gills (Lovoll et al. 2009). However, in this study the fish were i.p. challenged in order to control the dose of bacteria used for infection. Nevertheless, this was not expected to have an influence of the resulting pathological changes in the musculature compared with a natural route of infection because

the bacteria spread systemically through the blood (Høie 1999; Larsen and Pedersen 1999). Because of the small size of the fish when vaccinated against Y. ruckeri, an immersion vaccine was chosen. In contrast to the vaccine used against V. anguillarum, there was no effect of the Y. ruckeri vaccine, which is in line with earlier observations when fish are i.p. challenged following an immersion vaccine (Raida and Buchmann 2008b). In contrast, the fish were i.p. vaccinated against V. anguillarum due to a larger size at the time of vaccination, and for these fish with almost full protection of the vaccine was seen. The ELISA data showed no significant difference to any of the different groups either for V. anguillarum or Y. ruckeri. A survey on the occurrence of typical bacterial pathogens from Danish marine rainbow trout farms has revealed that both species of bacteria are isolated from fish in there (Pedersen et al. 2008). Hence, the observed antibody titres of the bacteria indicate that the fish have been primed from naturally occurring bacteria in the water and because of this have developed antibodies (Ellis 1988). The time from the last infection and antibiotic treatment and until slaughter was about 16 months. Hence, the long-term effects of disease were examined in this study. It could further be expected that the shorter-term effects would be more prominent because the signs of disease are expected to diminish over time (Lerfall et al. 2012).

Overall, from the results it should be concluded that disease has an impact on product quality-associated parameters. The results may be useful knowledge for the fish industry because knowledge about previous history of disease may be used in the choice of strategy regarding processing of the fish following slaughter. Thus, it might be beneficial to use previously infected fish for smoked products. Lastly, the findings underline the importance of effective vaccination in order to prevent subsequent quality changes.

ACKNOWLEDGMENTS

This article was funded by the Ministry of Food, Agriculture and Fisheries in Denmark, grant no. 3310-06-00116. Thanks to Inger Dalsgaard from The Veterinary Institute, Technical University of Denmark for providing the bacterial strains of *Y. ruckeri* and *V. anguillarum*.

REFERENCES

- ASHLEY, P.J. 2007. Fish welfare: Current issues in aquaculture. Appl. Anim. Behav. Sci. *104*, 199–235.
- AUNSMO, A., VALLE, P.S., SANDBERG, M., MIDTLYNG, P.J. and BRUHEIM, T. 2010. Stochastic modelling of direct costs of pancreas disease (PD) in Norwegian farmed Atlantic salmon (*Salmo salar* L.). Prev. Vet. Med. *93*, 233–241.

BENEDIKTSDOTTIR, E., HELGASON, S. and SIGURJONSDOTTIR, H. 1998. Vibrio spp. isolated from salmonids with shallow skin lesions and reared at low temperature. J. Fish Dis. *21*, 19–28.

- BONDAD-REANTASO, M.G., SUBASINGHE, R.P., ARTHUR, J.R., OGAWA, K., CHINABUT, S., ADLARD, R., TAN, Z. and SHARIFF, M. 2005. Disease and health management in Asian aquaculture. Vet. Parasitol. *132*, 249–272.
- COYNE, R., SMITH, P., DALSGAARD, I., NILSEN, H., KONGSHAUG, H., BERGH, O. and SAMUELSEN, O. 2006. Winter ulcer disease of post-smolt Atlantic salmon: An unsuitable case for treatment? Aquaculture 253, 171–178.
- DALSGAARD, I. and MADSEN, L. 2000. Bacterial pathogens in rainbow trout, *Oncorhynchus mykiss* (Walbaum), reared at Danish freshwater farms. J. Fish Dis. *23*, 199–209.
- DIEGELMANN, R.F. and EVANS, M.C. 2004. Wound healing: An overview of acute, fibrotic and delayed healing. Front. Biosci. *9*, 283–289.
- ELLIS, A.E. 1988. Fish Vaccination, Academic Press, San Diego, CA.
- FAERGEMAND, J., RONSHOLDT, B., ALSTED, N. and BORRESEN, T. 1995. Fillet Texture of rainbow-trout as affected by feeding strategy, slaughtering procedure and storage post-mortem. Water Sci. Technol. *31*, 225–231.
- FAO. 2008. The state of world fisheries and aquaculture. http://www.fao.org/docrep/011/i0250e/i0250e00.htm (accessed April 23, 2012).
- GREEN-PETERSEN, D.M.B. and HYLDIG, G. 2010. Variation in sensory profile of individual rainbow trout (*Oncorhynchus mykiss*) from the same production batch. J. Food Sci. 75, S499–S505.
- HATAE, K., NAKAI, H., TANAKA, C., SHIMADA, A. and WATABE, S. 1996. Taste and texture of abalone meat after extended cooking. Fish. Sci. *62*, 643–647.
- HØIE, S. 1999. Yersinose. In *Fiskehelse og Fiskesykdomme* (T. Poppe, ed.) pp. 94–96, Universitetsforlaget, Oslo, Norway.
- HYLDIG, G. 2009a. Sensory aspects of heat treated seafood. In *Handbook of Seafood and Seafood Products Analysis*, (L.M.L. Nollet and F. Toldra, eds.) pp. 499–512, CRC/Taylor & Francis, Boca Raton, FL.
- HYLDIG, G. 2009b. Sensory descriptors. In *Handbook of Seafood and Seafood Products Analysis*, (L.M.L. Nollet and F. Toldra, eds.) pp. 481–497, CRC/Taylor & Francis, Boca Raton, FL.
- INGERSLEV, H.C., LUNDER, T. and NIELSEN, M.E. 2010. Inflammatory and regenerative responses in salmonids following mechanical tissue damage and natural infection. Fish Shellfish Immun. *29*, 440–450.
- ISO 1988. Sensory analysis general guidance for the design of test rooms. Reference numer ISO 8589:1988(E). In.
- ISO 1993. Sensory analysis general guidance for the selection, training and monitoring of assessors. Reference number ISO 8586-1:1993(E). In.
- LARSEN, J.E. and PEDERSEN, K. 1999. Infeksjoner med Vibrio-bakterier. In *Fiskehelse og Fiskesykdommer* (T. Poppe, ed.) pp. 68–83, Universitetsforlaget, Oslo, Norway.
- LEPETIT, J. 2008. Collagen contribution to meat toughness: Theoretical aspects. Meat Sci. *80*, 960–967.

PREVIOUS INFECTION INFLUENCES THE TEXTURE OF FISH MEAT

H.-C. INGERSLEV ET AL.

- LERFALL, J., LARSSON, T., BIRKELAND, S., TAKSDAL, T., DALGAARD, P., AFANASYEV, S., BJERKE, M.T. and MORKORE, T. 2012. Effect of pancreas disease (PD) on quality attributes of raw and smoked fillets of Atlantic salmon (*Salmo salar* L.). Aquaculture 324–325, 209–217.
- LONERGAN, E.H., ZHANG, W.A. and LONERGAN, S.M. 2010. Biochemistry of postmortem muscle – lessons on mechanisms of meat tenderization. Meat Sci. *86*, 184–195.
- LOVOLL, M., WIIK-NIELSEN, C.R., TUNSJO, H.S., COLQUHOUN, D., LUNDER, T., SORUM, H. and GROVE, S. 2009. Atlantic salmon bath challenged with *Moritella viscosa* – pathogen invasion and host response. Fish Shellfish Immun. 26, 877–884.
- LUNDER, T., EVENSEN, O., HOLSTAD, G. and HASTEIN, T. 1995. Winter ulcer in the Atlantic salmon *Salmo salar* – pathological and bacteriological investigations and transmission experiments. Dis. Aquat. Organ. 23, 39–49.
- MCLOUGHLIN, M.F., NELSON, R.T., ROWLEY, H.M., COX, D.I. and GRANT, A.N. 1996. Experimental pancreas disease in Atlantic salmon *Salmo salar* post-smolts induced by salmon pancreas disease virus (SPDV). Dis. Aquat. Organ. 26, 117–124.
- MEILGAARD, M., CIVILLE, G.V. and CARR, T. 1999. Descriptive analysis techniques. In *Sensory Evaluation Techniques*, (M. Meilgaard, ed.) pp. 161–172, CRC Press, New York, NY.
- MERKIN, G.V., ROTH, B., GJERSTAD, C., DAHL-PAULSEN, E. and NORTVEDT, R. 2010. Effect of pre-slaughter procedures on stress responses and some quality parameters in sea-farmed rainbow trout (*Oncorhynchus mykiss*). Aquaculture 309, 231–235.
- MIKKELSEN, H., LINDENSTROM, T. and NIELSEN, M.E. 2006. Effects of temperature on production and specificity of antibodies in rainbow trout (*Oncorhynchus mykiss*). J. World Aquac. Soc. *37*, 518–522.
- MORAN, J.D.W., WHITAKER, D.J. and KENT, M.L. 1999. A review of the myxosporean genus *Kudoa Meglitsch*, 1947, and its impact on the international aquaculture industry and commercial fisheries. Aquaculture *172*, 163–196.
- MORKORE, T., NETTEBERG, C., JOHNSSON, L. and PICKOVA, J. 2007. Impact of dietary oil source on product quality of fanned Atlantic cod, *Gadus morhua*. Aquaculture 267, 236–247.
- PEDERSEN, K., KOFOD, H., DALSGAARD, I. and LARSEN, J.L. 1994. Isolation of oxidase-negative Aeromonas salmonicida from diseased turbot Scophthalmus maximus. Dis. Aquat. Organ. 18, 149–154.
- PEDERSEN, K., SKALL, H.F., LASSEN-NIELSEN, A.M., NIELSEN, T.F., HENRIKSEN, N.H. and OLESEN, N.J. 2008. Surveillance of health status on eight marine rainbow trout, *Oncorhynchus mykiss* (Walbaum), farms in Denmark in 2006. J. Fish Dis. 31, 659–667.

- PETTERSSON, A., JOHNSSON, L., BRANNAS, E. and PICKOVA, J. 2009. Effects of rapeseed oil replacement in fish feed on lipid composition and self-selection by rainbow trout (*Oncorhynchus mykiss*). Aquac. Nutr. 15, 577–586.
- POLI, B.M., PARISI, G., SCAPPINI, F. and ZAMPACAVALLO, G. 2005. Fish welfare and quality as affected by pre-slaughter and slaughter management. Aquac. Int. *13*, 29–49.
- POPPE, T. and BERGH, Ø. 1999. *Fiskehelse og Fiskesykdommer*, Universitetsforlaget, Oslo, Norway.
- RAIDA, M.K. and BUCHMANN, K. 2008a. Development of adaptive immunity in rainbow trout, Oncorhynchus mykiss (Walbaum) surviving an infection with *Yersinia ruckeri*. Fish Shellfish Immun. 25, 533–541.
- RAIDA, M.K. and BUCHMANN, K. 2008b. Bath vaccination of rainbow trout (*Oncorhynchus mykiss* Walbaum) against *Yersinia ruckeri*: Effects of temperature on protection and gene expression. Vaccine 26, 1050–1062.
- RILEY, D.G., JOHNSON, D.D., CHASE, C.C., WEST, R.L., COLEMAN, S.W., OLSON, T.A. and HAMMOND, A.C. 2005. Factors influencing tenderness in steaks from Brahman cattle. Meat Sci. 70, 347–356.
- SAITO, M., TAKENOUCHI, Y., KUNISAKI, N. and KIMURA, S. 2001. Complete primary structure of rainbow trout type I collagen consisting of alpha 1(I)alpha 2(I)alpha 3(I) heterotrimers. Eur. J. Biochem. 268, 2817–2827.
- SALTE, R., RORVIK, K.A., REED, E. and NORBERG, K. 1994. Winter ulcers of the skin in Atlantic salmon, *Salmo salar* L.: Pathogenesis and possible etiology. J. Fish Dis. *17*, 661–665.
- SIGHOLT, T., ERIKSON, U., RUSTAD, T., JOHANSEN, S., NORDTVEDT, T.S. and SELAND, A. 1997. Handling stress and storage temperature affect meat quality of farmed-raised Atlantic salmon (*Salmo salar*). J. Food Sci. 62, 898–905.
- SIGURGISLADOTTIR, S., TORRISSEN, O., LIE, Ø., THOMASSEN, M. and HAFSTEINSSON, H. 1997. Salmon quality: Methods to determine the quality parameters. Rev. Fish. Sci. 5, 223–252.
- STONE, M.A.B., MACDIARMID, S.C. and PHARO, H.J. 1997. Important Health Risk Analysis: Salmonids for Human Consumption. p. 269, MAF Regulatory Authority. New Zealand.
- TOBBACK, E., DECOSTERE, A., HERMANS, K., HAESEBROUCK, F. and CHIERS, K. 2007. Yersinia ruckeri infections in salmonid fish. I. Fish Dis. *30*, 257–268.
- TRIGO, D., ISLAM, R., JOSEFOWITZ, P. and HILDEBRANDT, G. 2008. Structure of muscle fibres in cooked ham products. Fleischwirtschaft 88, 95–98.
- WOLOWINY, J. 2008. Beef quality. Zywn-Nauk. Technol. Ja. 15, 5–22.

Journal of Food Quality •• (2012) ••-•• © 2012 Wiley Periodicals, Inc.

10

PAPER II

β-glucan enriched bath directly stimulates the wound healing process in common carp (*Cyprinus carpio* L.)

Dominika A. Przybylska, Jacob G. Schmidt, Natalia V. Vera-Jimenez, Dieter Steinhagen and Michael E. Nielsen

(Submitted to Fish and Shellfish Immunology)

Abstract

Wound healing is a complex and well-organized process in which physiological factors and immune mechanisms are involved. A number of different immune modulants have been found to enhance the non-specific defence system in vertebrates, among which β -glucans are the most powerful and extensively investigated.

The aim of the present study was to investigate the biological impact of two commercially available β -glucan enriched products on the wound healing process in carp. To investigate the direct effect of beta glucans on wound healing in carp, bath treatment was applied. Throughout the experiment fish were either kept untreated (control), or in a water supplemented with the two different β -glucans. The healing of the wounds was monitored using a visual system. The correlation between wound closure and immune response was investigated by measuring expression of IL-1 β , IL-6, IL-8 and Muc5b genes, and measurement of radical production. Furthermore, systemic versus local reaction was investigated by collection of internal controls. PAMPs/DAMPs stimulation resulted in an inflammatory response by activating IL-1b, IL-6 and IL-8 and local differences in expression pattern dependent on stimulation by DAMPs alone or DAMPs/PAMPs combination. Obtained results clearly showed that beta glucan enriched bath promotes the closure of wound in common carp.

1. Introduction

Skin, together with mucosa forms the first line of defence against pathogens and is an essential protective barrier in aquatic organisms. In skin epithelia, goblet cells, cells responsible for active mucus production are situated. In mucus, foreign particles and microorganisms are entrapped, before imping epithelial surface and cause damage [1]. The main mucus components are water and large filamentous, highly glycosylated glycoproteins called mucins. The additional compounds are proteins, salts, and enzymes.

Mucins and mucus have rheological properties (viscosity and elasticity), which are important for their physiological function [2]. Based on biochemical characterisation, 19 mucin genes have been identified in humans [3], and their gene structure and role in the infection process has been proved [4,5]. Up to date, only two mucin genes, Muc2c and Muc5b were cloned and sequenced in fish [6].

Wounds in fish can be caused by pathogens, such ectoparasites as *Ichthyophthirius multifillis* and lice, Gram-negative bacteria as *Moritella viscosus*, as well as mechanical trauma [7-9]. In vertebrates reduced skin integrity or mucus production facilitates entry of pathogens into the underlying tissue and vascular system, and therefore rapid recruitment of immune cells to the wound site is essential [10]. Wound healing is a complex, well-organized process which can be roughly divided into three overlapping phases: inflammation, tissue formation and remodeling, in which blood cells, soluble mediators, resident cells (fibroblasts, endothelial cells, goblet cells) and extracellular matrix components are involved [10-12].

The immune modulatory effect of β -glucans has been shown in several studies including fish [13-18]. These naturally occurring polysaccharides differ in length, molecular weight, extend of branching and bioactivity, and in vertebrates trigger different biological effects [19-24]. β -glucans have been shown to accelerate the wound healing process in mammals, however that matter has not yet been proven in fish [25, 26].

β-glucans, as pathogen-associated molecular pattern molecules (PAMPs) are recognized by pattern recognition receptors (PRRs) such as toll-like receptors (TLR) or nod-like receptors (NLRs), and activate transcription of pro-inflammatory genes [27]. Further, PRR signaling can also be activated by "danger signals" – damage associated molecular patterns (DAMPs), released endogenous structures such as nucleic acids, heat shock proteins or degraded hyaluronan products, which can bind to TLRs or NLRs [28, 29]. Hence, introducing PAMPs and DAMPs simultaneously may have a combined impact [30].

87

In mammals, in response to PAMPs and DAMPs stimulation, numerous cytokines are secreted, and selectively activate and guide leukocyte subsets during wound healing and tissue repair [12]. A few of cytokines are of high importance during wound healing process. Mammalian interleukin 1 (IL- 1β) was shown to be strongly up-regulated during the inflammatory phase of healing [31, 32]. Main IL- 1β source were polymorphonuclear leukocytes (constitute nearly 50% of all cells at the wound site) and macrophages, as well as some resident cells [12, 31]. In addition, IL-1 was reported to affect mucus release in mice [33]. Interleukin 8 (IL-8) is a known regulator of neutrophil trafficking [12, 32, 34, 35] and augments angiogenesis in mammals [36]. Carp possesses two CXCL8 lineages, which appear to be functional homologs to mammalian IL-8 [37]. Expression of IL-8 in mammals is correlated with interleukin 6 (IL-6); [38]. IL-6 is produced and secreted by macrophage, neutrophils, as well as resident cells [12, 31]. Experiments on diabetic or IL-6 knockout mice have shown drop in neutrophil and macrophage number, as well as a reduction in collagen production and deposition [39, 40]. Orthologous of the mammalian IL-6 members have been identified in fish [41-43] and recently, Varela et al., (2011) have confirmed that zebrafish IL-6 shares structural characteristic with IL-6 from other vertebrates and its expression follows immune stimulation [44].

In vertebrates, besides cytokines production, PAMPs and DAMPs stimulate production and secretion of hydrolases, complement components, and reactive oxygen and nitrogen species (ROS and RNS) by macrophages [45, 46]. Modulation of macrophage phagocytic and respiratory burst activity has been noted in fish in response to β -glucan [47, 48].

The objective of this study was to examine the direct biological effect of β -glucans on the wound healing process in common carp in a controlled environment. This effect was ascertained by monitoring wound contraction dynamics and expression of three pro-inflammatory cytokines: IL-1 β , IL-6 and IL-8, as well as measurement of radical production. Additionally, to enable the involvement of mucus in the tissue healing process, we investigated the expression of Muc5b, the major gel-forming mucin that provides mucus major protective matrix in mammals [49].

2. Materials and methods

2.1 Animals

One-year-old common carp (*Cyprinus carpio* L., R3xR8, WUR, The Netherlands) were kept in 301 tap water aquariums at 21 °C and fed a commercial carp feed (Trouvit, Nutreco) at 2% of their estimated body weight per day. Fish were exposed to a 12-hour light/dark cycle. Prior to experiment, fish were divided into 6 groups (Table 1.), and were either wounded or unwounded. Fish were kept constantly for two weeks in the treatments: tap water: control (C), wounded-control

(W-C), MacroGard supplemented tap water: MacoGard (MG), wounded-MacroGard (W-MG) or 6.3 kDa oat fiber supplemented water: 6.3 kDa (6.3), wounded-6.3 kDa (W-6.3).

2.2 Preparation of glucans

For stock solution preparation, 0.5 g of MacroGard (Biorigin, Brasil) and 6.3 kDa fiber (Scan Oat, Sweden) was dissolved in 500 ml MilliQ water. In order to dissolve soluble β -glucan fractions, solutions of both products were stirred for 1 h at 90 °C and autoclaved (121 °C, 15 min. 1atm.). In the experimental setup, water was changed daily and the β -glucan concentrations were adjusted to 0.1µg/ml.

2.3 Wounding and sampling procedure

Prior to bathing, carp were anesthetized by immersion in 0.01 % benzocaine solution (VetPharm, Belgium) and 5 mm biopsy punch (Miltex, Inc, USA, Figure 1.) was used to mechanically damage areas of the skin and musculature (left side, above the lateral line). At day one, three and 14 post wounding, four individuals from each group were anesthetized and by using an 8 mm biopsy punch, the following tissue samples were collected: 1. muscle tissue from the wounded area 2. muscle tissue from a non-wounded area 3. skin tissue from the wound area 4. skin tissue from non-wounded area 5. muscle and skin tissue from all control fish (Figure 1). Samples were collected and immediately frozen in liquid nitrogen for further analysis. Experiment was conducted by a FELASA accredited scientists and according to Danish legislation.

2.4 RNA isolation and cDNA synthesis

Sampled tissues (50 mg) were homogenized by sonication for 30 s (pulse 2, amplitude 70 %) using a soVCX-130 sonicator (CiAB, Sweden). RNA was isolated using GenElute MammalianTM Total RNA Miniprep Kit (Sigma-Aldrich, Denmark) and subsequently treated with DNase-I (Sigma-Aldrich, Denmark) to remove any genomic DNA. RNA purity and quantity was determined by OD260/280 measurements on a NanoDropTM 2000 spectrophotometer (Thermo Fisher Scientific, Germany). Thus, on 400 ng of total RNA in a 20 µl, random hexamer primed reverse transcription reactions were performed using TaqMan® Reverse Transcription reaction (Applied Biosystems, Denmark). The synthesized cDNA samples were diluted 1:10 in MilliQ water and stored at -20°C for further analysis.

2.5 Real-time quantitative PCR

PCR primer sets, specific for 40S, IL-1B, IL-6 subfamily-like cytokine M17 (further in the text IL-6) and IL-8, were designed using the program Primer3 [50] based on sequences deposited in the GeneBank (referred to as Table 2). The mucin 5b (Muc5b) primer sets were provided by Fish Disease Research Unit from Hannover. The subunit S11 of the ribosomal gene 40S was selected as a reference gene based on previous work (Gonzalez et al, 2006). The assays for examined genes were run using SYBR[®] Green JumpStartTM *Taq* ReadyMixTM (Sigma-Aldrich, Denmark). Quantitive RT-PCR was performed using a Stratagene MX3000PTM real-time PCR system (Integrated Sciences, Australia). Master mix for each PCR run was prepared as follows: 12.5 µl Brilliant SYBR[®] Green JumpStartTM *Tag* ReadyMixTM, 1 µl of each primer and 5.5 µl ultra pure water. Finally 5 µl of diluted cDNA were added to 20 µl of master mix. The cycling conditions for the reaction were: incubation step of 10 min at 95 °C, followed by 45 cycles of 95 °C for 30 s and 60 °C for 1 min. At the end of each reaction, DNA melting curve analysis was performed in order to confirm the specificity of the PCR products. For each primer set, control reactions without cDNA were included in order to detect any non-specific amplification. The expression results were analyzed using the $-2^{-\Delta\Delta Ct}$ method [51]. Data are shown as fold expression relative to non-wounded internal control site, and analysis carried out using Mx ProTM QPCR software (Integrated Sciences, Australia). The threshold values (Ct) were determined manually for each run.

2.6 Isolation of head-kidney (HK) cells and measurement of respiratory burst

Anesthetized fish were bled from the caudal vein. Both head-kidneys (HK) were excised and placed on a 100µm nylon cell strainer (BD Falcon). A cell suspension was obtained by pressing the head-kidneys with a plunger through the cell strainer, and rinsing them with phenol red-free Hank's balanced salt solution with 0.1% gelatine (g-HBSS). The HK cells were counted using a Bürker chamber. Cell viability was assessed by Trypan exclusion and cell concentration was brough to $3X10^{6}$ HKcells/ml in g-HBSS. Subsequently, the respiratory burst activity was measured using luminol-amplified chemiluminescence, modified from the protocol described by Vernho *et al.* (2005); [52]. To elicit the response of respiratory burst, *Aeromonas hydrophila* was incubated overnight. The bacterium was wash twice with HBSS and after its OD₆₀₀ was set to 0.5, 50µl of *A. hydrophila* were added to each well in a white 96-well plate (Sigma P8616), to amplify the chemiluminescence. Subsequently, 40µl of 10mM luminol in 0.2 M borate buffer (pH 9.0) and 100µl of HK cells were added next. The total well volume was adjusted with g-HBSS to 300µl. The chemiluminescence emission of the HK cells was measured with a luminometer (Synergy2, Biotek)

every 3 minutes during 210 minutes at 26°C. Results are expressed as the integral of the relative light units (RLU) recorded by the luminometer (Max RLU).

2.7 Visual analysis of wound closure

Multispectral images of the wounds were acquired using a VideometerLab (Videometer A/S, Hørsholm, Denmark). The wound edges on day 14 post-wounding were outlined and the resulting open wound area determined using MATLAB (The MathWorks Inc., Natick, MA, USA).

2.8 Statistics

A non-parametric Mann-Whitney test was used to compare results from gene expression and respiratory burst. Differences between groups were regarded as significant at p < 0.05. Results from visual wound closure were tested with a two-way ANOVA using the Prism software, version 4.03 (GraphPad Inc., La Jolla, CA, USA).

3. Results

3.1 The visual healing of the wounds

At day 14 after injury, the wound size was significantly (p <0.01) reduced in all β -glucan treated groups in comparison to W-C fish. Both, W-MC and W-6.3 kDa treated fish showed reduced wound size when compared to W-C fish (p<0.01), with higher wound closure ratio in MacroGard-treated group. Figure 3 shows measurements of the wound area two weeks after wounding from the experimental groups.

3.2 Gene expression during wound healing

No significant changes have been observed in non-wounded groups.

3.2.1 Interleukin 1 β (IL-1 β)

In skin collected from W-C group, IL-1 β showed a tendency to increase over time (Figure 4A). In muscle, one day post injury, high and significantly up-regulation (p < 0.05) was noted (Figure 4B).

In skin collected from W-MG group, significant up-regulation (p < 0.05) three days after injury was observed (Figure 4A). No significant changes in muscle were seen (Figure 4B).

In skin collected from W-6.3 group, IL-1 β expression was low but significant (p < 0.05) one day post wounding (Figure 4A). In muscle, high and significant up-regulation (p < 0.05) was observed one day post injury (Figure 4B).

3.2.2 Interleukin 6 subfamily-like cytokine M17 (IL-6)

In skin collected from W-C group, IL-6 showed a tendency for up-regulation three and 14 days post-wounding (Figure 4C). In muscle, significant up-regulation (p < 0.05) was noted one day post injury (Figure 4D).

In skin collected from W-MG fish, significant IL-6 up-regulation (p < 0.05) was seen three days post injury (Figure 4C). In musculature tissue, no significant changes were seen (Figure 4D). In skin collected from W-6.3 fish, a tendency for up-regulation was observed (Figure 4C). In muscle, high and significant up-regulation (p < 0.05) was detected one day after wounding (Figure 4D).

3.2.3 Interleukin 8 (IL-8)

In skin collected from W-C group, three and 14 days post-wounding, high and significant upregulation (p < 0.01) of IL-8 was observed (Figure 4E). In muscle, high and significant upregulation (p < 0.01) was noted one day after wounding (Figure 4F).

In skin collected from W-MG fish, IL-8 showed a tendency to be down-regulated overtime (Figure 4E). In muscle, high and significant up-regulation (p < 0.01) was detected three days post injury (Figure 4F).

In skin collected from W-6.3 fish, a tendency for IL-8 up-regulation was seen 14 days after wounding (Figure 4E). In muscle, significant up-regulation (p < 0.05) was detected three days after wounding (Figure 4F).

3.2.4 Mucin 5b (Muc5b)

In skin collected from W-C fish, a significant up-regulation (p < 0.05) of Muc5b was observed 14 days after wounding (Figure 4G). In muscle, 14 days after injury, a tendency for up-regulation was observed (Figure 4H).

In skin collected from W-MG group, no significant differences in Muc5b expression were observed between wounded and non-wounded sides (Figure 4G). In muscle, a tendency for up-regulation was noted 14 days post wounding (Figure 4H).

In skin collected from W-6.3 group, a significant up-regulation (p < 0.05) in Muc5b expression was observed three days post injury with tendency to be up-regulated 14 days post-wounding (Fig. 4G). In muscle a tendency for up-regulation was seen 14 days after injury (Fig. 4H).

3.3 Measurement of respiratory burst

No significant differences were observed within or between treatments and groups (referred as to Table 3).

4. Discussion

Fish mucosal immunity is dynamically investigated area and new captivating opportunities arise, as described by Zhang et al. (2011) [53]. In contrast to terrestrial vertebrates, fish skin is not keratinised and hence, metabolically active [54, 55]. Skin injury (DAMPs) activates mucosal immune cells as well as epithelial resident cells. It has been proven, not only "typical" immune cells of haematopoietic origin, such as neutrophils or macrophages are involved in wound closure and cytokines production [56]. Fibroblasts have been described in humans as the non-conventional cell type having immune regulation capabilities [57]. Recent studies in human have shown that fibroblasts can not only respond to cytokines, but also generate these signals (IL-1β, IL-4, IL-6, IL-8, IL-10, IL-12, TNF- α and IFN- γ) and influence the behaviour of immunocompetent cells trafficking to the injury site [57, 58]. Several fibroblast cell lines have been established and characterised in fish [59, 60]. Ingerslev et al., (2010) reported that fish fibroblasts are susceptible to DAMPs [56]. In the present study, high up-regulation of IL-1 β and IL-6 in skin samples from W-MG fish three days post injury could be explained by the ability of "moving" fibroblasts to express genes of immune functions before the more potent immune cells have been recruited. Presence of DAMPs/PAMPs could potentiate immune-like performance of fibroblasts, as well as having a direct effect on the resident cells present in skin, such as epithelial cells, resident macrophages and lymphocytes. That hypothesis has been strengthen by Wei et al., (2002) who have presented first evidences that glucan can directly stimulate human fibroblasts [61].

Few studies have emphasized the role of goblet cells and mucus in wound healing process in rodents [11, 62]. Goblet cells differentiation and functioning in mammalian system is affected by mucosal immunity and gel-forming mucins can be regulated by inflammatory cytokines such as interleukins (IL-1 β , -6, -17), tumour necrosis factor α (TNF- α) or exogenous factors [2, 6, 63, 64]. Van der Marel et al., (2012) have shown Muc5b is exclusively expressed in skin and gills of common carp, therefore, we expected, that bath supplemented with beta glucans will have an effect on Muc5b expression in skin [6]. High Muc5b up-regulation three and 14 days post-wounding has,

93

however not been restricted to the wound area, but was a general response of the skin mucosa to the injury. Hence, high local inflammation and IL-1β expression could hamper Muc5b expression at the site of injury. Possibly it could also explain why Muc5b expression was not higher in MacroGard treated group as would expect. Present results are in accordance with work presented by van der Marel co-workers (2010) where water with increased bacterial load did not induce clinical symptoms in carp, however rapid skin mucosa respond have been seen, even if the bacteria involved were considered to be non-pathogenic [65]. In vertebrates, beta glucans and lipopolysaccharide (LPS) work through the same receptors, hence, the mechanism by which LPS, MacroGard and 6.3 kDa oat fiber stimulate Muc5b expression might be alike. Moreover, in humans, Muc5b is one of the largest gel-forming glycoprotein in the body, however cells may differ in the density of mucin granules, suggesting a diversity of the mucins synthesized even by an individual cell [2, 49, 66]. Thus, we can not assure, that beta glucans bath did not have any effect on expression of other mucin granules present in mucous or goblet cells.

On the other hand, murine goblet cells can be stimulated by IL-1β directly, via specific receptors on goblet cells to induce mucus excretion, or indirectly, through stimulation of other cells, as it has been suggested by Cohan and co-workers (1991); [33]. In present study, surprisingly Muc5b expression has been detected in samples collected from muscle, 14 days post wounding, which is in contradiction to results presented by van der Marel et al. (2012); [6]. In vertebrates, cells responsible for the production and maintenance of the protective mucus blanket, goblet cells are exclusively lodged in epithelia [11]. Cheng and Leblond (1974), and further Paulus et al. (1993) and Kanter and Akpolat (2008) have postulated that columnar and goblet cells in mice and rats originate from a common stem cell located at the crypt base [11, 67, 68]. Moreover, Scheffer et al. (1984) have found goblet cells in rat are always lost in the presence of intense inflammation and regulation by mucogenesis in goblet cells was associated with the cessation of inflammation [69]. The same findings were confirmed in rat by Jin-Jang et al., (2003); [70]. In the present study PAMPs stimulated epidermal cells on the wounds edges to migrating on an uncovered surface and full wound closure in groups stimulated with beta glucans was observed 14 days post wounding.

DAMPs activate immune response, but danger signals such RNA/DNA or heat shock proteins are as well involved in myogenesis and have high constitutive expression in the heart and muscles in fish [71]. Immediate increase in IL-1 β , IL-6 and IL-8 cytokines day post-damage in muscle collected from W-C group is likely to be the inflammatory response caused by the release of DAMPs e.g. heat shock proteins. Interestingly, in skin samples from W-C fish, different pattern of pro-inflammatory cytokine expression has been noted. No or low IL-1 β , IL-6 and IL-8 expression has been detected one day after damage, however no gene expression was investigated at earlier

94

time points. It is possible that stored IL-1 β , IL-6 and IL-8 were quickly secreted (hours) in response to injury and subsequently, new amount of these cytokines needed to be produce. The temporally inversed pattern of gene expression in skin/muscle tissue has been seen in the β -glucan supplemented group as well, with change in expression three days post wounding. It is consistent with work done on rodents by Wolk and Danon (1985); [25]. According to authors, in rodents, the most pronounced differences in wound closure, between glucan treated and control took place between 48 and 96 h. These changes include proliferation and arrangement of fibroblasts as well as deposition of the collagenous matrix, and occurred earlier in the glucan treated groups.

Yeast β -glucan is known as a powerful immune modulator and many studies have described its positive effect on various vertebrates [14-18, 25]. In contrast, few studies have reported the effect of small fractions of β -glucans. Bohn and BeMiller (1995) described β -glucans as biological response modifiers, in which activity correlates with branching degree and size [24]. They have reported that less branched glucans with molecular weight of 5 kDa-10 kDa were non-active. Conversely, Tanioka et al. (2011) have shown that barley-derived β -glucan (~2 kDa) stimulates maturation of mouse dendritic cells [72]. In addition, up-regulation (~ x30) of IL-6 expression in marine bone marrow cells has been seen when compared to treatment with 40-70 kDa barely β glucan [72]. Present results show, that both β -glucans, MacroGard and 6.3 kDa oat fiber, promote wound healing process in common carp, however with higher wound closure ratio in group treated with MacroGard. Even though we used the same amount of beta glucans, we cannot say that final concentration of β -glucan was the same in both treatments and therefore, we cannot specify MacroGard as a better immune modifier compared to small beta glucan. Additionally, visual analysis clearly shows significant differences in healing rate between 6.3 kDa oat fiber and control group, suggesting it can be considered as bioactive.

To our knowledge, this is the first investigation of the effect of β -glucans on wound healing in fish, where during the entire experiment, fish have been kept in tap water supplemented with two different β -glucan enriched products, allowing open wounds to stay in direct and constant contact with the β -glucan. The mechanism by which glucan enhance wound closure remains unclear. However, many studies have described the effect of glucans on the fish immune system [14-18, 73]. Wolk and Danon (1985) have emphasized that potential application of glucans to promote wound healing in humans and other heterogenic populations of animals, would require an experimental model in which a wound on one side of the animal could be compared to a symmetrical wound on the other side of the same animal [25]. In our study, internal controls of non-damaged muscle tissue and skin have been taken, which allowed us to follow changes taking

place in wounded and non-wounded areas due to DAMPs/PAMPs stimulation, and subsequently, information about the local versus systemic response in carp.

Overall, our results show that both β-glucan treatments promoted faster wound healing compared to non-treated fish. We concluded that bathing in β-glucans, specifically MacroGard and 6.3 kDa oat fiber, have a direct positive effects on the wound closure in common carp and it can be related to high branching level. We showed the immunological and regenerative response following stimulation with PAMPs and DAMPs in controlled conditions, without the exposure to pathogens. PAMPs/DAMPs stimulation resulted in an inflammatory response by activating IL-1b, IL-6 and IL-8, and local differences in expression pattern dependent on stimulation by DAMPs alone or DAMPs/PAMPs combination. In addition, the absence of marked differences on the respiratory burst activity in head-kidney cells supports the notion of a localized immune response to the site of injury.

5. Acknowledgements

Research leading to this paper was supported by the European Community's Seventh Framework Programme (FP7/2007-2013) under Grant agreement number PITN-GA-2008-214505.

6. References

- Rogers DF. Airway goblet cells responsive and adaptable front-line defenders. European Respiratory Journal 1994; 7: 1690-1706.
- [2] Roussel P and Delmotte P. The diversity of epithelial secreted mucins. Current Organic Chemistry 2004; 8: 413-437.
- [3] Dekker J, Rossen JWA, Buller HA and Einerhand, AWC. The MUC family: An obituary 2002; 27: 126-131.
- [4] McAuley JL, Linden SK, Png CW, King RM, Pennington HL, Gendler SJ, Florin TH, Hill GR, Korolik V and McGuckin MA. MUC1 cell surface mucin is a critical element of the mucosal barrier to infection. Journal of Clinical Investigation 2007, 117: 2313-2324.
- [5] Thornton DJ, Rousseau K and McGuckin MA. Structure and Function of the Polymeric Mucins in Airways Mucus. Annual Review of Physiology 2008; 70: 459-486.
- [6] van der Marel M, Adamek M, Gonzalez SF, Frost P, Rombout JHWM, Wiegertjes GF, Savelkoul HFJ and Steinhagen D. Molecular cloning and expression of two b-defensin and two mucin genes in common carp (Cyprinus carpio L.) and their up-regulation after b-glucan feeding. Fish and Shellfish Immunology 2012; 32:494-501.
- [7] Gonzalez SF, Huising MO, Stakauskas R, Forlenza M, Verburg-van Kemenade LBM, Buchmann K. Real-time gene expression analysis in carp (*Cyprinus carpio* L.) skin: inflammatory responses to injury mimicking infection with ectoparasites. Developmental and Comparative Immunology 2007; 31: 244-254.
- [8] Krkosek M, Connors BM, Ford H, Peacock S, Mages P, Ford JS, Morton A, Volpe JP, Hilborn R, Dill LM and Lewis MA. Fish farms, parasites, and predators: implications for salmon population dynamics. Ecological applications. 2011; 21:897-914.
- [9] Ingerslev HC, Lunder T, Nielsen ME. Inflammatory and regenerative responses in salmonids following mechanical tissue damage and natural infection. Fish and Shellfish Immunology 2010; 29: 440-450.
- [10] Singer AJ and Clark RAF. Cutaneous wound healing. The New England Journal of Medicine 1999; 341: 738-746.
- [11] Kanter M and Akpolat M. Vitamin C protects against ionizing radiation damage to goblet cells of the ileum in rats. Acta Histochemica 2008; 110: 481-490.

- [12] Gillitzer R and Goebeler M. Chemokines in cutaneous wound Healing. Journal of Leukocyte Biology 2001; 69: 513-521.
- [13] Dalmo RA and Bøgwald J. β-glucans as conductors of immune symphonies. Fish Shellfish Immunology 2008; 25: 384-396.
- [14] Yano T, Matsuyama H and Mangindaan REP. Polysaccharide-induced protection of carp Cyprinus-Carpio L. against bacterial infection. Journal of Fish Diseases 1991; 14: 577-582.
- [15] Raa J, Roerstad G, Engstad R and Robertsen B. The use of immunostimulants to increase the resistance of aquatic organisms to microbial infections. In: Shariff IM, Subasinghe RP, Arthur JR, editors. Diseases in Asian Aquaculture II. Manila, Philipines: Fish Health Section, Asian Fisheries Society 1992: 39-50.
- [16] Duncan PL and Klesius PH. Dietary immunostimulants enhance nonspecific immune responses in channel catfish but not resistance to Edwardsiella ictaluri. Journal of Aquatic Animal Health 1996; 8: 241-248.
- [17] Rodríguez I, Chamorro R, Novoa B and Figueras A. β-Glucan administration enhances disease resistance and some innate immune responses in zebrafish (Danio rerio). Fish and Shellfish Immunology 2009; 27: 369-373.
- [18] Zhang Z, Swain T, Bøgwald J, Dalmo RA and Kumari J. Bath immunostimulation of rainbow trout (Oncorhynchus mykiss) fry induces enhancement of inflammatory cytokine transcripts, while repeated bath induce no changes. Fish and Shellfish Immunology 2009; 26: 677–684.
- [19] Williams DL, Mueller A and Browder W. Glucan-based macrophage stimulators Areview of their anti-infective potential. Clinical Immunotherapeutics 1996; 5: 392-399.
- [20] Akramiene D, Kondrotas A, Diziapetriene J and Kevelaitis E. Effects of β-glucans on the immune system. Medicina 2007; 43: 597-606.
- [21] Brown GD and Gordon S. Fungal β–Glucans and Mammalian Immunity. Immunity 2003; 19: 311-315.
- [22] Goodridge HS, Wolf AJ and Underhill DM. Beta-glucan recognition by the innate immune system. Immunological Reviews 2009; 230: 38-50.
- [23] Nakagawa Y, Ohno N and Murai T. Suppression by Candida albicans beta-glucan of cytokine release from activated human monocytes and from T cells in the presence of monocytes. The Journal of Infectious Diseases 2003; 187: 710-713.
- [24] Bohn JA and BeMiller JN. (1-3)-Beta-d-glucans as biological response modifiers: a review of structure-functional activity relationships. Carbohydrate Polymers 1995; 28: 3-14.
- [25] Wolk D and Danon D. Promotion of wound healing by yeast glucan evaluated on single animals. Medical Biology 1985; 63: 73-80.
- [26] Kwon A-H, Qiu Z, Hashimoto M, Yamamoto K and Kimura T. Effects of medicinal mushroom (Sparassis crispa) onwound healing in streptozotocin-induced diabetic rats. The American Journal of Surgery 2009, 197: 503-509.
- [27] Medzhitov R and Janeway C. Innate immune recognition: mechanisms and pathways. Immunological Reviews 2000; 173: 89-97.
- [28] Matzinger P. The danger model: a renewed sense of self. Science 2002; 296: 301-305.
- [29] Seong SY and Matzinger P. Hydrophobicity: an ancient damage-associated molecular pattern initiates innate immune response. Nature reviews. Immunology 2004; 4: 469-478.
- [30] Holtick U, Klein-Gonzalez N and von Bergwelt-Baildon MS. Potential of Toll-like receptor 9 agonists in combined anticancer immunotherapy strategies: synergy of PAMPs and DAMPs? Immunotherapy 2011; 3: 301-304.
- [31] Hübner G, Brauchle M, Smola H, Madlener M, Fässler R and Werner S. Differential regulation of proinflammatory cytokines during wound healing in normal and glucocorticoid-treated mice. Cytokine 1996; 8: 548-556.
- [32] Grellner W, Georg T and Wilske J. Quantitative analysis of proinflammatory cytokines (IL-1beta, IL-6, TNFalpha) in human skin wounds. Forensic Science International 2000; 1-3: 251-264.
- [33] Cohan VL, Scott AL, Dinarello CA and Prendergast RA. Interleukin-1 Is a Mucus Secretagogue. Cellular Immunology 1991; 136: 425-434.
- [34] Rennekampff H-O, Hansbrough JF, Kiessig V, Dore', Sticherling M and Schro J-M. Bioactive Interleukin-8 is expressed in wounds and enhances wound healing. Journal of Surgical Research 2000; 93: 41-54.
- [35] Werner S and Grose R. Regulation of Wound Healing by Growth Factors and Cytokines. Physiological Reviews 2003; 83: 835–870.
- [36] Belperio AJ, Keane MP, Arenberg DA, Addison CL, Ehlert JE, Burdick MD, and Strieter RM. CXC chemokines in angiogenesis. Journal of Leukocyte Biology 2000; 68: 1-8.

- [37] van der Aa LM, Chadzinska M, Tijhaar E, Boudinot P and Verburg-van Kemenade BML. CXCL8 Chemokines in Teleost Fish: Two Lineages with Distinct Expression Profiles during Early Phases of Inflammation. Plos One 2010; 5:Article No.: e12384.
- [38] Nielsen AR and Pedersen BK. The biological roles of exercise-induced cytokines: IL-6, IL-8, and IL-15. Applied Physiology, Nutrition and Metabolism 2007; 32: 833-839.
- [39] Lin Z-Q, Kondo T, Ishida Y, Takayasu T and Mukaida N. Essential involvement of IL-6 in the skin woundhealing process as evidenced by delayed wound healing in IL-6-deficient mice. Journal of Leukocyte Biology 2003; 73: 713-721.
- [40] Berdal M, Appelbom HI, Eikrem JH, Lund Å, Zykova S, Busund L-T, Seljelid R and Jenssen T. Aminated b-1,3-D-glucan improves wound healing in diabetic db/db mice. Wound Repair and Regeneration 2007; 15 825–832.
- [41] Bird S, Zou J, Savan R, Kono T, Sakai M, Woo J and Secombes C. Characterisation and expression analysis of an interleukin 6 homologue in the Japanese pufferfish, Fugu rubripes. Developmental and Comparative Immunology 2005; 29: 775-789.
- [42] Iliev DB, Castellana B, MacKenzie S, Planas JP and Goetz FW. Cloning and expression analysis of an IL-6 homolog in rainbow trout (Oncorhynchus mykiss). Molecular Immunology 2007; 44: 1803-1807.
- [43] Castellana B, Iliev DB, Sepulcre MP, MacKenzie S, Goetz FW, Mulero V, Planas JV. Molecular characterization of interleukin-6 in the gilthead seabream (Sparus aurata). Molecular Immunology 2008; 45: 3363-3370.
- [44] Varela M, Dios S and Novoa B. Characterisation, expression and ontogeny of interleukin-6 and its receptors in zebrafish (Danio rerio). Developmental and Comparative Immunology 2012; 37: 97-106.
- [45] Mosser DM and Edwards JP. Exploring the full spectrum of macrophage activation. Nature Reviews. Immunology 2008; 8: 958-969.
- [46] Novoa B, Figueras A, Ashton I and Secombs CJ. In vitro studies on the regulation of rainbow trout (Oncorhynchus mykiss) macrophage respiratory burst activity. Developmental and Comparative Immunology 1996; 20: 207-216.
- [47] Jørgensen JB, Lunde H and Robertsen B. Peritoneal and head kidney cell response to intraperitoneally injected yeast glucan in Atlantic salmon, Salmo salar L. Journal of Fish Diseases 1993; 16: 313-325.
- [48] Chettri JK, Holten-Andersen L and K Buchmann. Factors influencing in vitro respiratory burst assays with head kidney leucocytes from rainbow trout, Oncorhynchus mykiss (Walbaum). Journal of Fish Diseases 2010; 33: 593-602.
- [49] Kesimer M, Makhov AM, Griffith JD, Verdugo P, and Sheehan JK. Unpacking a gel-forming mucin: a view of MUC5B organization after granular release. American Journal of Physiology 2010; 298: 15-22.
- [50] Rozen S and Skaletsky H. Primer3 on the WWW for General Users and for Biologist Programmers. Methods in Molecular Biology 2000; 132: 365-386.
- [51] Livak KJ and Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2-ΔΔCT Method. Methods 2001; 25: 402-408.
- [52] Verho S, Jarvinen S, Nikoskelainen S and Lilius EM. Biological effect of vaccination can be assessed directly from diluted whole blood of rainbow trout using homologous blood phagocytes as immunosensors. Fish and Shellfish Immunology 2005; 19: 175-183.
- [53] Zhang Y-A, Salinas I, Li J, Parra D, Bjork S, Xu Z, LaPatra SE, Bartholomew J and Sunyer JO. IgT, a primitive immunoglobulin class specialized in mucosal immunity. Nature Immunology 2010; 11: 827-835.
- [54] Martin P. Wound Healing—Aiming for Perfect Skin Regeneration. Science 1997; 276: 75-81.
- [55] Press CM and Evensen Ø. The morphology of the immune system in teleost fishes. Fish and Shellfish Immunology 1999; 9: 309-318.
- [56] Ingerslev H-C, Ossum CG, Lindenstrøm T and Nielsen ME. Fibroblasts Express Immune Relevant Genes and Are Important Sentinel Cells during Tissue Damage in Rainbow Trout (Oncorhynchus mykiss). Plos One 2010; 5: Article No.: e9304.
- [57] Chen B, Tsui S and Smith TJ. IL-1β Induces IL-6 Expression in Human Orbital Fibroblasts: Identification of an Anatomic-Site Specific Phenotypic Attribute Relevant to Thyroid-Associated Ophthalmopathy. Journal of Immunology 2005; 175: 1310-1319.
- [58] Ruiz C, Pérez E, García-Martínez O, Díaz-Rodríguez L, Arroyo-Morales M and Reyes-Botella C. Expression of cytokines IL-4, IL-12, IL-15, IL-18, and IFNg and modulation by different growth factors in cultured human osteoblast-like cells. Journal of Bone and Mineral Metabolism 2007; 25: 286-292.
- [59] Collet B and Secombes CJ. The rainbow trout (Oncorhynchus mykiss) Mx1 promoter. Structural and functional characterization. European Journal of Biochemistry 2001; 268: 1577-1584.
- [60] Ossum CG, Hoffmann EK, Vijaya MM, Holte SE and Bols NC. Characterization of a novel fibroblast-like cell line from rainbow trout and responses to sublethal anoxia. Journal of Fish Biology 2004; 64: 1103-1116.

- [61] Wei D, Zhang L, Williams DL and Browder IW. Glucan stimulates human dermal fibroblast collagen biosynthesis through a nuclear factor-1 dependent mechanism. Wound Repair and Regeneration 2002; 10:161-168.
- [62] Geggel HS, Friend J, and Thofr RA. Conjunctival Epithelial Wound Healing. Investigative Ophthalmology and Visual Science 1984; 25: 860-863.
- [63] Ciacci C, Di Vizio D, Seth R, Insabato G, Mazzacca G, Podolsky DK and Mahide YR. Selective reduction of intestinal trefoil factor in untreated coeliac disease. Clinical and Experimental Immunology 2002; 130: 526-531.
- [64] Chakraborty S, Bonthu N, Swanson BJ and Batra SK. Role of mucins in the skin during benign and malignant conditions. Cancer Letters 2010; 301: 127-141.
- [65] van der Marel M, Caspari N, Neuhaus H, Meyer W, Enss M-Land Steinhagen D. Changes in skin mucus of common carp, Cyprinus carpio L., after exposure to water with a high bacterial load. Journal of Fish Dseases 2010; 33: 431-439.
- [66] Sheehan JK and Thornton DJ. Heterogeneity and Size Distribution of Gel-Forming Mucins. Methods in Molecular Biology 2000; 125: 87-9.
- [67] Cheng H and Leblond CP. Origin, differentiation and renewal of the four main epithelial cell types in the mouse small intestine. The American Journal of Anatomy 1974; 141:461-479.
- [68] Paulus U, Loeffler M, Zeidler J, Owen G, and Potten CS. The differentiation and lineage development of goblet cells in the murine small intestinal crypt: experimental and modelling studies. Journal of Cell Science 1993; 106: 473-484.
- [69] Tseng SCG, Hirst LW, Farazdaghi M and Green WR. Goblet Cell Density and Vascularization during Conjunctival Transdifferentiation. Investigative Ophthalmology and Visual Science 1984; 25: 1168-1176.
- [70] Jin-Jang G, Di-Seng W, Hung-Tu H. Spontaneous remission of edema and regranulation of goblet cells in rat tracheae after capsaicin-induced acute inflammation. Anatomy and Embryology 2003; 206: 301-309.
- [71] Sass JB, Weinberg ES and Krone PH. Specific localization of zebrafish hsp9Oa mFWA to myoD-expressing cells suggests a role for hsp9Ocr during normal muscle development. Mechanisms of Development 1996; 54: 195-204.
- [72] Tanioka A, An W-W, Kuge T, Tsubaki K and Nakaya K. Barley low molecular weight beta-glucan induces maturation of mouse dendritic cells. Anticancer Research 2011; 31: 1647-1652.
- [73] Dalmo RA, Bøgwald J.
 ß-glucans as conductors of immune symphonies. Fish Shellfish Immunology 2008; 25: 384-396.

Figure legends:

Figure 1. Illustration of wounding (A) and sampling (B) procedures, and biopsy punch. Fish were wounded on the left side above lateral line (red circle). During sampling, skin and muscle tissue were taken from the damaged site (red circle), as well as from non-damaged site (internal control, green circle).

Figure 2. Images of the wound area, 14 days post-wounding taken by VideometerLab. Top row – control, middle row – MacroGard, botton row – 6.3 kDa oat fiber. NS – no sample

Figure 3. Statistical representation of the wound sizes 14 days post injury from wounded-control, wounded-MacroGard and wounded-6.3 kDa groups. Bars show mean values (pixels) + SD of n=4 (n=3 14 days in wounded-MacroGard group). * Depicts statistical significance between treatments (* p < 0.05; ** p < 0.01).

Figure 4. Quantitative real-time PCR for mechanically damaged fish. Expression for genes interleukin 1 β (IL-1 β), interleukin 6-subfamily-like cytokine M17 (IL-6), interleukin 8 (IL-8) and mucin 5b (Muc5b) is shown in skin (A, C, E, F) and muscle (B, D, F, H) of common carp. Results are obtained by RQ-PCR and expressed relative to internal control at each time point. The data are normalised relative to 40S and analysed using the -2 $^{-\Delta\Delta Ct}$ method. Bars represent fold expression + SD relative to non-wounded side of n=4 (n=3 14 days in wounded-MacroGard group). cnt – control; ND – not detected. * Depicts statistical significance between site of injury and internal control site (*p < 0.05; **p < 0.01).



Figure 1.





Day 14 post wounding

Figure 3.



Table 1. Experimental groups.

		Water Condition			
		Control	MacroGard	6.3 kDa fiber	
ding	No wound	12	12	12	
Wour	5 mm Biopsy	12	12	12	

Table 2. Sequences of primers used for real-time PCR.

Gene	Primer	Sequence (5' - 3')	GeneBank acc. No.	
405	Forward	GTTGAAGGAAGTGGCAAGGA	A B012087	
405	Reverse	AGAATACGGCCTCTGATGGA	AD012087	
U 16	Forward	AAGGAGGCCAGTGGCTCTGT	A 1245635	
IL-IP	Reverse	CCTGAAGAAGAGGAGGCTGTCA	AJ245055	
IL-6-subfamily-like	Forward	CCGCACATGAAGACAGTGAT	AV102622	
cytokine M17	Reverse	GGGTATATTTGGCTGCAGGA	A 1 102032	
11.8	Forward	TGGAGCTCTTCCCTCCAAG	EU011242	
112-0	Reverse	AGGGTGCAGTAGGGTCCAG	E0011243	
Muc5h	Forward	CAGCCCTCTTCCTCTTTCATC	IE3/13/138	
Wide50	Reverse	CCACTCATCTTTCCTTTCTCTTC	JI ⁻ J4J4J0	
Table 3. Respiratory burst measurement by luminol-amplified chemiluminescence of carp head-kidney leukocytes. The gray colour indicates significance p < 0.05. 1) non-wounded control (1dpw); 2) non-wounded MacroGard (1dpw); 3) non-wounded 6.3 kDa (1dpw); 4) non-wounded control (3dpw); 5) non-wounded MacroGard (3dpw); 6) non-wounded 6.3 kDa (3dpw); 7) non-wounded control (14dpw); 8) non-wounded MacroGard (14dpw); 9) non-wounded 6.3 kDa (14dpw); 10) wounded-control (1dpw); 11) wounded-MacroGard (1dpw); 12) wounded-6.3 kDa (1dpw); 13) wounded-control (3dpw); 14) wounded-MacroGard (3dpw); 15) wounded-6.3 kDa (3dpw); 16) wounded-control (14dpw); 17) wounded-MacroGard (14dpw).

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1																		
2																		
3																		
4																		
5																		
6																		
7																		
8																		
9																		
10																		
11																		
12																		
13																		
14																		
15																		
16																		
17																		
18																		

PAPER III

Intravenous injection of β-glucan in a dose-related manner causes an immunoglobulin class change in skin of common carp (*Cyprinus carpio* L.)

Dominika A. Przybylska and Michael E. Nielsen

(in preparation)

Abstract

 β -glucans, naturally occurring polysaccharides are found in the cell wall of plants, bacteria and fungi. They have been proven to enhance the immune system and pharmacologically they are classified as biological response modifiers (BRM). However, there are also toxicological effects associated with the systemic administration of these agents.

The purpose of this study was to determinate if intravenous (i.v.) injection of Aeromonas hydrophila bacterin (BSK10) with two dosages of MacroGard has a biological effect on immune parameters in carp. Hence, 1 g/ml and 10 g/ml β -glucan in PBS were mixed with bacterin and i.v. injection was performed. To follow stress reaction caused by handling and injection, blood samples from all fish were collected every three days for measurement of cortisol level. Additionally, ratio for specific antibody against A. hdrophila was examined. Furthermore, 4, 12 and 23 days post i.v. injection, skin and head kidney (HK) samples were collected from seven random representatives of each group to determinate gene expression of IL-1β, Hsp70, IgM and IgZ1. Within two days post injection, 65% mortality in a group injected with bacterin + high dose of β -glucan (B-H β g) was noted. No mortality was observed in bacterin injected (B) and bacterin + low β -glucan dose (B-Lßg) injected group, or later during experiment. ELISA for specific antibody against A.s hydrophila showed no significant differences between experimental groups. Results obtained from cortisol assay showed high stress level in B-Hßg group which was correlated with high level of IL-1ß and Hsp70 in skin and head kidney. B-H β g and B-L β g groups showed high level of IgZ1 over IgM in skin. Obtained data indicate that the systemic administration of β-glucan affects immunoglobulin expression in carp skin. Moreover, high dose of MacroGard injected intravenously may contribute to elevating cortisol level and, subsequently to high mortality.

1. Introduction

β-glucans, are well known from their ability to modulate immune system in all organisms from plants to vertebrates, including fish [1-6]. Pharmacologically they are classified as biological response modifiers (BRM; [7]). Depends on the purpose, three ways of administration are available: oral, bath and injection. In the case of fish farms, the most convenient route for antibiotics and immune modulators administration is feed supplementation [3]. On the other hand, bath is an easy and fast route for β -glucans administration to fish in all sizes, and recently, it has been shown that bath in beta glucans supplemented water promotes faster wound closure [3, 8]. Injection is time-consuming and stressful for fish, however it is considered to be the most common and efficient route, due to fact it can reach all internal organs [2]. In fish, the immune system is linked to nervous system, due to double function of head kidney [9]. Fish head kidney is the main hematopoietic and endocrine organ; the major corticosteroid is cortisol, which is produced by the internal cells of the pronefros or head kidney [10-12]. Such organization suggests that the immune processes in fish may be under direct, paracrine hormonal control and, vice versa, chemical signals of the immune system e.g. cytokines could exert direct paracrine action on the interrenal (steroid) and chromaffin (catecholaminergic) cells [12]. That is why immune modulation administered to fish, not only promotes more effective immune response to infectious agents, but also acts upon effects of stress [13]. Teleost fish produce stress signals via their hypothalamus-pituitary-interrenal (HPI) axis that is analogous to the mammalian HPA axis [14]. In response to stressor, fish will undergo a series of biological and physiological changes [15]. In fish, glucocorticoids secretion cause changes in plasma and tissue ion and metabolite levels, hematological features and heat shock proteins (Hsp) and hampers the immune response, resulting in increase of susceptibility to diseases, suppression lymphocytes, macrophages and neutrophils function, decreased antibody production and inflammatory cytokine expression inhibition in vitro, [14, 16-25]. Upon duration and magnitude of stress, fish may be affected at all levels, from biochemical and molecular to population, including ultimately survival [26-28].

In the current study we investigated the biological effect of intravenously injected β -glucan on immune and stress response in common carp. Blood samples were collected from all fish to establish if handling and β -glucan injection together affect cortisol level, as elevated cortisol has profound and differential effects on the immune system [19, 29]. Additionally, ratio for specific anty-*Aeromonas hydrophila* antibody was measured to investigate if systemically introduced β glucan affects B cells in carp blood stream. Furthermore, gene expression of main proinflammatory cytokine IL-1β, damage associated molecular pattern (DAMP) and chaperone protein - heat shock protein 70 (Hsp70), major fish immunoglobulin IgM and recently discovered mucosal immunoglobulin IgZ1 were examined in collected skin and head kidney.

2. Material and methods

2.1. Animals

Sixty-nine, two-year-old common carp (*Cyprinus carpio* L., R3xR8, WUR, The Netherlands, 50 g mean weight) were kept in 120 l tap water aquariums at 25°C and fed a commercial carp feed (Tetra, Melle, Germany) at 2 % of their estimated body weight per day. Fish were exposed to a 12-hour light/dark cycle. Three experimental groups were kept in one aquarium. Additional fin clips were made.

2.1 Preparation of Aeromonas hydrophila bacterin

Aeromonas hydrophila strain BSK10 [30] was cultured in 150 ml of LB medium (Merck, Darmstadt, Germany) overnight at 28°C. Subsequently, 15 ml aliquots were spin at 1800 RCF for 10 min. Supernatant was discarded and pellet was dissolved in PBS. A bacterium was killed through heating for one hour at 60°C. Aliquots were spin down and pellet was re-suspended in low amount of PBS. Subsequently, aliquots were pooled and OD measured at 450nm determined bacterial concentration. Bacterin at final concentration of 10⁶ was injected into fish. A bacterin was plate on agar to confirm no presence of live *Aeromonas hydrophila* cells.

2.2 Preparation of β -glucan

Two doses of MacroGard were chosen: 1g/ml (low dose of β -glucan, L β g) and 10g/ml (high dose of β -glucan, H β g). To obtain high dose of β -glucan, 0.4 g of MacroGard (Biorigin, Sao Paulo, Brasil) was suspended in 10 ml MilliQ water. In order to obtain soluble fractions of β -glucan, solution was stirred for 1 h at 90°C and autoclaved (121°C, 15 min. 1 atm.). To obtain low β -glucan concentration, dilution of high dose β -glucan was made. β -glucan and bacterin solutions were mixed prior to injection.

2.3 Fish and samples recognition

All experimental groups were kept in one aquarium. In order to distinguish individuals during sampling procedure, mathematical program, MatLab has been used, which recognized individuals based on their scales shape.

2.4 Injection and sampling procedure

Fish were acclimatized one week prior to experiment. Two days before i.v. injection, blood samples were collected from all carp. Additionally, skin and head kidney (HK) samples from seven individuals were collected. Prior to injection, fish were anesthetized by immersion in 0.01 % benzocaine solution (VetPharm, Brussels, Belgium). The speciments were divided randomly into three groups of 23 fish and all fish were injected with 100 μ l of: 23 fish with 100 μ l of bacterin in sterile PBS (B); 23 fish with 100 μ l of bacterin + low dose of β -glucan (B-L β g) and, 23 fish with 100 μ l of bacterin + high dose of β -glucan (B-H β g). Blood samples (300 μ l) were collected from all individuals one, 4, 12 and 23 days post injection from caudal vein using heparinized syringes. Plasma was obtain after centrifugation at 15000 g x for 5 min. Tissue samples (skin and HK) were collected from seven random representatives of each group, 4, 12 and 23 days post injection. Samples were collected and kept in RNA later for further analysis. Experiment was conducted by a FELASA accredited scientists and according to Danish legislation.

2.5 RNA isolation and cDNA synthesis

Sampled tissues (50 mg) were homogenized by sonication for 30 s (pulse 2, amplitude 70 %) using a soVCX-130 sonicator (CiAB, Stocholm, Sweden). RNA was isolated using GenElute MammalianTM Total RNA Miniprep Kit (Sigma-Aldrich, Glostrup, Denmark) and subsequently treated with DNase-I (Sigma-Aldrich, Glostrup, Denmark) to remove any genomic DNA. RNA purity and quantity was determined by OD260/280 measurements on a NanoDropTM 2000 spectrophotometer (Thermo Fisher Scientific, Schwerke, Germany). Consequently, random hexamer primed reverse transcription reactions were performed on 400 ng of total RNA in a 20 μ l setup using TaqMan® Reverse Transcription reaction (Applied Biosystems, Naerum, Denmark). The synthesized cDNA samples were diluted 1:10 in MilliQ water and stored at -20°C for further work.

2.6 Real-time quantitative PCR

PCR primer sets, specific for 40S, IL-1β, IgM, and IgZ1 were designed using the program Primer3 [31] based on sequences deposited in the GeneBank (referred to as Table 2). The subunit S11 of the ribosomal gene 40S was selected as a reference gene based on previous work [32]. The assays for examined genes were run using SYBR[®] Green JumpStartTM *Taq* ReadyMixTM (Sigma-Aldrich, Glostrup, Denmark). Quantitive RT-PCR was performed using a Stratagene MX3000PTM real-time

PCR system (Integrated Sciences, Willoughby, Australia). Master mix for each PCR run was prepared as follows: 12.5 μ l Brilliant SYBR[®] Green JumpStartTM *Taq* ReadyMixTM, 1 μ l of each primer and 5.5 μ l ultra pure water. Finally 5 μ l of diluted cDNA were added to 20 μ l of master mix. The cycling conditions for the reaction were: incubation step of 10 min at 65 °C, followed by 45 cycles of 65 °C for 30s and 60 °C for 1 min. At the end of each reaction, DNA melting curve analysis was performed in order to confirm the specificity of the PCR products. For each primer set, control reactions without cDNA were included in order to detect any non-specific amplification. The expression results were analyzed using the -2 ^{- $\Delta\Delta$ Ct} method [33]. Data are shown as fold expression relative to non-wounded internal control site, and analysis carried out using Mx ProTM QPCR software (Integrated Sciences, Australia). The threshold values (Ct) were determined manually for each run. All samples were blind-tested.

2.7 ELISA assay for cortisol

Ninety-six well plates for cortisol ELISA (Demeditec Diagnostics GmbH, Kiel, Germany) were used following manufacturers' protocol. The absorbance was measured at 450 nm using a Synergy 2 Multi Mode microplate reader (BioTek Instruments, Winooski, USA). Samples were analyzed in triplicates and blind.

2.8 ELISA assay for detection of Aeromonas hydrophila

An ELISA assay was used to detect the presence of the antibody against *Aeromonas hydrophila* (BSK10) in the plasma collected from fish. All samples were blind-tested, run in triplicates according to protocol described by [34]. In the following, all steps in the protocol were carried out at room temperature. Antigen (*A. hydrophila*) was prepared as described in 2.2 section. Initially, the wells of microtiter plates (Immunosorb, Nunc, Roskilde, Denmark) were coated using 200 μ l of antigen solution diluted in coating buffer (10⁶ bacterin / well) and further incubated at 4 °C overnight. An unbound bacterin was removed by three washes in PBS-Tween. Subsequently, the plates were blocked using 200 μ l of 0.5% bovine serum albumin (BSA) in PBS and shaken for 15 minutes. The wells were then further washed three times in PBS-Tween and frozen in -20°C for further work.

To each well, 100 μ l of carp serum diluted 1:160 in PBS with 0.1% BSA was added and the plates were gently shaken for 30 minutes. Wells were then washed in order to remove unbound plasma. Further, 100 μ l of rabbit-anti carp Ig diluted 1:2000 in PBS with 0.1% BSA (Sigma Aldrich) was added to the wells and the plates were gently shaken for 10 minutes followed by washing step. Subsequently, 100 μ l of HRP conjugated goat-anti rabbit antibody (BioRad, Copenhagen,

Denmark) diluted 1:3000 in PBS with 0.1% BSA was added to the wells followed by shaking for 10 minutes. The wells were then washed three times. Lastly, 100 μ l of enzyme substrate (TMB; Sigma Aldrich, Glostrup, Denmark) was added to the wells and the plates were incubated for 5 minutes. The reaction was stopped using 50 μ l of stop solution (0.5M H₂SO₄). The absorbance was when measured at 450 nm using a Synergy 2 Multi Mode microplate reader (BioTek Instruments, Winooski, USA).

2.9 Statistics

Non-parametric Mann-Whitney test was used to compare experimental groups. Differences between groups were regarded as significant at p < 0.05. Survival analysis was done with Kaplan-Meier estimator [35].

3 Results

3.1 Mortality

One day post i.v. injection, in a group injected with high dose of β -glucan 26% mortality was observed and raised up to 65% two days post injection (Figure 1). No mortality was observed either in B-injected or B-L β g-injected group, or further during experiment.

3.2 ELISA assay for cortisol

Values from all experimental groups are presented in Table 2. Significantly elevated plasma cortisol was observed in all experimental groups (B-injected: $463,51 \pm 46,90$; B-L β g: 722,86 \pm 16,55; B-H β g: 1158,78 \pm 13,80) one day post i.v. injection (Figure 2). The significant difference (p < 0.001) in cortisol level between the B-injected and B-H β g group, as well as between B-L β g and B-H β g group (p < 0.01) was observed one day post injection (Figure 2). In all experimental groups, cortisol level went back to the basal level 12 days post injection and stayed stable till the end of experiment (Figure 2).

3.3 ELISA assay for detection of Aeromonas hydrophila

The level of specific antibodies against *Aeromonas hydrophila* was measured blind, in triplicates from every single fish from all experimental groups. Results have shown presence of antibodies against *Aeromonas hydrophila* in all experimental groups, with clear peak in antibody ratio, four days post i.v. injection, although no significant difference in antibody titer between treatments was observed (Figure 3).

3.4 Gene expression

3.4.1 Interleukin 1 β (IL-1 β)

In skin and HK samples collected from B-L β g group a tendency for up regulation was observed 23 days post injection (Figure 4A and Figure 4E). In skin and HK collected from B-H β g high, significant (p < 0.001) up regulation was detected 12 days post injection (Figure 5A and Figure 5F subsequently).

3.4.2 Heat shock protein 70 (Hsp70)

In skin collected from B-L β g injected group, no significant changes were observed at any time point, although tendency for up regulation was observed 12 days post injection (Figure 4B). In HK, significant up regulation was observed four (p < 0.01) and 12 days (p < 0.001) post injection (Figure 4F). In skin collected from B-H β g injected fish, significant (p < 0.001) up regulation was observed four and 12 days post injection (Figure 5B). In HK, significant (p < 0.001) up regulation was observed four days after injection (Figure 5F).

3.4.3 Immunoglobulin M (IgM)

In skin collected from B-L β g group no significant changes in IgM expression at any time point were observed (Figure 4C). In HK, significant (p < 0.01) down regulation was noted 12 days post injection (Figure 4G). In samples collected from B-H β g group, significant (p < 0.05) down regulation in HK four days after injection was noted (Figure 5G).

3.4.4 Immunoglobulin Z1 (IgZ1)

High and significant (p < 0.001) up regulation was observed 12 and 23 days post injection in skin and HK samples collected from B-L β g group (Figure 4D and 4H respectively). High and significant (p < 0.01) up regulation was observed four days post injection in skin and HK samples collected from B-H β g (Figure 5D and 5H respectively). Significant (p < 0.001) IgZ1 up regulation was seen in skin 12 days post injection (Figure 5D).

4. Discussion

An intriguing difference between mammals and fish is the tolerance of fish to high concentration of endotoxins and resistance to septic shock and [36-39]. Simply, fish live in water and therefore intimate contact with a potentially high amount of microorganisms. If a high inflammatory reaction

was triggered after each contact with external putative pathogens, fish simply could not survive [40]. Beta glucans are biological extracts and pharmacologically they are classified as biological response modifiers (BRMs; [7]). The immune modulatory effect of β -glucans has been shown in several studies including fish [2-6, 8, 41]. However, in vertebrates, there are some side and toxicological effects associated with the systemic administration of these agents. Long-term feeding regime of peptido-glucans decreased the immune response in rainbow trout when challenged with *Vibrio anguillarum*, as well as in catfish, indicating a negative feedback effect of β -glucan [42-44]. In the current study we observed 65% mortality two days post injection in B-Hβg-injected group. No mortality was observed either in B-injected or B-Lßg-injected group, or further during experiment. Such high mortality in only one experimental group suggests toxicological, doserelated effect of β-glucan. Strong stress reaction caused by handling and injection could be additionally boosted with high dose of injected β -glucan, which is recognized as pathogen associated molecular pattern (PAMP). In vertebrates, effects of immune modulation occur due to increased activity of macrophages, neutrophils and other immunocytes [3, 45-47]. High dose of injected β -glucan together with bacterin potentially overwhelmed the immune system of carp. Moreover, Williams and co-workers has showed in a mice model, that intravenous administration of glucan water-insoluble micro particulates is associated with hypertrophy and hyperplasia of macrophage-rich organs such as liver, lung and spleen [48-50]. Furthermore, many of the BRMs were proved to exhibit significant toxicity in mammals, including vasodilatation, microvascular hemorrhage and circulatory collapse [51-53]. Interestingly, it was reported that if water-insoluble forms of β-glucans are converted to a water-soluble form, the activity of murine immune system is preserved, but the undesirable side effects are eliminated [43, 54].

In the present study i.v. injection with high dose of β -glucan was found to be fish stressor factor which potentially could elevate cortisol level and in subsequently, contribute to high mortality seen in one of the experimental groups (B-H β g). It is in agreement with results presented by Williams et al., (1988) where i.v. injection of soluble glucan in doses ranging from 40 to 1000 mg/kg did not induce mortality, appearance or behavioral changes in mice or rats [54]. In fact, preliminary studies in that laboratory have indicated that the systemic administration of soluble glucan in the dose of 2000 mg/kg did not induce any side effects in mice and rats (unpublished data). In contrary to that are results presented by Maeda et al. (1984) that showed vascular dilatation and hemorrhage in mice 24 hours following a single i.p. injection of 10 mg/kg [51]. However, in rat model, intraperitoneal injection is being considered more effective route of drugs administration [55]. Results of the present study showed that dose 10 g/ml intravenously injected into carp elevates cortisol level and increase expression of Hsp70. Heat shock proteins, have been recognized among the mammalian primary defense cellular mechanisms during acute stress [56-59]. Moreover, in mammals, Hsp70 is one of the "danger signals" – damage associated molecular patterns (DAMPs), released endogenous structures that can activate immune reaction. [60]. We believe, that faster and higher reaction in B-Hβg injected group is dose-related.

Fish head kidney is an intriguing link between immune and endocrine systems [10-12]. HK is also main source of macrophages and neutrophilic granulocytes and therefore, actively participates in the clearance of antigens from bloodstream [61,62]. Chadzinska et al., (2008) reported migration of inflammatory leukocytes from HK into peritoneum during intraperitoneal injection of zymosan particles. Constitutive expression of IL-1ß in peritoneal leukocytes (PTL) was two-fold higher than in HK, however during zymosan-induced peritonitis, IL-1 β expression showed twice higher expression in peritoneal leukocytes (PTL) than in HK 6 hours after injection. IL-1β peak in HK was observed 96 hours post injection [63]. In our study, we observed a tendency for IL-1β up-regulation in B-Lßg injected group, and significant up-regulation 12 days post injection in B-Hßg injected group. Besides HK, we measured IL-1ß expression in skin and raising overtime expression in B-Lßg, as well as high expression in B-Hßg group, could strengthen hypothesis of leukocytes leaving HK and locate other tissues. Moreover, fish from B-H β g injected group appeared to have skin and muscle damage (personal observation) which could enhance IL-1ß expression in response to DAMPs. On the other hand, in vitro studies in common carp showed inhibition of inflammatory cytokine due to acute stress, to prevent damage due to an excessive inflammatory response [64]. In mammals, corticosteroids often have inhibitory effect on monocytes and macrophages, what could explain no changes in IL-1^β expression [65]. *In vitro* study by showed that proliferation of RTS11, a monocyte/macrophage rainbow trout cell line is inhibited by cortisol [66,67]. As macrophages are main source of IL-1β, inhibition in their proliferation could effectively inhibit IL-1β expression. Furthermore, it cannot be excluded that, IL-1ß is main pro-inflammatory cytokine, which reaches peak within minutes after stimulation, so measurement 96 hours after stimulation does not give clear picture.

Aeromonas hydrophila is the most common Gram-negative bacteria present in all freshwater environments that cause a wide range of diseases in fish [30]. There are more than 120 serotypes of *A. hydrophila* described, however it does not usually cause problems in fish populations under normal conditions: infection with *A. hydrophila* in fish appears as a secondary infection, under environmental or physiological stress, injury or infection with other pathogens [68, 69]. It is in accordance with work presented by van der Marel and co-workers (2010) where water with increased bacterial load did not induce clinical symptoms in carp, however rapid skin mucosa respond have been seen, even if the bacteria involved were considered to be non-pathogenic [70].

117

The present study revealed that intravenous injection of dead *A. hydrophila* bacteria alone or together with β -glucan did not elevate production of antibodies against *A.hydrophila* as was originally expected. Handling, injection and dose of injected β -glucan caused stress reaction in all groups, and we expected it would have an effect on specific antibody production. According to Selvaraj et al. (2005) the minimum amount of glucan required for enhanced production of antibodies against *A.hydrophila* was 500 µl/ fish [71]. Following that, we expected group injected with 10 g/ml/ fish, to have significantly higher antibody titer in comparison to bacterin or 1g/ml/ fish injected group. Surprisingly, the basal level of antibody (2 days pre to i.v. injection) was high in all three groups, with no difference among them. This draws a conclusion of fish being previously primed with *A.hydrophila* and, as a consequence, approximately high antibody titer was produced. Elevation in specific antibody level four days after injection, could be a short-term alert in response to injected PAMPs (bacterin and beta glucan) and slightly affect antibody production.

Up to date, only IgM has been shown to be functionally involved in protective immunity in fish [72,73]. B cells comprise 25-35% of all blood lymphocytes in rainbow trout, 50% in puffer fish and 70% in common carp [11,74,75]. Recently described IgT/IgZ seem to be involved in fish mucosal immunity [6, 76, 77]. Ryo et al. (2010) have shown that the highest induction of IgZ1 mRNA expression was seen during the blood parasite infection, T. borreli, as well as during vaccination with V. anguillarum antigens, which are rich in lipopolysaccharides [77]. In vertebrates, beta glucans and lipopolysaccharide (LPS) work through the same receptors, hence, we believe, the mechanism by which IgZ1 expression was triggered in T. borreli infection could be alike while injecting beta glucans. Moreover, Ryo et al. (2010) reported that IgZ1 functions as antibody that neutralizes foreign antigens in the fish blood [77]. It could explain high expression of IgZ1 in current study – if mechanisms triggered by LPS and β -glucan are alike, IgZ1 level should increase to clear circulation from injected β-glucan. In teleost, IgM and IgZ are mutually exclusive on Bcells as reported by Li et al (2010; [78]). More evidence is accumulating that, the antibodies in the mucus, although antigenically identical to the serum Igs, may differ in other aspects: Lobb and Clem (1981b) identified three types of Igs in the mucus of the Archosargus probatocephalus sheepshead bile: one type of tetrameric high molecular weight Ig, identical to the serum counterpart and two types of low molecular weight dimeric Ig [79]. The existence of three different types of Igs seems to suggest three distinct ways of producing them, systemically or locally. Our results indicate absence of IgM expression, with high IgZ1 expression in skin and HK samples collected from B- $L\beta g$ injected group. High expression of IgZ1 in skin, with lower but significant up-regulation in HK, indicates presence of resident B-cells and/or antibody-secreting cells (ASC) in skin of common carp, which were not affected by high cortisol level. The existence of two distinct B cell

lineages, with mutuall expression either IgM or IgZ, was suggested. In 2010, Zhang and co-workers provided evidence for the existence of a previously unrecognized B cell lineage that expressed only IgT [80]. During infection with rainbow trouts' gut parasite Ceratomyxa Shasta substantial upreguation (~733-fold) of IgT in gut was observed, with almost negligible upregulation of IgM expression. Such a high expression of IgT in gut and protective role against gut parasite indicates, that it is the first nontetrapod immunoglobulin specialized in mucosal immunity [80]. In the current study we observed high expression of IgZ1 in skin 12 and 23 days post injection (injection with B-L β g; ~390- and ~310-fold respectively) with no significant changes in IgM expression. Taking under consideration mucosal functions of IgT(Z) reported by Zhang el al. (2010) and further by Ryo et al., (2011), high IgZ1 expression detected in the injection experiment leads to the conclusion that $IgT(Z)^+$ B cells are major B cells in skin of common carp, with low cell count for IgM^+ B cells. Zhang et al., (2010) found the $IgT(Z)^+$ B cell population constituted 16-28% of all trout B cells in the blood, spleen, head kidney and peritoneal cavity, while in gut they represented the main B cell subset. Results obtained from the injection experiment show that IgZ1 expression was significantly higher in skin than in common carp head kidney, suggesting that $IgT(Z)^+ B$ cells have the majority among cells in carp skin, or β -glucan injection enhanced proliferation of these cells. Moreover, IgT(Z)⁺ B cells seem to respond to β -glucan stimulation, which suggests presence of the receptor(s) through which, any actions with β -glucan will be more pronounced in fish skin.

Three routes of antibiotics, vitamins, immune modulators and others administration are possible: injection, oral and bath. The best one should be chosen depends on the aim of experiment. However, we can observe an interesting pattern when it comes to beta glucans administration in common carp. Despite of the administration route, effects can be observed in carp skin. Van der Marel et al. (2012) have reported that β -glucan feeding regime had an effect on the expression of mucus-related genes in carp. In addition to the mucin encoding genes, β -glucan feeding affected the expression of carp β -defensins genes, which were significantly higher when compared to group fed without β -glucan [81]. This suggests that, by feeding β -glucans, not only gut mucosal system is affected, but also changes in skin mucosa might be an important to monitor fed-related improvement of fish health. Recently, Przybylska et al. (submitted) showed that β -glucan enriched bath, specifically MacroGard and 6.3 kDa oat fiber have a direct positive effects on the wound closure in common carp and promote faster wound healing compared to non-treated fish [8]. Results of the current study show that intravenous injection of β -glucan, once again affects skin mucosa.

In conclusion, we show that effect of direct β -glucan stimulation by intravenous injection in common carp is rigorously correlated with dosage. Moreover, high amount of particulate β -glucan

introduced to circulation system of carp elevates cortisol level, and at high dose can be toxic for fish. Furthermore, systemic administration of β -glucan affects immunoglobulin expression in carp skin with immunoglobulin switch-like process, where presence of main fish immunoglobulin IgM is overtaken by recently described, mucosal immunoglobulin IgZ1.

5 Acknowledgements

Research leading to this paper was supported by the European Community's Seventh Framework Programme (FP7/2007-2013) under Grant agreement number PITN-GA-2008-214505 and Ministry of Food, Agriculture and Fisheries in Denmark, grant no. 3310-FVFP-08-K-08-01.

6 Literature

- Rosenberger RF. Control of teichoic and teochuronic acid biosynthesis in Bacillus-subtilis 168TRP Evidence for repression of enzyme synthesis and inhibition of enzyme activity. Biochemica and Biophysica Acta, 1976; 428: 516-524.
- [2] Yano T, MatsuyamaH and Mangindaan REP. Polysaccharide-induced protection of carp Cyprinus-Carpio L. against bacterial infection. Journal of Fish Diseases 1991; 14: 577-582.
- [3] Raa J, Roerstad G, Engstad R and Robertsen B. The use of immunostimulants to increase the resistance of aquatic organisms to microbial infections. In: Shariff IM, Subasinghe RP, Arthur JR, editors. Diseases in Asian Aquaculture II. Manila, Philipines: Fish Health Section, Asian Fisheries Society 1992: 39-50.
- [4] Duncan PL and Klesius PH. Dietary immunostimulants enhance nonspecific immune responses in channel catfish but not resistance to Edwardsiella ictaluri. Journal of Aquatic Animal Health 1996; 8: 241-248.
- [5] Rodríguez I, Chamorro R, Novoa B and Figueras A. β-Glucan administration enhances disease resistance and some innate immune responses in zebrafish (Danio rerio). Fish and Shellfish Immunology 2009; 27: 369-373.
- [6] Zhang Z, Swain T, Bøgwald J, Dalmo RA and Kumari J. Bath immunostimulation of rainbow trout (Oncorhynchus mykiss) fry induces enhancement of inflammatory cytokine transcripts, while repeated bath induce no changes. Fish and Shellfish Immunology 2009; 26: 677–684.
- [7] Bohn JA and BeMiller JN. (1-3)-Beta-d-glucans as biological response modifiers: a review of structure-functional activity relationships. Carbohydrate Polymers 1995; 28: 3-14.
- [8] Przybylska DA, Schmidt JG, Vera-Jimenez NI, Steinchagen D and Nielsen ME. Submitted to Fish and Shellfish Immunology.
- [9] Manning MJ. Fishes. In: Immunology: A Comparative Approach (ed. by RT Turner), John Willey and Sons, Chichester, 1994; 69-100.
- [10] Weyts FAA, Flik G and Verburg-van Kemenade BML. Cortisol inhibits apoptosis in carp neutrophilic granulocytes. Developmental and Comparative Immunology, 1998a; 22: 563-572.
- [11] Weyts FAA, Flik G, Rombout JHWM and Verburg-van Kemenade BML. Cortisol induces apoptosis in activated B cells, not in other lymphoid cells of the common carp, Cyprinus carpio L. Developmental and Comparative Immunology, 1998; 22: 551-562.
- [12] Flik G, Klaren PHM, Van den Burg EH, Metz JR and Huising MO. Review: CRF and stress in fish. General and Comaprative Endocrinology, 2006; 146: 36–44.
- [13] Anderson DP. Immunostimulants, adjuvants and vaccine carriers in fish: application to aquaculture. In: Faisal M and Hetrick FM (eds) Annual review of fish diseases. Pergamon Press, 1992; 281–307.
- [14] Stolte EH, Nabuurs SB, Bury NR, Sturm A, Flik G, Savelkoul HFJ and Verburg-van Kemenade LBM. Stress and innate immunity in carp: Corticosteroid receptors and pro-inflammatory cytokines. Molecular Immunology 2008; 46: 70-79.
- [15] Barton BA. Stress in Fishes: A Diversity of Responses with Particular Reference to Changes in Circulating Corticosteroids. Integrative and Comparative Biology 2002; 42: 517-525.
- [16] Pickering AD. Endocrine-induced pathology in stressed salmonid fish. Fisheries Research 1993; 17: 35-50.
- [17] Iwama GK, Takemura A and Takano K. Oxygen consumption rates of tilapia in fresh water, sea water, and hypersaline sea water. Journal of Fish Biology 1997; 51: 886-894.
- [18] Iwama GK, Vijayan M, Mazur CF and Ackerman P. Heat shock proteins and physiological stress in fish. American Zoologist 1999; 39: 901-909.
- [19] Mommsen TP, Vijayan MM and Moon TW. Cortisol in teleosts: dynamics, mechanisms of action, and metabolic regulation. Reviews of Fish Biology and Fisheries 1999; 9: 211-268.

- [20] Ainsworth AJ, Dexiang C and Waterstrat PR. Changes of peripheral blood leukocytes percentages and function of neutrophils in stressed channel catfish. Journal of Aquatic Animal Health 1991; 3: 41-47.
- [21] Stolte EH, Chadzinska M, Przybylska D, Flik G, Savelkoul HFJ and Verburg-van Kemenade BML. The immune response differentially regulates Hsp70 and glucocorticoid receptor expression in vitro and in vivo in common carp (Cyprinus carpio L.). Fish and Shellfish Immunology 2009; 27: 9-16.
- [22] Verburg-van Kemenade BML, Nowak B, Engelsma MY and Weyts FAA. Differential effects of cortisol on apoptosis and proliferation of carp B-lymphocytes from head kidney, spleen and blood. Fish and Shellfish Immunology 1999; 9: 405-415.
- [23] Saelij JPJ, Verburg-van Kemenade LBM, van Muiswinkel WB and Wiegertjes GF. Daily handling stress reduces resistance of carp to Trypanoplasma borreli: in vitro modulatory effects of cortisol on leukocyte function and apoptosis. Developmental and Comparative Immunology 2003; 27: 233-245.
- [24] Elenkov IJ and Chrousos GP. Stress system-organization, physiology and immunoregulation. Neuroimmunomodulation 2006; 13: 257-267.
- [25] Fast MD, Johnson SC and Jones SRM. Differential expression of the pro-inflammatory cytokines IL-1β-1, TNFα-1 and IL-8 in vaccinated pink (Oncorhynchus gorbuscha) and chum (Oncorhynchus keta) salmon juveniles. Fish and Shellfish Immunology 2007; 22: 403-407.
- [26] Wedemeyer GA and McLeay DJ. Methods for determining the tolerance of fishes to environmental stressors. Academic Press, New York. Pickering (ed.) 1981, Stress and fish 247–275.
- [27] Wedemeyer GA, Barton BA, and McLeay DJ. Stress and acclimation. American Fisheries Society, Bethesda, Maryland In C. B. Schreck and P. B. Moyle (eds.) 1990, Methods for fish biology: 451–489.
- [28] Adams SM. Biological indicators of stress in fish. American Fisheries Society Symposium Series 8, Bethesda, Maryland. 1990.
- [29] Wendelaar BSE. 1997. The stress response in fish. Physiology Reviews 1997; 77: 591-625.
- [30] Nielsen ME, Høi L, Schmidt AS, Qian D, Shimada T, Shen JY and Larsen Jl. Is Aeromonas hydrophila the dominant motile Aeromonas species that causes disease outbreaks in aquaculture production in the Zhejiang Province of China? Diesase of Aquatic Organisms 2001; 46: 23–29.
- [31] Rozen S and Skaletsky H. Primer3 on the WWW for General Users and for Biologist Programmers. Methods in Molecular Biology 2000; 132: 365-386.
- [32] Gonzalez SF, Buchmann K and Nielsen ME. Real-time gene expression analysis in carp (*Cyprinus carpio* L.) skin: Inflammatory responses caused by the ectoparasite*Ichthyophthirius multifiliis*. Fish and Shellfish Immunology 2007; 22: 641-650.
- [33] Livak KJ and Schmittgen TD. Analysis of Realative gene Expression Data Using Real-Time Quantitative PCR and the 2- ΔΔCT method. Methods 2001; 25: 402-408.
- [34] Sigh J and Buchmann K. Comparison of immobilization assays and enzyme-linked immunosorbent assays for detection of rainbow trout antibody-titres against Ichthyophthirius multifiliis Fouquet, 1876. Journal of Fish Diseases 2001; 24: 49-51.
- [35] Kaplan EL and Meier P. Nonparametric estimation from incomplete observations. Journal of The American Statistical Association 1958; 53: 457–481.
- [36] Roher N, Callol A, Planas JV, Goetz FW and McKenzie SA. Endotoxin recognition in fish results in inflammatory cytokine secretion not gene expression. Innate Immunity 2011; 17: 16-28.
- [37] McKenzie S, Planas JV and Goetz FW. LPS-stimulated expression of a tumor necrosis factor-alpha mRNA in primary trout monocytes and in vitro differentiated macrophages. Developmental and Comparative Immunology 2003; 27: 393-400.
- [38] Swain P, Nayak SK, Nanda PK and Dash S. Biological effects of bacterial lipopolysaccharide (endotoxin) in fish: A review. Fish and Shellfish Immunology 2008; 25: 191-201.
- [39] Iliev DB, Roach JC, McKenzie S, Planas JV and Goetz FW. Endotoxin recognition: In fish or not in fish?. FEBS Letters 2005; 579: 6519-6528.
- [40] Novoa B, Bowman TV, Zon L and Figueras A. LPS response and tolerance in the zebrafish (Danio rerio). Fish and Shellfish Immunology 2009; 26: 326-331.
- [41] Dalmo RA and Bøgwald J. β-glucans as conductors of immune symphonies. Fish Shellfish Immunology 2008; 25: 384-396.
- [42] Matsuo K and Miyazono I. The influence of long-term administration of peptidoglycan on disease resistance and growth of juvenile rainbow trout. Nippon Suisan Gakkaishi 1993; 59: 1377-1379.
- [43] Yoshida T, Kruger R and Inglis V. Augmentation of non-specific protection in African catfish, Clarias gariepinus (Burchell), by the long-term oral administration of immunostimulants. Journal of Fish Diseases 1995; 18: 195-198.
- [44] Sakai M. Current research status of fish immunostimulants. Auqaculture 1999; 172: 63-92.
- [45] Siwicki AK, Morand M, Terech-Majewska E, Niemczuk W, Kazun K and Glabski E. Influence of immunostimulants on the effectiveness of vaccines in fish: in vitro and in vivo study. Journal of Applied Ichthyology 1998; 14: 225-227.
- [46] Brown GD and Gordon S. Minireview: Fungal β-glucans and Mammalian Immunity. Immunity, 2003; 19: 311-315.

- [47] Akramiene D, Kodrotas A, Didziapetriene J and Kevelaitis E. Effects of beta-glucans on the immune system. Medicina Lithuania 2007; 43: 597-606.
- [48] Williams DL, Pretus HA, McMamee RB, Jones EL, Ensley HE, Browder IW and DiLuzio NR. Development, physicochemical characterization and preclinical efficacy evaluation of a water soluble glucan sulfate derived from Saccharomyces cerevisiae. Immunopharmacology 1991; 22: 139-155.
- [49] Williams DL. Overview of (1 -> 3)-beta-D-glucan immunobiology. Mediators of Inflammation 1997; 6: 247-250.
- [50] Williams DL, Sherwood ER, McNamee RB, Jones EL and DiLuzio NRD. Therapeutic efficiency of glucan in a murine model of hepatic metastatic disease. Hepatology 1985; 5: 198-206.
- [51] Maeda YY, Watanabe ST, Chihara G and Rokutanda M. T-cell mediated vascular dilatation and hemmorrage induced by antitumor polysaccharides. International Journal of Immunopharmacology 1984; 6: 493-501.
- [52] Maeda YY, Watanabe ST, Chihara Ch and Rokutanda M. Denaturation and renaturation of a β-1,6; 1,3-glucan, lentinan, associated with expression of T-cell-mediated responses. Cancer Research 1988; 48: 671-675.
- [53] Seljelid R. Tumor regression after treatment with animated beta-1-3D polyglucose is initiated by circulatory failure. Scandibavian Journal of Immunology 1989; 29: 181-192.
- [54] Williams DL, Sherwood ER, William BI, McNamee, RB, Jones EL and DiLuzio NR. Pre-clinical safety evaluation of soluble glucan. International Journal of Immunopharmacology 1988; 10: 405-414.
- [55] Lo C-M, Xu M, Yang Q, Zheng S, Carey K, Tubb MR, Davidson WS, Liu M, Woods SC and Tso P. Effect of intraperitoneal and intravenous administration of cholecystokinin-8 and apolipoprotein AIV on intestinal lymphatic CCK-8 and apo AIV concentration. American Journal of Physiology – Regulatory Integrative and Comparative Physiology 2009; 296: R43-R50.
- [56] Li GC, Li L, Liu RY, Rehman M and Lee WM. Heat shock protein hsp70 protects cells from thermal stress even after deletion of its ATP-binding domain. Proceedings of The National Academy of Science of The United States of America 1992; 89: 2036-2040.
- [57] Trautinger F. Heat shock proteins in the photobiology of human skin. Journal of Photochemistry and Photobiology B: Biology 2001; 63: 70-77.
- [58] Guo S, Wharton W, Moseley P and Shi H. Heat shock protein 70 regulates cellular redox status by modulating glutathione-related enzyme activities. Cell Stress and Chaperones 2007; 12: 245-254.
- [59] Guzhova I and Margulis B. Hsp70 chaperone as a survival factor in cell pathology. International Review of Cytology – A Survey of Cell Biology 2006; 254: 101-149.
- [60] Seong SY and Matzinger P. Hydrophobicity: an ancient damage-associated molecular pattern initiates innate immune response. Nature reviews. Immunology 2004; 4: 469-478.
- [61] Fange R. Lymphoid organs in sturgeons acipenseridae. Veterinaty Immunology and Immunopathology 1986; 12: 153-162.
- [62] Ferguson HW, Claxton MJ, Moccia RD and Wilkie EJ. The quantitative clearance of bacteria from the bloodstream of rainbow trout (Salmo gairdneri). Veterinary Pathology 1982; 19: 687-699.
- [63] Chadzinska M, Leon-Kloosterziel KM, Plytycz B and Verburg-van Kemenade LBM. In vivo kinetics of cytokine expression during peritonitis in carp: Evidence for innate and alternative macrophage polarization. Developmental and Comparative Immunology 2008; 32: 509-518.
- [64] Saeij JPJ, Verburg-van Kemenade LBM, van Muiswinkel WB and Wiegertjes GF. Daily handling stress reduces resistance of carp to Trypanoplasma borreli: in vitro modulatory effects of cortisol on leukocyte function and apoptosis. Developmental and Comparative Immunology 2003; 27: 233-245.
- [65] Baybutt HN and Holsboer F. Inhibition of Macrophage differentiation and function by cortisol. Endocrinology 1990; 127: 476-480.
- [66] Ganassin RC and Bols NC. Development of a monocyte/macrophage-like cell line, RTS11, from rainbow trout spleen. Fish and Shellfish Immunology 1998; 8: 457-476.
- [67] Pagniello KB, Bols NC and Lee LEJ. Effect of corticosteroids on viability and proliferation of the rainbow trout monocyte/macrophage cell line, RTS11. Fish aand Shellfish Immunology 2002; 13: 199-214.
- [68] Pai R, Karunasagar I, Shetty HPC and Karunasagar I. The effect of some stress factors on infection of fish by Aeromonas hydrophila. Journal od Aquaculture in the topics 1995; 10: 29-35.
- [69] Fang HM, Ling KC, Ge R and Sin YM. Enhancement of protective immunity in blue gourami, Trichogaster trichopterus (Pallas), against Aeromonas hydrophila and Vibrio anguillarum by A. hydrophila major adhesin. Journal of Fish Diseases 2000; 23: 137-145.
- [70] van der Marel M, Caspari N, Neuhaus H, Meyer W, Enss ML and Steinhagen D. Changes in skin mucus of common carp, Cyprinus carpio L., after exposure to water with a high bacterial load. Journal of Fish Diseases 2010; 33: 431-439.
- [71] Selvaraj V, Sampath K and Seker V. Administration of yeast glucan enhances survival and some non-specific and specific immune parameters in carp (Cyprinus carpio) infected with Aeromonas hydrophila. Fish and Shellfish Immunology 2005; 19: 293-306.
- [72] Bromage ES, Kaattari IM, Zwollo P, Kaattari SL. Plasmablast and plasma cell production and distribution in trout immune tissues. Journal of Immunology 2004; 173: 7317-7323.
- [73] Zwollo P, Cole S, Bromage E and Kaattari S. B cell heterogeneity in the teleost kidney: Evidence for a maturation gradient from anterior to posterior kidney. Journal of Immunology 2005; 174: 6608-6616.

- [74] Hansen JD, Landis ED and Philips RB. Discovery of a unique lg heavy-chain isotype (lgT) in rainbow trout: Implications for a distinctive B cell developmental pathway in teleost fish. Proceedings – National Academy of Science USA 2005; 102: 6919-6924.
- [75] Miyadai T, Ootani M, Tahara D, Aoki M and Saitoh K. Monoclonal antibodies recognising serum immunoglobulins and surface immunoglobulin-positive cells of puffer fish, torafugu (Takifugu rubripes). Fish and Shellfish Immunology 2004; 17: 211-222.
- [76] Danilova N, Bussmann J, Jekosch K and Steiner LA. The immunoglobulin heavy-chain locus in zebrafish: identification and expression of a previously unknown isotype, immunoglobulin Z. Nature Immunology 2005; 6: 295-302
- [77] Ryo S, Wijdeven RHM, Tyagi A, Hermsen T, Kono T, Karunasagar I, Rombout JHWM, Sakai M, Verburg-van Kemenade BML and Savan R. Common carp have two subclasses of bonyfish specific antibody IgZ showing differential expression in response to infection. Developmental and Comparative Immunology2010; 34: 1183-1190.
- [78] Li J, Barreda DR, Zhang YA, Boshra H, Gelman AE, Lapatra S, Tort L and Sunyer JO. B lymphocytes from early vertebrates have potent phagocytic and microbicidal abilities. Nature Immunology 2006; 7: 1116–1124.
- [79] Lobb CJ and Clem LW. Phylogeny of immunoglobulin structure and function. 11 Secretory immunoglobulins in the cutaneous mucus of the sheepshead, Archosargus probatocephalus. Developmental and Comparative immunology, 1981b; Volume 5, 587-596.
- [80] Zhang Y-A, Salinas I, Li J, Parra D, Bjork S, Xu Z, LaPatra SE, Bartholomew J and Sunyer JO. IgT, a primitive immunoglobulin class specialized in mucosal immunity. Nature 2010; 11: 827-836.
- [81] van der Marel M, Adamek M, Gonzalez SF, Frost P, Rombout JHWM, Wiegertjes GF, Savelkoul HFJ and Steinhagen D. Molecular cloning and expression of two b-defensin and two mucin genes in common carp (Cyprinus carpio L.) and their up-regulation after b-glucan feeding. Fish and Shellfish Immunology 2012; 32:494-501.

Figure legends

Figure 1. Kaplan-Meier curve of survival percentage among all groups during experiment period. B - bacterin injected group; B-L β g - bacterin + low dose of β -glucan injected group; B-H β g - bacterin + high dose of β -glucan injected group.

Figure 2. Plasma cortisol concentrations for B-injected, B-L β g-injected and B-H β g-injected group. The results are shown as mean (n=7) ± standard deviation (SD). Significantly elevated level of plasma cortisol was observed in all experimental groups. The significant difference (Δ) in cortisol level between the B-injected and B-H β g groups was observed one day post injection. At the same time point, significant difference (***) between B-L β g and B-H β g group was noted. B - bacterin injected group; B-L β g - bacterin + low dose of β -glucan injected group; B-H β g - bacterin + high dose of β -glucan injected group. *** p < 0.001; Δ depicts significant difference between B and B-H β g injected groups.

Figure 3. Antibodies against *Aeromonas hydrophila*. The results are shown as mean absorbance at 450 nm + standard deviation (SD). Results have shown presence of antibodies against *Aeromonas hydrophila* in all experimental groups, although no significant differences in antibody titre between treatments were observed.

Figure 4. Quantitative real-time PCR in group injected with bacterin + low dose of β -glucan (B-L β g). Expression for genes interleukin 1 β (IL-1 β), Heat shock protein 70 (Hsp70), immunoglobulin M (IgM) and immunoglobulin Z1 (IgZ1) in skin (A-D) and head kidney (E-H) of common carp. Results are obtained by RQ-PCR and expressed relative to B-injected group at each time point. The data are normalised relative to 40S and analysed using the $-2^{-\Delta\Delta Ct}$ method. Bars represent fold expression (n = 7) + SD. * depicts significant differences in comparison to B-injected group (*p < 0.05; **p < 0.01; ***p < 0.001).

Figure 5. Quantitative real-time PCR in group injected with bacterin + high dose of β -glucan (B-H β g). Expression for genes interleukin 1 β (IL-1 β), Heat shock protein 70 (Hsp70), immunoglobulin M (IgM) and immunoglobulin Z1 (IgZ1) in skin (A-D) and head kidney (E-H) of common carp. Results are obtained by RQ-PCR and expressed relative to B-injected group at each time point. The data are normalised relative to 40S and analysed using the $-2^{-\Delta\Delta Ct}$ method. Bars represent fold expression (n = 7) + SD. * depicts significant differences in comparison to B-injected group (*p < 0.05; **p < 0.01; ***p < 0.001).



Time post injection (days)

Figure 1.



Days post injection

Figure 2.



Figure 3.



Figure 4.





Gene	Primer	Sequence (5' – 3')				
405	Forward	GTTGAAGGAAGTGGCAAGGA				
403	Reverse	AGAATACGGCCTCTGATGGA				
H 10	Forward	AAGGAGGCCAGTGGCTCTGT				
пс-тр	Reverse	CCTGAAGAAGAGGAGGCTGTCA				
H 70	Forward	GGCAGAAGGTGACAAATGCA				
Hsp70	Reverse	TGGGCTCGTTGATGTTCTCA				
	Forward	CACAAGGCGGGAAATGAAGA				
IgIVI	Reverse	GGAGGCACTATATCAACAGCA				
	Forward	ACTCCCTGGTGTGTGACCTC				
IgZ1	Reverse	TACAAACAGCATGAGCCAGCTA				

Table 1. Sequences of primers used for real-time PCR.

 Table 2. Plasma cortisol (nmol/L)

Days post injection										
	-2	1	4	7	12	23				
В	342,12	463,51	386,26	813,91	231,76	344,86				
	±77,25	±46,9	±63,46	±19,31	±99,30	±35,87				
B-Lβg	339,36	722,86	551,8	800,11	333,84	413,85				
	±104,84	±16,55	±63,46	±22,1	±80	±38,63				
B-Hβg	215,2	1158,78	510,42	744,93	257,9	342,12				
	±113,12	±13,8	±35,8	±13,8	±44,14	±8,28				

