



Hydrogen Bonding in Ion-pair Molecules in Vapors over ionic liquids, studied by Raman Spectroscopy and *ab initio* Calculations

Berg, Rolf W.

Publication date:
2010

[Link back to DTU Orbit](#)

Citation (APA):

Berg, R. W. (2010). *Hydrogen Bonding in Ion-pair Molecules in Vapors over ionic liquids, studied by Raman Spectroscopy and ab initio Calculations*. Paper presented at Conference on Molten Salts and Ionic Liquids 2010, Bamberg, Germany.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Hydrogen Bonding in Ion-pair Molecules in Vapors over ionic liquids, studied by Raman Spectroscopy and *ab initio* Calculations

Rolf W. Berg, *Department of Chemistry, Technical University of Denmark, Kemitorvet, Building 207, DK-2800 Lyngby, Denmark*

The hydrogen bonding interactions in selected archetypal vapor molecules formed in the gas phase over protic ionic liquids are discussed, based on Raman spectroscopy assisted with *ab initio* molecular orbital DFT-type quantum mechanical calculations (B3LYP with 6-311+G(d,p) basis sets) on assumed gaseous free state models. Some extreme examples are to be discussed: (1) The 1,1,3,3-tetramethylguanidinium chloride “molecule” [TMG-H-Cl] found¹ to exist in gaseous state over its corresponding liquid in evacuated ampouls heated at ~225°C (Fig. 1); (2) the analogous bromide “molecule” [TMG-H-Br]² and (3) the 1-methylimidazolium ethanoate, [mim-H-O₂CCH₃] found³ to have a less likely existence in the vapor of the corresponding liquid in ampouls at ~200°C (Fig. 2). Experimental Raman results will be compared to *ab initio* calculated spectra.





Fig. 1. Likely gaseous ion pair molecule of 1,1,3,3-tetramethylguanidinium chloride. The optimized N-H distance in the N-H \cdots Cl bond was 1.099 Å. The H-Cl distance was 1.832 Å to compare with the ~1.27 Å in HCl gas. The bromide behaved similarly².

Fig. 2. Not so likely 1-methylimidazolium ethanoate gas molecule. The optimized N-H and H-O distances in the N-H \cdots O bond were found as 1.730 and 1.005 Å. The comparable H-O distance in solid ethanoic acid is ~1.011 Å (neutron diffraction).



- 1 R. W. Berg, A. Riisager & R. Fehrmann, Formation of an ion pair molecule with a single NH⁺ \cdots Cl⁻ hydrogen bond: Raman spectra of 1,1,3,3-Tetramethylguanidinium chloride, *J. Phys. Chem. A*, 2008, 112, 8585-8592.
- 2 R. W. Berg, O. Nguyen van Buu, A. Riisager. & R. Fehrmann, In preparation.
- 3 R. W. Berg, J.N.C. Lopes, L.P.N. Rebelo, K.R. Seddon & A.A. Tomaszowska, A Raman Spectroscopic Study of the Vapour Phase of 1-Methylimidazolium Ethanoate, a Protic Ionic Liquid, To be submitted for *J. Phys. Chem. A* (2009).