



Radioactivity in the Risø District January-June 2010

Nielsen, Sven Poul; Andersson, Kasper Grann; Miller, Arne

Publication date:
2010

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Nielsen, S. P., Andersson, K. G., & Miller, A. (2010). *Radioactivity in the Risø District January-June 2010*. Danmarks Tekniske Universitet, Risø Nationallaboratoriet for Bæredygtig Energi. Denmark. Forskningscenter Risoe. Risoe-R No. 1756(EN)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Radioactivity in the Risø District January-June 2010

Risø-R-Report

Sven P. Nielsen, Kasper G. Andersson and Arne Miller
Risø-R-1756(EN)
December 2010

Risø DTU
National Laboratory for Sustainable Energy



Author: Sven P. Nielsen, Kasper G. Andersson and Arne Miller
Title: Radioactivity in the Risø District January-June 2010
Division: Radiation Research

Risø-R-1756(EN)
December 2010

Abstract (max. 2000 char.): The environmental surveillance of the Risø environment was continued in January - June 2010. The mean concentrations in air were: $0.99 \pm 1.11 \mu\text{Bq m}^{-3}$ of ^{137}Cs , $4.56 \pm 2.71 \text{ mBq m}^{-3}$ of ^7Be and $0.49 \pm 0.57 \text{ mBq m}^{-3}$ of ^{210}Pb (± 1 S.D.; $N = 26$). The depositions by precipitation at Risø in the first half of 2010 were: 0.095 Bq m^{-2} of ^{137}Cs , 521 Bq m^{-2} of ^7Be , 48.2 Bq m^{-2} of ^{210}Pb and $< 0.5 \text{ kBq m}^{-2}$ of ^3H . The average background dose rate (TLD) at Risø (Zone I) was 106 nSv h^{-1} compared with $91 \pm 4 \text{ nSv h}^{-1}$ (± 1 S.D.; $N = 4$) in the four zones around Risø.

ISSN 0106-2840
ISBN 978-87-550-3863-9

Contract no.:

Group's own reg. no.:
1400103-07

Sponsorship:

Cover :

Pages: 24
Tables: 14
References:

Information Service Department
Risø National Laboratory for
Sustainable Energy
Technical University of Denmark
P.O.Box 49
DK-4000 Roskilde
Denmark
Telephone +45 46774005
bibl@risoe.dtu.dk
Fax +45 46774013
www.risoe.dtu.dk

Contents

| | | |
|------------|---|----|
| Table 1. | Radionuclides in air | 5 |
| Table 2.1. | Radionuclides in precipitation | 6 |
| Table 2.2. | Radionuclides in precipitation | 6 |
| Table 2.3. | Tritium in precipitation | 7 |
| Table 2.4. | Tritium in precipitation | 7 |
| Table 3.1. | Radionuclides in sediment samples | 8 |
| Table 4.1. | Radionuclides in seawater | 8 |
| Table 4.2. | Tritium in seawater | 8 |
| Table 5.1. | Radionuclides in grass | 9 |
| Table 5.2. | Radionuclides in sea plants | 10 |
| Table 7.1. | Waste water | 11 |
| Table 8.1. | Background dose rates around the border of Risø (TLD) | 12 |
| Table 8.2. | Background dose rates around Risø (TLD) | 13 |
| Table 8.3. | Terrestrial dose rates at the Risø zones (NaI(Tl) detector) | 14 |
| Fig. 1. | Map of Risø | 15 |
| Fig. 1.1. | Caesium-137 in air | 16 |
| Fig. 1.2. | Beryllium-7 and lead-210 in air | 16 |
| Fig. 2.3.1 | Tritium in precipitation (1 m ² rain collector) | 17 |
| Fig. 2.3.2 | Tritium in precipitation (10 m ² rain collector) | 17 |
| Fig. 3.1 | Caesium-137 in sediment samples | 18 |
| Fig. 4.1 | Caesium-137 in seawater | 19 |
| Fig. 4.2 | Tritium in seawater | 19 |
| Fig. 7.1 | Total-beta radioactivity in waste water | 20 |
| Fig. 8.1. | Map of Risø with locations for TLD measurements | 21 |
| Fig. 8.2. | The environment of Risø | 22 |

Table 1. Radionuclides in ground level air collected at Risø (cf. Figs. 1, 1.1 and 1.2), January - June 2010. (Unit: $\mu\text{Bq m}^{-3}$)

| Date | ^7Be | ^{137}Cs | ^{210}Pb |
|------------------------|---------------|-------------------|-------------------|
| 28-Dec-09 – 04-Jan-10 | 2923 | 0.472 | 239 |
| 04-Jan-10 – 11-Jan-10 | 4289 | 1.193 | 759 |
| 11-Jan-10* – 18-Jan-10 | 9364 | 1.838 | 872 |
| 18-Jan-10* – 26-Jan-10 | 14080 | 4.618 | 1550 |
| 26-Jan-10* – 01-Feb-10 | 6610 | 2.287 | 684 |
| 01-Feb-10* – 08-Feb-10 | 7738 | 3.669 | 2293 |
| 08-Feb-10* – 15-Feb-10 | 5422 | 2.191 | 1525 |
| 15-Feb-10* – 22-Feb-10 | 3387 | 1.238 | 1262 |
| 22-Feb-10 – 01-Mar-10 | 3011 | 0.796 | 383 |
| 01-Mar-10 – 08-Mar-10 | 4779 | 0.864 | 330 |
| 08-Mar-10 – 15-Mar-10 | 4413 | 0.752 | 164 |
| 15-Mar-10 – 22-Mar-10 | 8337 | 1.241 | 468 |
| 22-Mar-10 – 29-Mar-10 | 3657 | 0.416 | 205 |
| 29-Mar-10 – 06-Apr-10 | 2344 | 0.263 | 103 |
| 06-Apr-10 – 13-Apr-10 | 2401 | 0.434 | 202 |
| 13-Apr-10 – 19-Apr-10 | 2615 | 0.565 | 171 |
| 19-Apr-10 – 26-Apr-10 | 3703 | 0.353 | 155 |
| 26-Apr-10 – 03-May-10 | 3707 | 0.431 | 188 |
| 03-May-10 – 10-May-10 | 3858 | 0.292 | 169 |
| 10-May-10 – 17-May-10 | 3820 | 0.295 | 159 |
| 17-May-10 – 25-May-10 | 2470 | 0.361 | 248 |
| 25-May-10 – 31-May-10 | 2105 | 0.240 | 87 |
| 31-May-10 – 07-Jun-10 | 3505 | 0.299 | 157 |
| 07-Jun-10 – 14-Jun-10 | 4547 | 0.150 | 185 |
| 14-Jun-10 – 22-Jun-10 | 2626 | 0.177 | 98 |
| 22-Jun-10 – 28-Jun-10 | 2934 | 0.316 | 157 |
| Mean | 4563 | 0.990 | 493 |
| SD | 2714 | 1.114 | 567 |

* Uncertain air volume determination due to frost

Table 2.1. Radionuclides in precipitation in the 10 m² rain collector at Risø (cf. Fig. 1), January - June 2010. (Unit: Bq m⁻³)

| Month | ⁷ Be | ¹³⁷ Cs | ²¹⁰ Pb |
|----------|-----------------|-------------------|-------------------|
| January | 2092 | 0.313 | 308 |
| February | 2509 | 0.285 | 221 |
| March | 1830 | 0.124 | 137 |
| April | 1805 | 0.331 | 69 |
| May | 821 | 0.126 | 38 |
| June | 5359 | 1.339 | 561 |

Table 2.2. Radionuclides in precipitation in the 10 m² rain collector at Risø (cf. Fig. 1), January - June 2010. (Unit: Bq m⁻²)

| Month | Precipitation (m) | ⁷ Be | ¹³⁷ Cs | ²¹⁰ Pb |
|----------|-------------------|-----------------|-------------------|-------------------|
| January | 0.016 | 34.5 | 0.0052 | 5.1 |
| February | 0.023 | 56.5 | 0.0064 | 5.0 |
| March | 0.054 | 99.0 | 0.0067 | 7.4 |
| April | 0.012 | 20.9 | 0.0038 | 0.8 |
| May | 0.052 | 43.1 | 0.0066 | 2.0 |
| June | 0.049 | 267.0 | 0.0667 | 27.9 |
| Sum | 0.206 | 521.0 | 0.0954 | 48.2 |

Table 2.3. Tritium in precipitation collected at Risø (cf. Figs. 1, 2.3.1 and 2.3.2). January - June 2010. (Unit: kBq m⁻³)

| Month | 1 m ² rain collector* | 10 m ² rain collector* |
|-------------------------|----------------------------------|-----------------------------------|
| January | < 2.0 | < 2.0 |
| February | < 2.0 | < 2.0 |
| March | < 2.0 | < 2.0 |
| April | < 2.0 | < 2.0 |
| May | < 2.0 | 2.5 |
| June | < 2.0 | 2.7 |
| Double determinations*. | | |

Table 2.4. Tritium in precipitation collected at Risø (cf. Fig. 1). January - June 2010. (Unit: kBq m⁻²)

| Month | Precipitation (m) | 1 m ² rain collector | 10 m ² rain collector |
|----------|-------------------|---------------------------------|----------------------------------|
| January | 0.016 | < 0.032 | < 0.032 |
| February | 0.023 | < 0.046 | < 0.046 |
| March | 0.054 | < 0.108 | < 0.108 |
| April | 0.012 | < 0.024 | < 0.024 |
| May | 0.052 | < 0.104 | 0.130 |
| June | 0.049 | < 0.098 | 0.132 |
| Sum | 0.206 | < 0.412 | < 0.472 |

Table 3.1. Radionuclides in sediment samples collected at Bolund in Roskilde Fjord.(cf. Fig. 3.1) January - June 2010. (Unit: Bq kg⁻¹ dry)

No samples in this period

Table 4.1. Radionuclides in seawater collected in Roskilde Fjord (cf. Fig. 4.1) January - June 2010. (Unit: Bq m⁻³)

No samples in this period

Table 4.2. Tritium in seawater collected in Roskilde Fjord (Risø pier) (cf. Fig. 4.2) January - June 2010.

| Month | kBq m ⁻³ |
|----------|---------------------|
| January | < 2.0 * |
| February | < 2.0 * |
| March | < 2.0 * |
| April | 2.3 * |
| May | < 2.0 * |
| June | < 2.0 * |

* Double determinations

Table 5.1. Radionuclides in grass (* snow) collected at Risø (near the Waste Treatment Station (cf. Fig. 1)), January - June 2010. (**Measured on bulked ash samples)

| Week no. or month | Date | K (g kg ⁻¹ fresh) | ¹³⁷ Cs (Bq kg ⁻¹ fresh) | ¹³⁷ Cs (Bq m ⁻²) |
|----------------------|--------------|---------------------------------|--|--|
| 1 | 4 January* | <0.9 | <0.3 | |
| 2 | 11 January* | - | <0.3 | |
| 3 | 18 January* | <0.2 | <0.4 | |
| 4 | 25 January* | <0.2 | <0.3 | |
| 5 | 1 February* | <1.6 | <0.4 | |
| 6 | 8 February* | <0.2 | <0.3 | |
| 7 | 15 February* | <0.1 | <0.3 | |
| 8 | 22 February* | <0.2 | <0.4 | |
| 9 | 1 March* | <0.1 | <0.2 | |
| 10 | 8 March* | <0.2 | <0.3 | |
| 11 | 15 March | 4.2 | <0.5 | |
| 12 | 22 March | 4.1 | <1.4 | |
| 13 | 29 March | 4.1 | <1.6 | |
| 14 | 6 April | 5.2 | <1.1 | |
| 15 | 12 April | 4.6 | <0.7 | |
| 16 | 19 April | 6.8 | <1.2 | |
| 17 | 26 April | 7.3 | <0.6 | |
| 18 | 3 May | 4.8 | <0.5 | |
| 19 | 10 May | 5.0 | <0.7 | |
| 20 | 17 May | 3.5 | <0.5 | |
| 21 | 25 May | 4.5 | <0.4 | |
| 22 | 31 May | 4.2 | <0.4 | |
| 23 | 7 June | 4.2 | <0.5 | |
| 24 | 14 June | 4.4 | <0.5 | |
| 25 | 21 June | 4.8 | <0.5 | |
| 26 | 28 June | 4.7 | <0.4 | |
| **January | | 1.3 | - | - |
| **February | | - | - | - |
| **March | | 4.2 | 0.197 | 0.021 |
| **April | | 5.9 | 0.238 | 0.037 |
| **May | | 4.9 | 0.130 | 0.050 |
| **June | | 4.8 | 0.040 | 0.037 |

Table 5.2. Radionuclides in Fucus vesiculosus collected at Bolund in Roskilde Fjord. January - June 2010. (Unit: Bq kg⁻¹ dry)

No samples in this period

Table 7.1. Waste water collected at Risø (cf. Fig. 1), January - June 2010.

| Week number | eqv. mg KCl l ⁻¹ | ¹³⁷ Cs (Bq m ⁻³) | ¹³¹ I (Bq m ⁻³) | ²²⁶ Ra (Bq m ⁻³) |
|-------------|--------------------------------|--|---|--|
| 1 | 32 | <128 | <136 | <243 |
| 2 | 58 | <113 | <118 | <230 |
| 3 | 63 | <107 | <115 | <218 |
| 4 | 77 | <112 | <117 | <221 |
| 5 | 79 | <113 | <124 | <229 |
| 6 | 90 | <110 | <117 | <216 |
| 7 | 96 | <123 | <119 | <224 |
| 8 | 89 | <113 | <120 | <229 |
| 9 | 74 | <115 | <120 | 484 |
| 10 | 78 | <132 | <138 | <249 |
| 11 | 126 | 185.7 | <127 | <223 |
| 12 | 75 | <123 | <120 | <222 |
| 13 | 103 | <112 | <117 | <219 |
| 14 | 51 | <124 | <120 | <242 |
| 15 | 57 | <112 | <117 | <221 |
| 16 | 66 | <113 | <118 | <220 |
| 17 | 95 | <125 | <112 | <215 |
| 18 | 49 | <126 | <123 | <229 |
| 19 | 93 | <114 | <120 | <220 |
| 20 | 75 | <135 | <141 | <261 |
| 21 | 60 | <116 | <121 | <226 |
| 22 | 70 | <114 | <117 | <224 |
| 23 | 81 | <128 | <129 | <237 |
| 24 | 49 | <109 | <112 | <213 |
| 25 | 72 | <112 | <119 | <217 |
| 26 | 129 | <113 | <120 | <226 |
| Mean | 76.4 | | | |
| SD | 22.6 | | | |

Table 8.1. Background dose rates around the border of Risø (cf. Fig. 8.1) measured with thermoluminescence dosimeters (TLD) in the period November 2009 – April 2010. (Results are normalized to nSv h⁻¹)

| Location | nSv h ⁻¹ |
|----------|---------------------|
| 1 | 95 |
| 2 | 87 |
| 3 | 78 |
| 4 | 86 |
| 5 | 94 |
| 6 | 105 |
| Mean | 91 |

Table 8.2. Background dose rates around Risø (cf. Fig. 8.2 and Fig. 1) measured with thermoluminescence dosimeters (TLD) in the period November 2009 – April 2010. (Results are normalized to nSv h^{-1})

| Risø zone | Location | nSv h^{-1} |
|-----------|----------|---------------------|
| I | 1 | 76 |
| I | 2 | 92 |
| I | 3 | 169 |
| I | 4 | 96 |
| I | 5 | 98 |
| Mean | | 106 |
| II | P1 | 95 |
| II | P2 | 89 |
| II | P3 | 76 |
| II | P4 | 104 |
| Mean | | 91 |
| III | P1 | 83 |
| III | P2 | 92 |
| III | P3 | 86 |
| Mean | | 87 |
| IV | P1 | 76 |
| IV | P2 | 83 |
| IV | P3 | 92 |
| IV | P4 | 94 |
| IV | P5 | 96 |
| IV | P6 | 85 |
| IV | P7 | 114 |
| Mean | | 91 |
| V | P1 | 101 |
| V | P2 | 94 |
| V | P3 | 115 |
| V | P4 | 80 |
| V | P5 | 103 |
| V | P6 | 81 |
| V | P7 | 86 |
| V | P8 | 110 |
| V | P9 | 96 |
| V | P10 | 104 |
| Mean | | 97 |

Table 8.3. Terrestrial dose rates at the Risø zones (cf. Fig. 8.2 and Fig. 1) January - June 2010. Measured with a NaI(Tl) detector. (Unit: nSv h⁻¹)

| Risø zone | Location | January | October |
|-----------|----------|---------|---------|
| I | P1 | 33 | 34 |
| I | P2 | 43 | 43 |
| I | P3 | 420 | 377 |
| I | P4 | 39 | 40 |
| I | P5 | 40 | 40 |
| Mean | | 115 | 107 |
| II | P1 | 43 | 41 |
| II | P2 | 38 | 37 |
| II | P3 | 36 | 39 |
| II | P4 | 39 | 37 |
| Mean | | 39 | 39 |
| III | P1 | | 41 |
| III | P2 | | 40 |
| III | P3 | | 38 |
| Mean | | | 40 |
| IV | P1 | | 35 |
| IV | P2 | | 42 |
| IV | P3 | | 38 |
| IV | P4 | | 40 |
| IV | P5 | | 36 |
| IV | P6 | | 40 |
| IV | P7 | | 42 |
| Mean | | | 39 |
| V | P1 | | 35 |
| V | P2 | | 44 |
| V | P3 | | 48 |
| V | P4 | | 42 |
| V | P5 | | 45 |
| V | P6 | | 40 |
| V | P7 | | 34 |
| V | P7a | | 37 |
| V | P8 | | 38 |
| V | P9 | | 47 |
| V | P10 | | 34 |
| Mean | | | 40 |




Fig. 1. Locations for measurements of gamma-background radiation Zone I and II (cf. Tables 8.2 and 8.3)




Fig. 1.1. Caesium-137 in ground level air collected at Risø in January-June 2010. (Unit: $\mu\text{Bq m}^{-3}$)




Fig. 1.2. Beryllium-7 and lead-210 in ground level air collected at Risø in January- June 2010. (Unit: $\mu\text{Bq m}^{-3}$)




Fig. 2.3.1. Tritium in precipitation collected at Risø (1 m^2 rain collector) 1980 - 2010. (Unit: kBq m^{-3} ; DL = detection limit)




Fig. 2.3.2. Tritium in precipitation collected at Risø (10 m^2 rain collector) 1980 - 2010. (Unit: kBq m^{-3} ; DL = detection limit)




Fig. 3.1. Caesium-137 in sediment samples collected at Bolund in Roskilde Fjord. 1980 – 2009. (Unit: Bq kg⁻¹ dry matter)




Fig. 4.1. Caesium-137 in seawater collected in Roskilde Fjord 1980 - 2009. (Unit: Bq m^{-3})





Fig. 4.2. Tritium in seawater collected in Roskilde Fjord 1980 - 2010. (Unit: kBq m^{-3} ; DL = detection limit)



*Fig. 7.1. Total-beta radioactivity in waste water collected at Risø 1994 - 2010.
(Unit: eqv. mg KCl l⁻¹)*




Fig. 8.1. Locations (1-6) for TLD measurements around the border of Risø (cf. Table 8.1).



Fig. 8.2. Locations for measurements of background radiation around Risø in Zones III, IV and V.

Risø DTU is the National Laboratory for Sustainable Energy. Our research focuses on development of energy technologies and systems with minimal effect on climate, and contributes to innovation, education and policy. Risø has large experimental facilities and interdisciplinary research environments, and includes the national centre for nuclear technologies.

Risø DTU
National Laboratory for Sustainable Energy
Technical University of Denmark

Frederiksborgvej 399
PO Box 49
DK-4000 Roskilde
Denmark
Phone +45 4677 4677
Fax +45 4677 5688

www.risoe.dtu.dk