



## Environmental radioactivity in the Faroes in 1981

Aarkrog, A.; Dahlgaard, Henning; Hallstadius, L.; Holm, E.; Lippert, Jørgen Emil

*Publication date:*  
1982

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Aarkrog, A., Dahlgaard, H., Hallstadius, L., Holm, E., & Lippert, J. E. (1982). *Environmental radioactivity in the Faroes in 1981*. Denmark. Forskningscenter Risoe. Risoe-R No. 470

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

RISO

Rise-R-470

# Environmental Radioactivity in the Faroes in 1981

A. Aarås og, H. Dahlgaard, L. Hallstadius,  
E. Holm, and J. Lippert

Rise National Laboratory, DK-4000 Roskilde, Denmark

July 1982

ENVIRONMENTAL RADIOACTIVITY IN THE FAROES IN 1981

A. Aarkrog, H. Danilgaard, L. Hallstadius\*, E. Holm\*,  
and J. Lippert

\*University of Lund, Sweden

Abstract. Measurements of fallout radioactivity in the Faroes in 1981 are presented. Strontium-90 (and  $^{137}\text{Cs}$  in most cases) was determined in regularly collected samples of precipitation, grass, milk, fish, sea water, bread and drinking water. In addition, analyses were made of spot samples of lamb, sea birds, potatoes, sea plants, vegetables, eggs, and human bone. Estimates are given of the mean contents of  $^{90}\text{Sr}$  and  $^{137}\text{Cs}$  in the human diet in the Faroes in 1981. In Appendix the results from samplings of sea water and biota along the Icelandic and the Norwegian coasts are given.

INIS Descriptors

- [0] DIET, ENVIRONMENT, EXPERIMENTAL DATA, FAROE ISLANDS, FISHES, FOOD, FOOD CHAINS, GLOBAL FALLOUT, MILK, PLANTS, RADIOACTIVITY, SEA WATER, SHEEP, TABLES
- [1] ATMOSPHERIC PRECIPITATIONS, BONE TISSUES, DRINKING WATER, MAN, STRONTIUM 90
- [2] CESIUM 137
- [3] PLUTONIUM

UDC 614.73 (491.2)

July 1982

Risø National Laboratory, DK-4000 Roskilde, Denmark

**ISBN 87-550-0856-9**

**ISSN 0106-2840**

**Risø Repro 1982**

## CONTENTS

	Page
1. INTRODUCTION .....	7
2. RESULTS AND DISCUSSION .....	9
2.1. Strontium-90 in precipitation .....	9
2.2. Strontium-90 and Cesium-137 in grass .....	10
2.3. Strontium-90 and Cesium-137 in milk .....	11
2.4. Strontium-90 and Cesium-137 in terrestrial animals .....	17
2.5. Strontium-90 and Cesium-137 in sea animals .....	18
2.6. Strontium-90 in drinking water .....	19
2.7. Strontium-90 and Cesium-137 in miscellaneous samples .....	21
2.7.1. Soil .....	21
2.7.2. Sea water .....	23
2.7.3. Sea plants .....	24
2.7.4. Vegetables .....	26
2.7.5. Bread .....	28
2.7.6. Eggs .....	28
2.8. Humans .....	28
2.8.1. Strontium-90 in human bone .....	28
3. ESTIMATE OF THE MEAN CONTENTS OF $^{90}\text{Sr}$ AND $^{137}\text{Cs}$ IN THE HUMAN DIET .....	30
4. CONCLUSION .....	37
ACKNOWLEDGEMENTS .....	38
REFERENCES .....	39
APPENDICES	
APPENDIX A. A comparison between observed and predicted levels in the human food chain in the Faroes in 1981 .....	40

	Page
<b>APPENDIX B: Sea plants collected in West-Norway in 1980 ...</b>	<b>41</b>
<b>APPENDIX C: Sea water and biota collected around Iceland and between Bergen, The Faroe Islands and Iceland in June-July 1981 .....</b>	<b>45</b>

## ABBREVIATIONS AND UNITS

J: joule: the unit of energy;  $1 \text{ J} = 1 \text{ Nm} (= 0.239 \text{ cal})$   
Gy: gray: the unit of absorbed dose  $= 1 \text{ J kg}^{-1} (= 100 \text{ rad})$   
Sv: sievert: the unit of dose equivalent  $= 1 \text{ J kg}^{-1} (= 100 \text{ rem})$   
Bq: becquerel: the unit of radioactivity  $= 1 \text{ s}^{-1} (= 27 \text{ pCi})$

cal: calorie  $= 4.186 \text{ J}$

rad:  $0.01 \text{ Gy}$

rem:  $0.01 \text{ Sv}$

Ci: curie:  $3.7 \cdot 10^{10} \text{ Bq} (= 2.22 \cdot 10^{10} \text{ dpm})$

T: tera:  $10^{12}$

G: giga:  $10^9$

M: mega:  $10^6$

m: milli:  $10^{-3}$

$\mu$ : mikro:  $10^{-6}$

n: nano:  $10^{-9}$

p: pico:  $10^{-12}$

f: femto:  $10^{-15}$

a: atto:  $10^{-18}$

cap: caput: (per individual)

TNT: trinitrotoluol; 1 Mt TNT: nuclear explosives equivalent to  $10^9 \text{ kg TNT}$ .

cpm: counts per minute

dpm: disintegrations per minute

OR: observed ratio

CF: concentration factor

FP: fission products

$\mu\text{R}$ : micro-roentgen,  $10^{-6} \text{ roentgen}$

S.U.:  $\text{pCi } ^{90}\text{Sr (g Ca)}^{-1}$

O.R.: observed ratio

M.U.:  $\text{pCi } ^{137}\text{Cs (g K)}^{-1}$

**V:** vertebrae  
**m:** male  
**f:** female  
**nSr:** natural (stable) Sr

**eqv. mg KCl:** equivalents mg KCl: activity as from 1 mg KCl  
(~ 0.88 dpm)

**S.D.:** standard deviation:  $\sqrt{\frac{\sum (\bar{x} - x_i)^2}{(n-1)}}$

**S.E.:** standard error:  $\sqrt{\frac{\sum (\bar{x} - x_i)^2}{n(n-1)}}$

**U.C.L.:** upper control level

**L.C.L.:** lower control level

**S.S.D.:** sum of squares of deviation:  $\sum (\bar{x} - x_i)^2$

**f:** degrees of freedom

**s<sup>2</sup>:** variance

**v<sup>2</sup>:** ratio between the variance in question and the  
residual variance

**P:** probability fractile of the distribution in question

**η:** coefficient of variation, relative standard deviation

**ANOVA:** analysis of variance

**A:** relative standard deviation 20-33%

**B:** relative standard deviation >33%, such results are  
not considered significantly different from zero  
activity

**B.D.L.:** below detection limit

In the significance test the following symbols were used:

**\*** : probably significant (P > 95%)

**\*\*** : significant (P > 99%)

**\*\*\*** : highly significant (P > 99.9%)



## 1. INTRODUCTION

### 1.1.

The fallout programme for the Faroes, which was initiated in 1962<sup>1)</sup> in close co-operation with the National Health Service and the chief physician of the Faroes, was continued in 1981. Samples of human bone were obtained in 1981 from Dronning Alexandrine's Hospital in Thorshavn.




Fig. 1. The Faroese Islands.

**1.2.**

The present report will not repeat information concerning sample collection and analysis already given in Risø Reports Nos. 64, 86, 108, 131, 155, 181, 202, 221, 246, 266, 292, 306, 324, 346, 361, 387, 404, 422 and 448<sup>1)</sup>.

**1.3.**

The estimated mean diet of the Faroese as used in this report is still based on the estimate given by Professor E. Hoff-Jørgensen, Ph.D., in 1962.

**1.4.**

The present investigation was carried out together with corresponding examinations of fallout levels in Denmark and Greenland, described in Risø Reports Nos. 469<sup>2)</sup> and 471<sup>3)</sup>, respectively.

## 2. RESULTS AND DISCUSSION

### 2.1. Strontium-90 in precipitation

Table 2.1 shows the  $^{90}\text{Sr}$  content in precipitation collected at Højvig (near Thorshavn) and Klaksvig in 1981. The amount of fall-out at Klaksvig was a factor of 1.2 greater than that found at Højvig.

The  $^{90}\text{Sr}$  fallout in 1981 was approximately 2.1 times of the 1980 levels in the Faroes. In Denmark the 1981 levels were 3.0 times the 1980 levels<sup>2)</sup>.

Table 2.1. Strontium-90 in precipitation in the Faroes in 1981 (sampling area = 0.02 m<sup>2</sup>)

	Højvig		Klaksvig	
	Bq m <sup>-3</sup>	Bq m <sup>-2</sup>	Bq m <sup>-3</sup>	Bq m <sup>-2</sup>
Jan-Feb	15.5	4.2	14.2	6.8
March-April	26	4.3	52	2.0
May-June	39	6.4	31	9.9
July-Aug	28	3.0	28	3.5
Sept-Oct	10.7	3.3	4.3	2.4
Nov-Dec	5.9	1.32	4.4	2.1
1981	18.6	$\Sigma 23$ $\Sigma_m 1.236$	13.5	$\Sigma 27$ $\Sigma_m 2.004$
1981	0.50 pCi l <sup>-1</sup>	$\Sigma 0.62 \text{ mCi km}^{-2}$	0.36 pCi l <sup>-1</sup>	$\Sigma 0.72 \text{ mCi km}^{-2}$




Fig. 2.1. Accumulated  $^{90}\text{Sr}$  at Klaksvig and Højvig calculated from precipitation measurements since 1962. The accumulated fallout by 1962 was estimated from the Danish fallout data (cf. Rise Report No. 403<sup>2</sup>), Appendix D) and from the ratio between the  $^{90}\text{Sr}$  fallout at the Faroese stations and the fallout in Denmark in the period 1962-1974.

## 2.2. Strontium-90 and Cesium-137 in grass

Grass samples were collected near Thorshavn in 1981. Table 2.2 shows the results. The 1981  $^{137}\text{Cs}$  mean level in grass was 2.1 times the 1980 level. As compared with Danish grass in 1981<sup>2</sup>) we found the  $^{90}\text{Sr}$  level in the Faroese grass to be higher by a factor of approximately 10 in the summer months.

Table 2.2. Strontium-90 and Cesium-137 in grass from Thorshavn 1981

Month	Bq $^{90}\text{Sr}$ kg <sup>-1</sup>	Bq $^{90}\text{Sr}$ (kg Ca) <sup>-1</sup>	Bq $^{137}\text{Cs}$ kg <sup>-1</sup>	Bq $^{137}\text{Cs}$ (kg K) <sup>-1</sup>	$^{137}\text{Cs}/^{90}\text{Sr}$
June	4.5	6300	6.9	1020	1.5
August	6.9	11400	19.5	8600	2.0

### 2.3. Strontium-90 and Cesium-137 in milk

As in previous years<sup>1)</sup>, weekly samples of fresh milk were obtained from Thorshavn, Klaksvig, and Tverå. Strontium-90 and  $^{137}\text{Cs}$  were determined in bulked monthly samples.

Table 2.3.1 shows the results and Tables 2.3.2, 2.3.3 and 2.3.4 the analysis of variance of the  $\text{Bq } ^{90}\text{Sr (kg Ca)}^{-1}$ ,  $\text{Bq } ^{137}\text{Cs (kg K)}^{-1}$ , and  $\text{Bq } ^{137}\text{Cs m}^{-3}$  figures respectively. As also observed in previous years, the variation between locations was significant for  $^{137}\text{Cs}$  as well as for  $^{90}\text{Sr}$ . The highest levels were found in the milk from Tverå.

Figure 2.3.1 shows the quarterly  $\text{Bq } ^{90}\text{Sr (kg Ca)}^{-1}$  values and Fig. 2.3.2 the quarterly  $\text{Bq } ^{137}\text{Cs m}^{-3}$  levels since 1962. The annual mean values for 1981 were  $250 \text{ Bq } ^{90}\text{Sr (kg Ca)}^{-1}$  (6.6 S.U.) and  $5100 \text{ Bq } ^{137}\text{Cs m}^{-3}$  ( $138 \text{ pCi } ^{137}\text{Cs l}^{-1}$ ), i.e. the  $^{90}\text{Sr}$  levels in 1981 were 89% of the 1980 concentration, while the  $^{137}\text{Cs}$  levels were approximately 119% of the 1980 mean levels. In Danish milk the  $^{90}\text{Sr}$  concentration did not change from 1980 to 1981, but the  $^{137}\text{Cs}$  1981 level was 120% of the 1980 content.

The annual mean values of the ratio:  $\text{Bq } ^{137}\text{Cs (kg K)}^{-1} / \text{Bq } ^{90}\text{Sr (kg Ca)}^{-1}$  in Faroese milk are shown in Fig. 2.3.3. The mean ratio in 1981 for the three locations was  $12.5 \pm 0.6$  (1 S.E.) during the grazing period (May-October), and in the winter time it was  $12.8 \pm 0.6$ .

Figure 2.3.4 shows a comparison between the  $^{90}\text{Sr}$  and  $^{137}\text{Cs}$  levels in Faroese- and Danish-produced milk. It is evident that indirect contamination plays an important role for the  $^{137}\text{Cs}$  levels in the Faroes, because the ratio between  $^{137}\text{Cs}$  in Faroese and Danish milk increases when the fallout rate decreases. The ratios between the  $^{90}\text{Sr}$  levels in Faroese and Danish milk have shown a slight tendency to decrease through the years.

Table 2.3.1. Strontium-90 and Cesium-137 in milk from the Faroes in 1981

Month	Thorshavn			Klaksvig			Tvørfá			Mean		
	Bq <sup>90</sup> Sr (kg Ca) <sup>-1</sup>	Bq <sup>137</sup> Cs m <sup>-3</sup>	Bq <sup>137</sup> Cs (kg K) <sup>-1</sup>	Bq <sup>90</sup> Sr (kg Ca) <sup>-1</sup>	Bq <sup>137</sup> Cs m <sup>-3</sup>	Bq <sup>137</sup> Cs (kg K) <sup>-1</sup>	Bq <sup>90</sup> Sr (kg Ca) <sup>-1</sup>	Bq <sup>137</sup> Cs m <sup>-3</sup>	Bq <sup>137</sup> Cs (kg K) <sup>-1</sup>	Bq <sup>90</sup> Sr (kg Ca) <sup>-1</sup>	Bq <sup>137</sup> Cs m <sup>-3</sup>	Bq <sup>137</sup> Cs (kg K) <sup>-1</sup>
Jan	169	2300	1320	230	5200	2900	220	5300	3000	210	4300	2400
Feb	118	2200	1510	220	5200	3100	250	5200	3500	195	4200	2700
March	183	2300	1440	137	2200	1410	210	5900	3700	175	3500	2200
April	166	2100	1230	200	6200	3500	250	5900	3900	210	4700	2900
May	180	2400	1450	220	5700	3300	300	6000	3500	240	4700	2800
June	184	2100	1580	190	5200	3600	340	7600	4900	240	5000	3400
July	260	2500	1600	230	6200	3400	430	9100	5200	310	5900	3400
Aug	280	4300	2800	360	5500	3600	490	12000	6900	380	7300	4400
Sept	270	3900	2300	270	6600	4100	400	12800	7800	320	7800	4700
Oct	210	2400	1530	290	7000	4100	350	6700	4600	280	5400	3400
Nov	200	2500	1630	168	4700	3000	210	6100	3600	192	4400	2700
Dec	220	2100	1380	220	4600	2900	230	3900	2100	220	3500	2400
Mean	200	2600	1600	230	5400	3200	310	7200	4400	250	5100	3100
Mean (pCi)	5.5 S.U.	70 pCi <sup>137</sup> Cs l <sup>-1</sup>	45 M.U.	6.2 S.U.	145 pCi <sup>137</sup> Cs l <sup>-1</sup>	88 M.U.	8.3 S.U.	195 pCi <sup>137</sup> Cs l <sup>-1</sup>	120 M.U.	6.6 S.U.	138 pCi <sup>137</sup> Cs l <sup>-1</sup>	86 M.U.

**Table 2.3.2.** Analysis of variance of  $\ln \text{Bq } ^{90}\text{Sr (kg Ca)}^{-1}$  in Faroese milk in 1981 (from Table 2.3.1)

Variation	SSD	f	s <sup>2</sup>	v <sup>2</sup>	P
Between months	1.800	11	0.164	6.263	> 99.95%
Between locations	1.000	2	0.500	19.127	> 99.95%
Remainder	0.575	22	0.026		

**Table 2.3.3.** Analysis of variance of  $\ln \text{Bq } ^{137}\text{Cs (kg K)}^{-1}$  in Faroese milk in 1981 (from Table 2.3.1)

Variation	SSD	f	s <sup>2</sup>	v <sup>2</sup>	P
Between months	1.795	11	0.163	4.520	> 99.5%
Between locations	6.200	2	3.100	85.862	> 99.95%
Remainder	0.794	22	0.036		

**Table 2.3.4.** Analysis of variance of  $\ln \text{Bq } ^{137}\text{Cs m}^{-3}$  in Faroese milk in 1981 (from Table 2.3.1)

Variation	SSD	f	s <sup>2</sup>	v <sup>2</sup>	P
Between months	1.874	11	0.170	3.621	> 99.5%
Between locations	6.457	2	3.228	68.627	> 99.95%
Remainder	1.035	22	0.047		




Fig. 2.3.1. Strontium-90 in Faroes milk, 1962-1981.






Fig. 2.3.2. Cesium-137 in Faroese milk, 1962-1981.




Fig. 2.3.3.  $\frac{\text{M.U.}}{\text{S.U.}}$  ratios in Faroese and Danish milk, 1963-1981.




Fig. 2.3.4. A comparison between Faroese and Danish milk levels, 1962-1981.

#### 2.4. Strontium-90 and Cesium-137 in terrestrial animals

The 1981 levels in one sample of mutton were  $0.32 \text{ Bq } ^{90}\text{Sr kg}^{-1}$ ,  $97 \text{ Bq } ^{137}\text{Cs kg}^{-1}$ ,  $3300 \text{ Bq } ^{90}\text{Sr (kg Ca)}^{-1}$  and  $30\,000 \text{ Bq } ^{137}\text{Cs (kg K)}^{-1}$ . The bone level was  $3300 \text{ Bq } ^{90}\text{Sr (kg Ca)}^{-1}$ .




Fig. 2.4.1. Strontium-90 ( $\text{Bq (kg Ca)}^{-1}$ ) in lamb bone collected in the Faroes, 1962-1981.




Fig. 2.4.2. Cesium-137 (Bq (kg K)<sup>-1</sup>) in lamb meat collected in the Faroes, 1962-1981.

## 2.5. Strontium-90 and Cesium-137 in sea animals

Table 2.5.1 shows the <sup>137</sup>Cs levels in fish collected in 1981 in the Faroes. The mean levels in *Gadus aeglefinus* and *Gadus callarias* were 0.34 Bq <sup>137</sup>Cs kg<sup>-1</sup> and 0.031 Bq <sup>90</sup>Sr kg<sup>-1</sup>.

Table 2.5.1. Strontium-90 and Cesium-137 in fish flesh from the Faroes in 1981

Sampling month	Species	Sample type	Bq <sup>90</sup> Sr kg <sup>-1</sup>	Bq <sup>90</sup> Sr (kg Ca) <sup>-1</sup>	Bq <sup>137</sup> Cs kg <sup>-1</sup>	Bq <sup>137</sup> Cs (kg K) <sup>-1</sup>
Jan	<i>Gadus callarias</i>	Cod flesh	0.022	220	0.31	86
March	- " -	- " -	0.032	340	0.31	90
Sept	- " -	- " -	0.036	330	0.22	60
Dec	- " -	- " -	0.031	310	0.89	240
Jan	<i>Gadus aeglefinus</i>	Haddock flesh	0.034	240	0.32	85
March	- " -	- " -	0.023	200	0.23	70
Sept	- " -	- " -	0.051	520	0.33	89
Dec	- " -	- " -	0.0170	122	0.141	45




Fig. 2.5.1. Cesium-137 levels in meat of cod (*Gadus callarias*) and Haddock (*Gadus aeglefinus*) collected in the Faroes, 1962-1981).

## 2.6. Strontium-90 in drinking water

Drinking-water samples were collected as previously but the samples were combined before the analysis as shown in Table 2.6.1. As in previous years, drinking water from Thorshavn contained more  $^{90}\text{Sr}$  than that from Tvarå (cf. the explanation in Risø Report No. 181<sup>1</sup>). The mean level in 1981 was  $4.4 \text{ Bq } ^{90}\text{Sr m}^{-3}$  ( $0.12 \text{ pCi l}^{-1}$ ), i.e. a little lower than in 1980.

Table 2.6.1. Strontium-90 in drinking water from the Faroes in 1981 (Unit:  $\text{Bq m}^{-3}$ )

	Thorshavn	Klaksvig	Tvarå
Jan-June	9.2	1.90	4.5
July-Nov	5.5	1.30	4.0
1981	7.4	1.60	4.2




Fig. 2.6.1. Strontium-90 in drinking water from the Faroes, 1962-1981.

Figure 2.6.1 shows the annual mean levels of  $^{90}\text{Sr}$  in drinking water from the three locations since 1962.

The tritium concentrations in Faroese drinking water (Table 2.6.2) did not follow the  $^{90}\text{Sr}$  levels. The mean tritium level in 1981 was  $3.9 \pm 1.1$  (1 S.D.)  $\text{kBq m}^{-3}$ , this is approximately 3/4 of the 1980 level.

Table 2.6.2. Tritium in drinking water from the Faroes in 1981. (Unit:  $\text{kBq m}^{-3}$ )

	Thorshavn	Klaksvig	Tværå
Jan-March	$3.1 \pm 0.2$	$3.0 \pm 0.4$	$4.1 \pm 0.4$
July-Sept	$5.2 \pm 1.5$	$5.0 \pm 0.2$	$2.8 \pm 0.5$

The error term is 1 S.E. of the mean of double determinations.

## 2.7. Strontium-90 and Cesium-137 in miscellaneous samples

### 2.7.1. Soil

In June we collected 3 soil cores (6.5 cm in diameter) at Höyvik. The samples were analysed for  $^{137}\text{Cs}$  and potassium. The two locations: F 13 and F 14 gave similar results down to 50 cm. The accumulated  $^{137}\text{Cs}$  level in these cores was  $4400 \text{ Bq m}^{-2}$  or  $118 \text{ mCi } ^{137}\text{Cs km}^{-2}$ . If the  $^{137}\text{Cs}/^{90}\text{Sr}$  ratio in fallout is 1.6 the calculated  $^{90}\text{Sr}$  fallout in these samples would have been  $2750 \text{ Bq m}^{-2}$  (or  $74 \text{ mCi km}^{-2}$ ). The third location (F 15) showed 50% higher  $^{137}\text{Cs}$  levels than the two others. In 1967<sup>1)</sup> we collected soil samples in the same area. At that time the accumulated fallout down to 50 cm was measured to  $8180 \text{ Bq } ^{137}\text{Cs m}^{-2}$  or  $221 \text{ mCi km}^{-2}$ . If we correct for decay and for new fallout since 1967 we would expect a level in 1981 similar to that found at F 15. We believe that this location has received some run off which makes the measured soil levels higher than that expected.

Figure 2.7.1. shows the distribution of the  $^{137}\text{Cs}$  activity at Höyvik as a function of depth in 1967 and 1981. Although the levels have decreased since 1967 it is remarkable that the distribution is nearly unchanged. Approximately 90% of the activity in the 50-cm layer is found in the upper 20-cm layer.

Table 2.7.1. Cesium-137 in soil collected at Höyvig at 3 locations, June 15, 1981

Depth in cm	F 13		F 15		F 14		F 13	F 15	F 14
	Bq kg <sup>-1</sup> dry w.	Bq m <sup>-2</sup>	Bq kg <sup>-1</sup> dry w.	Bq m <sup>-2</sup>	Bq kg <sup>-1</sup> dry w.	Bq m <sup>-2</sup>	g K kg <sup>-1</sup> dry w.		
0-10	128	2910	142	4320	-	-	1.21	1.43	-
10-20	27	870	45	1530	-	-	1.37	0.86	-
20-30	7.7	260	16	470	-	-	0.87	0.94	-
30-40	2.8	88	13	330	-	-	0.61	0.58	-
40-50	2.2	68	8.7	220	-	-	0.57	0.27	-
Σ 0-50		4196		6670	74	4560			1.04
50-70	2.4	138	7.1	320			0.72	0.62	
70-80	2.6	62	5.4	85			0.64	0.55	
Σ 50-80		200		405					




Fig. 2.7.1. Cesium-137 in Faroese soil samples collected in 1967 and 1981.

From earlier years' observations we estimate the accumulated fallout at Thorshavn to be  $2540 \text{ Bq } ^{90}\text{Sr m}^{-2}$  and that at Klaksvig to be  $4710 \text{ Bq } ^{90}\text{Sr m}^{-2}$  (cf. Fig. 2.1).



### 2.7.2. Sea water

Surface sea water was collected near Thorshavn on four occasions in 1981. The  $^{90}\text{Sr}$  mean level was  $3.0 \text{ Bq } ^{90}\text{Sr m}^{-3}$  and  $3.4 \text{ Bq } ^{137}\text{Cs m}^{-3}$ .

Table 2.7.2.1. Strontium-90 and Cesium-137 in surface sea water from the Faroes in 1981

Sampling month	Bq $^{90}\text{Sr m}^{-3}$	Bq $^{137}\text{Cs m}^{-3}$	Salinity o/oo
March	3.0	3.5	35.2
April	3.1	3.3	35.3
August	3.3	3.1	35.0
December	2.7	3.6	35.0

Figure 2.7.2 shows the  $^{90}\text{Sr}$  levels since 1962.




Fig. 2.7.2. Strontium-90 and Cesium-137 in Faroese sea water, 1962-1981.

The  $^{137}\text{Cs}/^{90}\text{Sr}$  ratio was  $1.13 \pm 0.17$  (1 S.D.), i.e. probably less than expected in ocean water.

Tritium was measured in Faroese sea water collected in 1981 (Table 2.7.2.2).

**Table 2.7.2.2. Tritium in surface sea water from the Faroes in 1981**

Month	$\text{kBq } ^3\text{H m}^{-3}$ $\pm 1 \text{ S.E.}$	Salinity o/oo
March	$1.48 \pm 0.37$	35.2
April	$1.66 \pm 0.56$	35.3
August	$0.92 \pm 0.18$	35.0
December	$1.85 \pm 0.37$	35.0

The error term is 1 S.E. of the mean of double determinations.

The mean concentration ( $\pm 1$  S.D.) was  $1.48 \pm 0.40 \text{ kBq } ^3\text{H m}^{-3}$ . This is lower than in 1980 where we found  $4.35 \pm 0.35$  ( $N = 1$ ). In North Sea water the tritium concentrations varied between 0.4 and 3.3  $\text{kBq m}^{-3}$ .

### **2.7.3. Sea plants**

Two samples of Laminaria were analysed in 1981. They contained 3.2 Bq  $^{90}\text{Sr kg}^{-1}$  ash (47 Bq  $^{90}\text{Sr (kg Ca)}^{-1}$ ) and 3.4 Bq  $^{137}\text{Cs kg}^{-1}$  ash (17 Bq  $^{137}\text{Cs (kg K)}^{-1}$ ). Table 2.7.3.2 shows the results of our own sampling in the Faroes in June 1981.

**Table 2.7.3.1. Strontium-90 and Cesium-137 in sea plants from the Faroes in 1981**

Species	Sampling month	Bq $^{90}\text{Sr (kg ash)}^{-1}$	Bq $^{90}\text{Sr (kg Ca)}^{-1}$	Bq $^{137}\text{Cs (kg ash)}^{-1}$	Bq $^{137}\text{Cs (kg K)}^{-1}$
Laminaria hyperborea	April	4.3	62	4.0	18.6
Laminaria saccharina	August	2.1	32	2.8	15.0
Rhodomenia palmata	August	1.24	53	5.1	27

Table 2.7.3.2. Radioisotopes in Farnese sea plants collected in June 1981. (Unit: Bq kg<sup>-1</sup> dry weight)

Location	Station No.	Species	<sup>70</sup> Se	<sup>80</sup> Se kg S kg <sup>-1</sup>	<sup>134</sup> Mn	<sup>95</sup> Zr	<sup>95</sup> Nb	<sup>106</sup> Ru	<sup>125</sup> Sb	<sup>137</sup> Cs	<sup>141</sup> Co	<sup>144</sup> Co	<sup>238</sup> Bu	<sup>239,240</sup> Pu	<sup>238</sup> Am 238,239Pu	<sup>241</sup> Am 240,241Pu	dry weight fraction
Bayish 62°02'N 6°45'W	P11	Fucus spiralis	30 (5)	0.032 (0.4)	1.4 (9)	10 (2)	30 (1)	14 (7)		1.01 (10)	4.8 (12)	25 (2)	1.5 (8)	0.31 (5)			4.10
- " -	P10	Laminaria digitata		0.071 (0.3)		7.2 (6)	16 (2)			0.63 (10)	4 (17)	26 (3)	2.1 (7)	0.050 (0)			0.70
Cydov 62°20'N 6°16'W	P17	Fucus disticus		0.042 (0.3)	3.4 (5)	24 (10)	56 (4)	10 (7)	2.0 (11)	1.11 (9)		44 (2)	2.6 (5)	0.076 (4)	0.04		4.17
- " -	P16	Laminaria digitata		0.092 (0.2)		15 (2)	22 (12)			0.00 (15)		35 (3)	2.1 (7)	0.040 (5)			0.16
Steymer-Systurøy 62°13'N 7°00'W	P18	Ascophyllus nodosum		0.023 (0.5)	0.53 (2)	10 (2)	21 (9)	7 (15)		0.63 (12)		11 (7)	1.5 (7)	0.160 (10)		0.08	0.20

In brackets: vol. S.D. in % due to counting.

Table 2.7.3.2 compared with Table C.1 in Appendix C shows that the concentration ratio:  $\frac{\text{Bq kg}^{-1} \text{ dry w. Fucus}}{\text{Bq l}^{-1} \text{ sea water}}$  is 300 for  $^{137}\text{Cs}$  and  $(0.4 - 1.0) \cdot 10^4$  for Pu. The ratios for laminaria were 200 and  $0.3 \cdot 10^4$ , respectively. In Danish waters<sup>2)</sup> the concentration ratio for  $^{137}\text{Cs}$  in Fucus varied between 200 and 700 showing an increase with decreasing salinity. In the case of Pu, Danish waters<sup>2)</sup> showed concentration factors to fucoids at  $(1-3) \cdot 10^4$ . The concentration ratios in Faroese waters were similar to those in high-salinity Danish waters (cf. also Appendix C).




Fig. 2.7.3. Strontium-90 ( $\text{Bq (kg Ca)}^{-1}$ ) in sea plants collected at Thorshavn, 1962-1981.

#### 2.7.4. Vegetables

One sample of potatoes was analysed in 1981. It contained  $0.141 \text{ Bq } ^{90}\text{Sr kg}^{-1}$  ( $3100 \text{ Bq } ^{90}\text{Sr (kg Ca)}^{-1}$ ) and  $0.89 \text{ Bq } ^{137}\text{Cs kg}^{-1}$  ( $200 \text{ Bq } ^{137}\text{Cs (kg K)}^{-1}$ ). Both levels were lower than those observed in 1980.




Fig. 2.7.4.1. Cesium-137 in Faroese potatoes, 1962-1981.




Fig. 2.7.4.2. Strontium-90 in Faroese potatoes, 1962-1981.

### 2.7.5. Bread

Rye bread and white bread were collected at Thorshavn in June. The levels in white bread were 0.19 Bq  $^{90}\text{Sr}$   $\text{kg}^{-1}$  and 0.10 Bq  $^{137}\text{Cs}$   $\text{kg}^{-1}$ . The rye bread collected in 1981 contained 0.37 Bq  $^{90}\text{Sr}$   $\text{kg}^{-1}$  and 0.25 Bq  $^{137}\text{Cs}$   $\text{kg}^{-1}$ , i.e. the bread levels were higher than the 1980 levels.

Table 2.7.5. Strontium-90 and Cesium-137 in Faroese bread in June 1981

Sort	Bq $^{90}\text{Sr}$ $\text{kg}^{-1}$	Bq $^{90}\text{Sr}$ (kg Ca) $^{-1}$	Bq $^{137}\text{Cs}$ $\text{kg}^{-1}$	Bq $^{137}\text{Cs}$ (kg K) $^{-1}$
White bread	0.192	560	0.100	57
Rye bread	0.37	280	0.25	114

The  $^{137}\text{Cs}$  and  $^{90}\text{Sr}$  (kg $^{-1}$ ) levels in Faroese bread were generally lower than the corresponding Danish<sup>2)</sup>.

### 2.7.6. Eggs

Eggs were collected from Thorshavn in June 1981. The levels of hens eggs were 0.030 Bq  $^{90}\text{Sr}$   $\text{kg}^{-1}$  (92 Bq (kg Ca) $^{-1}$ ) and < 0.04 Bq  $^{137}\text{Cs}$   $\text{kg}^{-1}$ .

## 2.8. Humans

### 2.8.1. Strontium-90 in human bone

In 1981 seven human bone samples representing 9 individuals from Dronning Alexandrine's Hospital in Thorshavn were analysed. Table 2.8.1 shows the results.

The mean level in bone of newborn infants was 95 Bq  $^{90}\text{Sr}$  (kg Ca) $^{-1}$  (2.6 pCi  $^{90}\text{Sr}$  (g Ca) $^{-1}$ ).

The adult bone samples were all femur and the mean content was 62 Bq  $^{90}\text{Sr}$  (kg Ca) $^{-1}$  (1.7 pCi  $^{90}\text{Sr}$  (g Ca) $^{-1}$ ).

The bone levels in 1981 were in general higher than those observed in 1980. Compared to Danish bones in 1981<sup>2)</sup> the Faroese bones contained approximately two times as much  $^{90}\text{Sr}$ .

**Table 2.8.1. Strontium-90 in human bone collected in the Faroes in 1981**

Age	Bone type		Sex	Bq $^{90}\text{Sr}$ (kg Ca) $^{-1}$	S.U.
0	Vertebrae		M	92 B	2.5
0	Vertebrae		M	98 A	2.6
61 years	Femur	Amputation	F	59	1.60
79 years	Femur	- " -	F	59	1.59
82 years	Femur	- " -	M	78	2.11
82 years	Femur	- " -	F	75	2.03
86 years	Femur	- " -	M	38	1.03

### 3. ESTIMATE OF THE MEAN CONTENTS OF $^{90}\text{Sr}$ AND $^{137}\text{Cs}$ IN THE HUMAN DIET

#### 3.1. Annual quantities

The annual quantities are still based on the estimate made by Professor E. Hoff-Jørgensen, Ph.D., in 1962<sup>1)</sup> assuming a daily per caput intake of approximately 3000 calories (12.6 MJ).

#### 3.2. Milk and cream

75% of the milk consumed in the Faroes is assumed to be of local origin, and 25% comes from Denmark. Hence the  $^{90}\text{Sr}$  content in milk consumed in the Faroes in 1981 was  $1.2 \cdot (0.75 \cdot 0.25 + 0.25 \cdot 0.108) = 0.290 \text{ Bq } ^{90}\text{Sr kg}^{-1}$ , and the  $^{137}\text{Cs}$  content was  $0.75 \cdot 5.1 + 0.25 \cdot 0.134 = 3.86 \text{ Bq } ^{137}\text{Cs kg}^{-1}$  (cf. 2.3 and ref. 2). 1 kg milk contains 1.2 g Ca.

#### 3.3. Cheese

Nearly all cheese consumed in the Faroes is of Danish origin, and the Danish figures from ref. 2 were used:  $0.92 \text{ Bq } ^{90}\text{Sr kg}^{-1}$  and  $0.10 \text{ Bq } ^{137}\text{Cs kg}^{-1}$ .

#### 3.4. Grain products

As most grain products are imported from Denmark, the Danish figures for 1981<sup>2)</sup> were used in the calculation of the Faroese levels. The mean daily consumption of grain products in the Faroes is, as in Denmark, 80 g rye flour, 120 g wheat flour, and 20 g grits. Hence the mean concentration of  $^{90}\text{Sr}$  in grain products consumed in the Faroes in 1981 is  $0.42 \text{ Bq } ^{90}\text{Sr kg}^{-1}$  and  $0.33 \text{ Bq } ^{137}\text{Cs kg}^{-1}$ .



### 3.5. Potatoes

All potatoes consumed in the Faroes are assumed to be of local origin. The values from Table 2.7.4 were used, i.e. 0.14 Bq  $^{90}\text{Sr}$   $\text{kg}^{-1}$  and 0.89 Bq  $^{137}\text{Cs}$   $\text{kg}^{-1}$ .

### 3.6. Other vegetables and fruit

As the amount of vegetables and fruit grown in the Faroes is limited, the Danish figures from 1981<sup>2)</sup> were used. Thus the mean contents in vegetables other than potatoes were 0.38 Bq  $^{90}\text{Sr}$   $\text{kg}^{-1}$  and 0.086 Bq  $^{137}\text{Cs}$   $\text{kg}^{-1}$ , and the mean contents in fruit were 0.044 Bq  $^{90}\text{Sr}$   $\text{kg}^{-1}$  and 0.09 Bq  $^{137}\text{Cs}$   $\text{kg}^{-1}$ .

### 3.7. Meat and eggs

Meat and egg consumption in the Faroes is estimated to consist of 50% locally produced mutton (or lamb), 25% local whale meat, and 25% sea birds and eggs.

The mutton contained 0.32 Bq  $^{90}\text{Sr}$   $\text{kg}^{-1}$  and 97 Bq  $^{137}\text{Cs}$   $\text{kg}^{-1}$  (cf. 2.4). Whale meat from 1980 contained 0.04 Bq  $^{90}\text{Sr}$   $\text{kg}^{-1}$  and 0.19 Bq  $^{137}\text{Cs}$   $\text{kg}^{-1}$ , sea birds from 1979 and eggs from 1981 (cf. 2.7.6): 0.007 Bq  $^{90}\text{Sr}$   $\text{kg}^{-1}$  and 0.030 Bq  $^{90}\text{Sr}$   $\text{kg}^{-1}$ , and 0.27 and  $\sim 0.04$  Bq  $^{137}\text{Cs}$   $\text{kg}^{-1}$  respectively.

Hence we estimate the mean content of  $^{90}\text{Sr}$  in meat and eggs consumed in 1981 to be

$$0.50 \cdot 0.32 + 0.25 \cdot 0.04 + 0.25 \cdot \left( \frac{0.007 + 0.030}{2} \right) = 0.175 \text{ Bq } ^{90}\text{Sr } \text{kg}^{-1}$$

and the  $^{137}\text{Cs}$  content to be

$$0.50 \cdot 97 + 0.25 \cdot 0.19 + 0.25 \cdot \left( \frac{0.27 + 0.04}{2} \right) = 48.6 \text{ Bq } ^{137}\text{Cs } \text{kg}^{-1}.$$

### 3.8. Fish

All fish consumed in the Faroes is of local origin, and the mean contents in fish, obtained from subsection 2.5, were 0.031 Bq  $^{90}\text{Sr}$   $\text{kg}^{-1}$  and 0.34 Bq  $^{137}\text{Cs}$   $\text{kg}^{-1}$ .

### 3.9. Coffee and tea

The Danish figures for 1981<sup>2)</sup> were used, i.e. 0.66 Bq  $^{90}\text{Sr}$   $\text{kg}^{-1}$  and 2.21 Bq  $^{137}\text{Cs}$   $\text{kg}^{-1}$ .

### 3.10. Drinking water

The mean value found in Table 2.6.1 was used, i.e. 0.0044 Bq  $^{90}\text{Sr}$   $\text{kg}^{-1}$ . The  $^{137}\text{Cs}$  content was estimated to be approximately one fourth (the ratio found in New York tap water in 1964<sup>4)</sup>) of the  $^{90}\text{Sr}$  content, i.e. 0.001 Bq  $^{137}\text{Cs}$   $\text{kg}^{-1}$ .

Tables 3.1 and 3.2 show the diet estimates of  $^{90}\text{Sr}$  and  $^{137}\text{Cs}$  respectively.

**Table 3.1. Estimate of the mean content of  $^{90}\text{Sr}$  in the human diet in the Faroe Islands in 1981**

Type of food	Annual quantity in kg	Bq $^{90}\text{Sr}$ per kg	Total Bq $^{90}\text{Sr}$	Percentage of total Bq $^{90}\text{Sr}$ in food
Milk and cream	146	0.29	42.34	35.2
Cheese	7.3	0.92	6.72	5.6
Grain products	80	0.42	33.60	27.9
Potatoes	91	0.14	12.74	10.6
Vegetables	20	0.38	7.60	6.3
Fruit	18	0.044	0.79	0.7
Meat and eggs	37	0.175	6.48	5.4
Fish	91	0.031	2.82	2.3
Coffee and tea	7.3	0.66	4.82	4.0
Drinking water	548	0.0044	2.41	2.0
Total			120.32	

The mean annual calcium intake is estimated to be 0.6 kg (approx. 200-250 g of creta praeparata). Hence the ratio: Bq  $^{90}\text{Sr}$  (kg Ca) $^{-1}$  in total Faroese diet was 200 (5.4 pCi  $^{90}\text{Sr}$  (g Ca) $^{-1}$ ).

**Table 3.2.** Estimate of the mean content of  $^{137}\text{Cs}$  in the human diet in the Faroe Islands in 1981

Type of food	Annual quantity in kg	Bq $^{137}\text{Cs}$ per kg	Total Bq $^{137}\text{Cs}$	Percentage of total Bq $^{137}\text{Cs}$ in food
Milk and cream	146	3.86	563.6	22.4
Cheese	7.3	0.10	0.7	0
Grain products	80	0.33	26.4	1.1
Potatoes	91	0.89	81.0	3.2
Vegetables	20	0.086	1.7	0.1
Fruit	18	0.09	1.6	0.1
Meat and eggs	37	48.6	1798.2	71.3
Fish	91	0.34	30.9	1.2
Coffee and tea	7.3	2.21	16.1	0.6
Drinking water	548	0.001	0.5	0
Total			2520.7	
The mean annual intake of potassium is estimated to be approx. 1.2 kg. Hence the ratio: Bq $^{137}\text{Cs}$ (kg K) $^{-1}$ becomes 2100 (56.7 pCi $^{137}\text{Cs}$ (g K) $^{-1}$ ).				

### 3.11. Discussion

Figures 3.1 and 3.2 show the Faroese diet levels since 1962.

The 1981  $^{90}\text{Sr}$  level in the total diet was nearly equal to the 1980 concentration, but the  $^{137}\text{Cs}$  level was 44% higher than that observed in 1980.

The main contributors to the  $^{90}\text{Sr}$  content in the Faroese diet were milk products, cereals and potatoes, which together accounted for approximately 79% of the total  $^{90}\text{Sr}$  content in the diet in 1981. As regards  $^{137}\text{Cs}$ , milk products, meat (lamb) and potatoes were the most important contributors. In 1981, 97% of the total  $^{137}\text{Cs}$  content in the diet originated from these products.




Fig. 3.1. Strontium-90 in Faroese diet, 1962-1981.




Fig. 3.2. Cesium-137 in Faroese diet, 1962-1981.

The Faroese mean diet contained 1.3 times as much  $^{90}\text{Sr}$  and approximately 16 times as much  $^{137}\text{Cs}$  as the Danish diet in 1981<sup>2)</sup>.

As earlier<sup>1)</sup> mentioned, the year-to-year variations in the  $^{137}\text{Cs}$  estimates for Faroese diet are markedly influenced by the mutton and potatoe samples obtained for analysis.

#### 4. CONCLUSION

##### 4.1.

The  $^{90}\text{Sr}$  fallout rate in the Faroes in 1981 was approximately 25 Bq  $^{90}\text{Sr m}^{-2}$  (0.67 mCi  $\text{km}^{-2}$ ). The accumulated fallout by the end of 1981 was estimated at approximately 3600 Bq  $^{90}\text{Sr m}^{-2}$  (98 mCi  $\text{km}^{-2}$ ) (the mean at Thorshavn and Klaksvig).

##### 4.2.

The mean level of  $^{90}\text{Sr}$  in Faroese milk was 250 Bq  $(\text{kg Ca})^{-1}$  (6.6 pCi  $(\text{g Ca})^{-1}$ ). The  $^{137}\text{Cs}$  concentration was 5100 Bq  $^{137}\text{Cs m}^{-3}$  (138 pCi  $\text{l}^{-1}$ ).

Mutton contained 0.32 Bq  $^{90}\text{Sr kg}^{-1}$  (8.6 pCi  $\text{kg}^{-1}$ ) and 97 Bq  $^{137}\text{Cs kg}^{-1}$  (2.6 nCi  $\text{kg}^{-1}$ ). Fish showed a mean level of 0.34 Bq  $^{137}\text{Cs kg}^{-1}$  (9.2 pCi  $\text{kg}^{-1}$ ).

The mean content of  $^{90}\text{Sr}$  in drinking water was 4.4 Bq  $\text{m}^{-3}$  (0.12 pCi  $\text{l}^{-1}$ ).

The mean daily per caput intakes resulting from the Faroese diet in 1981 were estimated at 0.33 Bq  $^{90}\text{Sr}$  (8.9 pCi  $\text{d}^{-1}$ ) and 6.9 Bq  $^{137}\text{Cs}$  (186 pCi  $\text{d}^{-1}$ ).

##### 4.3.

From the measurements on Faroese human bones, the Faroese bone level in 1981 was estimated at 70 Bq  $^{90}\text{Sr (kg Ca)}^{-1}$  (2 pCi  $(\text{g Ca})^{-1}$ ).

The mean content of  $^{137}\text{Cs}$  in the Faroese adult was estimated at approximately  $6000 \text{ Bq } ^{137}\text{Cs} (\text{kg K})^{-1}$  ( $160 \text{ pCi (g K)}^{-1}$ ). This estimate is based on the diet estimate.

#### ACKNOWLEDGEMENTS

The authors wish to thank Miss Anna Holm Petersen, Mrs. Jytte Clausen, Mrs. Else Sørensen, Mrs. Anna Madsen, Mrs. Oda Brandstrup and Mrs. Helle Porsdal for their conscientious performance of the analyses.

Our special thanks are due to the staff of the pharmacy "Tjaldur" in Thorshavn and to all other persons in the Faroes who have contributed by collecting samples.



## REFERENCES

- 1) Environmental Radioactivity in the Faroes 1962-1978. Riss Reports Nos. 64, 86, 108, 131, 155, 181, 202, 221, 246, 266, 292, 306, 324, 346, 362, 387, 404, 422 and 448 (1963-1981).
- 2) A. Aarkrog, L. Bøtter-Jensen, H. Dahlgaard, Heinz Hansen, J. Lippert, S.P. Nielsen and Karen Nilsson, Environmental Radioactivity in Denmark in 1981. Riss Report No. 469 (1981).
- 3) A. Aarkrog, Henning Dahlgaard, Heinz Hansen, Elis Holm, J. Lippert and Karen Nilsson, Environmental Radioactivity in Greenland in 1981. Riss Report No. 471 (1981).
- 4) E.P. Hardy, Jr., and Joseph Rivera, Fallout Program Quarterly Summary Report (March 1, 1965, through June 1, 1965). HASL-161 (1965).
- 5) B.G. Bennett and C.S. Klusek, Strontium-90 in human bone, EML-344 (1978).
- 6) A. Aarkrog, Environmental Studies on Radioecological Sensitivity and Variability. Riss-R-437 (June 1979).
- 7) G.C. Christensen, Radioaktivitet i blæretang (P. Ves) langs Norskekysten 1980-81 (in Norwegian). 3. Nordiske Radioøkologiseminar, Hyvinkää 11-13 May 1982.
- 8) A. Aarkrog, H. Dahlgaard and K. Nilsson, Studies of the Distribution of Transuranics in the Baltic Sea, The Danish Belts, The Kattegat and the North Sea. IAEA-TECDOC-265, p. 23-32 (1982).

## APPENDIX A

The models used for the predictions shown in Table A were based on data collected 1962-1976<sup>6)</sup>. If the predictions for previous years 1977-1980<sup>1)</sup> were considered too, we conclude that the model for  $^{90}\text{Sr}$  in milk overestimates the level and so do the models: for  $^{90}\text{Sr}$  in drinking water from Klaksvig, for  $^{137}\text{Cs}$  in milk from Tverå and for  $^{137}\text{Cs}$  in potatoes. The following models underestimate the concentrations:  $^{90}\text{Sr}$  in cod fish,  $^{90}\text{Sr}$  in newborn bone, and probably also  $^{137}\text{Cs}$  in milk from Klaksvig.

Table A. Comparison between observed and predicted  $^{90}\text{Sr}$  and  $^{137}\text{Cs}$  concentrations in Faroese samples collected in 1981

Sample	Unit	Observed ±1 S.E.	Number of samples	Predicted	Obs./pre. ±1 S.E.	Model is ref. 6
Drinking water, Thorshavn	Bq $^{90}\text{Sr}$ m <sup>-3</sup>	7.4 ±1.6	2	17.4	0.42±0.10	C.1.4.1 No. 9
- " - , Klaksvig	- " -	1.6 ±0.3	2	3.2	0.50±0.09	- " - No. 10
- " - , Tverå	- " -	4.2 ±0.3	2	4.8	0.88±0.06	- " - No. 11
Sea water	- " -	3.0 ±0.13	4	2.5	1.20±0.05	C.1.5.1 No. 3
Rye bread	Bq $^{90}\text{Sr}$ kg <sup>-1</sup>	0.37	1	0.78	0.45	C.2.3.1 No. 6
White bread	- " -	0.27	1	0.19	1.42	- " - No. 7
Rye bread	Bq $^{137}\text{Cs}$ kg <sup>-1</sup>	0.25	1	0.49	0.51	- " - No. 8
White bread	- " -	0.10	1	0.19	0.53	- " - No. 9
Grass	Bq $^{90}\text{Sr}$ (kg Ca) <sup>-1</sup>	8850 ±2550	2	9350	0.95±0.27	C.2.4.1 No. 4
- " -	Bq $^{137}\text{Cs}$ (kg K) <sup>-1</sup>	5200 ±3400	2	2275	2.29±1.50	C.2.4.2 No. 3
Potatoes	Bq $^{90}\text{Sr}$ kg <sup>-1</sup>	0.14	1	0.23	0.61	C.2.5.1 No. 11
- " -	Bq $^{137}\text{Cs}$ kg <sup>-1</sup>	0.9	1	7.0	0.13	C.2.5.3 No. 8
Brown algae	Bq $^{90}\text{Sr}$ (kg Ca) <sup>-1</sup>	47 ±15	2	116	0.40±0.13	C.2.7.1 No. 5
Milk	- " -	250 ±18	12	435	0.57±0.04	C.3.3.1 No. 1
Milk Thorshavn	Bq $^{137}\text{Cs}$ m <sup>-3</sup>	2370 ±265	12	2620	0.90±0.10	C.3.3.2 No. 7
Milk Klaksvig	- " -	5500 ±240	12	2700	2.04±0.08	- " - No. 9
Milk Tverå	- " -	6900 ±860	12	12650	0.55±0.07	- " - No. 11
Mutton	Bq $^{90}\text{Sr}$ (kg Ca) <sup>-1</sup>	3200	1	2000	1.60	C.3.4.1 No. 5
- " -	Bq $^{137}\text{Cs}$ (kg K) <sup>-1</sup>	30000	1	6000	5.00	C.3.4.2 No. 5
Sheep bone	Bq $^{90}\text{Sr}$ (kg Ca) <sup>-1</sup>	3300	1	2820	1.17	C.3.4.3 No. 1
Cod fish	- " -	220 ±40	8	30	7.3 ±1.3	C.3.5.1 No. 3
- " -	Bq $^{137}\text{Cs}$ kg <sup>-1</sup>	0.34 ±0.08	8	0.24	1.42±0.34	C.3.5.2 No. 2
Newborn bone	Bq $^{90}\text{Sr}$ (kg Ca) <sup>-1</sup>	95 ±3	2	9.5	10 ±0.3	C.4.3.1 No. 15

## APPENDIX B

### Algae from West Norway

In 1980, Gordon Christensen, of the Institute of Energy Technology, and T. Bertelsen, of the Norwegian Institute of Radiation Hygiene, sampled algae along the entire Norwegian coastline. A number of these samples have in an intercalibration effort been analysed at Risø. The results of the plutonium determinations are shown in Table B.1.

Risø participated at two of the Norwegian locations, Bud and Vågsøy, and made supplementary samplings there (cf. Figs. B.1 and B.2). Table B.2 shows the Pu determinations of these samples.

Table B.1. Plutonium-239,240 in *Fucus vesiculosus* collected along the Norwegian Westcoast in August-September 1980

Location (position)	Sample No.	Date	mBq kg <sup>-1</sup> dry w.	rel. S.D. in %	wet weight dry weight
Jacobselv (70°N, 31°E)	1-2	Aug 11	184	9	4.9
Hammerfest (71°N, 24°E)	4A-3	Aug 14	270	6	3.4
Vestvågøy (68°N, 14°E)	6-4	Aug 20	140	9	3.8
Kvaløya (65°N, 12°E)	7B-2	Aug 23	110	12	3.6
Lista (58°N, 7°E)	11-3	Sept 14	132	10	4.2
Tromøya (58°30'N, 9°E)	12-1	Sept 15	89	11	3.9
Tjøme (59°N, 10°E)	13-4	Sept 15	75	9	4.6
Hvaler (59°N, 11°E)	14-6	Sept 26	86	14	5.8

**Table B.2.** Plutonium-239,240 in sea plants collected in September 1980 at Bud, W-Norway (62°N, 6°E)

Sample No.	Species	mBq kg <sup>-1</sup> dry w.	rel. S.D. %
Vågsøy 2	<i>Fucus vesiculosus</i>	81	11
Bud 3	<i>Laminaria digitata</i>	112	7
Bud 6	<i>Fucus vesiculosus</i>	100	13
Bud 7	<i>Pelvetia</i>	84	14
Bud 16	<i>Pelvetia</i>	67	20
Bud 17	<i>Fucus spiralis</i>	98	18
Bud 18	<i>Fucus vesiculosus</i>	87	15
Bud 19	<i>Ascophyllum nodosum</i>	110	8
Bud 20	<i>Fucus serratus</i>	153	7

The plutonium concentrations in *Fucus vesiculosus* showed a decreasing tendency from north to south. A single location: Lista at the southwest corner of Norway did not follow the general pattern. Whether or not this was due to a contribution of plutonium from Windscale at this location is unclear. If the plutonium results are compared with those of Christensen<sup>7)</sup> on the same samples, the mean ratio of ours to Christensen's is  $1.18 \pm 0.27$  (1 S.D., N = 8). The relative standard deviations of double Pu determinations carried out by the two laboratories were thus 23%. These are the same deviations as found earlier in Nordic Pu-intercomparison runs<sup>8)</sup> on seaweed and sediments.

The relative SD at a single location: Bud (Table B2) between the 9 samples obtained at this location was 27%. The 9 samples represented 6 different species. The local variation along the entire coastline of *Fucus vesiculosus* alone was 48%, if we use the data in Table B.1.

The mean of all seaweed samples in Tables B1 and B2 was 116 mBq <sup>239,240</sup>Pu kg<sup>-1</sup> dry weight (1 SD: 50, N = 17). The corresponding fresh weight concentration was 25 mBq kg<sup>-1</sup> if the mean ratio of fresh to dry weight was 4.6 as found by Christensen in 1980<sup>7)</sup>. The mean of the 15 samples analysed by Christensen was 26 mBq kg<sup>-1</sup> wet weight (1 S.D.:11).





Fig. B.1. Sampling locations in West-Norway, Bud and Vågsøy, 1980.



**Fig. B.2.** Sampling locations in the Bud area, West Norway, 1980. Sample numbers are indicated.

## Appendix C

### Sea water and biota collected in Iceland and on the Bergen-Faroe Islands-Iceland route in June-July 1981

Surface sea water was collected from M/S "Smyril" on its way from Norway to Iceland and back again 18 days later. Table C.1 and Figs. C.1 and C.2 show the  $^{137}\text{Cs}$  and  $^{239,240}\text{Pu}$  concentrations, respectively. Between the Faroes and Norway the samples were collected twice at nearly the same locations. The upper figures are the concentrations found on the way out and the lower ones on the way back to Norway. It is evident that the  $^{137}\text{Cs}$  concentrations, in particular, reproduce convincingly.

Close to the Norwegian coast the surface water contained  $^{137}\text{Cs}$  levels which were nearly 30 times higher than in the open Atlantic Ocean. The presence of  $^{134}\text{Cs}$  suggested that the activity came from Sellafield. The concentrations decreased rapidly on moving westwards. At Iceland the lowest concentrations were found on the west coast (cesium-137 levels there were nearly 30% lower than those observed at the Faroe Islands. A single water sample collected east of Iceland (S 57) contained more  $^{137}\text{Cs}$  than expected. This sample may have been contaminated because the ship had just arrived from Scottish waters when this sample was collected.

The  $^{239,240}\text{Pu}$  concentrations were also lower in the waters around Iceland than between Norway and the Faroes, but the difference was less marked than for  $^{137}\text{Cs}$ , and the presence of any surplus plutonium originating from Sellafield was not demonstrated in these samples. On the average the  $^{241}\text{Am}$  concentration was 20% of the plutonium activity. A few samples were analysed for particulate Pu and Am (Table C.1 shows the results).

Table C.1. Radiocesium, plutonium and americium in surface sea water collected in June-July 1981 between Norway, The Faroe Islands and Iceland. (Unit: Bq m<sup>-3</sup>)

Latitude N	Longitude	No. & Location	Date	<sup>137</sup> Cs	<sup>134</sup> Cs <sup>137</sup> Cs	<sup>239,240</sup> Pu	<sup>241</sup> Am <sup>239,240</sup> Pu	β particulate activity Pu Am		Salinity in ‰
60°45'	0°41'E	S1	June 13	91 (10)	0.039 (80)	0.019 (160)	0.4 (310)			31.74
61°06'	2°00'E	S2	June 13	57 (10)	0.049 (100)	0.014 (160)	0.3 (200)			34.75
61°19'	0°45'E	S3	June 13	5.4 (60)	-	0.021 (230)	0.2 (460)			35.21
61°30'	1°22'W	S4	June 14	3.0 (90)	-	0.017 (110)	0.15 (310)			35.21
61°40'	3°23'W	S5	June 14	4.2 (80)	-	0.017 (150)	0.3 (310)			35.21
61°51'	5°20'W	S6	June 14	3.5 (100)	-	0.017 (260)	0.15 (390)			35.21
62°02'	6°45'W	P7-9 Boyvik	June 14	3.55 (30)	-	0.020 (110)	2.17 (250)	13	30	33.54
60°00'	16°25'W	120 Krossness	June 17			0.017 (130)	0.3 (200)			30.30
63°50'	22°27'W	122-24 Grindavik	June 19	2.77 (40)	-	0.010 (190)	0.2 (440)	9	15	32.36
66°07'	23°06'W	127, 35, 36 Hnífsdalur	June 22	2.92 (40)	-	0.014 (110)	0.1 (510)	6	4	34.10
66°09'	18°55'W	137, 138 Siglufjörður	June 24	3.19 (30)	-	0.010 (140)	0.15 (430)			32.55
66°20'	15°55'W	140, 149 Raufarhöfn	June 26	3.10 (40)	-	0.010 (120)	0.1 (1000)			31.21
65°10'	13°43'W	153, 154 Vestmanna- strandur	June 28	3.19 (40)	-	0.010 (120)	-			28.58
65°03'	12°57'W	S57	July 1	5.5 (40)	-	0.010 (230)	0.2 (460)			34.03
63°43'	9°52'W	S58	July 1	3.3 (80)	-	0.012 (190)	0.2 (530)			35.21
62°53'	0°00'W	S59	July 1	3.2 (90)	-	0.014 (160)	0.2 (470)			34.97
61°59'	6°02'W	S60	July 1	3.4 (100)	-	0.026 (340)	0.1 (440)			34.96
61°44'	3°47'W	S61	July 2	4.0 (90)	-	0.009 (170)	0.2 (430)			35.09
61°30'	1°31'W	S62	July 2	3.9 (80)	-	0.010 (210)	0.2 (400)			35.09
61°20'	0°42'E	S63	July 2	5.7 (70)	-	0.016 (160)	0.2 (410)			35.09
61°00'	2°55'E	S64	July 2	50 (10)	0.056 (120)	0.025 (250)	0.1 (560)			33.41
60°45'	4°36'E	S65	July 2	83 (10)	0.041 (70)	0.014 (160)	0.2 (300)			31.74

In brackets relative S.D. due to counting error.






Fig. C.1. Cesium-137 in surface sea water collected June-July 1981. (Unit: Bq m<sup>-3</sup>).

Fig. C.2. Plutonium-239,240 in surface sea water collected June-July 1981. (Unit: mBq m<sup>-3</sup>).

Algae were collected at the Faroe Islands (Table 2.7.3.2) and along the Icelandic coastline (Table C.2 and Fig. C.3). *Fucus distichus* and *Fucus vesiculosus* were the most frequent species. An anova showed that the radionuclide concentrations in these two species did not differ significantly. The anova also showed that samples in the eastern locations in Iceland had approximately 30% higher concentrations of radionuclides than those from the north and west. (In the anova we included 7 isotopes:  $^{54}\text{Mn}$ ,  $^{95}\text{Nb}$ ,  $^{106}\text{Ru}$ ,  $^{137}\text{Cs}$ ,  $^{144}\text{Ce}$ ,  $^{226}\text{Ra}$ , and  $^{239,240}\text{Pu}$ , and 3 areas: West: Grindavik, Skalanes, North: Hnifsdalur, Arnarnes, Siglufjordur, Raufarhöfn and East: Seydisfjordur, Neskaupstadur, Streithvarf). The interactions among species, nuclides, or locations were not significant.

The radionuclides  $^{95}\text{Zr}$ ,  $^{95}\text{Nb}$ ,  $^{106}\text{Ru}$ ,  $^{125}\text{Sb}$ , and  $^{144}\text{Ce}$  occurred in the same relative mutual concentrations in fucoids as in air collected in June 1981 at Pisa<sup>2)</sup>. This suggests that the contamination of the fucoids with these radionuclides was an adsorption of fallout particles suspended in the water.

Another pathway may have been uptake directly from the air if the plants have been exposed to the atmosphere during low tide. The relative concentrations of  $^{54}\text{Mn}$  and  $^{239,240}\text{Pu}$  compared with the other radionuclides were, respectively, 3-5 times higher in fucoids than in fresh fallout. This suggests that these nuclides were present in the water in dissolved form also, because only if a nuclide is in solution can we expect that the uptake by algae will differ for the various elements. The ratios:  $^{95}\text{Zr}/^{137}\text{Cs}$ ,  $^{95}\text{Nb}/^{137}\text{Cs}$ ,  $^{106}\text{Ru}/^{137}\text{Cs}$ ,  $^{125}\text{Sb}/^{137}\text{Cs}$ , and  $^{144}\text{Ce}/^{137}\text{Cs}$  in fucoids were on the average 50% higher than those in air. This suggests that  $^{137}\text{Cs}$  has been dissolved from the fallout particles absorbed by the algae in contrast to the other five fallout nuclides. Furthermore, it shows that the concentration factor is lower for  $^{137}\text{Cs}$  in solution than for dissolved  $^{54}\text{Mn}$  and  $^{239,240}\text{Pu}$ .

Table C.3 shows the concentration ratios for  $^{137}\text{Cs}$  and  $^{239,240}\text{Pu}$  in fucoids. These are slightly lower than, but not significantly different from, those calculated for the Faroe Islands in 2.7.3.

**Table C.3. Radionuclides in algae collected around the Icelandic coastline in June-July 1991.**  
(Unit: Bq kg<sup>-1</sup> dry weight)

Date	Location (position)	Station No.	Species	40K kg kg <sup>-1</sup>	54Mn	95Sr	99Mo	106Ru	135Ba	137Cs	144Ce	226Ra	230,232Pu	241Am 241Am, 241Pu	dry weight 770K-0.1241
June 17	Streitshvarf (64°45'N 16°00'W)	119	Fucus distichus	0.046 (0.40)		9 (270)	15 (150)			0.60 (100)	3.0 (200)	1.4 (100)	0.034 (70)		0.11
June 18	Vik (63°35'N 19°00'W)	121	Fucus distichus	0.041 (0.40)	0.0 (250)		20 (140)	8 (230)		0.67 (170)	4.7 (100)	2.0 (80)	0.007 (130)		0.20
June 18	Grindavik (63°50'N 22°27'W)	125	Fucus vesiculosus	0.034 (0.40)	0.6 (230)	8 (200)	16 (50)	4.3 (250)		0.00 (110)	9.7 (70)	2.3 (80)			0.19
June 21	Skalanen (65°30'N 22°30'W)	126	Fucus vesiculosus	0.020 (0.50)	0.0 (170)	32 (115)	70 (40)	10.2 (130)	0.0 (250)	1.07 (70)	30 (30)	3.2 (40)	0.130 (30)	0.10	0.22
June 22	Naifadalur (66°07'N 23°06'W)	130	Fucus vesiculosus	0.037 (0.40)	0.6 (340)		19 (100)			0.63 (150)	12 (60)	1.0 (90)	0.079 (130)		0.20
- " -	- " -	131	Fucus distichus	0.050 (0.40)			22 (100)			0.06 (140)	11 (80)	2.0 (80)	0.020 (70)		0.16
- " -	- " -	132	Ascophyllium nodosum	0.023 (0.60)		8 (250)	15 (110)			0.42 (220)	3 (200)	1.5 (80)	0.010 (100)		0.27
- " -	Arnarnes (66°07'N 23°06'W)	134	Fucus vesiculosus	0.024 (0.40)	0.40 (170)	0.0 (30)	15 (10)	2.7 (240)		0.45 (150)	5.2 (60)	1.7 (70)	0.055 (80)		0.25
June 24	Siglufjörður (66°09'N 18°55'W)	140	Fucus distichus	0.036 (0.40)	0.5 (330)	17 (110)	23 (50)	5 (200)		0.73 (140)	10.4 (40)	1.7 (80)			0.17
- " -	Siglufjörður pier (66°09'N 18°55'W)	141	Fucus distichus	0.042 (0.40)	0.6 (330)	21 (140)	41 (50)	7.0 (130)		0.70 (130)	31 (30)	1.4 (100)	0.052 (80)		0.11
June 25	Raufarhöfn (66°20'N 15°55'W)	142	Fucus distichus	0.020 (10)		8.5 (40)	11 (30)			0.50 (130)	3.0 (100)	1.5 (70)	0.025 (150)		0.20
- " -	- " -	144	Fucus vesiculosus	0.030 (0.40)	0.5 (340)	5.7 (160)	14 (50)	3 (200)		0.77 (130)	10.2 (40)	1.5 (80)			0.19
- " -	- " -	145	Fucus distichus	0.046 (0.40)			12 (160)			0.65 (170)	0.7 (100)	1.7 (100)	0.024 (90)		0.17
- " -	- " -	146	Laminaria digitata	0.044 (0.30)			16 (120)			0.81 (120)	6.3 (130)	1.6 (100)	0.0064 (110)		0.17
- " -	- " -	147	Laminaria saccharina	0.044 (0.40)			11 (100)			0.54 (210)	8.0 (90)	1.7 (100)	0.0051 (130)		0.17
June 28	Neskaupstaður (65°10'N 13°43'W)	150	Fucus distichus	0.041 (0.40)	0.0 (220)	20 (90)	50 (30)	9.7 (130)		1.20 (90)	32 (30)	2.5 (110)			0.15
- " -	- " -	151	Fucus vesiculosus	0.034 (0.40)	0.6 (240)	15 (140)	31 (50)	6.0 (170)		0.80 (90)	22 (40)	1.2 (120)	0.020 (60)		0.17
- " -	Seydísfjörður (65°17'N 13°50'W)	155	Fucus distichus	0.036 (0.40)	1.0 (150)	20 (90)	43 (30)	0.9 (140)	0.9 (230)	1.03 (90)	23 (40)	1.0 (80)			0.12
- " -	- " -	156	Fucus vesiculosus	0.023 (10)	1.0 (110)	21 (20)	30 (10)	9.5 (90)	0.9 (270)	0.04 (130)	30 (20)	1.7 (80)			0.15

The relative S.D. due to counting is shown in brackets.

**Fig. C.3. Cesium-137 and Plutonium-239,240 in algae collected around Iceland in June 1981.**

Mussels were collected at four locations in Iceland and one in the Faroes (Table C.4). The mean ratio between the concentrations of  $^{54}\text{Mn}$ ,  $^{95}\text{Nb}$ ,  $^{137}\text{Cs}$ ,  $^{144}\text{Ce}$ , and  $^{239,240}\text{Pu}$  in fucoids and *Mytilus* was 2.5 on a dry weight basis. Similar ratios have been found in Danish samples<sup>2)</sup>. Ruthenium-106 did not follow the same pattern as the above-mentioned nuclides. The mean concentration of  $^{106}\text{Ru}$  was thus nearly the same in *Mytilus* as in *Fucus*. The local variation for *Mytilus* was similar to that of *Fucus*, i.e. the eastern location showed the highest concentrations (except for Pu).

Table C.3. Concentration ratios (CR) in Icelandic fucoids

		$^{137}\text{Cs}$	$^{239,240}\text{Pu}$
Sea water	Bq $\text{l}^{-1}$ : $\bar{x}$ :	$3.05 \cdot 10^{-3}$	$0.012 \cdot 10^{-3}$
	SD:	$0.19 \cdot 10^{-3}$	$0.003 \cdot 10^{-3}$
	N:	5	6
Fucoids	Bq $\text{kg}^{-1}$ dry w.: $\bar{x}$ :	0.79	0.055
	SD:	0.21	0.037
	N:	16	10
CR $\pm$ 1 S.D.		260 $\pm$ 70	4600 $\pm$ 3300

A single moss sample were collected at Isafjordur (Table C.5). The sample consisted of 19.1 kg dry matter and covered an area of 5 m<sup>2</sup>. The accumulated  $^{137}\text{Cs}$  fallout in the moss was 2188 Bq m<sup>-2</sup> or 59 mCi  $^{137}\text{Cs}$  km<sup>-2</sup>. It is interesting to note that radio-nuclides such as  $^{60}\text{Co}$  and  $^{207}\text{Bi}$  occur. In order to determine the inhomogeneity of the sample we measured five aliquots ranging from 169 to 699 g. The mean deposition in these 5 samples was 2490 Bq  $^{137}\text{Cs}$  m<sup>-2</sup> (1 SD: 550 Bq m<sup>-2</sup>). The masses of four aliquots of 94 g dry matter each were analysed for  $^{239,240}\text{Pu}$ ,  $^{238}\text{Pu}$ , and  $^{241}\text{Am}$ ; the mean values  $\pm$ 1 S.D. were:  $9.87 \pm 0.20$  Bq  $\text{kg}^{-1}$ ,  $0.35 \pm 0.03$  Bq  $\text{kg}^{-1}$  and  $2.34 \pm 0.17$  Bq  $\text{kg}^{-1}$ , respectively. In accumulated fallout in 1981, we thus found the  $^{238}\text{Pu}/^{239,240}\text{Pu}$  ratio as  $0.035 \pm 0.003$  and the  $^{241}\text{Am}/^{239,240}\text{Pu}$  ratio, to  $0.24 \pm 0.002$ .

**Table C.4. Radionuclides in *Mytilus edulis* collected at the Faroe Islands and Iceland in June 1981. (Unit: Bq kg<sup>-1</sup> dry matter)**

Date	Location (position)	Station No.	<sup>7</sup> Be	<sup>40</sup> K kg K kg <sup>-1</sup>	<sup>54</sup> Mn	<sup>95</sup> Sr	<sup>95</sup> Nb	<sup>106</sup> Ru	<sup>125</sup> Sb	<sup>137</sup> Cs	<sup>144</sup> Ce	<sup>239,240</sup> Pu	dry weight fresh weight
June 14	Rayvík 62°02'N 6°45'W	P12	8 (200)	0.015 (0.00)	0.3 (350)	1.8 (200)	3.0 (80)	5.3 (100)		0.34 (300)	3.0 (120)	0.026 (90)	0.17
June 22	Arnanæs 66°07'N 23°06'W	I33	7 (210)	0.015 (0.00)		3.1 (100)	6.3 (40)	5.4 (210)		0.27 (330)	4.4 (100)	0.020 (100)	0.15
June 24	Siglufjörður 66°09'N 18°55'W	I39	17 (90)	0.016 (0.70)	0.4 (200)	3.5 (70)	6.3 (30)	10.2 (90)	0.5 (100)	0.33 (270)	6.4 (80)	0.023 (120)	0.15
June 25	Roulachófn 66°20'N 15°55'W	I43	6 (260)	0.016 (0.00)		1.4 (100)	3.3 (60)	4.0 (210)		0.39 (240)	4.3 (100)	0.015 (150)	0.17
June 28	Neskaupstaður 65°10'N 13°43'W	I52	13 (120)	0.016 (0.70)	0.4 (260)	3.6 (70)	7.2 (30)	15 (70)		0.40 (240)	6.0 (60)	0.012 (170)	0.17

The relative S.D. due to counting is shown in brackets.

Table C.5. Radionuclides in moss collected June 22, 1981 at 66°07'N, 23°06'W (Isafjörður) Iceland (I 29)

Nuclide	Bq kg <sup>-1</sup> fresh w.	Bq m <sup>-2</sup>	rel. S.D.
<sup>7</sup> Be	339	2900	10%
<sup>54</sup> Mn	0.57	5.1	3%
<sup>60</sup> Co	0.19	1.7	3%
<sup>95</sup> Zr	11.1	100	4%
<sup>95</sup> Nb	23	210	2%
<sup>106</sup> Ru	10	90	5%
<sup>125</sup> Nb	4.6	41	2%
<sup>137</sup> Cs	242	2190	0.1%
<sup>144</sup> Ce	23	210	2%
<sup>155</sup> Eu	3.3	30	2%
<sup>207</sup> Bi	0.31	2.8	3%
<sup>226</sup> Ra	0.17	1.6	15%
<sup>238</sup> Pu	0.35	3.2	8%
<sup>239,240</sup> Pu	9.87	89	2%
<sup>241</sup> Am	2.34	21	7%

rel. S.D. determined  
from analysis  
of four aliquots

0.37 g K kg<sup>-1</sup> fresh weight



**Sales distributors:**  
**Jul. Gjellerup, Sølvgade 87,**  
**DK-1307 Copenhagen K, Denmark**

**Available on exchange from:**  
**Risø Library, Risø National Laboratory,**  
**P.O.Box 49, DK-4000 Roskilde, Denmark**

**ISBN 87-550-0856-9**  
**ISSN 0106-2840**