



Sea level reconstruction from satellite altimetry and tide gauge data

Svendsen, Peter Limkilde; Andersen, Ole Baltazar; Nielsen, Allan Aasbjerg

Published in:
Geophysical Research Abstracts

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Svendsen, P. L., Andersen, O. B., & Nielsen, A. A. (2012). Sea level reconstruction from satellite altimetry and tide gauge data. *Geophysical Research Abstracts*, 14, EGU2012-2746-2.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Sea level reconstruction from satellite altimetry and tide gauge data

P. L. Svendsen, O. B. Andersen, and A. A. Nielsen

DTU Space, Technical University of Denmark, Kgs. Lyngby, Denmark

Ocean satellite altimetry has provided global sets of sea level data for the last two decades, allowing determination of spatial patterns in global sea level. For reconstructions going back further than this period, tide gauge data can be used as a proxy. We examine different methods of combining satellite altimetry and tide gauge data using optimal weighting of tide gauge data, linear regression and EOFs, including automatic quality checks of the tide gauge time series. We attempt to augment the model using various proxies such as climate indices like the NAO and PDO, and investigate alternative transformations such as maximum autocorrelation factors (MAF), which better take into account the spatio-temporal structure of the variation.

Rather than trying to maximize the amount of variance explained, the MAF transform considers noise to be uncorrelated with a spatially or temporally shifted version of itself, whereas the desired signal will exhibit autocorrelation. This will be applied to a global dataset, necessitating wrap-around consideration of spatial shifts.

Our focus is a timescale going back approximately 50 years, allowing reasonable global availability of tide gauge data. This allows for better sensitivity analysis with respect to spatial distribution, and tide gauge data are available around the Arctic Ocean, which may be important for a later high-latitude reconstruction.