Flow distortion at a dense forest edge

Dellwik, Ebba

Publication date:
2012

[Link back to DTU Orbit](#)

Citation (APA):
Dellwik, E. (2012). *Flow distortion at a dense forest edge*. Abstract from 12th EMS Annual Meeting and 9th European Conference on Applied Climatology (ECAC), ód, Poland.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Flow distortion at a dense forest edge

Results from a forest edge experiment with two masts and one horizontally pointed wind lidar are presented. The experiment was performed at a dense beech forest edge of the Tromnæs forest. This forest is a 24m tall mature beech forest on the island Falster, Denmark. The topography at the site is flat. The masts were placed approximately 1.5 canopy heights upwind and downwind of the edge and are two canopy heights tall. We present data showing how the forest edge distorts the flow when the flow is perpendicular to the edge and towards the forest during near-neutral atmospheric stratification. Despite that the wind gradient above the canopy is similar before and after the edge, the momentum flux is strongly reduced above the canopy. This is contrary to the results by standard Reynolds' averaged Navier Stokes models that predict an overshoot of the momentum flux. We explore theoretical frameworks that can explain this reduction of the momentum flux as well as the change of other terms in the covariance matrix. Another noteworthy result from the analysis was that the porosity of the canopy is wind-speed dependent. As a result, several aspects of the wind field at the edge depend on the mean wind speed.

The results are relevant for understanding the on-set and growth of the internal boundary layer of forested areas.