



Timing of the Little Ice Age in southern Greenland

Kjær, Kurt H.; Kjeldsen, Kristian K. ; Bjørk, Anders A.; Khan, Shfaqat Abbas; Korsgaard, Niels J.; Funder, Svend ; Larsen, Nicolaj K.; Vinther, Bo; Andresen, Camilla S.; Long, Anthony J.

Total number of authors:
13

Published in:
Geophysical Research Abstracts

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Kjær, K. H., Kjeldsen, K. K., Bjørk, A. A., Khan, S. A., Korsgaard, N. J., Funder, S., Larsen, N. K., Vinther, B., Andresen, C. S., Long, A. J., Woodroffe, S. A., Hansen, E. S., & Olsen, J. (2013). Timing of the Little Ice Age in southern Greenland. *Geophysical Research Abstracts*, 15, EGU2013-10410.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Timing of the Little Ice Age in southern Greenland

Kurt H. Kjær (1), Kristian K. Kjeldsen (1), Anders A. Bjørk (1), Shfaqat A. Khan (2), Niels J. Korsgaard (1), Svend Funder (1), Nicolaj K. Larsen (3), Bo Vinther (4), Camilla S. Andresen (5), Antony J. Long (6), Sarah A. Woodroffe (6), Eric Steen Hansen (7), and Jesper Olsen (8)

(1) Natural History Museum of Denmark, University of Copenhagen, Centre for GeoGenetics, Denmark (kkjeldsen@snm.ku.dk), (2) DTU Space, National Space Institute, Technical University of Denmark, (3) Department of Geoscience, Aarhus University, (4) Centre for Ice and Climate, The Niels Bohr Institute, University of Denmark, (5) Geological Survey of Denmark and Greenland, (6) Department of Geography, Durham University, UK, (7) Natural History Museum of Denmark, University of Copenhagen, Denmark, (8) Department of Physics and Astronomy

Northern hemisphere temperatures reached their Holocene minimum and most glaciers reached their maximum during The Little Ice Age (LIA), but the timing of specific cold intervals is site-specific. In southern Greenland, we have compiled data from organic matter incorporated in LIA sediments, used as a signal for ice-free terrain being overridden by LIA glacier advances, and data from threshold lakes showing the onset of glacier-fed lakes, thus revealing the advance-maximum phase initiating the LIA. Finally, we have compiled lichenometry results indicating the onset of bedrock vegetation succeeding ice retreat. Our results show that the advance of glaciers during the LIA occurs early after the Medieval Warm Period terminating soon after 1200 AD and culminates c. 1500-1600 AD. Historical maps also show that many glaciers on the western coast occupy a still-stand near the LIA maximum until 1900 AD before retreat commence.

Thus in southern Greenland, we define LIA as the period between the first signs of Late Holocene glacier readvance and the latest onset of retreat – i.e. from ca. 1200 to c. 1900. During this period northern hemisphere annual mean temperatures, although fluctuating, were generally below the 1961-1990 average, with the coldest interval between c. 1600 and 1800. Even though winter temperatures may have dominated the cooling, also the summer temperatures which are most closely correlated with glacier mass balances, dropped, to c. 0.6° below the average in the northern hemisphere including the Arctic. Furthermore, the glacier response seems to be mirrored by a oceanic cooling between 500-1000 AD, followed by onset of the LIA at 1150-1250 AD as seen in the relative strength of warm subsurface water and the influence of the East Greenland Current.