



On the electron to proton mass ratio and the proton structure

Trinhammer, Ole L.

Published in:
EPL

Link to article, DOI:
[10.1209/0295-5075/102/42002](https://doi.org/10.1209/0295-5075/102/42002)

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Trinhammer, O. L. (2013). On the electron to proton mass ratio and the proton structure. *EPL*, 102(4), 42002. <https://doi.org/10.1209/0295-5075/102/42002>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

On the electron to proton mass ratio and the proton structure

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2013 EPL 102 42002

(<http://iopscience.iop.org/0295-5075/102/4/42002>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 192.38.67.112

The article was downloaded on 11/06/2013 at 12:43

Please note that [terms and conditions apply](#).

On the electron to proton mass ratio and the proton structure

OLE L. TRINHAMMER

*Department of Physics, Technical University of Denmark - Fysikvej, Building 307,
DK-2800 Kongens Lyngby, Denmark, EU*

received 27 March 2013; accepted in final form 13 May 2013
published online 7 June 2013

PACS 21.60.Fw – Models based on group theory
PACS 14.20.Gk – Baryon resonances ($S=C=B=0$)
PACS 14.20.Dh – Protons and neutrons

Abstract – We derive an expression for the electron to nucleon mass ratio from a reinterpreted lattice gauge theory Hamiltonian to describe interior baryon dynamics. We use the classical electron radius as our fundamental length scale. Based on expansions on trigonometric Slater determinants for a neutral state, a specific numerical result is found to be less than three percent off the experimental value for the neutron. Via the exterior derivative on the Lie group configuration space $u(3)$ we derive approximate parameter-free parton distribution functions that compare rather well with those for the u and d valence quarks of the proton.

Copyright © EPLA, 2013

The mass ratio and the model. – The ratio we get between the electron mass m_e and the proton mass m_p is

$$\frac{m_e}{m_p} = \frac{\alpha}{\pi} \frac{1}{E}, \quad (1)$$

where $\alpha = e^2/(\hbar 4\pi\epsilon_0)$ is the fine-structure constant [1] and $E = E/\Lambda$ is the dimensionless ground-state eigenvalue of a reinterpreted lattice gauge theory Kogut-Susskind Hamiltonian [2],

$$\frac{\hbar c}{a} \left[-\frac{1}{2}\Delta + \frac{1}{2}\text{Tr}\chi^2 \right] \Psi(u) = E\Psi(u). \quad (2)$$

with Manton's action [3] used now as a potential for a configuration variable $u = e^{i\chi}$ in the Lie group $u(3)$ instead of a link variable U in the $SU(3)$ algebra [4]. The energy scale $\Lambda \equiv \hbar c/a$ corresponds to a fundamental length scale a , which we shall relate to the classical electron radius R_0 by

$$a\pi = R_0, \quad (3)$$

where R_0 is determined by the electron self-potential energy [5]

$$\frac{1}{4\pi\epsilon_0} \frac{e^2}{R_0} = m_e c^2. \quad (4)$$

We assume (2) to describe the baryon spectrum and identify the ground state with the proton. With $E = m_p c^2 = E\Lambda = E\hbar c/a$ and (4) applied in (3), eq. (1) follows directly. Our configuration space is “orthogonal” to the laboratory

space wherefore (2) describes a truly interior dynamics which may be projected on laboratory space parameters through the eigenangles θ_j parametrizing the eigenvalues $e^{i\theta_j}$ of u . The projection introduces the dimensionful scale a , thus

$$x_j = a\theta_j. \quad (5)$$

Now a shortest geodesic [6] to track along the full extension of the $u(3)$ maximal torus runs from the origin at $u = I$, where all eigenvalues are $1 = e^{i0}$, to $u = -I$, where all eigenvalues are $-1 = e^{i\pi}$, see also fig. 1. When, for instance, the neutron decays to the proton—and the electron is created as a “peel off”—the topological change in the interior baryon state maps by projection to laboratory space. It is not a new idea to suggest the classical electron radius as a fundamental length in elementary particle physics [7,8]. Here we specify the introduction of the scale (3) via the projection (5). Conjugate to the space (angle) parameters in (5) are canonical momentum (action) operators

$$p_j = -i\hbar \frac{1}{a} \frac{\partial}{\partial \theta_j} = \frac{\hbar}{a} T_j. \quad (6)$$

The toroidal generators T_j induce coordinate fields ∂_j according to the above-mentioned eigenangle parametrization of the $u(3)$ torus. In general the nine generators T_k of $u(3)$ namely induce coordinate fields as follows:

$$\partial_k = \frac{\partial}{\partial \theta} u e^{i\theta T_k} |_{\theta=0} = u i T_k. \quad (7)$$




Fig. 1: Projection from configuration space to laboratory space.

The remaining six off-toroidal generators are important in the baryon spectroscopy phenomenology resulting from (2) since they take care of spin and flavour degrees of freedom. With these parametrizations the Laplacian ($\hbar = 1$) in a polar decomposition reads [9,10]

$$\Delta = \sum_{j=1}^3 \frac{1}{J^2} \frac{\partial}{\partial \theta_j} J^2 \frac{\partial}{\partial \theta_j} - \sum_{\substack{i < j \\ k \neq i, j}}^3 \frac{K_k^2 + M_k^2}{8 \sin^2 \frac{1}{2}(\theta_i - \theta_j)}. \quad (8)$$

Here the Van de Monde determinant, the ‘‘Jacobian’’ of the parametrization is [11]

$$J = \prod_{i < j}^3 2 \sin \left(\frac{1}{2}(\theta_i - \theta_j) \right). \quad (9)$$

The operators K_k commute as body fixed angular-momentum operators and M_k ‘‘connect’’ the algebra by commuting into the subspace of K_k

$$[M_k, M_l] = [K_k, K_l] = -iK_m, \quad (10)$$

cyclic in k, l, m . The presence of the components of $\mathbf{K} = (K_1, K_2, K_3)$ and $\mathbf{M} = (M_1, M_2, M_3)$ in the Laplacian opens for the inclusion of spin and flavour. Interpreting \mathbf{K} as the interior spin operator is encouraged by the body fixed signature of the commutation relations. The relation between laboratory space and interior space is like the relation in nuclear physics between fixed coordinate systems and intrinsic body fixed coordinate systems for the description of rotational degrees of freedom. To derive

the spectrum for M^2 we use a coordinate representation [12],

$$\begin{aligned} K_1 &= a\theta_2 p_3 - a\theta_3 p_2 = \hbar \lambda_7, \\ K_2 &= a\theta_1 p_3 - a\theta_3 p_1 = \hbar \lambda_5, \\ K_3 &= a\theta_1 p_2 - a\theta_2 p_1 = \hbar \lambda_2 \end{aligned} \quad (11)$$

and

$$\begin{aligned} M_3/\hbar &= \theta_1 \theta_2 + \frac{a^2}{\hbar^2} p_1 p_2 = \lambda_1, \\ M_2/\hbar &= \theta_3 \theta_1 + \frac{a^2}{\hbar^2} p_3 p_1 = \lambda_4, \\ M_1/\hbar &= \theta_2 \theta_3 + \frac{a^2}{\hbar^2} p_2 p_3 = \lambda_6. \end{aligned} \quad (12)$$

The lambdas are corresponding Gell-Mann generators [13]. From these and

$$\begin{aligned} Y/\hbar &= \frac{1}{6}(\theta_1^2 + \theta_2^2 - 2\theta_3^2) + \frac{1}{6} \frac{a^2}{\hbar^2} (p_1^2 + p_2^2 - 2p_3^2) = \lambda_8/\sqrt{3}, \\ 2I_3/\hbar &= \frac{1}{2}(\theta_1^2 - \theta_2^2) + \frac{1}{2} \frac{a^2}{\hbar^2} (p_1^2 - p_2^2) = \lambda_3 \end{aligned} \quad (13)$$

we find by straightforward but tedious commutations the spectrum

$$M^2 = \frac{4}{3} \left(n + \frac{3}{2} \right)^2 - K(K+1) - 3 - \frac{1}{3}y^2 - 4i_3^2, \quad n = 0, 1, 2, 3, \dots, \quad (14)$$

where y and i_3 are hypercharge and isospin three-component quantum numbers. The minimum value for the positive definite M^2 is 13/4 in the case of spin 1/2, hypercharge 1 and isospin 1/2 as for the nucleon. To solve for the eigenvalues we factorize the wave function

$$\Psi(u) = \tau(\theta_1, \theta_2, \theta_3) \Upsilon(\alpha_4, \alpha_5, \alpha_6, \alpha_7, \alpha_8, \alpha_9), \quad (15)$$

insert it in (2) and then integrate over the off-toroidal degrees of freedom ($\alpha_4, \alpha_5, \alpha_6, \alpha_7, \alpha_8, \alpha_9$) to get for the measure scaled wave function $R = J\tau(\theta_1, \theta_2, \theta_3)$, for toroidal degrees of freedom

$$\left[- \sum_{j=1}^3 \frac{\partial^2}{\partial \theta_j^2} + V \right] R = 2ER. \quad (16)$$

Here

$$\begin{aligned} V &= -2 + \frac{1}{3}(K(K+1) + M^2) \sum_{i < j}^3 \frac{1}{8 \sin^2 \frac{1}{2}(\theta_i - \theta_j)} \\ &\quad + 2(v(\theta_1) + v(\theta_2) + v(\theta_3)), \end{aligned} \quad (17)$$

where (see fig. 2)

$$\begin{aligned} v(\theta) &= \frac{1}{2}(\theta - n\pi)^2, \quad \theta \in [(2n-1)\pi, (2n+1)\pi], \\ &\text{for } n \in \mathbb{Z}. \end{aligned} \quad (18)$$




Fig. 2: Periodic parametric potential (18) originating from the potential in (2). The dashed curve corresponds to the Wilson analogue [14] of the Manton action and is not considered in the present work.

By expansion on Slater determinants [10]

$$b_{pqr} = \epsilon_{ijk} \cos(p\theta_i) \sin(q\theta_j) \cos(r\theta_k) \quad (19)$$

with integer p, q, r , we can solve (16) by the Rayleigh-Ritz method [15] to yield the ground-state eigenvalue $E_0 = 4.38$ which corresponds to $m_e/m_0 = 1/1885$. This is less than three percent off the value $m_e/m_n = 1/1838.6\dots$ based on experimental data for the electron and neutron [16]. Note that b_{pqr} is antisymmetric in the colour degrees of freedom θ_j .

Parton distribution functions. – We can generate parton distribution functions by projections via the momentum form, *i.e.*, the exterior derivative on the $u(3)$ manifold. For this we expand the exterior derivative dR of the measure scaled toroidal wave function R on the torus forms $d\theta_j$,

$$dR = \psi_j d\theta_j. \quad (20)$$

The action of the torus forms on the toroidal coordinate fields expresses the generalization to the interior configuration space manifold of the quantization inherent in the conjugate variables in (5) and (6), thus

$$d\theta_i(\partial_j) = \delta_{ij} \Leftrightarrow [\partial_j, \theta_i] = \delta_{ij}. \quad (21)$$

Inspired by Bettini's elegant treatment of parton scattering [17], we generate distribution functions via our exterior derivative. The derivation runs like this (with $\hbar = c = 1$): Imagine a proton at rest with four-momentum $P = (\mathbf{0}, E_0)$. We boost it virtually to energy E by impacting upon it a massless four-momentum $q = (\mathbf{q}, E - E_0)$ which we assume to hit a parton xP . After impact the parton represents a virtual mass xE . Thus,

$$(xP_\mu + q_\mu) \cdot (xP^\mu + q^\mu) = x^2 E^2, \quad (22)$$

from which we get the parton momentum fraction

$$x = \frac{2E_0}{E + E_0}, \quad (23)$$

or the boost parameter

$$\xi \equiv \frac{E - E_0}{E} = \frac{2 - 2x}{2 - x}. \quad (24)$$




Fig. 3: The generators T_u and T_d given in (28) generate traces along the $u(3)$ torus as shown here in the first and third toroidal degrees of freedom with T_3 pointing to the left.

We can use the boost parameter for an angular track $\theta = \pi\xi$ on the manifold in the direction laid out by a specific toroidal generator

$$T = a_1 T_1 + a_2 T_2 + a_3 T_3. \quad (25)$$

With the toroidal generator T as introtangling momentum operator we namely have the qualitative correspondence $q_\mu \sim E - E_0 \sim (1-x)E \sim (1-x)T$. That is, we will project along $\xi T \sim (1-x)T$ in order to probe on xP_μ . With a probability amplitude interpretation of R we project on a fixed colour base iT_j and sum over the colour components for a specific generator T to get the corresponding distribution function $f_T(x)$ determined by

$$f_T(x)dx = \left(\sum_{j=1}^3 dR_{u=\exp(\theta iT)}(iT_j) \right)^2 d\theta. \quad (26)$$

By a pull-back operation [18] to parameter space we get

$$\begin{aligned} \sum_{j=1}^3 dR_{u=\exp(\theta iT)}(iT_j) &= dR_{u=\exp(\theta iT)}(i(T_1 + T_2 + T_3)) \\ &= \frac{d}{dt} R(ue^{ti(T_1+T_2+T_3)}) \Big|_{t=0} \\ &= \frac{d}{dt} R(a_1\theta + t, a_2\theta + t, a_3\theta + t) \Big|_{t=0} \\ &= \sum_{j=1}^3 \frac{\partial R}{\partial \theta_j} \Big|_{(\theta_1, \theta_2, \theta_3) = (\theta \cdot a_1, \theta \cdot a_2, \theta \cdot a_3)} \cdot \frac{\partial(a_j\theta + t)}{\partial t} \Big|_{t=0} \\ &= \left(\frac{\partial R}{\partial \theta_1} + \frac{\partial R}{\partial \theta_2} + \frac{\partial R}{\partial \theta_3} \right) \Big|_{(\theta_1, \theta_2, \theta_3) = (\theta \cdot a_1, \theta \cdot a_2, \theta \cdot a_3)} \\ &\equiv D(\theta \cdot a_1, \theta \cdot a_2, \theta \cdot a_3). \end{aligned} \quad (27)$$

Along the tracks shown in fig. 3, we generate distributions by

$$T_u = \begin{Bmatrix} 2/3 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{Bmatrix} \quad \text{and} \quad T_d = \begin{Bmatrix} -1/3 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{Bmatrix} \quad (28)$$

as shown in fig. 4 for a first-order approximation

$$b_{0, \frac{1}{2}, 1}(\theta_1, \theta_2, \theta_3) = \frac{1}{N} \begin{vmatrix} 1 & 1 & 1 \\ \sin \frac{1}{2}\theta_1 & \sin \frac{1}{2}\theta_2 & \sin \frac{1}{2}\theta_3 \\ \cos \theta_1 & \cos \theta_2 & \cos \theta_3 \end{vmatrix}, \quad (29)$$




Fig. 4: Parton distribution functions from a first-order approximation to the ground state of (2) as a function of the parton fraction x . The distributions are generated by impact along $T_u = (2/3, 0, -1)$ (solid, green line) and $T_d = (-1/3, 0, -1)$ (dashed, red line) with no fitting parameters. For comparison we show u and d valence quarks of the proton at 10 GeV^2 in the insert adapted from the Particle Data Group with other parton distributions erased [16].

to an expansion for a protonic state with normalization constant N . For instance,

$$xf_{T_u}(x) = x \left[D \left(\pi \frac{2-2x}{2-x} \cdot \frac{2}{3}, \pi \frac{2-2x}{2-x} \cdot 0, \pi \frac{2-2x}{2-x} \cdot (-1) \right) \right]^2 \cdot \frac{\pi \cdot 2}{(2-x)^2}, \quad (30)$$

where in its full glory

$$\begin{aligned} ND(\theta_1, \theta_2, \theta_3) = & \\ & -\frac{1}{2} \cos \frac{\theta_1}{2} \cdot (\cos \theta_3 - \cos \theta_2) - \sin \theta_1 \cdot \left(\sin \frac{\theta_3}{2} - \sin \frac{\theta_2}{2} \right) \\ & + \frac{1}{2} \cos \frac{\theta_2}{2} \cdot (\cos \theta_3 - \cos \theta_1) + \sin \theta_2 \cdot \left(\sin \frac{\theta_3}{2} - \sin \frac{\theta_1}{2} \right) \\ & - \frac{1}{2} \cos \frac{\theta_3}{2} \cdot (\cos \theta_2 - \cos \theta_1) - \sin \theta_3 \cdot \left(\sin \frac{\theta_2}{2} - \sin \frac{\theta_1}{2} \right). \end{aligned} \quad (31)$$

The distribution functions shown in fig. 4 contain no fitting parameters at all. Note that the T_u -parton momentum content of the approximate state (29) is close to the double of the T_d content,

$$\int_0^1 xf_{T_u}(x) dx \Big/ \int_0^1 xf_{T_d}(x) dx = 0.2722 / 0.1437 = 1.89 \approx 2. \quad (32)$$

Work is in progress to solve (2) for the charged proton by antisymmetrized parity definite expansions on Bloch states [19]

$$g_{pqr} = e^{i\kappa \cdot \theta} e^{ip\theta_1} e^{iq\theta_2} e^{ir\theta_3}, \quad (33)$$

where $\kappa = (\kappa_1, \kappa_2, \kappa_3)$ are appropriately chosen Bloch vectors and $\theta = (\theta_1, \theta_2, \theta_3)$.

Comments. – It is not clear whether the minor but still significant discrepancy in our value for m_e/m_n is due to disparate partial wave projections of the base set (19) or due to corrections needed in (3). It is somewhat surprising that when applied to a neutral state, expression (1) gives a result just three percent off the experimental value for the electron to neutron value. This is surprising since the classical electron radius on which we base our prediction is normally presented to be just an order of magnitude scale for strong interaction phenomena [20]. It should be noted though that in (5) we have introduced a well-defined projection to space parameters. However, at the present level of understanding it seems wise to quote Landau and Lifshitz: “...it is impossible within the framework of classical electrodynamics to pose the question whether the total mass of the electron is electrodynamic.” [5].

Conclusion. – We have derived an expression for the electron to nucleon mass ratio from a reinterpreted lattice gauge theory Hamiltonian. A specific calculation for the neutron from expansions on Slater determinants of indefinite spin-parity gives a result less than three percent off the experimental value. The proximity of prediction to observation should encourage further study within the framework of the Hamiltonian model presented. From the same model we have derived approximate parameter-free parton distribution functions that compare rather well with those for the u and d valence quarks of the proton. Work is in progress to establish a complete Bloch wave base for expansions with suitable symmetry requirements implied by the potential and the antisymmetry under interchange of colour degrees of freedom.

I am grateful to the late VICTOR F. WEISSKOPF for inspiration on m_e/m_p and to HOLGER BECH NIELSEN for clarifying discussions on the momentum form. I thank TORBEN AMTRUP for support through the years, JAKOB BOHR for advice and ERIK BOTH for help in preparing the manuscript for publication.

REFERENCES

- [1] SAKURAI J. J., *Advanced Quantum Mechanics* (Addison-Wesley, United Kingdom) 1967/1984, p. 12.
- [2] KOGUT J. B. and SUSSKIND L., *Phys. Rev. D*, **11** (1975) 395.
- [3] MANTON N. S., *Phys. Lett. B*, **96** (1980) 328.
- [4] TRINHAMMER O., *Phys. Lett. B*, **129** (1983) 234.

- [5] LANDAU L. D. and LIFSHITZ E. M., *The Classical Theory of Fields, Course of Theoretical Physics*, Vol. **2**, 4th edition (Elsevier Butterworth-Heinemann, Oxford) 2005, p. 97.
- [6] MILNOR J., *Ann. Math. Stud.*, **51** (1963) 1.
- [7] HEISENBERG W., *Ann. Phys. (Leipzig)*, **32** (1938) 20.
- [8] MEHRA J. and RECHENBERG H., *The Historical Development of Quantum Theory - The Conceptual Completion and the Extensions of Quantum Mechanics 1932-1941*, Vol. **6**, Part 2 (Springer, New York) 2001, p. 956.
- [9] TRINHAMMER O. L. and OLAFSSON G., *The Full Laplace-Beltrami operator on $U(N)$ and $SU(N)$* , arXiv:math-ph/9901002v2 (1999/2012).
- [10] WADIA S. R., *Phys. Lett. B*, **93** (1980) 403.
- [11] WEYL H., *The Classical Groups - Their Invariants and Representations*, 2nd edition (Princeton University Press) 1997, p. 197.
- [12] SCHIFF L. I., *Quantum Mechanics*, 3rd edition (McGraw-Hill) 1968, p. 211.
- [13] See ref. [12], p. 209.
- [14] WILSON K. G., *Phys. Rev. D*, **10** (1974) 2445.
- [15] BRUUN NIELSEN H., Technical University of Denmark (private communication, 1997).
- [16] PARTICLE DATA GROUP (BERINGER J. *et al.*), *Phys. Rev. D*, **86** (2012) 010001.
- [17] BETTINI A., *Introduction to Elementary Particle Physics* (Cambridge University Press, UK) 2008, p. 204.
- [18] GUILLEMIN W. and POLLACK A., *Differential Topology* (Prentice-Hall, NJ, USA) 1974, p. 163.
- [19] JACOBSEN KARSTEN W., Technical University of Denmark (private communication, June 2012).
- [20] See ref. [7], p. 20.