



Iridoid and Phenylethanoid Glucosides from *Veronica* (sect. *Hebe*) *lavaudiana*

Taskova, Rilka Mladenova; Kokubun, Tetsuo; Ryan, Ken G.; Garnock-Jones, Phil J.; Jensen, Søren Rosendal

Published in:
Journal of Natural Products

Link to article, DOI:
[10.1021/np200233p](https://doi.org/10.1021/np200233p)

Publication date:
2011

Document Version
Early version, also known as pre-print

[Link back to DTU Orbit](#)

Citation (APA):
Taskova, R. M., Kokubun, T., Ryan, K. G., Garnock-Jones, P. J., & Jensen, S. R. (2011). Iridoid and Phenylethanoid Glucosides from *Veronica* (sect. *Hebe*) *lavaudiana*. *Journal of Natural Products*, 74(6), 1477-1483. <https://doi.org/10.1021/np200233p>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Iridoid and Phenylethanoid Glucosides from *Veronica* *lavaudiana*

Rilka M. Taskova,[†] Tetsuo Kokubun,[‡] Ken G. Ryan,[†] Phil J. Garnock-Jones,[†] and Søren R. Jensen^{*, §}

School of Biological Sciences, Victoria University of Wellington, PO Box 600, Wellington 6140, New Zealand; Jodrell Laboratory, Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3DS, UK; and Department of Chemistry, Technical University of Denmark, DK-2800, Lyngby, Denmark.

* To whom correspondence should be addressed. Tel.: +45 45252103. Fax: +45 45933968. E-mail: srj@kemi.dtu.dk.

[†] Victoria University of Wellington

[‡] Royal Botanic Gardens, Kew

[§] Technical University of Denmark

From an extract of *Veronica* (sect. *Hebe*) *lavaudiana* we have identified mannitol and isolated 11 iridoid glucosides, the carbohydrate ester hebitol II, and four phenylethanoid glycoside esters. Five of the iridoid glycosides are new; of these, lavaudioside A, B, and C (**2a**, **3a**, and **4**) are 1-mannityl esters of 8-epiloganic acid while **7e** and **7f** are 6'-*O*-caffeoyl derivatives of catalpol. The new phenylethanoid glycoside esters, helioside A, B, and C (**8b-8d**), are 6'-xylosyl derivatives of aragoside. The structures of the new compounds were elucidated mainly by spectroscopic analysis, but also by chemical degradation. We also demonstrated that the structures of the known glycosides globularitol as well as hebitols I and II should be revised. These compounds are derivatives of mannitol and not glucitol as previously believed.

Veronica (sect. *Hebe*) *lavaudiana* Raoul is an endemic subshrub that grows on basalt outcrops and cliffs above 150 m altitude on Banks Peninsula on the east coast of the South Island of New Zealand.¹ The peninsula, reaching an altitude of about 900 m, is formed from two extinct volcanoes and has at times in the past been separated as an island. *V. lavaudiana* and its nearest relatives in the sect. *Hebe* belong in the sun hebe clade.² It has been formerly treated as the genus *Heliohebe*¹ or as part of *Hebe*³ or *Parahebe*,⁴ all of which are now included in a broad and monophyletic concept of *Veronica*.⁵ Little chemical work has been undertaken on *V. lavaudiana*. Thus, this species was included in an early chemotaxonomic survey of iridoids and flavonoids in *Veronica* and related genera,⁶ and in a recent LC/MS investigation on the flavonoid chemistry of the five species of *Heliohebe*.⁷

Fresh plant material was blended with cold MeOH and the water-soluble part of the extract was subjected to a series of chromatographic procedures. Seventeen compounds were isolated and identified including the hexitol, mannitol (**1**), the carbohydrate ester hebitol II (**5**), 11 iridoid glucosides (**2a**, **3a**, **4**, **6**, **7-7f**), and four phenylethanoid glycoside esters (**8**, **8b-8d**). Of these, the five iridoids **2a**, **3a**, **4**, **7e**, **7f** and the three phenylethanoids **8b-8d** are new.

The composition of the sugar fraction was deduced by interpretation of the ¹³C NMR data only. Structural elucidations of the new compounds were carried out by spectroscopic methods (1D and 2D NMR, MS) and chemical degradation. Despite intensive purification, most of the compounds contained minor concentrations of *cis*-caffeoyl or -feruloyl isomers. Compound **8d**, contained 8% of the *cis*-form, in accord with our previous experience that feruloyl esters readily isomerize.

Lavaudioside A (**2a**) was obtained as an amorphous solid $[\alpha]_D^{20}$ -30. The molecular formula was C₃₁H₄₂O₁₈, as deduced from the ¹³C NMR data and the quasimolecular ion obtained by LC-HRESIMS (observed *m/z* 720.2713 [M + NH₄]⁺). The ¹H NMR spectrum (Table 1) in methanol-*d*₄ showed five

resonances corresponding to the presence of a caffeoyl group (δ_{H} 7.51 to 6.24). An additional resonance at δ_{H} 7.53 (s, H-3), as well as two more at δ_{H} 5.55 (d, H-1) and δ_{H} 4.68 (d, H-1') suggested that **2a** was a carboxylated iridoid glucoside. Furthermore, the three-proton doublet at δ_{H} 1.10 showed the presence of a C-10 methyl group. In the ^{13}C NMR spectrum, 30 resonances were observed of which one (δ_{C} 169.0) was of double intensity. Nine of the resonances were consistent with the presence of a caffeoyl group. Of the remaining resonances in the spectrum, 16 were similar to those reported for a 7-*O*-acyl derivative of the iridoid glucoside epiloganic acid (**2**),⁸ including those from the 1-*O*- β -glucopyranosyl group. The remaining six resonances were all in the δ_{C} 65-73 region, which suggested the presence of a hexityl moiety. This was consistent with the ^1H NMR spectrum, where the signals not accounted for were observed in the shift interval δ_{H} 3.65-4.44. The HMBC spectrum allowed the carbons of attachment between the iridoid moiety and the peripheral parts to be discerned. Thus, the position of the ester group was confirmed by an HMBC correlation between H-7 (δ_{H} 4.9) and the carbonyl carbon of the caffeoyl group (δ_{C} 169.0). The 1'''-CH₂-resonances of the hexityl moiety at δ_{H} 4.44 and 4.21 showed correlations with the carbonyl carbon (δ_{C} 169.0) of the iridoid core and to a carbinol signal (δ_{C} 70.5), which could therefore be assigned as C-2'''. The HSQC spectrum designated the remaining resonances at δ_{C} 65.1, 72.8, 70.9 and 70.8, as one CH₂OH and three CHOH groups, respectively, thereby fulfilling the molecular formula determined by the MS. The first three shift values closely matched those of mannitol (**1**). Confirmation that the hexitol was indeed mannitol was achieved by an in-NMR tube hydrolysis of **2a** with ammonia in D₂O. After 48 h at room temperature, ^{13}C NMR resonances corresponding to mannitol were observed at δ_{C} 64.5 (C-1 and 6), 72.1 (C-2 and 5), and 70.5 (C-3 and 4).⁹ These increased in intensity following the addition of authentic mannitol (see Supporting Information).

Lavaudioside B (**3a**) was obtained as an amorphous solid [α]_D²² -24. The molecular formula was C₃₁H₄₀O₁₈, as deduced by the ^{13}C NMR data and LC-HRESIMS (observed *m/z* 718.2558 [M + NH₄]⁺). The NMR data (Table 1) in methanol-*d*₄ was assigned as above. The ^1H NMR spectrum was similar to

that of **2a**, including a caffeoyl and a mannityl group, as well as the resonance at δ_{H} 7.62 (1H, s, H-3) and others at δ_{H} 5.41 and 4.70, assigned to H-1 and H-1', respectively. The major differences were in the iridoid moiety. Thus, additional resonances (δ_{H} 5.48 and 5.45) were present while those from a methyl group were missing. In the ^{13}C NMR spectrum of **3a**, the expected 31 resonances were observed. When compared with the spectrum of **2a**, only the resonances for the iridoid aglucone differed. The two resonances at δ_{C} 148.9 and 116.5 suggested an iridoid with an 8,10-double bond, in accordance with the ^1H NMR data. A comparison with the data reported⁸ for a 7-*O*-acyl derivative of gardoside (**3**) showed a satisfactory coincidence for the chemical shift values.

Lavaudioside C (**4**) was isolated as an amorphous solid $[\alpha]_{\text{D}}^{22}$ -48, and the molecular formula was $\text{C}_{46}\text{H}_{58}\text{O}_{26}$, deduced by the ^{13}C NMR data and LC-HRESIMS (observed m/z 1027.3269 $[\text{M} + \text{H}]^+$). The ^1H NMR spectrum (Table 1) was similar to that of **2a**, but resonances from two caffeoyl groups were present. Also, two β -hexopyranosyl groups were considered present due to the two doublets at δ_{H} 4.69 and 4.36. In the ^{13}C NMR spectrum (Table 1) of **4**, 39 resonances were observed, of which five were of double and one (δ_{C} 71.6) was of triple intensity, as also confirmed by the HSQC data. Eighteen of the resonances could be assigned to the two caffeoyl groups and comparing with the data for **2a**, a further 16 could be assigned to an epiloganlyl moiety including its 1-*O*-glucosyl group. This left 12 ^{13}C NMR resonances to be accounted for. That at δ_{C} 105.0 was assigned to an anomeric carbon atom and the remaining 11, resonating at δ_{C} 64-78, suggested the presence of a hexopyranosyl and a hexityl moiety. 2D NMR data suggested that the hexopyranosyl moiety was an additional β -glucopyranosyl group but the full structure of **4** could only be resolved in relation to that of hebitol II (**5**).

The sugar ester hebitol II (**5**), was considered to be composed of a 6-*O*-caffeoyl- β -D-glucopyranose moiety with a 1-glucityl group as the aglucone.¹⁰ The structure of **5** was originally elucidated by comparison of the NMR data of an analogue, the feruloyl ester globularitol (**5b**) isolated from

Globularia orientalis.¹¹ The structure of **5b** was in turn inferred by deacylation and comparison by TLC of the resulting disaccharide with authentic β -D-glucopyranosyl-(1 \rightarrow 6)-glucitol (**5d**), obtained by NaBH₄ reduction of gentiobiose. However, because the corresponding ¹³C NMR shift values of **5** (for C-1''' to C-3''' in Table 1) compared better with those of β -D-glucopyranosyl-(1 \rightarrow 6)-mannitol (**5c**)¹² than with those of the glucitol derivative, we therefore decided to re-examine the structure of **5**. Thus, treatment of **5** with ammonia in D₂O gave a disaccharide with a ¹³C NMR spectrum (see Supporting Information) identical to that reported for **5c**. The similar levorotatory activity of **5** and **5a-5c** pointed to the β -D-form of the glucose moiety in the four compounds (See Supporting Information). Consequently, the structures of the three analogues hebitol II (**5**), hebitol I (**5a**),¹⁰ and globularitol (**5b**)¹¹ must be revised to the β -D-glucosylmannitol derivatives shown in the formula chart.

Returning to the structure of **4**, the resonances of the caffeoyl and β -glucopyranosyl groups found in **4**, but not in **2a**, were coincident with the corresponding resonances of **5**. In addition, those from C-1''' through C-3''' for **4** were almost coincident with the resonances of the corresponding carbon atoms of **2a**, and the resonances of C-4''' to C-6''' were coincident with those of **5**. This provided the hypothetical structure for **4**. In the HMBC spectrum, correlations could be seen between the 1'''-CH₂ group (δ_{H} 4.42 and 4.20) and C-2''' (δ_{C} 70.7) as well as C-11 of the iridoid core (δ_{C} 169.0) since a cross-peak with H-5 (δ_{H} 3.13) was also found at this shift value. Other correlations were seen between the 6'''-CH₂ group (δ_{H} 4.14 and 3.76) and C-5''' (δ_{C} 71.6) as well as the anomeric carbon atom C-1''' of the internal β -glucopyranosyl moiety (δ_{C} 105.0). Also, H-5''' (δ_{H} 3.55) displayed a correlation with C-1''', and furthermore, cross-peaks could be found between 6'''-CH₂ (δ_{H} 4.22 and 4.28) and C-5''' (δ_{C} 75.5) as well as one of the low field carbonyl carbon atoms (δ_{C} 169.1). Therefore, it was concluded that the latter could be assigned to one of the caffeic acid moieties. The structure of lavaudioside C has therefore been established as that shown by **4**.

Compound **7e** was obtained as an amorphous solid $[\alpha]_D^{22} -70$, and the molecular formula was $C_{24}H_{28}O_{13}$, determined by the ^{13}C NMR data and LC-HRESIMS (observed m/z 523.1465 $[M - H]^-$). The 1H NMR data (Table 2) were somewhat similar to those of verminoside (**7b**)¹³ both containing a caffeoyl and a catalpol moiety. However, in **7e** H-6 (δ_H 3.78) resonated 1.2 ppm upfield while the H-6' signals (δ_H 4.41 and 4.49) were 0.6 and 1.1 ppm downfield compared to those of **7b**. This indicated that in **7e** the caffeoyl moiety was located at the 6'-*O*-position. The conclusion was supported by the relative changes observed in the ^{13}C NMR data. In the HMBC spectrum, correlations between 6'-CH₂ (δ_H 4.49 and 4.41) and the carbonyl carbon atom of the caffeoyl group (δ_C 169.0) confirmed that the structure of **7e** was 6'-*O*-caffeoylcatalpol.

Compound **7f** was also an amorphous solid $[\alpha]_D^{20} -20$, with the molecular formula $C_{33}H_{34}O_{16}$, deduced by the ^{13}C NMR data and LC-HRESIMS (observed m/z 709.1791 $[M + Na]^+$). The 1H NMR data (Table 2) showed that the molecule contained one catalpol and two caffeoyl units. Comparison with the NMR data of **7b** and **7e** showed that in **7f**, the resonances for the catalpol aglucone moiety were very similar to those of the former, while the resonances of the β -D-glucopyranosyl unit were similar to those of the latter. This suggested that the two caffeoyl groups were placed at the 6-*O*- and the 6'-*O*-positions of catalpol. Analysis of the HMBC spectrum confirmed this. Thus, the positions of the ester groups were confirmed by correlations between H-6 (δ_H 4.89) as well as by one of the 6'-CH₂ protons (δ_H 4.47), and the carbonyl carbon atoms of the caffeoyl groups (δ_C 169.0 and 168.9), respectively. The structure of compound **7f** was therefore 6,6'-di-*O*-caffeoylcatalpol.

Helioside A (**8b**), $[\alpha]_D^{20} -23$, had a molecular formula $C_{39}H_{52}O_{24}$, as deduced by the ^{13}C NMR data and LC-HRESIMS (observed m/z 922.3200 $[M+NH_4]^+$). The NMR data (Table 3) were assigned by 2D NMR spectroscopy and by comparison with the model compounds aragoside (**8**)¹⁴ and its 6'- β -glucopyranosyl derivative chionoside D (**8a**) from *V. thomsonii* (Buchanan) Cheeseman.¹⁵ The ^{13}C NMR spectrum showed 37 resonances of which two (δ_C 115.3 and 77.8) were of double intensity. This

included four signals at δ_C 103-106 arising from anomeric carbon atoms showing correlations with protons in the δ_H 4.2-4.7 region (HSQC). The NMR data (Table 2) were almost superimposable with those of **8a**, a 3,4-dihydroxyphenylethyl β -D-glucopyranoside esterified with a caffeoyl group at C-4' and substituted with an α -arabinopyranosyl at C-2' and β -glucopyranosyl groups at C-3' and C-6'. However, one of the peripheral glucosyl groups of **8a** was replaced by a pentosyl moiety in **8b**, identified as a β -xylopyranosyl group by the NMR data. The correlations in the HMBC spectrum confirmed the sites of attachment of the peripheral units. Thus, a cross-peak between H-1' (δ_H 4.51) of the central glucosyl moiety and C-8 (δ_C 72.2) proved the position of the aglycone. A correlation between H-4' (δ_H 4.96) and the carbonyl carbon atom of the caffeoyl ester (δ_C 168.6) showed the position of the acyl moiety. Cross-peaks between H-1''' (δ_H 4.52) of the arabinosyl group and C-2' (δ_C 82.5), as well as between H-1'' (δ_H 4.63) of the peripheral glucosyl group and C-3' (δ_C 81.2), demonstrated the positions of these sugar moieties. Finally, a correlation between H-1'''' of the xylosyl moiety (δ_H 4.24) and the 6'-CH₂-group (δ_C 69.1), proved the structure as shown by **8b**.

Helioside B (**8c**), $[\alpha]_D^{20}$ -25, had a molecular formula C₄₈H₅₈O₂₇ as deduced by the ¹³C NMR data and LC-HRESIMS (observed m/z 1084.3451 [M+NH₄]⁺). The NMR data (Table 3) were assigned as above. When the NMR and MS data were compared with those of **8b** it became apparent that **8c** was a caffeoyl ester of the former. Beside the additional caffeoyl group, the 6''-CH₂ resonances (δ_H 4.34, 4.27) of the peripheral glucopyranosyl group were found 0.6 and 0.8 ppm, respectively, downfield in the ¹H NMR spectrum. Similarly, in the ¹³C NMR spectrum of **8c**, C-6'' (δ_C 64.7) and C-5'' (75.1) of the peripheral β -glucopyranosyl group were found 1.6 ppm downfield and 2.7 ppm upfield, respectively, when compared to the data for **8b**. Therefore, the additional caffeoyl group was positioned at the C-6'' oxygen atom in helioside B. The correlations in the HMBC spectrum were consistent with this, including the cross-peak between one of the C-6'' protons (δ_H 4.27) and the ester carbonyl carbon atom assigned to the additional caffeoyl group (δ_C 169.1). Helioside B is therefore the 6''-caffeoyl ester of helioside A.

Helioside C (**8d**), $[\alpha]_D^{20}$ -22, had the molecular formula $C_{40}H_{54}O_{24}$, deduced by the ^{13}C NMR data and LC-HRESIMS (observed m/z 936.3333 $[M+NH_4]^+$). The NMR data (Table 3) were coincident with those of **8b**, except for the resonances of the aroyl moiety and for the presence of a methoxy group in **8d**. The chemical shift values (δ_H 3.90; δ_C 56.5), as well as a cross-peak between the *O*-methyl protons (δ_H 3.90) and C-3'''' (δ_C 149.4) were consistent with the presence of a feruloyl substituent in **8d**. Consequently, helioside C is the feruloyl analogue of helioside A.

Our study confirms the previous report⁶ of aucubin (**6**), catalpol (**7**), and the two catalpol esters **7a** and **7b** in *V. laevigata*. Epiloganic acid (**2**) and gardoside (**3**), and their methyl esters are often found in plants containing **6** and **7** and have been reported from several *Veronica* species.^{10, 16-20} However, iridoid 11-carboxylic acids esterified with alditols like **2a**, **3a**, and **4**, found in the present study, have not been reported before. Hebitol II (**5**), isolated in the present work, adds to the variety of sugar esters found in this *Veronica* sect. *Hebe*.^{10, 18-21}

Experimental Section

General Experimental Procedures. Optical rotations were measured on a Perkin-Elmer 241 polarimeter. One-dimensional 1H , ^{13}C NMR and 2D DQF-COSY, gHSQC, gHMBC and NOESY NMR spectra were recorded on a Varian Unity Inova 500 MHz spectrometer in methanol- d_4 and the chemical shifts are given as δ values with reference to the solvent peaks (δ_H 3.30 or δ_C 49.0, respectively). LC-HRESIMS was performed on an Agilent HP 1100 HPLC equipped with a BDS-C₁₈ reversed-phase column running a H₂O-MeCN (50 ppm TFA in H₂O) gradient. The LC was coupled to a LCT of a TOF MS (Micromass, Manchester, UK) operated in the positive electrospray ion mode using 5-leucineenkephalin as lock mass. UV spectra were recorded on a Shimadzu UV-1601 instrument. Preparative HPLC was performed on an Agilent 1100 Series LC System (Agilent, Santa Clara, CA, USA) with a guarded Luna C₁₈ column (10 × 250 mm, 5 μ m, Phenomenex) kept at 40 °C and a Waters

system (Watford, UK) with Genesis C₁₈ column (10 × 250 mm, 5 μm, Jones Chromatography, Mid Glamorgan, UK) at 30 °C, with MeOH-H₂O mixtures as eluents at a flow rate of 4 mL/min. The known compounds were identified by NMR and compared with published data: mannitol (**1**);⁹ epiloganic acid (**2**), gardoside (**3**), aucubin (**6**) and catalpol (**7**);²² verproside (**7a**);²³ verminoside (**7b**) and minecoside (**7d**);¹³ 6-*O*-feruloylcatalpol (**7c**).²⁴

Plant Material. *Veronica lavaudiana* Raoul was collected from plants cultivated in the grounds of Landcare Research, Lincoln, New Zealand. A voucher specimen has been deposited at the Allan Herbarium, Landcare Research (CHR).

Extraction and Isolation. Plant material (40 g, fresh wt.) was blended with MeOH and filtered. The concentrated extract (14 g) was partitioned between Et₂O-H₂O and the aqueous phase loaded on a 4 × 45 cm Diaion HP-20 (Supelco, Bellefonte, US) column and eluted with H₂O-MeOH mixtures. The fraction eluted with H₂O (5.5 g dry wt.) contained mainly mannitol (**1**). The fraction eluted with 10% MeOH (780 mg) contained mainly aucubin (**6**) and catalpol (**7**).

The fraction eluted with 30% MeOH (250 mg) was separated on silica gel column eluted with CHCl₃-MeOH-H₂O (60:15:4, 60:22:4) followed by HPLC (Luna C₁₈, linear gradient from 25 to 40% MeOH over 10 min) to give hebitol II (**5**, 4.2 mg).

The fraction eluted with 50% MeOH (2.2 g) was separated on a Sephadex LH-20 (Amersham Biosciences, Uppsala, Sweden; 3 × 30 cm) column eluted with 90% MeOH to give: Fr. A (550 mg), further separated by HPLC (Luna C₁₈, 3 to 50% MeOH over 10 min) to give helioside A (**8b**, 191 mg) and helioside C (**8d**, 25 mg). The final purification of these was performed by HPLC on Genesis C₁₈ with 30% MeOH for **8b** (135 mg) and 35% MeOH for **8d** (4.1 mg). Fr. B (492 mg), further separated by HPLC (Luna C₁₈, 30 to 60% MeOH over 10 min) to give aragoside (**8**, 63.4 mg); lavaudioside B (**3a**,

12.6 mg); helioside B (**8c**, 24.0 mg); lavaudioside A (**2a**, 41.6 mg); and 6'-*O*-caffeoylcatalpol (**7e**, 2.7 mg), all additionally purified by HPLC (Genesis C₁₈, 35% MeOH). Fr. C (545 mg), further separated by HPLC (Luna C₁₈, 30 to 50% MeOH over 10 min) to give 6'-*O*-caffeoylcatalpol (**7e**, 3.3 mg) and verminoside (**7b**, 268 mg).

The fraction eluted with 60% MeOH (3.9 g) was separated on a Sephadex LH-20 column (3 × 40 cm) eluted with 90% MeOH to give: Fr. D (215 mg), further separated by HPLC (Luna C₁₈, 35 to 45% MeOH over 15 min) to give aragoside (**8**, 11.0 mg), lavaudioside B (**3a**, 23.1 mg), verminoside (**7b**, 39.1 mg) and a fraction (35.6 mg), purified by HPLC (Genesis C₁₈, 42% MeOH) to give lavaudioside A (**2a**, 17.1 mg). Fr. E (1.0 g), further separated on silica gel column (CHCl₃-MeOH-H₂O, 60:22:4) to give verminoside (**7b**, 645 mg).

The fraction eluted with MeOH (1.8 g) was separated on Sephadex LH-20 column (2.5 × 50 cm) eluted with MeOH: Fr. F (866 mg) was further separated by HPLC (Luna C₁₈, 45% MeOH over 10 min) to give verminoside (**7b**, 202 mg), 6-*O*-feruloylcatalpol (**7c**, 71.5 mg), minecoside (**7d**, 31 mg), and a fraction (50.5 mg), separated on silica gel (CHCl₃-MeOH-H₂O, 60:22:4) and additionally purified by HPLC (Genesis C₁₈ with 45% MeOH) to give lavaudioside C (**4**, 2.0 mg). Fr. G (589 mg) was further separated by HPLC (Luna C₁₈, 40-50% MeOH over 20 min) to give verminoside (**7b**, 311 mg). Fr. H (131 mg) was separated by HPLC on Luna C₁₈ (35-60 % MeOH over 20 min) and additionally purified on Genesis C₁₈ (48% MeOH) to give 6,6'-di-*O*-caffeoylcatalpol (**7f**, 1.4 mg).

Lavaudioside A (2a). Colorless amorphous solid: $[\alpha]_D^{20}$ -30 (*c* 0.4; MeOH); UV (MeOH) λ_{\max} (log ϵ) 330 (4.24), 294 (4.10, sh), 239 (4.30); ¹H and ¹³C NMR, Table 1; LC-HR ESIMS *m/z* 720.2713 [M + NH₄]⁺ (calcd for C₃₁H₄₆NO₁₈, 720.2715).

Lavaudioside B (3a). Colorless amorphous solid: $[\alpha]_D^{22}$ -24 (*c* 0.5; MeOH); UV (MeOH) λ_{\max} (log ϵ) 330 (4.16), 293 (4.02, sh), 236 (4.22); ^1H and ^{13}C NMR, Table 1; LC-HR ESIMS m/z 718.2558 $[\text{M} + \text{NH}_4]^+$ (calcd for $\text{C}_{31}\text{H}_{44}\text{NO}_{18}$, 718.2558).

Lavaudioside C (4). Colorless amorphous solid: $[\alpha]_D^{22}$ -48 (*c* 0.1; MeOH); UV (MeOH) λ_{\max} (log ϵ) 329 (4.26), 297 (4.14, sh), 240 (4.25); ^1H and ^{13}C NMR, Table 1; LC-HR ESIMS m/z 1027.3269 $[\text{M} + \text{H}]^+$ (calcd for $\text{C}_{46}\text{H}_{59}\text{O}_{26}$, 1027.3289).

6'-*O*-Caffeoylcatalpol (7e). Colorless amorphous solid: $[\alpha]_D^{22}$ -70 (*c* 0.2; MeOH); UV (MeOH) λ_{\max} (log ϵ) 332 (4.12), 300 (3.99, sh), 246 (3.89); ^1H and ^{13}C NMR, Table 2; LC-HR ESIMS m/z 523.1465 $[\text{M} - \text{H}]^-$ (calcd for $\text{C}_{24}\text{H}_{27}\text{O}_{13}$, 523.1452).

6,6'-*Di-O*-caffeoylcatalpol (7f). Colorless amorphous solid: $[\alpha]_D^{20}$ -20 (*c* 0.2; MeOH); UV (MeOH) λ_{\max} (log ϵ) 331 (4.43), 289 (4.33, sh), 246 (4.21); ^1H and ^{13}C NMR, Table 2; LC-HR ESIMS m/z 709.1791 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{33}\text{H}_{34}\text{NaO}_{16}$, 709.1741).

Helioside A (8b). Colorless amorphous solid: $[\alpha]_D^{20}$ -23 (*c* 0.5; MeOH); UV (MeOH) λ_{\max} (log ϵ) 332 (4.21), 292 (4.08), 249 (3.94, sh); ^1H and ^{13}C NMR, Table 3; LC-HR ESIMS m/z 922.3200 $[\text{M} + \text{NH}_4]^+$ (calcd for $\text{C}_{39}\text{H}_{56}\text{NO}_{24}$, 922.3175).

Helioside B (8c). Colorless amorphous solid: $[\alpha]_D^{20}$ -25 (*c* 0.1; MeOH); UV (MeOH) λ_{\max} (log ϵ) 331 (4.14), 290 (4.00, sh), 247 (3.93, sh); ^1H and ^{13}C NMR, Table 3; LC-HR ESIMS m/z 1084.3451 $[\text{M} + \text{NH}_4]^+$ (calcd for $\text{C}_{48}\text{H}_{62}\text{NO}_{27}$, 1084.3490).

Helioside C (8d). Colorless amorphous solid: $[\alpha]_D^{20}$ -22 (*c* 0.4; MeOH); UV (MeOH) λ_{\max} (log ϵ) 328 (4.25), 293 (4.08), 233 (4.09, sh); ^1H and ^{13}C NMR, Table 3; LC-HR ESIMS m/z 936.3333 $[\text{M} + \text{NH}_4]^+$ (calcd for $\text{C}_{40}\text{H}_{58}\text{NO}_{24}$, 936.3331).

Treatment of 2a with Ammonium Hydroxide. Lavaudioside A (**2a**) (17 mg) was dissolved in D_2O (0.7 mL) in an NMR tube (5 mm) and 25 % ammonium hydroxide (70 μL) was added. After 48 h at room temperature the ^{13}C NMR spectrum was recorded (D_2O , 75 MHz): δ_{C} 64.5 (C-1 and 6), 72.1 (C-2 and 4), 70.5 (C-3 and 5); after addition of excess mannitol, the three relevant resonances increased in size (see Supporting Info.).

Treatment of 5 with Ammonium Hydroxide. Hebitol II (**5**) (13 mg) was dissolved in D_2O (0.7 mL) in an NMR tube (5 mm) and 25 % ammonium hydroxide (70 μL) was added. After 72 h at room temperature the ^{13}C NMR spectrum was recorded (D_2O , 75 MHz): δ_{C} 64.5 (C-1), 72.1 (C-2), 70.3 (C-3), 70.9 (C-4), 70.9 (C-5), 73.1 (C-6), 104.2 (C-1'), 74.5 (C-2'), 76.9 (C-3'), 70.4 (C-4'), 77.3 (C-5'), 62.0 (C-6') (see Supporting Info.) coinciding with that reported for **5c**.¹²

Acknowledgement. We thank Dr. K. F. Nielsen, BioCentrum, DTU, DK, for providing high-resolution ESI-MS data. The School of Chemical and Physical Sciences and the School of Psychology at VUW provided access to some laboratory equipment. The research was supported by grants from Victoria University of Wellington's University Research Fund and from the New Zealand Foundation for Research Science and Technology through the OBI "Defining New Zealand's Land Biota". We thank Melissa Hutchison for permission to use her photograph of *V. lavaudiana*.


Supporting Information Available: NMR spectra (^1H , ^{13}C) of lavaudiosides A-C (**2a**, **3a**, and **4**), the catalpol esters (**7d** and **7e**), heliosides A-C (**8b-8d**), and hydrolysis products of **2a** and **5** are available free of charge via the Internet at <http://pubs.acs.org>.

References and Notes


- (1) Garnock-Jones, P.J. *New Zealand J. Bot.* **1993**, *31*, 323–339.
- (2) Albach, D. C.; Meudt, H. M. *Mol. Phylogenet. Evol.* **2010**, *54*, 457–471.
- (3) Allan, H.H. *Flora of New Zealand*, vol. 1. Government Printer, Wellington, 1961.
- (4) Heads, M.J. *Bot. J. Linn. Soc.* **1994**, *115*, 65–89.
- (5) Garnock-Jones, P.; Albach, D.; Briggs, B. G. *Taxon* **2007**, *56*, 571–582.
- (6) Grayer-Barkmeijer, R. J. *Chemosystematic Investigations in Veronica L. (Scrophulariaceae) and Related Genera*; PhD Thesis, Univ. of Leiden, The Netherlands; **1979**.
- (7) Taskova, R. M.; Kokubun, T.; Grayer, R. J.; Ryan, K. G.; Garnock-Jones, P. J. *Biochem. Syst. Ecol.* **2008**, *36*, 110–116.
- (8) Lahloub, M.-F.; Zagloul, M. G.; Afifi, M. S.; Sticher, O. *Phytochemistry* **1992**, *33*, 401–405.
- (9) Bock, K.; Pedersen, C. *Adv. Carbohydr. Chem. Biochem.* **1983**, *41*, 27–66.

- (10) Pedersen, P.; Gotfredsen C. H.; Wagstaff S. J.; Jensen S. R. *Biochem. Syst. Ecol.* **2007**, *35*, 777–784 .
- (11) Calis, I.; Kirmizibekmez, H.; Tasdemir, D.; Sticher, O.; Ireland, C. M. *Z. Naturforsch.* **2002**, *57c*, 591–596.
- (12) Usov, A. I.; Chizhov, A. O. *Izv. Akad. Nauk. Ser. Khim.* **1993**, 1817–1820.
- (13) Sticher, O.; Afifi-Yazar, F. U. *Helv. Chim. Acta* **1979**, *62*, 535–539.
- (14) Rønsted, N.; Bello, M. A.; Jensen S. R. *Phytochemistry* **2003**, *64*, 529–533.
- (15) Taskova, R. M.; Kokubun, T.; Ryan, K. G.; Garnock-Jones, P. J.; Jensen, S. R. *Chem. Pharm. Bull.* **2010**, *58*, 703–711.
- (16) Jensen, S. R.; Albach, D. C.; Ohno, T., Grayer, R. J. *Biochem. Syst. Ecol.* **2005**, *33*, 1031–1047.
- (17) Taskova, R. M.; Gotfredsen, C. H.; Jensen S. R. *Phytochemistry* **2006**, *67*, 286–301.
- (18) Johansen, M.; Larsen, T. S.; Matthebjerg, M. A.; Gotfredsen, C. H.; Jensen S. R. *Biochem. Syst. Ecol.* **2007**, *35*, 614–620.
- (19) Jensen, S. R.; Gotfredsen, C. H.; Grayer, R. J. *Biochem. Syst. Ecol.* **2008**, *36*, 207–215.
- (20) Maggi, A.; Taskova, R.M.; Gotfredsen, C.H.; Bianco, A.; Jensen, S.R. *Biochem. Syst. Ecol.* **2009**, *37*, 731–736.
- (21) Taskova, R. M.; Kokubun, T.; Ryan, K. G.; Garnock-Jones, P. J.; Jensen, S. R. *Chem. Pharm. Bull.* **2010**, *58*, 703–711.
- (22) Rønsted, N.; Göbel, E.; Franzyk, H.; Jensen, S. R.; Olsen, C. E. *Phytochemistry* **2000**, *55*, 337–348.
- (23) Afifi-Yazar, F. U.; Sticher, O. *Helv. Chim. Acta* **1980**, *63*, 1905–1907.


(24) Stuppner, H.; Wagner, H. *Planta Med.* **1989**, *55*, 467–469.




1 mannitol




2 R''=R'''=H; epiloganic acid
2a R''=caffeoyl; R'''=1-mannityl;
lavaudioside A




3 R''=R'''=H; gardoside
3a R''=caffeoyl; R'''=1-mannityl;
lavaudioside B





6 aucubin




4 lavaudioside C




5 R=caffeoyl; hebitol II
5a R= benzoyl; hebitol I
5b R=feruloyl; globularitol



5c R=OH; R¹=H
5d R=H; R¹=OH



7 R=R⁶=H; catalpol
7a R=3,4-di-OH-benzoyl; R⁶=H; verproside
7b R=caffeoyl; R⁶=H; verminoside
7c R=feruloyl; R⁶=H
7d R=isoferuloyl; R⁶=H; minecoside
7e R=H; R⁶=caffeoyl
7f R=R⁶=caffeoyl



8 R=caffeoyl; R¹=R²=H; aragoside
8a R=caffeoyl; R¹=H; R²=Glc; chionoside D
8b R=caffeoyl; R¹=H; R²=Xyl; helioside A
8c R=R¹=caffeoyl; R²=Xyl; helioside B
8d R=feruoyl; R¹=H; R²=Xyl; helioside C

SYNOPSIS TOC




Table 1. ¹H (500 MHz) and ¹³C NMR (125 MHz) Data of Lavaudiosides A-C and Model Compounds in Methanol-*d*₄ (or D₂O).

	Lavaudioside A (2a)		Lavaudioside B (3a)		Lavaudioside C (4)		Hebitol II (5)		glu-mannitol (5c)
position	δ _C	δ _H (<i>J</i> in Hz)	δ _C	δ _H (<i>J</i> in Hz)	δ _C	δ _H (<i>J</i> in Hz)	δ _C	δ _H (<i>J</i> in Hz)	δ _C (D ₂ O) ^a
Irid. agluc									
1	95.9	5.55, d (4.3)	96.3	5.41, d (5.6)	96.0	5.54, d (4.5)			
3	152.9	7.53, s	154.2	7.62, <i>s</i>	152.9	7.52, <i>s</i>			
4	113.7		111.2		113.6				
5	31.4	3.14, br q (8)	33.2	3.28, obsc	31.5	3.13, br q (8)			
6	39.0	2.27, ddd (14, 8, 3) 2.00, dt (14, 7)	38.6	2.23, dt (13, 6) 2.12, dt (13, 6)	39.0	2.26, ddd (14, 8, 3) 1.99, dt (14, 6.5)			
7	82.6	4.90, obsc	76.4	5.58, br t (5)	82.6	4.90, obsc			
8	43.2	2.44, m	148.9		43.2	2.43, dq (14, 8)			
9	43.1	2.62, dt (4.3, 8)	44.8	3.00, br t (6)	43.1	2.60, dt (8, 4.5)			
10	14.3	1.10, d (7.3)	116.5	5.48, 5.45, br s's	14.3	1.09, d (7.4)			
11	169.0		169.0*		169.0				
Irid. Glc									
1'	99.7	4.68, d (7.9)	99.9	4.70, d (7.9)	99.7	4.69, d (7.9)			
2'	74.7	3.20, dd (9.2, 7.9)	74.8	3.23, dd (9.2, 7.9)	74.8	3.20, dd (9.2, 7.9)			
3'	77.9	3.38, t (9.1)	78.0	3.38, t (9.0)	78.0	3.37, obsc			
4'	71.6	3.25, t (9.1)	71.7	3.26, obsc	71.6	3.24, obsc			
5'	78.3	3.32, obsc	78.5	3.32, obsc	78.4	3.32, obsc			
6'	62.8	3.92, obsc 3.66, obsc	62.9	3.90, obsc 3.64, obsc	62.9	3.90, dd (11.9, 1.7) 3.64, dd (11.9, 6.4)			
Caffeoyl									
CO"	169.0		168.9*		169.0*/169.1*		169.1		
α"	115.3	6.24, d (15.9)	115.3	6.26, d (15.9)	115.3/114.8	6.24/6.29, d's (15.9)	114.8	6.29, d (15.9)	
β"	146.9	7.51, d (15.9)	147.1	7.53, d (15.9)	147.3/146.9	7.51/7.57, d's (15.9)	146.8	7.57, d (15.9)	
1"	127.6		127.7		127.7/127.7		127.6		
2"	115.1	7.04, d (1.6)	115.1	7.05, br s	115.2/115.1	2H, 7.04, m	115.1	7.05, d (1.9)	
3"	146.8		146.8		146.8/147.3		147.3		
4"	149.5		149.7		149.6/149.6		149.6		
5"	116.5	6.77, d (8.2)	116.5	6.77, d (8.2)	116.5/116.5	2H, 6.77, d (8.2)	116.5	6.77, d (8.2)	
6"	123.0	6.94, dd (8.2, 1.6)	123.1	6.94, br d (8.2)	123.0/123.1	2H, 6.94, m	123.1	6.95, dd (8.2, 1.9)	
Hexitol									
1'''	67.4	4.44, dd (11.5, 2.1) 4.21, dd (11.5, 5.9)	67.5	4.45, dd (11.6, 2.4) 4.21, dd (11.6, 5.9)	67.4	4.42, dd (11.5, 2.3) 4.20, dd (11.5, 5.8)	65.2	3.80, obsc 3.61, dd (11.1, 6.0)	65.1
2'''	70.5	3.91, obsc	70.5	3.91, obsc	70.7	3.87, obsc	72.9	3.69, m	72.8
3'''	70.8*	3.81, obsc	70.8*	3.81, obsc	70.7*	3.80, obsc	70.9*	3.80, obsc	71.0
4'''	70.9*	3.81, obsc	71.0*	3.81, obsc	71.6*	3.80, obsc	71.1*	3.82, obsc	71.6
5'''	72.8	3.69, obsc	72.9	3.70, obsc	71.6	3.75, obsc	71.6	3.80, obsc	71.6
6'''	65.1	3.82, obsc	65.2	3.82, obsc	73.6	4.14, dd (10.5, 1.3)	73.7	4.15, dd (10.5, 2.3)	73.7

Centr. Glc	3.65, obsc	3.64, obsc	3.76, m Glc	3.72, dd (10.5, 6.1) Glc
1 ^{'''}		105.0	4.36, d (7.8)	105.0
2 ^{'''}		75.1	3.25, obsc	75.2
3 ^{'''}		77.7	3.41, t (9.2)	77.6
4 ^{'''}		71.6	3.38, obsc	71.6
5 ^{'''}		75.5	3.55, m	75.5
6 ^{'''}		64.6	4.52, dd (11.9, 1.7)	64.6
			4.28, dd (11.9, 5.8)	62.6
				4.29, dd (11.9, 5.7)

Obsc: signal obscured. * Interchangeable within column.

^a Data from ref. 12. For direct comparison, C-6^{'''} was aligned with C-6^{'''} in **5**.

Table 2. ¹H (500 MHz) and ¹³C NMR (125 MHz) Data of Catalpol Esters (**7e** and **7f**) in Methanol-*d*₄.

position	6'- <i>O</i> -Caffeoylcatalpol (7e)		6,6'-Di <i>O</i> -caffeoylcatalpol (7f)	
	δ _C	δ _H (<i>J</i> in Hz)	δ _C	δ _H (<i>J</i> in Hz)
Agluc				
1	96.3	4.86, d (9.9)	95.2	4.96, d (9.7)
3	141.8	6.31, dd (6.0, 1.5)	142.4	6.33, dd (5.9, 1.5)
4	104.1	5.02, dd (6.0, 4.5)	103.1	4.92, dd (5.9, 4.8)
5	39.1	2.22, m	36.8	2.53, m
6	79.7	3.78, br d (8.1)	81.3	4.89, br d (9.0)
7	62.4	3.40, br s	61.6	3.67, br s
8	66.1		66.7	
9	43.3	2.53, dd (9.9, 7.8)	43.0	2.61, dd (9.7, 7.8)
10	61.9	4.15, d (13.0) 3.64, d (13.0)	60.2	4.19, d (13.1) 3.70, d (13.1)
Glc				
1'	99.7	4.77, d (7.9)	99.8	4.79, d (7.9)
2'	74.7	3.29, obsc	74.8	3.30, obsc
3'	77.4	3.41, m	77.5	3.43, m
4'	71.5	3.41, m	71.7	3.39, m
5'	75.9	3.53, m	76.1	3.55, m
6'	63.9	4.49, dd (11.9, 2.0) 4.41, dd (11.9, 5.6)	63.9	4.50, dd (12.0, 2.7) 4.47, dd (12.0, 5.9)
Caffeoyl				
CO"	169.0		169.0/168.9	
α"	114.8	6.27, d (15.9)	114.8/114.8	6.30/6.28, d (15.9)
β"	146.8	7.54, d (15.9)	146.8/146.8	2H, 7.58, d (15.9)
1"	127.7		127.7/127.6	
2"	115.1	7.04, d (1.7)	115.2/115.3	2H, 7.06, br s
3"	147.3		147.4/147.7	
4"	149.7		149.7/149.8	
5"	116.5	6.77, d (8.2)	116.5/116.5	6.78/6.73, d (8.2)
6"	123.1	6.94, dd (8.2, 1.7)	123.2/123.3	6.97/6.95, dd (8, 2)

Obsc: signal obscured.

Table 3. ¹H (500 MHz) and ¹³C NMR (125 MHz) Data of Heliosides A-C (**8b-8d**) in Methanol-*d*₄.

position	Helioside A (8b)		Helioside B (8c)		Helioside C (8d)	
	δ _C	δ _H (<i>J</i> in Hz)	δ _C	δ _H (<i>J</i> in Hz)	δ _C	δ _H (<i>J</i> in Hz)
Aglucone						
1	131.9		132.0		131.9	
2	117.5	6.74, d (1.8)	117.5	6.72, d (1.9)	117.5	6.74, d (1.9)
3	145.8		146.0		146.0	
4	144.1		144.7		144.7	
5	116.4	6.67, d (8.0)	116.3	6.67, d (8.1)	116.3	6.68, d (7.9)
6	121.6	6.57, dd (8.0, 1.8)	121.5	6.56, dd (8.0, 1.9)	121.5	6.57, dd (7.9, 1.9)
7	36.6	2H, 2.77, m	36.7	2H, 2.74, m	36.7	2H, 2.77, m
8	72.2	4.06/3.68, m's	72.2	4.01/3.62, m's	72.3	4.06/3.67, m's
Central Glc						
1'	103.2	4.51, d (7.9)	103.3	4.40, d (7.7)	103.3	4.51, d (7.9)
2'	82.5	3.70, obsc	82.8	3.69, dd (9.5, 7.7)	82.8	3.69, obsc
3'	81.2	4.10, t (9.5)	81.6	3.94, t (9.5)	81.4	4.09, t (9.7)
4'	70.2	4.96, t (9.5)	70.0	4.98, t (9.5)	70.3	4.97, t (9.7)
5'	74.1	3.77, obsc	74.5	3.65, obsc	74.4	3.78, obsc
6'	69.1	3.87, dd (11.4, 1.5) 3.60, dd (11.5, 6.4)	69.1	3.81, dd (11.5, 1.5) 3.60, obsc	69.4	3.87, dd (11.5, 1.5) 3.61, dd (11.5, 6.1)
2'-Ara						
1'''	104.7	4.52, d (7.4)	104.7	4.51, d (7.3)	104.3	4.52, d (7.6)
2'''	73.1	3.56, dd (9.1, 7.4)	73.2	3.57, obsc	73.2	3.56, dd (9.0, 7.4)
3'''	73.9	3.46, obsc	74.0	3.46, obsc	74.0	3.46, obsc
4'''	69.8	3.73, obsc	69.8	3.74, obsc	69.8	3.73, obsc
5'''	67.3	3.76, obsc 3.16, br d (11.7)	67.4	3.74, obsc 3.09, obsc	67.4	3.78, obsc 3.16, obsc
3'-Glu						
1''	104.1	4.63, d (7.9)	104.5	4.67, d (7.9)	104.9	4.64, d (7.8)
2''	75.2	3.08, dd (9.2, 7.9)	75.1*	3.14, obsc	74.8	3.08, dd (9.0, 7.8)
3''	77.8*	3.28, obsc.	78.0*	3.34, obsc	78.0*	3.27, obsc
4''	72.2	3.02, t (9.3)	71.7	3.20, t (9.3)	72.0	3.00, t (9.3)
5''	77.8	3.21, obsc	75.1	3.45, obsc	78.1*	3.22, obsc
6''	63.1	3.77, obsc 3.44, obsc	64.7	4.34, dd (1.9, 11.5) 4.27, dd (6.4, 11.5)	63.2	3.77, obsc 3.41, obsc
4'-Acyl						
CO''''	168.6		168.7		168.6	
α''''	115.3	6.33, d (15.9)	115.0	6.29, d (15.9)	115.7	6.44, d (15.9)
β''''	147.3	7.56, d (15.9)	147.5	7.52, d (15.9)	147.2	7.62, d (15.9)
1''''	127.6		127.6		127.7	
2''''	115.3	7.07, d (1.9)	115.3	7.00, br s	111.6	7.24, d (1.6)
3''''	146.6		146.7		149.3	
4''''	149.5		149.6		150.6	
5''''	116.6	6.77, d (8.2)	116.6	6.77, d (8.2)	116.5	6.81, br d (8.1)
6''''	123.2	6.98, dd (8.2, 1.9)	123.1	6.90, br d (8.2)	124.3	7.10, dd (8.1, 1.6)
OMe						
6'-Xyl						
1'''''	105.1	4.24, d (7.4)	105.2	4.20, d (7.4)	105.3	4.24, d (7.4)
2'''''	74.7	3.19, obsc	75.3*	3.20, dd (9.0, 7.4)	75.3	3.19, dd (9.0, 7.4)
3'''''	77.3*	3.39, obsc	77.5*	3.30, obsc	77.5*	3.30, obsc
4'''''	70.9	3.45, obsc	71.0	3.46, obsc	71.1	3.45, obsc
5'''''	66.7	3.82, dd (11.5, 5.3) 3.14, obsc	66.8	3.81, obsc 3.13, obsc	66.9	3.82, dd (11.5, 5.4) 3.15, obsc
6''-cafffeoyl						
CO''''''			169.1			
α''''''			115.0	6.18, d (15.9)		
β''''''			147.1	7.50, d (15.9)		
1''''''			127.6			

2''''	115.5*	7.00, br s
3''''	146.6*	
4''''	149.5*	
5''''	116.5	6.69, d (8.2)
6''''	123.1	6.90, br d (8.2)

Obsc: signal obscured. * Interchangable within column.

