



## Dynamics of Catalyst Nanoparticles

Hansen, Thomas Willum; Cavalca, Filippo; Wagner, Jakob Birkedal

*Publication date:*  
2013

[Link back to DTU Orbit](#)

*Citation (APA):*

Hansen, T. W., Cavalca, F., & Wagner, J. B. (2013). *Dynamics of Catalyst Nanoparticles*. Abstract from XXIV Brazilian Congress of Microscopy and Microanalysis, Caxambu, Brazil.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Dynamics of Catalyst Nanoparticles


**Thomas W. Hansen**<sup>1</sup>, Filippo Cavalca<sup>1</sup> and Jakob B. Wagner<sup>1</sup>

<sup>1</sup>Center for Electron Nanoscopy, Technical University of Denmark, DK-2800, Lyngby,  
Denmark  
[twh@cen.dtu.dk](mailto:twh@cen.dtu.dk)

### Abstract

Transmission electron microscopy (TEM) is extensively used in catalysis research. Recent developments in aberration correction allows imaging surface structures with unprecedented resolution. Using these correctors in conjunction with environmental TEM (ETEM), where imaging of materials can be done under gas exposure, dynamic phenomena such as sintering and growth can be observed with sub-Ångström resolution.

Metal nanoparticles contain the active sites in heterogeneous catalysts, which are important for many industrial applications including the production of clean fuels, chemicals and pharmaceuticals, and the cleanup of exhaust from automobiles and stationary power plants. Sintering, or thermal deactivation, is an important mechanism for the loss of catalyst activity. In order to initiate a systematic study of the dynamics and sintering of nanoparticles, various catalytic systems have been studied in atmospheres relevant for their operation. By studying growth patterns, we found that changes in the atmospheres changes heavily effects not only the rate of sintering but also the mechanism through which it occurs. In many cases, anomalously large particles were observed indicating that particle sintering is not solely governed by the mechanisms previously proposed. These results are divided into the different phases of the catalyst lifetime.



ETEM image sequence of Au/graphene in 200 Pa H<sub>2</sub> at 104°C. The image series shows different stages of two Au particles coalescing resulting in a single Au particle. Image dimensions: 18x18 nm.