DTU Library



A Bayesian Clinical Decision Support for Early Detection of Classical Swine Fever in Individual Pigs - Evaluation of the Sensitivity and Specificity of the Model

Elbers, Armin; van der Gaag, Linda; Schmeiser, Stefanie; Uttenthal, Åse; Lohse, Louise; Nielsen, Jens; Crooke, Helen; Blome, Sandra; Loefffen, Willie

Publication date: 2011

Document Version
Publisher's PDF, also known as Version of record

Link back to DTU Orbit

Citation (APA):

Elbers, A., van der Gaag, L., Schmeiser, S., Uttenthal, A., Lohse, L., Nielsen, J., Crooke, H., Blome, S., & Loefffen, W. (2011). A Bayesian Clinical Decision Support for Early Detection of Classical Swine Fever in Individual Pigs - Evaluation of the Sensitivity and Specificity of the Model. Abstract from 8th ESVV International Pestivirus Symposium, Hanover, Lower Saxony, Germany. http://www.pestivirus2011.de/

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Session 2: Evolution, epidemiology and control - poster ID 11

A Bayesian Clinical Decision Support System for Early Detection of Classical Swine Fever in Individual Pigs -Evaluation of the Sensitivity and Specificity of the Model[#]

Armin Elbers¹, Linda van der Gaag², Stefanie Schmeiser³, Åse Uttenthal⁴, Louise Lohse⁴, Jens Nielsen⁴, Helen Crooke⁵, Sandra Blome⁶, and Willie Loeffen⁷

¹ Department of Epidemiology, Crisis organization and Diagnostics, Central Veterinary Institute, part of Wageningen UR, Lelystad, The Netherlands

² Decision-support Systems group, Department of Information and Computing Sciences, Utrecht University, Utrecht, The Netherlands

³ Institute of Virology, University of Veterinary Medicine Hannover, Germany.

⁴ Department of Virology, National Veterinary Institute, Technical University of Denmark, Lindholm,
Denmark

Virology Department, Veterinary Laboratory Agencies, Addlestone, United Kingdom
 Institute of Diagnostic Virology, Friedrich Loeffler Institute, Insel Riems, Germany
 Department of Virology, Central Veterinary Institute, Lelystad, The Netherlands

Detection of CSF-suspect cases based on clinical signs is difficult. Low specificity of the clinical signs causes the CSF-suspicion to be highly uncertain for a long period.

In order to supply practitioners and state-veterinary-officers with an objective tool to early identify CSF-suspect situations, we developed a clinical decision-support system (CDSS). The system builds upon a probabilistic network, which essentially is a graphical model of a probability distribution over a collection of stochastic variables. The network models the pathogenesis of the disease and includes stochastic variables related to infection and to clinical signs that may occur as a consequence of infection. A prototype of the system is currently operational.

The ability of the CDSS to detect CSF-infected pigs was evaluated using inoculation experiments from Germany, United Kingdom, Denmark and the Netherlands. A total of 128 pigs in 23 experimental groups were inoculated with CSF-virus varying in virulence. For highly virulent CSF-strains, the CDSS was able to detect infected piglets 5-7 days post infection (dpi). For less virulent CSF-strains, infected piglets were detected 14-17 dpi.

The specificity of the model was evaluated using field data, collected by practitioners from the Netherlands, Belgium, Denmark, Italy and Germany. Data was collected from 408 sick pigs for which CSF was in the differential diagnosis but not the first to think of as a possible cause. Given the threshold values of alpha=0.005, 0.01 and 0.05 (a priori probability of CSF being present in the country), the specificity of the CDSS was 96.6%, 97.8% and 99.5%, respectively.