

Bioactivity, Chemical Profiling, and 16S rRNA-Based Phylogeny of Pseudoalteromonas Strains Collected on a Global Research Cruise

Vynne, Nikolaj Grønnegaard; Månsson, Maria; Nielsen, Kristian Fog; Gram, Lone

Published in: Marine Biotechnology

Link to article, DOI: 10.1007/s10126-011-9369-4

Publication date: 2011

Link back to DTU Orbit

Citation (APA):

Vynne, N. G., Månsson, M., Nielsen, K. F., & Gram, L. (2011). Bioactivity, Chemical Profiling, and 16S rRNA-Based Phylogeny of Pseudoalteromonas Strains Collected on a Global Research Cruise. *Marine Biotechnology*, *13*(6), 1062-1073. https://doi.org/10.1007/s10126-011-9369-4

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1	Bioactivity, chemical profiling and 16S rRNA based phylogeny of
2	Pseudoalteromonas strains collected on a global research cruise
3	
4	Nikolaj G. Vynne ^{1*} , Maria Månsson ² , Kristian F. Nielsen ² and Lone Gram ¹
5	
6	¹ Technical University of Denmark, National Food Institute, Søltofts Plads, bldg. 221, DK-2800
7	Kgs. Lyngby, Denmark
8	² Technical University of Denmark, Department of Systems Biology, Søltofts Plads, bldg. 221, DK-
9	2800 Kgs. Lyngby, Denmark
10	
11	* corresponding author:
12	phone: +45 4525 2571
13	fax: +45 4588 4774
14	e-mail: ngvy@food.dtu.dk
15	
16	
17	December 13 th 2010
18	
19	

1 ABSTRACT

2 One hundred one antibacterial Pseudoalteromonas strains that inhibited growth of a Vibrio 3 anguillarum test strain were collected on a global research cruise (Galathea 3), and 51 of the strains 4 repeatedly demonstrated antibacterial activity. Here, we profile secondary metabolites of these 5 strains to determine if particular compounds serve as strain or species markers and to determine if the secondary metabolite profile of one strain represents the bioactivity of the entire species. 16S 6 7 rRNA gene similarity divided the strains into two primary groups: One group (51 strains) consisted 8 of bacteria which retained antibacterial activity, 48 of which were pigmented, and another group (50 9 strains) of bacteria which lost antibacterial activity upon sub-culturing, two of which were 10 pigmented. The group that retained antibacterial activity consisted of six clusters in which strains 11 were identified as Pseudoalteromonas luteoviolacea, Pseudoalteromonas aurantia, 12 Pseudoalteromonas phenolica, Pseudoalteromonas ruthenica, Pseudoalteromonas rubra and 13 Pseudoalteromonas piscicida. HPLC-UV/VIS analyses identified key peaks, such as violacein in P. 14 *luteoviolacea*. Some compounds, such as a novel bromoalterochromide were detected in several species. HPLC-UV/VIS detected systematic intra-species differences for some groups and testing 15 16 several strains of a species was required to determine these differences. The majority of non-17 antibacterial, non-pigmented strains were identified as Pseudoalteromonas agarivorans and HPLC-18 UV/VIS did not further differentiate this group. Pseudoalteromonas retaining antibacterial were 19 more likely to originate from biotic or abiotic surfaces in contrast to planktonic strains. Hence, the pigmented, antibacterial Pseudoalteromonas have a niche specificity, and sampling from marine 20 21 biofilm environments is a strategy for isolating novel marine bacteria that produce antibacterial 22 compounds. 23

24 Words: 252

25

26 KEYWORDS: Pseudoalteromonas, antibacterial activity, secondary metabolites, bioprospecting,

27 Galathea 3

1 INTRODUCTION

2

3 Compounds of relevance for the pharmaceutical and biotechnology industries are produced by 4 marine microorganisms (Burgess et al., 1999) and it has been suggested that some compounds of 5 pharmacological interest previously attributed to macroorganisms may in fact be of microbial origin (Bewley and Faulkner, 1998;Simmons et al., 2008;Sudek et al., 2006). The emergence of 6 7 multiresistant pathogenic bacterial strains and the failure of combinatorial and diversity oriented 8 chemistry to adequately supply the drug discovery pipeline (Newman, 2008) has re-invigorated 9 natural product chemistry as a path for discovery and development of new antibiotics. With this in 10 mind, we isolated marine bacteria with antibacterial activity during the Danish Galathea 3 marine 11 research expedition (Gram et al., 2010). The antibacterial strains were tentatively identified using 16S rRNA-similarity and one of the major groups of isolated bacteria was identified as 12 13 Pseudoalteromonas (Gram et al., 2010).

14

15 The genus Pseudoalteromonas consists of Gram negative marine bacteria belonging to the y-16 proteobacteria and are present globally in marine waters where they constitute 0.5 to 6% of the total 17 bacterioplankton (Wietz et al., 2010). They are heterotrophic aerobes, non-fermentative and the 18 cells are motile by one or more polar flagella. The genus divides into two groups; pigmented and 19 non-pigmented species. The pigmented species are often producers of bioactive secondary 20 metabolites (Bowman, 2007) displaying cytotoxic (Zheng et al., 2006), antibacterial (Gauthier, 21 1976b;Gauthier and Flatau, 1976;Isnansetyo and Kamei, 2003;Jiang et al., 2000;McCarthy et al., 22 1994), antifungal (Franks et al., 2006; Kalinovskaya et al., 2004) or antifouling (Egan et al., 23 2001;Holmström et al., 2002) effects. It has been hypothesized that bioactive Pseudoalteromonas 24 are primarily associated with higher organisms (Holmström and Kjelleberg, 1999), suggesting an 25 ecological role in which some bioactive species might play an active part in host defense against pathogens and fouling organisms (Holmström et al., 1996;Armstrong et al., 2001;Egan et al., 2008). 26

A link between surface colonization and antibacterial activity has not been experimentally verified,
although several studies have successfully isolated epiphytic bacteria with antibacterial activity
from algae and other marine organisms (Armstrong et al., 2001;Boyd et al., 1999;James et al.,
1996;Penesyan et al., 2009). The group of non-pigmented species have highly similar 16S rRNA
gene sequences (Ivanova et al., 2004) and is rarely inhibitory against other microorganisms,
although e.g. *P. haloplanktis* strain INH produces isovaleric acid and 2-methylbutyric acid showing
a broad spectrum of bacterial inhibition (Hayashida-Soiza et al., 2008).

8

9 Phylogeny and differentiation of bacterial species rely heavily on 16S rRNA gene similarity 10 (Stackebrandt et al., 2002) however, 16S rRNA gene similarity does not provide sufficient 11 differentiation below species level (Fox et al., 1992). Comparison of secondary metabolite 12 production has supported species delineation within the Actinomycete genus Salinispora, where 13 strains with high 16S rRNA similarity were shown to belong to distinct species each with different 14 specific metabolite profiles (Jensen et al., 2007). In mycology, comparison of chemical profiles 15 (e.g. TLC, direct-infusion mass spectrometry, and HPLC with various detectors such as UV/VIS 16 and/or mass spectrometry) of secondary metabolites has been widely used to identify and 17 differentiate filamentous fungi (chemotaxonomy) also at sub-species level (Frisvad et al., 2008), 18 and the chemophylogeny correlates well with phylogenetic analysis of sequences of specific 19 housekeeping genes (e.g. chitin synthase, β-tubulin and calmodulin) (Geiser et al., 2007). Since 20 several *Pseudoalteromonas* species produce a range of secondary metabolites, we hypothesized that 21 chemical profiling and specific marker compounds could be indicative of bioactive potential and at 22 the same time be useful in species identification or differentiation (Jensen et al., 2007).

23

The aim of the present study was to profile the secondary metabolites of these strains to determine if particular compounds serve as markers of strains or species with antibacterial activity, and to determine if several strains of each species must be tested to assess the full bioactivity potential. As

1	part of this we provide accurate identification and phylogeny of these organisms by detailed 16S
2	rRNA gene sequence comparative analysis. Since the bacteria were isolated from different sample
3	types, our collection also allows us to address aspects of <i>Pseudoalteromonas</i> ecology such as the
4	possible link between surface or planktonic lifestyle and antibacterial activity.
5	
6	
7	

MATERIALS AND METHODS

4	Strain isolation. Approx. 500 marine bacterial strains with antagonist activity against V.
5	anguillarum strain 90-11-287 (Skov et al., 1995) were isolated during the Danish Galathea 3
6	research expedition (Gram et al., 2010). One hundred and one of these strains tentatively identified
7	as Pseudoalteromonas species were included in the present study. V. anguillarum 90-11-287 was
8	used as target strain since the expedition ship was not equipped to handle e.g. potential human
9	pathogens, and this Vibrio strain is in our experience very sensitive to antibacterial compounds from
10	other marine bacteria (Hjelm et al., 2004).
11	
12	Growth media and culture conditions. Pseudoalteromonas strains were grown in marine broth
13	(MB) 2216 (Difco, Detroit, USA) and on marine agar (MA) 2216 (Difco, Detroit, USA) prepared in
14	accordance with the manufacturer's instructions. Broth cultures were incubated under stagnant
15	conditions at 25°C. Pigment production was determined by visual inspection of 48 hour old culture
16	broths (MB) and colonies grown on MA for 24 to 48 hours.
17	
18	Antibacterial activity. Instant Ocean (IO) bioassay agar plates were prepared as described by
19	Hjelm et al. (Hjelm et al., 2004). Ten g/l agar, 3.3 g/l casamino acids (Difco 223050, Detroit, USA)
20	and 30 g/l Instant Ocean aquatic salts (Instant Ocean® Aquarium systems Inc., Sarrebourg, France)
21	were added to distilled water and autoclaved. Glucose (0.4%) and 10 μ l/ml of V. anguillarum
22	overnight culture was added to the cooled (44°C) IO and plates poured. The plates were allowed to
23	dry for 15 minutes and if used for well diffusion agar assays (WDAA), wells (diameter 6 mm) were
24	punched. The inhibitory activity of live <i>Pseudoalteromonas</i> bacterial cells was tested by spotting 48
25	hour old MA grown colonies on freshly prepared IO-agar plates containing V. anguillarum. Plates
26	were incubated at 25°C and inspected for clearing zones in the growth of V. anguillarum after 24h.
27	

1 Cell-free supernatants were prepared to test for the presence of water soluble antibacterial 2 compounds secreted to the broth and ethyl acetate extracts were prepared to test for production of 3 non-polar antibacterial compounds. Each strain was grown in 20 ml of MB for 48 h. A 1.5 ml 4 sample was withdrawn for 0.2 µm filtering and subsequently the remainder of the culture was 5 extracted with an equal volume of ethyl acetate. The ethyl acetate fraction was transferred to a new 6 vessel, evaporated to dryness and redissolved in 2×0.5 ml ethyl acetate. The 1.5 ml cell-free 7 sterile filtered supernatant and the ethyl acetate extracts were stored at -20°C until tested in the 8 WDAA (50 µl sample per well) based on IO-agar plates containing V. anguillarum. Controls 9 (sterile MB and pure ethyl acetate) did not cause any inhibition zones.

10

The number of antibacterial *Pseudoalteromonas* strains in surface samples (e.g. algae, driftwood, fish and sediment samples) was compared to their numbers in water samples by the Fisher's exact test (Fisher, 1958). A 2 × 2 contingency table was used to test the hypothesis that presumed antibacterial strains with stable antibacterial activity were equally likely to be isolated from water samples and surface samples.

16

17 16S rRNA gene sequence analyses. A detailed phylogenetic analysis was performed on 16S 18 rRNA sequences obtained in a previous study (Gram et al., 2010). For the analysis in this study, we 19 conducted a BLAST (http://blast.ncbi.nlm.nih.gov) search against a compilation of 20 Pseudoalteromonas type strain sequences retrieved from GenBank (list of type strains obtained 21 from http://www.bacterio.cict.fr) and sequences of the type strains with a BLAST match in our 22 strain collection were included in 16S rRNA sequence analysis to obtain a robust phylogenetic tree. 23 Sequences from two additional *Pseudoalteromonas* strains were included: The genus type strain *P*. 24 haloplanktis and the bioactive P. tunicata. Salinispora arenicola CNS-205 was used as out-group. 25 The sequences were aligned by the MAFFT online software (http://www.ebi.ac.uk/Tools/mafft/) 26 (Katoh et al., 2002) and curated with the Gblocks software on its least stringent settings

(Castresana, 2000;Talavera and Castresana, 2007). The resulting alignment was processed using the
MEGA4 software (Tamura et al., 2007) to create neighbor joining and minimum evolution trees.
PhyML 3.0 was used to generate a maximum likelihood tree (Guindon and Gascuel, 2003).
Phylogenetic trees were generated under default parameters with 1,000 bootstrap replications for
neighbor joining and minimum evolution trees, and 100 bootstrap replications for the maximum
likelihood tree. GenBank accession numbers for the *Pseudoalteromonas* strains used in this study
are included in supplementary table 1.

8

9 HPLC-UV/VIS analysis of secondary metabolites. The strains were grown in static cultures in 10 10 ml MB for 3 days at 25°C, and for each species one strain was cultured in triplicate to establish 11 extraction and growth variation. Cultures were extracted with equal volumes of ethyl acetate, 12 centrifuged and the ethyl acetate and evaporated under N₂ to dryness. Samples were re-dissolved in 13 $300 \,\mu$ l acetonitrile-water (1:1 v/v), and filtered through a 13 mm ID PFTE syringe filter. A 14 subsample of 2 µl was then analyzed by reversed phase HPLC on an Agilent 1100 System equipped 15 with a UV/VIS photo diode array detector (DAD, scanning 200-600 nm). Separation was done on a 16 100 mm \times 2 mm i.d., 3 µm Gemini C₆-phenyl column (Phenomenex, Torrance, CA), running at 40°C 17 using a binary linear solvent system of water (A) and acetonitrile (B) (both buffered with 50 μ l/l 18 trifluoroacetic acid) at a flow of 300 μ l/min. The gradient profile was: t= 0 min, 5% B; t =22 min, 70% 19 B; t=24.5,100% B, t= 27 min, 100% B; t =29 min, B= 5%; holding this for 8 minutes prior to the next 20 injection. The chromatographic profiles were compared, subtracting peaks present in media blank 21 extracts. Samples were analyzed in random order, and 6 of the first extracts were analyzed several 22 times during the sequence to determine any retention time shifts. Cluster analysis was done on a matrix 23 of detected / non-detected peaks (1/0) using NTSYSpc 2.20q (Exeter Software, Setauket, NY). SAHN 24 clustering was used by UPGMA (unweighted pair-group method) and simple distance measurement. 25 Representative extracts were also analyzed by HPLC-UV/VIS-TOFMS in both positive and 26 negative electro spray (Nielsen and Smedsgaard, 2003). Peaks were tentatively identified by UV-

- spectra and accurate mass data by matching in Antibase 2009 (35 930 microbial secondary
- metabolites) (Wiley & Sons, Hoboken, NJ) (Nielsen et al., 2006).

2 **RESULTS**

3

1

4 Pigmentation and antibacterial activity. The one hundred and one *Pseudoalteromonas* strains 5 were originally isolated for their ability to inhibit V. anguillarum (Gram et al., 2010). However, on 6 re-cultivation and re-testing for the ability to inhibit V. anguillarum after storage at -80°C for 7 several months, only 51 strains retained inhibitory activity (Table 1). These 51 strains were all 8 inhibitory when tested as live cultures in the 'spot test' assay. Twenty of the strains produced water-9 soluble, diffusible antibacterial substances as indicated by the ability of cell-free sterile filtered 10 supernatant to inhibit growth of V. anguillarum in the WDAA. These 20 strains were identified as 11 P. phenolica (1 strain), P. luteoviolacea (5 strains), P. rubra (9 strains), P. citrea (1 strain) and P. 12 aurantia (4 strains). Ethyl acetate extraction of culture broths resulted in 19 crude extracts which 13 inhibited growth of v. anguillarum in the WDAA. Four of these were identified as P. luteoviolacea 14 and were the only strains where both cell-free supernatant and crude ethyl acetate extracts inhibited 15 Vibrio growth. The remaining 15 inhibitory crude extracts all originated from strains identified as P. 16 ruthenica. The crude ethyl acetate extracts of 26 strains were intensely colored, however only some 17 of these extracts inhibited growth of *Vibrio* indicating that the pigments were not universally 18 antibacterial. Cell-free culture supernatants and ethyl acetate crude extracts of strains with no 19 growth inhibition of V. anguillarum in the 'spot test' assay were also tested but showed no growth 20 inhibition.

Forty eight of the fifty one antibacterial strains were pigmented, while two (S3431 and S3655) of the fifty non-active strains were pigmented (Table 1). Antibacterial activity was significantly more likely to be produced by pigmented strains as determined by Fisher's exact test (2-tailed *p*-value of 0.0000). In total, 70 strains were isolated from surface swabs and 31 from water samples. Of the surface associated strains 45 remained active in the spot-assay in comparison to 6 of the water sample strains. The Fisher's exact test demonstrated a significant relation between surface association and stable antibacterial activity (2-tailed *p*-value of 0.0000).

1

Pseudoalteromonas strains were isolated on all parts of the global cruise in both tropical and
temperate waters (Figure 1). Our strain collection is not large enough to allow for complete
biogeographic analysis but pigmented strains appeared to be more frequent in coastal areas whereas
the non-pigmented strains appeared associated with open waters (Figure 1).

6

16S rRNA gene sequence analyses. We initially performed a BLAST search querying the 16S rRNA gene sequence of each strain against the GenBank database. The results of this analysis were ambiguous, as some sequences returned more than 40 hits all with identical scores in the BLAST results (data not shown), frequently including the sequences of several different *Pseudoalteromonas* species. Therefore, a BLAST analysis was carried out querying the sequences against a complete set of *Pseudoalteromonas* type strains and hence each strain is matched with the best type strain BLAST match (Suppl Table 1).

14

The 16S rRNA gene sequences were used to cluster the strains by constructing a neighbor-joining tree and branch support was verified by comparison to minimum evolution and maximum likelihood trees (Figure 2). Nodes supported by an 80% bootstrap cut-off were collapsed when 3 or more strains were included in the cluster (Figure 2). An exception was made for clusters VI and VII, which are shown as separate clusters due to obvious differences in phenotype (pigment, bioactivity, secondary metabolite profile). Ninety-nine of the strains fell into one of 8 primary clusters.

22

23 Clusters I and III consisted of non-pigmented non-inhibitory strains (Table 1). Cluster I included 38

24 strains and the type strains of *P. haloplanktis*, *P. agarivorans*, *P. tetraodonis*, *P. paragorgicola*, *P.*

25 distincta, P. arctica, P. nigrifaciens, P. elyakovii, P. carrageenovora, P. marina and P. aliena.

26 Strain S3431 – a black pigmented strain in cluster I – did show more than 97% similarity even when

1 compared to the full GenBank database which suggests that S3431 could represent a novel

Pseudoalteromonas species. Despite the low BLAST similarity score, phylogenetic analysis and tree construction placed strain S3431 in the diverse cluster I (the non-collapsed cluster I is shown in supplementary figure 1). Cluster III contained no type strains which supported the BLAST analysis where the strains in cluster III were 97% similar to the best type strain match (Supplementary table 1).

7

8 The remaining six of the eight clusters contained pigmented strains. Pale yellow strains clustered 9 with the type strains of *P. citrea* and *P. aurantia* (cluster II) and four intensely purple strains 10 grouped in cluster V with the type strain of P. luteoviolacea. Cluster VI contained nine red-11 pigmented strains and the type strain of *P. rubra* and cluster VII consisted of 12 intensely yellow 12 strains, 1 pale yellow strain and the P. flavipulchra, P. maricaloris and P. piscicida type strains. 13 Fifteen strains and their nearest BLAST match, P. ruthenica, formed cluster VII. These strains all 14 produced a pale brown pigment. Cluster IV contained six strains and the type strain of *P. phenolica*. 15 Four of the strains in this cluster had *P. phenolica* as their best type strain BLAST match; however 16 strain S1093 had *P. luteoviolacea* as its best match at 98% identity, while *P. rubra* and *P.* 17 *luteoviolacea* type strains scored identically (97%) as the best matches for S2724. The strains in 18 cluster IV were heterogeneous with respect to pigmentation, some were non-pigmented and others 19 appeared brown.

20

Profiling of secondary metabolites. The 101 strains and select type strains were separable in discrete groups by HPLC-UV/VIS (Figure 3) and the triplicate profiles from an isolate of each species were very reproducible and could be superimposed on each other (data not shown). All of the 38 strains of the 16S rRNA cluster I fell into group A, in which no UV/VIS peaks were unique compared to the media blanks indicating that no secondary metabolites were produced. This large group also included all nine strains from 16S rRNA cluster III and strains of *P. phenolica* and *P*. 1 *ruthenica* less proficient in secondary metabolite production. A summary of the detected

2 compounds is shown in table 2 and the producer organisms are shown by 16S rRNA cluster in table
3 3.

4

5 Based on their production of specific metabolites, the majority of pigmented bacteria formed six main groups not including four *P. rubra* strains (Figure 3). In the pigmented bacteria, a total of 26 6 7 distinct peaks were detected and included in the cluster analysis. We identified indolmycin, 8 violacein, and prodigiosin among the significant peaks based on reference standards. Furthermore, 9 9 peaks could be tentatively identified based on HPLC-UV/VIS-TOFMS results and data in 10 Antibase2009. These nine included two likely novel bromoalterochromides and a brominated indole 11 (Table 2). 12 13 Comparing the 16S rRNA gene sequence clusters with the chemical profiling revealed several 14 patterns. Some compounds were exclusively produced by strains belonging to the same cluster whereas other compounds were produced across several strains from different clusters. All strains in 15 16 cluster II (*P. aurantia / P. citrea*) shared production of compound B (retention time, RT 12.31 min) 17 whereas the production of 4 other compounds F, Q, T and U (RT 15.47, 16.60, 17.96 and 18.28

18 min), were scattered in the group. This included a novel bromoalterochromide (compound Q, RT

19 16.60 min) that was also found in cluster VII (P. flavipulchra/P. piscicida). Three of the other

20 compounds were identified as quinolins based on distinct UV spectra and accurate mass.

21

P. luteoviolacea strains (cluster V) shared production of compound D (violacein, RT 14.29) in all
four strains and the type strain, but were sub-divided by compound A (indolmycin, RT 11.21)
produced by 2 strains and compound Z (pentabromopseudilin, RT 22.65) produced by the 2 other
strains and the type strain. This division is visible in figure 4, which shows chromatograms of the 4
strains in cluster V. Interestingly, also two *P. phenolica* strains produced compound Z.

2 Cluster VII (P. flavipulchra and P. piscicida) was chemically very homogeneous. All strains except 3 one produced three bromoalterochromides P, Q and R (RT 16.49, 16.60 and 17.10) of which Q and 4 R were novel compounds. In contrast to cluster V and compound D, the production of P, Q and R 5 was not a unique marker for strains of this cluster as P, Q and R were also detected in one strain 6 from cluster II and one strain from cluster VI. 7 8 Thirteen of the 15 strains in cluster VIII, identified as P. ruthenica, shared a unique chemical profile 9 and produced the compounds H, K and O (RT 15.78, 15.98, 16.30) with characteristic UV spectra. 10 None of these matched compounds in Antibase2009 and potentially constitute novel antibacterials. 11 These compounds were not detected among strains from other clusters, yet they were not suitable as 12 a distinct chemical marker for cluster VIII since no secondary metabolites were detected in the 13 remaining two strains in this cluster or the type strain. 14 15

The strains in cluster VI were identified as *P. rubra* and were chemically very heterogeneous. Five of nine strains produced the red pigment prodigiosin (compound M, RT 16.00) which was not detected in strains of other clusters. Additionally, a multitude of known and non-identified compounds were detected in one or more strains in the cluster. In total, 16 compounds were detected within the cluster and 12 of these were unique for this cluster. Only two strains shared an identical production of secondary metabolites, further stressing the chemical diversity among the strains in this cluster.

23

- 24
- 25
- 26

2 **DISCUSSION**

3

1

4 We demonstrate in this study, in agreement with earlier findings (Bowman, 2007), that species 5 within the *Pseudoalteromonas* genus produce a range of secondary metabolites, some with 6 antibacterial activity. Several species of the genus are intensely pigmented and it is hypothesized 7 that pigmentation co-occur with antibacterial activity (Egan et al., 2002). In our global collection of 8 Pseudoalteromonas strains that demonstrated antibacterial activity on initial isolation, strains that 9 were pigmented were significantly more likely to retain antibacterial activity on re-growth than 10 non-pigmented strains. Several intensely coloured organic extracts were not inhibitory against V. 11 anguillarum and hence we do not believe that the pigments, in general, are the cause of the 12 antibacterial activity although e.g. the purple pigment violacein is a known antibiotic compound 13 (Lichstein and Vandesand, 1945).

14

15 Nearly half of the isolated strains did not retain any antibacterial effect after frozen storage and 16 subculturing despite being isolated on the original plates due to antibacterial activity. The observed 17 loss of antibacterial activity may be due to a requirement for factors specific to local seawater, as 18 initial tests for antibacterial activity were carried out using 50% local seawater (Gram et al., 2010). 19 Furthermore, loss of antibacterial activity may be due to repression or inhibition of gene clusters 20 encoding products that are required for secondary metabolite synthesis (e.g. by catabolite 21 repression). A reduction in antibiotic production when the producer organism is grown in excess of 22 a carbon source is a known phenomenon (Sanchez et al., 2010) and suppression of secondary 23 metabolite production by excess concentrations of other substrate components is demonstrated in 24 Streptomyces (Doull and Vining, 1990). This could suggest that culturing the strains under nutrient 25 limited conditions may reestablish production of antibacterial compounds. Also, during the original 26 sampling and screening procedure, the agar plates may have harbored co-cultured microorganisms 27 which potentially induce antibacterial activity as has been demonstrated by Mearns-Spragg et al.

(Mearns-Spragg et al., 1998). Hence, it may be possible to re-induce the antibacterial activity if the
 right conditions can be created.

3

4 Several bioactive *Pseudoalteromonas* have been isolated from higher organisms and it has been 5 hypothesised that antibacterial compounds may play a role in bacterial competition or as protective 6 agents beneficial for the host organism (Holmström and Kjelleberg, 1999). We provide statistical 7 evidence that surface associated presumed antibacterial pseudoalteromonads are significantly more 8 likely to show stable production of antibacterial compounds than Pseudoalteromonas species 9 isolated as planktonic cells. This suggests that production of antibacterial compounds may play an 10 important role in the ability of *Pseudoalteromonas* strains to colonize and persist on surfaces 11 submerged in the marine environment, as previously suggested for *P. tunicata* strain D2 (Rao et al., 2005). 12

13

14 The analysis of 16S rRNA gene sequences from our global collection of Pseudoalteromonas 15 confirms that phylogenetic analysis results in a number of clusters encompassing predominantly 16 pigmented species or non-pigmented species (Ivanova et al., 2004). Strain S3431 was the single 17 pigmented strain in the so-called non-pigmented clusters. Novel diversity might be represented in 18 cluster III which consisted of strains with 98% or less 16S rRNA gene sequence similarity to 19 *Pseudoalteromonas* type strains and formed a separate cluster in the phylogenetic analysis. 20 However, these strains showed no antibacterial activity and no small molecule metabolites were 21 detected. Such novel diversity could still represent untapped biotechnological potential, producing 22 e.g. enzymes or peptides with biological activity, as known for other non-pigmented 23 Pseudoalteromonas (Hoyoux et al., 2001; Violot et al., 2005).

24

Chemical profiling of the strains detected an array of secondary metabolites. In addition to
 complementing our analysis of 16S rRNA gene sequences, it also demonstrated that some

1 compounds (e.g. violacein, prodigiosin) were characteristic of a species, other compounds were 2 produced by several species and we also detected intra-species clusters of different secondary 3 metabolite profiles. In a broad sense, the clustering based on 16S rRNA gene similarity agreed with 4 the groups resulting from the chemophylogenetic analysis. However, some compounds were 5 produced by organisms of different species that then clustered together using the secondary 6 metabolites as basis. The chemical analysis separated the 4 isolated P. luteoviolacea strains into two 7 distinct sub-groups, showing intra-species chemical diversity. The P. luteoviolacea strains 8 produced violacein and pentabromopseudilin which are active against gram-positive and gram-9 negative bacteria, and the anti-staphylococcal agent indolmycin (Hornemann et al., 1971;Hurdle et 10 al., 2004). Violacein and pentabromopseudilin have previously been detected in *Pseudoalteromonas* 11 luteoviolacea (Gauthier, 1976a; Laatsch and Pudleiner, 1989), but to our knowledge this is the first 12 report of *Pseudoalteromonas* strains producing indolmycin (Månsson et al., 2010).

13

14 Within some species, all strains were consistently antibacterial. However, in others such activity did 15 not appear to be a consistent trait of the species. For instance, strains of the 16S cluster VI (P. 16 phenolica) were heterogeneous in their ability to inhibit Vibrio in our assays, while all but one 17 strain in the homogeneus cluster VII had identical metabolite profiles and all were inhibitory. Even 18 more obvious was the heterogeneous chemical profiles within the P. rubra strains. All except one 19 strain shared a chemical marker prodigiosin and/or RT 15.99 min, but had major variations in 14 other compounds. This may in part be due to loss of ability to produce a compound. For instance, 20 21 strain S2471 over time lost ability to produce the brominated indole (RT 13.78 min). Also, we note 22 that the type strain DSM 6842 (ATCC 29570) did not in our culture produce prodigiosin which has 23 been observed previously (Gauthier, 1976b;Gauthier and Flatau, 1976). The consistent 24 bromoalterochromide production in the two species P. piscicida and P. flavipulchra/maricaloris 25 (cluster VII) was expected (Speitling et al., 2007) and supported the high DNA sequence similarity 26 between the two. This emphasizes the need to isolate and screen multiple strains from each species

when bioprospecting within the genus *Pseudoalteromonas*, as even the homogeneous cluster VII
 contains one strain with a metabolite profile that does not share a single compound with the other
 strains in this cluster.

4

5 Several of the 26 detected peaks were known substances, with a majority known as antibacterials. 6 These included: violacein (Lichstein and Vandesand, 1945), two bromopseudilins (Lovell, 1966), 7 two indolmycins (Werner, 1980), four quinolins (Wratten et al., 1977); and prodigiosin (Kalesperis 8 et al., 1975). Due to its very low aqueous solubility violacein probably protects against predation 9 rather than acts as a true antibiotic, and it has been shown to induce cell death in grazing organisms 10 (Matz et al., 2008). Such compounds would be very beneficial for protection of a biofilm, which is 11 likely how surface associated *Pseudoalteromonas* would grow. The 14 compounds that could not be identified were mainly not identified due to poor ionization in ESI⁺ and ESI⁻ and/or several 12 13 plausible candidates in Antibase2009. However, for chemotaxonomic studies identity of the 14 compounds is not necessary as long as they can be unambiguously identified between samples 15 (Frisvad et al., 2008).

16

17 Within cluster VIII (P. ruthenica) and cluster II (aurantia/citrea) we found examples where strains 18 with highly similar 16S rRNA gene sequences (>99%) and with identical chemotaxonomy 19 originated from geographically distinct locations. This latter observation is in agreement with 20 studies on *Salinispora* biogeography and secondary metabolite production in which the authors 21 show how strains of the marine bacterium Salinispora arenicola isolated from worldwide locations 22 are highly related and produce identical patterns of secondary metabolites (Jensen and Mafnas, 23 2006; Jensen et al., 2007). In contrast, the P. luteoviolacea and P. rubra strains showed both local 24 and global variations in their secondary metabolite profile, which one might speculate is due to 25 adaptation to local specific niches.

- 1 In conclusion, we believe sampling from specific niches, e.g. biofilms on surfaces, to be of
- 2 importance in discovery of novel secondary metabolites from the genus *Pseudoalteromonas*. While
- 3 differences in metabolite patterns among species encourage isolation and screening of novel
- 4 diversity, bioprospecting known *Pseudoalteromonas* species should not be ruled out. Investigation
- 5 of multiple strains of one *Pseudoalteromonas* species can yield novel compounds due to intra-
- 6 species variations within secondary metabolite profiles.
- 7
- 8
- 9

1

2 Acknowledgements

- 3 We acknowledge Dr. Jesper B. Bruhn for valuable input during the early phase of this study. This
- 4 study was supported by the Programme Commission on Health, Food and Welfare under the Danish
- 5 Council for Strategic Research. The present work was carried out as part of the Galathea 3
- 6 expedition under the auspices of the Danish Expedition Foundation. This is Galathea 3 contribution
- 7 no. pXXXXXX (to be added if/when accepted)

- 9
- 10

1 2		Reference List
3 4	1.	Armstrong, E., Yan, L.M., Boyd, K.G., Wright, P.C., and Burgess, J.G. (2001). The symbiotic role of marine microbes on living surfaces. Hydrobiologia 461:37-40.
5 6	2.	Bewley, C.A. and Faulkner, D.J. (1998). Lithistid sponges: Star performers or hosts to the stars. Angew Chem Int Ed 37:2162-2178.
7 8	3.	Bowman, J.P. (2007). Bioactive compound synthetic capacity and ecological significance of marine bacterial genus <i>Pseudoalteromonas</i> . Mar Drugs 5:220-241.
9 10	4.	Boyd,K.G., Adams,D.R., and Burgess,J.G. (1999). Antibacterial and repellent activities of marine bacteria associated with algal surfaces. Biofouling 14:227-236.
11 12	5.	Burgess, J.G., Jordan, E.M., Bregu, M., Mearns-Spragg, A., and Boyd, K.G. (1999). Microbial antagonism: a neglected avenue of natural products research. J Biotechnol 70:27-32.
13 14	6.	Castresana, J. (2000). Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. Mol Biol Evol 17:540-552.
15 16 17	7.	Doull,J.L. and Vining,L.C. (1990). Nutritional control of actinorhodin production by <i>Streptomyces coelicolor</i> A3(2) - Suppressive effects of nitrogen and phosphate. Appl Microbiol Biotechnol 32:449-454.
18 19 20	8.	Egan,S., Holmström,C., and Kjelleberg,S. (2001). <i>Pseudoalteromonas ulvae</i> sp nov., a bacterium with antifouling activities isolated from the surface of a marine alga. Int J Syst Evol Microbiol 51:1499-1504.
21 22 23	9.	Egan, S., James, S., Holmström, C., and Kjelleberg, S. (2002). Correlation between pigmentation and antifouling compounds produced by <i>Pseudoalteromonas tunicata</i> . Environ Microbiol 4:433-442.
24 25 26	10.	Egan,S., Thomas,T., and Kjelleberg,S. (2008). Unlocking the diversity and biotechnological potential of marine surface associated microbial communities. Curr Opin Microbiol 11:219-225.
27	11.	Fisher, R.A. (1958). Statistical methods for research workers. New York: Hafner.
28 29	12.	Fox,G.E., Wisotzkey,J.D., and Jurtshuk,P. (1992). How close is close: 16S rRNA sequence identity may not be sufficient to guarantee species identity. Int J Syst Evol Microbiol 42:166.
30 31 32	13.	Franks,A., Egan,S., Holmstrom,C., James,S., Lappin-Scott,H., and Kjelleberg,S. (2006). Inhibition of fungal colonization by <i>Pseudoalteromonas tunicata</i> provides a competitive advantage during surface colonization. Appl Environ Microbiol 72:6079-6087.
33 34	14.	Frisvad,J.C., Andersen,B., and Thrane,U. (2008). The use of secondary metabolite profiling in chemotaxonomy of filamentous fungi. Mycol Res 112:231-240.
35 36	15.	Gauthier, M.J. (1976a). Morphological, physiological, and biochemical characteristics of some violet-pigmented bacteria isolated from seawater. Can J Microbiol 22:138-149.
37 38	16.	Gauthier, M.J. (1976b). <i>Alteromonas rubra</i> sp. nov., a new marine antibiotic-producing bacterium. Int J Syst Bacteriol 26:459-466.

- Gauthier, M.J. and Flatau, G.N. (1976). Antibacterial activity of marine violet-pigmented
 Alteromonas with special reference to the production of brominated compounds. Can J
 Microbiol 22:1612-1619.
- 4 18. Geiser, D.M., Klich, M.A., Frisvad, J.C., Peterson, S.W., Varga, J., and Samson, R.A. (2007). The
 5 current status of species recognition and identification in *Aspergillus*. Stud Mycol 59:1.
- Gram,L., Melchiorsen,J., and Bruhn,J.B. (2010). Antibacterial activity of marine culturable
 bacteria collected from a global sampling of ocean surface waters and surface swabs of marine
 organisms. Mar Biotechnol 12:439-451.
- 9 20. Guindon,S. and Gascuel,O. (2003). A simple, fast, and accurate algorithm to estimate large
 10 phylogenies by maximum likelihood. Syst Biol 52:696.
- Hayashida-Soiza,G., Uchida,A., Mori,N., Kuwahara,Y., and Ishida,Y. (2008). Purification and characterization of antibacterial substances produced by a marine bacterium
 Pseudoalteromonas haloplanktis strain. J Appl Microbiol 105:1672-1677.
- Hjelm,M., Bergh,Ø., Riaza,A., Nielsen,J., Melchiorsen,J., Jensen,S., Duncan,H., Ahrens,P.,
 Birkbeck,H., and Gram,L. (2004). Selection and identification of autochthonous potential
 probiotic bacteria from turbot larvae (*Scophthalmus maximus*) rearing units. Syst Appl
 Microbiol 27:360-371.
- Holmström, C., Egan, S., Franks, A., McCloy, S., and Kjelleberg, S. (2002). Antifouling
 activities expressed by marine surface associated *Pseudoalteromonas* species. FEMS
 Microbiol Ecol 41:47-58.
- Holmström, C., James, S., Egan, S., and Kjelleberg, S. (1996). Inhibition of common fouling
 organisms by marine bacterial isolates with special reference to the role of pigmented bacteria.
 Biofouling 10:251-259.
- 24 25. Holmström, C. and Kjelleberg, S. (1999). Marine *Pseudoalteromonas* species are associated
 25 with higher organisms and produce biologically active extracellular agents. FEMS Microbiol
 26 Ecol 30:285-293.
- 26. Hornemann,U., Hurley,L.H., Speedie,M.K., and Floss,H.G. (1971). Biosynthesis of
 indolmycin. J Am Chem Soc 93:3028-3035.
- 27. Hoyoux, A., Jennes, I., Dubois, P., Genicot, S., Dubail, F., Francois, J.M., Baise, E., Feller, G., and
 30 Gerday, C. (2001). Cold-adapted beta-galactosidase from the Antarctic psychrophile
 31 *Pseudoalteromonas haloplanktis*. Appl Environ Microbiol 67:1529-1535.
- Hurdle, J.G., O'Neill, A.J., and Chopra, I. (2004). Anti-staphylococcal activity of indolmycin, a
 potential topical agent for control of staphylococcal infections. J Antimicrob Chemothe
 54:549-552.
- Isnansetyo,A. and Kamei,Y. (2003). MC21-A, a bactericidal antibiotic produced by a new
 marine bacterium, *Pseudoalteromonas phenolica* sp. nov. O-BC30T, against methicillin resistant *Staphylococcus aureus*. Antimicrob Agents Chemother 47:480-488.
- 38 30. Ivanova,E.P., Flavier,S., and Christen,R. (2004). Phylogenetic relationships among marine
 39 *Alteromonas*-like proteobacteria: emended description of the family *Alteromonadaceae* and
 40 proposal of *Pseudoalteromonadaceae* fam. nov., *Colwelliaceae* fam. nov., *Shewanellaceae*

- fam. nov., *Moritellaceae* fam. nov., *Ferrimonadaceae* fam. nov., *Idiomarinaceae* fam. nov.
 and *Psychromonadaceae* fam. nov. Int J Syst Evol Microbiol 54:1773-1788.
- 31. James,S.G., Holmström,C., and Kjelleberg,S. (1996). Purification and characterization of a
 novel antibacterial protein from the marine bacterium D2. Appl Environ Microbiol 62:2783 2788.
- G 32. Jensen, P.R. and Mafnas, C. (2006). Biogeography of the marine actinomycete *Salinispora*.
 Finite Free Provided Action (2006). Environ Microbiol 8:1881-1888.
- 33. Jensen, P.R., Williams, P.G., Oh, D.C., Zeigler, L., and Fenical, W. (2007). Species-specific
 secondary metabolite production in marine actinomycetes of the genus *Salinispora*. Appl
 Environ Microbiol 73:1146-1152.
- Jiang,Z., Boyd,K.G., Mearns-Spragg,A., Adams,D.R., Wright,P.C., and Burgess,J.G. (2000).
 Two diketopiperazines and one halogenated phenol from cultures of the marine bacterium,
 Pseudoalteromonas luteoviolacea. Nat Prod Let 14:435-440.
- Kalesperis,G.S., Prahlad,K.V., and Lynch,D.L. (1975). Toxigenic studies with the antibiotic
 pigments from *Serratia marcescens*. Canadian Journal of Microbiology 21:213.
- Kalinovskaya,N.I., Ivanova,E.P., Alexeeva,Y.V., Gorshkova,N.M., Kuznetsova,T.A.,
 Dmitrenok,A.S., and Nicolau,D.V. (2004). Low-molecular-weight, biologically active
 compounds from marine *Pseudoalteromonas* species. Curr Microbiol 48:441-446.
- 19 37. Katoh,K., Misawa,K., Kuma,K., and Miyata,T. (2002). MAFFT: a novel method for rapid
 20 multiple sequence alignment based on fast Fourier transform. Nucleic Acids Res 30:3059 21 3066.
- 38. Laatsch,H. and Pudleiner,H. (1989). Marine bakterien, I: Synthese von pentabrompseudilin,
 einem phenylpyrrol aus *Alteromonas luteoviolaceus*. Liebigs Ann Chem863-881.
- 24 39. Lichstein,H.C. and Vandesand,V.F. (1945). Violacein, an antibiotic pigment produced by
 25 *Chromobacterium violaceum*. J Infect Dis 76:47-51.
- 40. Lovell,F.M. (1966). Structure of a bromine-rich marine antibiotic. J Am Chem Soc 88:4510&.
- 41. Månsson,M., Phipps,R.K., Gram,L., Munro,M.H., Larsen,T.O., and Nielsen,K.F. (2010).
 Explorative Solid-Phase Extraction (E-SPE) for Accelerated Microbial Natural Product
 Discovery. J Nat Prod 73:1126-1132.
- 42. Matz, C., Webb, J.S., Schupp, P.J., Phang, S.Y., Penesyan, A., Egan, S., Steinberg, P., and
 Kjelleberg, S. (2008). Marine biofilm bacteria evade eukaryotic predation by targeted chemical
 defense. PLoS ONE 3.
- 43. McCarthy,S.A., Johnson,R.M., and Kakimoto,D. (1994). Characterization of an antibiotic
 produced by *Alteromonas luteoviolacea* Gauthier 1982, 85 isolated from Kinko Bay, Japan. J
 Appl Microbiol 77:426-432.
- 44. Mearns-Spragg,A., Bregu,M., Boyd,K.G., and Burgess,J.G. (1998). Cross-species induction
 and enhancement of antimicrobial activity produced by epibiotic bacteria from marine algae
 and invertebrates, after exposure to terrestrial bacteria. Lett Appl Microbiol 27:142-146.

1 45. Newman, D.J. (2008). Natural products as leads to potential drugs: An old process or the new 2 hope for drug discovery? J Med Chem 51:2589-2599. 3 46. Nielsen, K.F. and Smedsgaard, J. (2003). Fungal metabolite screening: database of 474 4 mycotoxins and fungal metabolites for dereplication by standardised liquid chromatography-5 UV-mass spectrometry methodology. J Chromatogr A 1002:111-136. 6 47. Nielsen, K.F., Sumarah, M.W., Frisvad, J.C., and Miller, J.D. (2006). Production of metabolites from the Penicillium roqueforti complex. J Agricult Food Chem 54:3756-3763. 7 8 48. Penesyan, A., Marshall-Jones, Z., Holmstrom, C., Kjelleberg, S., and Egan, S. (2009). 9 Antimicrobial activity observed among cultured marine epiphytic bacteria reflects their 10 potential as a source of new drugs. FEMS Microbiol Ecol 69:113-124. 11 49. Rao, D., Webb, J.S., and Kjelleberg, S. (2005). Competitive interactions in mixed-species 12 biofilms containing the marine bacterium Pseudoalteromonas tunicata. Appl Environ 13 Microbiol 71:1729-1736. 14 50. Sanchez, S., Chavez, A., Forero, A., Garcia-Huante, Y., Romero, A., Sanchez, M., Rocha, D., 15 Sanchez, B., Avalos, M., Guzman-Trampe, S., Rodriguez-Sanoja, R., Langley, E., and Ruiz, B. 16 (2010). Carbon source regulation of antibiotic production. The Journal of Antibiotics 63:442-17 459. 18 51. Simmons, T.L., Coates, R.C., Clark, B.R., Engene, N., Gonzalez, D., Esquenazi, E., 19 Dorrestein, P.C., and Gerwick, W.H. (2008). Biosynthetic origin of natural products isolated 20 from marine microorganism-invertebrate assemblages. P Natl A Sci USA 105:4587-4594. 21 52. Skov, M.N., Pedersen, K., and Larsen, J.L. (1995). Comparison of pulsed-field gel 22 electrophoresis, ribotyping, and plasmid profiling for typing of Vibrio anguillarum serovar 23 O1. Appl Environ Microbiol 61:1540-1545. 53. Speitling, M., Smetanina, O.E., Kuznetsova, T.A., and Laatsch, H. (2007). Marine bacteria. 24 25 XXXV. Bromoalterochromides A and A ', unprecedented chromopeptides from a marine 26 Pseudoalteromonas maricaloris strain KMM 636. J Antibiot 60:36-42. 27 54. Stackebrandt, E., Frederiksen, W., Garrity, G.M., Grimont, P.A.D., Kampfer, P., Maiden, M.C.J., 28 Nesme, X., Rossello-Mora, R., Swings, J., Truper, H.G., Vauterin, L., Ward, A.C., and 29 Whitman, W.B. (2002). Report of the ad hoc committee for the re-evaluation of the species 30 definition in bacteriology. Int J Syst Evol Microbiol 52:1043-1047. 31 55. Sudek, S., Lopanik, N.B., Waggoner, L.E., Hildebrand, M., Anderson, C., Liu, H., Patel, A., 32 Sherman, D.H., and Haygood, M.G. (2006). Identification of the putative bryostatin polyketide 33 synthase gene cluster from "Candidatus Endobugula sertula", the uncultivated microbial 34 symbiont of the marine bryozoan Bugula neritina. J Nat Prod 70:67-74. 35 56. Talavera, G. and Castresana, J. (2007). Improvement of phylogenies after removing divergent 36 and ambiguously aligned blocks from protein sequence alignments. Syst Biol 56:564-577. 37 57. Tamura, K., Dudley, J., Nei, M., and Kumar, S. (2007). MEGA4: Molecular Evolutionary 38 Genetics Analysis (MEGA) software version 4.0. Mol Biol Evol 24:1596. 39 58. Violot, S., Aghajari, N., Czjzek, M., Feller, G., Sonan, G.K., Gouet, P., Gerday, C., Haser, R., and 40 Receveur-Brechot, V. (2005). Structure of a full length psychrophilic cellulase from

- 1 Pseudoalteromonas haloplanktis revealed by X-ray diffraction and small angle X-ray 2 scattering. J Mol Biol 348:1211-1224. 3 59. Werner, R.G. (1980). Uptake of indolmycin in gram-positive bacteria. Antimicrob Agents 4 Chemother 18:858. 5 60. Wietz, M., Schramm, A., Jørgensen, B., and Gram, L. (2010). Latitudinal patterns in the abundance of major marine bacterioplankton groups. Aquat Microb Ecol (Accepted). 6 7 61. Wratten, S.J., Wolfe, M.S., Andersen, R.J., and Faulkner, D.J. (1977). Antibiotic metabolites 8 from a marine Pseudomonad. Antimicrob Agents Chemother 11:411-414.
- 62. Zheng,L., Yan,X.J., Han,X.T., Chen,H.M., Lin,W., Lee,F.S.C., and Wang,X.R. (2006).
 Identification of norharman as the cytotoxic compound produced by the sponge
 (*Hymeniacidon perleve*)-associated marine bacterium *Pseudoalteromonas piscicida* and its
 apoptotic effect on cancer cells. Biotechnol Appl Biochem 44:135-142.
- 13
- 14

Figure 1: Isolation sites of 101 *Pseudoalteromonas* strains with each color code representing one of the 8 clusters as indicated in Table 1. Each circle represents the isolation of one or more strains belonging to the given cluster. White = cluster I, yellow = cluster II, grey = cluster III, green = cluster IV, violet = cluster V, red = cluster VI, orange = cluster VII, pink = cluster VII, and blue = non-clustered strain.

- 6
- 7



1 Figure 2: Phylogenetic tree based on 16S-rRNA gene sequences of the Galathea 3

Pseudoalteromonas strains. Sequences were aligned by MAFFT (default options) and the resulting
alignment was used to generate a Neighbor Joining tree in the MEGA4 software package (default
settings, 1000 bootstraps). *Salinispora arenicola* CNS-205, GenBank accession number CP000850,
GeneID: 5705939 was used as outgroup. Clusters containing 2 or more non-type strain sequences
were collapsed. The scale bar represents 0.02 amino acid substitutions per nucleotide position. •:
Nodes also occurring in minimum evolution and maximum likelihood trees.

- 8
- 9





0.02

Figure 3: Dendrogram from cluster analysis of detected peaks from HPLC-UV/VIS detection of compounds in the ethyl acetate extracted broth and biomass. Data was processed in NTSYSpc 2.20q, with SAHN clustering by UPGMA and simple distance measurement.



Figure 4: HPLC-UV/VIS profiles of the ethyl acetate extracted broth and biomass of four *P*. *luteoviolacea* with the tailing violacein peak at 14-15 min, and in the two upper traces pentabromopseudilin at 18 min in contrast to indolmycin at 11 min in the two lower traces.



16S-	No. of	strains	Related type	No. of stra	ains from	No. of strai	Inhibition of <i>Vibrio</i>	
rKNA	Diamontod	Non-	strain ¹	Surface complex	Water samples	Diamonted	Non-	by EtAc
cluster	I Ignienieu	pigmented		Surface samples	water samples	Tigmented	pigmented	extracts
Ι	1	37	P. agarivorans	16	22	0	0	0
II	5	0	P. aurantia	2	3	5	0	0
III	0	9	P. prydzensis	8	1	0	0	0
IV	3	3	P. phenolica	3	3	3	2	0
V	4	0	P. luteoviolacea	4	0	4	0	4
VI	9	0	P. rubra	9	0	9	0	0
VII	13	0	P. flavipulchra	13	0	13	0	0
VIII	15	0	P. ruthenica	14	1	15	0	15
S1727	0	1	P. mariniglutinosa	0	1	0	0	0
S3655	1	0	P. spongiae	1	0	0	0	0
Total	51	50		70	31	49	2	19

Table 1: Identity, antibacterial activity and pigmentation of *Pseudoalteromonas* strains from a global collection

3 ¹: The type strain which the majority of the strains in the cluster were most closely related to

Table 2: Ethyl acetate extractable secondary metabolites produced by pigmented
 1

Pseudoalteromonas strains 2

2
.)

Compound	$MI(Da)^1$	UV-max data	RT	Code
	MI (Du)	0 v max data	(min)	coue
Indolmycin ²	257		11.21	А
2-Pentyl-4-quinolinol ³	215		12.31	В
Novel mono brominated indole	280		13.78	С
Violacein ²	343		14.29	D
Unidentified	244	212nm (100%), 250nm (48%)	15.21	Е
2-n-Heptyl-(1H)-quinolin-4-one ³	243		15.47	F
Unidentified	NI	228nm (45%), 308 nm (100%)	15.70	G
Unidentified	386	< 200 nm	15.78	Н
Unidentified	316	286nm (100%)	15.80	Ι
Unidentified	NI^4	< 200 nm	15.81	J
Unidentified	NI	228nm (45%), 308nm (100%)	15.98	Κ
Unindentified	676	< 200nm	15.99	L
Prodigiosin ²	323		16.00	М
Unidentified	NI	< 200nm	16.26	Ν
Unidentified	NI	228nm (45%), 308nm (100%)	16.30	0
Bromoalterochromide A ³	843		16.49	Р
Novel bromoalterochromide, 2 bromine	921		16.60	Q
Novel bromoalterochromide, 1 bromine	857		17.10	R
Unidentified	333	310nm (100%)	17.20	S
2-n-Nonyl-(1H)-quinolin-4-one ³	271		17.96	Т
Nonyl-quinolin-one analogue ³	271		18.28	U
Unidentified	315	362nm (100%)	18.90	V
Unidentified	244	218nm (100%), 280nm (82%)	19.42	W
Unidentified	NI	218nm (100%), 288nm (82%)	19.78	Х
Unidentified	NI	250nm (100), 280nm (86)	19.92	Y
Pentabromopseudilin ³	549		22.65	Ζ

¹ Mono-isotopic mass 4

5

² validated reference standard used for verification.
³ Accurate mass and UV data fits data from Antibase2009

⁴ NI No ionization or MI could not be assigned using ESI⁺ and ESI⁻ 7

8

Table 3. Secondary metabolites produced by *Pseudoalteromonas* species clustered by 16S rRNA

gene similarity. Identification of peak by capital letter in table 2

16S cluster # strains none A B C D E G H I J K L N O P Q R S T V W X Y Z I 38 x x X X X Y Z X			Pea	ak at :	rete	ent	ior	n tii	ne p	ore	sei	nt in	n I	Pse	ud	oal	lte	ror	no	na	s s	tra	in/	org	ani	isn	n
1 38 x P.tetraodonis DSM9166 x P.prydzensis LMG21428 x I 1 x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 2 x x x 2 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x	16S cluster	# strains	none	A B	С	D	E	F	GΗ	Ι	J	K	L	Μ	Ν	0	Р	Q	R	S	Т	U	V	W	Х	Y	Ζ
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	Ι	38	Х																								
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	P.tetraodonis	DSM9166	х																								
II 1 x x x x 2 x x x x x x 1 x x x x x x 1 x x x x x x P. aurantia DSM6057 x x x x x P. citrea DSM8771 x x x x x II 9 x x x x x III 9 x x x x x III 9 x x x x x IV 3 x x x x x I x x x x x x VI 1 x x x x x x I x	P.prydzensis I	LMG21428	х																								
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	II	1		х				х													Х						
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		2		х				Х													Х	Х					
1 x x x x x x P. aurantia DSM6057 x x x x x x P.citrea DSM8771 x x x x x x III 9 x x x x x x III 9 x x x x x x 1 x x x x x x x 2 x x x x x x x V 2 x x x x x x x 1 x		1		х				Х										х				Х					
P. aurantia DSM6057xxxP.citrea DSM8771xxxIII9xxIV3xx21x x x2xxP.phenolica DSM21460xxV2xx2xxP.luteoviolacea ATCC33492xxXxxx1xxx		1		х				Х										х				х					
P.citrea DSM8771 x x x x III 9 x x x IV 3 x x x 2 1 x x x 2 x x x x P.phenolica DSM21460 x x x x Q x x x x x P.phenolica DSM21460 x x x x x Q x x x x x x P.luteoviolacea ATCC33492 x x x x x 1 x x x x x x 1 x x x x x x 1 x x x x x x 1 x x x x x x 1 x x x x x x 1 x x x x x x	P. aurantia D.	SM6057		х																	х						
III9xIV3x1x x x2x2xV2x2x2xN2VI11x1 <tr< td=""><td>P.citrea DSM</td><td>8771</td><td></td><td>х</td><td></td><td></td><td></td><td>Х</td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td>х</td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></tr<>	P.citrea DSM	8771		х				Х										х									
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	III	9	х																								
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	IV	3	х																								
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		1															х	х	х								
P.phenolica DSM21460 x V 2 x x 2 x x x P.luteoviolacea ATCC33492 x x VI 1 x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 3 x x x 1 x x x 1 x x x 3 x x x 1 x <td></td> <td>2</td> <td></td> <td>х</td>		2																									х
V2xx2xx2xxP.luteoviolacea ATCC33492xXXx1x1x1x1x1x1x1x1x1x1x1x1x1x1x1x1x1x1x3x1x3x1x2x1x3x1x1x2x1x3x1x1x2x1x3x1x1x1x1x3x1<	P.phenolica D	SM21460	х																								
2 x x P.luteoviolacea ATCC33492 x x VI 1 x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 3 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x x 1 x x	V	2				х																					Х
P.luteoviolacea ATCC33492 x x x VI 1 x x x x x 1 x x x x x x x 1 x x x x x x x 1 x x x x x x x 1 x x x x x x x x 1 x		2		х		х																					
VI 1 x 1 x x x x 1 x x x x x 1 x x x x x x 1 x x x x x x 1 x x x x x x 1 x x x x x x 1 x x x x x x 1 x x x x x x 1 x x x x x x x 1 x <td>P.luteoviolace</td> <td>a ATCC33</td> <td>492</td> <td></td> <td></td> <td>х</td> <td></td> <td>Х</td>	P.luteoviolace	a ATCC33	492			х																					Х
1 x	VI	1											х														
1 x <td></td> <td>1</td> <td></td> <td>Х</td> <td></td> <td></td> <td>х</td> <td>х</td> <td>х</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>х</td> <td></td>		1												Х			х	х	х							х	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$		1			х		Х			х														Х	Х		
1 x x x x 2 x 1 x 1 x x 1 x x P.rubra DSM6842 x VII 9 3 x x x 1 x x x 1 x x x P.flavipulchra LMG20361 x x x		1									х			х						х			х				
2 x 1 x 1 x x 1 x x P.rubra DSM6842 x VII 9 3 x x x 1 x x 1 x x 1 x x 1 x x 1 x x 1 x x		1							Х				х	Х	х												
1 x 1 x x P.rubra DSM6842 x VII 9 3 x x x 1 x x 1 x x P.flavipulchra LMG20361 x x x		2											х														
1 x x P.rubra DSM6842 x VII 9 x x x 3 x x x x 1 x x x x P.flavipulchra LMG20361 x x x		1												Х													
P.rubra DSM6842 x VII 9 x x x 3 x x x x 1 x x x x <i>P.flavipulchra</i> LMG20361 x x x		1											х	Х													
VII 9 x x x 3 x x x x 1 x x x x P.flavipulchra LMG20361 x x x	P.rubra DSM	6842											х														
3 x x x x 1 x x <i>P.flavipulchra</i> LMG20361 x x x	VII	9															х	х	x								
1 x x P.flavipulchra LMG20361 x x		3															х	х	х							х	
P.flavipulchra LMG20361 x x x		1		х				Х																			
	P.flavipulchra	1 LMG2036	51														х	х	х								
VIII 2 X	VIII	2	х																								
13 x x x		13							х			Х				х											
P.ruthenica LMG19699 x	P.ruthenica L	MG19699	х																								
S1727 1 x	S1727	1	х																								
S3655 1 x	S3655	1	x																								

Figure and table legends

Supplementary Figure 1: The phylogenetic tree also shown in figure 2, here with cluster I not collapsed to provide an overview of the large number of strains contained therein. The filled black triangle marks strain S3431, which in the BLAST analysis showed a low (97%) level of similarity to *Pseudoalteromonas* type strains.

Supplementary Table 1: Origin and anti-bacterial activity of *Pseudoalteromonas* strains included in this study. The cluster designation refers to figure 1. Sequence identity was determined using the BLAST algorithm. Where two hits are listed they scored equally in BLAST analysis.

Supplementary Figure 1: The phylogenetic tree also shown in figure 2, here with cluster I not collapsed to provide an overview of the large number of strains contained therein. The filled black triangle marks strain S3431, which in the BLAST analysis showed a low (97%) level of similarity to *Pseudoalteromonas* type strains.



Supplementary Table 1: Origin and anti-bacterial activity of *Psedoalteromonas* strains included in this study. The cluster designation refers to figure 1. Sequence identity was determined using the BLAST algorithm. Where two hits are listed they scored equally in the BLAST analysis.

		Latitude	Longitude	Water		%	Cluster			Inhibition	of V. an	guillarum
Strain	GenBank acc. #	+=N	+= E	or surface	BLAST type strain match	BLASI ident.	16S	Chem.	Pigment	Spot-assay	Sup	EtAc ex
S185	FJ457121	62.03815	-9.99592	W	P. arctica	98	1	А	-	-	-	-
S186	FJ457122	62.03815	-9.99592	W	P. distinct	98	1	А	-	-	-	-
S187	FJ457123	62.03815	-9.99592	W	P. prydzensis	97	3	А	-	-	-	-
S201	FJ457124	62.03815	-9.99592	W	P. agarivorans P. distincta	99	1	А	-	-	-	-
S326	FJ457125	62.2603	-51.6539	S	P. agarivorans	99	1	А	-	-	-	-
S327	FJ457126	62.2603	-51.6539	S	P. agarivorans	99	1	А	-	-	-	-
S407	FJ457127	66.90525	-53.2894	W	P. nigrifaciens	98	1	А	-	-	-	-
S408	FJ457128	66.90525	-53.2894	W	P. nigrifaciens	98	1	А	-	-	-	-
S409	FJ457129	66.90525	-53.2894	W	P. nigrifaciens	98	1	А	-	-	-	-
S410	FJ457130	66.90525	-53.2894	W	P. nigrifaciens	98	1	А	-	-	-	-
S411	FJ457131	66.90525	-53.2894	W	P. nigrifaciens	98	1	А	-	-	-	-
S554	FJ457133	42.6047	-29.9597	S	P. marina	99	1	А	-	-	-	-
S558	FJ457134	42.6047	-29.9597	S	P. aliena	99	1	А	-	-	-	-
S569	FJ457136	40.67227	-28.8374	W	P. tetraodonis	99	1	А	-	-	-	-
S570	FJ457137	40.67227	-28.8374	W	P. tetraodonis	99	1	А	-	-	-	-
S815	FJ457139	23.07925	-24.0523	W	P. agarivorans	99	1	А	-	-	-	-
S816	FJ457140	23.07925	-24.0523	W	P. agarivorans	99	1	А	-	-	-	-
S979	FJ457141	23.07918	-24.0542	W	P. agarivorans	99	1	А	-	-	-	-
S980	FJ457142	23.07918	-24.0542	W	P. agarivorans	99	1	А	-	-	-	-
S981	FJ457143	23.07918	-24.0542	W	P. agarivorans	99	1	А	-	-	-	-
S983	FJ457144	4.568133	-14.7097	W	P. agarivorans	99	1	А	-	-	-	-
S1093	FJ457145	4.570467	-1.72975	W	P. luteoviolacea	98	4	А	-	+	+	-
S1189	FJ457146	4.570467	-1.72975	W	P. phenolica	98	4	А	Brown	+	+	-

S1607	FJ457149	-38.407	66.3738	S	P. flavipulchra	99	7	G	Yellow	+	-	-
S1608	FJ457150	-38.407	66.3738	S	P. agarivorans	99	1	А	-	-	-	-
S1609	FJ457151	-38.407	66.3738	S	P. tetraodonis	100	1	А	-	-	-	-
S1610	FJ457152	-38.407	66.3738	S	P. agarivorans P. paragorgicola	99	1	А	-	-	-	-
S1612	FJ457153	-37.2168	72.7091	S	P. agarivorans P. paragorgicola	99	1	А	-	-	-	-
S1649	FJ457154	-35.3413	79.9641	S	P. prydzensis	98	3	А	-	-	-	-
S1650	FJ457155	-35.3413	79.9641	S	P. prydzensis	98	3	А	-	-	-	-
S1688	FJ457157	-31.4061	91.17758	W	P. elyakovii	99	1	А	-	-	-	-
S1690	FJ457158	-31.4061	91.17758	W	P. agarivorans	99	1	А	-	-	-	-
S1691	FJ457159	-31.4061	91.17758	W	P. agarivorans	99	1	А	-	-	-	-
S1692	FJ457160	-31.4061	91.17758	W	P. agarivorans	99	1	А	-	-	-	-
S1727	FJ457161	-19.7461	114.8573	W	P. mariniglutinosa	99	-	А	-	-	-	-
S1731	FJ457162	-19.7461	114.8573	W	P. agarivorans	99	1	А	-	-	-	-
S1845	FJ457163	-17.7746	121.8656	S	P. flavipulchra	99	7	G	Yellow	+	-	-
S1925	FJ457165	-17.0038	120.7788	S	P. flavipulchra	99	7	G	Yellow	+	-	-
S1941	FJ457166	-17.2706	121.1293	W	P. agarivorans	99	1	А	-	-	-	-
S1944	FJ457168	-17.0038	120.7788	S	P. rubra	99	6	В	Red	+	+	-
S1946	FJ457169	-17.0038	120.7788	S	P. rubra	99	6	G	Red	+	+	-
S1947	FJ457170	-17.0038	120.7788	S	P. flavipulchra	99	7	G	Yellow	+	-	-
S1948	FJ457171	-17.0038	120.7788	S	P. flavipulchra	99	7	G	Yellow	+	-	-
S2040	FJ457173	-16.0604	119.3541	S	P. flavipulchra	99	7	G	Yellow	+	-	-
S2047	FJ457174	-16.0604	119.3541	S	P. flavipulchra	99	7	G	Yellow	+	-	-
S2048	FJ457175	-16.0604	119.3541	S	P. piscicida	99	7	G	Yellow	+	-	-
S2049	FJ457176	-16.0604	119.3541	S	P. piscicida	99	7	G	Yellow	+	-	-
S2050	FJ457177	-16.0604	119.3541	S	P. piscicida	99	7	G	Yellow	+	-	-
S2053	FJ457179	-16.0604	119.3541	S	P. piscicida	99	7	G	Yellow	+	-	-
S2231	FJ457180	-33.3267	127.6573	S	P. aurantia P. citrea	99	2	F	Pale yellow	+	+	-
S2233	FJ457181	-33.3267	127.6573	S	P. citrea	99	2	F	Pale yellow	+	+	-
S2471	FJ457184	-22.9636	153.9486	S	P. rubra	99	6	-	Red	+	+	-

S2472	FJ457185	-22.9636	153.9486	S	P. rubra	99	6	-	Red	+	+	-
S2599	FJ457186	-10.3454	157.7956	S	P. rubra	99	6	-	Red	+	+	-
S2607	FJ457187	-10.3454	157.7956	S	P. luteoviolacea	99	5	С	Purple	+	+	+
S2676	FJ457188	-8.0692	155.8781	S	P. rubra	99	6	В	Red	+	+	-
S2678	FJ457189	-8.0692	155.8781	S	P. rubra	99	6	В	Red	+	+	-
S2717	FJ457190	-7.8244	156.0689	W	P. ruthenica	99	8	А	Brown	+	-	+
S2721	FJ457191	-8.1005	156.8451	S	P. prydzensis	97	3	А	-	-	-	-
S2722	FJ457192	-8.1005	156.8451	S	P. rubra	99	6	-	Red	+	+	-
S2724	FJ457193	-8.1005	156.8451	S	P. rubra P. luteoviolacea	97	4	G	Brown	+	-	-
S2755	FJ457196	-9.108	156.8595	S	P. flavipulchra	99	7	G	Yellow	+	-	-
S2756	FJ457197	-15.2329	156.665	S	P. ruthenica	99	8	Е	Brown	+	-	+
S2893	FJ457198	-48.915	178.1056	S	P. agarivorans	99	1	А	-	-	-	-
S2897	FJ457199	-48.915	178.1056	S	P. ruthenica	99	8	Е	Brown	+	-	+
S2898	FJ457200	-48.915	178.1056	S	P. ruthenica	99	8	Е	Brown	+	-	+
S2899	FJ457201	-48.915	178.1056	S	P. ruthenica	99	8	Е	Brown	+	-	+
S3136	FJ457202	-66.7706	-76.4383	S	P. ruthenica	99	8	Е	Brown	+	-	+
S3137	FJ457203	-66.7706	-76.4383	S	P. ruthenica	99	8	Е	Brown	+	-	+
S3138	FJ457204	-66.7706	-76.4383	S	P. ruthenica	99	8	Е	Brown	+	-	+
S3173	FJ457205	-67.4876	-89.1285	S	P. tetraodonis	98	1	А	-	-	-	-
S3177	FJ457206	-67.4876	-89.1285	S	P. ruthenica	99	8	Е	Brown	+	-	+
S3178	FJ457207	-67.4876	-89.1285	S	P. carrageenovora	99	1	А	-	-	-	-
S3244	FJ457208	-67.4876	-89.1285	S	P. ruthenica	99	8	Е	Brown	+	-	+
S3245	FJ457209	-67.4876	-89.1285	S	P. ruthenica	99	8	А	Brown	+	-	+
S3257	FJ457210	-64.0947	-62.8228	S	P. ruthenica	99	8	Е	Brown	+	-	+
S3258	FJ457211	-64.0947	-62.8228	S	P. ruthenica	99	8	Е	Brown	+	-	+
S3260	FJ457212	-64.0947	-62.8228	S	P. agarivorans	99	1	А	-	-	-	-
S3431	FJ457214	-50.4498	-74.8912	S	P. arctica	97	1	А	Black	-	-	-
S3655	FJ457216	-20.0568	-70.755	S	P. spongiae	99	-	А	Orange	-	-	-
S3663	FJ457217	-20.0568	-70.755	S	P. phenolica	97	4	С	-	+	-	-
S3776	FJ457222	-14.2295	-76.6073	S	P. agarivorans	99	1	А	-	-	-	-
S3785	FJ457223	-14.2295	-76.6073	S	P. agarivorans	99	1	А	-	-	-	-

S3788	FJ457224	-17.0857	-72.419	W	P. aurantia	98	2	F	Pale yellow	+	+	-
S3790	FJ457225	-17.0857	-72.419	W	P. aurantia	98	2	F	Pale yellow	+	+	-
S3895	FJ457226	-14.1631	-77.4286	W	P. aurantia	98	2	F	Pale yellow	+	+	-
S3898	FJ457228	-13.818	-76.7648	W	P. phenolica	97	4	А	-	-	-	-
S3944	FJ457229	-5.3492	-81.4284	S	P. mariniglutinosa	98	3	А	-	-	-	-
S4047	FJ457230	2.9817	-86.6892	S	P. luteoviolacea	98	5	D	Purple	+	+	+
S4048	FJ457231	2.9817	-86.6892	S	P. phenolica	98	4	С	Brown	+	-	-
S4054	FJ457234	2.9817	-86.6892	S	P. luteoviolacea	99	5	D	Purple	+	+	+
S4059	FJ457237	2.9817	-86.6892	S	P. rubra	99	6	В	Red	+	+	-
S4060	FJ457238	2.9817	-86.6892	S	P. luteoviolacea	98	5	С	Purple	+	+	+
S4382	FJ457239	26.5042	-66.9964	S	P. ruthenica	99	8	E	Brown	+	-	+
S4388	FJ457240	26.5042	-66.9964	S	P. ruthenica	99	8	E	Brown	+	-	+
S4389	FJ457241	26.5042	-66.9964	S	P. mariniglutinosa	97	3	А	-	-	-	-
S4488	FJ457243	24.9963	-67.0246	S	P. prydzensis	97	3	А	-	-	-	-
S4491	FJ457244	24.9963	-67.0246	S	P. prydzensis	97	3	А	-	-	-	-
S4492	FJ457245	24.9963	-67.0246	S	P. mariniglutinosa	98	3	А	-	-	-	-
S4498	FJ457247	24.9963	-67.0246	S	P. flavipulchra P. maricaloris	99	7	F	-	+	-	-
S4741	FJ457248	58.8041	-3.0564	S	P. elyakovii	99	1	А	-	-	-	-

Supplementary figure S1: The phylogenetic tree also shown in figure 2, here with cluster I not collapsed to provide an overview of the large number of strains contained therein. The filled black triangle marks strain S3431, which in the BLAST analysis showed a low (97%) level of similarity to *Pseudoalteromonas* type strains.



0.02