



Molecular-like Redox Activity and Size-dependent Electrocatalysis of Inorganic Hybrid Nanoparticles

Chi, Qijin; Zhu, Nan; Ulstrup, Jens

Publication date:
2011

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Chi, Q., Zhu, N., & Ulstrup, J. (2011). *Molecular-like Redox Activity and Size-dependent Electrocatalysis of Inorganic Hybrid Nanoparticles*. Abstract from The 62nd Annual Meeting of the International Society of Electrochemistry, Niigata, Japan.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Molecule-like Redox Activity and Size-dependent Electrocatalysis of Inorganic Hybrid Nanoparticles

Qijin Chi,* Nan Zhu and Jens Ulstrup

Department of Chemistry and NanoDTU, Technical University of Denmark, DK-2800 Kgs. Lyngby, Denmark. (*E-mail: cq@kemi.dtu.dk)

The development of low-cost, robust and high-efficient nanoscale electrocatalysts is arguably a dream approach to the use of nanomaterials as key building blocks in design and construction of chemical and biological sensing devices as well as fuel cells. Electroactive nanoparticles are a type of nanoparticles that have intrinsic electroactivity. One of representative examples is the nanoparticles composed of *Prussian Blue* or/and its analogues. This type of nanoparticles has advantages over classic electrocatalysts in several regards. In this communication, we present some of our recent efforts on synthesis, characterization, and electrocatalytic function of *Prussian Blue* nanoparticles (PBNPs). Molecule-like redox activity and size-dependent electrocatalysis are clearly revealed, which could offer crucial clues for further optimization of design of nanoscale electrocatalysts and their applications in sensors.




Fig. 1 Cyclic voltammograms of PBNPs immobilized on an Au(111) surface. Scan rates 0.1 to 1.0 V s⁻¹.




Fig. 2 Distance-dependent ET kinetics clearly showing the feature of tunnelling mechanism.

Acknowledgements: This project is supported in part by the Danish Research Council for Technology and Production Sciences (to Q.C and J.U.).