



## Activity and stability of feruloyl esterase A from *Aspergillus niger* in ionic liquid systems

Zeuner, Birgitte; Riisager, Anders; Meyer, Anne S.

*Publication date:*  
2011

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Zeuner, B., Riisager, A., & Meyer, A. S. (2011). *Activity and stability of feruloyl esterase A from Aspergillus niger in ionic liquid systems*. Abstract from 6th Danish Conference on Biotechnology and Molecular Biology, Vejle, Denmark.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Activity and stability of feruloyl esterase A from *Aspergillus niger* in ionic liquid systems

Birgitte Zeuner<sup>1\*</sup>, Anders Riisager<sup>2</sup>, Anne S. Meyer<sup>1</sup>

<sup>1</sup>Center for BioProcess Engineering, Department of Chemical and Biochemical Engineering, Technical University of Denmark, DK-2800 Lyngby, Denmark.

<sup>2</sup>Centre for Catalysis and Sustainable Chemistry, Department of Chemistry, Technical University of Denmark, DK-2800, Denmark.

\*biz@kt.dtu.dk

Feruloyl esterases (FAEs; EC 3.1.1.73) are accessory plant cell wall-degrading enzymes, which catalyse the hydrolysis of the ester bond between ferulic acid and the monosaccharide to which it is covalently linked. FAEs can however also be brought to catalyse the (trans)esterification reaction in solvents that favour synthesis over hydrolysis, i.e. systems with low water content such as organic solvents or ionic liquids (ILs). The esterification of sinapic acid with glycerol catalysed by FAE A from *Aspergillus niger* (AnFaeA) in a series of ILs containing 15% (v/v) buffer showed that AnFaeA stability – and hence activity – was highly dependent on the anion nature: AnFaeA was stable and active for more than 2 hours in [PF<sub>6</sub>]<sup>-</sup>-based ILs, but rapidly lost activity in [BF<sub>4</sub>]<sup>-</sup>-based systems. This effect can be explained in terms of hydrogen bonding capacity of the two anions: As predicted by the quantum chemistry-based COSMO-RS method, [BF<sub>4</sub>]<sup>-</sup> has a tendency to form hydrogen bonds and thus interfere with the secondary structure of the enzyme, while [PF<sub>6</sub>]<sup>-</sup> is unlikely to form hydrogen bonds and therefore does not cause denaturation of the enzyme. Similar results have been obtained for lipases [1], but this is the first report on FAE stability in ILs [2]. COSMO-RS, which is now widely used for solvent screening in the complex IL systems [3], may be a valuable tool for fast enzyme stability predictions and/or solvent screening in the future.

### References:

[1]: Zhao, H. J. *Chem. Technol. Biotechnol.*, **2010**, 85, 891-907.

[2]: Zeuner, B., Ståhlberg, T., van Buu, O.N., Kunov-Kruse, A.J., Riisager, A., and Meyer, A.S. *Green Chem.*, **2011** (accepted April 2011).

[3]: Klamt, A., Eckert, F., and Arlt, W. *Annu. Rev. Chem. Biomol. Eng.*, **2010**, 1, 101-122.