



## Radioactivity in the Risø District January-June 2011

**Nielsen, Sven Poul; Andersson, Kasper Grann; Miller, Arne**

*Publication date:*  
2011

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

### Citation (APA):

Nielsen, S. P., Andersson, K. G., & Miller, A. (2011). *Radioactivity in the Risø District January-June 2011*. Danmarks Tekniske Universitet, Risø Nationallaboratoriet for Bæredygtig Energi. Denmark. Forskningscenter Risoe. Risoe-R No. 1800(EN)

---


### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Radioactivity in the Risø District January-June 2011



Sven P. Nielsen, Kasper G. Andersson and Arne Miller  
Risø-R-1800(EN)  
December 2011



**Author:** Sven P. Nielsen, Kasper G. Andersson and Arne Miller  
**Title:** Radioactivity in the Risø District January-June 2011  
**Division:** Radiation Research

Risø-R-1800(EN)  
December 2011

**Abstract (max. 2000 char.):** The environmental surveillance of the Risø environment was continued in January-June 2011. The mean concentrations in air were:  $7.97 \pm 20.7 \text{ } \mu\text{Bq m}^{-3}$  of  $^{137}\text{Cs}$ ,  $2.82 \pm 1.18 \text{ mBq m}^{-3}$  of  $^7\text{Be}$  and  $0.16 \pm 0.14 \text{ mBq m}^{-3}$  of  $^{210}\text{Pb}$  ( $\pm 1 \text{ S.D.}; N = 26$ ). The unusually high mean value and standard deviation over the period for  $^{137}\text{Cs}$  were due to highly enhanced concentrations in a period in March-May caused by the releases from the Fukushima accident in Japan. This is also reflected in the values for  $^{134}\text{Cs}$  and  $^{131}\text{I}$ . The depositions by precipitation at Risø in the first half of 2011 were:  $0.527 \text{ Bq m}^{-2}$  of  $^{137}\text{Cs}$  (correspondingly unusually high),  $481 \text{ Bq m}^{-2}$  of  $^7\text{Be}$ ,  $40.4 \text{ Bq m}^{-2}$  of  $^{210}\text{Pb}$  and  $< 1.3 \text{ kBq m}^{-2}$  of  $^3\text{H}$ . The average background dose rate (TLD) at Risø (Zone I) was  $64 \text{ nSv h}^{-1}$  compared with  $50 \pm 11 \text{ nSv h}^{-1}$  ( $\pm 1 \text{ S.D.}; N = 4$ ) in the four zones around Risø.

**ISSN 0106-2840**  
**ISBN 978-87-550-3951-3**

**Contract no.:**

**Group's own reg. no.:**  
1400103-07

**Sponsorship:**

**Cover :**

**Pages: 24**  
**Tables: 14**  
**References:**

Information Service Department  
Risø National Laboratory for  
Sustainable Energy  
Technical University of Denmark  
P.O.Box 49  
DK-4000 Roskilde  
Denmark  
Telephone +45 46774005  
[bibl@risoe.dtu.dk](mailto:bibl@risoe.dtu.dk)  
Fax +45 46774013  
[www.risoe.dtu.dk](http://www.risoe.dtu.dk)

# Contents

Table 1.	Radionuclides in air	5
Table 2.1.	Radionuclides in precipitation	6
Table 2.2.	Radionuclides in precipitation	6
Table 2.3.	Tritium in precipitation	7
Table 2.4.	Tritium in precipitation	7
Table 3.1.	Radionuclides in sediment samples	8
Table 4.1.	Radionuclides in seawater	8
Table 4.2.	Tritium in seawater	8
Table 5.1.	Radionuclides in grass	9
Table 5.2.	Radionuclides in sea plants	10
Table 7.1.	Waste water	11
Table 8.1.	Background dose rates around the border of Risø (TLD)	12
Table 8.2.	Background dose rates around Risø (TLD)	13
Table 8.3.	Terrestrial dose rates at the Risø zones (NaI(Tl) detector)	14
Fig. 1.	Map of Risø	15
Fig. 1.1.	Caesium-137 in air	16
Fig. 1.2.	Beryllium-7 and lead-210 in air	16
Fig. 2.3.1	Tritium in precipitation (1 m <sup>2</sup> rain collector)	17
Fig. 2.3.2	Tritium in precipitation (10 m <sup>2</sup> rain collector)	17
Fig. 3.1	Caesium-137 in sediment samples	18
Fig. 4.1	Caesium-137 in seawater	19
Fig. 4.2	Tritium in seawater	19
Fig. 7.1	Total-beta radioactivity in waste water	20
Fig. 8.1.	Map of Risø with locations for TLD measurements	21
Fig. 8.2.	The environment of Risø	22



*Table 1. Radionuclides in ground level air collected at Risø (cf. Figs. 1, 1.1 and 1.2), January - June 2011. (Unit:  $\mu\text{Bq m}^{-3}$ )*

Date		$^7\text{Be}$	$^{137}\text{Cs}$	$^{134}\text{Cs}$	$^{131}\text{I}$	$^{210}\text{Pb}$
27-Dec-10	-	03-Jan-11	1229	0.610	< 0.42	< 0.88
03-Jan-11	-	10-Jan-11	2296	0.445	< 0.36	< 0.88
10-Jan-11	-	17-Jan-11	1834	0.216	< 0.42	< 0.97
17-Jan-11	-	24-Jan-11	1173	0.354	< 0.35	< 0.77
24-Jan-11	-	31-Jan-11	1778	0.619	< 0.31	< 0.78
31-Jan-11	-	07-Feb-11	3062	0.204	< 0.45	< 1.20
07-Feb-11	-	14-Feb-11	1935	0.211	< 0.26	< 0.68
14-Feb-11	-	21-Feb-11	3665	1.132	< 0.61	< 1.57
21-Feb-11	-	28-Feb-11	2585	1.265	< 0.24	< 0.68
28-Feb-11	-	07-Mar-11	1735	0.704	< 0.42	< 1.01
07-Mar-11	-	14-Mar-11	1516	0.219	< 0.43	< 1.05
14-Mar-11	-	21-Mar-11	2167	0.515	< 0.89	< 1.75
21-Mar-11	-	28-Mar-11	2879	5.516	33.3	379
28-Mar-11	-	04-Apr-11	1925	100.218	76.5	484
04-Apr-11	-	11-Apr-11	1482	35.182	35.7	183
11-Apr-11	-	18-Apr-11	2750	28.548	28.4	84.2
18-Apr-11	-	26-Apr-11	4528	11.755	13.0	20.9
26-Apr-11	-	02-May-11	4027	11.733	13.0	14.8
02-May-11	-	09-May-11	3584	4.775	5.87	4.7
09-May-11	-	16-May-11	2556	0.841	< 0.72	< 1.7
16-May-11	-	23-May-11	2757	0.525	< 0.84	< 1.9
23-May-11	-	30-May-11	5275	0.599	< 0.88	< 2.3
30-May-11	-	06-Jun-11	3272	0.333	< 0.95	< 2.3
06-Jun-11	-	14-Jun-11	4435	0.294	< 0.74	< 2.0
14-Jun-11	-	20-Jun-11	4426	0.196	< 0.43	< 1.1
20-Jun-11	-	27-Jun-11	4597	0.214	< 0.73	< 1.7
Mean		2826	7.97			162
SD		1178	20.8			136

*Table 2.1. Radionuclides in precipitation in the 10 m<sup>2</sup> rain collector at Risø (cf. Fig. 1), January - June 2011. (Unit: Bq m<sup>-3</sup>)*

Month	<sup>7</sup> Be	<sup>137</sup> Cs	<sup>210</sup> Pb
January	1155	0.149	164
February	602	-	6
March	1749	5.852	208
April	1456	6.967	131
May	4614	3.790	412
June	2507	0.381	91

*Table 2.2. Radionuclides in precipitation in the 10 m<sup>2</sup> rain collector at Risø (cf. Fig. 1), January - June 2011. (Unit: Bq m<sup>-2</sup>)*

Month	Precipitation (m)	<sup>7</sup> Be	<sup>137</sup> Cs	<sup>210</sup> Pb
January	0.048	55.8	0.0072	7.9
February	0.028	16.7	-	0.2
March	0.031	55.1	0.1843	6.6
April	0.022	31.3	0.1498	2.8
May	0.045	210.0	0.1722	18.8
June	0.045	112.0	0.0136	4.1
Sum	0.219	480.9	0.5271	40.4

*Table 2.3. Tritium in precipitation collected at Risø (cf. Figs. 1, 2.3.1 and 2.3.2). January - June 2011. (Unit: kBq m<sup>-3</sup>)*

Month	1 m <sup>2</sup> rain collector*	10 m <sup>2</sup> rain collector*
January	< 1.7	< 2.3
February	< 1.7	< 1.7
March	2.7	3.1
April	3.0	20.0
May	< 1.9	5.5
June	3.2	6.5
Double determinations*.		

*Table 2.4. Tritium in precipitation collected at Risø (cf. Fig. 1). January - June 2011. (Unit: kBq m<sup>-2</sup>)*

Month	Precipitation (m)	1 m <sup>2</sup> rain collector	10 m <sup>2</sup> rain collector
January	0.048	< 0.082	< 0.110
February	0.028	< 0.048	< 0.048
March	0.031	0.084	0.096
April	0.022	0.066	0.440
May	0.045	< 0.086	0.248
June	0.045	0.144	0.293
Sum	0.219	< 0.508	< 1.234

*Table 3.1. Radionuclides in sediment samples collected at Bolund in Roskilde Fjord.(cf. Fig. 3.1) January - June 2011. (Unit: Bq kg<sup>-1</sup> dry)*

No samples in this period.

*Table 4.1. Radionuclides in seawater collected in Roskilde Fjord (cf. Fig. 4.1) January - June 2011. (Unit: Bq m<sup>-3</sup>)*

No samples in this period.

*Table 4.2. Tritium in seawater collected in Roskilde Fjord (Risø pier) (cf. Fig. 4.2) January - June 2011.*

Month	kBq m <sup>-3</sup>
January	< 1.7 *
February	2.7 *
March	1.9 *
April	< 1.7 *
May	< 1.7 *
June	2.7 *

\* Double determinations

*Table 5.1. Radionuclides in grass (\* snow) collected at Risø (near the Waste Treatment Station (cf. Fig. 1)), January - June 2011. (\*\*Measured on bulked ash samples)*

Week no. or month	Date	K (g kg <sup>-1</sup> fresh)	<sup>137</sup> Cs (Bq kg <sup>-1</sup> fresh)	<sup>137</sup> Cs (Bq m <sup>-2</sup> )
1	3 January*	<0.1	<0.2	
2	10 January*	<0.1	<0.2	
3	17 January	3.3	<0.4	
4	24 January	2.5	<0.4	
5	31 January*	<0.1	<0.3	
6	7 February	1.8	<0.4	
7	14 February	1.6	<0.9	
8	21 February	2.1	<0.9	
9	28 February	2.0	<1.0	
10	7 March	2.5	<0.4	
11	14 March	1.0	<0.4	
12	21 March	1.8	<1.0	
13	28 March	2.2	<0.2	
14	4 April	3.7	<1.0	
15	11 April	5.1	<1.0	
16	18 April	5.5	<0.7	
17	26 April	5.9	<0.9	
18	2 May	4.4	<0.6	
19	9 May	5.2	<0.5	
20	16 May	4.9	<0.5	
21	23 May	4.3	<0.4	
22	30 May	4.5	<0.4	
23	6 June	5.8	<0.4	
+25	20 June	4.8	<0.5	
**January		2.8	0.070	0.037
**February		1.7	0.139	0.043
**March		2.5	0.312	0.076
**April		5.1	0.855	0.166
**May		5.0	0.187	0.097
**June		5.3	0.068	0.059

<sup>+</sup> Note: After week 23 of 2011, the sampling frequency is fortnightly.

*Table 5.2. Radionuclides in *Fucus vesiculosus* collected at Bolund in Roskilde Fjord. January - June 2011. (Unit: Bq kg<sup>-1</sup> dry)*

No samples in this period.

*Table 7.1. Waste water collected at Risø (cf. Fig. 1), January - June 2011.*

Week number	eqv. mg KCl l <sup>-1</sup>	<sup>137</sup> Cs (Bq m <sup>-3</sup> )	<sup>131</sup> I (Bq m <sup>-3</sup> )	<sup>226</sup> Ra (Bq m <sup>-3</sup> )
1	48	<128	<135	<278
2	44	<115	<117	<234
3	22	<118	<116	<214
4	45	<121	<114	<221
5	37	<71	<73	<140
6	36	<105	<111	<209
7	45	<106	<106	<228
8	39	<105	<109	<206
9	57	<107	<112	<215
10	67	<109	<110	<214
11	74	<105	<108	<210
12	83	<112	<115	<217
13	85	<118	<121	<229
14	57	<109	<115	<230
15	67	<111	<116	<209
16	78	<74	<62	347
17	60	<116	<119	<227
18	82	<113	<119	<223
19	88	<92	<101	<187
20	85	<120	<117	<215
21	86	<111	<111	<213
22	78	<115	<118	<220
23	79	<112	<114	<225
24	88	<104	<106	<198
25	95	<65	<77	<131
26	97	<69	<67	312
Mean	66.2			
SD	21.2			

*Table 8.1. Background dose rates around the border of Risø (cf. Fig. 8.1) measured with thermoluminescence dosimeters (TLD) in the period November 2010 – April 2011. (Results are normalized to nSv h<sup>-1</sup>)*


Location	nSv h <sup>-1</sup>
1	55
2	43
3	39
4	46
5	51
6	63
Mean	50

*Table 8.2. Background dose rates around Risø (cf. Fig. 8.2 and Fig. 1) measured with thermoluminescence dosimeters (TLD) in the period November 2010 – April 2011. (Results are normalized to nSv h<sup>-1</sup>)*

Risø zone	Location	nSv h <sup>-1</sup>
I	1	42
I	2	53
I	3	107
I	4	57
I	5	61
Mean		64
II	P1	53
II	P2	53
II	P3	34
II	P4	59
Mean		50
III	P1	45
III	P2	51
III	P3	42
Mean		46
IV	P1	41
IV	P2	41
IV	P3	48
IV	P4	57
IV	P5	50
IV	P6	46
IV	P7	61
Mean		49
V	P1	53
V	P2	49
V	P3	64
V	P4	39
V	P5	57
V	P6	36
V	P7	45
V	P8	62
V	P9	46
V	P10	55
Mean		51

*Table 8.3. Terrestrial dose rates at the Risø zones (cf. Fig. 8.2 and Fig. 1) January - June 2011. Measured with a NaI(Tl) detector. (Unit: nSv h<sup>-1</sup>)*

Risø zone	Location	January	April
I	P1	38	34
I	P2	45	41
I	P3	344	311
I	P4	40	38
I	P5	45	48
Mean		102	94
II	P1	35	45
II	P2	34	39
II	P3	33	34
II	P4	34	35
Mean		34	38
III	P1		46
III	P2		47
III	P3		41
Mean			45
IV	P1		35
IV	P2		46
IV	P3		42
IV	P4		38
IV	P5		36
IV	P6		37
IV	P7		40
Mean			39
V	P1		33
V	P2		43
V	P3		49
V	P4		44
V	P5		43
V	P6		41
V	P7		34
V	P7a		36
V	P8		45
V	P9		44
V	P10		34
Mean			41



*Fig. 1. Locations for measurements of gamma-background radiation Zone I and II  
(cf. Tables 8.2 and 8.3)*




Fig. 1.1. Caesium-137 in ground level air collected at Risø in January-June 2011.  
(Unit:  $\mu\text{Bq m}^{-3}$ )




Fig. 1.2. Beryllium-7 and Lead-210 in ground level air collected at Risø in January-June 2011. (Unit:  $\mu\text{Bq m}^{-3}$ )




Fig. 2.3.1. Tritium in precipitation collected at Risø (  $1 \text{ m}^2$  rain collector )  
1980 - 2011. (Unit:  $\text{kBq m}^{-3}$ ; DL = detection limit )




Fig. 2.3.2. Tritium in precipitation collected at Risø (  $10 \text{ m}^2$  rain collector )  
1980 - 2011. (Unit:  $\text{kBq m}^{-3}$ ; DL = detection limit )






Fig. 3.1. Caesium-137 in sediment samples collected at Bolund in Roskilde Fjord. 1980 – 2011. (Unit:  $\text{Bq kg}^{-1}$  dry matter)



*Fig. 4.1. Caesium-137 in seawater collected in Roskilde Fjord 1980 - 2011.  
(Unit:  $\text{Bq m}^{-3}$ )*



*Fig. 4.2. Tritium in seawater collected in Roskilde Fjord 1980 - 2011.  
(Unit:  $\text{kBq m}^{-3}$ ; DL = detection limit)*




Fig. 7.1. Total-beta radioactivity in waste water collected at Risø 1994 - 2011.  
(Unit: eqv. mg KCl l<sup>-1</sup>)





Fig. 8.1. Locations (1-6) for TLD measurements around the border of Risø (cf. Table 8.1).



*Fig. 8.2. Locations for measurements of background radiation around Risø in Zones III, IV and V.*



Risø DTU is the National Laboratory for Sustainable Energy. Our research focuses on development of energy technologies and systems with minimal effect on climate, and contributes to innovation, education and policy. Risø has large experimental facilities and interdisciplinary research environments, and includes the national centre for nuclear technologies.

---

**Risø DTU**  
**National Laboratory for Sustainable Energy**  
**Technical University of Denmark**

Frederiksborgvej 399  
PO Box 49  
DK-4000 Roskilde  
Denmark  
Phone +45 4677 4677  
Fax +45 4677 5688

[www.risoe.dtu.dk](http://www.risoe.dtu.dk)