



Food and soil-borne *Penicillia* in Arctic environments: Chemical diversity

Frisvad, Jens Christian

Publication date:
2011

Document Version
Early version, also known as pre-print

[Link back to DTU Orbit](#)

Citation (APA):
Frisvad, J. C. (2011). *Food and soil-borne Penicillia in Arctic environments: Chemical diversity*. Abstract from Polar and Alpine Microbiology, Ljubljana, Slovenia. <http://www.polaralpinemicrobiology.com/>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Food and soil-borne *Penicillia* in Arctic environments: chemical diversity

Jens Christian Frisvad

^b*Department of Systems Biology, Technical University of Denmark, Søltofts Plads, Building 221, DK-2800 Kgs. Lyngby, Denmark;*

E-mail: jcf @bio.dtu.dk

Keywords: extrolites, secondary metabolites, biodiversity, Greenland

Penicillia are very common inhabitants of cold environments, including arctic soil, plants, animals, and foods. We have investigated the mycobiota of Greenland inland ice and soil, and found a very unique and pronounced diversity among the *Penicillia*. Nearly all species were new to science, and only few well known *Penicillium* species were found. Most of the soil-borne *Penicillia* grew very poorly, if at all, at 25°C and could not sporulate at that temperature. Only one species of this kind has yet been described, *Penicillium jamesonlandense*, but there are at least 20 new species of this kind. The species found in inland ice were both of the soil-borne type, and *Penicillia* that grow and sporulate well at 25°C. The latter group of *Penicillia* have been found earlier in refrigerated foods, including *P. nordicum*, and in glacier ice and melting water from Svalbard (se Sonjak et al., this conference). This “food-borne group” of arctic fungi also contained some new species, but not as many as in arctic soil. The chemical diversity of the *Penicillium* species was remarkably high and in most cases even larger than the chemical diversity of *Penicillia* in the tropics. Several new secondary metabolites were found and known secondary metabolites with pronounced bioactivity were also found in most species, such as chrysogine, compactin, cyclopenin, kojic acid, mycophenolic acid, penicillic acid, pseurotins, pyripyropens etc.