



The link between atmospheric radicals and newly formed particles at a spruce forest site in Germany

Bonn, B.; Bourtsoukidis, E.; Sun, T. S.; Bingemer, H.; Rondo, L.; Javed, U.; Li, J.; Axinte, R.; Li, X.; Brauers, T.

Total number of authors:
15

Published in:
Atmospheric Chemistry and Physics Discussions

Link to article, DOI:
[10.5194/acpd-13-27501-2013](https://doi.org/10.5194/acpd-13-27501-2013)

Publication date:
2013

[Link back to DTU Orbit](#)

Citation (APA):

Bonn, B., Bourtsoukidis, E., Sun, T. S., Bingemer, H., Rondo, L., Javed, U., Li, J., Axinte, R., Li, X., Brauers, T., Sonderfeld, H., Koppmann, R., Sogachev, A., Jacobi, S., & Spracklen, D. V. (2013). The link between atmospheric radicals and newly formed particles at a spruce forest site in Germany. *Atmospheric Chemistry and Physics Discussions*, 13(10), 27501-27560. <https://doi.org/10.5194/acpd-13-27501-2013>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

The link between atmospheric radicals and newly formed particles at a spruce forest site in Germany

B. Bonn^{1,*}, E. Bourtsoukidis^{1,2}, T. S. Sun^{1,2}, H. Bingemer¹, L. Rondo¹, U. Javed², J. Li², R. Axinte², X. Li³, T. Brauers³, H. Sonderfeld⁴, R. Koppmann⁴, A. Sogachev⁵, S. Jacobi⁶, and D. V. Spracklen⁷

¹Institute for Atmospheric and Environmental Sciences, Goethe University, Frankfurt, Germany

²Air Chemistry Department, Max-Planck-Institute for Chemistry, Mainz, Germany

³Institute for Energy and Climate Research, IEK-8, Research Center, Jülich, Germany

⁴Physics Department, University of Wuppertal, Wuppertal, Germany

⁵Wind Energy Department, Technical University of Denmark, Roskilde, Denmark

⁶Hessian Agency for the Environment and Geology, Wiesbaden, Germany

⁷Institute for Climate and Atmospheric Science, School of Earth and Environment, University of Leeds, Leeds, UK

*now at: Institute for Advanced Sustainable Studies, Potsdam, Germany

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Received: 5 September 2013 – Accepted: 13 October 2013 – Published: 24 October 2013

Correspondence to: B. Bonn (boris.bonn@iass-potsdam.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

ACPD

13, 27501–27560, 2013

**Link between
radicals and particle
formation**

B. Bonn et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

It has been claimed for more than a century that atmospheric new particle formation is primarily influenced by the presence of sulphuric acid. However, the activation process of sulphuric acid related clusters into detectable particles is still an unresolved topic.

In this study we focus on the PARADE campaign measurements conducted during August/September 2011 at Mt. Kleiner Feldberg in central Germany. During this campaign a set of radicals, organic and inorganic compounds and oxidants and aerosol properties were measured or calculated. We compared a range of organic and inorganic nucleation theories, evaluating their ability to simulate measured particle formation rates at 3 nm in diameter (J_3) for a variety of different conditions. Nucleation mechanisms involving only sulphuric acid tentatively captured the observed noon-time daily maximum in J_3 , but displayed an increasing difference to J_3 measurements during the rest of the diurnal cycle. Including large organic radicals, i.e. organic peroxy radicals (RO_2) deriving from monoterpenes and their oxidation products in the nucleation mechanism improved the correlation between observed and simulated J_3 . This supports a recently proposed empirical relationship for new particle formation that has been used in global models. However, the best match between theory and measurements for the site of interest was found for an activation process based on large organic peroxy radicals and stabilized Criegee intermediates (sCI). This novel laboratory derived algorithm simulated the daily pattern and intensity of J_3 observed in the ambient data. In this algorithm organic derived radicals are involved in activation and growth and link the formation rate of smallest aerosol particles with OH during daytime and NO_3 during nighttime. Because of the RO_2 s lifetime is controlled by HO_2 and NO we conclude that peroxy radicals and NO seem to play an important role for ambient radical chemistry not only with respect to oxidation capacity but also for the activation process of new particle formation. This is supposed to have significant impact of atmospheric radical species on aerosol chemistry and should to be taken into account when studying the impact of new particles in climate feedback cycles.

Link between radicals and particle formation

B. Bonn et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

The formation of new particles by gaseous molecules has been observed in and above a variety of different forest ecosystems (Kulmala et al., 2004d). Newly formed particles are frequently able to grow to sizes at which they can be activated as cloud droplets (particle diameter $D_p > 60$ nm), so influencing the brightness and lifetime of clouds. It has been suggested that biogenic emissions from forests are involved in new particle formation and growth, meaning that forests could impact climate through aerosol-cloud interactions. Previous studies have suggested that the magnitude of this effect could be substantial, with Spracklen et al. (2008) estimating that monoterpene emissions from boreal forests cause between 1.8 and 6.4 W m⁻² of regional cooling through the aerosol indirect effect. Therefore a new climate feedback process linking forest emissions, aerosol and climate effects has been proposed (Kulmala et al., 2004d; Bonn et al., 2008, 2009; Carslaw et al., 2010; Paasonen et al., 2013). For both steps involved in new particle formation, i.e. nucleation and first steps of growth, knowledge of the chemical composition and the controlling parameters for the initial size range (particle diameter $D_p < 10$ nm) are essential but so far unidentified.

A number of studies focusing on detailed understanding and explanation of atmospheric nucleation processes have been conducted in order to elucidate the key parameters driving and controlling new particle formation (NPF) (Hyyönen et al., 2005; Kulmala et al., 2004a, b, c; Bonn and Moortgat, 2003; Bonn et al., 2008, 2009; Mauldin III. et al., 2013). However, the current impossibility to chemically analyze the tiny and chemically unstable aerosol particles at their smallest size (≈ 1.5 nm in diameter) complicates the identification of the nucleation initiating molecules or compounds (NIM) and the compounds needed for the initial activation and growth. This initial activation is thought to be controlled by the presence of air ions (Laakso et al., 2006) or by strong chemical interactions of participating molecules such as sulphuric acid. Some previous studies have focused on sulphuric acid related explanations only (Weber et al., 1999; Laakso et al., 2006; Riipinen et al., 2007; Paasonen et al., 2012; Mauldin III. et al.,

[Link between radicals and particle formation](#)

B. Bonn et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Link between
radicals and particle
formation**B. Bonn et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2013), whereas other theories rely on iodine oxides (Burkholder et al., 2004; O'Dowd and Hoffmann, 2005), organic substances (Bonn and Moortgat, 2003), or a mixed approach of organic and inorganic species (Kulmala et al., 2004a, b; Bonn et al., 2008). For example, Bonn and Moortgat (2003) have suggested that sesquiterpenoid products initiate nucleation.

To find indications of the nucleation mechanism, indirect approaches involving correlations between particle formation rate (typically either at 3 nm (J_3) or at 1 nm (J_1) in diameter) and ambient gas-phase concentrations have been applied. A number of studies have explored the relationship between particle formation rate and the concentration of gaseous sulphuric acid for the SMEAR II boreal forest site in Hyytiälä (Southern Finland) as well as for other locations (Kulmala et al., 2006; Sihto et al., 2006; Riipinen et al., 2007; Zhang et al., 2012). The correlations appear to be environment dependent, with relationships linking sulphuric acid concentration [H_2SO_4] and J_3 involving sulphuric acid to the power of between unity and two (Kulmala et al., 2006; Sihto et al., 2006; Riipinen et al., 2007; Nieminen et al., 2012) multiplied by empirically derived pre-factors (i.e. A or K) that vary spatially and temporally by orders of magnitude (Kulmala et al., 2006; Sihto et al., 2006; Riipinen et al., 2007). These empirically derived pre-factors are assumed to include the lumped kinetics of particle formation as well as further species contributing to the initial steps of particle growth (Bonn et al., 2008). The most recent empirical formulation for the boreal forest site in Hyytiälä (Finland) is provided by Nieminen et al. (2012). Nonetheless, the detailed meaning and understanding of the formation process remains an open question.

One challenge in this context is the correlation between a variety of different substances especially those linked through a similar method of production or destruction. For example, sulphuric acid is highly correlated with the atmospheric oxidant OH, which is directly linked to solar radiation intensity. OH controls the oxidation of most atmospheric compounds, therefore linking a host of organic compounds to the concentration of OH and sulphuric acid. This complicates correlations between different atmospheric species, with individual correlations being difficult to interpret without observations of

an entire set atmospheric species. Therefore a good correlation is neither a proof nor a disproof of a compounds particular role in the nucleation process.

Here we investigate a novel potential link between the new particle formation process and ambient radicals. These radicals are expected to increase the nucleus size through chemical reactions that enable the cluster to overcome the critical nucleation size. The primary candidates for these heterogeneous reactions (NIM or cluster + radical), are organic peroxy radicals from terpene oxidation (Bonn et al., 2007; Wolf et al., 2009, 2011). If this is the case, it is likely that the entire formation process is controlled by a number of different compounds, i.e. ambient inorganic and organic radicals such as hydroxyl radical (OH), hydroperoxy radicals (HO₂), nitrogen monoxide (NO), the nitrate radical (NO₃) and different organic peroxy radicals (RO₂). These are key species for representing different atmospheric chemistry situations, i.e. nitrogen oxides (NO_x) and VOC control, the impact of radiation and temperature and of biosphere atmosphere interactions.

2 Theoretical background

New aerosol particles are formed from gaseous conversion forming an amorphous new phase (Pankow, 1994). Therefore a non-volatile and supersaturated gas-mixture is usually considered (Friedlander, 2000; Jacobson, 2005; Seinfeld and Pandis, 2006). In this context we will distinguish between two different rates, i.e. the nucleation and the formation rate of new particles. As nucleation describes the novel production of a new phase with a distinct boundary between the phases the corresponding rate is named nucleation rate. On the contrary, ultrafine particles at larger sizes form by the split up of larger ones or by the growth of smaller ones to the apparent size. The increase in number in a certain size interval is called the formation rate and the exact value depends on the particular size of interest. As the exact starting size of a new particle relies on the compounds involved, we will focus on the formation rate at 3 nm in particle diameter J_3 . At this size a new phase (i.e. a particle) is definitely formed independent

Link between radicals and particle formation

B. Bonn et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of its chemical composition and is detectable with a commercial condensation particle counter.

The formation of clusters and particles from gaseous molecules in the atmosphere needs to overcome several nucleation hampering effects. Those include the surface production for a novel droplet or crystal (Kelvin effect) and a sufficient growth, before the clusters or smallest particles become captured by preexisting particles. This capturing process depends on the growth rate GR (diameter per time) and the condensation (in m^{-2} , CS') or coagulation sink (in m^{-2} , CoagS'). The larger GR (substantial mass supply), and therefore the shorter the time to survive before becoming detectable, and the smaller CS' or CoagS' , the higher the survival rate. These two sink terms are commonly provided in two different units, i.e. (a) in s^{-1} to represent the ambient lifetime of the clusters or particles before getting lost on preexisting surfaces (Pirjola and Kulmala, 1998; Dal Maso et al., 2005) or in (b) m^{-2} (here denoted with an $'$), i.e. related to the available aerosol surface area for attachment due to diffusional collisions (Kerminen and Kulmala, 2002).

Since homogeneous homomolecular nucleation by a single compound generally requires extremely high supersaturation for any of the relevant species to overcome the Kelvin effect, it is not relevant for stable compounds except radicals at atmospheric background conditions (Friedlander, 2000; McFiggans et al., 2004). Homogeneous nucleation by multiple compounds (heteromolecular) requires a substantial interaction between compounds such as present in the binary or ternary nucleation of sulphuric acid-water and of sulphuric acid-water-ammonia or amines (Friedlander, 2000; Seinfeld and Pandis, 2006; Kirkby et al., 2011; Paasonen et al., 2012). These intermolecular interactions such as acid-base stabilization (Kirkby et al., 2011; Ryding et al., 2012) lower the critical energy that is required to form the new particle phase. There are two further physico-chemical effects that may cause a drop in the nucleation barrier energy, i.e. (i) the solution (Raoult) effect like in the case of water-soluble species and (ii) the ionic (Thompson) effect (Friedlander, 2000; Jacobson, 2005; Seinfeld and Pandis, 2006).

**Link between
radicals and particle
formation**

B. Bonn et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Link between
radicals and particle
formation**B. Bonn et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

In the size range of interest the solvent mass is extremely small and even water vapor is not able to join the aerosol phase due to the Kelvin effect without considering chemical effects. One theory including this effect is the so-called nano-Koehler theory (Kulmala et al., 2004a), which assumes sulphuric acid and water containing clusters become activated by water-soluble organic compounds. But organic compounds of notable mass are primarily less water-soluble and the solubility effects commonly become predominant at sizes beyond 50 nm in diameter, i.e. sizes substantially greater than the nucleation size. Earlier investigations on the ionic effect within the vegetated boundary layer revealed a contribution of less than $0.1 \text{ cm}^{-3} \text{ s}^{-1}$ (Hirsikko et al., 2011) to the particle formation rate at 2 nm (J_2). This effect declines further still towards larger particle sizes because of a multitude of sink terms. Laakso et al. (2006) quantified its relative contribution to less than 10%. Despite these nontrivial theories, numerous discrepancies exist and so far none of these schemes is capable of adequately explaining the details of observed particle formation and growth rates in the boundary layer above vegetation. There is a need for modified or novel approaches and explanations for the growth of NIM to detectable sizes of 3 nm in particle diameter.

The remaining possibility is of reactive nature. Herein gas phase molecules interact with cluster surfaces and form new macromolecules (Zappoli et al., 1999; McFiggans et al., 2004; Kalberer et al., 2004; O'Dowd and Hoffmann, 2005). The process is not affected by the Kelvin effect and is controlled only by the reactivity and concentration of cluster and corresponding molecules. This reaction controlled activation (RCA) requires the presence of trace gases, possessing a high reactivity and notable molar mass, i.e. larger radicals, which basically initiate a catalytic reaction chain. In this RCA, radicals react with clusters maintaining the reactive site and contributing to particulate mass (Burkholder et al., 2004). The process is similar to the behavior of OIO that initiates a polymerization to form new clusters and particles. This is capable of explaining new particle formation observed at coastal sites (O'Dowd et al., 2001; McFiggans et al., 2004; O'Dowd and Hoffmann, 2005). But halogen radicals are not available in substantial amounts for nucleation at forest sites.

**Link between
radicals and particle
formation**B. Bonn et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

In forested areas large amounts of biogenic volatile organic compounds (BVOCs) are emitted (Guenther et al., 1995, 2006; Bourtsoukidis et al., 2012). These compounds have a variety of different reactivities, functionalities, molecular sizes (Kesselmeier and Staudt, 1999) and driving forces (Guenther et al., 1995; Niinemets et al., 2010; Bourtsoukidis et al., 2012). During daytime most of the BVOCs are oxidized by the hydroxyl radical (OH) to form an organic peroxy radical (RO₂) in the first stage. This radical proceeds reacting with one of the other available radicals such as HO₂, NO, RO₂ and NO₃ in the gas-phase to form stable intermediate products such as alcohols, nitrates, organic hydroperoxides, carbonyl compounds or organic acids. Those compounds contribute to secondary organic aerosol mass after partitioning to the preexisting organic aerosol (Pankow, 1994; Hoffmann et al., 1997; Barsanti and Pankow, 2004, 2005, 2006; Donahue et al., 2006, 2011). A further heterogeneous reaction was postulated by Bonn et al. (2007) and revealed under laboratory conditions by Wolf et al. (2009, 2011) for organic peroxy radicals consisting of more than five carbon atoms (RO₂(C > 5)). These large organic peroxy radicals are thought to react with aerosol constituents producing a variety of different macromolecules that have been seen in infrared spectra (Wolf et al., 2009). Organic peroxy radicals with less than six carbon atoms seem to split after formation of a very short-lived intermediate caused by the release of reaction energy, undergoing transfer to stable carbonyl compounds. Thus, they cannot contribute to particle mass and may seal the reactive sites preventing further reactions. Organic peroxy radicals with six C atoms or more (RO₂(C > 5)) tend to distribute the elevated energy over a sufficiently large molecule and form a tentatively stable macromolecule. The proposed growth process is independent of the cluster chemical nature, i.e. inorganic or organic, and is only dependent on its reactivity with respect to ambient radicals. This distinction between small and large organic compounds has important implications for BVOC emissions from vegetation: temperate forests with high emissions of isoprene will yield small (C ≤ 5) RO₂ in high amounts, preventing activation and thus particle formation (Kiendler-Scharr et al., 2009), while coniferous forests with dominant

(mono)terpene emissions act as proposed by Bonn et al. (2007) and Wolf et al. (2009, 2011).

Support for this process has been found recently in laboratory studies (Hummel, 2010; Ebach, 2010; Kestel, 2013). These studies used alkene-ozone reactions in the presence of nopinone (C₉H₁₄O), a ketone product of β-pinene oxidation. Neither ethene- nor propene- or cis-2-butene-ozone reactions are expected to create new particles at moderate concentrations in the range of less than 10¹² cm⁻³ (< 40 ppb_v). However, adding nopinone (2.5 × 10¹²–1.5 × 10¹³ cm⁻³, that is 100–1000 ppb_v, i.e. ca. two to three orders of magnitude below saturation level) they cause significant formation of new particles and the activation process can be explained solely by the formed OH radicals as a product of the ozonolysis and the organic peroxy radicals involved.

For the present study three different nucleation hypotheses are considered:

- (A) sulphuric acid only based on Nieminen et al. (2012),
- (B) interaction of organic compounds with sulphuric acid based on Bonn et al. (2008) and Metzger et al. (2010) and
- (C) organic only based on studies of Bonn et al. (2007); Wolf et al. (2009, 2011); Hummel (2010); Ebach (2010) and Kestel (2013).

(A) Sulphuric acid only

This hypothesis is solely focused on the presence of sulphuric acid, obtained from fits to measurement data at the Finnish boreal station Hyytiälä and is scaled to the acids concentration the following way:

$$J_{3,H_2SO_4} = 2 \times 10^{-11} \text{ cm}^{-3} \text{ s}^{-1} ([H_2SO_4] \text{ cm}^3)^{1.78} \quad (1)$$

The pre-factor 2 × 10⁻¹¹ cm⁻³ s⁻¹ includes all unknowns and the reactivity of interacting, partially unknown compounds.

(B) Interaction of organic compounds with sulphuric acid

The nucleation is described empirically as a product of monoterpenes, OH, condensation sink in the unit s^{-1} (CS, Eq. 3) and sulphuric acid concentration assigned with a scaling factor K (Kulmala et al., 2006; Sihto et al., 2006; Riipinen et al., 2007) and has been found to agree in a certain range for different ecosystems around the globe:

$$J_{3,\text{mix}} = K \cdot \frac{[\text{MT}][\text{OH}]}{\text{CS}} \cdot [\text{H}_2\text{SO}_4] \quad (2)$$

This formation approach (Eq. 2) uses the sulphuric acid concentration, an empirical fitting parameter K derived from ambient measurements and includes a monoterpene-OH-oxidation product that is assumed non-volatile and sticks to any aerosol surface at collision. For the present study a fit of PARADE data results in a K value of $(5 \pm 1) \times 10^{-25} \text{ cm}^6 \text{ s}^{-2}$ that is used for the intercomparison. For the formation approach (B) the lifetime of smallest clusters is assumed as $1/\text{CS}$, i.e. the ambient lifetime before collision with the next particle surface. CS in s^{-1} is used as formulated by Pirjola and Kulmala (1998):

$$\text{CS} = 2 \int_{D_{p0}}^{\infty} \beta n(D_p) \beta_M dD_p \approx 2\pi D_i \sum_i \frac{1 + \frac{2\lambda_{\text{air}}}{D_p}}{1 + \left(0.377 + \frac{3}{4\alpha}\right) \cdot \frac{2\lambda_{\text{air}}}{D_p} + \frac{3}{4\alpha} \left(\frac{2\lambda_{\text{air}}}{D_p}\right)^2} \cdot D_{pi} N_i \quad (3)$$

$$\text{CS}' = \frac{\text{CS}}{4\pi D_i} \quad (4)$$

These two equations include several parameters: the diffusion coefficient of the condensing molecules D_i commonly assumed to be sulphuric acid, the collision rate β of gas molecules and particles which changes for different sizes, the particle number density $n(D_p)$, the transition correction factor β_M for different particle sizes that includes the Knudsen number $Kn (= \frac{2\lambda_{\text{air}}}{D_p})$, the mean free path of air λ_{air} , the particle diameter

[Link between radicals and particle formation](#)

B. Bonn et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



D_p , the sticking coefficient α commonly assumed to be 1 and finally the particle number concentration N_i in a certain size section i . In order to calculate the entire loss of condensing molecules the loss must be integrated over the entire particle size distribution with the largest particles contributing most because of the difference in speed and because of the large surface area.

Analysing the particle formation rate description in Eq. (2), it may be concluded that the proposed interaction of large ($C > 5$) organic peroxy radicals is already included. Those are formed in the first step of monoterpene oxidation by OH. The reciprocal condensation sink displays a lifetime that is in the same range as the common RO_2 sinks such as NO or the hydroperoxy radical HO_2 . The K factor can be interpreted as well in two different ways, i.e. as described by Bonn et al. (2008) as a lumped constant from several reactions including ozone, terpenes and water vapour or as the reaction rate constant between organic peroxy radicals from monoterpenes and OH times the former reaction rate constant of the monoterpene-OH product and sulphuric acid.

(C) Organic only

This new particle formation theory is described by a three step process (see scheme provided in Fig. 1): (1) first the nucleation initiating molecule (NIM) with weak chemical bonds such as the secondary ozonide (SOZ) is formed (Bonn et al., 2002; Bonn and Moortgat, 2003). It provides insufficient supersaturation to allow homogeneous nucleation to occur. (2) The activation step occurs essentially as a catalytic chain reaction including the large organic peroxy radicals ($RO_2(C > 5)$) as well as stabilized Criegee intermediates (sCI, Criegee (1975)), during which the weak O-O bond is continuously split and reformed. This process is supposed to bridge the gap between the NIM size of around 1 nm and the step 3. (3) Onset of partitioning of organic compounds (Pankow, 1994). In terms of mass production, this step becomes efficient above about 5 nm and dominant above about 10 nm in particle diameter. This third step actually includes the partitioning and the condensation of low volatile compounds and heterogeneous reactions such as acid-catalyzed reactions (Barsanti and Pankow, 2004, 2005, 2006).

[Link between radicals and particle formation](#)

B. Bonn et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Link between
radicals and particle
formation**

B. Bonn et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The formation rate J_3 can be summarized by the Eqs. (5) and (6). It approximates the formation process by the limiting reaction between the SOZ and either a ($\text{RO}_2(\text{C} > 5)$) or a sCI. For this purpose SOZ, ($\text{RO}_2(\text{C} > 5)$) and sCI are gained by steady-state approaches (i.e. sources balance sinks), which is feasible for short-lived gases in the ambient with respect to the time resolution. However for the SOZ this may result in a moderate overestimation because of its larger lifetime of several minutes up to a quarter of an hour. The empirically shortened formulation results from a set of laboratory studies (Bonn et al., 2002; Bonn and Moortgat, 2003; Hummel, 2010; Ebach, 2010; Kestel, 2013) conducted earlier on the impact of a variety of trace gases such as NO, CO, water vapour and large ketones:

$$J_{3,\text{org}} = [\text{SOZ}] \cdot \min(2.5 \times 10^{-13} \text{ cm}^3 \text{ molec.}^{-1} \text{ s}^{-1} \cdot [\text{RO}_2](\text{terp.}), 2 \times 10^{-12} \text{ cm}^3 \text{ molec.}^{-1} \text{ s}^{-1} \cdot [\text{sCI}]) \quad (5)$$

$$\cdot \frac{2.5 \times 10^{-13} \text{ cm}^3 \text{ molec.}^{-1} \text{ s}^{-1} \cdot [\text{RO}_2](\text{terp.}) + 2 \times 10^{-12} \text{ cm}^3 \text{ molec.}^{-1} \text{ s}^{-1} \cdot [\text{sCI}]}{k_{\text{RO}_2}^{\text{HO}_2}[\text{HO}_2] + k_{\text{RO}_2}^{\text{NO}}[\text{NO}] + k_{\text{RO}_2}^{\text{NO}_3}[\text{NO}_3] + k_{\text{RO}_2}^{\text{RO}_2}[\text{RO}_2]}$$

$$\text{SOZ} = \frac{k_{\text{SQT}}^{\text{O}_3}[\text{SQT}][\text{O}_3] \cdot Y_{\text{stab}}}{80 \text{ s}^{-1} + k_{\text{sCI}}^{\text{H}_2\text{O}}[\text{H}_2\text{O}]} \cdot \frac{80 \text{ s}^{-1}}{k_{\text{SOZ}}^{\text{OH}}[\text{OH}] + \text{CS}} \quad (6)$$

[SOZ] abbreviates the secondary ozonide concentration formed by the intramolecular formation during sesquiterpene (SQT) ozone (O_3) reactions and approximated in Eq. (6). The kinetic speed of this intramolecular reaction has been quantified by Beck et al. (2011) for α -humulene and was assumed identical for β -caryophyllene, farnesene and the other sesquiterpenes present at Taunus Observatory (Bourtsoukidis et al., 2012). The reaction rate constant of the stabilized Criegee intermediates (sCI) with water, which is currently not accurately determined ($10^{-19} \text{ cm}^{-3} \text{ s}^{-1} < k_{\text{sCI}}^{\text{H}_2\text{O}} < 10^{-15} \text{ cm}^{-3} \text{ s}^{-1}$, Atkinson et al., 2006), is assumed as $2 \times 10^{-17} \text{ cm}^3 \text{ s}^{-1}$ (Großmann, 1999) and Y_{stab} is the stabilization ratio of the sesquiterpene mixture (here: as-

Link between radicals and particle formationB. Bonn et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

sumed as 0.8). $RO_2(\text{terp.})$ represents the sum of organic peroxy radicals formed from (mono)terpene oxidation by OH and NO_3 . It is important to note that while the sesquiterpenes are assumed to contribute primarily to the SOZ formation, monoterpenes and oxidation products are considered to control the activation process by sCI and $RO_2(\text{terp.})$. Although monoterpenes react much slower with ambient ozone, their sCIs do not form SOZ intramolecularly and therefore the concentration is essentially dependent on the primary sink reaction with ambient water vapour.

The mean reaction rate $k_{\text{SQT}}^{\text{O}_3}$ of the sesquiterpene mixture (Bourtsoukidis et al., 2012) with ambient ozone was calculated from the product of individual contributions and reaction rate constants taken as available from the literature. Missing reaction rate constants for individual SQTs have been approximated by structure activity relationships (Kourtchev, I., personal communication, 2012) based on the detected mixture determined by GC-MS analysis. The value of $1/\text{CS}$ again represents the approximated lifetime of SOZ molecules before being lost to an aerosol surface, which is controlled by preexisting aerosol surface area. The chemical destruction of SOZ is assumed to occur by OH ($k_{\text{SOZ}}^{\text{OH}} = 7.29 \times 10^{-11} \text{ cm}^3 \text{ s}^{-1}$) (Saunders et al., 2003; Jenkin et al., 2003), which leads to a minimum chemical lifetime of about 10^4 s assuming a hydroxyl radical concentration $[\text{OH}]$ of 10^6 cm^{-3} . If compared to the sink on aerosol surfaces CS the destruction by OH contributes negligibly to the gas-phase loss of SOZ. Finally, the minimum expression of two reaction rates of $RO_2(\text{C} > 5)$ and of sCI with the SOZ or the cluster, respectively represents the activation step (2) with the slower one controlling the entire process. These two reaction rate constants are taken from laboratory experiments of Hummel (2010) and simulations of Ebach (2010) representing the activation cycle summarized in a single step. While the reaction rate constant of RO_2 and NIM is basically the $RO_2 + RO_2$ reaction rate constant for larger peroxy radicals (Atkinson et al., 2006), the reaction rate constant of sCI and NIM is about 1.5×10^5 times the one of sCI with water. This was found to be rather consistent for a range of different conditions.

At least two of the nucleation hypotheses (B) and (C) are linked to organic peroxy radicals, while the situation remains unclear in case (A) because of the lumping of a multitude of processes in a single empirical fitting constant.

In order to test this hypothesized link between radicals and particle formation rates and to evaluate the potential importance of the individual pathways for atmospheric conditions at Mt. Kleiner Feldberg, we require the knowledge of the concentrations of individual terpene groups, OH, NO₃, ozone, water vapor, large organic peroxy radicals (RO₂(C > 5)), HO₂, NO, stabilized Criegee intermediates (sCI) and of the condensation sink CS of NIM at approximately 1.2–1.5 nm.

3 Measurements and methods

The “Particles And RADicals: Diel observations of the impact of urban and biogenic Emissions” (PARADE) campaign measurements were conducted at Taunus Observatory (50°13′ N, 8°26′ E, 825 m.a.s.l.). It is located at the hilltop of the second highest elevation of the Taunus ridge within a managed spruce forest of about 60 to 80 yr in age (Handisides et al., 2003). While the forest predominantly consists of spruce, smaller amounts of pines are present. The hill top was cleared of any forests about a century ago for meteorological measurements (circle of around 160 m in diameter) and kept free of coniferous trees since then. Several invasive smaller trees or bushes have grown there since then. A full set of trace gas, radical, particle and photolysis measurements were performed between the 14 August and the 9 September 2011.

3.1 Gaseous measurements

Gas-phase measurements were performed at two different locations (Fig. 2) due to the campaign focus and the measurement capacities:

Link between radicals and particle formation

B. Bonn et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(a) at the hill top including the following compounds were measured: Ozone, NO, NO₂, NO₃, N₂O₅, PAN, HONO, CO, HO₂, H₂O₂, SO₂, H₂SO₄, HCHO, glyoxal, acetone, methanol, toluene and monoterpenes.

(b) The second point was located close to the forest (find further details in Bourtsoukidis et al., 2012) measuring at a dynamical plant cuvette, which allowed detection of the ambient mixing ratio when a budget approach was applied. The compounds investigated were HCHO, acetone, benzene, isoprene, mono- and sesquiterpenes, major primary terpene oxidation products such as nopinone and pinonaldehyde and linalool. A highly time-resolved measurement of volatile organic compound (VOC) pattern was obtained continuously by Proton Transfer Reaction-Mass Spectrometry (HS-PTR-MS, Ionicon, Bourtsoukidis et al., 2013) in a plant cuvette at 4 m above soil level at a sunlit branch at the edge of the forest. This was essential to obtain sesquiterpene emissions and ambient mixing ratios above the detection limit. More details on the method can be found in Bourtsoukidis et al. (2012, 2013). Because of the insensitive PTR-MS approach for compounds with identical molar masses the detailed composition of mono- and sesquiterpenes were derived by (a) exemplary branch samples analyzed by gas chromatography-mass spectrometry (GC-MS) (FMI, Helsinki, FI) (Hakola et al., 2006) during the season and (b) by air samples analyzed by GC-MS (MPI Mainz). OH-reactivity was obtained by a novel method from the Max-Planck Institute in Mainz (Nölscher et al., 2012, 2013) for the same cuvette as used for the VOC measurements using the PTR-MS and at the hill top too. The individual instruments and methods as well as the corresponding groups responsible for data acquisition are listed in Table 1.

3.2 Particle measurements

Particles were measured by different sets of instruments. One set measured the particle size distribution by a nano-DMPS set-up focusing on the smallest particle size range between 3 and 20 nm in particle diameter at the hill top container. Particle formation rates J_3 at 3 nm were considered as the positive change in concentration of the particles between 3 and 5 nm per 10 min interval. For condensation sink calculations

**Link between
radicals and particle
formation**

B. Bonn et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a second set-up, a FMPS and an OPC were deployed in a Mobile Laboratory (MoLa) (Drewnick et al., 2012) about 20 m away from the cuvette measurement site in eastward direction (Fig. 2). With this set-up particle size distributions between 5.6 nm and 32 μm were detected, which is sufficient for aerosol sink estimates. The condensation sink itself was calculated according to Pirjola and Kulmala (1998); Kulmala et al. (2001) as given in Eq. (3). All particle measurements were conducted at a height of about 5 to 6 m above soil level similar to gas-phase and wind speed measurements. The detailed instrumentation and references to their methods can be found in Table 1.

3.3 Boundary layer mixing and dilution

As the hill top site is not homogeneous, with the surrounding vegetation diverse in height, mixture and density, a distinct horizontal variation of the individual parameters, especially VOCs and related radicals has to be expected. Therefore we will describe how forest edge concentration measurements have been modified to account for dilution during advection to the aerosol measurement location at the hill top center. This step is necessary in order to allow intercomparison with the aerosol measurements and the calculations based on the different hypotheses (A)–(C). Important to remember is the different location of the measurements due to optimum measurement conditions. VOC measurements were conducted at the spruce forest edge, while most of the other measurements were performed at the hill-top at a distance between 20–60 m away from the forest edge (Sect. 3.1). Therefore the derived VOC concentrations need a dilution correction as a function of wind speed and direction as well as of boundary layer characteristics to obtain the concentration at the aerosol measurement location.

This was done using the numerical atmospheric boundary layer model SCADIS applied to a very detailed 3-D structure of the present surface. The model accounts for topographic variations and considers the vegetation as a multi-layer medium. Implementing parameterizations for drag forces on leaves and stomatal conductance enables SCADIS to describe the exchange between the vegetative canopy and the atmosphere. The full description of the model, equations, and numerical details can be

The wind speed u_{meas} was obtained by HLOG at 4 m above the surface, half way between the forest edge and hill top and was interpolated for 10 min intervals.

3.4 Estimation of OH-concentration

One of the essential points for atmospheric particle formation is the quantification of atmospheric radicals such as OH. Therefore different approaches were used: as planned OH measurements failed because of a detector malfunction, the following calculation methods were applied: (1) the correlation of the photolysis rate of $\text{O}(^1\text{D})$ $\text{J}(\text{O}^1\text{D})$ with OH (Rohrer and Berresheim, 2006), (2) the budget approach of sources and sinks of OH as quantified by gaseous measurements and (3) the budget approach using quantified sources and OH reactivity measurements at the hill top.

With respect to approach (1) $\text{J}(\text{O}^1\text{D})$ was measured by spectroradiometer as described by Bohn et al. (2008). Data were available during the entire period for daytime. Rohrer and Berresheim (2006) have found a linear correlation with a mean slope of $2.4 \times 10^{11} \text{ s}$ for Hohenpeißenberg, Southern Germany. Available values for different comparable areas and pollution impacts were found between $1.4 \times$ and $3.9 \times 10^{11} \text{ s}$, this range is also to be expected at the Taunus Observatory. As the site of interest is influenced by anthropogenic and biogenic sources neither of the assumptions of Rohrer and Berresheim (2006) is valid and the provided equations were used in an extensive version (see supporting online information). With this, a correlation of $(1.8 \pm 0.1) \times 10^{11} \text{ s}$ was found (see supporting online information, Supplement Fig. S8). This was used for photolysis rates above 10^{-6} s^{-1} . Since this approach uses a limited amount of data and includes a notable uncertainty of individual parameters we assume a maximum uncertainty of the estimated OH concentration of a factor of 2. For the budget approaches (approaches 2 and 3) a chemical box model has been used assuming a steady-state condition between all source and sink terms for OH. All significant tropospheric production terms were considered, which include the photolysis of ozone in the presence of water vapor, hydrogen peroxide, HONO, HNO_3 and the reactions of HO_2 with NO, ozone, and the nitrate radical. HO_2 was estimated in a similar interactive

[Link between radicals and particle formation](#)

B. Bonn et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Link between
radicals and particle
formation**B. Bonn et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



way as OH through applying a chemical box model including the established tropo-
spheric background chemistry (see the Master Chemical Mechanism v3.2, Saunders
et al., 2003; Jenkin et al., 1997, 2003, 2012). Therefore initial OH and HO₂ concen-
trations were assumed as follows: 10⁴ cm⁻³ for OH and 10⁷ cm⁻³ for HO₂ and iterated
100 times until steady-state conditions were obtained. The OH destruction (sink) was
taken into account by two different approaches, i.e. (approach 2) by considering all
known sink terms of the radical (list of sinks, Supplement Table S1 of supporting online
information, section (B) and (approach 3) by taking into account the local OH reactivity
measurements (Nölscher et al., 2013).

Intercomparison of the three different approaches (Fig. 4) yielded similar results.
While approach (2) underestimated the OH concentration calculated by approach (1)
during daytime (median: -37 %), approach (3) overestimated it (median: +39 %, values
above 10⁷ cm⁻³ are excluded). This difference is caused by the different sink terms
only. However, the sink calculated is notably larger (+53 %) than measured by the OH
reactivity method of Nölscher et al. (2013) leading to higher OH estimations using
the reactivity method. Most likely horizontal changes in VOC mixing ratios and further
sinks do matter significantly as the OH reactivity changes notably from the forest edge
towards the cleared hill cap. However, further investigations are beyond the scope of
this study. More information can be found in the supporting online information (sections
B and C).

For formation rate calculations the measurements have been used where available
and the estimations using approach (1) ([OH] = f(J(O¹D))) at daytime and the budget
approach (2) at nighttime, where J(O¹D) is not available.

3.5 Particle formation rate at 3 nm and 5 nm

In order to test the effect of different sizes, particle formation rates have additionally
been calculated for 5 nm in diameter using the same nano-DMPS set-up. Both parti-
cle formation rates, i.e. at 3 and at 5 nm are related by the growth rate and the loss
rate. As this is a purely dynamical process Kerminen and Kulmala (2002) have derived

a simplified equation (Eq. 8):

$$J_5 = J_3 \cdot \exp\left(\gamma \frac{\text{CoagS}'(3 \text{ nm})}{\text{GR}} \cdot \left(\frac{1}{5 \text{ nm}} - \frac{1}{3 \text{ nm}}\right)\right) \text{CoagS}'(3 \text{ nm}) = \frac{\text{CoagS}(3 \text{ nm})}{4\pi D_i} \quad (8)$$

Here J_3 and J_5 represent the formation rates at 3 and 5 nm, γ is a lumped factor of $0.23 \text{ nm}^2 \text{ m}^2 \text{ h}^{-1}$ deriving from diffusive speeds, $\text{CoagS}'(3 \text{ nm})$ is the coagulation sink of 3 nm particles (unit s^{-1} , $\text{CoagS}(3 \text{ nm})$ divided by 4π and the diffusion constant D_i of the growing cluster). In the case of gases CoagS' is replaced by CS' and D_i is the diffusion constant of the condensing species, here to be assumed as sulphuric acid as is commonly assumed (Kulmala et al., 2001).

The CoagS' value in the unit of m^2 (Kulmala et al., 2001; Dal Maso et al., 2005) has been calculated from the FMPS and OPC derived particle size distribution. The GR at such small sizes is difficult to access directly especially above an inhomogeneous surface. For this a novel approach was applied. It is commonly assumed that new particle formation is linked to a rise in sulphuric acid concentration (Riipinen et al., 2007; Sihto et al., 2006) and thus in OH concentration or vice versa. The time difference in sulphuric acid or OH rise and the rise in particle number at a certain size can thus be interpreted as growing time. Thus the GR results from the size difference divided by the time shift. Here we cross-correlated the particle formation rate at 3 nm J_3 with the OH concentration (measurement, or estimation where not available) and looked for the time of maximum correlation within five hours of a particular day ($\text{GR}_{\text{min}} = 0.5 \text{ nm h}^{-1}$ assumed). This limitation is set because of a limit in data at larger times. The cross-correlation displays the time-shift between both quantities that displays the highest correlation. The application of the Kerminen–Kulmala (Kerminen and Kulmala, 2002) Eq. (8) is used in this context to check for the loss contribution and to check the sensitivity of individual hypothesis parameterizations.

[Link between radicals and particle formation](#)

B. Bonn et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



4 Results

4.1 Boundary layer transport dilution

As noted earlier, the surface information on vegetation, its height and leaf area density was used to initialize the backtrajectory boundary layer model. The outcome was treated as a function of wind direction and was subsequently scaled to the present wind speeds at 4 m above soil level. The resulting relative reduction for VOCs during the transport without any reactions is displayed in Fig. 5 for two different locations, i.e. the hill top, where formation rates and gases have been acquired as well as for the MoLa location, at which the entire particle size distribution has been measured. It is obvious that there were several periods of minor dilution (small transport times): at the start of measurements, at the day of the year (DOY) 233 and at the initial days of the second half (241–244). During these periods relative reduction was minor, while during the other times dilutions up to a tenth of the original values have occurred.

4.2 VOCs and tropospheric basic trace gases

This dilution certainly impacted on the VOC mixing ratios at the hill top. In Fig. 6 the volume mixing ratios of isoprene, mono- and sesquiterpenes are displayed at the forest edge close to the cuvette (top) and at the hill top in the center of the clearing, where most of the measurements were performed and the formation of new particles was investigated. The calculated dilution matches the three different VOC measurements of Bourtsoukidis et al. (2013) and by the MPI Mainz (not shown) performed at the hill top reasonably. A clear decline is apparent for sesquiterpenes since these compounds are additionally reduced by the reaction with ambient ozone. Nevertheless, notable amounts were still present at the center of the hill. The mean and extreme concentrations are listed in Table 3.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



results and thus the sink terms that depend on the water uptake of soluble VOCs and the hydronium ion (H_3O^+) concentration, used for subsequent calculations. Those days were the most humid days during the entire seasonal measurements in 2011 and those exceptional periods occurred during these particular days only. The nitrate radicals (NO_3) were measured using Cavity Ring-Down spectroscopy (Crowley et al., 2010, 2011), and the data are displayed in the lower graph of Fig. 8. These exhibit the opposite behaviour with maximum concentrations during the night. Mean nighttime NO_3 mixing ratios were (21 ± 16) ppt_v with maximum values during the night between DOY 227 and 228, 231 and 232 as well as between 243 and 244. The total maximum of 215 ppt_v was measured at DOY 244.1.

4.5 VOC based radicals

Ten minute-averaged concentrations of sesquiterpenes, ozone and water vapour have been used to approximate sCl concentrations and large RO_2 concentrations based on the mass balance between sources and sinks (bottom graph in Fig. 9) according to Eqs. (9)–(12):

$$[sCl] \approx \frac{k_{MT}^{O_3} \cdot [MT][O_3] \cdot Y_{stab,MT} + k_{LI}^{O_3} \cdot [Linalool][O_3] \cdot Y_{stab,LI}}{k_{sCl}^{H_2O} \cdot [H_2O]} \quad (9)$$

$$[RO_2(\text{terp}, \text{day})] = \frac{k_{SQT}^{O_3} \cdot [SQT][O_3] \cdot Y_{stab,SQT} + \frac{80 \text{ s}^{-1}}{k_{MT}^{OH} \cdot [MT] + k_{SQT}^{OH} \cdot [SQT] + \sum [TOX,d] \cdot k_{TOX,d}^{OH}} \cdot [OH]}{k_{RO_2}^{HO_2} \cdot [HO_2] + k_{RO_2}^{NO} \cdot [NO] + k_{RO_2}^{RO_2} \cdot [RO_2] + k_{RO_2}^{NO_3} \cdot [NO_3]} \quad (10)$$

$$\approx \frac{\left(k_{MT}^{OH} \cdot [MT] + k_{SQT}^{OH} \cdot [SQT] + \sum [TOX,d] \cdot k_{TOX,d}^{OH} \right) \cdot [OH]}{k_{RO_2}^{HO_2} \cdot [HO_2] + k_{RO_2}^{NO} \cdot [NO]}$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Link between
radicals and particle
formation**

B. Bonn et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$$[\text{RO}_2(\text{terp, night})] = \frac{(k_{\text{MT}}^{\text{NO}_3} \cdot [\text{MT}] + k_{\text{SQT}}^{\text{NO}_3} \cdot [\text{SQT}] + \sum[\text{TOX},n] \cdot k_{\text{TOX},n}^{\text{NO}_3}) \cdot [\text{NO}_3]}{k_{\text{RO}_2}^{\text{HO}_2} \cdot [\text{HO}_2] + k_{\text{RO}_2}^{\text{NO}} \cdot [\text{NO}] + k_{\text{RO}_2}^{\text{RO}_2} \cdot [\text{RO}_2] + k_{\text{RO}_2}^{\text{NO}_3} \cdot [\text{NO}_3]} \quad (11)$$

$$\approx \frac{(k_{\text{MT}}^{\text{NO}_3} \cdot [\text{MT}] + k_{\text{SQT}}^{\text{NO}_3} \cdot [\text{SQT}] + \sum[\text{TOX},n] \cdot k_{\text{TOX},n}^{\text{NO}_3}) \cdot [\text{NO}_3]}{k_{\text{RO}_2}^{\text{HO}_2} \cdot [\text{HO}_2] + k_{\text{RO}_2}^{\text{NO}} \cdot [\text{NO}] + k_{\text{RO}_2}^{\text{NO}_3} \cdot [\text{NO}_3]}$$

$$[\text{RO}_2(\text{terp})] = [\text{RO}_2(\text{terp, day})] + [\text{RO}_2(\text{terp, night})] \quad (12)$$

5 For both radical groups, i.e. sCI and $\text{RO}_2(\text{terp.})$ monoterpenes were the most important source class. Equation (9) displays the steady-state approach based on the ozonolysis reactions of monoterpenes, linalool and sesquiterpenes source and the predominant sink reaction with ambient water vapour for monoterpene and linalool derived sCI, while the intramolecular SOZ-formation of sesquiterpene derived sCI is the major sink of the

10 latter. k_X^Y constants represent reaction rate constants as provided by Bourtsoukidis et al. (2012); Shu and Atkinson (1994) and Atkinson et al. (2006) and $Y_{\text{stab},i}$ abbreviate the stabilization ratios of linalool or the sum of individual mono- and sesquiterpenes weighted by their contributions to the corresponding terpene class. The following values have been used: $Y_{\text{stab, MT}} = 0.4$, $Y_{\text{stab, LI}} = 0.4$ and $Y_{\text{stab, SQT}} = 0.8$.

15 The sum of individual terpene oxidation product concentrations is abbreviated by $([\text{TOX},i])$ with a “d” for daytime and an “n” for nighttime’ conditions. This sum times the corresponding reaction rate constants with OH ($k_{\text{TOX},i}^{\text{OH}}$) included the following measured VOCs: nopinone/sabinaketone (mass 138 g mole^{-1}), methyl chavicol (mass 148 g mole^{-1}) (Bouvier-Brown et al., 2009), linalool (mass 154 g mole^{-1}) and pinonaldehyde/limonaldehyde (mass 168 g mole^{-1}). Nopinone and sabinaketone were not

20 considered at nighttime because of their insignificant reaction rate constant. Since the molar mass was detected only and no structural information was available, the reaction rate constants of nopinone/sabinaketone as well as of pinonaldehyde/limonaldehyde

were taken as the sum of the individual reaction rate constants times the individual contribution of the parent monoterpene to emission (see supp. online information).

With respect to the organic peroxy radicals two aspects are worth mentioning:

- (1) Isoprene forms smaller peroxy radicals ($\text{RO}_2(\text{C} \leq 5)$) reducing the lifetime and thus the concentration of (very) large RO_2 and therefore counteracting the activation (Kiendler-Scharr et al., 2009). Thus, this RO_2 was considered only for the total sum of RO_2 but not for the $\text{RO}_2(\text{terp.})$ provided in Eqs. (10)–(12).
- (2) Investigating the individual contributions to $[\text{RO}_2(\text{C} > 5)]$ the contributions of sesquiterpenes and primary oxidation products of monoterpenes (ketones at 138 g mole^{-1} and aldehydes at 168 g mole^{-1}) are generally minor (10%). Thus the production of $[\text{RO}_2(\text{C} > 5)]$ is found to be primarily controlled by the monoterpene oxidation.

The calculated $[\text{sCl}]$ and $[\text{RO}_2(\text{C} > 5)]$ can be seen in Fig. 9.

4.6 Particle formation rate at 3

Next, Eqs. (1)–(3) were used to calculate the formation rate at 3 nm based on the theories (A), (B) and (C). The apparent J_3 was calculated as the change in interval number concentration between 3 and 5 nm and divided by 600 s due to the time resolution of 10 min of all measurements. Negative values as well as values below $10^{-3} \text{ cm}^{-3} \text{ s}^{-1}$ were excluded because of the focus on formation and because of the high uncertainty below the smallest value (count rate smaller than 1). A similar approach was done for obtaining J_5 that was derived from WCPC measurements (Table 1).

All derived values are displayed in Fig. 10 (top) and zoomed in the bottom plot for the third week with best quantitative measurements available. It is obvious that the total formation rate of particles is rather similar in structure for most of the days with about $0.1 \text{ cm}^{-3} \text{ s}^{-1}$ during midnight and a maximum of about $100 \text{ cm}^{-3} \text{ s}^{-1}$ during daytime. During the four weeks of campaign the first ten days were found to have the greatest

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Link between
radicals and particle
formation**B. Bonn et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

J_3 with smaller values between DOY 238 and 240 followed by a subsequent rise towards the end of the measurement period. None of the three different theories could be compared to the full dataset, since data gaps occurred particularly for sulphuric acid and VOCs. Therefore we concentrate on the periods of data availability for nearly all of the compounds, i.e. the start and the second half of the campaign. The peaks in J_3 at daily noon-time were calculated by all of the theories, since the entire process seems to be controlled by OH and the condensation sink. However, individual theories indicate a different daily variation. While the organic only hypothesis (C) displays a moderate daily variability due to the permanent presence of VOCs, the availability of ozone and the rather cloudy or rainy conditions, the hypotheses (B) and (A) show substantially greater variability between night- and day-time. The more the hypothesis is bound to sulphuric acid only the stronger the difference between night and day. This is shown in the bottom graph in Fig. 10. In matching the diurnal cycle, theory (C) performs best. In order to check if the exact corresponding size of the particle formation rate matters for the intercomparison, results of particle formation rates at 3 and 5 nm are both shown.

This different behaviour is obvious in the mean daily pattern by the measured and the calculated formation rates in Fig. 11. Both approaches (A) and (B) display a strong daily variation, while the organic based approach (C) does not. The reason for this difference is actually the different origins of large organic peroxy radicals at day- (OH) and nighttime (NO_3) and the different limitations in the minimum term (Sect. E in supporting online information). This minimum function in Eq. (5) results in a sCI limitation at nighttime with notable amounts of large RO_2 due to high levels of NO_3 , while at daytime the approach (C) is limited by the large monoterpene derived RO_2 .

A rather important aspect is the inhomogeneity of the surface resulting in a notable change in local concentrations and sinks, which are measurement point dependent. As different gases and the aerosol size distribution have been measured at different places (Fig. 2) because of logistical issues including availability of instrumentation and tubing losses in case of longer sampling lines, the exact merging of the entire dataset is affected by meteorological transport within the boundary layer. The approach of cal-

culating the transport dilution of terpenes and in this case especially the highly reactive sesquiterpenes is a reasonable approach but is limited by notable uncertainties in their loss intensity during the transport. We propose that it may serve as an order of magnitude estimate for cases of larger dilution as a function of the wind distribution and of longer transport times. Most of the deviations of the organic only hypothesis (C) actually occur exactly in situations of remarkable dilution and notable changes across the hill top. Therefore a spatially resolved 3-D simulation might be useful in future work. Other compounds such as sulphuric acid should be less dependent on local dilution, because of the production of sulphuric acid by sulfur dioxide that has been advected.

4.7 Loss between 3 and 5 nm

The next section will be used to investigate the size sensitivity of the formation rates that have been measured and calculated. As stated in Eq. (8) the loss between two sizes depends on CS' and GR. Both describe all sink and growth terms of distinct particles i.e. the loss onto present aerosol surface area (CS) and the uptake of gas molecules that is the acquisition of mass (GR).

CS' in the unit s^{-1} (CS) is shown in Fig. 14 (top) as well as GR in the bottom graph. A nearly uniform structure of CS between 10^{-2} and $10^{-3} s^{-1}$ is visible for Taunus Observatory. Please note the coagulation sink (CoagS) term for 3 nm particles is shown too. It is slightly smaller due to the reduced diffusivity of smallest particles compared to gas molecules. GR is more difficult to obtain as the growth rate at smallest sizes is hard to achieve. Here we use a novel approach and cross-correlate the OH-concentration that is related to the initial formation process and the formation rate at a certain size. For completeness the correlation with RO_2 and H_2SO_4 is shown as well (Fig. 12, top). The maximum correlation within four hours is assumed as the time taken to grow from nucleation size (ca. 1.2 nm) to the formation rate size, e.g. 3 nm, thus a minimum $GR(1.2-3\text{ nm})$ of 0.5 nm h^{-1} needs to be present for resolving it.

The cross-correlation analysis works as follows: two parameters are checked for similarity in structures i.e. pattern, maxima and minima locations. A perfect fit would result

**Link between
radicals and particle
formation**B. Bonn et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

in a value of unity, a perfect anticyclic behavior in -1 . Next, the correlation of parameters is calculated by shifting one parameter dataset in time (time shift displayed), while the second parameter dataset is kept constant. In this context the second parameter is the formation rate J_3 while the gases datasets are shifted in time. The larger the difference the earlier the same pattern must have appeared at the corresponding trace gas, thus the earlier the gas must have participated in the nucleation and formation process. Apparently the time shift is largest for OH. Thus, it is the most likely relevant species for the first step of formation. The shift is gently smaller for monoterpene derived RO_2 and nearly no time shift is visible for H_2SO_4 . This might suggest that sulphuric acid participates in the formation process at the latest stage. However since the process is non-linear and small amounts of sulphuric acid may be sufficient this is not a disproof for hypothesis (A), only an indication. In order to check how the OH estimation impacts on the results the bottom graph in Fig. 13 is added in order to display both structures, i.e. of OH calculation by using OH reactivity for all sink terms and by using the detailed chemistry calculations of all compounds considered. Only a minor change is obvious.

The derived GRs are shown in Fig. 12 (bottom). Applying those values in Eq. (6) the loss between 3 and 5 nm was calculated (Fig. 14). A loss between 1 and 16% was found. The measurement time is evidently split into two periods, the first part with a negligible loss and notable growth rates and the second part after DOY 240 with relative losses above 10%.

J_5 is shown as well. It can be seen that sometimes the organic only hypothesis formation rate slows down to the J_5 value. That might be caused by the notable variety of input parameters for calculation, i.e. the exact dilution especially at elevated wind speeds and the corresponding variability of the terpenes and thus the $RO_2(\text{terp})$ as mentioned above. However, at the mean daily behaviour, i.e. averaging all values at identical times throughout the campaign days, the organic hypothesis (C) matches best with a notable uncertainty range because of the multitude of parameters used for the calculations. Both other hypotheses (A) and (B) fit well at noon but display an increasing deviation towards the nighttime (Fig. S9, Supplement).

4.8 The link of particle formation rates to radicals

Finally the link between formation rates and different radicals, i.e. organic ($\text{RO}_2(\text{terp.})$ and sCl) and inorganic ones (OH), will be investigated and discussed. In order to allow intercomparison with other theories, sulphuric acid is displayed as well. An overview is presented in Fig. 15. It is apparent that all radicals and sulphuric acid display a linear relationship. both upper graphs of sulphuric acid and OH show a rather similar structure as the presence of sulphuric acid is highly connected with the presence of OH . The behaviour is seen for the stabilized Criegee intermediates primarily formed by monoterpenes and ozone and the large organic peroxy radicals primarily produced from monoterpene reactions with OH . It is apparent that there is more scatter in the lower graphs (sCl and $\text{RO}_2(\text{C} > 5)$) compared to the upper graphs of sulphuric acid and OH . This is caused by two aspects: (i) measurements of sulphuric acid and OH were available only for the last ten days of the campaign with improved meteorological conditions and (ii) sCl and $\text{RO}_2(\text{C} > 5)$ depend on a significant number of measurements and parameters such as VOC , dilution and oxidation speed of the potentially varying mixture in time. However H_2SO_4 , OH and $\text{RO}_2(\text{C} > 5)$ display a clear relationship (see table in supporting online information, Table S1). Correlations are also apparent for stabilised CI but the uncertainty of the derived relationship is significant. The general link to the particle formation rate at 3 nm is obvious as except sCl intermediates all depend on OH and thus provide a similar slope structure. To give a further indication for this Fig. 16 displays the relationship between $\text{RO}_2(\text{terp.})$ and sulphuric acid measurements (upper plot) as well as (b) the relationship of the reaction controlled activation term of Eq. (3) and the particle formation rates (lower plot). A link is visible.

5 Conclusions

In this study the potential link between ambient organic radical species and new particle formation rates was investigated. Observations of gas-phase and aerosol species

Link between radicals and particle formation

B. Bonn et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Link between
radicals and particle
formation**B. Bonn et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



were made during a four week intensive field campaign at the Taunus Observatory in Central Germany. Different particle formation hypotheses have been intercompared with observed particle formation rates at 3 nm particle size. While the simplified theories based only on concentrations of gas phase sulphuric acid were able to reproduce the maximum formation rates observed around local noon they were unable to reproduce the observed daily variability. Nucleation theories that involved the concentrations of organic radical species better matched the observed daily cycle in particle formation rate. Based on previous laboratory findings, the observations have been interpreted by the interaction of biogenic sesqui- and monoterpenes with ozone, OH and NO₃. It is the first time that the particle formation rate at or above a forest has been explained by a detailed nucleation scheme that is based on laboratory data and with a predominant role of ambient organic radicals. We have suggested that different biogenic alkenes may take different roles in the formation process: sesquiterpenes may form nucleation cores by reacting with ambient ozone, while monoterpenes may form activating radicals by reaction with OH and NO₃. This link between ambient radicals and new particle formation has not been reported so far and may explain the diversity of formation and growth rates observed at different locations and chemical conditions (NO_x, HO_x and RO₂). Our proposed nucleation scheme is also capable of explaining the isoprene suppression effect on particle formation (Kiendler-Scharr et al., 2009), since isoprene reacts with OH and NO₃ producing smaller organic peroxy radicals that do not activate but affect the larger radicals lifetime. If so this behavior would have consequences for the climate change feedback cycles of different ecosystems (Spracklen et al., 2008) and would provide an important key to understanding the atmospheric impacts of biome change under changing climatic conditions.

Supplementary material related to this article is available online at
**[http://www.atmos-chem-phys-discuss.net/13/27501/2013/
acpd-13-27501-2013-supplement.pdf](http://www.atmos-chem-phys-discuss.net/13/27501/2013/acpd-13-27501-2013-supplement.pdf)**

Acknowledgements. We would like to thank all participants and supporters of the PARADE field campaign 2011 at Taunus Observatory not included in the list of authors (B. Bohn, B. Bühner, J. Crowley, W. Haunold, A. Kürten, J. Curtius, K. Hens, H. Harder, A. Novelli, J. Fachinger, F. Drewnick, H. Fischer, G. Phillips, J. Thieser, A. Reiffs, A. Nölscher, J. Williams and R. Sitals) for sharing their data and comments and a wet but nice field campaign. Especially Birger Bohn, Hartwig Harder and Korbinian Hens are acknowledged for the intensive discussion about the OH derivation and reliability.

References

- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., Troe, J., and IUPAC Subcommittee: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II – gas phase reactions of organic species, *Atmos. Chem. Phys.*, 6, 3625–4055, doi:10.5194/acp-6-3625-2006, 2006. 27513, 27514, 27525
- Barsanti, K. C. and Pankow, J. F.: Thermodynamics of the formation of atmospheric organic particulate matter by accretion reactions – part 1: Aldehydes and ketones, *Atmos. Environ.*, 38, 4371–4382, 2004. 27509, 27512
- Barsanti, K. C. and Pankow, J. F.: Thermodynamics of the formation of atmospheric organic particulate matter by accretion reactions – 2. dialdehydes, methylglyoxal, and diketones, *Atmos. Environ.*, 39, 6597–6607, 2005. 27509, 27512
- Barsanti, K. C. and Pankow, J. F.: Thermodynamics of the formation of atmospheric organic particulate matter by accretion reactions – part 3: Carboxylic and dicarboxylic acids, *Atmos. Environ.*, 40, 6676–6686, 2006. 27509, 27512
- Beck, M., Winterhalter, R., Hermann, F., and Moortgat, G. K.: The gas-phase ozonolysis of α -humulene, *Phys. Chem. Chem. Phys.*, 13, 10970–11001, 2011. 27513
- Bohn, B., Corlett, G. K., Gillmann, M., Sanghavi, S., Stange, G., Tensing, E., Vrekoussis, M., Bloss, W. J., Clapp, L. J., Kortner, M., Dorn, H.-P., Monks, P. S., Platt, U., Plass-Dülmer, C., Mihalopoulos, N., Heard, D. E., Clemmshaw, K. C., Meixner, F. X., Prevot, A. S. H., and Schmitt, R.: Photolysis frequency measurement techniques: results of a comparison within the ACCENT project, *Atmos. Chem. Phys.*, 8, 5373–5391, doi:10.5194/acp-8-5373-2008, 2008. 27519

[Link between radicals and particle formation](#)

B. Bonn et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Link between
radicals and particle
formation**B. Bonn et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Bonn, B. and Moortgat, G. K.: Sesquiterpene ozonolysis: Origin of atmospheric new particle formation from biogenic hydrocarbons, *Geophys. Res. Lett.*, 30, 1585, doi:10.1029/2003GL017000, 2003. 27504, 27505, 27512, 27513
- Bonn, B., Korhonen, H., Petäjä, T., Boy, M., and Kulmala, M.: Understanding the formation of biogenic secondary organic aerosol from α -pinene in smog chamber studies: role of organic peroxy radicals, *Atmos. Chem. Phys. Discuss.*, 7, 3901–3939, doi:10.5194/acpd-7-3901-2007, 2007. 27506, 27509, 27510
- Bonn, B., Schuster, G., and Moortgat, G. K.: Influence of water vapor on the process of new particle formation during monoterpene ozonolysis, *J. Phys. Chem. A*, 106, 2869–2881, 2002. 27512, 27513
- Bonn, B., Kulmala, M., Riipinen, I., Sihto, S.-L., and Ruuskanen, T. M.: How biogenic terpenes govern the correlation between sulfuric acid concentrations and new particle formation, *J. Geophys. Res.*, 113, D12209, doi:10.1029/2007JD009327, 2008. 27504, 27505, 27510, 27512
- Bonn, B., Boy, M., Kulmala, M., Groth, A., Trawny, K., Borchert, S., and Jacobi, S.: A new parametrization for ambient particle formation over coniferous forests and its potential implications for the future, *Atmos. Chem. Phys.*, 9, 8079–8090, doi:10.5194/acp-9-8079-2009, 2009. 27504
- Bourtsoukidis, E., Bonn, B., Dittmann, A., Hakola, H., Hellén, H., and Jacobi, S.: Ozone stress as a driving force of sesquiterpene emissions: a suggested parameterisation, *Biogeosciences*, 9, 4337–4352, doi:10.5194/bg-9-4337-2012, 2012. 27509, 27513, 27514, 27516, 27525
- Bourtsoukidis, E., Williams, J., Kesselmeier, J., Jacobi, S., and Bonn, B.: From emissions to ambient mixing ratios: on-line seasonal field measurements over a Norway spruce dominated forest in central Germany, *Atmos. Chem. Phys. Discuss.*, submitted, 2013. 27516, 27518, 27522
- Bouvier-Brown, N. C., Holzinger, R., Palitzsch, K., and Goldstein, A. H.: Large emissions of sesquiterpenes and methyl chavicol quantified from branch enclosure measurements, *Atmos. Environ.*, 43, 389–401, 2009. 27525
- Burkholder, J. B., Curtius, J., Ravishankara, A. R., and Lovejoy, E. R.: Laboratory studies of the homogeneous nucleation of iodine oxides, *Atmos. Chem. Phys.*, 4, 19–34, doi:10.5194/acp-4-19-2004, 2004. 27505, 27508



**Link between
radicals and particle
formation**B. Bonn et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Carslaw, K. S., Boucher, O., Spracklen, D. V., Mann, G. W., Rae, J. G. L., Woodward, S., and Kulmala, M.: A review of natural aerosol interactions and feedbacks within the Earth system, *Atmos. Chem. Phys.*, 10, 1701–1737, doi:10.5194/acp-10-1701-2010, 2010. 27504
- Criegee, R.: Mechanismus der Ozonolyse, *Angew. Chem.*, 87, 765–771, 1975. 27512
- 5 Crowley, J. N., Schuster, G., Pouvesle, N., Parchatka, U., Fischer, H., Bonn, B., Bingemer, H., and Lelieveld, J.: Nocturnal nitrogen oxides at a rural mountain-site in south-western Germany, *Atmos. Chem. Phys.*, 10, 2795–2812, doi:10.5194/acp-10-2795-2010, 2010. 27524
- Crowley, J. N., Thieser, J., Tang, M. J., Schuster, G., Bozem, H., Beygi, Z. H., Fischer, H., Diesch, J.-M., Drewnick, F., Borrmann, S., Song, W., Yassaa, N., Williams, J., Pöhler, D.,
10 Platt, U., and Lelieveld, J.: Variable lifetimes and loss mechanisms for NO₃ and N₂O₅ during the DOMINO campaign: contrasts between marine, urban and continental air, *Atmos. Chem. Phys.*, 11, 10853–10870, doi:10.5194/acp-11-10853-2011, 2011. 27524
- Dal Maso, M., Kulmala, M., Riipinen, I., Wagner, R., Hussein, T., Aalto, P. P., and Lehtinen, K. E. J.: Formation and growth of fresh atmospheric aerosols: eight years of aerosol size distribution data from SMEAR II, Hyytiälä, Boreal Environ. Res., 5, 323–336, 2005. 27507, 27521
- 15 Donahue, N. M., Robinson, A. L., Stanier, C. O., and Pandis, S. N.: Coupled partitioning, dilution, and chemical aging of semivolatile organics, *Environ. Sci. Technol.*, 40, 2635–2643, 2006. 27509
- 20 Donahue, N. M., Trump, E. R., Pierce, J. R., and Riipinen, I.: Theoretical constraints on pure vapor pressure driven condensation of organics to ultrafine particles, *Geophys. Res. Lett.*, 38, L16801, doi:10.1029/2011GL048115, 2011. 27509
- Drewnick, F., Böttger, T., von der Weiden-Reinmüller, S.-L., Zorn, S. R., Klimach, T., Schneider, J., and Borrmann, S.: Design of a mobile aerosol research laboratory and data processing tools for effective stationary and mobile field measurements, *Atmos. Meas. Tech.*, 5, 1443–1457, doi:10.5194/amt-5-1443-2012, 2012. 27517
- 25 Ebach, F.: Simulation der Grössenverteilung von organischen Aerosolen während der Alken-Ozon-Reaktion unter dem Einfluss von NO, Master's thesis, J. W. Goethe-University, Frankfurt (Main), 2010. 27510, 27513, 27514
- 30 Friedlander, S. K.: *Smoke, Dust and Haze: Fundamentals of Aerosol Dynamics*, Oxford Univ. Press, New York, 2000. 27506, 27507

**Link between
radicals and particle
formation**

B. Bonn et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Großmann, D.: Die Gasphasenozonolyse von Alkenen in Gegenwart von Wasserdampf als Quelle für Wasserstoffperoxid und organische Peroxide in der Atmosphäre, PhD thesis, J. Gutenberg-University, Mainz, 1999. 27513

5 Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W. A., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.: A global model of natural volatile organic compound emissions, *J. Geophys. Res.*, 100, 8873–8892, 1995. 27509

10 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, doi:10.5194/acp-6-3181-2006, 2006. 27509

Hakola, H., Tarvainen, V., Bäck, J., Ranta, H., Bonn, B., Rinne, J., and Kulmala, M.: Seasonal variation of mono- and sesquiterpene emission rates of Scots pine, *Biogeosciences*, 6, 93–101, 2006,
15 <http://www.biogeosciences.net/6/93/2006/>. 27516

Handisides, G. M., Plass-Dülmer, C., Gilge, S., Bingemer, H., and Berresheim, H.: Hohenpeissenberg Photochemical Experiment (HOPE 2000): Measurements and photostationary state calculations of OH and peroxy radicals, *Atmos. Chem. Phys.*, 3, 1565–1588, doi:10.5194/acp-3-1565-2003, 2003. 27515

20 Hirsikko, A., Nieminen, T., Gagné, S., Lehtipalo, S. K., Manninen, H. E., Ehn, M., Hörrak, U., Kerminen, V.-M., Laakso, L., McMurry, P. H., Mirme, A., Mirme, S., Petäjä, T., Tammet, H., Vakkari, V., Vana, M., and Kulmala, M.: Atmospheric ions and nucleation: a review of observations, *Atmos. Chem. Phys.*, 11, 767–798, 2011,
<http://www.atmos-chem-phys.net/11/767/2011/>. 27508

25 Hoffmann, T., Odum, J., Bowman, F., Collins, D., Klockow, D., Flagan, R. C., and Seinfeld, J. H.: Formation of organic aerosols from the oxidation of biogenic hydrocarbons, *J. Atmos. Chem.*, 26, 189–222, 1997. 27509

Hummel, M.: Laborstudie zum Beitrag organischer Peroxyradikale (RO₂) bei der Partikelneubildung während der Ethen–Ozon–Reaktion, Master's thesis, J. W. Goethe-University, Frankfurt (Main), 2010. 27510, 27513, 27514

30 Hyvönen, S., Junninen, H., Laakso, L., Dal Maso, M., Grönholm, T., Bonn, B., Keronen, P., Aalto, P., Hiltunen, V., Pohja, T., Launiainen, S., Hari, P., Mannila, H., and Kulmala, M.:

**Link between
radicals and particle
formation**

B. Bonn et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A look at aerosol formation using data mining techniques, *Atmos. Chem. Phys.*, 5, 3345–3356, doi:10.5194/acp-5-3345-2005, 2005. 27504

Jacobson, M. Z.: *Fundamentals of Atmospheric Modeling*, 2nd edn., Cambridge Univ. Press, New York, 2005. 27506, 27507

5 Jenkin, M. E., Saunders, S. M., and Pilling, M. J.: The tropospheric degradation of volatile organic compounds: a protocol for mechanism development, *Atmos. Environ.*, 31, 81–104, 1997. 27520

Jenkin, M. E., Saunders, S. M., Wagner, V., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic volatile organic compounds, *Atmos. Chem. Phys.*, 3, 181–193, doi:10.5194/acp-3-181-2003, 2003. 27514, 27520

10 Jenkin, M. E., Wyche, K. P., Evans, C. J., Carr, T., Monks, P. S., Alfarra, M. R., Barley, M. H., McFiggans, G. B., Young, J. C., and Rickard, A. R.: Development and chamber evaluation of the MCM v3.2 degradation scheme for β -caryophyllene, *Atmos. Chem. Phys.*, 12, 5275–5308, doi:10.5194/acp-12-5275-2012, 2012. 27520

15 Kalberer, M., Paulsen, D., Sax, M., Steinbacher, M., Dommen, J., Prevot, A. S. H., Fisseha, R., Weingartner, E., Frankevich, V., Zenobi, R., and Baltensperger, U.: Identification of polymers as major components of atmospheric organic aerosols, *Science*, 303, 1659–1662, 2004. 27508

20 Kerminen, V. M., and Kulmala, M.: Analytical formulae connecting the “real” and the “apparent” nucleation rate and the nuclei number concentration for atmospheric nucleation events, *J. Aerosol Sci.*, 33, 609–622, 2002. 27507, 27520, 27521

Kesselmeier, J. and Staudt, M. A.: Biogenic volatile organic compounds (VOC): an overview on emission, physiology and ecology, *J. Atmos. Chem.*, 33, 23–88, 1999 27509

25 Kestel, P.: Nukleationsstudie zur Ozonolyse von cis-2-Buten, Master’s thesis, J. W. Goethe-University, Frankfurt (Main), 2013. 27510, 27513

Kiendler-Scharr, A., Wildt, J., Dal Maso, M., Hohaus, T., Kleist, E., Mentel, T. F., Tillmann, R., Uerlings, R., Schurr, U., and Wahner, A.: New particle formation in forests inhibited by isoprene emissions, *Nature*, 461, 381–384, 2009. 27509, 27526, 27531

30 Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagné, S., Ickes, L., Kürten, A., Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Tsagkogeorgas, G., Wimmer, D., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Downard, A., Ehn, M., Flagan, R. C., Haider, S., Hansel, A., Hauser, D., Jud, W.,

**Link between
radicals and particle
formation**

B. Bonn et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Junninen, H., Kreissl, F., Kvashin, A., Laaksonen, A., Lehtipalo, K., Lima, J., Lovejoy, E. R., Makhmutov, V., Mathot, S., Mikkilä, J., Minginette, P., Mogo, S., Nieminen, T., Onnela, A., Pereira, P., Petäjä, T., Schnitzhofer, R., Seinfeld, J. H., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P. E., Walther, H., Weingartner, E., Wex, H., Winkler, P. M., Carslaw, K. S., Worsnop, D. R., Baltensperger, U., and Kulmala, M.: Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation, *Nature*, 476, 429–433, 2011. 27507

Kulmala, M., Dal Maso, M., Mäkelä, J. M., Pirjola, L., Väkevä, M., Aalto, P., Miiikkulainen, P., Hämeri, K., and O'Dowd, C. D.: On the formation, growth and composition of nucleation mode particles, *Tellus B*, 53, 479–490, 2001. 27517, 27521

Kulmala, M., Kerminen, V.-M., Anttila, T., Laaksonen, A., and O'Dowd, C.: Organic aerosol formation via sulphate cluster activation, *J. Geophys. Res.*, 109, D04205, doi:10.1029/2003JD003961, 2004a. 27504, 27505, 27508

Kulmala, M., Laakso, L., Lehtinen, K. E. J., Riipinen, I., Dal Maso, M., Anttila, T., Kerminen, V.-M., Hörrak, U., Vana, M., and Tammet, H.: Initial steps of aerosol growth, *Atmos. Chem. Phys.*, 4, 2553–2560, doi:10.5194/acp-4-2553-2004, 2004b. 27504, 27505

Kulmala, M., Suni, T., Lehtinen, K. E. J., Dal Maso, M., Boy, M., Reissell, A., Rannik, Ü., Aalto, P., Keronen, P., Hakola, H., Bäck, J., Hoffmann, T., Vesala, T., and Hari, P.: A new feedback mechanism linking forests, aerosols, and climate, *Atmos. Chem. Phys.*, 4, 557–562, doi:10.5194/acp-4-557-2004, 2004c. 27504

Kulmala, M., Vehkamäki, H., Petäjä, T., Dal Maso, M., Lauri, A., Kerminen, V.-M., Birmili, W., and McMurry, P. H.: Formation and growth rates of ultrafine atmospheric particles: a review of observations, *J. Aerosol Sci.*, 35, 143–176, 2004d. 27504

Kulmala, M., Lehtinen, K. E. J., and Laaksonen, A.: Cluster activation theory as an explanation of the linear dependence between formation rate of 3nm particles and sulphuric acid concentration, *Atmos. Chem. Phys.*, 6, 787–793, doi:10.5194/acp-6-787-2006, 2006. 27505, 27511

Laakso, L., Anttila, T., Lehtinen, K. E. J., Aalto, P. P., Kulmala, M., Hörrak, U., Paatero, J., Hanke, M., and Arnold, F.: Kinetic nucleation and ions in boreal forest particle formation events, *Atmos. Chem. Phys.*, 4, 2353–2366, doi:10.5194/acp-4-2353-2004, 2004. 27504, 27508

**Link between
radicals and particle
formation**

B. Bonn et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Mauldin III., R. L., Berndt, T., Sipilä, M., Paasonen, P., Petäjä, T., Kim, S., Kurtén, T., Stratmann, F., Kerminen, V.-M., and Kulmala, M.: A new atmospherically relevant oxidant of sulphur dioxide, *Nature*, 488, 193–197, 2013. 27504

5 McFiggans, G., Coe, H., Burgess, R., Allan, J., Cubison, M., Alfarra, M. R., Saunders, R., Saiz-Lopez, A., Plane, J. M. C., Wevill, D., Carpenter, L., Rickard, A. R., and Monks, P. S.: Direct evidence for coastal iodine particles from *Laminaria* macroalgae – linkage to emissions of molecular iodine, *Atmos. Chem. Phys.*, 4, 701–713, doi:10.5194/acp-4-701-2004, 2004. 27507, 27508

10 Metzger, A., Verheggen, B., Dommen, J., Duplissy, J., Prevot, A. S. H., Weingartner, E., Riipinen, I., Kulmala, M., Spracklen, D. V., Carslaw, K. S., and Baltensperger, U.: Evidence for the role of organics in aerosol particle formation under atmospheric conditions, *P. Natl. Acad. Sci. USA*, 107, 6646–6651, 2010. 27510

15 Nieminen, T., Dal Maso, M., Petäjä, T., Aalto, P. P., Kerminen, V.-M., Hari, P., and Kulmala, M.: Trends on atmospheric new particle formation – 16 years of observations in boreal forest, *Geophysical Research Abstracts*, vol. 14, EGU2012-10036, EGU General Assembly, 2012. 27505, 27510

Niinemets, Ü., Arneth, A., Kuhn, U., Monson, R. K., Peñuelas, J., and Staudt, M.: The emission factor of volatile isoprenoids: stress, acclimation, and developmental responses, *Biogeosciences*, 7, 2203–2223, doi:10.5194/bg-7-2203-2010, 2010. 27509

20 Nölscher, A. C., Williams, J., Sinha, V., Custer, T., Song, W., Johnson, A. M., Axinte, R., Bozem, H., Fischer, H., Pouvesle, N., Phillips, G., Crowley, J. N., Rantala, P., Rinne, J., Kulmala, M., Gonzales, D., Valverde-Canossa, J., Vogel, A., Hoffmann, T., Ouwensloot, H. G., Vilà-Guerau de Arellano, J., and Lelieveld, J.: Summertime total OH reactivity measurements from boreal forest during HUMPPA-COPEC 2010, *Atmos. Chem. Phys.*, 12, 8257–8270, doi:10.5194/acp-12-8257-2012, 2012. 27516

25 Nölscher, A.C., Bourtsoukidis, E., Bonn, B., Kesselmeier, J., Lelieveld, J., and Williams, J.: Seasonal measurements of total OH reactivity emission rates from Norway spruce in 2011, *Biogeosciences*, 10, 4241–4257, doi:10.5194/bg-10-4241-2013, 2013. 27516, 27520

O'Dowd, C. D. and Hoffmann, T.: Coastal new particle formation: a review of the current state of the art, *Environ. Chem.*, 2, 245–255, 2005. 27505, 27508

30 O'Dowd, C. D., Jimenez, J. L., Bahreini, R., Flagan, R. C., Seinfeld, J. H., Hämeri, K., Pirjola, L., Kulmala, M., Jennings, S. G., and Hoffmann, T.: Marine aerosol formation from biogenic iodine emissions, *Nature*, 417, 632–636, 2001. 27508

[Link between radicals and particle formation](#)

B. Bonn et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Paasonen, P., Olenius, T., Kupiainen, O., Kurtén, T., Petäjä, T., Birmili, W., Hamed, A., Hu, M., Huey, L. G., Plass-Duelmer, C., Smith, J. N., Wiedensohler, A., Loukonen, V., McGrath, M. J., Ortega, I. K., Laaksonen, A., Vehkamäki, H., Kerminen, V.-M., and Kulmala, M.: On the formation of sulphuric acid – amine clusters in varying atmospheric conditions and its influence on atmospheric new particle formation, *Atmos. Chem. Phys.*, 12, 9113–9133, doi:10.5194/acp-12-9113-2012, 2012. 27504, 27507
- Paasonen, P., Asmi, A., Petäjä, T., Kajos, M. K., Äijälä, M., Junninen, H., Holst, T., Abbatt, J. P. D., Arneth, A., Birmili, W., Denier van der Gon, H., Hamed, A., Hoffer, A., Laakso, L., Laaksonen, A., Leaitch, W. R., Plass-Dülmer, C., Pryor, S. C., Räisänen, P., Swietlicki, E., Wiedensohler, A., Worsnop, D. R., Kerminen, V.-M., and Kulmala, M.: Warming-induced increase in aerosol number concentration likely to moderate climate change, *Nat. Geosci.*, 6, 438–442, 2013. 27504
- Pankow, J. F.: An absorption model of gas/particle partitioning of organic compounds in the atmosphere, *Atmos Environ.*, 28, 185–188, 1994. 27506, 27509, 27512
- Pirjola, L. and Kulmala, M.: Modelling the formation of $\text{H}_2\text{SO}_4\text{--H}_2\text{O}$ particles in rural, urban and marine conditions, *Atmos. Res.*, 46, 321–347, 1998. 27507, 27511, 27517
- Riipinen, I., Sihto, S.-L., Kulmala, M., Arnold, F., Dal Maso, M., Birmili, W., Saarnio, K., Teinilä, K., Kerminen, V.-M., Laaksonen, A., and Lehtinen, K. E. J.: Connections between atmospheric sulphuric acid and new particle formation during QUEST III–IV campaigns in Heidelberg and Hyytiälä, *Atmos. Chem. Phys.*, 7, 1899–1914, doi:10.5194/acp-7-1899-2007, 2007. 27504, 27505, 27511, 27521
- Rohrer, F. and Berresheim, H.: Strong correlation between levels of tropospheric hydroxyl radicals and solar ultraviolet radiation, *Nature*, 442, 184–187, 2006. 27519
- Ryding, M. J., Å. M. Jonsson, Zatulala, A. S., Andersson, P. U., and Uggerud, E.: Reactions of $\text{H}^+(\text{pyridine})_m(\text{H}_2\text{O})_n$ and $\text{H}^+(\text{NH}_3)_1(\text{pyridine})_m(\text{H}_2\text{O})_n$ with NH_3 : experiments and kinetic modelling, *Atmos. Chem. Phys.*, 12, 2809–2822, doi:10.5194/acp-12-2809-2012, 2012. 27507
- Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, *Atmos. Chem. Phys.*, 3, 161–180, doi:10.5194/acp-3-161-2003, 2003. 27514, 27520
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, 2nd edn., J. Wiley & Sons, 2006. 27506, 27507

**Link between
radicals and particle
formation**

B. Bonn et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Shu, Y. and Atkinson, R.: Rate constants for the gas-phase reactions of O₃ with a series of terpenes and OH radical formation from the O₃ reactions with sesquiterpenes at 296 ± 2 K, *Int. J. Chem. Kinet.*, 26, 1193–1205, 1994. 27525
- 5 Sihto, S.-L., Kulmala, M., Kerminen, V.-M., Dal Maso, M., Petäjä, T., Riipinen, I., Korhonen, H., Arnold, F., Janson, R., Boy, M., Laaksonen, A., and Lehtinen, K. E. J.: Atmospheric sulphuric acid and aerosol formation: implications from atmospheric measurements for nucleation and early growth mechanisms, *Atmos. Chem. Phys.*, 6, 4079–4091, doi:10.5194/acp-6-4079-2006, 2006. 27505, 27511, 27521
- 10 Sogachev, A., Menzhulin, G., Heimann, M., and Lloyd, J.: A simple three dimensional canopy-planetary boundary layer simulation model for scalar concentrations and fluxes, *Tellus B*, 54, 784–819, 2002. 27518
- Sogachev, A., Rannik, Ü., and Vesala, T.: Flux footprints over complex terrain covered by heterogeneous forest, *Agr. Forest Meteorol.*, 127, 143–158, 2004. 27518
- 15 Sogachev, A., Leclerc, M. Y., Zhang, G., Rannik, Ü., and Vesala, T.: CO₂ fluxes near a forest edge: a numerical study, *Ecol. Appl.*, 18, 1454–1469, 2008. 27518
- Sogachev, A., Kelly, M., and Leclerc, M. Y.: Consistent two-equation closure modeling for atmospheric research: buoyancy and vegetation implementations, *Bound.-Lay. Meteorol.*, 145, 307–327, 2012. 27518
- 20 Spracklen, D. V., Bonn, B., and Carslaw, K. S.: Boreal forests, aerosols and the impacts on clouds and climate, *Philos. T. R. Soc. A*, 366, 4613–4626, 2008. 27504, 27531
- Weber, R. J., McMurry, P. H., Mauldin III, R. L., Tanner, D., Eisele, F. L., Clarke, A. D., and Kapustin, V. N.: New particle formation in the remote troposphere: a comparison of observations at various sites, *Geophys. Res. Lett.*, 26, 307–310, 1999. 27504
- 25 Wolf, J. L., Suhm, M., and Zeuch, T.: Suppressed particle formation by kinetically controlled ozone removal: revealing the role of transient-species chemistry during alkene ozonolysis, *Angew. Chem.*, 48, 2231–2235, 2009. 27506, 27509, 27510
- Wolf, L., Richters, S., Pecher, J., and Zeuch, T.: Pressure dependent mechanistic branching in the formation pathways of secondary organic aerosol from cyclic-alkene gas-phase ozonolysis, *Phys. Chem. Chem. Phys.*, 13, 10952–10964, 2011. 27506, 27509, 27510
- 30 Zappoli, S., Andracchio, A., Fuzzi, S., Facchini, M. C., Gelencser, A., Kiss, G., Krivacsy, Z., Molnar, A., Meszaros, E., Hansson, H. C., Rosman, K., and Zebuhr, Y.: Inorganic, organic and macromolecular components of fine aerosol in different areas of Europe in relation to their water solubility, *Atmos. Environ.*, 33, 2733–2743, 1999. 27508

Zhang, R., Khalizov, A., Wang, L., Hu, M., and Xu, W.: Nucleation and growth of nanoparticles in the atmosphere, Chem. Rev., 112, 1957–2011, 2012. 27505

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

ACPD

13, 27501–27560, 2013

Link between radicals and particle formation

B. Bonn et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Measured parameter, instrumentation and groups responsible during the campaign.

Measured parameter	Instrumental techn. and instrument	Group responsible
O ₃	UV absorption, TE49	Fischer, MPI Mainz, D
NO, NO ₂	chemiluminescence,	Fischer, MPI Mainz, D
CO	RT-QCLS,	Fischer, MPI Mainz, D
H ₂ O ₂	Dual enzyme method, AL 2021	Fischer, MPI Mainz, D
HCHO	Hantzsch method, AL 4021	Fischer, MPI Mainz, D
NO ₂	GANDALF (LIF-NO ₂)	Harder, MPI Mainz, D
OH, HO ₂	IPI-FAGE-LIF	Harder, MPI Mainz, D
PAN, PAA	Iodide CIMS	Crowley, MPI Mainz, D
NO ₃ , N ₂ O ₅	CRD	Crowley, MPI Mainz, D
NO ₂ , alkyl nitrates	TD-CRDS	Crowley, MPI Mainz, D
Methanol, toluene	PTR-ToF-MS	Koppmann, BU Wuppertal, D
Isoprene, terpenes, div. VOCs	HS-PTR-MS, Ionicon, AUT	Bonn, GU Frankfurt/M., D
ind. terpene quantification	GC-MS	Hakola, FMI, Helsinki, D
OH reactivity	PTR-MS	Williams, MPI Mainz, D
H ₂ SO ₄	CIMS	Curtius, GU Frankfurt/M., D
J ₅	WCPC (TSI 3785)	Bingemer, GU Frankfurt/M., D
Aerosol size distribution	FMPS, APS	Drewnick, MPI Mainz, D
Particle size distribution (3–20 nm)	nano-SMPS	Kürten, GU Frankfurt/M., D
Photolysis rates	CCD spectrometer	Bohn, FZ Jülich, D
HONO	LOPAP	Li, FZ Jülich, D
T, rH, O ₃ , global radiation, NO _x	div.	Jacobi, HLUG, Wiesbaden, D

**Link between
radicals and particle
formation**

B. Bonn et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Link between radicals and particle formation

B. Bonn et al.

Table 2. Relative dilution at standard wind speed for four different wind directions between the hill top and the closed forest edge opposite to wind direction.

Wind direction	Relative reduction (nano-DMPS, gases)	Relative reduction (FMPS, OPS)
Southward (0°)	−(5 ± 1) %	−(0 ± 1) %
Westward (90°)	−(5 ± 1) %	−(25 ± 5) %
Northward (180°)	−(50 ± 5) %	−(65 ± 5) %
Eastward (270°)	−(50 ± 5) %	−(65 ± 5) %

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Link between radicals and particle formation

B. Bonn et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Measured mean, minimum and maximum isoprene, mono- and sesquiterpene concentrations at the forest edge and at the 2011.

VOC, forest edge	Mean conc. [molec. cm ⁻³]	Min. conc. [molec. cm ⁻³]	Max. conc. [molec. cm ⁻³]
Isoprene	$(4.5 \pm 2.5) \times 10^9$	0.8×10^9	2.8×10^{10}
Monoterpenes	$(6.5 \pm 3.6) \times 10^9$	1.7×10^9	2.6×10^{10}
Sesquiterpenes	$(1.1 \pm 0.6) \times 10^9$	2.3×10^8	3.1×10^9
VOC, hill top	Mean conc. [molec. cm ⁻³]	Min. conc. [molec. cm ⁻³]	Max. conc. [molec. cm ⁻³]
Isoprene	$(2.8 \pm 2.1) \times 10^9$	0.3×10^9	1.4×10^{10}
Monoterpenes	$(3.6 \pm 3.1) \times 10^9$	0.7×10^9	1.9×10^{10}
Sesquiterpenes	$(0.6 \pm 0.5) \times 10^9$	0.1×10^8	3.0×10^9

Link between radicals and particle formation

B. Bonn et al.




Fig. 1. Particle formation scheme of hypothesis (C) by organic compounds only. Three essential steps, of which the first two can both limit nucleation, are marked.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Link between radicals and particle formation

B. Bonn et al.



Fig. 2. Campaign site and location of individual measurements.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Link between radicals and particle formation

B. Bonn et al.




Fig. 3. Exemplary relative dilution simulation of a passive scalar at the height of 4 m for a southward wind direction (0°). The figures display: (top, left) the topography, (top right) the leaf area index distribution, (bottom left) concentration difference scaled to the concentration at the forest edge and (bottom right) relative trace gas flux reduction (Eq. (4), see Fig. 2). The measurement container for gases and some aerosol parameters at the hill top is located at (480 m, 430 m). Further plots can be found in supporting online information.

[Link between radicals and particle formation](#)

B. Bonn et al.




Fig. 4. Concentrations of ambient OH calculated through different approaches for a period during PARADE (DOY 240–247): (1) $J(O^1D)$ based formulation, (2) budget approach of measured individual sources and sinks as well as (3) budget approach of measured individual sources and OH reactivity. The extended version can be found in supporting online information.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

[Link between radicals and particle formation](#)

B. Bonn et al.




Fig. 5. Relative reduction of concentrations during transport from the forest edge to the hill top depending on the present wind direction and speed. This factor is used for scaling the measured VOC concentrations to obtain values after transport.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Link between
radicals and particle
formation**

B. Bonn et al.




Fig. 6. Measured volume mixing ratios at the forest edge (top) and calculated mixing ratios at the hill top (bottom) of isoprene, mono- and sesquiterpenes.

Link between radicals and particle formation

B. Bonn et al.




Fig. 7. Detailed behavior of detected ambient ozone and NO mixing ratios. Note, detection limit of NO was 21 ppt_v.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

[Link between radicals and particle formation](#)

B. Bonn et al.




Fig. 8. (top) Calculated OH concentrations at the hill top. (bottom) Detected NO_3 radical mixing ratios at the same location.




Fig. 9. (top) Estimated ambient stabilized Criegee Intermediates (sCI) and (bottom) $\text{RO}_2(\text{terp.})$ radicals.

Link between radicals and particle formation

B. Bonn et al.




Fig. 10. (top) Measured and derived particle formation rates at 3 nm in particle diameter and measured one at 5 nm. (bottom) Zoom to five days only in order to show the behavior at different times of the day.

[Link between radicals and particle formation](#)

B. Bonn et al.




Fig. 11. Mean daily pattern of measured and calculated particle formation rates at 3 nm diameter.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Link between
radicals and particle
formation

B. Bonn et al.




Fig. 12. (top) Calculated condensation (CS) and coagulation sinks (CoagS(3 nm)) and (bottom) mean growth rates (GR) of clusters or particles between 1.5 and 5 nm in diameter.

Link between radicals and particle formation

B. Bonn et al.




Fig. 13. (top) Cross-correlation of particle formation at 5 nm with RO₂(terp.), OH, NO₃ and H₂SO₄. (bottom) Cross-correlation between the particle formation rate at 5 nm and the estimated OH-concentration using two different approaches. The maximum is found at τ of 5.5 h (6 h for the budget approach), resulting in a GR of 3.5 nm/5.5 h = 0.6 nm/h.

Link between radicals and particle formation

B. Bonn et al.




Fig. 14. Calculated relative survival of particles during growth from 3 to 5 nm.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

[Link between radicals and particle formation](#)

B. Bonn et al.




Fig. 15. Scatter plots of sulphuric acid, OH, stabilised Criegee intermediates and large organic peroxy radicals by monoterpene oxidation ($RO_2(\text{terp})$) with the particle formation rate at 3 nm.




Fig. 16. Scatter plot of the activation term in Eq. (5) vs. the formation rate.