



## Physical Retracking of Jason-1 LRM data for ocean surface height/gravity field determination

Jain, Maulik; Andersen, Ole Baltazar; Stenseng, Lars; Dall, Jørgen

*Published in:*  
Geophysical Research Abstracts

*Publication date:*  
2013

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Jain, M., Andersen, O. B., Stenseng, L., & Dall, J. (2013). Physical Retracking of Jason-1 LRM data for ocean surface height/gravity field determination. *Geophysical Research Abstracts*, 15, Article EGU2013-8036.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



## **Physical Retracking of Jason-1 LRM data for ocean surface height/gravity field determination**

Maulik Jain, Ole Baltazar Andersen, Lars Stenseng, and Jørgen Dall

Technical University of Denmark, National Space Institute, DTU Space, Denmark (jain@space.dtu.dk)

Jason-1 Low Resolution Mode (LRM) waveforms can be processed to extract accurate heights of the ocean surface. These heights are adjusted taking into account various corrections available. Further the along surface slope/variation of these ocean heights can be used to make an estimation of the gravity field. An important part of this gravity field estimation is dependent on the way the LRM waveform is processed. Thus a physical model based on an error function is used, and the LRM waveforms are fit to this model. A processing system made up of 2 parameter and 3 parameter fitting models are used in order to extract the most reliable ocean surface heights. The quality of the processing system is judged by evaluating the standard deviation of the sea surface anomaly obtained after all corrections and the mean sea surface/geoid are removed. The lower the value of the standard deviation of the sea surface anomaly, the better the quality of processing is. Hence, different processing schemes are considered and evaluated in order to conclude towards the best retracking procedure which would eventually result in high accuracy gravity field estimations.