



Helheim 2006: Integrated Geophysical Observations of Glacier Flow

Nettles, M.; Ahlstrøm, A.; Elosegui, P.; Hamilton, G.; Kahn, S.; Langer, M.; Stearns, L.; Stenseng, Lars; Davis, J.; Rkström, G.

Total number of authors:
13

Publication date:
2006

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Nettles, M., Ahlstrøm, A., Elosegui, P., Hamilton, G., Kahn, S., Langer, M., Stearns, L., Stenseng, L., Davis, J., Rkström, G., Forsberg, R., Jørgensen, T., & Larsen, T. (2006). *Helheim 2006: Integrated Geophysical Observations of Glacier Flow*. Abstract from American Geophysical Union Fall Meeting 2006, San Francisco, CA, United States. <http://abstractsearch.agu.org/meetings/2006/FM/sections/S/sessions/S44A/abstracts/S44A-08.html>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

[AGU Abstract Browser](#) Beta

- [About](#)
- [Meetings](#)
- [Sections](#)
- [Index Terms](#)
- [Advanced Search](#)

Helheim 2006: Integrated Geophysical Observations of Glacier Flow

Details

Meeting [2006 Fall Meeting](#)

Section [Seismology](#)

Session [Bipolar Seismology II](#)

Identifier S44A-08

[Nettles, M*, Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY 10964 United States](#)

[Ahlstrøm, A, Geological Survey of Denmark and Greenland, \(GEUS\), Copenhagen, DK-1350 Denmark](#)

[Elósegui, P, Institute for Space Sciences, CSIC/IEEE, Barcelona, 08034 Spain](#)

[Hamilton, G, Climate Change Institute, University of Maine, Orono, ME 04469 United States](#)

[Khan, S, Danish National Space Center, \(DRC\), Copenhagen, DK-2100 Denmark](#)

[Langer, M, Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY 10964 United States](#)

[Stearns, L, Climate Change Institute, University of Maine, Orono, ME 04469 United States](#)

Authors

[Stenseng, L, Danish National Space Center, \(DRC\), Copenhagen, DK-2100 Denmark](#)

[Davis, J, Harvard-Smithsonian Center for Astrophysics, Space Geodesy Group, Cambridge, MA 02138 United States](#)

[Ekström, G, Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY 10964 United States](#)

[Forsberg, R, Danish National Space Center, \(DRC\), Copenhagen, DK-2100 Denmark](#)

[Jørgensen, T, Geological Survey of Denmark and Greenland, \(GEUS\), Copenhagen, DK-1350 Denmark](#)

[Larsen, T, Geological Survey of Denmark and Greenland, \(GEUS\), Copenhagen, DK-1350 Denmark](#)

[Glaciers \[0720\]](#)

Index [GEODESY AND GRAVITY \[1200\]](#)

Terms [Integrations of techniques \[1295\]](#)

[Abrupt/rapid climate change \[1605\]](#)
[Earthquake source observations \[7215\]](#)

Abstract

During the summer field season, 2006, we undertook a pilot geophysical experiment at Helheim Glacier, East Greenland, in which we deployed a network of GPS instruments on and around the glacier to measure the ice deformation field as a function of time. The experiment was motivated by the discovery of a new class of earthquakes occurring at glaciers in Alaska, Antarctica, and Greenland (Ekström, Nettles, and Abers, 2003). Teleseismic analysis indicates that these glacial earthquakes may result from the rapid sliding of the glacial ice over the glacier bed, and recent evidence (Ekström, Nettles, and Tsai, 2006) suggests a link to the hydrological cycle. However, little is understood about the mechanism by which the earthquakes occur. We installed sixteen GPS receivers on Helheim glacier, in a network spanning an upglacier distance of ~25~km from a point ~10~km behind the calving front. We also installed three GPS receivers at nearby rock sites to help define a stable reference frame. The stations were deployed in late June, 2006, and retrieved in late August, 2006. The GPS receivers recorded at a rate of at least 5~samples/sec. In addition, we operated several receivers for a few days each just behind the calving front during field visits in late June, late July, and late August, and we recorded the tidal stage using a pressure sensor near the end of Helheim Fjord for ~3~weeks during the experiment. Initial results show a variation in flow speed from about 25~m/day near the calving front to about 6~m/day at a location ~35~km behind the front. The horizontal flow speeds are tidally modulated, and an abrupt spatial change in vertical displacements due to the water tide gives the probable location of the glacier grounding line. We will present our geodetic results, and combine these results with seismological and glaciological observations to place constraints on the conditions under which glacial earthquakes are generated.

Cite as: Author(s) (2006), Title, *Eos Trans. AGU*, 87(52), Fall Meet. Suppl., Abstract S44A-08