

### Identifying added value in two high-resolution climate simulations over Scandinavia

Mayer, S.; Fox Maule, C.; Sobolowski, S.; Christensen, O. B.; Sørup, Hjalte Jomo Danielsen; Sunyer Pinya, Maria Antonia; Arnbjerg-Nielsen, Karsten; Barstad, I.

Publication date: 2014

Document Version Publisher's PDF, also known as Version of record

Link back to DTU Orbit

Citation (APA):

Mayer, S., Fox Maule, C., Sobolowski, S., Christensen, O. B., Sørup, H. J. D., Sunyer Pinya, M. A., Arnbjerg-Nielsen, K., & Barstad, I. (2014). *Identifying added value in two high-resolution climate simulations over Scandinavia*. Poster session presented at 3rd International Lund Regional-Scale Climate Modelling Workshop, Lund, Sweden.

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.





# Identifying added value in two high-resolution climate simulations over Scandinavia

### S. Mayer<sup>1,2</sup>, C. Fox Maule<sup>3</sup>, S. Sobolowski<sup>1,2</sup>, O.B. Christensen<sup>3</sup>, H.J.D. Sørup<sup>3,4</sup>, M.A. Sunyer<sup>4</sup>, K. Arnbjerg-Nielsen<sup>4</sup>, I. Barstad<sup>5</sup>

1) Uni Climate, Uni Research AS, Bergen, Norway, 2) Bjerknes Centre for Climate Research, Bergen, Norway, 3) Danish Climate Centre, Danish Meteorological Institute, Copenhagen, Denmark, 4) DTU Environment, Technical University of Denmark, Lyngby, Denmark, 5) Uni Computing, Uni Research AS, Bergen, Norway

- High resolution data are needed in order to assess potential impacts of extreme events on infrastructure in the mid latitudes.
- Dynamical downscaling offers one way to obtain this information.
- Prior to implementation in any impacts assessment scheme, model output must be validated and determined



fit-for-purpose.

 Results from two perfect-boundary experiments downscaling ERA-interim with HIRHAM5 and WRF3.3.1 on an 8 km grid over Scandinavia are shown.



Figure 1: Biases of seasonal temperature in K (simulation minus E-OBS data) in the WRF simulation for a) winter and b) summer season; c) and d) for the HIRHAM5 simulation; e) and f) for ERA interim. Model data is re-projected to the E-OBS 0.22 rotated grid. Yellow dots indicate grid points where the model biases lie outside an interval of 2. standard error which is given within the E-OBS data set.

Figure 2: Relative precipitation bias in %, in the WRF simulation for a) winter and b) summer season; c) and d) for the HIRHAM5 simulation; e) and f) for ERA interim. Yellow dots indicate grid points where the model biases lie outside an interval of 2 standard error which is given within the E-OBS data set. The absolute error ranges between ±3 mm/day.

Figure 3: Biases of wind speed in m/s (simulation minus Qscat) in a) and b) WRF, c) and d) HIRHAM, e) and f) ERA interim. The winter season is shown in the left column and the summer season in the right column. Gray areas indicate no data.

Table 1. Seasonal biases of wet-day mean precipitation in % were calculated for selected locations. A wet day is defined as a day when the precipitation amount exceeds 1 mm.

		DJF	MAM	JJA	SON
	ERA	-23.2	-28.8	-36.3	-33.6
Bergen	WRF	+10.5	-0.6	-16.6	-1.4
	HIRHAM	+14.1	+0.2	-28.9	-6.2
	ERA	-15.0	-18.6	-17.0	-26.5
Oslo	WRF	+18.8	+7.8	+7.3	+9.7
	HIRHAM	+66.6	+24.4	+1.2	+14.8
	ERA	-13.0	-26.6	-32.0	-25.5
Copenhagen	WRF	+1.3	-7.2	-11.6 -12.6 -1.7 +5.3	
	HIRHAM	+7.6	-6.1		

Table 2. Estimated e-folding distances in km of extreme precipitation for the duration of 3 hours and 24 hours. As in Gregersen et al. (2013) the estimates are derived from the fitted exponential models shown in Figure 4.

3	24 hr	Figure 4: Spatio-temporal correlation structure of observed (SVK), ERA interim (70 km) and
SVK	13	downscaled (8 km) mean intensities of extreme precipitation for 3 hour (a) and 24 hour (b) duration. To highlight the tendencies an exponential function is used for fitting by using least
WRF 2	42	square method (Gregersen et al., 2013).
HIRHAM 3	32	
ERA 1	128	

## 2.) Local-scale precipitation





- **Results** The models exhibit systematic cold and wet biases on seasonal time scales (-1 K and +50-100%, respectively). However, frequency based skill scores for daily precipitation and temperature are high (not shown), indicating that the distributions of these variables are generally well captured. Wind speeds over the North and Norwegian Seas were simulated more realistically in the models than ERA-interim reanalysis (Figure 3). Most important for impacts assessments, however, is that the high-resolution models are better capable of capturing the spatio-temporal behavior of short-duration extreme precipitation (Figure 4). In this respect both models outperform the reanalysis over Denmark, where recent pluvial floods led to costly damages to infrastructure.
- Gregersen, I., Sørup, H., Madsen, H., Rosbjerg, D., Mikkelsen, P., and Arnbjerg-Nielsen, K. (2013). Assessing future climatic changes of rainfall extremes at small spatio-temporal scales. Clim. *Change*, **118**(4), 783-797.

Contact: S. Mayer, Uni Research AS, Uni Climate, Allégaten 70, 5007 Bergen, Norway. Email: stephanie.mayer@uni.no