



Agriculture, Forestry and Other Land Use (AFOLU)

Smith, Pete ; Bustamante, Mercedes ; Ahammad, Helal ; Clark, Harry ; Dong, Hongmin ; Elsiddig, Elnour A. ; Haberl, Helmut ; Harper, Richard ; House, Joanna ; Jafari, Mostafa

Total number of authors:
19

Published in:
Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Smith, P., Bustamante, M., Ahammad, H., Clark, H., Dong, H., Elsiddig, E. A., Haberl, H., Harper, R., House, J., Jafari, M., Masera, O., Mbow, C., Ravindranath, N. H., Rice, C. W., Abad, C. R., Romanovskaya, A., Sperling, F., Tubiello, F. N., & Bolwig, S. (2014). Agriculture, Forestry and Other Land Use (AFOLU). In *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 811-922). Cambridge University Press.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Agriculture, Forestry and Other Land Use (AFOLU)

Coordinating Lead Authors:

Pete Smith (UK), Mercedes Bustamante (Brazil)

Lead Authors:

Helal Ahammad (Australia), Harry Clark (New Zealand), Hongmin Dong (China), Elnour A. Elsiddig (Sudan), Helmut Haberl (Austria), Richard Harper (Australia), Joanna House (UK), Mostafa Jafari (Iran), Omar Masera (Mexico), Cheikh Mbow (Senegal), Nijavalli H. Ravindranath (India), Charles W. Rice (USA), Carmenza Robledo Abad (Switzerland/Colombia), Anna Romanovskaya (Russian Federation), Frank Sperling (Germany/Tunisia), Francesco N. Tubiello (FAO/USA/Italy)

Contributing Authors:

Göran Berndes (Sweden), Simon Bolwig (Denmark), Hannes Böttcher (Austria/Germany), Ryan Bright (USA/Norway), Francesco Cherubini (Italy/Norway), Helena Chum (Brazil/USA), Esteve Corbera (Spain), Felix Creutzig (Germany), Mark Delucchi (USA), Andre Faaij (Netherlands), Joe Fargione (USA), Gesine Hänsel (Germany), Garvin Heath (USA), Mario Herrero (Kenya), Richard Houghton (USA), Heather Jacobs (FAO/USA), Atul K. Jain (USA), Etsushi Kato (Japan), Oswaldo Lucon (Brazil), Daniel Pauly (France/Canada), Richard Plevin (USA), Alexander Popp (Germany), John R. Porter (Denmark/UK), Benjamin Poulter (USA), Steven Rose (USA), Alexandre de Siqueira Pinto (Brazil), Saran Sohi (UK), Benjamin Stocker (USA), Anders Strømman (Norway), Sangwon Suh (Republic of Korea/USA), Jelle van Minnen (Netherlands)

Review Editors:

Thelma Krug (Brazil), Gert-Jan Nabuurs (Netherlands)

Chapter Science Assistant:

Marina Molodovskaya (Canada/Uzbekistan)

This chapter should be cited as:

Smith P., M. Bustamante, H. Ahammad, H. Clark, H. Dong, E.A. Elsiddig, H. Haberl, R. Harper, J. House, M. Jafari, O. Masera, C. Mbow, N.H. Ravindranath, C.W. Rice, C. Robledo Abad, A. Romanovskaya, F. Sperling, and F. Tubiello, 2014: Agriculture, Forestry and Other Land Use (AFOLU). In: *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Edenhofer, O., R. Pichs-Madruga, Y. Sokona, E. Farahani, S. Kadner, K. Seyboth, A. Adler, I. Baum, S. Brunner, P. Eickemeier, B. Kriemann, J. Savolainen, S. Schlömer, C. von Stechow, T. Zwickel and J.C. Minx (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Contents

Executive Summary	816
11.1 Introduction	818
11.2 New developments in emission trends and drivers	819
11.2.1 Supply and consumption trends in agriculture and forestry	822
11.2.2 Trends of GHG emissions from agriculture	822
11.2.3 Trends of GHG fluxes from forestry and other land use	825
11.3 Mitigation technology options and practices, and behavioural aspects	829
11.3.1 Supply-side mitigation options	829
11.3.2 Mitigation effectiveness (non-permanence: saturation, human and natural impacts, displacement)	832
11.4 Infrastructure and systemic perspectives	836
11.4.1 Land: a complex, integrated system	836
11.4.2 Mitigation in AFOLU—feedbacks with land-use competition	837
11.4.3 Demand-side options for reducing GHG emissions from AFOLU	838
11.4.4 Feedbacks of changes in land demand	841
11.4.5 Sustainable development and behavioural aspects	842
11.5 Climate change feedback and interaction with adaptation (includes vulnerability)	843
11.5.1 Feedbacks between AFOLU and climate change	845
11.5.2 Implications of climate change on terrestrial carbon pools and mitigation potential of forests	845
11.5.3 Implications of climate change on peatlands, grasslands, and croplands	845
11.5.4 Potential adaptation options to minimize the impact of climate change on carbon stocks in forests and agricultural soils	846
11.5.5 Mitigation and adaptation synergies and tradeoffs	846

11.6	Costs and potentials	847
11.6.1	Approaches to estimating economic mitigation potentials	848
11.6.2	Global estimates of costs and potentials in the AFOLU sector	848
11.6.3	Regional disaggregation of global costs and potentials in the AFOLU sector	849
11.7	Co-benefits, risks, and spillovers	852
11.7.1	Socio-economic effects	853
11.7.2	Environmental effects	855
11.7.3	Public perception	857
11.7.4	Spillovers	858
11.8	Barriers and opportunities	858
11.8.1	Socio-economic barriers and opportunities	858
11.8.2	Institutional barriers and opportunities	858
11.8.3	Ecological barriers and opportunities	859
11.8.4	Technological barriers and opportunities	859
11.9	Sectoral implications of transformation pathways and sustainable development	859
11.9.1	Characterization of transformation pathways	860
11.9.2	Implications of transformation pathways for the AFOLU sector	862
11.9.3	Implications of transformation pathways for sustainable development	862
11.10	Sectoral policies	862
11.10.1	Economic incentives	864
11.10.2	Regulatory and control approaches	864
11.10.3	Information schemes	868
11.10.4	Voluntary actions and agreements	868
11.11	Gaps in knowledge and data	868

11.12	Frequently Asked Questions	869
11.13	Appendix Bioenergy: Climate effects, mitigation options, potential and sustainability implications	870
11.13.1	Introduction	870
11.13.2	Technical bioenergy potential	870
11.13.3	Bioenergy conversion: technologies and management practices	873
11.13.4	GHG emission estimates of bioenergy production systems	877
11.13.5	Aggregate future potential deployment in integrated models	882
11.13.6	Bioenergy and sustainable development	883
11.13.7	Tradeoffs and synergies with land, water, food, and biodiversity	883
	References	887

Executive Summary

11

Agriculture, Forestry, and Other Land Use (AFOLU) is unique among the sectors considered in this volume, since the mitigation potential is derived from both an enhancement of removals of greenhouse gases (GHG), as well as reduction of emissions through management of land and livestock (*robust evidence; high agreement*). The land provides food that feeds the Earth's human population of ca. 7 billion, fibre for a variety of purposes, livelihoods for billions of people worldwide, and is a critical resource for sustainable development in many regions. Agriculture is frequently central to the livelihoods of many social groups, especially in developing countries where it often accounts for a significant share of production. In addition to food and fibre, the land provides a multitude of ecosystem services; climate change mitigation is just one of many that are vital to human well-being (*robust evidence; high agreement*). Mitigation options in the AFOLU sector, therefore, need to be assessed, as far as possible, for their potential impact on all other services provided by land. [Section 11.1]

The AFOLU sector is responsible for just under a quarter (~10–12 GtCO₂eq/yr) of anthropogenic GHG emissions mainly from deforestation and agricultural emissions from livestock, soil and nutrient management (*robust evidence; high agreement*) [11.2]. Anthropogenic forest degradation and biomass burning (forest fires and agricultural burning) also represent relevant contributions. Annual GHG emissions from agricultural production in 2000–2010 were estimated at 5.0–5.8 GtCO₂eq/yr while annual GHG flux from land use and land-use change activities accounted for approximately 4.3–5.5 GtCO₂eq/yr. Leveraging the mitigation potential in the sector is extremely important in meeting emission reduction targets (*robust evidence; high agreement*) [11.9]. Since publication of the IPCC Fourth Assessment Report (AR4), emissions from the AFOLU sector have remained similar but the share of anthropogenic emissions has decreased to 24% (in 2010), largely due to increases in emissions in the energy sector (*robust evidence, high agreement*). In spite of a large range across global Forestry and Other Land Use (FOLU) flux estimates, most approaches indicate a decline in FOLU carbon dioxide (CO₂) emissions over the most recent years, largely due to decreasing deforestation rates and increased afforestation (*limited evidence, medium agreement*). As in AR4, most projections suggest declining annual net CO₂ emissions in the long run. In part, this is driven by technological change, as well as projected declining rates of agriculture area expansion, which, in turn, is related to the expected slowing in population growth. However, unlike AR4, none of the more recent scenarios projects growth in the near-term [11.9].

Opportunities for mitigation include supply-side and demand-side options. On the supply side, emissions from land-use change (LUC), land management and livestock management can be reduced, terrestrial carbon stocks can be increased by sequestration in soils and biomass, and emissions from energy production can be saved through

the substitution of fossil fuels by biomass (*robust evidence; high agreement*) [11.3]. On the demand side, GHG emissions could be mitigated by reducing losses and wastes of food, changes in diet and changes in wood consumption (*robust evidence; high agreement*) [11.4] though quantitative estimates of the potential are few and highly uncertain. Increasing production without a commensurate increase in emissions also reduces emission intensity, i.e., the GHG emissions per unit of product that could be delivered through sustainable intensification; another mechanism for mitigation explored in more detail here than in AR4. Supply-side options depend on the efficacy of land and livestock management (*medium evidence; high agreement*) [11.6]. Considering demand-side options, changes in human diet can have a significant impact on GHG emissions from the food production lifecycle (*medium evidence; medium agreement*) [11.4]. There are considerably different challenges involved in delivering demand-side and supply-side options, which also have very different synergies and tradeoffs.

The nature of the sector means that there are potentially many barriers to implementation of available mitigation options, including accessibility to AFOLU financing, poverty, institutional, ecological, technological development, diffusion and transfer barriers (*medium evidence; medium agreement*) [11.7, 11.8]. Similarly, there are important feedbacks to adaptation, conservation of natural resources, such as water and terrestrial and aquatic biodiversity (*robust evidence; high agreement*) [11.5, 11.8]. There can be competition between different land uses if alternative options to use available land are mutually exclusive, but there are also potential synergies, e.g., integrated systems or multi-functionality at landscape scale (*medium evidence; high agreement*) [11.4]. Recent frameworks, such as those for assessing environmental or ecosystem services, provide one mechanism for valuing the multiple synergies and tradeoffs that may arise from mitigation actions (*medium evidence; medium agreement*) [11.1]. Sustainable management of agriculture, forests, and other land is an underpinning requirement of sustainable development (*robust evidence; high agreement*) [11.4].

AFOLU emissions could change substantially in transformation pathways, with significant mitigation potential from agriculture, forestry, and bioenergy mitigation measures (*medium evidence; high agreement*). Recent multi-model comparisons of idealized implementation transformation scenarios find land emissions (nitrous oxide, N₂O; methane, CH₄; CO₂) changing by –4 to 99% through 2030, and 7 to 76% through 2100, with the potential for increased emissions from land carbon stocks. Land-related mitigation, including bioenergy, could contribute 20 to 60% of total cumulative abatement to 2030, and 15 to 40% to 2100. However, policy coordination and implementation issues are challenges to realizing this potential [11.9]. Large-scale biomass supply for energy, or carbon sequestration in the AFOLU sector provide flexibility for the development of mitigation technologies in the energy supply and energy end-use sectors, as many technologies already exist and some of them are commercial (*limited evidence; medium agreement*) [11.3], but there are potential implications for biodiversity, food security, and other services provided by land (*medium evidence, high*

agreement) [11.7]. Implementation challenges, including institutional barriers and inertia related to governance issues, make the costs and net emission reduction potential of near-term mitigation uncertain. In mitigation scenarios with idealized comprehensive climate policies, agriculture, forestry, and bioenergy contribute substantially to the reduction of global CO₂, CH₄, and N₂O emissions, and to the energy system, thereby reducing policy costs (*medium evidence; high agreement*) [11.9]. More realistic partial and delayed policies for global land mitigation have potentially significant spatial and temporal leakage, and economic implications, but could still be cost-effectively deployed (*limited evidence; limited agreement*) [11.9].

Economic mitigation potential of supply-side measures in the AFOLU sector is estimated to be 7.18 to 10.60 (full range: 0.49–10.60) GtCO₂eq/yr in 2030 for mitigation efforts consistent with carbon prices up to 100 USD/tCO₂eq, about a third of which can be achieved at < 20 USD/tCO₂eq (*medium evidence; medium agreement*) [11.6]. These estimates are based on studies that cover both forestry and agriculture and that include agricultural soil carbon sequestration. Estimates from agricultural sector-only studies range from 0.3 to 4.6 GtCO₂eq/yr at prices up to 100 USD/tCO₂eq, and estimates from forestry sector-only studies from 0.2 to 13.8 GtCO₂eq/yr at prices up to 100 USD/tCO₂eq (*medium evidence; medium agreement*) [11.6]. The large range in the estimates arises due to widely different collections of options considered in each study, and because not all GHGs are considered in all of the studies. The composition of the agricultural mitigation portfolio varies with the carbon price, with the restoration of organic soils having the greatest potential at higher carbon prices (100 USD/tCO₂eq) and cropland and grazing land management at lower (20 USD/tCO₂eq). In forestry there is less difference between measures at different carbon prices, but there are significant differences between regions, with reduced deforestation dominating the forestry mitigation potential in Latin America and Caribbean (LAM) and Middle East and Africa (MAF), but very little potential in the member countries of the Organisation for Economic Co-operation and Development (OECD-1990) and Economies in Transition (EIT). Forest management, followed by afforestation, dominate in OECD-1990, EIT, and Asia (*medium evidence, strong agreement*) [11.6]. Among demand-side measures, which are under-researched compared to supply-side measures, changes in diet and reductions of losses in the food supply chain can have a significant, but uncertain, potential to reduce GHG emissions from food production (0.76–8.55 GtCO₂eq/yr by 2050), with the range being determined by assumptions about how the freed land is used (*limited evidence; medium agreement*) [11.4]. More research into demand-side mitigation options is merited. There are significant regional differences in terms of mitigation potential, costs, and applicability, due to differing local biophysical, socioeconomic, and

cultural circumstances, for instance between developed and developing regions, and among developing regions (*medium evidence; high agreement*) [11.6].

The size and regional distribution of future mitigation potential is difficult to estimate accurately because it depends on a number of inherently uncertain factors. Critical factors include population (growth), economic and technological developments, changes in behaviour over time (depending on cultural and normative backgrounds, market structures and incentives), and how these translate into demand for food, fibre, fodder and fuel, as well as development in the agriculture, aquaculture and forestry sectors. Other factors important to mitigation potential are potential climate change impacts on carbon stocks in soils and forests including their adaptive capacity (*medium evidence; high agreement*) [11.5]; considerations set by biodiversity and nature conservation requirements; and interrelations with land degradation and water scarcity (*robust evidence; high agreement*) [11.8].

Bioenergy can play a critical role for mitigation, but there are issues to consider, such as the sustainability of practices and the efficiency of bioenergy systems (*robust evidence, medium agreement*) [11.4.4, Box 11.5, 11.13.6, 11.13.7]. Barriers to large-scale deployment of bioenergy include concerns about GHG emissions from land, food security, water resources, biodiversity conservation and livelihoods. The scientific debate about the overall climate impact related to land use competition effects of specific bioenergy pathways remains unresolved (*robust evidence, high agreement*) [11.4.4, 11.13]. Bioenergy technologies are diverse and span a wide range of options and technology pathways. Evidence suggests that options with low lifecycle emissions (e.g., sugar cane, Miscanthus, fast growing tree species, and sustainable use of biomass residues), some already available, can reduce GHG emissions; outcomes are site-specific and rely on efficient integrated 'biomass-to-bioenergy systems', and sustainable land-use management and governance. In some regions, specific bioenergy options, such as improved cookstoves, and small-scale biogas and biopower production, could reduce GHG emissions and improve livelihoods and health in the context of sustainable development (*medium evidence, medium agreement*) [11.13].

Policies governing practices in agriculture and in forest conservation and management need to account for both mitigation and adaptation. One of the most visible current policies in the AFOLU sector is the implementation of REDD+ (see Annex I), that can represent a cost-effective option for mitigation (*limited evidence; medium agreement*) [11.10], with economic, social, and other environmental co-benefits (e.g., conservation of biodiversity and water resources).

11.1 Introduction

Agriculture, Forestry, and Other Land Use (AFOLU¹) plays a central role for food security and sustainable development (Section 11.9). Plants take up carbon dioxide (CO₂) from the atmosphere and nitrogen (N) from the soil when they grow, re-distributing it among different pools, including above and below-ground living biomass, dead residues, and soil organic matter. The CO₂ and other non-CO₂ greenhouse gases (GHG), largely methane (CH₄) and nitrous oxide (N₂O), are in turn released to the atmosphere by plant respiration, by decomposition of dead plant biomass and soil organic matter, and by combustion (Section 11.2). Anthropogenic land-use activities (e.g., management of croplands, forests, grasslands, wetlands), and changes in land use/cover (e.g., conversion of forest lands and grasslands to cropland and pasture, afforestation) cause changes superimposed on these natural fluxes. AFOLU activities lead to both sources of CO₂ (e.g., deforestation, peatland drainage) and sinks of CO₂ (e.g., afforestation, management for soil carbon sequestration), and to non-CO₂ emissions primarily from agriculture (e.g., CH₄ from livestock and rice cultivation, N₂O from manure storage and agricultural soils and biomass burning (Section 11.2).

The main mitigation options within AFOLU involve one or more of three strategies: *reduction/prevention* of emissions to the atmosphere by conserving existing carbon pools in soils or vegetation that would otherwise be lost or by reducing emissions of CH₄ and N₂O (Section 11.3); *sequestration*—enhancing the uptake of carbon in terrestrial reservoirs, and thereby removing CO₂ from the atmosphere (Section 11.3); and reducing CO₂ emissions by *substitution* of biological products for fossil fuels (Appendix 1) or energy-intensive products (Section 11.4). Demand-side options (e.g., by lifestyle changes, reducing losses and wastes of food, changes in human diet, changes in wood consumption), though known to be difficult to implement, may also play a role (Section 11.4).

Land is the critical resource for the AFOLU sector and it provides food and fodder to feed the Earth's population of ~7 billion, and fibre and fuel for a variety of purposes. It provides livelihoods for billions of people worldwide. It is finite and provides a multitude of goods and ecosystem services that are fundamental to human well-being (MEA, 2005). Human economies and quality of life are directly dependent on the services and the resources provided by land. Figure 11.1 shows the many provisioning, regulating, cultural and supporting services provided by land, of which climate regulation is just one. Implementing mitigation options in the AFOLU sector may potentially affect other services provided by land in positive or negative ways.

In the Intergovernmental Panel on Climate Change (IPCC) Second Assessment Report (SAR) (IPCC, 1996) and in the IPCC Fourth Assess-

ment Report (AR4) (IPCC, 2007a), agricultural and forestry mitigation were dealt with in separate chapters. In the IPCC Third Assessment Report (TAR) (IPCC, 2001), there were no separate sectoral chapters on either agriculture or forestry. In the IPCC Fifth Assessment Report (AR5), for the first time, the vast majority of the terrestrial land surface, comprising agriculture, forestry and other land use (AFOLU) (IPCC, 2006), is considered together in a single chapter, though settlements (which are important, with urban areas forecasted to triple in size from 2000 global extent by 2030; Section 12.2), are dealt with in Chapter 12. This approach ensures that all land-based mitigation options can be considered together; it minimizes the risk of double counting or inconsistent treatment (e.g., different assumptions about available land) between different land categories, and allows the consideration of systemic feedbacks between mitigation options related to the land surface (Section 11.4). Considering AFOLU in a single chapter allows phenomena common across land-use types, such as competition for land (Smith et al., 2010; Lambin and Meyfroidt, 2011) and water (e.g., Jackson et al., 2007), co-benefits (Sandor et al., 2002; Venter et al., 2009), adverse side-effects (Section 11.7) and interactions between mitigation and adaptation (Section 11.5) to be considered consistently. The complex nature of land presents a unique range of barriers and opportunities (Section 11.8), and policies to promote mitigation in the AFOLU sector (Section 11.10) need to take account of this complexity.

In this chapter, we consider the competing uses of land for mitigation and for providing other services (Sections 11.7; 11.8). Unlike the chapters on agriculture and forestry in AR4, impacts of sourcing bioenergy from the AFOLU sector are considered explicitly in a dedicated appendix (Section 11.13). Also new to this assessment is the explicit consideration of food/dietary demand-side options for GHG mitigation in the AFOLU sector (Section 11.4), and some consideration of freshwater fisheries and aquaculture, which may compete with the agriculture and forestry sectors, mainly through their requirements for land and/or water, and indirectly, by providing fish and other products to the same markets as animal husbandry.

This chapter deals with AFOLU in an integrated way with respect to the underlying scenario projections of population growth, economic growth, dietary change, land-use change (LUC), and cost of mitigation. We draw evidence from both 'bottom-up' studies that estimate mitigation potentials at small scales or for individual options or technologies and then scale up, and multi-sectoral 'top-down' studies that consider AFOLU as just one component of a total multi-sector system response (Section 11.9). In this chapter, we provide updates on emissions trends and changes in drivers and pressures in the AFOLU sector (Section 11.2), describe the practices available in the AFOLU sector (Section 11.3), and provide refined estimates of mitigation costs and potentials for the AFOLU sector, by synthesising studies that have become available since AR4 (Section 11.6). We conclude the chapter by identifying gaps in knowledge and data (Section 11.11), providing a selection of Frequently Asked Questions (Section 11.12), and presenting an Appendix on bioenergy to update the IPCC Special Report on Renewable Energy Sources and Climate Change Mitigation (SRREN) (IPCC, 2011; see Section 11.13).

¹ The term AFOLU used here consistent with the (IPCC, 2006) Guidelines is also consistent with Land Use, Land-Use Change and Forestry (LULUCF) (IPCC, 2003), and other similar terms used in the scientific literature.




Figure 11.1 | Multiple ecosystem services, goods and benefits provided by land (after MEA, 2005; UNEP-WCMC, 2011). Mitigation actions aim to enhance climate regulation, but this is only one of the many functions fulfilled by land.

11.2 New developments in emission trends and drivers

Estimating and reporting the anthropogenic component of gross and net AFOLU GHG fluxes to the atmosphere, globally, regionally, and at country level, is difficult compared to other sectors. First, it is not always possible to separate anthropogenic and natural GHG fluxes from land. Second, the input data necessary to estimate GHG emissions globally and regionally, often based on country-level statistics or on remote-sensing information, are very uncertain. Third, methods for estimating GHG emissions use a range of approaches, from simple default methodologies such as those specified in the IPCC GHG Guide-

lines² (IPCC, 2006), to more complex estimates based on terrestrial carbon cycle modelling and/or remote sensing information. Global trends in total GHG emissions from AFOLU activities between 1971 and 2010 are shown in Figure 11.2; Figure 11.3 shows trends of major drivers of emissions.

² Parties to the United Nations Framework Convention on Climate Change (UNFCCC) report net GHG emissions according to IPCC methodologies (IPCC, 2006). Reporting is based on a range of methods and approaches dependent on available data and national capacities, from default equations and emission factors applicable to global or regional cases and assuming instantaneous emissions of all carbon that will be eventually lost from the system following human action (Tier 1) to more complex approaches such as model-based spatial analyses (Tier 3).





Figure 11.2 | Top: AFOLU emissions for the last four decades. For the agricultural sub-sectors emissions are shown for separate categories, based on FAOSTAT, (2013). Emissions from crop residues, manure applied to soils, manure left on pasture, cultivated organic soils, and synthetic fertilizers are typically aggregated to the category ‘agricultural soils’ for IPCC reporting. For the Forestry and Other Land Use (FOLU) sub-sector data are from the Houghton bookkeeping model results (Houghton et al., 2012). Emissions from drained peat and peat fires are, for the 1970s and the 1980s, from JRC/PBL (2013), derived from Hooijer et al. (2010) and van der Werf et al. (2006) and for the 1990s and the 2000s, from FAOSTAT, 2013. Bottom: Emissions from AFOLU for each RCS region (see Annex II.2) using data from JRC/PBL (2013), with emissions from energy end-use in the AFOLU sector from IEA (2012a) included in a single aggregated category, see Annex II.9, used in the AFOLU section of Chapter 5.7.4 for cross-sectoral comparisons. The direct emission data from JRC/PBL (2013; see Annex II.9) represents land-based CO₂ emissions from forest and peat fires and decay that approximate to CO₂ flux from anthropogenic emission sources in the FOLU sub-sector. Differences between FAOSTAT/Houghton data and JRC/PBL (2013) are discussed in the text. See Figures 11.4 and 11.6 for the range of differences among available databases for AFOLU emissions.




Figure 11.3 | Global trends from 1971 to 2010 in (top) area of land use (forest land—available only from 1990; 1000 Mha) and amount of N fertilizer use (million tonnes), and (bottom) number of livestock (million heads) and poultry (billion heads). Data presented by regions: 1) Asia, 2) LAM, 3) MAF, 4) OECD-1990, 5) EIT (FAOSTAT, 2013). The area extent of AFOLU land-use categories, from FAOSTAT, (2013): ‘Cropland’ corresponds to the sum of FAOSTAT categories ‘arable land’ and ‘temporary crops’ and coincides with the IPCC category (IPCC, 2003); ‘Forest’ is defined according to FAO (2010); countries reporting to UNFCCC may use different definitions. ‘Permanent meadows and pasture’, are a subset of IPCC category ‘grassland’ (IPCC, 2003), as the latter, by definition, also includes unmanaged natural grassland ecosystems.

11.2.1 Supply and consumption trends in agriculture and forestry

In 2010 world agricultural land occupied 4889 Mha, an increase of 7% (311 Mha) since 1970 (FAOSTAT, 2013). Agricultural land area has decreased by 53 Mha since 2000 due to a decline of the cropland area (Organisation for Economic Co-operation and Development (OECD)-1990, Economies in Transition (EIT)) and a decrease in permanent meadows and pastures (OECD-1990 and Asia). The average amount of cropland and pasture land per capita in 1970 was 0.4 and 0.8 ha and by 2010 this had decreased to 0.2 and 0.5 ha per capita, respectively (FAOSTAT, 2013).

Changing land-use practices, technological advancement and varietal improvement have enabled world grain harvests to double from 1.2 to 2.5 billion tonnes per year between 1970 and 2010 (FAOSTAT, 2012). Average world cereal yields increased from 1600 to 3030 kg/ha over the same period (FAOSTAT, 2012) while there has also been a 233% increase in global fertilizer use from 32 to 106 Mt/yr, and a 73% increase in the irrigated cropland area (FAOSTAT, 2013).

Globally, since 1970, there has been a 1.4-fold increase in the numbers of cattle and buffalo, sheep and goats (which is closely linked to the trend of CH₄ emissions in the sector; Section 11.2.2), and increases of 1.6- and 3.7-fold for pigs and poultry, respectively (FAOSTAT, 2013). Major regional trends between 1970 and 2010 include a decrease in the total number of animals in Economies in Transition (EIT) and OECD-1990 (except poultry), and continuous growth in other regions, particularly Middle East and Africa (MAF) and Asia (Figure 11.3, bottom panel). The soaring demand for fish has led to the intensification of freshwater and marine fisheries worldwide, and an increased freshwater fisheries catch that topped 11 Mt in 2010, although the marine fisheries catch has slowly declined (78 Mt in 2010; FAOSTAT, 2013). The latter is, how-

ever, compensated in international markets by tremendous growth of aquaculture production to 60 Mt wet weight in 2010, of which 37 Mt originate from freshwater, overwhelmingly in Asia (FAOSTAT, 2013).

Between 1970 and 2010, global daily per capita food availability, expressed in energy units, has risen from 10,008 to 11,850 kJ (2391 to 2831 kcal), an increase of 18.4%; growth in MAF (10,716 kJ in 2010) has been 22%, and in Asia, 32% (11,327 kJ in 2010; FAOSTAT, 2013). The percentage of animal products in daily per capita total food consumption has increased consistently in Asia since 1970 (7 to 16%), remained constant in MAF (8%) and, since 1985, has decreased in OECD-1990 countries (32 to 28%), comprising, respectively, 1,790, 870 and 3,800 kJ in 2010 (FAOSTAT, 2013).

11.2.2 Trends of GHG emissions from agriculture

Organic and inorganic material provided as inputs or output in the management of agricultural systems are typically broken down through bacterial processes, releasing significant amounts of CO₂, CH₄, and N₂O to the atmosphere. Only agricultural non-CO₂ sources are reported as anthropogenic GHG emissions, however. The CO₂ emitted is considered neutral, being associated to annual cycles of carbon fixation and oxidation through photosynthesis. The agricultural sector is the largest contributor to global anthropogenic non-CO₂ GHGs, accounting for 56% of emissions in 2005 (U.S. EPA, 2011). Other important, albeit much smaller non-CO₂ emissions sources from other AFOLU categories, and thus not treated here, include fertilizer applications in forests. Annual total non-CO₂ GHG emissions from agriculture in 2010 are estimated to be 5.2–5.8 GtCO₂eq/yr (FAOSTAT, 2013; Tubiello et al., 2013) and comprised about 10–12% of global anthropogenic emissions. Fossil fuel CO₂ emissions on croplands added another




Figure 11.4 | Data comparison between FAOSTAT (2013), U.S. EPA (2006), and EDGAR (JRC/PBL, 2013) databases for key agricultural emission categories, grouped as agricultural soils, enteric fermentation, manure management systems, and rice cultivation, for 2005 | Whiskers represent 95% confidence intervals of global aggregated categories, computed using IPCC guidelines (IPCC, 2006) for uncertainty estimation (from Tubiello et al., 2013).

0.4–0.6 GtCO₂eq/yr in 2010 from agricultural use in machinery, such as tractors, irrigation pumps, etc. (Ceschia et al., 2010; FAOSTAT, 2013), but these emissions are accounted for in the energy sector rather than the AFOLU sector. Between 1990 and 2010, agricultural non-CO₂ emissions grew by 0.9%/yr, with a slight increase in growth rates after 2005 (Tubiello et al., 2013).

Three independent sources of disaggregated non-CO₂ GHG emissions estimates from agriculture at global, regional, and national levels are available. They are mostly based on FAOSTAT activity data and IPCC Tier 1 approaches (IPCC, 2006; FAOSTAT, 2012; JRC/PBL, 2013; U.S. EPA, 2013). EDGAR and FAOSTAT also provide data at country level. Estimates of global emissions for enteric fermentation, manure management and manure, estimated using IPCC Tier 2/3 approaches are also available (e.g., Herrero et al., 2013). The FAOSTAT, EDGAR and U.S. EPA estimates are slightly different, although statistically consistent given the large uncertainties in IPCC default methodologies (Tubiello et al., 2013). They cover emissions from enteric fermentation, manure deposited on pasture, synthetic fertilizers, rice cultivation, manure management, crop residues, biomass burning, and manure applied to soils. Enteric fermentation, biomass burning, and rice cul-

tivation are reported separately under IPCC inventory guidelines, with the remaining categories aggregated into ‘agricultural soils’. According to EDGAR and FAOSTAT, emissions from enteric fermentation are the largest emission source, while US EPA lists emissions from agricultural soils as the dominant source (Figure 11.4).

The following analyses refer to annual total non-CO₂ emissions by all categories. All three databases agree that that enteric fermentation and agricultural soils represent together about 70 % of total emissions, followed by paddy rice cultivation (9–11 %), biomass burning (6–12 %) and manure management (7–8 %). If all emission categories are disaggregated, both EDGAR and FAOSTAT agree that the largest emitting categories after enteric fermentation (32–40 % of total agriculture emissions) are manure deposited on pasture (15 %) and synthetic fertilizer (12 %), both contributing to emissions from agricultural soils. Paddy rice cultivation (11 %) is a major source of global CH₄ emissions, which in 2010 were estimated to be 493–723 MtCO₂eq/yr. The lower end of the range corresponds to estimates by FAO (FAOSTAT, 2013), with EDGAR and US EPA data at the higher end. Independent analyses suggest that emissions from rice may be at the lower end of the estimated range (Yan et al., 2009).




Figure 11.5 | Regional data comparisons for key agricultural emission categories in 2010 | Whiskers represent 95% confidence intervals computed using IPCC guidelines (IPCC, 2006; Tubiello et al., 2013). The data show that most of the differences between regions and databases are of the same magnitude as the underlying emission uncertainties. [FAOSTAT, 2013; JRC/PBL, 2013; U.S. EPA, 2013]

Enteric Fermentation. Global emissions of this important category grew from 1.4 to 2.1 GtCO₂eq/yr between 1961 and 2010, with average annual growth rates of 0.70% (FAOSTAT, 2013). Emission growth slowed during the 1990s compared to the long-term average, but became faster again after the year 2000. In 2010, 1.0–1.5 GtCO₂eq/yr (75% of the total emissions), were estimated to come from developing countries (FAOSTAT, 2013). Over the period 2000–2010, Asia and the Americas contributed most, followed by Africa and Europe (FAOSTAT, 2013); see Figure 11.5). Emissions have grown most in Africa, on average 2.4%/yr. In both Asia (2.0%/yr) and the Americas (1.1%/yr), emissions grew more slowly, and decreased in Europe (–1.7%/yr). From 2000 to 2010, cattle contributed the largest share (75% of the total), followed by buffalo, sheep and goats (FAOSTAT, 2013).

Manure. Global emissions from manure, as either organic fertilizer on cropland or manure deposited on pasture, grew between 1961 and 2010 from 0.57 to 0.99 GtCO₂eq/yr. Emissions grew by 1.1%/yr on average. Manure deposited on pasture led to far larger emissions than manure applied to soils as organic fertilizer, with 80% of emissions from deposited manures coming from developing countries (FAOSTAT, 2013; Herrero et al., 2013). The highest emitting regions from 2000–2010 were the Americas, Asia and Africa. Growth over the same period was most pronounced in Africa, with an average of 2.5%/yr, followed by Asia (2.3%/yr), and the Americas (1.2%/yr), while there was a decrease in Europe of –1.2%/yr. Two-thirds of the total came from grazing cattle, with smaller contributions from sheep and goats. In this decade, emissions from manure applied to soils as organic fertilizer were greatest in Asia, then in Europe and the Americas. Though the continent with the highest growth rates of 3.4%/yr, Africa's share in total emissions remained small. In this sub-category, swine and cattle contributed more than three quarters (77%) of the emissions. Emissions from manure management grew from 0.25 to 0.36 GtCO₂eq/yr, resulting in average annual growth rates of only 0.6%/yr during the period 1961–2010. From 2000–2010 most emissions came from Asia, then Europe, and the Americas (Figure 11.5).

Synthetic Fertilizer. Emissions from synthetic fertilizers grew at an average rate of 3.9%/yr from 1961 to 2010, with absolute values increasing more than 9-fold, from 0.07 to 0.68 GtCO₂eq/yr (Tubiello et al., 2013). Considering current trends, synthetic fertilizers will become a larger source of emissions than manure deposited on pasture in less than 10 years and the second largest of all agricultural emission categories after enteric fermentation. Close to three quarters (70%) of these emissions were from developing countries in 2010. In the decade 2000–2010, the largest emitter by far was Asia, then the Americas and then Europe (FAOSTAT, 2012). Emissions grew in Asia by 5.3%/yr, in Africa by 2.0%/yr, and in the Americas by 1.5%/yr. Emissions decreased in Europe (–1.8%/yr).

Rice. Emissions from rice are limited to paddy rice cultivation. From 1961 to 2010, global emissions increased with average annual growth rates of 0.4%/yr (FAOSTAT, 2013) from 0.37 to 0.52 GtCO₂eq/yr. The growth in global emissions has slowed in recent decades, consistent with trends in rice cultivated area. During 2000–2010, the largest share of emissions (94%) came from developing countries, with Asia being responsible for almost 90% of the total (Figure 11.5). The largest growth of emissions took place in in Africa (2.7%/yr), followed by Europe (1.4%/yr). Growth rates in Asia and the Americas were much smaller over the same period (0.4–0.7%/yr).




Figure 11.6 | Global net CO₂ emission estimates from FOLU including LUC. Black line: Houghton bookkeeping model approach updated to 2010 as in (Houghton et al., 2012), including LUC and forest management but no peatlands. Red lines: EDGAR 'LULUCF' emissions derived from the GFED 2.0 database (van der Werf et al., 2006) of emissions due to all forest fires (includes both FOLU and non-FOLU fires), with (solid line) and without (dotted line) peat fires and decay. Green lines: emissions from land-use change and management from FAO agricultural and forest inventory data (FAOSTAT, 2013), shown with (solid line) and without (dotted line) peat fires and peat degradation. Dark red line: deforestation and degradation fires only based on satellite fire data from GFED 3.0 database (van der Werf et al., 2010). Light blue lines: a selection of process-based vegetation model results, updated for WGI Chapter 6; (Le Quéré et al., 2013) include LUC, some include forest management, none include peatlands. LPJ-wsl: (Poulter et al., 2010); BernCC: (Stocker et al., 2011); VISIT: (Kato et al., 2011); ISAM: (Jain et al., 2013), IMAGE 2.4 (Van Minnen et al., 2009, deforestation only). The symbols and transparent rectangles represent mean values for the tropics only. Circles: tropical deforestation and forest management (Pan et al., 2011), using the Houghton (2003) bookkeeping model approach and FAO data. Triangle: tropical deforestation only, based on satellite forest area and biomass data (Baccini et al., 2012; Harris et al., 2012). Square: tropical deforestation and forest management, based on satellite forest area and biomass data and FAO data using bookkeeping model (Baccini et al., 2012; Harris et al., 2012).

Table 11.1 | Net global CO₂ flux from AFOLU.

	1750–2011	1980–1989	1990–1999	2000–2009
	Cumulative GtCO ₂	GtCO ₂ /yr	GtCO ₂ /yr	GtCO ₂ /yr
IPCC WGI Carbon Budget, Table 6.1^a:				
Net AFOLU CO ₂ flux ^b	660 ± 293	5.13 ± 2.93	5.87 ± 2.93	4.03 ± 2.93
Residual terrestrial sink ^c	-550 ± 330	-5.50 ± 4.03	-9.90 ± 4.40	-9.53 ± 4.40
Fossil fuel combustions and cement production ^d	1338 ± 110	20.17 ± 1.47	23.47 ± 1.83	28.60 ± 2.20
Meta-analyses of net AFOLU CO₂ flux:				
WGI, Table 6.2 ^e		4.77 ± 2.57	4.40 ± 2.20	2.93 ± 2.20
Houghton et al., 2012 ^f		4.18 ± 1.83	4.14 ± 1.83	4.03 ± 1.83

Notes: Positive fluxes represent net emissions and negative fluxes represent net sinks.

^(a) Selected components of the carbon budget in IPCC WGI AR5, Chapter 6, Table 6.1.

^(b) From the bookkeeping model accounting method of Houghton (2003) updated in Houghton et al., (2012), uncertainty based on expert judgement; 90 % confidence uncertainty interval.

^(c) Calculated as residual of other terms in the carbon budget.

^(d) Fossil fuel flux shown for comparison (Boden et al., 2011).

^(e) Average of estimates from 12 process models, only 5 were updated to 2009 and included in the 2000–2009 mean. Uncertainty based on standard deviation across models, 90 % confidence uncertainty interval (WGI Chapter 6).

^(f) Average of 13 estimates including process models, bookkeeping model and satellite/model approaches, only four were updated to 2009 and included in the 2000–2009 mean. Uncertainty based on expert judgment.

11.2.3 Trends of GHG fluxes from forestry and other land use³

This section focuses on the most significant non-agricultural GHG fluxes to the atmosphere for which there are global trend data. Fluxes resulting directly from anthropogenic FOLU activity are dominated by CO₂ fluxes, primarily emissions due to deforestation, but also uptake due to reforestation/regrowth. Non-CO₂ greenhouse gas emissions from FOLU are small in comparison, and mainly arise from peat degradation through drainage and biomass fires (Box 11.1; Box 11.2).

FOLU accounted for about a third of anthropogenic CO₂ emissions from 1750 to 2011 and 12 % of emissions in 2000 to 2009 (Table 11.1). At the same time, atmospheric measurements indicate the land as a whole was a net sink for CO₂, implying a 'residual' terrestrial sink offsetting FOLU emissions (Table 11.1). This sink is confirmed by inventory measurements in both managed and unmanaged forests in temperate and tropical regions (Phillips et al., 1998; Luysaert et al., 2008; Lewis et al., 2009; Pan et al., 2011). A sink of the right order of magnitude has been accounted for in models as a result of the indirect effects of human activity on ecosystems, i.e., the fertilizing effects of increased levels of CO₂ and N in the atmosphere and the effects of climate change (WGI Chapter 6; (Le Quéré et al., 2013), although some of it may be due to direct AFOLU activities not accounted for in current estimates (Erb et al., 2013). This sink capacity of forests is relevant to AFOLU mitigation through forest protection.

Global FOLU CO₂ flux estimates (Table 11.1 and Figure 11.6) are based on a wide range of data sources, and include different processes, definitions, and different approaches to calculating emissions (Houghton et al., 2012; Le Quéré et al., 2013; Pongratz et al., 2013). This leads to a large range across global FOLU flux estimates. Nonetheless, most approaches agree that there has been a decline in FOLU CO₂ emissions over the most recent years. This is largely due to a decrease in the rate of deforestation (FAO, 2010; FAOSTAT, 2013).

Regional trends in FOLU CO₂ emissions are shown in Figure 11.7. Model results indicate FOLU emissions peaked in the 1980s in Asia and LAM regions and declined thereafter. This is consistent with a reduced rate of deforestation, most notably in Brazil⁴, and some areas of afforestation, the latter most notably in China, Vietnam and India (FAOSTAT, 2013). In MAF the picture is mixed, with the Houghton model (Houghton et al., 2012) showing a continuing increase from the 1970s to the 2000s, while the VISIT model (Kato et al., 2011) indicates a small sink in the 2000s. The results for temperate and boreal areas represented by OECD and EIT regions are very mixed ranging from large net sources (ISAM) to small net sinks. The general picture in temperate and boreal regions is of declining emissions and/or increasing sinks. These regions include large areas of managed forests subjected to harvest and regrowth, and areas of reforestation (e.g., following cropland abandonment in the United States and Europe). Thus results are sensitive to whether and how the models include forest management and environmental effects on regrowing forests.

³ The term 'forestry and other land use' used here, is consistent with AFOLU in the (IPCC, 2006) Guidelines and consistent with LULUCF (IPCC, 2003).

⁴ For annual deforestation rates in Brazil see <http://www.obt.inpe.br/prodes/index.php>




Figure 11.7 | Regional trends in net CO₂ fluxes from FOLU (including LUC). Houghton bookkeeping model approach updated to 2010 as in Houghton et al., (2012) and five process-based vegetation models updated to 2010 for WGI Chapter 6; (Le Quéré et al., 2013): LPJ-wsl: (Poulter et al., 2010); BernCC: (Stocker et al., 2011); VISIT: (Kato et al., 2011); ISAM: (Jain et al., 2013), IMAGE 2.4: ((Van Minnen et al., 2009), deforestation only). Only the FAO estimates (FAOSTAT, 2013) include peatlands.

The bookkeeping model method (Houghton, 2003; Houghton et al., 2012) uses regional biomass, growth and decay rates from the inventory literature that are not varied to account for changes in climate or CO₂. It includes forest management associated with shifting cultivation in tropical forest regions as well as global wood harvest and regrowth cycles. The primary source of data for the most recent decades is FAO forest area and wood harvest (FAO, 2010). FAOSTAT (2013) uses the default IPCC methodologies to compute stock-difference to estimate emissions and sinks from forest management, carbon loss associated with forest conversion to other land uses as a proxy for emissions from deforestation, GFED4 data on burned area to estimate emissions from peat fires, and spatial analyses to determine emissions from drained organic soils (IPCC, 2007b). The other models in Figures 11.6 and 11.7 are process-based terrestrial ecosystem models that simulate changing plant biomass and carbon fluxes, and include climate and CO₂ effects, with a few now including the nitrogen cycle (Zaehle et al., 2011; Jain et al., 2013). Inclusion of the nitrogen cycle results in much higher modelled net emissions in the ISAM model (Jain et al., 2013) as N limitation due to harvest removals limits forest regrowth rates, particularly in temperate and boreal forests. Change in land cover in the process models is from the HYDE dataset (Goldewijk et al., 2011; Hurtt et al., 2011), based on FAO cropland and pasture area change data. Only some process models include forest management in terms of shifting cultivation (VISIT) or wood harvest and forest degradation (ISAM); none account for emissions from peatlands (see Box 11.1).

Satellite estimates of change in land cover have been combined with model approaches to calculate tropical forest emissions (Hansen et al., 2010). The data is high resolution and verifiable, but only covers recent decades, and does not account for fluxes due to LUC that occurred prior to the start of the study period (e.g., decay or regrowth). Satellite data alone cannot distinguish the cause of change in land use (deforestation, natural disturbance, management), but can be used in conjunction with activity data for attribution (Baccini et al., 2012). A recent development is the use of satellite-based forest biomass estimates (Saatchi et al., 2011) together with satellite land cover change in the tropics to estimate 'gross deforestation' emissions (Harris et al., 2012) or further combining it with FAO and other activity data to estimate net fluxes from forest area change and forest management (Baccini et al., 2012).

A detailed breakdown of the component fluxes in (Baccini et al., 2012) is shown in Figure 11.8. Where there is temporary forest loss through management, 'gross' forest emissions can be as high as for permanent forest loss (deforestation), but are largely balanced by 'gross' uptake in regrowing forest, so net emissions are small. When regrowth does not balance removals, it leads to a degradation of forest carbon stocks. In Baccini et al. (2012) this degradation was responsible for 15% of total net emissions from tropical forests (Houghton, 2013; Figure 11.8). Huang and Asner (2010) estimated that forest degradation in the Amazon, particularly from selective logging, is responsible for 15–19% higher C emissions than reported from deforestation alone. Pan et al.




Figure 11.8 | Breakdown of mean annual CO₂ fluxes from deforestation and forest management in tropical countries (GtCO₂/yr). Pan et al. (2011) estimates are based on FAO data and the Houghton bookkeeping model (Houghton, 2003). Baccini et al. (2012) estimates are based on satellite land cover change and biomass data with FAO data, and the Houghton (2003) bookkeeping model, with the detailed breakdown of these results shown in Houghton, (2013). Harris et al. (2012) estimates are based on satellite land cover change and biomass data.

Box 11.1 | AFOLU GHG emissions from peatlands and mangroves

Undisturbed waterlogged peatlands (organic soils) store a large amount of carbon and act as small net sinks (Hooijer et al., 2010). Drainage of peatlands for agriculture and forestry results in a rapid increase in decomposition rates, leading to increased emissions of CO₂ and N₂O, and vulnerability to further GHG emissions through fire. The FAO emissions database estimates globally 250,000 km² of drained organic soils under cropland and grassland, with total GHG emissions of 0.9 GtCO₂eq/yr in 2010—with the largest contributions from Asia (0.44 GtCO₂eq/yr) and Europe (0.18 GtCO₂eq/yr) (FAOSTAT, 2013). Joosten (2010), estimated that there are > 500,000 km² of drained peatlands in the world including under forests, with CO₂ emissions having increased from 1.06 GtCO₂/yr in 1990 to 1.30 GtCO₂/yr in 2008, despite a decreasing trend in Annex I countries, from 0.65 to 0.49 GtCO₂/yr,

primarily due to natural and artificial rewetting of peatlands. In Southeast Asia, CO₂ emissions from drained peatlands in 2006 were 0.61 ± 0.25 GtCO₂/yr (Hooijer et al., 2010). Satellite estimates indicate that peat fires in equatorial Asia emitted on average 0.39 GtCO₂ eq/yr over the period 1997–2009 (van der Werf et al., 2010), but only 0.2 GtCO₂ eq/yr over the period 1998–2009. This lower figure is consistent with recent independent FAO estimates over the same period and region. Mangrove ecosystems have declined in area by 20% (36 Mha) since 1980, although the rate of loss has been slowing in recent years, reflecting an increased awareness of the value of these ecosystems (FAO, 2007). A recent study estimated that deforestation of mangroves released 0.07 to 0.42 GtCO₂/yr (Donato et al., 2011).

Box 11.2. | AFOLU GHG emissions from fires

Burning vegetation releases CO₂, CH₄, N₂O, ozone-precursors and aerosols (including black carbon) to the atmosphere. When vegetation regrows after a fire, it takes up CO₂ and nitrogen. Anthropogenic land management or land conversion fire activities leading to permanent clearance or increasing levels of disturbance result in net emissions to the atmosphere over time. Satellite-detection of fire occurrence and persistence has been used to estimate fire emissions (e.g., GFED 2.0 database; (van der Werf et al., 2006). It is hard to separate the causes of fire as natural or anthropogenic, especially as the drivers are often combined. An update of the GFED methodology now distinguishes FOLU deforestation and degradation fires from other management fires (GFED 3.0 database; (van der Werf et al., 2010); Figure 11.6). The estimated tropical deforestation and degradation fire emissions

were 1.39 GtCO₂eq/yr during 1997 to 2009 (total carbon including CO₂, CH₄, CO and black carbon), 20% of all fire emissions. Carbon dioxide FOLU fire emissions are already included as part of the global models results such as those presented in Table 1.1 and Figures 11.6 and 11.7. According to (FAOSTAT, 2013)¹, in 2010 the non-CO₂ component of deforestation and forest degradation fires totalled 0.1 GtCO₂eq/yr, with forest management and peatland fires (Box 11.1) responsible for an additional 0.2 GtCO₂eq/yr.

¹ FOLU GHG emissions by fires include, as per IPCC GHG guidelines, all fires on managed land. Most current FOLU estimates are limited however to fires associated to deforestation, forest management and peat fires. Emissions from prescribed burning of savannahs are reported under agriculture. Both CO₂ and non-CO₂ emissions are accounted under these FOLU components, but CO₂ emissions dominate.

(2011) separated 'gross emissions' from deforestation and forest management on the one hand, from uptake in regrowing vegetation on the other. Deforestation emissions decline from the 1990s to 2000–2007, and uptake in regrowing vegetation increases, both contributing to the decline in net tropical CO₂ emissions.

Satellite fire data have also been used to estimate FOLU emissions (van der Werf et al., 2006); Box 11.2). The EDGAR⁵ database 'Land-

Use Change and Forestry' emissions are based on forest and peat fire data from GFED 2.0 (van der Werf et al., 2006), with additional estimates of post-burn decay, and emissions from degraded peatlands based on (Joosten, 2010); Box 11.1). However, GFED 2.0 fire data does not distinguish anthropogenic AFOLU fires from other fires, unlike GFED 3.0 (van der Werf et al., 2010); Box 11.2). Fire data also does not capture significant additional AFOLU fluxes due to land clearing and forest management that is by harvest rather than fire (e.g., deforestation activities outside the humid tropics) or regrowth following clearing. Thus EDGAR data only approximates the FOLU flux.

⁵ <http://edgar.jrc.ec.europa.eu/index.php>

FAO estimates AFOLU GHG emissions (FAOSTAT, 2013)⁶ based on IPCC Tier 1 methodology⁷. With reference to the decade 2001–2010, total GHG FOLU emissions were 3.2 GtCO₂eq/yr including deforestation (3.8 GtCO₂eq/yr), forest degradation and forest management (–1.8 GtCO₂eq/yr), biomass fires including peatland fires (0.3 GtCO₂eq/yr), and drained peatlands (0.9 GtCO₂eq/yr). The FAO estimated total mean net GHG FOLU flux to the atmosphere decreased from 3.9 GtCO₂eq/yr in 1991–2000 to 3.2 GtCO₂eq/yr in 2001–2010 (FAOSTAT, 2013).

11.3 Mitigation technology options and practices, and behavioural aspects

Greenhouse gases can be reduced by supply-side mitigation options (i.e., by reducing GHG emissions per unit of land/animal, or per unit of product), or by demand-side options (e.g., by changing demand for food and fibre products, reducing waste). In AR4, the forestry chapter (Nabuurs et al., 2007) considered some demand-side options, but the agriculture chapter focused on supply-side options only (Nabuurs et al., 2007; Smith et al., 2007). In this section, we discuss only supply-side options (Section 11.3.1). Demand-side options are discussed in Section 11.4.

Mitigation activities in the AFOLU sector can reduce climate forcing in different ways:

- Reductions in CH₄ or N₂O emissions from croplands, grazing lands, and livestock.
- Conservation of existing carbon stocks, e.g., conservation of forest biomass, peatlands, and soil carbon that would otherwise be lost.
- Reductions of carbon losses from biota and soils, e.g., through management changes within the same land-use type (e.g., reducing soil carbon loss by switching from tillage to no-till cropping) or by reducing losses of carbon-rich ecosystems, e.g., reduced deforestation, rewetting of drained peatlands.
- Enhancement of carbon sequestration in soils, biota, and long-lived products through increases in the area of carbon-rich ecosystems such as forests (afforestation, reforestation), increased carbon storage per unit area, e.g., increased stocking density in

forests, carbon sequestration in soils, and wood use in construction activities.

- Changes in albedo resulting from land-use and land-cover change that increase reflection of visible light.
- Provision of products with low GHG emissions that can replace products with higher GHG emissions for delivering the same service (e.g., replacement of concrete and steel in buildings with wood, some bioenergy options; see Section 11.13).
- Reductions of direct (e.g., agricultural machinery, pumps, fishing craft) or indirect (e.g., production of fertilizers, emissions resulting from fossil energy use in agriculture, fisheries, aquaculture, and forestry or from production of inputs); though indirect emission reductions are accounted for in the energy end-use sectors (buildings, industry, energy generation, transport) so are not discussed further in detail in this chapter.

11.3.1 Supply-side mitigation options

Mitigation potentials for agricultural mitigation options were given on a ‘per-area’ and ‘per-animal’ in AR4 (Nabuurs et al., 2007; Smith et al., 2007). All options are summarized in Table 11.2 with impacts on each GHG noted, and a categorization of technical mitigation potential, ease of implementation, and availability (supported by recent references). These mitigation options can have additive positive effects, but can also work in opposition, e.g., zero tillage can reduce the effectiveness of residue incorporation. Most mitigation options were described in detail in AR4 so are not described further here; additional practices that were not considered in AR4, i.e., biochar, reduced emissions from aquaculture, and bioenergy are described in Boxes 11.3, 11.4, and 11.5, respectively.

In addition to the per-area and per-animal mitigation options described in AR4, more attention has recently been paid to options that reduce emissions intensity by improving the efficiency of production (i.e., less GHG emissions per unit of agricultural product; (Burney et al., 2010; Bennetzen et al., 2012); a reduction in emissions intensity has long been a feature of agricultural emissions reduction and is one component of a process more broadly referred to as sustainable intensification (Tilman et al., 2009; Godfray et al., 2010; Smith, 2013; Garnett et al., 2013). This process does not rely on reducing inputs *per se*, but relies on the implementation of new practices that result in an increase in product output that is larger than any associated increase in emissions (Smith, 2013). Even though per-area emissions could increase, there is a net benefit since less land is required for production of the same quantity of product. The scope to reduce emissions intensity appears considerable since there are very large differences in emissions intensity between different regions of the world (Herrero et al., 2013). Sustainable intensification is discussed further in Section 11.4.2, and trends in changes in emissions intensity are discussed further in Section 11.6.

⁶ <http://faostat.fao.org/>

⁷ Parties to the UNFCCC report net GHG emissions according to IPCC methodologies (IPCC, 2003, 2006). Reporting is based on a range of methods and approaches dependent on available data and national capacities, from default equations and emission factors applicable to global or regional cases and assuming instantaneous emissions of all carbon that will be eventually lost from the system following human action (Tier 1) to more complex approaches such as model-based spatial analyses (Tier 3).

Table 11.2 | Summary of supply-side mitigation options in the AFOLU sector. Technical Mitigation Potential: Area = (tCO₂e/ha)/yr; Animal = percent reduction of enteric emissions. Low = < 1; < 5 % (white), Medium = 1–10; 5–15 % (light grey), High = > 10, > 15 % (grey); Ease of Implementation (acceptance or adoption by land manager): Difficult (white), Medium (light grey), Easy, i.e., universal applicability (grey); Timescale for Implementation: Long-term (at research and development stage; white), Mid-term (trials in place, within 5–10 years; light grey), Immediate (technology available now, grey).

Categories	Practices and Impacts	Technical Mitigation Potential	Ease of Implementation	Timescale for implementation	References
Forestry					
Reducing deforestation	C: Conservation of existing C pools in forest vegetation and soil by controlling deforestation protecting forest in reserves, and controlling other anthropogenic disturbances such as fire and pest outbreaks. Reducing slash and burn agriculture, reducing forest fires.				1
	CH ₄ , N ₂ O: Protection of peatland forest, reduction of wildfires.				2
Afforestation/Reforestation	C: Improved biomass stocks by planting trees on non-forested agricultural lands. This can include either monocultures or mixed species plantings. These activities may also provide a range of other social, economic, and environmental benefits.				3, 4, 5
Forest management	C: Management of forests for sustainable timber production including extending rotation cycles, reducing damage to remaining trees, reducing logging waste, implementing soil conservation practices, fertilization, and using wood in a more efficient way, sustainable extortion of wood energy				6, 7, 8, 9
	CH ₄ , N ₂ O: Wildfire behaviour modification.				10, 11, 12
Forest restoration	C: Protecting secondary forests and other degraded forests whose biomass and soil C densities are less than their maximum value and allowing them to sequester C by natural or artificial regeneration, rehabilitation of degraded lands, long-term fallows.				13, 14
	CH ₄ , N ₂ O: Wildfire behaviour modification.				
Land-based agriculture					
<i>Cropland management</i>					
Croplands—plant management	C: High input carbon practices, e.g., improved crop varieties, crop rotation, use of cover crops, perennial cropping systems, agricultural biotechnology.				15, 16, 17
	N ₂ O: Improved N use efficiency.				18
Croplands—nutrient management	C: Fertilizer input to increase yields and residue inputs (especially important in low-yielding agriculture).				19, 20
	N ₂ O: Changing N fertilizer application rate, fertilizer type, timing, precision application, inhibitors.				21, 22, 23, 24, 25, 105, 106
Croplands—tillage/residues management	C: Reduced tillage intensity; residue retention.				17, 24, 26, 27
	N ₂ O:				28, 96, 97
	CH ₄ :				96
Croplands—water management	C: Improved water availability in cropland including water harvesting and application.				29
	CH ₄ : Decomposition of plant residues.				
	N ₂ O: Drainage management to reduce emissions, reduce N runoff leaching.				
Croplands—rice management	C: Straw retention.				30
	CH ₄ : Water management, mid-season paddy drainage.				31, 32, 98
	N ₂ O: Water management, N fertilizer application rate, fertilizer type, timing, precision application.				32, 98, 99
Rewet peatlands drained for agriculture	C: Ongoing CO ₂ emissions from reduced drainage (but CH ₄ emissions may increase).				33
Croplands—set-aside and LUC	C: Replanting to native grasses and trees. Increase C sequestration.				34, 35, 36, 37, 38
	N ₂ O: N inputs decreased resulting in reduced N ₂ O.				
Biochar application	C: Soil amendment to increase biomass productivity, and sequester C (biochar was not covered in AR4 so is described in Box 11.3).				39, 40, 41
	N ₂ O: Reduced N inputs will reduce emissions.				39, 42



Categories	Practices and Impacts	Technical Mitigation Potential	Ease of Implementation	Timescale for implementation	References
<i>Grazing Land Management</i>					
Grazing lands—plant management	C: Improved grass varieties/sward composition, e.g., deep rooting grasses, increased productivity, and nutrient management. Appropriate stocking densities, carrying capacity, fodder banks, and improved grazing management.				43, 44, 45
	N ₂ O				46
Grazing lands—animal management	C: Appropriate stocking densities, carrying capacity management, fodder banks and improved grazing management, fodder production, and fodder diversification.				43, 47
	CH ₄				
	N ₂ O: Stocking density, animal waste management.				
Grazing land—fire management	C: Improved use of fire for sustainable grassland management. Fire prevention and improved prescribed burning.				
Revegetation					
Revegetation	C: The establishment of vegetation that does not meet the definitions of afforestation and reforestation (e.g., <i>Atriplex</i> spp.).				48
	CH ₄ : Increased grazing by ruminants may increase net emissions.				
	N ₂ O: Reduced N inputs will reduce emissions.				
Other					
Organic soils—restoration	C: Soil carbon restoration on peatlands; and avoided net soil carbon emissions using improved land management.				49
	CH ₄ : May increase.				
Degraded soils—restoration	Land reclamation (afforestation, soil fertility management, water conservation soil nutrients enhancement, improved fallow).				100, 101, 102, 103, 104
Biosolid applications	C: Use of animal manures and other biosolids for improved management of nitrogen; integrated livestock agriculture techniques.				26
	N ₂ O				
Livestock					
Livestock—feeding	CH ₄ : Improved feed and dietary additives to reduce emissions from enteric fermentation; including improved forage, dietary additives (bioactive compounds, fats), ionophores/antibiotics, propionate enhancers, archaea inhibitors, nitrate and sulphate supplements.				50, 51, 52, 53, 54, 55, 56, 57, 58, 59
Livestock—breeding and other long-term management	CH ₄ : Improved breeds with higher productivity (so lower emissions per unit of product) or with reduced emissions from enteric fermentation; microbial technology such as archaeal vaccines, methanotrophs, acetogens, defaunation of the rumen, bacteriophages and probiotics; improved fertility.				54, 55, 56, 58, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71
Manure management	CH ₄ : Manipulate bedding and storage conditions, anaerobic digesters; biofilters, dietary additives.				56, 58, 72, 73
	N ₂ O: Manipulate livestock diets to reduce N excreta, soil applied and animal fed nitrification inhibitors, urease inhibitors, fertilizer type, rate and timing, manipulate manure application practices, grazing management.				56, 58, 72, 74, 75, 76, 77, 78
Integrated systems					
Agroforestry (including agropastoral and agrosilvopastoral systems)	C: Mixed production systems can increase land productivity and efficiency in the use of water and other resources and protect against soil erosion as well as serve carbon sequestration objectives.				79, 80, 81, 82, 83, 84, 85, 86, 87, 88
	N ₂ O: Reduced N inputs will reduce emissions.				
Other mixed biomass production systems	C: Mixed production systems such as double-cropping systems and mixed crop-livestock systems can increase land productivity and efficiency in the use of water and other resources as well as serve carbon sequestration objectives. Perennial grasses (e.g., bamboo) can in the same way as woody plants be cultivated in shelter belts and riparian zones/buffer strips provide environmental services and supports C sequestration and biomass production.				82, 89, 90
	N ₂ O: Reduced N inputs will reduce emissions.				



Categories	Practices and Impacts	Technical Mitigation Potential	Ease of Implementation	Timescale for Implementation	References
Integration of biomass production with subsequent processing in food and bioenergy sectors	C: Integrating feedstock production with conversion, typically producing animal feed that can reduce demand for cultivated feed such as soy and corn and can also reduce grazing requirements. Using agricultural and forestry residues for energy production.				91, 92, 93, 94, 95
	N ₂ O: Reduced N inputs will reduce emissions.				
Bioenergy (see Box 11.5 and Section 11.13)					

¹Van Bodegom et al., 2009; ²Malmshheimer et al., 2008; ³Reyer et al., 2009; ⁴Sochacki et al., 2012; ⁵IPCC, 2000; ⁶DeFries and Rosenzweig, 2010; ⁷Takimoto et al., 2008; ⁸Masera et al., 2003; ⁹Silver et al., 2000; ¹⁰Dezzeb et al., 2005; ¹¹Ito, 2005; ¹²Sow et al., 2013; ¹³Reyer et al., 2009; ¹⁴Palm et al., 2004; ¹⁵Godfray et al., 2010; ¹⁶Burney et al., 2010; ¹⁷Conant et al., 2007; ¹⁸Huang and Tang, 2010; ¹⁹Lemke et al., 2010; ²⁰Eagle and Olander, 2012; ²¹Snyder et al., 2007; ²²Akiyama et al., 2010; ²³Barton et al., 2011; ²⁴Powlson et al., 2011; ²⁵van Kessel et al., 2013; ²⁶Farage et al., 2007; ²⁷Smith, 2012; ²⁸Abdalla et al., 2013; ²⁹Bayala et al., 2008; ³⁰Yagi et al., 1997; ³¹Tyagi et al., 2010; ³²Feng et al., 2013; ³³Lohilla et al., 2004; ³⁴Seaquist et al., 2008; ³⁵Mbow, 2010; ³⁶Assogbadjo et al., 2012; ³⁷Laganriere et al., 2010; ³⁸Bayala et al., 2011; ³⁹Singh et al., 2010; ⁴⁰Woolf et al., 2010; ⁴¹Lehmann et al., 2003; ⁴²Taghizadeh-Toosi et al., 2011; ⁴³Franzluebbers and Stuedemann, 2009; ⁴⁴Follett and Reed, 2010; ⁴⁵McSherry and Ritchie, 2013; ⁴⁶Saggar et al., 2004; ⁴⁷Thornton and Herrero, 2010; ⁴⁸Harper et al., 2007; ⁴⁹Smith and Wollenberg, 2012; ⁵⁰Beauchemin et al., 2008; ⁵¹Beauchemin et al., 2009; ⁵²Martin et al., 2010; ⁵³Grainger and Beauchemin, 2011; ⁵⁴Clark, 2013; ⁵⁵Cottle et al., 2011; ⁵⁶Eckard et al., 2010; ⁵⁷Sauvant and Giger-Reverdin, 2007; ⁵⁸Hristov et al., 2013; ⁵⁹Bryan et al., 2013; ⁶⁰Attwood and McSweeney, 2008; ⁶¹Attwood et al., 2011; ⁶²Hegarty et al., 2007; ⁶³Hook et al., 2010; ⁶⁴Janssen and Kirs, 2008; ⁶⁵Martin et al., 2010; ⁶⁶Morgavi et al., 2008; ⁶⁷Morgavi et al., 2010; ⁶⁸Place and Mitloehner, 2010; ⁶⁹Waghorn and Hegarty, 2011; ⁷⁰Wright and Klieve, 2011; ⁷¹Yan et al., 2010; ⁷²Chadwick et al., 2011; ⁷³Petersen and Sommer, 2011; ⁷⁴de Klein et al., 2010; ⁷⁵de Klein and Eckard, 2008; ⁷⁶Dijkstra et al., 2011; ⁷⁷Schils et al., 2013; ⁷⁸VanderZaag et al., 2011; ⁷⁹Oke and Odebiyi, 2007; ⁸⁰Rice, 2008; ⁸¹Takimoto et al., 2008; ⁸²Lott et al., 2009; ⁸³Sood and Mitchell, 2011; ⁸⁴Assogbadjo et al., 2012; ⁸⁵Wollenberg et al., 2012; ⁸⁶Semroc et al., 2012; ⁸⁷Souza et al., 2012; ⁸⁸Luedeling and Neufeldt, 2012; ⁸⁹Heggenstaller et al., 2008; ⁹⁰Herrero et al., 2010; ⁹¹Dale et al., 2009; ⁹²Dale et al., 2010; ⁹³Sparovek et al., 2007; ⁹⁴Sood and Mitchell, 2011; ⁹⁵Vermeulen et al., 2012; ⁹⁶Metay et al., 2007; ⁹⁷Rochette, 2008; ⁹⁸Ma et al., 2009; ⁹⁹Yao et al., 2010; ¹⁰⁰Arnalds, 2004; ¹⁰¹Batjes, 2004; ¹⁰²Hardner et al., 2000; ¹⁰³May et al., 2004; ¹⁰⁴Zhao et al., 2005; ¹⁰⁵Huang and Tang, 2010; ¹⁰⁶Kim et al., 2013.

11.3.2 Mitigation effectiveness (non-permanence: saturation, human and natural impacts, displacement)

Since carbon sequestration in soil and vegetation and the retention of existing carbon stocks forms a significant component of the mitigation potential in the AFOLU sector, this section considers the factors affecting this strategy compared to avoided GHG emissions.

Non-permanence/reversibility. Reversals are the release of previously sequestered carbon, which negates some or all of the benefits from sequestration that has occurred in previous years. This issue is sometimes referred to as 'non-permanence' (Smith, 2005). Various types of carbon sinks (e.g., afforestation/reforestation, agricultural soil C) have an inherent risk of future reversals.

Certain types of mitigation activities (e.g., avoided N₂O from fertilizer, emission reductions from changed diet patterns or reduced food-chain losses) are effectively permanent since the emissions, once avoided, cannot be re-emitted. The same applies to the use of bioenergy to displace fossil-fuel emissions (Section 11.13) or the use of biomass-based products to displace more emissions-intensive products (e.g., wood in place of concrete or steel) in construction.

Reversals may be caused by natural events that affect yields/growth. In some cases (e.g., frost damage, pest infestation, or fire; (Reichstein

et al., 2013), these effects may be temporary or short-term. Although these events will affect the annual increment of C sequestration, they may not result in a permanent decline in carbon stocks. In other cases, such as stand replacing forest fires, insect or disease outbreaks, or drought, the declines may be more profound. Although a substantial loss of above-ground stored carbon could occur following a wildfire, whether this represents a loss depends on what happens following the fire and whether the forest recovers, or changes to a lower carbon-storage state (see Box 11.2). Similarly, some systems are naturally adapted to fire and carbon stocks will recover following fire, whereas in other cases the fire results in a change to a system with a lower carbon stock (e.g., Brown and Johnstone, 2011). For a period of time following fire (or other disruptive event), the stock of carbon will be less than that before the fire. Similarly, emissions of non-CO₂ gases also need to be considered.

The permanence of the AFOLU carbon stock relates to the longevity of the stock, i.e., how long the increased carbon stock remains in the soil or vegetation. This is linked to consideration of the reversibility of the increased carbon stock (Smith, 2005), as discussed in Section 11.5.2.

Saturation. Substitution of fossil fuel and material with biomass, and energy-intensive building materials with wood can continue in perpetuity. In contrast, it is often considered that carbon sequestration in soils (Guldea et al., 2008) or vegetation cannot continue indefinitely. The carbon stored in soils and vegetation reaches a new equilibrium (as the trees mature or as the soil carbon stock saturates). As the

Box 11.3 | Biochar

This box summarizes the mitigation potential for biochar technologies, which were not considered in AR4. Biomass C stabilization could be combined with (or substitute) bioenergy capture as part of a land-based mitigation strategy (Lehmann, 2007). Heating biomass with air excluded (pyrolysis) generates energy-containing volatiles and gases. Hydrogen and O are preferentially eliminated, creating a stable (biologically recalcitrant) C-rich co-product (char). By adding char to soil as 'biochar' a system can be established that may have a higher carbon abatement than typical bioenergy alternatives (Woolf et al., 2010). The gain is probably highest where efficient bioenergy is constrained by a remote, seasonal, or diffuse biomass resource (Shackley et al., 2012). The benefit of pyrolysis-biochar systems (PBS) is increased considerably if allowance is made for the indirect effects of using biochar *via* the soil. These effects include increased crop and biomass production and decreased N₂O and CH₄ emissions. Realizing the mitigation potential for biochar technologies will be constrained by the need for sustainable feedstock acquisition, competing biomass use options are an important influence of the production process on biochar properties. Considering sustainable feedstock production and targeting biochar deployment on less fertile land, Woolf et al. (2010) calculated maximum global abatement of 6.6 GtCO₂eq/yr from 2.27 Gt biomass C. Allowing for competition for virgin non-waste biomass the value was lower (3.67 GtCO₂eq/yr from 1.01 Gt biomass C), accruing 240–480 GtCO₂eq abatement within 100 years.

Meta-analysis shows that in experimental situations crop productivity has, on average, been enhanced by circa 15 % near-term, but with a wide range of effects (Jeffery et al., 2011; Biederman and Harpole, 2013). This range is probably explained by the nature and extent of pre-existing soil constraints. The Woolf et al. (2010) analysis accordingly assumed crop yield increases of 0–90 % (relative). Relaxing this assumption by one-half decreased projected abatement by 10 %. Decreasing an assumed 25 % suppression on soil N₂O flux by the same proportion had a smaller impact. Beneficial interactions of biochar and the soil N cycle are beginning

to be understood with effects on mineralization, nitrification, denitrification, immobilization and adsorption persisting at least for days and months after biochar addition (Nelissen et al., 2012; Clough et al., 2013). Although the often large suppression of soil N₂O flux observed under laboratory conditions can be increasingly explained (Cayuela et al., 2013), this effect is not yet predictable and there has been only limited validation of N₂O suppression by biochar in planted field soils (Liu et al., 2012; Van Zwieten et al., 2013) or over longer timeframes (Spokas, 2013). The potential to gain enhanced mitigation using biochar by tackling gaseous emissions from manures and fertilizers before and after application to soil are less well-explored (Steiner et al., 2010; Angst et al., 2013). The abatement potential for PBS remains most sensitive to the absolute stability of the C stored in biochar. Estimates of 'half-life' have been inferred from wildfire charcoal (Lehmann, 2007) or extrapolated from direct short-term observation. These give values that range from < 50 to > 10,000 years, but predominantly between 100–1000 years (Singh et al., 2012; Spokas, 2013). Nonetheless, the assumption made by Woolf et al. (2010) for the proportion of biochar C that is stable long-term (85 %) is subject to refinement and field validation.

Demonstration of the equipment and infrastructure required for effective use of energy products from biomass pyrolysis is still limited, especially across large and small unit scales. Preliminary analyses shows, however, that the break-even cost of biochar production is likely to be location- and feedstock-specific (Shackley et al., 2012; Field et al., 2013). Until economic incentives are established for the stabilization of C, biochar adoption will depend on predictable, positive effects on crop production. This requires more research on the use of biochar as a regular low-dose soil input, rather than single applications at rates > 10t/ha, which have so far been the norm (Sohi, 2012). Product standards are also required, to ensure that biochar is produced in a way that does not create or conserve problematic concentrations of toxic contaminants, and to support regulated deployment strategies (IBI Biochar, 2012; Downie et al., 2012).

soils/vegetation approach the new equilibrium, the annual removal (sometimes referred to as the sink strength) decreases until it becomes zero at equilibrium. This process is called saturation (Smith, 2005; Körner, 2006, 2009; Johnston et al., 2009b), and the uncertainty associated with saturation has been estimated (Kim and McCarl, 2009). An alternative view is that saturation does not occur, with studies from old-growth forests, for example, showing that they can continue to sequester C in soil and dead organic matter even if net living biomass

increment is near zero (e.g., Luysaert et al., 2008). Peatlands are unlikely to saturate in carbon storage, but the rate of C uptake may be very slow (see Box 11.1).

Human and natural impacts. Soil and vegetation carbon sinks can be impacted upon by direct human-induced, indirect human-induced, and natural changes (Smith, 2005). All of the mitigation practices discussed in Section 11.3.1 arise from direct human-induced impacts (deliberate

Box 11.4 | Aquaculture

Aquaculture is defined as the farming of fish, shellfish, and aquatic plants (Hu et al., 2013). Although it is an ancient practice in some parts of world, this sector of the food system is growing rapidly. Since the mid-1970s, total aquaculture production has grown at an average rate of 8.3 % per year (1970–2008; (Hu et al., 2013). The estimated aquaculture production in 2009 was 55.10 Mt, which accounts for approximately 47 % of all the fish consumed by humans (Hu et al., 2013). The sector is diverse, being dominated by shellfish and herbivorous and omnivorous pond fish, either entirely or partly utilizing natural productivity, but globalizing trade and favourable economic conditions are driving intensive farming at larger scales (Bostock et al., 2010). Potential impacts of aquaculture, in terms emissions of N₂O, have recently been considered (Williams and Crutzen, 2010; Hu et al., 2012). Global N₂O emissions from aquaculture in 2009 were estimated to be 93 ktN₂O-N (~43 MtCO₂eq), and will increase to 383 ktN₂O-N (~178 MtCO₂eq) by 2030, which could account for 5.7 % of anthropogenic N₂O-N emissions if aquaculture continues to grow at the present growth rate (~7.1 %/yr; Hu et al., 2012).

Some studies have focused on rice-fish farming, which is a practice associated with wet rice cultivation in Southeast Asia, providing protein, especially for subsistence-oriented farmers (Bhattacharyya et al., 2013). Cultivation of fish along with rice

increases emissions of CH₄ (Frei et al., 2007; Bhattacharyya et al., 2013), but decreases N₂O emissions, irrespective of the fish species used (Datta et al., 2009; Bhattacharyya et al., 2013). Although rice-fish farming systems might be globally important in terms of climate change, they are also relevant for local economy, food security, and efficient water use (shared water), which makes it difficult to design appropriate mitigation measures, because of the tradeoffs between mitigation measures and rice and fish production (Datta et al., 2009; Bhattacharyya et al., 2013). Feeding rate and dissolved oxygen (DO) concentration could affect N₂O emissions from aquaculture systems significantly, and nitrification and denitrification processes were equally responsible for the emissions of N₂O in these systems. Measures to control N₂O from aquaculture are described by Hu et al. (2012), and include the maintenance of optimal operating conditions of the system, such as appropriate pH and temperature, sufficient DO and good quality feed. Additionally, two potential ways to minimize N₂O emissions from aquaculture systems include 'Aquaponic Aquaculture' (polyculture consisting of fish tanks (aquaculture) and plants that are cultivated in the same water cycle (hydroponic)), and Bioflocs Technology (BFT) Aquaculture (which involves the development and control of heterotrophic bacteria in flocs within the fish culture component), where the growth of heterotrophic bacteria is stimulated, leading to nitrogen uptake (Hu et al., 2012).

management). Both sink processes and carbon stocks can be affected by natural factors such as soil and hydrological conditions. Indirect human-induced changes can impact carbon sinks and are influenced by human activity, but are not directly related to the management of that piece of land; examples include climate change and atmospheric nitrogen deposition. For some tree species, rising concentrations of tropospheric ozone caused by human activities may counteract the effects of increased atmospheric CO₂ or N deposition on tree growth (Sitch et al., 2008; Matyssek et al., 2010). Natural changes that threaten to impact the efficacy of mitigation measures are discussed in Section 11.5.

Displacement/leakage. Displacement/leakage arises from a change in land use or land management that causes a positive or negative change in emissions elsewhere. This can occur within or across national boundaries, and the efficacy of mitigation practices must consider the leakage implications. For example, if reducing emissions in one place leads to increased emissions elsewhere, no net reduction occurs; the emissions are simply displaced (Powlson et al., 2011; Kastner et al., 2011b; a). However, this assumes a one-to-one correspondence. Murray et al. (2004) estimated the leakage from different forest carbon programmes and this varied from < 10 % to > 90 % depending on the nature of the

activity. West et al. (2010a) examined the impact of displaced activities in different geographic contexts; for example, land clearing in the tropics will release twice the carbon, but only produce half the crop yield of temperate areas. Indirect land-use change is an important component to consider for displaced emissions and assessments of this are an emerging area. Indirect land-use change is discussed further in Section 11.4 and in relation to bioenergy in Section 11.13.

The timing of mitigation benefits from actions (e.g., bioenergy, forest management, forest products use/storage) can vary as a result both of the nature of the activity itself (e.g., from the temporal pattern of soil or forest sequestration compared to biomass substitution), and rates of adoption. Timing thus needs to be considered when judging the effectiveness of a mitigation action. Cherubini et al. (2012) modelled the impact of timing of benefits for three different wood applications (fuel, non-structural panels, and housing construction materials) and showed that the options provide mitigation over different timeframes, and thus have different impacts on CO₂ concentrations and radiative forcing. The temporal pattern of emissions and removals is especially important in mitigating emissions of short-lived gases through carbon sequestration (Lauder et al., 2013).

Box 11.5 | Bioenergy

Bioenergy deployment offers significant potential for climate change mitigation, but also carries considerable risks. The SRREN (IPCC, 2011) suggested potential bioenergy deployment levels to be between 100–300 EJ. This assessment agrees on a technical bioenergy potential of around 100 EJ, and possibly 300 EJ and higher. Integrated models project between 15–245 EJ/yr deployment in 2050, excluding traditional bioenergy. Achieving high deployment levels would require, amongst others, extensive use of agricultural residues and second-generation biofuels to mitigate adverse impacts on land use and food production, and the co-processing of biomass with coal or natural gas with carbon dioxide capture and storage (CCS) to produce low net GHG-emitting transportation fuels and/or electricity. Integration of crucial sectoral research (albedo effects, evaporation, counterfactual land carbon sink assumptions) into transformation pathways research, and exploration of risks of imperfect policy settings (for example, in absence of a global CO₂ price on land carbon) is subject of further research (Sections 11.9, 11.13.2, 11.13.4). Small-scale bioenergy systems aimed at meeting rural energy needs synergistically provide mitigation and energy access benefits. Decentralized deployment of biomass for energy, in combination with improved cookstoves, biogas, and small-scale biopower, could improve livelihoods and health of around 2.6 billion people. Both mitigation potential and sustainability hinges crucially on the protection of land carbon (high-density carbon ecosystems), careful fertilizer application, interaction with food markets, and good land and water management. Sustainability and livelihood concerns might constrain beneficial deployment of dedicated biomass plantations to lower values (Sections 11.13.3, 11.13.5, 11.13.7).

Lifecycle assessments for bioenergy options demonstrate a plethora of pathways, site-specific conditions and technologies that produce a wide range of climate-relevant effects. Specifically, LUC emissions, N₂O emissions from soil and fertilizers, co-products, process design and process fuel use, end-use technology, and reference system can all influence the total attributional lifecycle emissions of bioenergy use. The large variance for specific pathways points to the importance of management decisions in reducing the lifecycle emissions of bioenergy use. The total marginal global warming impact of bioenergy can only be evaluated

in a comprehensive setting that also addresses equilibrium effects, e.g., indirect land-use change (iLUC) emissions, actual fossil fuel substitution, and other effects. Structural uncertainty in modelling decisions renders such evaluation exercises uncertain. Available data suggest a differentiation between options that offer low lifecycle emissions under good land-use management (e.g., sugarcane, Miscanthus, and fast-growing tree species) and those that are unlikely to contribute to climate change mitigation (e.g., corn and soybean), pending new insights from more comprehensive consequential analyses (Sections 8.7, 11.13.4).

Coupling bioenergy and CCS (BECCS) has attracted particular attention since AR4 because it offers the prospect of negative emissions. Until 2050, the economic potential is estimated to be between 2–10 GtCO₂ per year. Some climate stabilization scenarios see considerable higher deployment towards the end of the century, even in some 580–650 ppm scenarios, operating under different time scales, socioeconomic assumptions, technology portfolios, CO₂ prices, and interpreting BECCS as part of an overall mitigation framework. Technological challenges and potential risks of BECCS include those associated with the provision of the biomass feedstock as well as with the capture, transport and long-term underground storage of CO₂. BECCS faces large challenges in financing and currently no such plants have been built and tested at scale (Sections 7.5.5, 7.9, 11.13.3).

Land demand and livelihoods are often affected by bioenergy deployment. Land demand for bioenergy depends on (1) the share of bioenergy derived from wastes and residues; (2) the extent to which bioenergy production can be integrated with food and fibre production, and conservation to minimize land-use competition; (3) the extent to which bioenergy can be grown on areas with little current production; and (4) the quantity of dedicated energy crops and their yields. Considerations of tradeoffs with water, land, and biodiversity are crucial to avoid adverse effects. The total impact on livelihood and distributional consequences depends on global market factors, impacting income and income-related food security, and site-specific factors such as land tenure and social dimensions. The often site-specific effects of bioenergy deployment on livelihoods have not yet been comprehensively evaluated (Section 11.13.7).

Additionality: Another consideration for gauging the effectiveness of mitigation is determining whether the activity would have occurred anyway, with this encompassed in the concept of 'additionality' (see Glossary).

Impacts of climate change: An area of emerging activity is predicting the likely impacts of climate change on mitigation potential, both in terms of impacts on existing carbon stocks, but also on the rates of carbon sequestration. This is discussed further in Section 11.5.

11.4 Infrastructure and systemic perspectives

Only supply-side mitigation options are considered in Section 11.3. In this section, we consider infrastructure and systemic perspectives, which include potential demand-side mitigation options in the AFOLU sector. Since infrastructure is a minor issue in AFOLU compared to energy end-use sectors, this section focusses on systemic perspectives.

11.4.1 Land: a complex, integrated system

Mitigation in the AFOLU sector is embedded in the complex interactions between socioeconomic and natural factors simultaneously affecting land systems (Turner et al., 2007). Land is used for a variety of purposes, including housing and infrastructure (Chapter 12), production of goods and services through agriculture, aquaculture and forestry, and absorption or deposition of wastes and emissions (Dunlap and Catton, Jr., 2002). Agriculture and forestry are important for rural livelihoods and employment (Coelho et al., 2012), while aquaculture and fisheries can be regionally important (FAO, 2012). More than half of the planet's total land area




Figure 11.9 | Global land use and biomass flows arising from human economic activity in 2000 from the cradle to the grave. Values in Gt dry matter biomass/yr. Figure source: (Smith et al., 2013b). If a source reported biomass flows in energy units, the numbers were converted to dry matter assuming a gross energy value of 18.5 MJ/kg. The difference between inputs and outputs in the consumption compartment is assumed to be released to the atmosphere (respiration, combustion); small differences may result from rounding. Note that data sources a) area: (Erb et al., 2007; Schneider et al., 2009; FAO, 2010) ; b) biomass flows: (Wirsenius, 2003; Sims et al., 2006; Krausmann et al., 2008; FAOSTAT, 2012; Kummu et al., 2012) are incomplete; more research is needed to close data gaps between different statistical sources such as agricultural, forestry, and energy statistics (Section 11.11). 'Unused forests' are pristine forests not harvested or otherwise used.

(134 Mkm²) is used for urban and infrastructure land, agriculture, and forestry. Less than one quarter shows relatively minor signs of direct human use (Erb et al., 2007; Ellis et al., 2010; Figure 11.9). Some of the latter areas are inhabited by indigenous populations, which depend on the land for the supply of vitally important resources (Read et al., 2010).

Land-use change is a pervasive driver of global environmental change (Foley et al., 2005, 2011). From 1950 to 2005, farmland (cropland plus pasture) increased from 28 to 38% of the global land area excluding ice sheets and inland waters (Hurt et al., 2011). The growth of farmland area (+33%) was lower than that of population, food production, and gross domestic product (GDP) due to increases in yields and biomass conversion efficiency (Krausmann et al., 2012). In the year 2000, almost one quarter of the global terrestrial net primary production (one third of the above-ground part) was 'appropriated' by humans. This means that it was either lost because the net primary productivity (the biomass production of green plants, net primary production, NPP) of agro-ecosystems or urban areas was lower than that of the vegetation they replaced or it was harvested for human purposes, destroyed during harvest or burned in human-induced fires (Imhoff et al., 2004; Haberl et al., 2007). The fraction of terrestrial NPP appropriated by humans doubled in the last century (Krausmann et al., 2013), exemplifying the increasing human domination of terrestrial ecosystems (Ellis et al., 2010). Growth trajectories of the use of food, energy, and other land-based resources, as well as patterns of urbanization and infrastructure development are influenced by increasing population and GDP, as well as the on-going agrarian-industrial transition (Haberl et al., 2011b; Kastner et al., 2012).

Growing resource use and land demand for biodiversity conservation and carbon sequestration (Soares-Filho et al., 2010), result in increasing competition for land (Harvey and Pilgrim, 2011; Section 11.4.2). Influencing ongoing transitions in resource use is a major challenge (WBGU, 2011; Fischer-Kowalski, 2011). Changes in cities, e.g., in terms of infrastructure, governance, and demand, can play a major role in this respect (Seto et al., 2012b; Seitzinger et al., 2012; Chapter 12).

Many mitigation activities in the AFOLU sector affect land use or land cover and, therefore, have socioeconomic as well as ecological consequences, e.g., on food security, livelihoods, ecosystem services or emissions (Sections 11.1; 11.4.5; 11.7). Feedbacks involved in implementing mitigation in AFOLU may influence different, sometimes conflicting, social, institutional, economic, and environmental goals (Madlener et al., 2006). Climate change mitigation in the AFOLU sector faces a complex set of interrelated challenges (Sections 11.4.5; 11.7):

- Full GHG impacts, including those from feedbacks (e.g., iLUC) or leakage, are often difficult to determine (Searchinger et al., 2008).
- Feedbacks between GHG reduction and other important objectives such as provision of livelihoods and sufficient food or the maintenance of ecosystem services and biodiversity are not completely understood.
- Maximizing synergies and minimizing negative effects involves multi-dimensional optimization problems involving various social,

economic, and ecological criteria or conflicts of interest between different social groups (Martinez-Alier, 2002).

- Changes in land use and ecosystems are scale-dependent and may proceed at different speeds, or perhaps even move in different directions, at different scales.

11.4.2 Mitigation in AFOLU—feedbacks with land-use competition

Driven by economic and population growth, increased demand for food and bioenergy as well as land demand for conservation and urbanization (e.g., above-ground biomass carbon losses associated with land-clearing from new urban areas in the pan-tropics are estimated to be 5% of the tropical deforestation and land-use change emissions, (Seto et al., 2012a; Section 12.2), competition for land is expected to intensify (Smith et al., 2010; Woods et al., 2010). Maximization of one output or service (e.g., crops) often excludes, or at least negatively affects, others (e.g., conservation; (Phalan et al., 2011). Mitigation in the AFOLU sector may affect land-use competition. Reduced demand for AFOLU products generally decreases inputs (fertilizer, energy, machinery) and land demand. The ecological feedbacks of demand-side options are mostly beneficial since they reduce competition for land and water (Smith et al., 2013b).

Some supply-side options, though not all, may intensify competition for land and other resources. Based on Figure 11.9 one may distinguish three cases:

- **Optimization of biomass-flow cascades;** that is, increased use of residues and by-products, recycling of biogenic materials and energetic use of wastes (WBGU, 2009). Such options increase resource use efficiency and may reduce competition, but there may also be tradeoffs. For example, using crop residues for bioenergy or roughage supply may leave less C and nutrients on cropland, reduce soil quality and C storage in soils, and increase the risk of losses of carbon through soil erosion. Residues are also often used as forage, particularly in the tropics. Forest residues are currently also used for other purposes, e.g., chipboard manufacture, pulp and paper production (González-Estrada et al., 2008; Blanco-Canqui and Lal, 2009; Muller, 2009; Ceschia et al., 2010).
- **Increases in yields** of cropland (Burney et al., 2010; Foley et al., 2011; Tilman et al., 2011; Mueller et al., 2012; Lobell et al., 2013), grazing land or forestry and improved livestock feeding efficiency (Steinfeld et al., 2010; Thornton and Herrero, 2010) can reduce land competition if yield increases relative to any additional inputs and the emission intensity (i.e., GHG emissions per unit of product) decreases. This may result in tradeoffs with other ecological, social, and economic costs (IAASTD, 2009) although these can to some extent be mitigated if intensification is sustainable (Tilman et al., 2011). Another caveat is that increases in yields may result in rebound effects that increase consumption (Lambin and Meyfroidt, 2011; Erb, 2012) or provide incentives to farm more land (Matson

and Vitousek, 2006), and hence may fail to spare land (Section 11.10).

- **Land-demanding options** reduce GHG emissions by harnessing the potential of the land for either C sequestration or growing energy crops (including food crops used as feedstocks for bioenergy production). These options result in competition for land (and sometimes other resources such as water) that may have substantial social, economic, and ecological effects (positive or negative; UNEP, 2009; WBGU, 2009; Chum et al., 2011; Coelho et al., 2012). Such options may increase pressures on ecosystems (e.g., forests) and GHG emissions related to direct and indirect LUC, contribute to price increases of agricultural products, or negatively affect livelihoods of rural populations. These possible impacts need to be balanced against possible positive effects such as GHG reduction, improved water quality (Townsend et al., 2012), restoration of degraded land (Harper et al., 2007), biodiversity protection (Swingland et al., 2002), and job creation (Chum et al., 2011; Coelho et al., 2012).

Therefore, an integrated energy/agriculture/land-use approach for mitigation in AFOLU can help to optimize synergies and mitigate negative effects (Popp et al., 2011; Smith, 2012; Creutzig et al., 2012a; Smith et al., 2013b).

11.4.3 Demand-side options for reducing GHG emissions from AFOLU

Some changes in demand for food and fibre can reduce GHG emissions in the production chain (Table 11.3) through (i) a switch to the consumption of products with higher GHG emissions in the process chain to products with lower GHG emissions and (ii) by making land available for other GHG reduction activities e.g., afforestation or bioenergy (Section 11.4.4). Food demand change is a sensitive issue due to the prevalence of hunger, malnutrition, and the lack of food security in many regions (Godfray et al., 2010). Sufficient production of, and equitable access to, food are both critical for food security (Misselhorn et al., 2012). GHG emissions may be reduced through changes in food demand without

jeopardizing health and well-being by (1) reducing losses and wastes of food in the supply chain as well as during final consumption; (2) changing diets towards less GHG-intensive food, e.g., substitution of animal products with plant-based food, while quantitatively and qualitatively maintaining adequate protein content, in regions with high animal product consumption; and (3) reduction of overconsumption in regions where this is prevalent. Substituting plant-based diets for animal-based diets is complex since, in many circumstances, livestock can be fed on plants not suitable for human consumption or growing on land with high soil carbon stocks not suitable for cropping; hence, food production by grazing animals contributes to food security in many regions of the world (Wirsenius, 2003; Gill et al., 2010).

Reductions of losses in the food supply chain—Globally, rough estimates suggest that ~30–40% of all food produced is lost in the supply chain from harvest to consumption (Godfray et al., 2010). Energy embodied in wasted food is estimated at ~36 EJ/yr (FAO, 2011). In developing countries, up to 40% is lost on farm or during distribution due to poor storage, distribution, and conservation technologies and procedures. In developed countries, losses on farm or during distribution are smaller, but the same amount is lost or wasted in service sectors and at the consumer level (Foley et al., 2005; Parfitt et al., 2010; Godfray et al., 2010; Gustavsson et al., 2011; Hodges et al., 2011). However, uncertainties related to losses in the food supply chain are large and more research is needed.

Not all losses are (potentially) avoidable because losses in households also include parts of products normally not deemed edible (e.g., peels of some fruits and vegetables). According to Parfitt et al. (2010), in the UK, 18% of the food waste is unavoidable, 18% is potentially avoidable, and 64% is avoidable. Data for Austria, Netherlands, Turkey, the United Kingdom, and the United States, derived with a variety of methods, show that food wastes at the household level in industrialized countries are 150–300 kg per household per year (Parfitt et al., 2010). According to a top-down mass-flow modelling study based on FAO commodity balances completely covering the food supply chain, but excluding non-edible fractions, food loss values range from 120–170 kg/cap/yr in Sub-Saharan Africa to 280–300 kg/cap/yr in Europe and North America.

Table 11.3 | Overview of demand-side mitigation options in the AFOLU sector.

Measure	Description	References
Reduced losses in the food supply chain	Reduced losses in the food supply chain and in final consumption reduces energy use and GHG emissions from agriculture, transport, storage and distribution, and reduce land demand.	(Godfray et al., 2010; Gustavsson et al., 2011), see text.
Changes in human diets towards less emission-intensive products	Where appropriate, reduced consumption of food items with high GHG emissions per unit of product, to those with low GHG products can reduce GHG emissions. Such demand changes can reduce energy inputs in the supply chain and reduces land demand.	(Stehfest et al., 2009; FAO, 2011), see text
Demand-side options related to wood and forestry	Wood harvest in forests releases GHG and at least temporarily reduces forest C stocks. Conservation of wood (products) through more efficient use or replacement with recycled materials and replacing wood from illegal logging or destructive harvest with wood from certified sustainable forestry (Section 11.10) can save GHG emissions. Substitution of wood for non-renewable resources can reduce GHG emissions, e.g., when wood is substituted for emission-intensive materials such as aluminium, steel, or concrete in buildings. Integrated optimization of C stocks in forests and in long-lived products, as well as the use of by-products and wastes for energy, can deliver the highest GHG benefits.	(Gustavsson et al., 2006; Werner et al., 2010; Ingerson, 2011), see text.

Losses ranging from 20% in Sub-Saharan Africa to more than 30% in the industrialized countries were calculated (Gustavsson et al., 2011).

A range of options exist to reduce wastes and losses in the supply chain: investments into harvesting, processing and storage technologies in the developing countries, awareness raising, taxation and other incentives to reduce retail and consumer-related losses primarily in the developed countries. Different options can help to reduce losses (i.e., increase efficiency) in the supply chain and at the household level. Substantial GHG savings could be realized by saving one quarter of the wasted food according to (Gustavsson et al., 2011); see Table 11.4.

Changes in human diets—Land use and GHG effects of changing diets require widespread behavioural changes to be effective; i.e., a strong deviation from current trajectories (increasing demand for food, in particular for animal products). Cultural, socioeconomic and behavioural aspects of implementation are discussed in Sections 11.4.5 and 11.7.

Studies based on Lifecycle Assessment (LCA) methods show substantially lower GHG emissions for most plant-based food than for animal products (Carlsson-Kanyama and González, 2009; Pathak et al., 2010; Bellarby et al., 2012; Berners-Lee et al., 2012), although there

are exceptions, e.g., vegetables grown in heated greenhouses or transported by airfreight (Carlsson-Kanyama and González, 2009). A comparison of three meals served in Sweden with similar energy and protein content based on (1) soy, wheat, carrots, and apples, (2) pork, potatoes, green beans, and oranges, and (3) beef, rice, cooked frozen vegetables, and tropical fruits revealed GHG emissions of 0.42 kgCO₂eq for the first option, 1.3 kgCO₂eq for the second, and 4.7 kgCO₂eq for the third, i.e., a factor of > 10 difference (Carlsson-Kanyama and González, 2009). Most LCA studies quoted here use attributional LCA; differences to results from consequential LCA (see Annex II) are generally not large enough to reverse the picture (Thomassen et al., 2008). The GHG benefits of plant-based food over animal products hold when compared per unit of protein (González et al., 2011). In addition to plant-based foods having lower emissions than animal-based ones, GHG emissions of livestock products also vary considerably; emissions per unit of protein are highest for beef and lower for pork, chicken meat, eggs and dairy products (de Vries and de Boer, 2010) due to their feed and land-use intensities. Figure 11.10 presents a comparison between milk and beef for different production systems and regions of the world (Herrero et al., 2013). Beef production can use up to five times more biomass for producing 1 kg of animal protein than dairy. Emissions intensities for the same livestock product also




Figure 11.10 | Biomass use efficiencies for the production of edible protein from (top) beef and (bottom) milk for different production systems and regions of the world (Herrero et al., 2013).

Table 11.4 | Changes in global land use and related GHG reduction potentials in 2050 assuming the implementation of options to increase C sequestration on farmland, and use of spared land for either biomass production for energy or afforestation. Afforestation and biomass for bioenergy are both assumed to be implemented only on spare land and are mutually exclusive (Smith et al., 2013b).

Cases	Food crop area	Livestock grazing area	C sink on farmland ¹	Afforestation of spare land ^{2,3}	Biomass for bioenergy on spare land ^{2,4}	Total mitigation potential	Difference in mitigation from reference case
	[Gha]		GtCO ₂ eq/yr				
Reference	1.60	4.07	3.5	6.1	1.2–9.4	4.6–12.9	0
Diet change	1.38	3.87	3.2	11.0	2.1–17.0	5.3–20.2	0.7–7.3
Yield growth	1.49	4.06	3.4	7.3	1.4–11.4	4.8–14.8	0.2–1.9
Feeding efficiency	1.53	4.04	3.4	7.2	1.4–11.1	4.8–14.5	0.2–1.6
Waste reduction	1.50	3.82	3.3	10.1	1.9–15.6	5.2–18.9	0.6–6.0
Combined	1.21	3.58	2.9	16.5	3.2–25.6	6.1–28.5	1.5–15.6

Notes:

- ¹ Potential for C sequestration on cropland for food production and livestock grazing land with improved soil C management. The potential C sequestration rate was derived from Smith et al., (2008).
- ² Spare land is cropland or grazing land not required for food production, assuming increased but still sustainable stocking densities of livestock based on Haberl et al., (2011); Erb et al., (2012).
- ³ Assuming 11.8 (tCO₂eq/ha)/yr (Smith et al., 2000).
- ⁴ Assumptions were as follows. High bioenergy value: short-rotation coppice or energy grass directly replaces fossil fuels, energy return on investment 1:30, dry-matter biomass yield 190 GJ/ha/yr (WBGU, 2009). Low bioenergy value: ethanol from maize replaces gasoline and reduces GHG by 45%, energy yield 75 GJ/ha/yr (Chum et al., 2011). Some energy crops may, under certain conditions, sequester C in addition to delivering bioenergy; the effect is context-specific and was not included. Whether bioenergy or afforestation is a better option to use spare land for mitigation needs to be decided on a case-by-case basis.

vary largely between different regions of the world due to differences in agro-ecology, diet quality, and intensity of production (Herrero et al., 2013). In overall terms, Europe and North America have lower emissions intensities per kg of protein than Africa, Asia, and Latin America. This shows that the highest potential for improving emissions intensities lies in developing countries, if intensification strategies can be matched to local resources and contexts.

Studies based on integrated modelling show that changes in diets strongly affect future GHG emissions from food production (Stehfest et al., 2009; Popp et al., 2010; Davidson, 2012). Popp et al. (2010) estimated that agricultural non-CO₂ emissions (CH₄ and N₂O) would triple by 2055 to 15.3 GtCO₂eq/yr if current dietary trends and population growth were to continue. Technical mitigation options on the supply side, such as improved cropland or livestock management, alone could reduce that value to 9.8 GtCO₂eq/yr, whereas emissions were reduced to 4.3 GtCO₂eq/yr in a 'decreased livestock product' scenario and to 2.5 GtCO₂eq/yr if both technical mitigation and dietary change were assumed. Hence, the potential to reduce GHG emissions through changes in consumption was found to be substantially higher than that of technical mitigation measures. Stehfest et al., (2009) evaluated effects of dietary changes on CO₂ (including C sources/sinks of ecosystems), CH₄, and N₂O emissions. In a 'business-as-usual' scenario largely based on FAO (2006), total GHG emissions were projected to reach 11.9 GtCO₂eq/yr in 2050. The following changes were evaluated: no ruminant meat, no meat, and a diet without any animal products. Changed diets resulted in GHG emission savings of 34–64% compared to the 'business-as-usual' scenario; a switch to a 'healthy diet' recommended by the Harvard Medical School would save 4.3 GtCO₂eq/yr

(–36%). Adoption of the 'healthy diet' (which includes a meat, fish and egg consumption of 90 g/cap/day) would reduce global GHG abatement costs to reach a 450 ppm CO₂eq concentration target by ~50% compared to the reference case (Stehfest et al., 2009). The analysis assumed nutritionally sufficient diets; reduced supply of animal protein was compensated by plant products (soy, pulses, etc.). Considerable cultural and social barriers against a widespread adoption of dietary changes to low-GHG food may be expected (Davidson, 2012; Smith et al., 2013, 11.4.5).

A limitation of food-related LCA studies is that they have so far seldom considered the emissions resulting from LUC induced by changing patterns of food production (Bellarby et al., 2012). A recent study (Schmidinger and Stehfest, 2012) found that cropland and pastures required for the production of beef, lamb, calf, pork, chicken, and milk could annually sequester an amount of carbon equivalent to 30–470% of the GHG emissions usually considered in LCA of food products if the land were to be reforested. Land-related GHG costs differ greatly between products and depend on the time horizon (30–100 yr) assumed (Schmidinger and Stehfest, 2012). If cattle production contributes to tropical deforestation (Zaks et al., 2009; Bustamante et al., 2012; Houghton et al., 2012), land-use related GHG emissions are particularly high (Cederberg et al., 2011). These findings underline the importance of diets for GHG emissions in the food supply chain (Garnett, 2011; Bellarby et al., 2012). A potential co-benefit is a reduction in diet-related health risks in regions where overconsumption of animal products is prevalent (McMichael et al., 2007).

Demand-side options related to wood and forestry—A comprehensive global, long-term dataset on carbon stocks in long-lived wood

products in use (excluding landfills) shows an increase from approximately 2.2 GtC in 1900 to 6.9 GtC in 2008 (Lauk et al., 2012). Per capita, carbon stored in wood products amounted to ~1.4 tC/cap in 1900 and ~1.0 tC/cap in 2008. The net yearly accumulation of long-lived wood products in use varied between 35 and 91 MtC/yr in the period 1960–2008 (Lauk et al., 2012). The yearly accumulation of C in products and landfills was ~200 MtC/yr in the period 1990–2008 (Pan et al., 2011). If more long-lived wood products were used, C sequestration and mitigation could be enhanced.

Increased wood use does not reduce GHG emissions under all circumstances because wood harvest reduces the amount of carbon stored in the forest, at least temporarily, and increases in wood harvest levels may result in reduced long-term carbon storage in forests (Ingerson, 2011; Böttcher et al., 2012; Holtsmark, 2012; Lamers and Junginger, 2013). Reducing wood consumption, e.g., through paper recycling, can reduce GHG emissions (Acuff and Kaffine, 2013), as may the use of wood from sustainable forestry in place of emission-intensive materials such as concrete, steel, or aluminium. Recent studies suggest that, where technically possible, substitution of wood from sustainably managed forests for non-wood materials in the construction sector (concrete, steel, etc.) in single-family homes, apartment houses, and industrial buildings, reduces GHG emissions in most cases (Werner et al., 2010; Sathre and O'Connor, 2010; Ximenes and Grant, 2013). Most of the emission reduction results from reduced production emissions, whereas the role of carbon sequestration in products is relatively small (Sathre and O'Connor, 2010). Werner et al. (2010) show that GHG benefits are highest when wood is primarily used for long-lived products, the lifetime of products is maximized, and energy use of woody biomass is focused on by-products, wood wastes, and end-of-lifecycle use of long-lived wood products.

11.4.4 Feedbacks of changes in land demand

Mitigation options in the AFOLU sector, including options such as biomass production for energy, are highly interdependent due to their direct and indirect impacts on land demand. Indirect interrelationships, mediated *via* area demand for food production, which in turn affects the area available for other purposes, are difficult to quantify and require systemic approaches. Table 11.4 (Smith et al., 2013b) shows the magnitude of possible feedbacks in the land system in 2050. It first reports the effect of single mitigation options compared to a reference case, and then the combined effect of all options. The reference case is similar to the (FAO, 2006a) projections for 2050 and assumes a continuation of on-going trends towards richer diets, considerably higher cropland yields (+54%) and moderately increased cropland areas (+9%). The diet change case assumes a global contract-and-converge scenario towards a nutritionally sufficient low animal product diet (8% of food calories from animal products). The yield growth case assumes that yields in 2050 are 9% higher than those in the reference case, according to the 'Global Orchestration' scenario in (MEA, 2005). The feeding efficiency case assumes on aver-

age 17% higher livestock feeding efficiencies than the reference case. The waste reduction case assumes a reduction of the losses in the food supply chain by 25% (Section 11.4.3). The combination of all options results in a substantial reduction of cropland and grazing areas (Smith et al., 2013b), even though the individual options cannot simply be added up due to the interactions between the individual compartments.

Table 11.4 shows that demand-side options save GHG by freeing up land for bioenergy or afforestation and related carbon sequestration. The effect is strong and non-linear, and more than cancels out reduced C sequestration potentials on farmland. Demand-side potentials are substantial compared to supply-side mitigation potentials (Section 11.3), but implementation may be difficult (Sections 11.7; 11.8). Estimates of GHG savings from bioenergy are subject to large uncertainties related to the assumptions regarding power plants, utilization pathway, energy crop yields, and effectiveness of sustainability criteria (Sections 11.4.5; 11.7; 11.13).

The systemic effects of land-demanding mitigation options such as bioenergy or afforestation depend not only on their own area demand, but also on land demand for food and fibre supply (Chum et al., 2011; Coelho et al., 2012; Erb et al., 2012b). In 2007, energy crops for transport fuels covered about 26.6 Mha or 1.7% of global cropland (UNEP, 2009). Assumptions on energy crop yields (Section 11.13) are the main reason for the large differences in estimates of future area demand of energy crops in the next decades, which vary from < 100 Mha to > 1000 Mha, i.e., 7–70% of current cropland (Sims et al., 2006; Smeets et al., 2007; Pacca and Moreira, 2011; Coelho et al., 2012). Increased pressure on land systems may also emerge when afforestation claims land, or forest conservation restricts farmland expansion (Murtaugh and Schlap, 2009; Popp et al., 2011).

Land-demanding mitigation options may result in feedbacks such as GHG emissions from land expansion or agricultural intensification, higher yields of food crops, higher prices of agricultural products, reduced food consumption, displacement of food production to other regions and consequent land clearing, as well as impacts on biodiversity and non-provisioning ecosystem services (Plevin et al., 2010; Popp et al., 2012).

Restrictions to agricultural expansion due to forest conservation, increased energy crop area, afforestation and reforestation may increase costs of agricultural production and food prices. In a modelling study, conserving C-rich natural vegetation such as tropical forests was found to increase food prices by a factor of 1.75 until 2100, due to restrictions of cropland expansion, even if no growth of energy crop area was assumed (Wise et al., 2009). Food price indices (weighted average of crop and livestock products) are estimated to increase until 2100 by 82% in Africa, 73% in Latin America, and 52% in Pacific Asia if large-scale bioenergy deployment is combined with strict forest conservation, compared to a reference scenario without forest conservation and bioenergy (Popp et al., 2011). Further trade liberalization can

lead to lower costs of food, but also increases the pressure on land, especially on tropical forests (Schmitz et al., 2011).

Increased land demand for GHG mitigation can be partially compensated by higher agricultural yield per unit area (Popp et al., 2011). While yield increases can lead to improvements in output from less land, generate better economic returns for farmers, help to reduce competition for land, and alleviate environmental pressures (Burney et al., 2010; Smith et al., 2010), agricultural intensification if poorly implemented incurs economic costs (Lotze-Campen et al., 2010) and may also create social and environmental problems such as nutrient leaching, soil degradation, pesticide pollution, impact on animal welfare, and many more (IAASTD, 2009). Maintaining yield growth while reducing negative environmental and social effects of agricultural intensification is, therefore, a central challenge, requiring sustainable management of natural resources as well as the increase of resource efficiency (DeFries and Rosenzweig, 2010), two components of sustainable intensification (Garnett et al., 2013).

Additional land demand may put pressures on biodiversity, as LUC is one of the most important drivers of biodiversity loss (Sala et al., 2000). Improperly managed large-scale agriculture (or bioenergy) may negatively affect biodiversity (Groom et al., 2008), which is a key prerequisite for the resilience of ecosystems, i.e., their ability to adapt to changes such as climate change, and to continue to deliver ecosystem services in the future (Díaz et al., 2006; Landis et al., 2008). However, implementing appropriate management, such as establishing bioenergy crops or plantations for carbon sequestration in already degraded ecosystems areas represents an opportunity where bioenergy can be used to achieve positive environmental outcomes (e.g., Hill et al., 2006; Semere and Slater, 2007; Campbell et al., 2008; Nijssen et al., 2012). Because climate change is also an important driver of biodiversity loss (Sala et al., 2000), bioenergy for climate change mitigation may also be beneficial for biodiversity if it is planned with biodiversity conservation in mind (Heller and Zavaleta, 2009; Dawson et al., 2011; Section 11.13).

Tradeoffs related to land demand may be reduced through multifunctional land use, i.e., the optimization of land to generate more than one product or service such as food, animal feed, energy or materials, soil protection, wastewater treatment, recreation, or nature protection (de Groot, 2006; DeFries and Rosenzweig, 2010; Section 11.7). This also applies to the potential use of ponds and other small water bodies for raising fish fed with agricultural waste (Pullin et al., 2007).

11.4.5 Sustainable development and behavioural aspects

The assessment of impacts of AFOLU mitigation options on sustainable development requires an understanding of a complex multilevel system where social actors make land-use decisions aimed at various development goals, one of them being climate change mitigation. Depending on the specific objectives, the beneficiaries of a particular land-use

choice may differ. Thus tradeoffs between global, national, and local concerns and various stakeholders need to be considered (see also Section 4.3.7 and WGII Chapter 20). The development context provides opportunities or barriers for AFOLU (May et al., 2005; Madlener et al., 2006; Smith and Trines, 2006; Smith et al., 2007; Angelsen, 2008; Howden et al., 2008; Corbera and Brown, 2008; Cotula et al., 2009; Cattaneo et al., 2010; Junginger et al., 2011; Section 11.8 and Figure 11.11).

Further, AFOLU measures have additional effects on development, beyond improving the GHG balance (Foley et al., 2005; Alig et al., 2010; Calfapietra et al., 2010; Busch et al., 2011; Smith et al., 2013b; Branca et al., 2013; Albers and Robinson, 2013). These effects can be positive (co-benefits) or negative (adverse side-effects) and do not necessarily overlap geographically, socially or in time (Section 11.7 and Figure 11.11). This creates the possibility of tradeoffs, because an AFOLU measure can bring co-benefits to one social group in one area (e.g., increasing income), while bringing adverse side-effects to others somewhere else (e.g., reducing food availability).

Table 11.5 summarizes the issues commonly considered when assessing the above-mentioned interactions at various levels between sustainable development and AFOLU.

Social complexity: Social actors in the AFOLU sector include individuals (farmers, forest users), social groups (communities, indigenous groups), private companies (e.g., concessionaires, food-producer multinationals), subnational authorities, and national states (see Table 11.6).




Figure 11.11 | Dynamic interactions between the development context and AFOLU.

Spatial scale refers on the one hand to the size of an intervention (e.g., in number of hectares) and on the other hand to the biophysical characterization of the specific land (e.g., soil type, water availability, slope). Social interactions tend to become more complex the bigger the area of an AFOLU intervention, on a social-biophysical continuum: family/farm—neighbourhood—community—village—city—province—country—region—globe. Impacts from AFOLU measures on sustainable development are different along this spatial-scale continuum (Table 11.6). The challenge is to provide landscape governance that responds to societal needs as well as biophysical capacity at different spatial scales (Görg, 2007; Moilanen and Arponen, 2011; van der Horst and Vermeylen, 2011).

Temporal scale: As the concept of sustainable development includes current and future generations, the impacts of AFOLU over time need to be considered (see Chapter 4). Positive and negative impacts of AFOLU measures can be realized at different times. For instance, while reducing deforestation has an immediate positive impact on reducing GHG emissions, reforestation will have a positive impact on C sequestration over time. Further, in some circumstances, there is the risk of reversing current emission reductions in the future (see Section 11.3.2 on non-permanence).

Behavioural aspects: Level of education, cultural values and tradition, as well as access to markets and technology, and the decision power of individuals and social groups, all influence the perception of potential impacts and opportunities from AFOLU measures, and consequently have a great impact on local land management decisions (see Chapters 2, 3, and 4; Guthinga, 2008; Durand and Lazos, 2008; Gilg, 2009; Bhuiyan et al., 2010; Primmer and Karppinen, 2010; Durand and Vázquez, 2011). When decisions are taken at a higher administrative level (e.g., international corporations, regional authorities or national states), other factors or values play an important role, including national and international development goals and priorities, policies and commitments, international markets or corporate image (see Chapters 3 and 4). Table 11.7 summarizes the emerging behavioural aspects regarding AFOLU mitigation measures.

Land-use policies (Section 11.10) have the challenge of balancing impacts considering these parameters: social complexity, spatial scale, temporal scale, and behavioural aspects. Vlek and Keren (1992) and Vlek (2004) indicate the following dilemmas relevant to land-management decisions: Who should take the risks, when (this generation or future generations) and where (specific place) co-benefits and potential adverse effects will take place, and how to mediate between individual vs. social benefits. Addressing these dilemmas is context-specific. Nevertheless, the fact that a wide range of social actors need to face these dilemmas explains, to a certain extent, disagreements about environmental decision making in general, and land-management decisions in particular (Villamor et al., 2011; Le et al., 2012; see Section 11.10).

11.5 Climate change feedback and interaction with adaptation (includes vulnerability)

When reviewing the inter-linkages between climate change mitigation and adaptation within the AFOLU sector the following issues need to be considered: (i) the impact of climate change on the mitigation potential of a particular activity (e.g., forestry and agricultural soils) over time, (ii) potential synergies/tradeoffs within a land-use sector between mitigation and adaptation objectives, and (iii) potential tradeoffs across sectors between mitigation and adaptation objectives.

Mitigation and adaptation in land-based ecosystems are closely inter-linked through a web of feedbacks, synergies, and tradeoffs (Section 11.8). The mitigation options themselves may be vulnerable to climatic change (Section 11.3.2) or there may be possible synergies or tradeoffs between mitigation and adaptation options within or across AFOLU sectors.

Table 11.5 | Issues related to AFOLU measures and sustainable development.

Dimensions	Issues
Social and human assets	Population growth and migration, level of education, human capacity, individual skills, indigenous and traditional knowledge, cultural values, equity and health, animal welfare, organizational capacity
Natural assets	Availability of natural resources (land, forest, water, agricultural land, minerals, fauna), GHG balance, ecosystem integrity, biodiversity conservation, ecosystem services, the productive capacity of ecosystems, ecosystem health and resilience
State of infrastructure and technology	Availability of infrastructure and technology and industrial capacity, technology development, appropriateness, acceptance
Economic factors	Credit capacity, employment creation, income, wealth distribution/distribution mechanisms, carbon finance, available capital/investments, market access
Institutional arrangements	Land tenure and land-use rights, participation and decision making mechanisms (e.g., through Free, Prior and Informed Consent), sectoral and cross-sectoral policies, investment in research, trade agreements and incentives, benefit sharing mechanisms, existence and forms of social organization

Based on Madlener et al. (2006), Sneddon et al. (2006), Pretty (2008), Corbera and Brown (2008), Macauley and Sedjo (2011), and de Boer et al. (2011).

Table 11.6 | Characterization of social actors in AFOLU.

Social actors	Characterization
Individuals (legal and illegal forest users, farmers)	Rather small-scale interventions, although some can be medium-scale Decisions taken rather at the local level
Social groups (communities, indigenous peoples)	Small to medium interventions Decisions taken at the local or regional levels
Sub-national authorities (provinces, states)	Medium to large interventions Decisions taken at the national or sub-national level, depending on the governance structure
State (national level)	Rather large interventions Decisions taken at the national level, often in line with international agreements
Corporate (at the national or multinational levels)	Rather large interventions. Decisions can be taken within a specific region/country, in another country, or at global level (e.g., for multinational companies). National and international markets play a key role in decision making

Table 11.7 | Emerging behavioural aspects relevant for AFOLU mitigation measures.

Change in	Emerging behavioural aspects in AFOLU
Consumption patterns	<p>Dietary change: Several changes in diet can potentially reduce GHG emissions, including reduction of food waste and reduction of or changes in meat consumption (especially in industrialized countries). On the other hand, increasing income and evolving lifestyles with increasing consumption of animal protein in developing countries are projected to increase food-related GHG emissions.</p> <p>The potential of reducing GHG emissions in the food sector needs to be understood in a wider and changing socio-cultural context that determines nutrition.</p> <p>Potential drivers of change: Health awareness and information, income increase, lifestyle</p> <p>References 1, 2,3, 4, 5</p>
Production patterns	<p>Large-scale land acquisition: The acquisition of (long-term rights) of large areas of farmland in lower-income countries, by transnational companies, agribusiness, investments funds or government agencies. There are various links between these acquisitions and GHG emissions in the AFOLU sector. On one hand because some acquisitions are aimed at producing energy crops (through non-food or 'flex-crops'), on the other because these can cause the displacement of peoples and activity, increasing GHG leakage.</p> <p>Impacts on livelihood, local users rights, local employment, economic activity, or on biodiversity conservation are of concern.</p> <p>Potential drivers of change: International markets and their mechanisms, national and international policies</p> <p>References 6, 7, 8</p>
Production and consumption patterns	<p>Switching to low-carbon products: Land managers are sensitive to market changes. The promotion of low-carbon products as a means for reducing GHG emissions can increase the land area dedicated to these products. Side-effects from this changes in land management (positive and negative), and acceptability of products and technologies at the production and consumption sides are context-related and cannot be generalized</p> <p>Potential drivers of change: International agreements and markets, accessibility to rural energy, changes in energy demand</p> <p>References 9, 10, 11</p>
Relation between producers and consumers	<p>Certification: Labelling, certification, or other information-based instruments have been developed for promoting behavioural changes towards more sustainable products (Section 11.10). Recently, the role of certification in reducing GHG while improving sustainability has been explored, especially for bioenergy (Section 11.13).</p> <p>Potential drivers of change: Consumer awareness, international agreements, cross-national sector policies and initiatives.</p> <p>References 11, 12, 13, 14</p>
Management priorities	<p>Increasing interest in conservation and sustainable (land) management: Changing management practices towards more sustainable ones as alternative for gaining both environmental and social co-benefits, including climate change mitigation, is gaining recognition. Concerns about specific management practices, accountability methods of co-benefits, and sharing mechanisms seem to be elements of concerns when promoting a more sustainable management of natural resources.</p> <p>Potential drivers of change: Policies and international agreements and their incentive mechanisms, schemes for payments for environmental services.</p> <p>References 15, 16, 17, 18, 19</p>

¹Stehfest et al. (2009); ²Roy et al. (2012); ³González et al. (2011); ⁴Popp et al. (2010); ⁵Schneider et al. (2011); ⁶Cotula (2012); ⁷Messerli et al. (2013); ⁸German et al. (2013); ⁹Muys et al. (2014); ¹⁰MacMillan Uribe et al. (2012); ¹¹Chakrabarti (2010); ¹²Karipidis et al. (2010); ¹³Auld et al. (2008); ¹⁴Diaz-Chavez (2011); ¹⁵Calegari et al. (2008); ¹⁶Deal et al. (2012); ¹⁷DeFries and Rosenzweig (2010); ¹⁸Hein and van der Meer (2012); ¹⁹Lippke et al. (2003).

The IPCC WGI presents feedbacks between climate change and the carbon cycle (WGI Chapter 6; Le Quéré et al., 2013), while WGII assesses the impacts of climate change on terrestrial ecosystems (WGII Chapter 4) and crop production systems (WGII Chapter 7), including vulnerability and adaptation. This section focuses particularly on the impacts of climate change on mitigation potential of land-use sectors and interactions that arise with adaptation, linking to the relevant chapters of WGI and WGII reports.

11.5.1 Feedbacks between AFOLU and climate change

AFOLU activities can either reduce or accelerate climate change by affecting biophysical processes (e.g., evapotranspiration, albedo) and change in GHG fluxes to and from the atmosphere (WGI). Whether a particular ecosystem is functioning as sink or source of GHG emission may change over time, depending on its vulnerability to climate change and other stressors and disturbances. Hence, mitigation options available today (Section 11.3) in the AFOLU sectors may no longer be available in the future.

There is *robust evidence* that human-induced land-use changes have led to an increased surface albedo (WGI Chapter 8; Myhre and Shindell, 2013). Changes in evapotranspiration and surface roughness may counteract the effect of changes in albedo. Land-use changes affect latent heat flux and influence the hydrological cycle. Biophysical climate feedbacks of forest ecosystems differ depending on regional climate regime and forest types. For example, a decrease in tropical forests has a positive climate forcing through a decrease in evaporative cooling (Bala et al., 2007; Bonan, 2008). An increase in coniferous-boreal forests compared to grass and snow provides a positive climate forcing through lowering albedo (Bala et al., 2007; Bonan, 2008; Swann et al., 2010). There is currently *low agreement* on the net biophysical effect of land-use changes on the global mean temperature (WGI Chapter 8; Myhre and Shindell, 2013). By contrast, the biogeochemical effects of LUC on radiative forcing through emissions of GHG is positive (WGI Chapter 8; Sections 11.2.2; 11.2.3).

11.5.2 Implications of climate change on terrestrial carbon pools and mitigation potential of forests

Projections of the global carbon cycle to 2100 using 'Coupled Model Intercomparison Project Phase 5 (CMIP5) Earth System Models' (WGI Chapter 6; Le Quéré et al., 2013) that represent a wider range of complex interactions between the carbon cycle and the physical climate system consistently estimate a positive feedback between climate and the carbon cycle, i.e., reduced natural sinks or increased natural CO₂ sources in response to future climate change. Implications of climate change on terrestrial carbon pools biomes and mitigation potential of forests.

Rising temperatures, drought, and fires may lead to forests becoming a weaker sink or a net carbon source before the end of the century (Sitch et al., 2008). Pervasive droughts, disturbances such as fire and insect outbreaks, exacerbated by climate extremes and climate change put the mitigation benefits of the forests at risk (Canadell and Raupach, 2008; Phillips et al., 2009; Herawati and Santoso, 2011). Forest disturbances and climate extremes have associated carbon balance implications (Millar et al., 2007; Kurz et al., 2008; Zhao and Running, 2010; Potter et al., 2011; Davidson, 2012; Reichstein et al., 2013). Allen et al. (2010) suggest that at least some of the world's forested ecosystems may already be responding to climate change.

Experimental studies and observations suggest that predicted changes in temperature, rainfall regimes, and hydrology may promote the die-back of tropical forests (e.g., Nepstad et al., 2007). The prolonged drought conditions in the Amazon region during 2005 contributed to a decline in above-ground biomass and triggered a release of 4.40 to 5.87 GtCO₂ (Phillips et al., 2009). Earlier model studies suggested Amazon die-back in the future (Cox et al., 2013; Huntingford et al., 2013). However, recent model estimates suggest that rainforests may be more resilient to climate change, projecting a moderate risk of tropical forest reduction in South America and even lower risk for African and Asian tropical forests (Gumpenberger et al., 2010; Cox et al., 2013; Huntingford et al., 2013).

Arcidiacono-Bársony et al., (2011) suggest that the mitigation benefits from deforestation reduction under REDD+ (Section 11.10.1) could be reversed due to increased fire events, and climate-induced feedbacks, while Gumpenberger et al., (2010) conclude that the protection of forests under the forest conservation (including REDD) programmes could increase carbon uptake in many tropical countries, mainly due to CO₂ fertilization effects, even under climate change conditions.

11.5.3 Implications of climate change on peatlands, grasslands, and croplands

Peatlands: Wetlands, peatlands, and permafrost soils contain higher carbon densities relative to mineral soils, and together they comprise extremely large stocks of carbon globally (Davidson and Janssens, 2006). Peatlands cover approximately 3% of the Earth's land area and are estimated to contain 350–550 Gt of carbon, roughly between 20 to 25% of the world's soil organic carbon stock (Gorham, 1991; Fenner et al., 2011). Peatlands can lose CO₂ through plant respiration and aerobic peat decomposition (Clair et al., 2002) and with the onset of climate change, may become a source of CO₂ (Koehler et al., 2010). Large carbon losses are likely from deep burning fires in boreal peatlands under future projections of climate warming and drying (Flannigan et al., 2009). A study by Fenner et al. (2011) suggests that climate change is expected to increase the frequency and severity of drought in many of the world's peatlands which, in turn, will release far more GHG emissions than thought previously. Climate change is projected to have a severe impact on the peatlands in northern regions where

most of the perennially frozen peatlands are found (Tarnocai, 2006). According to Schuur et al. (2008), the thawing permafrost and consequent microbial decomposition of previously frozen organic carbon, is one of the most significant potential feedbacks from terrestrial ecosystems to the atmosphere in a changing climate. Large areas of permafrost will experience thawing (WGI Chapter 12), but uncertainty over the magnitude of frozen carbon losses through CO₂ or CH₄ emissions to the atmosphere is large, ranging between 180 and 920 GtCO₂ by the end of the 21st century under the Representative Concentration Pathways (RCP) 8.5 scenario (WGI Chapter 6; Le Quéré et al., 2013).

Grasslands: Tree cover and biomass in savannah has increased over the past century (Angassa and Oba, 2008; Witt et al., 2009; Lunt et al., 2010; Rohde and Hoffman, 2012) leading to increased carbon storage per hectare (Hughes et al., 2006; Liao et al., 2006; Throop and Archer, 2008; Boutton et al., 2009), which has been attributed to land management, rising CO₂, climate variability, and climate change. Climate change and CO₂ may affect grazing systems by altering species composition; for example, warming will favour tropical (C4) species over temperate (C3) species but CO₂ increase would favour C3 grasses (Howden et al., 2008).

Croplands: Climate change impacts on agriculture will affect not only crop yields, but also soil organic carbon (SOC) levels in agricultural soils (Rosenzweig and Tubiello, 2007). Such impacts can be either positive or negative, depending on the particular effect considered, which highlights the uncertainty of the impacts. Elevated CO₂ concentrations alone are expected to have positive effects on soil carbon storage, because of increased above- and below-ground biomass production in agro-ecosystems. Similarly, the lengthening of the growing season under future climate will allow for increased carbon inputs into soils. Warmer temperatures could have negative impacts on SOC, by speeding decomposition and by reducing inputs by shortening crop lifecycles (Rosenzweig and Tubiello, 2007), but increased productivity could increase SOC stocks (Gottschalk et al., 2012).

11.5.4 Potential adaptation options to minimize the impact of climate change on carbon stocks in forests and agricultural soils

Forests: Forest ecosystems require a longer response time to adapt. The development and implementation of adaptation strategies is also lengthy (Leemans and Eickhout, 2004; Ravindranath, 2007). Some examples of the adaptation practices (Murthy et al., 2011) are as follows: anticipatory planting of species along latitude and altitude, assisted natural regeneration, mixed-species forestry, species mix adapted to different temperature tolerance regimes, fire protection and management practices, thinning, sanitation and other silvicultural practices, *in situ* and *ex situ* conservation of genetic diversity, drought and pest resistance in commercial tree species, adoption of sustainable forest management practices, increase in Protected Areas and

linking them wherever possible to promote migration of species, forests conservation and reduced forest fragmentation enabling species migration, and energy-efficient fuel-wood cooking devices to reduce pressure on forests.

Agricultural soils: On current agricultural land, mitigation and adaptation interaction can be mutually re-enforcing, particularly for improving resilience to increased climate variability under climate change (Rosenzweig and Tubiello, 2007). Many mitigation practices implemented locally for soil carbon sequestration will increase the ability of soils to hold soil moisture and to better withstand erosion and will enrich ecosystem biodiversity by establishing more diversified cropping systems, and may also help cropping systems to better withstand droughts and floods, both of which are projected to increase in frequency and severity under a future warmer climate (Rosenzweig and Tubiello, 2007).

11.5.5 Mitigation and adaptation synergies and tradeoffs

Mitigation choices taken in a particular land-use sector may further enhance or reduce resilience to climate variability and change within or across sectors, in light of the multiple, and often competing, pressures on land (Section 11.4), and shifting demographics and consumption patterns (e.g., O'Brien et al., 2004; Sperling et al., 2008; Hunsberger and Evans, 2012). Land-use choices driven by mitigation concerns (e.g., forest conservation, afforestation) may have consequences for adaptive responses and/or development objectives of other sectors (e.g., expansion of agricultural land). For example, reducing emissions from deforestation and degradation may also yield co-benefits for adaptation by maintaining biodiversity and other ecosystem goods and services, while plantations, if they reduce biological diversity may diminish adaptive capacity to climate change (e.g., Chum et al., 2011). Primary forests tend to be more resilient to climate change and other human-induced environmental changes than secondary forests and plantations (Thompson et al., 2009). The impact of plantations on the carbon balance is dependent on the land-use system they replace. While plantation forests are often monospecies stands, they may be more vulnerable to climatic change (see IPCC WGII Chapter 4). Smith and Olesen (2010) identified a number of synergies between options that deliver mitigation in agriculture while also enhancing resilience to future climate change, the most prominent of which was enhancement of soil carbon stocks.

Adaptation measures in return may help maintain the mitigation potential of land-use systems. For example, projects that prevent fires and restore degraded forest ecosystems also prevent release of GHGs and enhance carbon stocks (CBD and GiZ, 2011). Mitigation and adaptation benefits can also be achieved within broader-level objectives of AFOLU measures, which are linked to sustainable development considerations. Given the exposure of many livelihoods and communities to multiple stressors, recommendations from case stud-

ies suggest that climate risk-management strategies need to appreciate the full hazard risk envelope, as well as the compounding socio-economic stressors (O'Brien et al., 2004; Sperling et al., 2008). Within this broad context, the potential tradeoffs and synergies between mitigation, adaptation, and development strategies and measures need to be considered. Forest and biodiversity conservation, protected area formation, and mixed-species forestry-based afforestation are practices that can help to maintain or enhance carbon stocks, while also providing adaptation options to enhance resilience of forest ecosystems to climate change (Ravindranath, 2007). Use of organic soil amendments as a source of fertility could potentially increase soil carbon (Gattinger et al., 2012). Most categories of adaptation options for climate change have positive impacts on mitigation. In the agriculture sector, cropland adaptation options that also contribute to mitigation are 'soil management practices that reduce fertilizer use and increase crop diversification; promotion of legumes in crop rotations; increasing biodiversity, the availability of quality seeds and integrated crop/livestock systems; promotion of low energy production systems; improving the control of wildfires and avoiding burning of crop residues; and promoting efficient energy use by commercial agriculture and agro-industries' (FAO, 2008, 2009a). Agroforestry is an example of mitigation-adaptation synergy in the agriculture sector, since trees planted sequester carbon and tree products provide livelihood to communities, especially during drought years (Verchot et al., 2007).

11.6 Costs and potentials

This section deals with economic costs and potentials of climate change mitigation (emission reduction or sequestration of carbon) within the AFOLU sector. Economic mitigation potentials are distinguished from technical or market mitigation potentials (Smith, 2012). Technical mitigation potentials represent the full biophysical potential of a mitigation option, without accounting for economic or other constraints. These estimates account for constraints and factors such as land availability and suitability (Smith, 2012), but not any associated costs (at least explicitly). By comparison, economic potential refers to mitigation that could be realized at a given carbon price over a specific period, but does not take into consideration any socio-cultural (for example, lifestyle choices) or institutional (for example, political, policy, and informational) barriers to practice or technology adoption. Economic potentials are expected to be lower than the corresponding technical potentials. Also, policy incentives (e.g., a carbon price; see also Section 11.10) and competition for resources across various mitigation options, tend to affect the size of economic mitigation potentials in the AFOLU sector (McCarl and Schneider, 2001). Finally, market potential is the realized mitigation outcome under current or forecast market conditions encompassing biophysical, economic, socio-cultural, and institutional barriers, as well as policy incentives for, technological and/or practice adoption, specific to a sub-national, national or

supra-national market for carbon. Figure 11.12 (Smith, 2012) provides a schematic view of the three types of mitigation potentials.

Economic (as well as market) mitigation potentials tend to be context-specific and are likely to vary across spatial and temporal scales. Unless otherwise stated, in the rest of this section, economic potentials are expressed in million tonnes (Mt) of mitigation in carbon dioxide equivalent (CO₂eq) terms, that can arise from an individual mitigation option or from an AFOLU sub-sector at a given cost per tonne of CO₂eq. (USD/tCO₂eq) over a given period to 2030, which is 'additional' to the corresponding baseline or reference case levels.

Various supply-side mitigation options within the AFOLU sector are described in Section 11.3, and Section 11.4 considers a number of potential demand-side options. Estimates for costs and potentials are not always available for the individual options described. Also, aggregate estimates covering both the supply- and demand-side options for mitigation within the AFOLU sector are lacking, so this section mostly focuses on the supply-side options. Key uncertainties and sensitivities around mitigation costs and potentials in the AFOLU sector are (1) carbon price, (2) prevailing biophysical and climatic conditions, (3) existing management heterogeneity (or differences in the baselines), (4) management interdependencies (arising from competition or co-benefits across tradition production, environmental outcomes and mitigation strategies or competition/co-benefits across mitigation options), (5) the extent of leakage, (6) differential impact on different GHGs associated with a particular mitigation option, and (7) timeframe for abatement activities and the discount rate. In this section, we (a) provide aggregate mitigation potentials for the AFOLU sector (because these were provided separately for agriculture and forestry in AR4), (b) provide estimates of




Figure 11.12 | Relationship between technical, economic, and market potential (based on Smith, 2012).

global mitigation costs and potentials published since AR4, and (c) provide a regional disaggregation of the potentials to show how potential, and the portfolio of available options, varies in different world regions.

11.6.1 Approaches to estimating economic mitigation potentials

Bottom-up and top-down modelling approaches are used to estimate AFOLU mitigation potentials and costs. While both approaches provide useful estimates for mitigation costs and potentials, comparing bottom-up and top-down estimates is not straightforward.

Bottom-up estimates are typically derived for discrete abatement options in agriculture at a specific location or time, and are often based on detailed technological, engineering and process information, and data on individual technologies (DeAngelo et al., 2006). These studies provide estimates of how much technical potential of particular AFOLU mitigation options will become economically viable at certain carbon dioxide-equivalent prices. Bottom-up mitigation responses are typically restricted to input management (for example, changing practices with fertilizer application and livestock feeding) and mitigation costs estimates are considered 'partial equilibrium' in that the relevant input-output prices (and, sometimes, quantities such as area or production levels) are held fixed. As such, unless adjusted for potential overlaps and tradeoffs across individual mitigation options, adding up various individual estimates to arrive at an aggregate for a particular landscape or at a particular point in time could be misleading.

With a 'systems' approach, top-down models (described in Chapter 6; Section 11.9) typically take into account possible interactions between individual mitigation options. These models can be sector-specific or economy-wide, and can vary across geographical scales: sub-national, national, regional, and global. Mitigation strategies in top-down models may include a broad range of management responses and practice changes (for example, moving from cropping to grazing or grazing to forestry) as well as changes in input-output prices (for example, land and commodity prices). Such models can be used to assess the cost competitiveness of various mitigation options and implications across input-output markets, sectors, and regions over time for large-scale domestic or global adoption of mitigation strategies. In top-down modelling, dynamic cost-effective portfolios of abatement strategies are identified incorporating the lowest cost combination of mitigation strategies over time from across sectors, including agricultural, forestry, and other land-based sectors across the world that achieve the climate stabilization target (see Chapter 6). Top-down estimates for 2030 are included in this section, and are revisited in Section 11.9 when considering the role of the AFOLU sector in transformation pathways.

Providing consolidated estimates of economic potentials for mitigation within the AFOLU sector as a whole is complicated because of complex interdependencies, largely stemming from competing demands on land for various agricultural and forestry (production and mitigation) activi-

ties, as well as for the provision of many ecosystem services (Smith et al., 2013a). These interactions are discussed in more detail in Section 11.4.

11.6.2 Global estimates of costs and potentials in the AFOLU sector

Through combination of forestry and agriculture potentials from AR4, total mitigation potentials for the AFOLU sector are estimated to be ~3 to ~7.2 GtCO₂eq/yr in 2030 at 20 and 100 USD/tCO₂eq, respectively (Figure 11.13), including only supply-side options in agriculture (Smith et al., 2007) and a combination of supply- and demand-side options for forestry (Nabuurs et al., 2007).

Estimates of global economic mitigation potentials in the AFOLU sector published since AR4 are shown in Figure 11.14, with AR4 estimates shown for comparison (IPCC, 2007a).

Table 11.8 summarizes the ranges of global economic mitigation potentials from AR4 (Nabuurs et al., 2007; Smith et al., 2007), and studies published since AR4 that are shown in full in Figure 11.14, for agriculture, forestry, and AFOLU combined.

As described in Section 11.3, since AR4, more attention has been paid to options that reduce emissions intensity by improving the efficiency of production (i.e., less GHG emissions per unit of agricultural product; Burney et al., 2010; Bennetzen et al., 2012). As agricultural and silvicultural efficiency have improved over recent decades, emissions intensities have declined (Figure 11.15). Whilst emissions intensity has increased (1960s to 2000s) by 45% for cereals, emissions intensities have decreased by 38% for milk, 50% for rice, 45% for pig meat, 76% for chicken, and 57% for eggs.

The implementation of mitigation measures can contribute to further decrease emission intensities of AFOLU commodities (Figure 11.16; which shows changes of emissions intensities when a commodity-specific mix of mitigation measures is applied). For cereal production, mitigation measures considered include improved cropland agronomy, nutrient and fertilizer management, tillage and residue management, and the establishment of agro-forestry systems. Improved rice management practices were considered for paddy rice cultivation. Mitigation measures applied in the livestock sector include improved feeding and dietary additives. Countries can improve emission intensities of AFOLU commodities through increasing production at the same level of input, the implementation of mitigation measures, or a combination of both. In some regions, increasing current yields is still an option with a significant potential to improve emission intensities of agricultural production. Foley et al. (2011) analyzed current and potential yields that could be achieved for 16 staple crops using available agricultural practices and technologies and identified large 'yield gaps', especially across many parts of Africa, Latin America, and Eastern Europe. Better crop management practices can help to close yield gaps and improve emission intensities if measures are selected that also have a mitigation potential.




Figure 11.13 | Mitigation potential for the AFOLU sector, plotted using data from AR4 (Nabuurs et al., 2007; Smith et al., 2007). Whiskers show the range of estimates (± 1 standard deviation) for agricultural options for which estimates are available.

Mitigation potentials and costs differ largely between AFOLU commodities (Figure 11.16). While average abatement costs are low for roundwood production under the assumption of perpetual rotation, costs of mitigation options applied in meat and dairy production systems have a wide range (1:3 quartile range: 58–856 USD/tCO₂eq). Calculations of emission intensities are based on the conservative assumption that production levels stay the same after the application of the mitigation option. However, some mitigation options can increase production. This would not only improve food security but could also increase the cost-effectiveness of mitigation actions in the agricultural sector.

Agriculture and forestry-related mitigation could cost-effectively contribute to transformation pathways associated with long-run climate change management (Sections 11.9 and 6.3.5). Transformation pathway modelling includes LUC, as well as land-management options that reduce emissions intensities and increase sequestration intensities. However, the resulting transformation pathway emissions (sequestration) intensities are not comparable to those discussed here. Transformation pathways are the result of integrated modelling and the resulting intensities are the net result of many effects. The intensities capture mitigation technology adoption, but also changes in levels of production, land-cover change, mitigation technology

competition, and model-specific definitions for sectors/regions/and assigned emissions inventories. Mitigation technology competition, in particular, can lead to intensification (and increases in agricultural emissions intensities) that support cost-effective adoption of other mitigation strategies, such as afforestation or bioenergy (Sections 11.9 and 6.3.5).

11.6.3 Regional disaggregation of global costs and potentials in the AFOLU sector

Figure 11.17 shows the economically viable mitigation opportunities in AFOLU in 2030 by region and by main mitigation option at carbon prices of up to 20, 50, and 100 USD/tCO₂eq. The composition of the agricultural mitigation portfolio varies greatly with the carbon price (Smith, 2012), with low cost options such as cropland management being favoured at low carbon prices, but higher cost options such as restoration of cultivated organic soils being more cost-effective at higher prices. Figure 11.17 also reveals some very large differences in mitigation potential, and different ranking of most effective options, between regions. Across all AFOLU options, Asia has the largest mitigation potential, with the largest mitigation in both forestry and agriculture, followed by LAM, OECD-1990, MAF, and EIT.




Figure 11.14 | Estimates of economic mitigation potentials in the AFOLU sector published since AR4, (AR4 estimates shown for comparison, denoted by arrows), including bottom-up, sectoral studies, and top-down, multi-sector studies. Some studies estimate potential for agriculture and forestry, others for one or other sector. Supply-side mitigation potentials are estimated for around 2030, but studies range from estimates for 2025 (Rose et al., 2012) to 2035 (Rose and Sohngen, 2011). Studies are collated for those reporting potentials at carbon prices of up to ~20 USD/tCO₂eq (actual range 1.64–21.45), up to ~50 USD/tCO₂eq (actual range 31.39–50.00), and up to ~100 USD/tCO₂eq (actual range 70.0–120.91). Demand-side options (shown on the right-hand side of the figure) are for ~2050 and are not assessed at a specific carbon price, and should be regarded as technical potentials. Smith et al. (2013) values are mean of the range. Not all studies consider the same options or the same GHGs; further details are given in the text.




Figure 11.15 | GHG emissions intensities of selected major AFOLU commodities for decades 1960s–2000s, based on (Tubiello et al., 2012). i) Cattle meat, defined as GHG (enteric fermentation + manure management of cattle, dairy and non-dairy)/meat produced; ii) Pig meat, defined as GHG (enteric fermentation + manure management of swine, market and breeding)/meat produced; iii) Chicken meat, defined as GHG (manure management of chickens)/meat produced; iv) Milk, defined as GHG (enteric fermentation + manure management of cattle, dairy)/milk produced; v) Eggs, defined as GHG (manure management of chickens, layers)/egg produced; vi) Rice, defined as GHG (rice cultivation)/rice produced; vii) Cereals, defined as GHG (synthetic fertilizers)/cereals produced; viii) Wood, defined as GHG (carbon loss from harvest)/roundwood produced. Data Source: (FAOSTAT, 2013).




Figure 11.16 | Potential changes of emission intensities of major AFOLU commodities through implementation of commodity-specific mitigation measures (left panel) and related mitigation costs (right panel). Commodities and GHG emission sources are defined as in Figure 11.15, except for roundwood, expressed as the amount of carbon sequestered per unit roundwood from reforestation and afforestation within dedicated plantation cycles. Agricultural emission intensities represent regional averages, calculated based on 2000–2010 data (FAOSTAT, 2013) for selected commodities. Data on mitigation potentials and costs of measures are calculated using the mean values reported by (Smith et al., 2008) and the maximum and minimum are defined by the highest and lowest values for four climate zones for cereals and rice, or five geographical regions for milk and cattle meat. Emission intensities and mitigation potentials of roundwood production are calculated using data from Sathaye et al. (2005; 2006), FAO (2006), and IPCC (2006); maximum and minimum values are defined by the highest and lowest values for 10 geographical regions. The right panel shows the mitigation costs (in USD/tCO₂eq) of commodity-specific mitigation measures (25th to 75th percentile range).

Table 11.8 | Ranges of global mitigation potential (GtCO₂eq/yr) reported since AR4 | All values are for 2030 except demand-side options that are for ~2050 (full data shown in Figure 11.14).

	up to 20 USD/tCO ₂ eq	up to 50 USD/tCO ₂ eq	up to 100 USD/tCO ₂ eq	Technical potential only
Agriculture only¹	0–1.59	0.03–2.6	0.26–4.6	-
Forestry only	0.01–1.45	0.11–9.5	0.2–13.8	-
AFOLU total^{1,2}	0.12–3.03	0.5–5.06	0.49–10.6	-
Demand-side options	-	-	-	0.76–8.55

Notes:

¹ All lower range values for agriculture are for non-CO₂ GHG mitigation only and do not include soil C sequestration

² AFOLU total includes only estimates where both agriculture and forestry have been considered together.




Figure 11.17 | Economic mitigation potentials in the AFOLU sector by region. Agriculture values are from Smith et al. (2007). Forestry values are from Nabuurs et al. (2007). For forestry, 20 USD values correspond to 'low', and 100 USD values correspond to 'high' values from Nabuurs et al. (2007). Values of 50 USD represent the mean of the 'high' and 'low' values from Nabuurs et al. (2007).

Differences between the most effective forestry options in each region (Figure 11.18) are particularly striking, with reduced deforestation dominating the forestry mitigation potential LAM and MAF, but very little potential in OECD-1990 and EIT. Forest management, followed by afforestation, dominate in OECD-1990, EIT, and Asia (Figure 11.18). Among agricultural options, among the most striking of regional differences are the rice management practices for which almost all of the global potential is in Asia, and the large potential for restoration of organic soils also in Asia (due to cultivated Southeast Asian peats), and OECD-1990 (due to cultivated northern peatlands; Figure 11.18).

11.7 Co-benefits, risks, and spillovers

Implementation of AFOLU mitigation measures (Section 11.3) will result in a range of outcomes beyond changes in GHG balances with respect to institutional, economic, social, and environmental objectives. To the extent these effects are positive, they can be deemed 'co-benefits'; if adverse and uncertain, they imply risks.⁸ A global assessment of

the co-benefits and adverse side-effects of AFOLU mitigation measures is challenging for a number of reasons. First, co-benefits and adverse side-effects depend on the development context and the scale of the intervention (size), i.e., implementing the same AFOLU mitigation measure in two different areas (different countries or different regions within a country) can have different socio-economic, institutional, or environmental effects (Forner et al., 2006; Koh and Ghazoul, 2008; Trabucchi et al., 2008; Zomer et al., 2008; Alves Finco and Doppler, 2010; Alig et al., 2010, p. 201; Colfer, 2011; Davis et al., 2013; Albers and Robinson, 2013; Muys et al., 2014). Thus the effects are site-specific and generalizations are difficult. Second, these effects do not necessarily overlap geographically, socially, or over the same time scales (Section 11.4.5). Third, there is no general agreement on attribution of co-benefits and adverse side-effects to specific AFOLU mitigation measures; and fourth there are no standardized metrics for quantifying many of these effects. Modelling frameworks are being developed that allow an integrated assessment of multiple outcomes at landscape (Bryant et al., 2011), project (Townsend et al., 2012), and smaller (Smith et al., 2013a) scales. Table 11.9 presents an overview of the potential effects from AFOLU mitigation measures, while the text presents the most relevant co-benefits and potential adverse side-effects from the recent literature.

Maximizing co-benefits of AFOLU mitigation measures can increase efficiency in achieving the objectives of other international agreements, including the United Nations Convention to Combat Desertification (UNCCD, 2011), or the Convention on Biological Diversity (CBD), and mitigation actions may also contribute to a broader global sus-

⁸ Co-benefits and adverse side-effects describe effects in non-monetary units without yet evaluating the net effect on overall social welfare. Please refer to the respective sections in the framing chapters as well as to the glossary in Annex I for concepts and definitions—particularly Sections 2.4, 3.6.3, and 4.8.




Figure 11.18 | Regional differences in forestry options, shown as a proportion of total potential available in forestry in each region. Global forestry activities (annual amount sequestered or emissions avoided above the baseline for forest management, reduced deforestation and afforestation), at carbon prices up to 100 USD/tCO₂ are aggregated to regions from results from three models of global forestry and land use: the Global Timber Model (GTM; Sohngen and Sedjo, 2006), the Generalized Comprehensive Mitigation Assessment Process (Sathaye et al., 2006), and the Dynamic Integrated Model of Forestry and Alternative Land Use (Benítez et al., 2007).

tainability agenda (Harvey et al., 2010; Gardner et al., 2012; see Chapter 4). In many cases, implementation of these agendas is limited by capital, and mitigation may provide a new source of finance (Tubiello et al., 2009).

11.7.1 Socio-economic effects

AFOLU mitigation measures can affect institutions and living conditions of the various social groups involved. This section includes potential effects of AFOLU mitigation measures on three dimensions of sustainable development: institutional, social, and economic (Section 11.4.5).

AFOLU mitigation measures may have impacts on *land tenure and land-use rights* for several social groups including indigenous peoples, local communities and other social groups, dependant on natural assets. Co-benefits from AFOLU mitigation measures can be clarification of land tenure and harmonization of rights, while adverse side-effects can be lack of recognition of customary rights, loss of tenure or possession rights, and even displacement of social groups (Sunderlin et al., 2005; Chhatre and Agrawal, 2009; Blom et al., 2010; Sikor et al., 2010; Robinson et al., 2011; Rosemary, 2011; Larson, 2011; Rosendal and Andresen, 2011). Whether an impact on land tenure and use rights is positive or negative depends upon two factors: (a) the institutions regulating land tenure and land-use rights (e.g., laws, policies), and (b) the level of enforcement by such institutions (Corbera and Brown, 2008; Araujo et al., 2009; Rosemary, 2011; Larson et al., 2013; Albers and Robinson, 2013). More research is needed on specific tenure forms (e.g., individual property, state ownership or community rights), and on the specific effects from tenure and rights options, on enabling

AFOLU mitigation measures and co-benefits in different regions under specific circumstances (Sunderlin et al., 2005; Katila, 2008; Chhatre and Agrawal, 2009; Blom et al., 2010; Sikor et al., 2010; Robinson et al., 2011; Rosemary, 2011; Larson, 2011; Rosendal and Andresen, 2011).

AFOLU mitigation measures can support *enforcement of sectoral policies* (e.g., conservation policies) as well as *cross-sectoral coordination* (e.g., facilitating a landscape view for policies in the agriculture, energy, and forestry sectors (Brockhaus et al., 2013). However, AFOLU mitigation activities can also introduce or reduce clashes with existing policies in other sectors (e.g., if a conservation policy covers a forest area, where agricultural land is promoted by another policy (Madlener et al., 2006; Halsnæs and Verhagen, 2007; Smith et al., 2007; Beach et al., 2009; Alig et al., 2010; Jackson and Baker, 2010; DeFries and Rosenzweig, 2010; Pettenella and Brotto, 2011; Section 11.10).

An area of increasing concern since AR4 is the potential impact of AFOLU mitigation measures on *food security*. Efforts to reduce hunger and malnutrition will increase individual food demand in many developing countries, and population growth will increase the number of individuals requiring secure and nutritionally sufficient food production. Thus, a net increase in food production is an essential component for securing sustainable development (Ericksen et al., 2009; FAO, WFP, and IFAD, 2012). AFOLU mitigation measures linked to increases in food production (e.g., agroforestry, intensification of agricultural production, or integrated systems) can increase food availability and access especially at the local level, while other measures (e.g., forest or energy crop plantations) can reduce food production at least locally (Foley et al., 2005; McMichael et al., 2007;

Pretty, 2008; Godfray et al., 2010; Jackson and Baker, 2010; Graham-Rowe, 2011; Jeffery et al., 2011).

Regarding *human health* reduced emissions from agriculture and forestry may also improve air, soil, and water quality (Smith et al., 2013a), thereby indirectly providing benefits to human health and well-being. Demand-side measures aimed at reducing the proportion of livestock products in human diets that are high in animal products are also

associated with multiple health benefits (McMichael et al., 2007; Stehfest et al., 2009; Marlow et al., 2009). AFOLU mitigation measures, particularly in the livestock sector, can have an impact on *animal welfare* (Sundrum, 2001; Lund and Algers, 2003; Keeling et al., 2011; Kehl-bacher et al., 2012; Koknaroglu and Akunal, 2013).

A major area of concern is related to the potential impacts of AFOLU mitigation measures on *equity* (Sections 3.3; 4.2; 4.7;

Box 11.6 | Challenges for mitigation in developing countries in the AFOLU sector

Mitigation challenges related to the AFOLU sector

The contribution of developing countries to future GHG emissions is expected to be very significant due to projected increases in food production by 2030 driving short-term land conversion in these countries. Mitigation efforts in the AFOLU sector rely mainly on reduction of GHG emissions and an increase in carbon sequestration (Table 11.2). Potential activities include reducing deforestation, increasing forest cover, agroforestry, agriculture and livestock management, and production of sustainable renewable energy (Sathaye et al., 2005; Smith et al., 2013b). Although agriculture and forestry are important sectors for GHG abatement (Section 11.2.3), it is likely that technology alone will not be sufficient to deliver the necessary transitions to a low-GHG future (Alig et al., 2010; Section 11.3.2). Other barriers include access to market and credits, technical capacities to implement mitigation options, including accurate reporting of emission levels and emission factors based on activity data, and institutional frameworks and regulations (Corbera and Schroeder, 2011; Mbow et al., 2012; Sections 11.7; 11.8). Additionally, the diversity of circumstances among developing countries makes it difficult to establish the modelled relationships between GDP and CO₂ emissions per capita found by using the Kaya identity. This partly arises from the wide gap between rural and urban communities, and the difference in livelihoods (e.g., the use of fuel wood, farming practices in various agro-ecological conditions, dietary preferences with a rising middle class in developing countries, development of infrastructure, and behavioural change, etc.; Lambin and Meyfroidt, 2011). Also, some mitigation pathways raise the issue of non-permanence and leakage that can lead to the transfer activities to non-protected areas, which may threaten conservation areas in countries with low capacities (Lippke et al., 2003; Jackson and Baker, 2010; Section 11.3.2).

Critical issues to address are the co-benefits and adverse side-effects associated with changed agricultural production, the necessary link between mitigation and adaptation, and how to manage incentives for a substantial GHG abatement initiative without compromising food security (Smith and Wollenberg, 2012; Sections 11.5; 11.7). The challenge is to strike a balance between emissions reductions/adaptation and development/poverty

alleviation priorities, or to find policies that co-deliver. Mitigation pathways in developing countries should address the dual need for mitigation and adaptation through clear guidelines to manage multiple options (Section 11.5.4). Prerequisites for the successful implementation of AFOLU mitigation projects are ensuring that (a) communities are fully engaged in implementing mitigation strategies, (b) any new strategy is consistent with ongoing policies or programmes, and (c) *a priori* consent of small holders is given. Extra effort is required to address equity issues including gender, challenges, and prospects (Mbow et al., 2012).

Mitigation challenges related to the bioenergy sector

Bioenergy has a significant mitigation potential, provided that the resources are developed sustainably and that bioenergy systems are efficient (Chum et al., 2011; Section 11.9.1). Bioenergy production can be integrated with food production in developing countries, e.g., through suitable crop rotation schemes, or use of by-products and residues (Berndes et al., 2013). If implemented sustainably this can result in higher food and energy outcomes and hence reduce land-use competition. Some bioenergy options in developing countries include perennial cropping systems, use of biomass residues and wastes, and advanced conversion systems (Beringer et al., 2011; Popp et al., 2011; Box 7.1). Agricultural and forestry residues can provide low-carbon and low-cost feedstock for bioenergy. Biomass from cellulosic bioenergy crops feature substantially in future energy systems, especially in the framework of global climate policy that aims at stabilizing CO₂ concentration at low levels (Popp et al., 2011; Section 11.13). The large-scale use of bioenergy is controversial in the context of developing countries because of the risk of reducing carbon stocks and releasing carbon to the atmosphere (Bailis and McCarthy, 2011), threats to food security in Africa (Mbow, 2010), and threats to biodiversity *via* the conversion of forests to biofuel (e.g., palm oil) plantations. Several studies underline the inconsistency between the need for bioenergy and the requirement for, e.g., Africa, to use its productive lands for sustainable food production (Cotula et al., 2009). Efficient biomass production for bioenergy requires a range of sustainability requirements to safeguard food production, biodiversity, and terrestrial carbon storage.

4.8). Depending on the actual and perceived distribution of socio-economic benefits, responsibilities (burden sharing), as well the access to decision making, financing mechanisms, and technology, AFOLU mitigation measures can promote inter- and intra-generational equity (Di Gregorio et al., 2013). Conversely, depending on the policy instruments and the implementation schemes of these mitigation measures, they can increase inequity and land conflicts, or marginalize small-scale farm/forest owners or users (Robinson et al., 2011; Kiptot and Franzel, 2012; Huettner, 2012; Mattoo and Subramanian, 2012). Potential impacts on equity and benefit-sharing mechanisms arise for AFOLU activities using forestry measures in developing countries including conservation, restoration, reduced deforestation and degradation, as well as sustainable management and afforestation/reforestation (Combes Motel et al., 2009; Cattaneo et al., 2010; Rosemary, 2011).

Large-scale land acquisition (often referred to as 'land grabbing') related to the promotion of AFOLU mitigation measures (especially for production of bioenergy crops) and its links to sustainable development in general, and equity in particular, are emerging issues in the literature (Cotula et al., 2009; Scheidel and Sorman, 2012; Mwakaje, 2012; Messerli et al., 2013; German et al., 2013).

In many cases, the implementation of agricultural and forestry systems with positive impacts mitigating climate change are limited by capital, and *carbon payments or compensation mechanisms* may provide a new source of finance (Tubiello et al., 2009). For instance, in some cases, mitigation payments can help to make production of non-timber forest products (NTFP) economically viable, further *diversifying income* at the local level (Singh, 2008). However, depending on the accessibility of the financing mechanisms (payments, compensation, or other) economic benefits can become concentrated, marginalizing many local stakeholders (Combes Motel et al., 2009; Alig et al., 2010; Asante et al., 2011; Asante and Armstrong, 2012; Section 11.8). The realization of economic co-benefits is related to the design of the specific mechanisms and depends upon three main variables: (a) the amount and coverage of these payments, (b) the recipient of the payments, and (c) timing of payments (*ex-ante* or *ex-post*; Corbera and Brown, 2008; Skutsch et al., 2011). Further considerations on financial mechanisms and carbon payments, both within and outside UNFCCC agreements, are described in Section 11.10.

Financial flows supporting AFOLU mitigation measures (e.g., those resulting from the REDD+) can have positive effects on conserving biodiversity, but could eventually create conflicts with conservation of biodiversity hotspots, when their respective carbon stocks are low (Gardner et al., 2012; Section 11.10). Some authors propose that carbon payments can be complemented with biodiversity payments as an option for reducing tradeoffs with biodiversity conservation (Phelps et al., 2010a). Bundling of ecosystem service payments, and links to carbon payments, is an emerging area of research (Deal and White, 2012).

11.7.2 Environmental effects

Availability of land and land competition can be affected by AFOLU mitigation measures. Different stakeholders may have different views on what land is available, and when considering several AFOLU mitigation measures for the same area, there can be different views on the importance of the goods and ecosystem services provided by the land, e.g., some AFOLU measures can increase food production but reduce water availability or other environmental services. Thus decision makers need to be aware of potential site-specific tradeoffs within the sector. A further potential adverse side-effect is that of increasing land rents and food prices due to a reduction in land availability for agriculture in developing countries (Muller, 2009; Smith et al., 2010, 2013b; Rathmann et al., 2010; Godfray et al., 2010; de Vries and de Boer, 2010; Harvey and Pilgrim, 2011; Amigun et al., 2011; Janzen, 2011; Cotula, 2012; Scheidel and Sorman, 2012; Haberl et al., 2013a).

AFOLU mitigation options can promote conservation of *biological diversity* (Smith et al., 2013a) both by reducing deforestation (Chhatre et al., 2012; Murdiyarto et al., 2012; Putz and Romero, 2012; Visseren-Hamakers et al., 2012), and by using reforestation/afforestation to restore biodiverse communities on previously developed farmland (Harper et al., 2007). However, promoting land-use changes (e.g., through planting monocultures on biodiversity hot spots) can have adverse side-effects, reducing biodiversity (Koh and Wilcove, 2008; Beringer et al., 2011; Pandit and Grumbine, 2012; Ziv et al., 2012; Hertwich, 2012; Gardner et al., 2012).

In addition to potential climate impacts, land-use intensity drives the three main N loss pathways (nitrate leaching, denitrification, and ammonia volatilization) and typical *N balances* for each land use indicate that total N losses also increase with increasing land-use intensity (Stevenson et al., 2010). Leakages from the N cycle can cause air (e.g., ammonia (NH₃⁺), nitrogen oxides (NO_x)⁹), soil nitrate (NO₃⁻) and water pollution (e.g., eutrophication), and agricultural intensification can lead to a variety of other adverse environmental impacts (Smith et al., 2013a). Combined strategies (e.g., diversified crop rotations and organic N sources) or single-process strategies (e.g., reduced N rates, nitrification inhibitors, and changing chemical forms of fertilizer) can reduce N losses (Bambo et al., 2009; Gardner and Drinkwater, 2009). Integrated systems may be an alternative approach to reduce leaching (Section 11.10).

AFOLU mitigation measures can have either positive or negative *impacts on water resources*, with responses dependant on the mitigation measure used, site conditions (e.g., soil thickness and slope, hydrological setting, climate; Yu et al., 2013) and how the particular mitigation measure is managed. There are two main components: water yield and water quality. Water yields can be manipulated with forest management, through afforestation, reforestation, for-

⁹ Please see Section 7.9.2 and WGII Section 11.9 for a discussion of health effects related to air pollution.

Table 11.9 | Summary of potential co-benefits (green arrows) and adverse side-effects (orange arrows) from AFOLU mitigation measures; arrows pointing up/down denote positive/negative effect on the respective issue. These effects depend on the specific context (including bio-physical, institutional, and socio-economic aspects) as well as on the scale of implementation. For an assessment of macroeconomic, cross-sectoral effects associated with mitigation policies (e.g., on energy prices, consumption, growth, and trade), see Sections 3.9, 6.3.6, 13.2.2.3, and 14.4.2. Note: Co-benefits/adverse side-effects of bioenergy are discussed in Section 11.13.

	Issue	Potential co-benefit or adverse side-effect	Scale	AFOLU mitigation measure
Institutional	Land tenure and use rights	Improving (↑) or diminishing (↓) tenure and use rights for local communities and indigenous peoples, including harmonization of land tenure and use regimes (e.g., with customary rights)	Local to national	Forestry (4, 5, 6, 8, 9, 12, 20)
	Sectoral policies	Promoting (↑) or contradicting (↓) the enforcement of sectoral (forest and/or agriculture) policies	National	Forestry (2, 5, 6, 9, 20); land-based agriculture (7, 11, 20)
	Cross-sectoral policies	Cross-sectoral coordination (↑) or clashes (↓) between forestry, agriculture, energy, and/or mining policies	Local to national	Forestry (7, 20); agriculture (7, 11, 20)
	Participative mechanisms	Creation/use of participative mechanisms (↑) for decision making regarding land management (including participation of various social groups, e.g., indigenous peoples or local communities)	Local to national	Forestry (4, 5, 6, 8, 9, 14, 20); agriculture (20, 32); integrated systems (20, 34)
	Benefit sharing mechanisms	Creation/use of benefits-sharing mechanisms (↑) from AFOLU mitigation measures	Local to national	Forestry (4, 5, 6, 8, 20)
Social	Food security	Increase (↑) or decrease (↓) on food availability and access	Local to national	Forestry (18, 19); agriculture (7, 15, 18, 19, 23, 28, 30); livestock (2, 3, 19, 35, 36); integrated systems (18, 19); biochar (17, 26)
	Local/traditional knowledge	Recognition (↑) or denial (↓) of indigenous and local knowledge in managing (forest/agricultural) land	Local/sub-national	Forestry (4, 5, 6, 8, 20); agriculture (20, 28); integrated systems (2); livestock (2, 3, 35); biochar (2)
	Animal welfare	Changes in perceived or measured animal welfare (perceived due to cultural values or measured, e.g., through amount of stress hormones)	Local to national	Livestock (2, 31, 35, 37, 38)
	Cultural values	Respect and value cultural habitat and traditions (↑), reduce (↓), or increase (↑) existing conflicts or social discomfort (4, 5, 6, 20, 8)	Local to trans-boundary	Forestry (4, 5, 6, 9, 20)
	Human health	Impacts on health due to dietary changes, especially in societies with a high consumption of animal protein (↓)	Local to global	Changes in demand patterns (31, 36)
	Equity	Promote (↑) or not (↓) equal access to land, decision making, value chain, and markets as well as to knowledge- and benefit-sharing mechanisms	Local to global	Forestry (4, 5, 6, 8, 9, 10, 20); agriculture (11, 23, 32)
Economic	Income	Increase (↑) or decrease (↓) in income. There are concerns regarding income distribution (↑)	Local	Forestry (6, 7, 8, 16, 20, 21, 22); agriculture (16, 19, 20, 23, 28); livestock (2, 3); integrated systems (7, 20); biochar (24); changes in demand patterns (2)
	Employment	Employment creation (↑) or reduction of employment (especially for small farmers or local communities) (↓)	Local	Forestry (8, 20); agriculture (20, 23); livestock (2, 3); integrated systems (7, 20)
	Financing mechanisms	Access (↑) or lack of access (↓) to new financing schemes	Local to global	Forestry (6, 8, 16, 20); agriculture (16, 20); livestock (2, 3)
	Economic activity	Diversification and increase in economic activity (↑) while concerns on equity (↑)	Local	Forestry (6, 7, 8, 20); land-based agriculture (16, 19, 20, 23, 28); livestock (2, 3)
Environmental	Land availability	Competition between land uses and risk of activity or community displacement (↑)	Local to trans-boundary	Forestry and land-based-agriculture (5, 6, 15, 18, 20, 29, 30); livestock (2, 3, 29, 40)
	Biodiversity	Monocultures can reduce biodiversity (↓). Ecological restoration increases biodiversity and ecosystem services (↑) by 44 and 25% respectively (28). Conservation, forest management, and integrated systems can keep biodiversity (↑) and/or slow desertification (↓)	Local to trans-boundary	Forestry (1, 19, 20, 27); on conservation and forest management (1, 19, 21, 27, 30); agriculture and integrated systems (15, 19, 20, 28, 30)
	Albedo	Positive impacts (↑) on albedo and evaporation and interactions with ozone	Local to global	See Section 11.5
	N and P cycles	Impacts on N and P cycles in water (↓/↑) especially from monocultures or large agricultural areas	Local to trans-boundary	Agriculture (19, 23, 30, 35); livestock (2, 3, 30)
	Water resources	Monocultures and/or short rotations can have negative impacts on water availability (↓). Potential water depletion due to irrigation (↓). Some management practices can support regulation of the hydrological cycle and protection of watersheds (↑)	Local to trans-boundary	Forestry (1, 19, 20, 27); land-based agriculture (30, 44); integrated systems (2, 30, 44)
	Soil	Soil conservation (↑) and improvement of soil quality and fertility (↑). Reduction of erosion. Positive or negative carbon mineralization priming effect (↑/↓)	Local	Forestry (44, 45); land-based agriculture (13, 19, 23, 28, 30); integrated systems biochar (39, 40)
	New products	Increase (↑) or decrease (↓) on fibre availability as well as non-timber/non-wood products output	Local to national	Forestry (18, 19, 41, 42); agriculture (7, 15, 18, 19, 23, 28, 30); integrated systems (18, 19)
Ecosystem resilience	Increase (↑) or reduction (↓) of resilience, reduction of disaster risks (↓)	Local to trans-boundary	Forestry, integrated systems (11, 33; see Section 11.5)	



	Issue	Potential co-benefit or adverse side-effect	Scale	AFOLU mitigation measure
Technology	Infrastructure	Increase (↑) or decrease (↓) in availability of and access to infrastructure. Competition for infrastructure for agriculture (↑), can increase social conflicts	Local	Agriculture (20, 46, 47)
	Technology innovation and transfer	Promote (↑) or delay (↓) technology development and transfer	Local to global	Forestry (7, 13, 25); agriculture (23); livestock (2, 3)
	Technology Acceptance	Can facilitate acceptance of sustainable technologies (↑)	Local to national	Forestry (7, 13, 25); livestock (2, 3, 35)

Notes: AFOLU mitigation measures are grouped following the structure given in Table 11.2

Sources: ¹Trabucco et al., 2008; ²Steinfeld et al., 2010; ³Gerber et al., 2010; ⁴Sikor et al., 2010; ⁵Rosemary, 2011; ⁶Pettenella and Brotto, 2011; ⁷Jackson and Baker, 2010; ⁸Corbera and Schroeder, 2011; ⁹Colfer, 2011; ¹⁰Blom et al., 2010; ¹¹Halsnæs and Verhagen, 2007; ¹²Larson, 2011; ¹³Lichtfouse et al., 2009; ¹⁴Thompson et al., 2011; ¹⁵Graham-Rowe, 2011; ¹⁶Tubiello et al., 2009; ¹⁷Barrow, 2012; ¹⁸Godfray et al., 2010; ¹⁹Foley et al., 2005; ²⁰Madlener et al., 2006; ²¹Strassburg et al., 2012; ²²Canadell and Raupach, 2008; ²³Pretty, 2008; ²⁴Galinato et al., 2011; ²⁵Macauley and Sedjo, 2011; ²⁶Jeffery et al., 2011; ²⁷Benayas et al., 2009; ²⁸Foley et al., 2011; ²⁹Haberl et al., 2013; ³⁰Smith et al., 2013; ³¹Stehfest et al., 2009; ³²Chhatre et al., 2012; ³³Seppälä et al., 2009; ³⁴Murdiyarso et al., 2012; ³⁵de Boer et al., 2011; ³⁶McMichael et al., 2007; ³⁷Koknaroglu and Akunal, 2013; ³⁸Kehlbacher et al., 2012; ³⁹Zimmerman et al., 2011; ⁴⁰Luo et al., 2011; ⁴¹Mirle, 2012; ⁴²Albers and Robinson, 2013; ⁴³Smith et al., 2013a; ⁴⁴Chatterjee and Lal, 2009; ⁴⁵Smith, 2008; ⁴⁶Ziv et al., 2012; ⁴⁷Beringer et al., 2011; ⁴⁸Douglas et al., 2009

est thinning, or deforestation. In general, reduction in water yields in afforestation/reforestation projects has been reported in both groundwater or surface catchments (Jackson et al., 2005), or where irrigation water is used to produce bioenergy crops. For water supply security, it is important to consider the relative yield reduction and this can have severe consequences in dry regions with inherent water shortages (Wang et al., 2011c). Where there is a water imbalance, however, this additional water use can be beneficial by reducing the efflux of salts (Jackson et al., 2005). Another aspect of water yield is the reduction of flood peaks, and also prolonged periods of water flow, because discharge is stabilized (Jackson et al., 2005), however low flows can be reduced because of increased forest water use. Water quality can be affected by AFOLU in several ways. For example, minimum tillage systems have been reported to reduce water erosion and thus sedimentation of water courses (Lal, 2011). Deforestation is well known to increase erosion and thus efflux of silt; avoiding deforestation will prevent this. In other situations, watershed scale reforestation can result in the restoration of water quality (e.g., Townsend et al., 2012). Furthermore, strategic placement of tree belts in lands affected by dryland salinity can remediate the affected lands by lowering the water table (Robinson et al., 2004). Various types of AFOLU mitigation can result in degradation of water sources through the losses of pesticides and nutrients to water (Smith et al., 2013a).

AFOLU mitigation measures can have several *impacts on soil*. Increasing or maintaining carbon stocks in living biomass (e.g., through forest or agroforestry systems) will reduce wind erosion by acting as wind breaks and may increase crop production; and reforestation, conservation, forest management, agricultural systems, or bioenergy systems can be used to restore degraded or abandoned land (Smith et al., 2008; Stickler et al., 2009; Chatterjee and Lal, 2009; Wicke et al., 2011b; Sochacki et al., 2012). Silvo-pastoral systems can help to reverse land degradation while providing food (Steinfeld et al., 2008, 2010; Janzen, 2011). Depending on the soil type, production temperature regimes,

the specific placement and the feedstock tree species, biochar can have positive or negative carbon mineralization priming effects over time (Zimmerman et al., 2011; Luo et al., 2011).

AFOLU mitigation options can promote innovation, and many technological supply-side mitigation options outlined in Section 11.3 also increase agricultural and silvicultural efficiency. At any given level of demand for agricultural products, intensification increases output per unit area and would therefore, if all else were equal, allow the reduction in farmland area, which would in turn free land for C sequestration and/or bioenergy production (Section 11.4). For example, a recent study calculated potentially large GHG reductions from global agricultural intensification by comparing the past trajectory of agriculture (with substantial yield improvements), with a hypothetical trajectory with constant technology (Burney et al., 2010). However, in real-world situations increases in yield may result in feedbacks such as increased consumption ('rebound effects'; see Section 11.4; Lambin and Meyfroidt, 2011; Erb, 2012).

11.7.3 Public perception

Mitigation measures that support sustainable development are likely to be viewed positively in terms of public perception, but a large-scale drive towards mitigation without inclusion of key stakeholder communities involved would likely not be greeted favourably (Smith and Wollenberg, 2012). However, there are concerns about competition between food and AFOLU outcomes, either because of an increasing use of land for biofuel plantations (Fargione et al., 2008; Alves Finco and Doppler, 2010), or afforestation/reforestation (Mitchell et al., 2012), or by blocking the transformation of forest land into agricultural land (Harvey and Pilgrim, 2011).

Further, lack of clarity regarding the architecture of the future international climate regime and the role of AFOLU mitigation measures is perceived as a potential threat for long-term planning and long-term investments (Streck, 2012; Visseren-Hamakers et al., 2012). Certain tech-

nologies, such as animal feed additives and genetically modified organisms are banned in some jurisdictions due to perceived health and/or environmental risks. Public perception is often as important as scientific evidence of hazard/risk in considering government policy regarding such technologies (Royal Society, 2009; Smith and Wollenberg, 2012).

11.7.4 Spillovers

Emerging knowledge on the importance of ecosystems services as a means for addressing climate change mitigation and adaptation have brought attention to the role of ecosystem management for achieving several development goals, beyond climate change adaptation and mitigation. This knowledge has enhanced the creation of ecosystem markets (Section 11.10). In some jurisdictions ecosystem markets are developing (MEA, 2005; Engel et al., 2008; Deal and White, 2012; Wünscher and Engel, 2012) and these allow valuation of various components of land-use changes, in addition to mitigation (Mayrand and Paquin, 2004; Barbier, 2007). Different approaches are used; in some cases the individual components (both co-benefits and adverse side-effects) are considered singly (bundled), in other situations they are considered together (stacked) (Deal and White, 2012). Ecosystem market approaches can serve as a framework to assess the benefits of mitigation actions from project, to regional and national level (Farley and Costanza, 2010). Furthermore, designing ecosystem market approaches yields methodologies for the evaluation of individual components (e.g., water quality response to reforestation, timber yield), and other types of ecosystem service (e.g., biodiversity, social amenity; Bryan et al., 2013).

11.8 Barriers and opportunities

Barriers and opportunities refer to the conditions provided by the development context (Section 11.4.5). These conditions can enable and facilitate (opportunities) or hinder (barriers) the full use of AFOLU mitigation measures. AFOLU programmes and policies can help to overcome barriers, but countries being affected by many barriers will need time, financing, and capacity support. In some cases, international negotiations have recognized these different circumstances among countries and have proposed corresponding approaches (e.g., a phased approach in the REDD+, Green Climate Fund; Section 11.10). Corresponding to the development framework presented in Section 11.4.5, the following types of barriers and benefits are discussed: socio-economic, environmental, institutional, technological, and infrastructural.

11.8.1 Socio-economic barriers and opportunities

The *design and coverage of the financing mechanisms* is key to successful use of the AFOLU mitigation potential (Section 11.10; Chapter

16). Questions remain over which costs will be covered by such mechanisms. If financing mechanisms fail to cover at least transaction and monitoring costs, they will become a barrier to the full implementation of AFOLU mitigation. According to some studies, opportunity costs also need to be fully covered by any financing mechanism for the AFOLU sector, especially in developing countries, as otherwise AFOLU mitigation measures would be less attractive compared to returns from other land uses (Angelsen, 2008; Cattaneo et al., 2010; Böttcher et al., 2012). Conversely, if financing mechanisms are designed to modify economic activity, they could provide an opportunity to leverage a larger proportion of AFOLU mitigation potential.

Scale of financing sources can become either a barrier (if a relevant financial volume is not secured) or create an opportunity (if financial sources for AFOLU suffice) for using AFOLU mitigation potential (Streck, 2012; Chapter 16). Another element is the *accessibility to AFOLU financing* for farmers and forest stakeholders (Tubiello et al., 2009, p. 200; Havemann, 2011; Colfer, 2011). Financial concerns, including reduced access to loan and credits, high transaction costs or reduced income due to price changes of carbon credits over the project duration, are potential risks for AFOLU measures, especially in developing countries, and when land holders use market mechanisms (e.g., Afforestation and Reforestation (A/R) Clean Development Mechanism (CDM); Madlener et al., 2006).

Poverty is characterized not only by low income, but also by insufficient food availability in terms of quantity and/or quality, limited access to decision making and social organization, low levels of education and reduced access to resources (e.g., land or technology; UNDP International Poverty Centre, 2006). High levels of poverty can limit the possibilities for using AFOLU mitigation options, because of short-term priorities and lacking resources. In addition, poor communities have limited skills and sometimes lack of social organization that can limit the use and scaling up of AFOLU mitigation options, and can increase the risk of displacement, with other potential adverse side-effects (Smith and Wollenberg, 2012; Huettner, 2012). This is especially relevant when forest land sparing competes with other development needs e.g., increasing land for agriculture or promoting some types of mining (Forner et al., 2006), or when large-scale bioenergy compromises food security (Nonhebel, 2005; Section 11.13).

Cultural values and social acceptance can determine the feasibility of AFOLU measures, becoming a barrier or an opportunity depending of the specific circumstances (de Boer et al., 2011).

11.8.2 Institutional barriers and opportunities

Transparent and accountable governance and swift institutional establishment are very important for a sustainable implementation of AFOLU mitigation measures. This includes the need to have *clear land tenure and land-use rights* regulations and a certain level of enforcement, as well as clarity about carbon ownership (Palmer, 2011; Thompson et al.,

2011; Markus, 2011; Rosendal and Andresen, 2011; Murdiyarso et al., 2012 Sections 11.4.5; 11.10; Chapters 14; 15).

Lack of institutional capacity (as a means for securing creation of equal institutions among social groups and individuals) can reduce feasibility of AFOLU mitigation measures in the near future, especially in areas where small-scale farmers or forest users are the main stakeholders (Laitner et al., 2000; Madlener et al., 2006; Thompson et al., 2011a). *Lack of an international agreement* that supports a wide implementation of AFOLU measures can become a major barrier for realizing the mitigation potential from the sector globally (Section 11.10; Chapter 13).

11.8.3 Ecological barriers and opportunities

Mitigation potential in the agricultural sector is highly site-specific, even within the same region or cropping system (Baker et al., 2007; Chatterjee and Lal, 2009). *Availability of land and water* for different uses need to be balanced, considering short- and long-term priorities, and global differences in resource use. Consequently, limited resources can become an ecological barrier and the decision of how to use them needs to balance ecological integrity and societal needs (Jackson, 2009).

At the local level, the *specific soil conditions, water availability, GHG emission-reduction potential as well as natural variability and resilience* to specific systems will determine the level of realization of mitigation potential of each AFOLU measure (Baker et al., 2007; Halvorson et al., 2011). Frequent droughts in Africa and changes in the hydro-meteorological events in Asia and Central and South America are important in defining the specific regional potential (Bradley et al., 2006; Rotenberg and Yakir, 2010). Ecological saturation (e.g., soil carbon or yield) means that some AFOLU mitigation options have their own limits (Section 11.5). The fact that many *AFOLU measures can provide adaptation benefits* provides an opportunity for increasing ecological efficiency (Guariguata et al., 2008; van Vuuren et al., 2009; Robledo et al., 2011; Section 11.5).

11.8.4 Technological barriers and opportunities

Technological barriers refer to the limitations in generating, procuring, and applying science and technology to identify and solve an environmental problem. Some mitigation technologies are already applied now (e.g., afforestation, cropland, and grazing land management, improved livestock breeds and diets) so there are no technological barriers for these options, but others (e.g., some livestock dietary additives, crop trait manipulation) are still at the development stage (see Table 11.2).

The *ability to manage and re-use knowledge assets* for scientific communication, technical documentation and learning is lacking in many areas where mitigation could take place. Future developments pres-

ent opportunities for additional mitigation to be realized if efforts to deliver ease-of-use and range-of-use are guaranteed. There is also a need to adapt technology to local needs by focusing on existing local opportunities (Kandji et al., 2006), as proposed in Nationally Appropriate Mitigation Actions (NAMAs) (Section 11.10).

Barriers and opportunities related to *monitoring, reporting, and verification* of the progress of AFOLU mitigation measures also need be considered. Monitoring activities, aimed at reducing uncertainties, provide the opportunity of increasing credibility in the AFOLU sector. However there are technical challenges. For instance, monitoring carbon in forests with high spatial variability in species composition and tree density can pose a technical barrier to the implementation of some AFOLU activities (e.g., REDD+; Baker et al., 2010; Section 11.10). The IPCC National Greenhouse Gas Inventory Guidelines (Paustian et al., 2006) also provide an opportunity, because they offer standard scientific methods that countries already use to report AFOLU emissions and removals under the UNFCCC. Also, field research in high-biomass forests (Gonzalez et al., 2010) shows that remote sensing data and Monte Carlo quantification of uncertainty offer a technical opportunity for implementing REDD+ (Section 11.10). Exploiting the existing *human skills* within a country is essential for realizing full AFOLU potential. A lack of trained people can therefore become a barrier to implementation of appropriate technologies (Herold and Johns, 2007).

Technology improvement and technology transfer are two crucial components for the sustainable increase of agricultural production in developed and developing regions with positive impacts in terms of mitigation, soil, and biodiversity conservation (Tilman et al., 2011). International and national policy instruments are relevant to foster technology transfer and to support research and development (Section 11.10.4), overcoming technological barriers.

11.9 Sectoral implications of transformation pathways and sustainable development

Some climate change management objectives require large-scale transformations in human societies, in particular in the production and consumption of energy and the use of the land resource. Chapter 6 describes alternative 'transformation pathways' of societies over time from now into the future, consistent with different climate change outcomes. Many pathways that foresee large efforts in mitigation will have implications for sustainable development, and corrective actions to move toward sustainability may be possible. However, impacts on development are context specific and depend upon scale and institutional agreements of the AFOLU options, and not merely on the type of option (see Sections 11.4 for development

context and systemic view, 11.7 for potential co-benefits and adverse effects, and 11.8 for opportunities and challenges). To evaluate sectoral implications of transformation pathways, it is useful to first characterize the pathways in terms of mitigation technologies and policy assumptions.

11.9.1 Characterization of transformation pathways

Uncertainty about reference AFOLU emissions is significant both historically (Section 11.2) and in projections (Section 6.3.1.3). The transformation projections of the energy system, AFOLU emissions and land-use are characterized by the reference scenario, as well as the abatement policy assumptions regarding eligible abatement options, regions covered, and technology costs over time. Many mitigation scenarios suggest a substantial cost-effective mitigation role for land related mitigation assuming idealized policy implementation, with immediate, global, and comprehensive availability of land-related mitigation options. However, policy implementation of large-scale land-based mitigation will be challenging. In addition, the transformation

pathways often ignore, or only partially cover, important mitigation risks, costs, and benefits (e.g., transaction costs or Monitoring Reporting and Verification (MRV) costs), and other developmental issues including intergenerational debt or non-monetary benefits (Ackerman et al., 2009; Lubowski and Rose, 2013).

In recent idealized implementation scenarios from a model comparison study, land-related changes can represent a significant share of emissions reductions (Table 11.10). In these scenarios, models assume an explicit terrestrial carbon stock incentive, or a global forest protection policy, as well as an immediate global mitigation policy in general. Bioenergy is consistently deployed (because it is considered to reduce net GHG emissions over time; see Section 6.3.5), and agricultural emissions are priced. Note that bioenergy related mitigation is not captured in Table 11.10. The largest land emission reductions occur in net CO₂ emissions, which also have the greatest variability across models. Some models exhibit increasing land CO₂ emissions under mitigation, as bioenergy feedstock production leads to LUC, while other models exhibit significant reductions with protection of existing terrestrial carbon stocks and planting of new trees to increase carbon stocks. Land-related CO₂ and N₂O mitigation is more important in the nearer-term

Table 11.10 | Cumulative land-related emissions reductions, land reduction share of global reductions, and percent of baseline land emissions reduced for CH₄, CO₂, and N₂O in idealized implementation 550 and 450 ppm CO₂eq scenarios. The number of scenarios is indicated for each GHG and atmospheric concentration goal. Negative values represent increases in emissions (Kriegler et al., 2013). Bioenergy-related mitigation is not captured in the table.

			550 ppm			450 ppm		
			2010–2030	2010–2050	2010–2100	2010–2030	2010–2050	2010–2100
Cumulative global land-related emissions reductions (GtCO ₂ eq)	CH ₄ (n = 5/5)	min	3.5	17.5	51.4	0.0	4.5	52.3
		max	9.8	46.0	201.7	12.7	50.5	208.6
	CO ₂ (n = 11/10)	min	-20.2	-43.2	-129.8	-20.3	-50.8	-153.9
		max	280.9	543.0	733.4	286.6	550.5	744.6
	N ₂ O (n = 4/4)	min	3.1	8.4	25.5	3.1	8.4	25.5
		max	8.2	27.7	96.6	9.7	29.3	96.8
	Sum (n = 4/4)	min	-8.7	2.5	53.9	-3.7	5.6	69.7
max		295.2	587.7	903.5	301.4	596.9	940.3	
Land reductions share of total global emissions reductions	CH ₄	min	25 %	20 %	20 %	22 %	20 %	16 %
		max	37 %	40 %	42 %	30 %	31 %	36 %
	CO ₂	min	-43 %	-12 %	-4 %	-20 %	-8 %	-4 %
		max	74 %	48 %	17 %	73 %	47 %	15 %
	N ₂ O	min	52 %	61 %	65 %	53 %	61 %	65 %
		max	95 %	90 %	87 %	78 %	83 %	85 %
	Sum	min	-11 %	0 %	1 %	-2 %	1 %	1 %
		max	70 %	47 %	19 %	69 %	46 %	17 %
Percent of baseline land emissions reduced	CH ₄	min	3 %	8 %	10 %	0 %	2 %	10 %
		max	8 %	16 %	28 %	10 %	18 %	30 %
	CO ₂	min	-42 %	-89 %	0 %	-42 %	-104 %	0 %
		max	373 %	417 %	504 %	381 %	423 %	512 %
	N ₂ O	min	4 %	6 %	8 %	4 %	6 %	8 %
		max	10 %	16 %	22 %	12 %	17 %	22 %
	Sum	min	-4 %	1 %	7 %	-2 %	1 %	8 %
		max	97 %	100 %	73 %	99 %	101 %	76 %

for some models. Land-related N_2O and CH_4 reductions are a significant part of total N_2O and CH_4 reductions, but only a small fraction of baseline emissions, suggesting that models have cost-effective reasons to keep N_2O and CH_4 emissions. Emissions reductions from land increase only slightly with the stringency of the atmospheric concentration goal, as energy and industry emission reductions increase faster with target stringency. This result is consistent with previous studies (Rose et al., 2012). Land-based CO_2 reductions can be over 100% of baseline emissions, from the expansion of managed and unmanaged forests for sequestration.

Emissions reductions from individual land-related technologies, especially bioenergy, are not generally reported in transformation pathway studies. In part, this is due to emphasis on the energy system, but also other factors that make it difficult to uniquely quantify mitigation by technology. An exception is Rose et al. (2012) who reported agriculture, forest carbon, and bioenergy abatement levels for various atmospheric concentration goals. Cumulatively, over the century, bioenergy was the dominant strategy, followed by forestry, and then agriculture. Bioenergy cumulatively generated approximately 5 to 52 Gt CO_2 eq and 113 to 749 Gt CO_2 eq mitigation by 2050 and 2100, respectively. In total, land-related strategies contributed 20 to 60% of total cumulative abatement to 2030, 15 to 70% to 2050, and 15 to 40% to 2100.

Within models, there is a positive correlation between emissions reductions and GHG prices. However, across models, it is less clear, as some estimate large reductions with a low GHG price, while others estimate low reductions despite a high GHG price (Rose et al., 2012). For the most part, these divergent views are due to differences in model assumptions and are difficult to disentangle. Overall, while a tighter target and higher carbon price results in a decrease in land-use emissions, emissions decline at a decreasing rate. This is indicative of the rising relative cost of land mitigation, the increasing demand for bioenergy, and subsequent increasing need for overall energy system GHG abatement and energy consumption reductions. For additional discussion of land's potential role in transformation pathways, especially regarding physical land-use and bioenergy, see sections 6.3.2.4 and 6.3.5.

Models project increased deployment of, and dependence on, modern bioenergy (i.e., non-traditional bioenergy that is produced centrally to service communities rather than individual household production for heat and cooking), with some models projecting up to 95 EJ per year by 2030, and up to 245 EJ per year by 2050. Models universally project that the majority of agriculture and forestry mitigation, and bioenergy primary energy, will occur in developing and transitional economies (Section 6.3.5).

More recently, the literature has begun analyzing more realistic policy contexts. This work has identified a number of policy coordination and implementation issues. There are many dimensions to policy coordination: technologies, sectors, regions, climate and non-climate policies,

and timing. There are three prominent issues. First, there is coordination between mitigation activities. For instance, increased bioenergy incentives without global terrestrial carbon stock incentives or global forest protection policy, could result in substantial land conversion and emissions with large-scale deployment of energy crops. The projected emissions come primarily from the displacement of pasture, grassland, and natural forest (Sections 6.3.5 and 11.4.3). Energy cropland expansion also results in non-energy cropland conversion. These studies find that ignoring land conversion emissions with energy crop expansion, results in the need for deeper emissions reductions in the fossil and industrial sectors, and increased total mitigation costs. However, illustrative scenarios by (Calvin et al., 2013a) suggest that extensive forest protection policies may be needed for managing bioenergy driven deforestation. Note that providing energy crops, especially while protecting terrestrial carbon stocks, could result in a significant increase in food prices, potentially further exacerbated if also expanding forests (Wise et al., 2009; Popp et al., 2011; Reilly et al., 2012; Calvin et al., 2013a; see also Sections 11.4.3 and 11.13.7). In addition to competition between energy crops and forest carbon strategies, there is also competition between avoided deforestation and afforestation mitigation strategies, but synergies between forest management and afforestation (Rose and Sohngen, 2011). Bioenergy sustainability policies across sectors also need to be coordinated (Frank et al., 2013).

The second major concern is coordination of mitigation activity over time. The analyses noted in the previous paragraph assume the ability to globally protect or incentivize all, or a portion, of forest carbon stocks. A few studies to date have evaluated the implications of staggered forest carbon incentives—across regions and forest carbon activities. For instance, (Calvin et al., 2009) estimate land CO_2 emissions increases of 4 and 6 Gt CO_2 /yr in 2030 and 2050, respectively, from scenarios with staggered global regional climate policies that include forest carbon incentives. And, Rose and Sohngen (2011) find that fragmented or delayed forest carbon policy could accelerate deforestation. They project 60–100 Gt CO_2 of leakage by 2025 with a carbon price of 15 USD₂₀₁₀/t CO_2 that rises at 5% per year. Regional agriculture and forestry mitigation supply costs are also affected by regional participation/non-participation, with non-participating regions potentially increasing the mitigation costs for participating regions (Golub et al., 2009). Staggered adoption of land-mitigation policies will likely have institutional and socioeconomic implications as well (Madlener et al., 2006). Institutional issues, especially clarification of land tenure and property rights and equity issues (Section 11.7), will also be critical for successful land mitigation in forestry over time (Palmer, 2011; Gupta, 2012; Karsenty et al., 2014).

Finally, the type of incentive structure has implications. International land-related mitigation projects are currently regarded as high risk carbon market investments, which may affect market appeal. Also, mitigation scenarios assume that all emissions and sequestration changes are priced (similar to capping all emissions). However, mitigation, especially in agriculture and forestry, may be sought through volun-

tary markets, where mitigation suppliers choose whether to participate (Section 11.10). For instance, Rose et al. (2013) estimate reduced mitigation potential, as well as over-crediting, for United States agriculture and forestry with voluntary mitigation supply incentives, e.g., mitigation decreased 25–55 % at 15 USD₂₀₁₀/tCO₂eq due to non-participant leakage and non-additional crediting.

11.9.2 Implications of transformation pathways for the AFOLU sector

Transformation pathways indicate that a combination of forces can result in very different projected landscapes relative to today, even in baseline scenarios (Section 6.3.5). For instance, Popp et al. (2013) evaluate three models, and show that projected 2030 baseline changes from today alone vary sharply across models in all regions (Figure 11.19). See Section 6.3.5 for global land cover change results for a broader set of studies and policy contexts. In the examples in Figure 11.19, projections exhibit growth and reductions in both non-energy cropland (e.g., ASIA), and energy cropland (e.g., ASIA, OECD-1990, EIT). Furthermore, different kinds of land are converted when baseline cropland expands (e.g., MAF). Mitigation generally induces greater land cover changes than in baseline scenarios, but there are very different potential transformation visions. Overall, it is difficult to generalize on regional land cover effects of mitigation. For the same atmospheric concentration goal, some models convert significant area, some do not. There is energy cropland expansion in many regions that supports the production of bioenergy. Less consistent is the response of forest land, primarily due to differences in the land carbon options/policies modelled (Section 6.3.5). Finally, there is relatively modest additional land conversion in the 450 ppm, compared to the 550 ppm, scenarios, which is consistent with the declining role of land-related mitigation with policy stringency.

The implications of transformation pathway scenarios with large regional expansion of forest cover for carbon sequestration, depends in part on how the forest area increases (Figure 11.19; Popp et al., 2013). If forest areas increase through the expansion of natural vegetation, biodiversity and a range of other ecosystem services provided by forests could be enhanced. If afforestation occurs through large-scale plantation, however, some negative impacts on biodiversity, water, and other ecosystem services could arise, depending on what land cover the plantation replaces and the rotation time (Section 11.7). Similar issues arise with large-scale bioenergy, and environmental impacts of energy crop plantations, which largely depend upon where, how, and at what scale they are implemented, and how they are managed (Davis et al., 2013; see Section 11.13.6). Not surprisingly, the realistic policy coordination and implementation issues discussed in Section 11.9.1 could have significant land-use consequences, and additional policy design research is essential to better characterize mitigation costs, net emissions, and other social implications.

11.9.3 Implications of transformation pathways for sustainable development

The implications of the transformation pathways on sustainable development are context- and time-specific. A detailed discussion of the implications of large-scale LUC, competition between different demands for land, and the feedbacks between LUC and other services provided by land is provided in Section 11.4, potential co-benefits and adverse side-effects are discussed in Section 11.7, and Section 6.6 compares potential co-benefits and adverse side-effects across sectors, while Section 11.8 presents the opportunities and barriers for promoting AFOLU mitigation activities in the future. Finally, Section 11.13 discusses the specific implications of increasing bioenergy crops.

11.10 Sectoral policies

Climate change and different policy and management choices interact. The interrelations are particularly strong in agriculture and forestry: climate has a strong influence on these sectors that also constitute sources of GHG as well as sinks (Golub et al., 2009). The land provides a multitude of ecosystem services, climate change mitigation being just one of many services that are vital to human well-being. The nature of the sector means that there are, potentially, many barriers and opportunities as well as a wide range of potential impacts related to the implementation of AFOLU mitigation options (Sections 11.7 and 11.8). Successful mitigation policies need to consider how to address the multi-functionality of the sector. Furthermore, physical environmental limitations are central for the implementation of mitigation options and associated policies (Pretty, 2013). The cost-effectiveness of different measures is hampered by regional variability. National and international agricultural and forest climate policies have the potential to redefine the opportunity costs of international land-use in ways that either complement or hinder the attainment of climate change mitigation goals (Golub et al., 2009). Policy interactions could be synergistic (e.g., research and development investments and economic incentives for integrated production systems) or conflicting (e.g., policies promoting land conversion vs. conservation policies) across the sector (see Table 11.11). Additionally, adequate policies are needed to orient practices in agriculture and in forestry toward global sharing of innovative technologies for the efficient use of land resources to support effective mitigation options (see Table 11.2).

Forty-three countries in total (as of December 2010) have proposed NAMAs to the UNFCCC. Agriculture and forestry activities were considered as ways to reduce their GHG emissions in 59 and 94 % of the proposed NAMAs. For the least developed countries, the forestry sector was quoted in all the NAMAs, while the agricul-




Figure 11.19 | Regional land cover change by 2030 from 2005 from three models for baseline (left) and idealized policy implementation 550 ppm CO₂eq (centre) and 450 ppm CO₂eq (right) scenarios. (Popp et al., 2013).

tural sector was represented in 70% of the NAMAs (Bockel et al., 2010). Policies related to the AFOLU sector that affect mitigation are discussed below according to the instruments through which they may be implemented (economic incentives, regulatory and control approaches, information, communication and outreach, research and development). Economic incentives (e.g., special credit lines for low-carbon agriculture, sustainable agriculture and forestry practices, tradable credits, payment for ecosystem services) and regulatory approaches (e.g., enforcement of environmental law to reduce

deforestation, set-aside policies, air and water pollution control reducing nitrate load and N₂O emissions) have been effective in different cases. Investments in research, development, and diffusion (e.g., improved fertilizer use efficiency, livestock improvement, better forestry management practices) could result in positive and synergistic impacts for adaptation and mitigation (Section 11.5). Emphasis is given to REDD+, considering its development in recent years, and relevance for the discussion of mitigation policies in the forestry sector.

Table 11.11 | Some regional and global programs and partnerships related to illegal logging, forest management and conservation and REDD+.

Programme/Institution/Source	Context	Objectives and Strategies
Forest Law Enforcement and Governance (FLEG)/ World Bank/ www.worldbank.org/eapfleg	Illegal logging and lack of appropriate forest governance are major obstacle to countries to alleviate poverty, to develop their natural resources and to protect global and local environmental services and values	Support regional forest law enforcement and governance (FLEG)
Improving Forest Law Enforcement and Governance in the European Neighbourhood Policy East Countries and Russia (ENPI-FLEG)/EU/ www.enpi-fleg.org	Regional cooperation in the European Neighbourhood Policy Initiative East Countries (Armenia, Azerbaijan, Belarus, Georgia, Moldova, and Ukraine), and Russia following up on the St. Petersburg Declaration	Support governments, civil society, and the private sector in participating countries in the development of sound and sustainable forest management practices, including reducing the incidence of illegal forestry activities.
Forest Law Enforcement, Governance and Trade (FLEGT)/European Union/ www.euflegt.efi.int/	Illegal logging has a devastating impact on some of the world's most valuable forests. It can have not only serious environmental, but also economic and social consequences.	Exclude illegal timber from markets, to improve the supply of legal timber and to increase the demand for responsible wood products. Central elements are trade accords to ensure legal timber trade and support good forest governance in the partner countries. There are a number of countries in Africa, Asia, South and Central America currently negotiating FLEGT Voluntary Partnership Agreements (VPAs) with the European Union.
Program on Forests (PROFOR)/multiple donors including the European Union, European countries, Japan and the World Bank/ www.profor.info	Well-managed forests have the potential to reduce poverty, spur economic development, and contribute to a healthy local and global environment	Provide in-depth analysis and technical assistance on key forest questions related to livelihoods, governance, financing, and cross-sectoral issues. PROFOR activities comprise analytical and knowledge generating work that support the strategy's objectives of enhancing forests' contribution to poverty reduction, sustainable development and the protection of environmental services.
UN-REDD Programme/United Nations/ www.un-redd.org	The UN collaborative initiative on Reducing Emissions from Deforestation and forest Degradation (REDD) in developing countries was launched in 2008 and builds on the convening role and technical expertise of the FAO, UNDP, and the UNEP.	The Programme supports national REDD+ readiness efforts in 46 partner countries (Africa, Asia-Pacific, and Latin America) through (i) direct support to the design and implementation of REDD+ National Programmes; and (ii) complementary support to national REDD+ action (common approaches, analyses, methodologies, tools, data, and best practices).
REDD+ Partnership/International effort (50 different countries)/ www.reddpluspartnership.org	The UNFCCC has encouraged the Parties to coordinate their efforts to reduce emissions from deforestation and forest degradation. As a response, countries attending the March 2010 International Conference on the Major Forest Basins, hosted by the Government of France, agreed on the need to forge a strong international partnership on REDD+.	The REDD+ Partnership serves as an interim platform for its partner countries to scale up actions and finance for REDD+ initiatives in developing countries (including improving the effectiveness, efficiency, transparency, and coordination of REDD+ and financial instruments), to facilitate knowledge transfer, capacity enhancement, mitigation actions and technology development, and transfer among others.
Forest Investment Program (FIP)/Strategic Climate Fund (a multi-donor Trust Fund within the Climate Investment Funds) www.climateinvestmentfunds.org/cif/	Reduction of deforestation and forest degradation and promotion of sustainable forest management, leading to emission reductions and the protection of carbon terrestrial sinks.	Support developing countries' efforts to REDD and promote sustainable forest management by providing scaled-up financing to developing countries for readiness reforms and public and private investments, identified through national REDD readiness or equivalent strategies.
Forest Carbon Partnership (FCPF)/World Bank/ www.forestcarbonpartnership.org	Assistance to developing countries to implement REDD+ by providing value to standing forests.	Builds the capacity of developing countries to reduce emissions from deforestation and forest degradation and to tap into any future system of REDD+.
Indonesia-Australia Forest Carbon Partnership/ www.iafcpc.or.id	Australia's assistance on climate change and builds on long-term practical cooperation between Indonesia and Australia.	The Partnership supports strategic policy dialogue on climate change, the development of Indonesia's National Carbon Accounting System, and implementing demonstration activities in Central Kalimantan.

11.10.1 Economic incentives

Emissions trading: Carbon markets occur under both compliance schemes and as voluntary programmes. A review of existing offset programmes was provided by Kollmuss et al. (2010). More details are also presented in Section 15.5.3. Compliance markets (Kyoto offset mechanisms, mandatory cap-and-trade systems, and other mandatory GHG systems) are created and regulated by mandatory national, regional, or international carbon reduction regimes (Kollmuss et al., 2010). The three Kyoto Protocol mechanisms are very important for the regulatory market: CDM, Joint Implementation (JI) and the Emissions Trading System (ETS). Currently, AFOLU projects in CDM only include specific types of projects: for agriculture—methane avoid-

ance (manure management), biogas projects, agricultural residues for biomass energy; for forestry—reforestation and afforestation. By June 2013, the total number of registered CDM projects was 6989, 0.6 and 2.5 % of this total being related to afforestation/reforestation and agriculture, respectively (UNFCCC—CDM); therefore, finance streams coming from A/R CDM Projects are marginal from the global perspective. An analysis of A/R CDM projects suggests crucial factors for the performance of these projects are initial funding support, design, and implementation guided by large organizations with technical expertise, occurrence on private land (land with secured property rights attached), and that most revenue from Certified Emission Reductions (CERs) is directed back to local communities (Thomas et al., 2010).

There are compliance schemes outside the scope of the Kyoto Protocol, but these are carried out exclusively at the national level, with no relation to the Protocol. In 2011, Australia started the Carbon Farming Initiative (CFI) that allows farmers and investors to generate tradable carbon offsets from farmland and forestry projects. This followed several years of state-based and voluntary activity that resulted in 65,000 ha of A/R projects (Mitchell et al., 2012). Another example is The Western Arnhem Land Fire Abatement Project (WALFA), a fire management project in Australia initiated in 2006 that produces a tradable carbon offset through the application of improved fire management using traditional management practices of indigenous land owners (Whitehead et al., 2008; Bradstock et al., 2012). Alberta's offset credit system is a compliance mechanism for entities regulated under the province's mandatory GHG emission intensity-based regulatory system (Kollmuss et al., 2010). In the case of N₂O emissions from agriculture, the Alberta Quantification Protocol for Agricultural N₂O Emissions Reductions issues C offset credits for on-farm reductions of N₂O emissions and fuel use associated with the management of fertilizer, manure, and crop residues for each crop type grown. Other N₂O emission reduction protocols (e.g., Millar et al., 2010) are being considered for the Verified Carbon Standard, the American Carbon Registry, and the Climate Action Reserve (Robertson et al., 2013).

Agriculture and Forestry activities are not covered by the European Union Emissions Trading Scheme (EU ETS), which is by far the largest existing carbon market. Forestry entered the New Zealand Kyoto Protocol compliant ETS in 2008, and mandatory reporting for agriculture began in 2012, although full entry of agriculture into the scheme has been delayed indefinitely. Agricultural participants include meat processors, dairy processors, nitrogen fertilizer manufacturers and importers, and live animal exporters, although some exemptions apply (Government of New Zealand). California's Cap-and-Trade Regulation took effect on January 1, 2012, with amendments to the Regulation effective September 1, 2012. The enforceable compliance obligation began on January 1, 2013. Four types of projects were approved as eligible to generate carbon credits to regulated emitters in California: avoidance of methane emissions from installation of anaerobic digesters on farms, carbon sequestration in urban and rural forestry, and destruction of ozone depleting substances (California Environmental Protection Agency).

Voluntary carbon markets operate outside of the compliance markets. By enabling businesses, governments, non-governmental organizations (NGOs), and individuals to purchase offsets that were created either in the voluntary market or through the CDM, they can offset their emissions (Verified or Voluntary Emissions Reductions (VERs)). The voluntary offset market includes a wide range of programmes, entities, standards, and protocols (e.g., Community & Biodiversity Standards, Gold Standard, Plan Vivo among others) to improve the quality and credibility of voluntary offsets. The most common incentives for the quantity buyers of carbon credits in the private sector are corporate social responsibility and public relations. Forest projects are increasing in the voluntary markets. Transactions of carbon credits from this

sector totalled 133 million USD in 2010, 95% of them in voluntary markets (Peters-Stanley et al., 2011).

Reducing emissions from deforestation; reducing emissions from forest degradation; conservation of forest carbon stocks; sustainable management of forests; and enhancement of forest carbon stocks (REDD+): REDD+ consists of forest-related activities implemented voluntarily by developing countries that may, in isolation or jointly lead to significant climate change mitigation¹⁰. REDD+ was introduced in the agenda of the UNFCCC in 2005, and has since evolved to an improved understanding of the potential positive and negative impacts, methodological issues, safeguards, and financial aspects associated with REDD+ implementation. Here, we first address the REDD+ discussions under the UNFCCC, but also introduce other REDD+-related initiatives. The novel aspects of REDD+ under the Convention, relative to previous forest-related mitigation efforts by developing countries under the UNFCCC are its national and broader coverage, in contrast to project-based mitigation activities¹¹ (e.g., under the CDM of the Kyoto Protocol). Its main innovation is its results-based approach, in which payments are done *ex post* in relation to a mitigation outcome already achieved, as opposed to project-based activities, where financing is provided *ex ante* in relation to expected outcomes. A phased approach to REDD+ was agreed at the UNFCCC, building from the development of national strategies or action plans, policies and measures, and evolving into results-based actions that should be fully measured, reported, and verified—MRV (UNFCCC Dec. 1/16). REDD+ payments are expected for results-based actions, and although the UNFCCC has already identified potential ways to pay for these¹², the financing architecture for the REDD+ mechanism is still under negotiation under the UNFCCC.

Meanwhile, and as a result to the explicit request from the UNFCCC for early actions in REDD+, different regional and global programmes and partnerships address forest management and conservation and readiness for REDD+ (Table 11.11), while some REDD+ strategies have started in countries with significant forest cover (see Box 11.7 for examples). Initiatives include multilateral activities (e.g., UN-REDD

¹⁰ Decision 1/CP.16 (FCCC/CP/2010/7/Add.1, paragraph 70) "Encourages developing countries to contribute to mitigation actions in the forest sector by undertaking the following activities, as deemed appropriate by each Party and in accordance with their respective capabilities and national circumstances—reducing emissions from deforestation; reducing emissions from forest degradation; conservation of forest carbon stocks; sustainable management of forests; and enhancement of forest carbon stocks".

¹¹ Decision 1/CP.16 (FCCC/CP/2010/7/Add.1, paragraph 73) "Decides that the activities undertaken by Parties referred to in paragraph 70 above should be implemented in phases, beginning with the development of national strategies or action plans, policies and measures, and capacity-building, followed by the implementation of national policies and measures and national strategies or action plans that could involve further capacity-building, technology development and transfer and results-based demonstration activities, and evolving into results-based actions that should be fully measured, reported and verified".

¹² Decision 2/CP.17 (FCCC/CP/2011/9/Add.1, paragraph 65) "Agrees that results-based finance provided to developing country Parties that is new, additional and predictable may come from a wide variety of sources, public and private, bilateral and multilateral, including alternative sources".

Programme, Forest Carbon Partnership Facility, Forest Investment Program), bilateral activities (e.g., Tanzania-Norway, Indonesia-Norway), country driven initiatives (in addition to 16 UN-REDD Programme countries, the Programme also supports 31 other partner countries across Africa, Asia-Pacific, and Latin America and the Caribbean; UN-REDD Programme—Support to Partner Countries).

REDD+ can be a very cost-effective option for mitigating climate change and could supply a large share of global abatement of emissions from the AFOLU sector from the extensive margin of forestry, especially through reducing deforestation in tropical regions (Golub et al., 2009). Issues of concern for REDD+ implementation have been captured under REDD+ safeguards in line with the UNFCCC Cancun Agreement. To respond to the requirements outlined in the UNFCCC agreement, a number of steps need to be considered in the development of country-level safeguard information systems for REDD+ including defining social and environmental objectives, assessing potential benefits and risks from REDD+, assessing current safeguard systems, drafting a strategic plan or policy, and establishing a governance system.

A growing body of literature has analyzed different aspects related to the implementation, effectiveness, and scale of REDD+, as well as the interactions with other social and environmental co-benefits (e.g., Angelsen et al., 2008; Levin et al., 2008; Larson, 2011; Gardner et al., 2012). Results-based REDD+ actions, which are entitled to results-based finance, require internationally agreed rules for MRV. Measuring and monitoring the results will most likely rely on a combination of remotely-sensed data with ground-based inventories. The design of a REDD policy framework (and specifically its rules) can have a significant

impact on monitoring costs (Angelsen et al., 2008; Böttcher et al., 2009). Forest governance is another central aspect in recent studies, including debate on decentralization of forest management, logging concessions in public-owned commercially valuable forests, and timber certification, primarily in temperate forests (Agrawal et al., 2008). Although the majority of forests continue to be formally owned by governments, there are indications that the effectiveness of forest governance is increasingly independent of formal ownership (Agrawal et al., 2008). However, there are widespread concerns that REDD+ will increase costs on forest-dependent peoples and in this context, stakeholder rights, including rights to continue sustainable traditional land-use practices, appear as a precondition for REDD development (Phelps et al., 2010b).

Some studies have addressed the potential displacement of emissions, i.e., a reduction of emissions in one place resulting in an increase of emissions elsewhere (or leakage) (Santilli et al., 2005; Forner et al., 2006; Nabuurs et al., 2007; Strassburg et al., 2008, 2009; Section 11.3.2). The national coverage of REDD+ might ameliorate the issue of emissions displacement, a major drawback of project-based approaches (Herold and Skutsch, 2011). To minimize transnational displacement of emissions, REDD+ needs to stimulate the largest number of developing countries to engage voluntarily. There are also concerns about the impacts of REDD+ design and implementation options on biodiversity conservation, as areas of high C content and high biodiversity are not necessarily coincident. Some aspects of REDD+ implementation that might affect biodiversity include site selection, management strategies, and stakeholder engagement (Harvey et al., 2010). From a conservation biology perspective, it is also relevant where the displacement occurs, as deforestation and exploitation of natural

Box 11.7 | Examples of REDD+ initiatives at national scale in different regions with significant extension of forest cover

Amazon Fund: The Amazon Fund in Brazil was officially created in 2008 by a presidential decree. The Brazilian Development Bank (BNDES) was given the responsibility of managing it. The Norwegian government played a key role in creating the fund by donating funds to the initiative in 2009. Since then, the Amazon Fund has received funds from two more donors: the Federal Republic of Germany and Petrobrás, Brazil's largest oil company. As of February 2013, 1.03 billion USD has been pledged, with 227 million USD approved for activities (Amazon Fund).

UN-REDD Democratic Republic of Congo: The Congo Basin rainforests are the second largest after Amazonia. In 2009, Democratic Republic of the Congo (DRC), with support of UN-REDD Programme and Forest Carbon Partnership Facility (FCPC), started planning the implementation stages of REDD+ readiness. The initial DRC National Programme transitioned into the full National Programme (Readiness Plan) after it was approved by

the UN-REDD Programme Policy Board in 2010 (UN-REDD Programme). The budget comprises 5.5 million USD₂₀₁₀ and timeframe is 2010–2013.

Indonesia-Norway REDD+ Partnership: In 2010, the Indonesia-Norway REDD+ Partnership was established through an agreement between governments of the two countries. The objective was to 'support Indonesia's efforts to reduce emissions from deforestation and degradation of forests and peatlands. Indonesia agreed to take systematic and decisive action to reduce its forest and peat-related GHG emissions, whereas Norway agreed to support those efforts by making available up to 1 billion USD₂₀₁₀, exclusively on a payment-for-results basis over the next few years' (UN-REDD Programme). In 2013, Indonesia's government has extended the moratorium on new forest concessions for a further two years, protecting an additional 14.5 Mha of forest.

resources could move from areas of low conservation value to those of higher conservation value, or to other natural ecosystems, threatening species native to these ecosystems (Harvey et al., 2010). Additionally, transnational displacement could cause deforestation to move into relatively intact areas of high biodiversity value, or into countries that currently have little deforestation (Putz and Redford, 2009).

Taxes, charges, subsidies: Financial regulations are another approach to pollution control. A range of instruments can be used: pollution charges, taxes on emission, taxes on inputs, and subsidies (Jakobsson et al., 2002). Nitrogen taxes are one possible instrument, since agricultural emissions of N₂O mainly derive from the use of nitrogenous fertilizers. An analysis of the tax on the nitrogen content of synthetic fertilizers in Sweden indicated that direct N₂O emissions from agricultural soils in Sweden (the tax abolished in 2010) would have been on average 160 tons or 2% higher without the tax (Mohlin, 2013). Additionally, the study showed that removal of the N tax could completely counteract the decreases in CO₂ emissions expected from the future tax increase on agricultural CO₂. The mitigation potential of GHG-weighted consumption taxes on animal food products was estimated for the EU using a model of food consumption (Wirsenius et al., 2011). A 7% reduction of current GHG emission in European Union (EU) agriculture was estimated with a GHG-weighted tax on animal food products of 79 USD₂₀₁₀/tCO₂eq (60 EUR₂₀₁₀/tCO₂eq). Low-interest loans can also support the transition to sustainable agricultural practices as currently implemented in Brazil, the second largest food exporter, through the national programme (launched in 2010; Plano ABC).

11.10.2 Regulatory and control approaches

Deforestation control and land planning (protected areas and land sparing/set-aside policies): The rate of deforestation in the tropics and relative contribution to anthropogenic carbon emissions has been declining (Houghton, 2012; see Section 11.2 for details). Public policies have had a significant impact by reducing deforestation rates in some tropical countries (see, e.g., Box 11.8).

Since agricultural expansion is one of the drivers of deforestation (especially in tropical regions), one central question is if intensification of agriculture reduces cultivated areas and results in land sparing by concentrating production on other land. Land sparing would allow released lands to sequester carbon, provide other environmental services, and protect biodiversity (Fischer et al., 2008). In the United States, over 13 Mha of former cropland are enrolled in the US Conservation Reserve Program (CRP), with biodiversity, water quality, and carbon sequestration benefits (Gelfand et al., 2011). In 1999, China launched the Grain for Green Program or Sloping Land Conversion Program as a national measure to increase vegetation cover and reduce erosion. Cropland and barren land were targeted and over 20 Mha of land were converted into mostly tree-based plantations. Over its first 10 years between ~800 to 1700 MtCO₂eq (Moberg, 2011) were sequestered.

Environmental regulation (GHG and their precursors emissions control): In many developed countries, environmental concerns related to water and air pollution since the mid-1990s led to the adoption of laws and regulations that now mandate improved agricultural nutrient management planning (Jakobsson et al., 2002). Some policy initiatives deal indirectly with N leakages and thus promote the reduction of N₂O emissions. The EU Nitrates Directive (1991) sets limits on the use of fertilizer N and animal manure N in nitrate-vulnerable zones. Across the 27 EU Member States, 39.6% of territory is subject to related action programmes. However, in terms of the effectiveness of environmental policies and agriculture, there has been considerable progress in controlling point pollution, but efforts to control non-point pollution of nutrients have been less successful, and potential synergies from various soil-management strategies could be better exploited. Emission targets for the AFOLU sector were also introduced by different countries (e.g., Climate Change Acts in UK and Scotland; European Union).

Bioenergy targets: Many countries worldwide, by 2012, have set targets or mandates or both for bioenergy, to deliver to multiple policy objectives, such as climate change mitigation, energy security, and rural development. The bulk of mandates continue to come from the EU-27 but 13 countries in the Americas, 12 in Asia-Pacific, and 8 in Africa have mandates or targets in place (Petersen, 2008; www.biofuelsdigest.com). For the sustainability of biofuels implementation, land-use planning and governance are central (Tilman et al., 2009), as related policy and legislation, e.g., in agriculture, forestry, environment and trade, can strongly influence the development of bioenergy programmes (Jull et al., 2007). A recent study analyzed the consequences of renewable targets of EU member states on the CO₂ sink of EU forests, and indicated a decrease in the forest sink by 4–11% (Böttcher et al., 2012). Another possible tradeoff of biofuel targets is related to international trade. Global trade in biofuels might have a major impact on other commodity markets (e.g., vegetable oils or animal fodder) and has already caused a number of trade disputes, because of subsidies and non-tariff barriers (Oosterveer and Mol, 2010).

Box 11.8 | Deforestation control in Brazil

The Brazilian Action Plan for the Prevention and Control of Deforestation in the Legal Amazon (PPCDAm) includes coordinated efforts among federal, state, and municipal governments, and civil organizations, remote-sensing monitoring, significant increase of new protected areas (Soares-Filho et al., 2010), and combination of economic and regulatory approaches. For example, since 2008 federal government imposed sanctions to municipalities with very high deforestation rates, subsidies were cut and new credit policies made rural credit dependent on compliance with environmental legislation (Macedo et al., 2012; Nolte et al., 2013).

11.10.3 Information schemes

Acceptability by land managers and practicability of mitigation measures (Table 11.2) need to be considered, because the efficiency of a policy is determined by the cost of achieving a given goal (Sections 11.4.5; 11.7). Therefore, costs related to education and communication of policies should be taken into account (Jakobsson et al., 2002). Organizations created to foster the use of science in environmental policy, management, and education can facilitate the flow of information from science to society, increasing awareness of environmental problems (Osmond et al., 2010). In the agriculture sector, non-profit conservation organizations (e.g., The Sustainable Agriculture Network (SAN)) and governments (e.g., Farming for a Better Climate, Scotland) promote the social and environmental sustainability of activities by developing standards and educational campaigns.

Certification schemes also support sustainable agricultural practices (Sections 11.4.5; 11.7). Climate-friendly criteria reinforce existing certification criteria and provide additional value. Different certification systems also consider improvements in forest management, reduced deforestation and carbon uptake by regrowth, reforestation, agroforestry, and sustainable agriculture. In the last 20 years, forest certification has been developed as an instrument for promoting sustainable forest management. Certification schemes encompass all forest types, but there is a concentration in temperate forests (Durst et al., 2006). Approximately 8% of global forest area has been certified under a variety of schemes and 25% of global industrial roundwood comes from certified forests (FAO, 2009b). Less than 2% of forest area in African, Asian, and tropical American forests are certified, and most certified forests (82%) are large and managed by the private sector (ITTO, 2008). In the forestry sector, many governments have worked towards a common understanding of sustainable forest management (Auld et al., 2008). Certification bodies certify that farms or groups comply with standards and policies (e.g., Rainforest Alliance Certified). In some, specific voluntary climate change adaptation and mitigation criteria are included.

Forest certification as an instrument to promote sustainable forest management (SFM) and biodiversity maintenance was evaluated by (Rametsteiner and Simula, 2003) they indicated that standards used for issuing certificates upon compliance are diverse, but often include elements that set higher than minimum standards.

Further, independent audits are an incentive for improving forest management. In spite of many difficulties, forest certification was considered successful in raising awareness, disseminating knowledge on the SFM concept worldwide, and providing a tool for a range of applications other than the assessment of sustainability, e.g., verifying carbon sinks. Another evaluation of certification schemes for conserving biodiversity (Harvey et al., 2008) indicated some constraints that probably also apply to climate-friendly certification: weakness of compliance or enforcement of standards, transaction costs and paperwork often limit participation, and incentives are insufficient to

attract high levels of participation. Biofuel certification is a specific case as there are multiple actors and several successive segments of biofuel production pathways: feedstock production, conversion of the feedstock to biofuels, wholesale trade, retail, and use of biofuels in engines (Gnansounou, 2011). Because of the length and the complexity of biofuel supply chains assessing sustainability is challenging (Kaphengst et al., 2009).

11.10.4 Voluntary actions and agreements

Innovative agricultural practices and technologies can play a central role in climate change mitigation and adaptation, with policy and institutional changes needed to encourage the innovation and diffusion of these practices and technologies to developing countries. Under the UNFCCC, the 2007 Bali Action Plan identified technology development and transfer as a priority area. A Technology Mechanism was established by Parties at the COP16 in 2010 "to facilitate the implementation of enhanced action on technology development and transfer, to support action on mitigation and adaptation, in order to achieve the full implementation of the Convention" (UNFCCC). For agriculture, Burney et al., (2010) indicated that investment in yield improvements compared favourably with other commonly proposed mitigation strategies.

Additionally, adaptation measures in agriculture can also generate significant mitigation effects. Lobell et al. (2013) investigated the co-benefits of adaptation measures on farm level that reduced GHG emissions from LUC. The study focused on investments in research for developing and deploying new technologies (e.g., disease-resistant or drought-tolerant crops, or soil-management techniques). It concluded that broad-based efforts to adapt agriculture to climate change have mitigation co-benefits that are associated with lower costs than many activities focusing on mitigation, especially in developed countries.

11.11 Gaps in knowledge and data

Data and knowledge gaps include:

- Improved global high-resolution data sets of crop production systems (including crop rotations, variety selection, fertilization practices, and tillage practices), grazing areas (including quality, intensity of use, management), and freshwater fisheries and aquaculture, also comprising subsistence farming.
- Globally standardized and homogenized data on soil as well as forest degradation and a better understanding of the effects of degradation on carbon balances and productivity.

- Improved understanding of the mitigation potential, interplay, and costs as well as environmental and socio-economic consequences of land use-based mitigation options such as improved agricultural management, forest conservation, bioenergy production, and afforestation on the national, regional, and global scale.
- Better understanding of the effect of changes in climate parameters, rising CO₂ concentrations and N deposition on productivity and carbon stocks of different types of ecosystems, and the related consequences for land-based climate change mitigation potentials.

11.12 Frequently Asked Questions

FAQ 11.1 How much does AFOLU contribute to GHG emissions and how is this changing?

Agriculture and land-use change, mainly deforestation of tropical forests, contribute greatly to anthropogenic greenhouse gas emissions and are expected to remain important during the 21st century. Annual GHG emissions (mainly CH₄ and N₂O) from agricultural production in 2000–2010 were estimated at 5.0–5.8 GtCO₂eq/yr, comprising about 10–12% of global anthropogenic emissions. Annual GHG flux from land use and land-use change activities accounted for approximately 4.3–5.5 GtCO₂eq/yr, or about 9–11% of total anthropogenic greenhouse gas emissions. The total contribution of the AFOLU sector to anthropogenic emissions is therefore around one quarter of the global anthropogenic total.

FAQ 11.2 How will mitigation actions in AFOLU affect GHG emissions over different timescales?

There are many mitigation options in the AFOLU sector that are already being implemented, e.g., afforestation, reducing deforestation, cropland and grazing land management, fire management, and improved livestock breeds and diets. These can be implemented now. Others (such as some forms of biotechnology and livestock dietary additives) are still in development and may not be applicable for a number of years. In terms of the mode of action of the options, in common with other sectors, non-CO₂ greenhouse gas emission reduction is immediate and permanent. However, a large portion of the mitigation potential in the AFOLU sector is carbon sequestration in soils and vegetation. This mitigation potential differs, in that the options are time-limited (the potential saturates), and the enhanced carbon stocks created are reversible and non-permanent. There is, therefore, a significant time

component in the realization and the duration of much of the mitigation potential available in the AFOLU sector.

FAQ 11.3 What is the potential of the main mitigation options in AFOLU for reducing GHG emissions?

In general, available top-down estimates of costs and potentials suggest that AFOLU mitigation will be an important part of a global cost-effective abatement strategy. However, potentials and costs of these mitigation options differ greatly by activity, regions, system boundaries, and the time horizon. Especially, forestry mitigation options—including reduced deforestation, forest management, afforestation, and agro-forestry—are estimated to contribute 0.2–13.8 GtCO₂/yr of economically viable abatement in 2030 at carbon prices up to 100 USD/tCO₂eq. Global economic mitigation potentials in agriculture in 2030 are estimated to be up to 0.5–10.6 GtCO₂eq/yr. Besides supply-side-based mitigation, demand-side mitigation options can have a significant impact on GHG emissions from food production. Changes in diet towards plant-based and hence less GHG-intensive food can result in GHG emission savings of 0.7–7.3 GtCO₂eq/yr in 2050, depending on which GHGs and diets are considered. Reducing food losses and waste in the supply chain from harvest to consumption can reduce GHG emissions by 0.6–6.0 GtCO₂eq/yr.

FAQ 11.4 Are there any co-benefits associated with mitigation actions in AFOLU?

In several cases, the implementation of AFOLU mitigation measures may result in an improvement in land management and therefore have socio-economic, health, and environmental benefits: For example, reducing deforestation, reforestation, and afforestation can improve local climatic conditions, water quality, biodiversity conservation, and help to restore degraded or abandoned land. Soil management to increase soil carbon sequestration may also reduce the amount of wind and water erosion due to an increase in surface cover. Further considerations on economic co-benefits are related to the access to carbon payments either within or outside the UNFCCC agreements and new income opportunities especially in developing countries (particularly for labour-intensive mitigation options such as afforestation).

FAQ 11.5 What are the barriers to reducing emissions in AFOLU and how can these be overcome?

There are many barriers to emission reduction. Firstly, mitigation practices may not be implemented for economic reasons (e.g., market failures, need for capital investment to realize recurrent savings), or a

range of factors including risk-related, political/bureaucratic, logistical, and educational/societal barriers. Technological barriers can be overcome by research and development; logistical and political/bureaucratic barriers can be overcome by better governance and institutions; education barriers can be overcome through better education and extension networks; and risk-related barriers can be overcome, for example, through clarification of land tenure uncertainties.

11.13 Appendix Bioenergy: Climate effects, mitigation options, potential and sustainability implications

11.13.1 Introduction

SRREN (IPCC, 2011) provided a comprehensive overview on bioenergy (Chum et al., 2011). However, a specific bioenergy Appendix in the context of the WGIII AR5 contribution is necessary because (1) many of the more stringent mitigation scenarios (resulting in 450 ppm, but also 550 ppm CO₂eq concentration by 2100, see Section 11.9.1) heavily rely on a large-scale deployment of bioenergy with carbon dioxide capture and storage (BECCS); (2) there has been a large body of literature published since SRREN, which complements and updates the analysis presented in this last report; (3) bioenergy is important for many chapters (Chapters 6; 7; 8; 10; 11), which makes it more useful to treat it in a single section instead of in many scattered chapter sections throughout the report. Chapter 11 is the appropriate location for the Appendix, as bioenergy analysis relies crucially on land-use assessments.

Bioenergy is energy derived from biomass, which can be deployed as solid, liquid, and gaseous fuels for a wide range of uses, including transport, heating, electricity production, and cooking (Chum et al., 2011). Bioenergy has a significant mitigation potential, but there are issues to consider, such as the sustainability of practices and the efficiency of bioenergy systems (Chum et al., 2011). Bioenergy systems can cause both positive and negative effects and their deployment needs to balance a range of environmental, social, and economic objectives that are not always fully compatible. The consequences of bioenergy implementation depend on (1) the technology used; (2) the location, scales, and pace of implementation; (3) the land category used (forest, grassland, marginal lands, and crop lands); and (4) the business models and practices adopted—including how these integrate with or displace the existing land use.

As an update to the SRREN, this report presents (1) a more fine-grained assessment of the technical bioenergy potential reflecting diverse per-

spectives in the literature; (2) recent potential estimates on technological solutions such as BECCS; (3) an in-depth description of different lifecycle emission accounting methods and their results; (4) a small increase in uncertainty on the future economic bioenergy potential; (5) a comprehensive assessment of diverse livelihood and sustainability effects of bioenergy deployment, identifying the need for systematic aggregation.

11.13.2 Technical bioenergy potential

The technical bioenergy potential, also known as the technical primary biomass potential for bioenergy, is the amount of the theoretical bioenergy output obtainable by full implementation of demonstrated technologies or practices (IPCC, 2011). Unfortunately there is no standard methodology to estimate the technical bioenergy potential, which leads to diverging estimates. Most of the recent studies estimating technical bioenergy potentials assume a 'food/fibre first principle' and exclude deforestation, eventually resulting in an estimate of the 'environmentally sustainable bioenergy potential' when a comprehensive range of environmental constraints is considered (Batidzirai et al., 2012).

Recently published estimates that are based in this extended definition of global technical bioenergy potentials in 2050 span a range of almost three orders of magnitude, from < 50 EJ/yr to > 1,000 EJ/yr (Smeets et al., 2007; Field et al., 2008; Haberl et al., 2010; Batidzirai et al., 2012). For example, Chum et al. reported global technical bioenergy potentials of 50–500 EJ/yr for the year 2050 (IPCC, 2011), and the Global Energy Assessment gave a range of 160–270 EJ/yr (Johansson et al., 2012). The discussion following the publication of these global reports has not resulted in a consensus on the magnitude of the future global technical bioenergy potential, but has helped to better understand some of its many structural determinants (Wirsenius et al., 2011; Berndes, 2012; Erb et al., 2012a). How much biomass for energy is technically available in the future depends on the evolution of a multitude of social, political, and economic factors, e.g., land tenure and regulation, trade, and technology (Dornburg et al., 2010).

Figure 11.20 shows estimates of the global technical bioenergy potential in 2050 by resource categories. Ranges were obtained from assessing a large number of studies based on a food/fibre first principle and various restrictions regarding resource limitations and environmental concerns but no explicit cost considerations (Hoogwijk et al., 2005; Smeets et al., 2007; Smeets and Faaij, 2007; van Vuuren et al., 2009; Hakala et al., 2009; Dornburg et al., 2010; Haberl et al., 2010, 2011a; Gregg and Smith, 2010; Chum et al., 2011; GEA, 2012; Rogner et al., 2012). Most studies agree that the technical bioenergy potential in 2050 is at least approximately 100 EJ/yr with some modelling assumptions leading to estimates exceeding 500 EJ/yr (Smeets et al., 2007). As stated, different views about sustainability and socio-ecological constraints lead to very different estimates, with some studies reporting much lower figures.

As shown in Figure 11.20, the total technical bioenergy potential is composed of several resource categories that differ in terms of their absolute potential, the span of the ranges—which also reflect the relative agreement/disagreement in the literature—and the implications of utilizing them. Regional differences—which are not addressed here—are also important as the relative size of each biomass resource within the total potential and its absolute magnitude vary widely across countries and world regions.

Forest and Agriculture residues. Forest residues (Smeets and Faaij, 2007; Smeets et al., 2007; Dornburg et al., 2010; Haberl et al., 2010; Gregg and Smith, 2010; Rogner et al., 2012) include residues from silvicultural thinning and logging; wood processing residues such as sawdust, bark, and black liquor; and dead wood from natural disturbances, such as storms and insect outbreaks (irregular source). The use of these resources is in general beneficial and any adverse side-effects can be mitigated by controlling residue removal rates considering biodiversity, climate, topography, and soil factors. There is a near-term tradeoff, particularly within temperate and boreal regions, in that organic matter retains organic C for longer if residues are left to decompose slowly instead of being used for energy. Agricultural residues (Smeets et al., 2007; Hakala et al., 2009; Haberl et al., 2010, 2011a; Gregg and Smith, 2010; Chum et al., 2011; Rogner et al., 2012) include manure, harvest residues (e.g., straw), and processing residues (e.g., rice husks from rice milling) and are also in general beneficial. However, mitigating

potential adverse side-effects—such as the loss of soil C—associated to harvesting agriculture residues is more complex as they depend on the different crops, climate, and soil conditions (Kochsiek and Knops, 2012; Repo et al., 2012). Alternative uses of residues (bedding, use as fertilizer) need to be considered. Residues have varying collection and processing costs (in both agriculture and forestry) depending on residue quality and dispersal, with secondary residues often having the benefits of not being dispersed and having relatively constant quality. Densification and storage technologies would enable cost-effective collections over larger areas. Optimization of crop rotation for food and bioenergy output and the use of residues in biogas plants may result in higher bioenergy yields from residues without food-energy competition.

Optimal forest harvesting is defined as the fraction of sustainable harvest levels (often set equal to net annual increment) in forests available for wood extraction, which is additional to the projected biomass demand for producing other forest products. This includes both biomass suitable for other uses (e.g., pulp and paper production) and biomass that is not used commercially (Smeets and Faaij, 2007; Chum et al., 2011). The resource potential depends on both environmental and socio-economic factors. For example, the change in forest management and harvesting regimes due to bioenergy demand depends on forest ownership and the structure of the associated forest industry. Also, the forest productivity—and C stock—response to changes in forest management and harvesting depends on the character of the forest ecosystem, as shaped by historic forest management and events such as fires, storms, and insect outbreaks, but also on the management scheme (e.g., including replanting after harvest, soil protection, recycling of nutrients, and soil types (Jonker et al., 2013; Lamers et al., 2013). In particular, optimizing forest management for mitigation is a complex issue with many uncertainties and still subject to scientific debate. Intensive forest management activities of the early- to mid-twentieth century as well as other factors such as recovery from past overuse, have led to strong forest C sinks in many OECD regions (Pan et al., 2011; Loudermilk et al., 2013; Nabuurs et al., 2013; Erb et al., 2013). However, the capacity of these sinks is being reduced as forests approach saturation (Smith, 2005; Körner, 2006; Guldea et al., 2008; Nabuurs et al., 2013; Sections 11.2.3, 11.3.2). Active forest management, including management for bioenergy, is therefore important for sustaining the strength of the forest carbon sink well into the future (Nabuurs et al., 2007, 2013; Canadell and Raupach, 2008; Ciais et al., 2008), although countries should realize that for some old forest areas, conserving carbon stocks may be preferential, and that the actively managed forests may for some time (decades) act as sources.




Figure 11.20 | Global Technical Bioenergy Potential by main resource category for the year 2050 | The figure shows the ranges in the estimates by major resource category of the global technical bioenergy potential. The color grading is intended to show qualitatively the degree of agreement in the estimates, from blue (large agreement in the literature) to purple (medium agreement) to red (small agreement). In addition, reducing traditional biomass demand by increasing its use efficiency could release the saved biomass for other energy purposes with large benefits from a sustainable development perspective.

Organic wastes include waste from households and restaurants, discarded wood products such as paper, construction, and demolition wood waste, and waste waters suitable for anaerobic biogas production (Haberl et al., 2010; Gregg and Smith, 2010). Organic waste may be dispersed and also heterogeneous in quality but the health and environmental gains from collection and proper management through




Figure 11.21 | Production pathways to liquid and gaseous fuels from biomass and, for comparison from fossil fuels (adapted from GEA, 2012; Turkenburg et al., 2012).

combustion or anaerobic digestion can be significant. Competition with alternative uses of the wastes may limit this resource potential.

Dedicated biomass plantations include annual (cereals, oil, and sugar crops) and perennial plants (e.g., switchgrass, *Miscanthus*) and tree plantations (both coppice and single-stem plantations (e.g., willow, poplar, eucalyptus, pine; (Hoogwijk et al., 2005, 2009; Smeets et al., 2007; van Vuuren et al., 2009; Dornburg et al., 2010; Wicke et al., 2011b; Haberl et al., 2011a). The range of estimates of technical bioenergy potentials from that resource in 2050 is particularly large (< 50 to > 500 EJ/yr). Technical bioenergy potentials from dedicated biomass plantations are generally calculated by multiplying (1) the area deemed available for energy crops by (2) the yield per unit area and year (Batidzirai et al., 2012; Coelho et al., 2012). Some studies have identified a sizable technical potential (up to 100 EJ) for bioenergy production using marginal and degraded lands (e.g., saline land) that are currently not in use for food production or grazing (Nijsen et al., 2012). However, how much land is really unused and available is contested (Erb et al., 2007; Haberl et al., 2010; Coelho et al., 2012). Contrasting views on future technical bioenergy potentials from dedicated biomass plantations can be explained by differences in assumptions regarding feasible future agricultural crop yields, livestock feeding efficiency, land availability for energy crops and yields of energy crops (Dornburg et al., 2010; Batidzirai et al., 2012; Erb et al., 2012a). Most scientists agree that increases in food crop yields and higher feeding efficiencies and lower consumption of animal products results in higher technical bioenergy potential. Also, there is a large agreement that careful policies for implementation focused on land-use zoning approaches

(including nature conservation and biodiversity protection), multifunctional land use, integration of food and energy production, avoidance of detrimental livelihood impacts, e.g., on livestock grazing and subsistence farming, and consideration of equity issues, and sound management of impacts on water systems are crucial for sustainable solutions.

Reduced traditional biomass demand. A substantial quantity of biomass will become available for modern applications by improving the end-use efficiency of traditional biomass consumption for energy, mostly in households but also within small industries (such as charcoal kilns, brick kilns, etc.). Traditional bioenergy represents approximately 15 % of total global energy use and 80 % of current bioenergy use (≈ 35 EJ/yr) and helps meeting the cooking needs of ~ 2.6 billion people (Chum et al., 2011; IEA, 2012b). Traditional bioenergy use covers several end-uses including cooking, water, and space heating, and small-industries (such as brick and pottery kilns, bakeries, and many others). Cooking is the dominant end use; it is mostly done in open fires and rudimentary stoves, with approximately 10–20 % conversion efficiency, leading to very high primary energy consumption. Advanced woodburning and biogas stoves can potentially reduce biomass fuel consumption by 60 % or more (Jetter et al., 2012) and further lower the atmospheric radiative forcing, reducing CO₂ emissions, and in many cases black carbon emissions, by up to 90 % (Anenberg et al., 2013). Assuming that actual savings reach on average 30–60 % of current consumption, the total bioenergy potential from reducing traditional bioenergy demand can be estimated at 8–18 EJ/yr. An unknown fraction of global traditional biomass is consumed in a non-environmentally sustainable way, leading to forest degradation and deforestation.

Detailed country studies have estimated the fraction of non-renewable biomass from traditional bioenergy use to vary widely, e.g., from 1.6 % for the Democratic Republic of Congo to 73 % for Burundi (CDM-SSC WG, 2011) with most countries in the range between 10–30 % (i.e., meaning that 70–90 % of total traditional bioenergy use is managed sustainably). Thus a fraction of the traditional biomass saved through better technology, should not be used for other energy purposes but simply not consumed to help restore the local ecosystems.

11.13.3 Bioenergy conversion: technologies and management practices

Numerous conversion technologies can transform biomass to heat, power, liquid, and gaseous fuels for use in the residential, industrial, transport, and power sectors (see Chum et al., 2011; GEA, 2012) for a comprehensive coverage of each alternative, and Figure 11.21 for the pathways concerning liquid and gaseous fuels). Since SRREN, the major advances in the large-scale production of bioenergy include the increasing use of hybrid biomass-fossil fuel systems. For example, current commercial coal and biomass co-combustion technologies are the lowest-cost technologies for implementing renewable energy policies, enabled by the large-scale pelletized feedstocks trade (REN21, 2013; Junginger et al., 2014). Direct biopower use is also increasing commercially on a global scale (REN21, 2013, p. 21). In fact, using biomass for electricity and heat, for example, co-firing of woody biomass with coal in the near term and large heating systems coupled with networks for district heating, and biochemical processing of waste biomass, are among the most cost-efficient and effective biomass applications for GHG emission reduction in modern pathways (Sterner and Fritsche, 2011).

Integrated gasification combined cycle (IGCC) technologies for co-production of electricity and liquid fuels from coal and biomass with higher efficiency than current commercial processes are in demonstration phase to reduce cost (Williams et al., 2011; GEA, 2012; Larson et al., 2012). Coupling of biomass and natural gas for fuels is another option for liquid fuels (Baliban et al., 2013) as the biomass gasification technology development progresses. Simulations suggest that integrated gasification facilities are technically feasible (with up to 50 % biomass input; Meerman et al., 2011), and economically attractive with a CO₂ price of about 66 USD₂₀₁₀/tCO₂ (50 EUR₂₀₁₀/tCO₂) (Meerman et al., 2012). Many gasification technology developments around the world are in pilot, demonstration, operating first commercial scale for a variety of applications (see examples in Bacovsky et al., 2013; Balan et al., 2013).

Many pathways and feedstocks (Figure 11.21) can lead to biofuels for aviation. The development of biofuel standards started and enabled testing of 50 % biofuel in jet fuel for commercial domestic and transatlantic flights by consortia of governments, aviation industry, and associations (IEA, 2010; REN21, 2013). Advanced 'drop in' fuels, such as iso-butanol, synthetic aviation kerosene from biomass gasification

or upgrading of pyrolysis liquids, can be derived through a number of possible conversion routes such as hydro treatment of vegetable oils, iso-butanol, and Fischer-Tropsch synthesis from gasification of biomass (Hamelinck and Faaij, 2006; Bacovsky et al., 2010; Meerman et al., 2011, 2012; Rosillo-Calle et al., 2012); see also Chapter 8). In specific cases, powering electric cars with electricity from biomass has higher land-use efficiency and lower global-warming potential (GWP) effects than the usage of bioethanol from biofuel crops for road transport across a range of feedstocks, conversion technologies, and vehicle classes (Campbell et al., 2009; Schmidt et al., 2011)¹³, though costs are likely to remain prohibitive for considerable time (van Vliet et al., 2011a; b; Schmidt et al., 2011).

The number of routes from biomass to a broad range of biofuels, shown in Figure 11.21, includes hydrocarbons connecting today's fossil fuels industry in familiar thermal/catalytic routes such as gasification (Williams et al., 2011; Larson et al., 2012) and pyrolysis (Brown et al., 2011; Bridgwater, 2012; Elliott, 2013; Meier et al., 2013). In addition, advances in genomic technology, the emphasis in systems approach, and the integration between engineering, physics, chemistry, and biology bring together many new approaches to biomass conversion (Liao and Messing, 2012) such as (1) biomolecular engineering (Li et al., 2010; Favaro et al., 2012; Peralta-Yahya et al., 2012; Lee et al., 2013; Yoon et al., 2013); (2) deconstruction of lignocellulosic biomass through combinations of mild thermal and biochemical routes in multiple sequential or consolidated steps using similar biomolecular engineering tools (Rubin, 2008; Chundawat et al., 2011; Beckham et al., 2012; Olson et al., 2012; Tracy et al., 2012; Saddler and Kumar, 2013; Kataeva et al., 2013); and (3) advances in (bio)catalysis and basic understanding of the synthesis of cellulose are leading to routes for many fuels and chemicals under mild conditions (Serrano-Ruiz et al., 2010; Carpita, 2012; Shen et al., 2013; Triantafyllidis et al., 2013; Yoon et al., 2013). Fundamental understanding of biofuel production increased for microbial genomes by forward engineering of cyanobacteria, microalgae, aiming to arrive at minimum genomes for synthesis of biofuels or chemicals (Chen and Blankenship, 2011; Eckert et al., 2012; Ungerer et al., 2012; Jones and Mayfield, 2012; Kontur et al., 2012; Lee et al., 2013).

Bioenergy coupled with CCS (Spath and Mann, 2004; Liu et al., 2010) is seen as an option to mitigate climate change through negative emissions if CCS can be successfully deployed (Cao and Caldeira 2010; Lenton and Vaughan 2009). BECCS features prominently in long-run mitigation scenarios (Sections 6.3.2 and 6.3.5) for two reasons: (1) The potential for negative emissions may allow shifting emissions in time; and (2) in scenarios, negative emissions from BECCS compensate for residual emissions in other sectors (most importantly transport) in the second half of the 21st century. As illustrated in Figure 11.22, BECCS is markedly different than fossil CCS because it not only reduces CO₂ emissions by storing C in long-term geological sinks, but it continu-

¹³ Biomass can be used for electric transport and biofuels within one pathway (Macedo et al., 2008)




Figure 11.22 | Illustration of the sum of CO₂eq (GWP₁₀₀)* emissions from the process chain of alternative transport and power generation technologies both with and without CCS. (*Differences in C-density between forest biomass and switchgrass are taken into account but not calorific values (balance-of-plant data are for switchgrass, ref. Larson et al., 2012). Specific emissions vary with biomass feedstock and conversion technology combinations, as well as lifecycle GHG calculation boundaries. For policy relevant purposes, counterfactual and market-mediated aspects (e.g., iLUC), changes in soil organic carbon, or changes in surface albedo need also to be considered, possibly leading to significantly different outcomes, quantitatively (Section 11.13.4, Figures 11.23 and 11.24). Unit: gCO₂eq/MJ_{el} (left y-axis, electricity); gCO₂eq/MJ combusted (right y-axis, transport fuels). Direct CO₂ emissions from energy conversion ('vented' and 'stored') are adapted from the mean values in Tables 12.7, 12.8, and 12.15 of ref. [1], which are based on the work of refs. [2, 3], and characterized with the emission metrics in ref. [4]. Impacts upstream in the supply chain associated with feedstock procurement (i.e., sum of GHGs from mining/cultivation, transport, etc.) are adapted from refs. [5, 6] and Figure 11.23 (median values).

¹Larson, et al. (2012); ²Woods, et al., (2007) ; ³Liu et al. (2010); ⁴Guest et al. (2013); ⁵Turconi et al. (2013); ⁶Jaramillo et al. (2008)

Notes:

* Global Warming Potential over 100 years. See Glossary and Section 1.2.5.

ally sequesters CO₂ from the air through regeneration of the biomass resource feedstock.

BECCS deployment is in the development and exploration stages. The most relevant BECCS project is the 'Illinois Basin—Decatur Project' that is projected to inject 1 MtCO₂/yr (Gollakota and McDonald, 2012; Senel and Chugunov, 2013). In the United States, two ethanol fuel production by fermentation facilities are currently integrated commercially with carbon dioxide capture, pipeline transport, and use in enhanced oil recovery in nearby facilities at a rate of about 0.2 MtCO₂/yr (DiPietro et al., 2012). Altogether, there are 16 global BECCS projects in exploration stage (Karlsson and Byström, 2011).

Critical to overall CO₂ storage is the realization of a lignocellulosic biomass supply infrastructure for large-scale commodity feedstock production and efficient advanced conversion technologies at scale; both benefit from cost reductions and technological learning as does the integrated system with CCS, with financial and institutional conditions that minimize the risks of investment and facilitate dissemination (Eranki and Dale, 2011; IEA, 2012c, 2013). Integrated analysis is needed to capture system and knock-on effects for bioenergy potentials. A nascent feedstock infrastructure for densified biomass trading globally could indicate decreased pressure on the need for closely co-located storage and production (IEA, 2011; Junginger et al., 2014).

The overall technical potential is estimated to be around 10 GtCO₂ storage per year for both Integrated Gasification Combined Cycle (IGCC)-CCS co-firing (IGCC with co-gasification of biomass), and Biomass Integrated Gasification Combined Cycle (BIGCC)-CCS dedicated, and around 6 GtCO₂ storage for biodiesel based on gasification and Fischer-Tropsch synthesis (FT diesel), and 2.7 GtCO₂ for biomethane production (Koorneef et al., 2012, 2013). Another study estimates the potential capacity (similar to technical potential) to be between 2.4 and 10 GtCO₂ per year for 2030–2050 (McLaren, 2012). The economic potential, at a CO₂ price of around 70 USD/t is estimated to be around 3.3 GtCO₂, 3.5 GtCO₂, 3.1 GtCO₂ and 0.8 GtCO₂ in the corresponding four cases, judged to be those with highest economic potential (Koorneef et al., 2012, 2013). Potentials are assessed on a route-by-route basis and cannot simply be added, as they may compete and substitute each other. Practical figures might be not much higher than 2.4 GtCO₂ per year at 70–250 USD/tCO₂ (McLaren, 2012). Altogether, until 2050, the economic potential is anywhere between 2–10 GtCO₂ per year. Some climate stabilization scenarios see considerable higher deployment towards the end of the century, even in some 580–650 ppm scenarios, operating under different time scales, socioeconomic assumptions, technology portfolios, CO₂ prices, and interpreting BECCS as part of an overall mitigation framework (e.g., Rose et al., 2012; Kriegler et al., 2013; Tavoni and Socolow, 2013).

Possible climate risks of BECCS relate to reduction of land carbon stock, feasible scales of biomass production and increased N₂O emis-

sions, and potential leakage of CO₂, which has been stored in deep geologic reservoirs (Rhodes and Keith, 2008). The assumptions of sufficient spatially appropriate CCS capture, pipeline, and storage infrastructure are uncertain. The literature highlights that BECCS as well as CCS deployment is dependent on strong financial incentives, as they are not cost competitive otherwise (Sections 7.5.5; 7.6.4; 7.9; 7.12).

Figure 11.22 illustrates some GHG effects associated with BECCS pathways. Tradeoffs between CO₂ capture rate and feedstock conversion efficiency are possible. Depicted are pathways with the highest removal rate but not necessarily with the highest feedstock conversion rate. Among all BECCS pathways, those based on integrated gasification combined cycle produce most significant geologic storage potential from biomass, alone (shown in Figure 11.23, electricity) or coupled with coal. Fischer-Tropsch diesel fuel production with biomass as feedstock and CCS attached to plant facilities could enable BECCS for transport; uncertainties in input factors, and output metrics warrant further research (van Vliet et al., 2009). Fischer-Tropsch diesel would also allow net removal but at lower rates than BIGCC.

Economics of scale in power plant size are crucial to improve economic viability of envisaged BECCS projects. Increasing power plant size requires higher logistic challenges in delivering biomass.

Scales of 4,000 to 10,000 Mg/day needed for > 600 MW power plants could become feasible as the biomass feedstock supply logistic development with manageable logistic costs if biomass is derived from high-yield monocrops; logistical costs are more challenging when biomass is derived from residues (e.g., Argo et al., 2013; Junginger et al., 2014). Large-scale biomass production with flexible integrated poly-generation facilities for fuels and/or power can improve the techno-economic performance, currently above market prices to become more economically competitive over time (Meerman et al., 2011). In the future, increased operating experience of BECCS IGCC-CCS through technological improvements and learning could enable carbon neutral electricity and, in combination with CCS, could result in net removal of CO₂ (Figure 11.22). BECCS is among the lowest cost CCS options for a number of key industrial sectors (Meerman et al., 2013). It should be noted that primary empiric cost and performance data for dedicated bioenergy plants are not yet available and needed for comprehensively assessing BECCS. The current status of CCS and on-going research issues are discussed in Sections 7.5.5 and 7.6.4. Social concerns constitute a major barrier for implement demonstration and deployment projects.

Integrated bio-refineries continue to be developed; for instance, 10% of the ethanol or corresponding sugar stream goes into bio-products in Brazil (REN21, 2012) including making ethylene for polymers (IEA-ETSAP and IRENA, 2013). Multi product bio-refineries could produce a wider variety of co-products to enhance the economics of the overall process, facilitating learning in the new industry (IEA, 2011); Lifecycle Analyses (LCAs) for these systems are complex (Pawelzik et al., 2013).




Figure 11.23 | Direct CO₂eq (GWP₁₀₀) emissions from the process chain or land-use disturbances of major bioenergy product systems, not including impacts from LUC (see Figure 11.24). The interpretation of values depends also on baseline assumption about the land carbon sink when appropriate and the intertemporal accounting frame chosen, and should also consider information from Figure 11.24. The lower and upper bounds of the bars represent the minimum and the maximum value reported in the literature. Whenever possible, peer-reviewed scientific literature published post SRREN is used (but results are comparable). Note that narrow ranges may be an artefact of the number of studies for a given case. Results are disaggregated in a manner showing the impact of *Feedstock* production (in gCO₂eq/MJ lower heating value (LHV) of feedstock) and the contributions from end product/conversion technology. Results from conversion into final energy products *Heat, Power, and Transport fuels* include the contribution from *Feedstock* production and are shown in gCO₂eq/MJ of final product. For some pathways, additional site-specific climate forcing agents apply and are presented as separate values to be added or subtracted from the value indicated by the median in the *Feedstock* bar (dark grey). Final products are also affected by these factors, but this is not displayed here. References: Corn 1–7; Oil crops 1, 8, 8–12; Crop residues 1, 4, 13–24; Sugarcane 2, 3, 5, 6, 25–27; Palm Oil 2, 3, 10, 28–31; Perennial grasses 1, 3, 11, 18, 22, 32–40; Short Rotation Woody Crops 1, 3, 6, 12, 22, 33, 35, 37, 38, 41–53; Forestry 5, 6, 38, 49, 54–66; Biogas, open storage: 67–69; Biogas, closed storage 69–71; Waste cooking oil: 22, 72–74. Note that the biofuels technologies for transport from lignocellulosic feedstocks, short rotation woody crops, and crop residues, including collection and delivery, are developing so larger ranges are expected than for more mature commercial technologies such as sugarcane ethanol and waste cooking oil (WCO) biodiesel. The biogas electricity bar represents scenarios using LCAs to explore treating mixtures of a variety of lignocellulosic feedstocks (e.g., ensiled grain or agricultural residues or perennial grasses) with more easily biodegradable wastes (e.g., from animal husbandry), to optimize multiple outputs. Some of the scenarios assume CH₄ leakage, which leads to very high lifecycle emissions.

¹Gelfand et al. (2013); ²Nemecek et al. (2012); ³Hoefnagels et al. (2010); ⁴Kaufman et al. (2010); ⁵Cherubini et al. (2009); ⁶Cherubini (2012); ⁷Wang et al. (2011b); ⁸Milazzo et al. (2013); ⁹Goglio et al. (2012); ¹⁰Stratton et al. (2011); ¹¹Fazio and Monti (2011); ¹²Börjesson and Tufvesson (2011); ¹³Cherubini and Ulgiati (2010); ¹⁴Li et al. (2012); ¹⁵Luo et al. (2009); ¹⁶Gabrielle and Gagnaire (2008); ¹⁷Smith et al. (2012b); ¹⁸Anderson-Teixeira et al. (2009); ¹⁹Nguyen et al. (2013); ²⁰Searcy and Flynn (2008); ²¹Giuntoli et al. (2013); ²²Whittaker et al. (2010); ²³Wang et al. (2013a); ²⁴Patrizi et al. (2013); ²⁵Souza et al. (2012a); ²⁶Seabra et al. (2011); ²⁷Walter et al. (2011); ²⁸Choo et al. (2011); ²⁹Harsono et al. (2012); ³⁰Sian-gjaeo et al. (2011); ³¹Silalertruksa and Gheewala (2012); ³²Smeets et al. (2009b); ³³Tiwary and Colls (2010); ³⁴Wilson et al. (2011); ³⁵Brandão et al. (2011); ³⁶Cherubini and Jungmeier (2010); ³⁷Don et al. (2012); ³⁸Pucker et al. (2012); ³⁹Monti et al. (2012); ⁴⁰Bai et al. (2010); ⁴¹Bacenetti et al. (2012); ⁴²Budsberg et al. (2012); ⁴³González-García et al. (2012a); ⁴⁴González-García (2012b); ⁴⁵Stephenson et al. (2010); ⁴⁶Hennig and Gawor (2012); ⁴⁷Buonocore et al. (2012); ⁴⁸Gabrielle et al. (2013); ⁴⁹Dias and Arroja (2012); ⁵⁰González-García et al. (2012b); ⁵¹Roedl (2010); ⁵²Djomo et al. (2011); ⁵³Njakou Djomo et al. (2013); ⁵⁴McKechnie et al. (2011); ⁵⁵Pa et al. (2012); ⁵⁶Puettmann et al. (2010); ⁵⁷Guest et al. (2011); ⁵⁸Valente et al. (2011); ⁵⁹Whittaker et al. (2011); ⁶⁰Bright and Strømman (2009); ⁶¹Felder and Dones (2007); ⁶²Solli et al. (2009); ⁶³Lindholm et al. (2011); ⁶⁴Mallia and Lewis (2013); ⁶⁵Bright et al. (2010); ⁶⁶Bright and Strømman (2010); ⁶⁷Rehl et al. (2012); ⁶⁸Blengini et al. (2011); ⁶⁹Boulamanti et al. (2013); ⁷⁰Lansche and Müller (2012); ⁷¹De Meester et al. (2012); ⁷²Sunde et al. (2011); ⁷³Thamsiroj and Murphy (2011); ⁷⁴Talens Peiró et al. (2010)

There are alternatives to land-based bioenergy. Microalgae, for example, offer a high-end technical potential. However, it might be compromised by water supply, if produced in arid land, or by impacts on ocean ecosystems. To make microalgae cost competitive, maximizing algal lipid content (and then maximizing growth rate) requires technological breakthroughs (Davis et al., 2011a; Sun et al., 2011; Jonker and Faaij, 2013). The market potential depends on the co-use of products for food, fodder, higher value products, and fuel markets (Chum et al., 2011).

Similarly, lignocellulosic feedstocks produced from waste or residues, or grown on land unsupportive of food production (e.g., contaminated land for remediation as in previously mined land) have been suggested to reduce socio-environmental impact. In addition, lignocellulosic feedstocks can be bred specifically for energy purposes, and can be harvested by coupling collection and pre-processing (densification and others) in depots prior to final conversion, which could enable delivery of more uniform feedstocks throughout the year (Eranki and Dale, 2011; U.S. DOE, 2011; Argo et al., 2013).

Various conversion pathways are in research and development (R&D), near commercialization, or in early deployment stages in several countries (see Section 2.6.3 in Chum et al., 2011). More productive land is also more economically attractive for cellulosic feedstocks, in which case competition with food production is more likely. Depending on the feedstock, conversion process, prior land use, and land demand, lignocellulosic bioenergy can be associated with high or low GHG emissions (e.g., Davis et al., 2011b). Improving agricultural lands and reducing non-point pollution emissions to watersheds remediate nitrogen run off and increase overall ecosystems' health (Van Dam et al., 2009a; b; Gopalakrishnan et al., 2012). Also regeneration of saline lands by salt-tolerant tree and grass species can have a large potential on global scale as demonstrated by Wicke et al. (2011).

A range of agro-ecological options to improve agricultural practices such as no/low tillage conservation, agroforestry, etc., have potential to increase yields (e.g., in sub-Saharan Africa), while also providing a range of co-benefits such as increased soil organic matter. Such options require a much lower level of investment and inputs and are thus more readily applicable in developing countries, while also holding a low risk of increased GHG emissions (Keating et al., 2013).

Substantial progress has also been achieved in the last four years in small-scale bioenergy applications in the areas of technology innovation, impact evaluation and monitoring, and in large-scale implementation programmes. For example, advanced combustion biomass cookstoves, which reduce fuel use by more than 60% and hazardous pollutant as well as short-lived climate pollutants by up to 90%, are now in the last demonstration stages or commercial (Kar et al., 2012; Anenberg et al., 2013). Innovative designs include micro-gasifiers, stoves with thermoelectric generators to improve combustion efficiency and provide electricity to charge LED lamps while cooking, stoves with advanced combustion chamber designs, and multi-use stoves (e.g.,


cooking and water heating for bathing (Ürge-Vorsatz et al., 2012; Anenberg et al., 2013). Biogas stoves, in addition to providing clean combustion, help reduce the health risks associated with the disposal of organic wastes. There has also been a boost in cookstove dissemination efforts ranging from regional (multi-country) initiatives (Wang et al., 2013b) to national, and project-level interventions. In total, more than 200 large-scale cookstove projects are in place worldwide, with several million efficient cookstoves installed each year (Cordes, 2011). A Global Alliance for Clean Cookstoves has been launched that is promoting the adoption of 100 million clean and efficient cookstoves per year by 2030 and several countries have launched National Cookstove Programs in recent years (e.g., Mexico, Peru, Honduras, and others). Many cookstove models are now manufactured in large-scale industrial facilities using state-of-the-art materials and combustion design technology. Significant efforts are also in place to develop international standards and regional stove testing facilities. In addition to providing tangible local health and other sustainable benefits, replacing traditional open fires with efficient biomass cookstoves has a global mitigation potential estimated to be between 0.6 and 2.4 GtCO₂/yr (Ürge-Vorsatz et al., 2012).

Small-scale decentralized biomass power generation systems based on biomass combustion and gasification and biogas production systems have the potential to meet the electricity needs of rural communities in the developing countries. The biomass feedstocks for these small-scale systems could come from residues of crops and forests, wastes from livestock production, and/or from small-scale energy plantations (Faaij, 2006).

11.13.4 GHG emission estimates of bioenergy production systems

The combustion of biomass generates gross GHG emissions roughly equivalent to the combustion of fossil fuels. If bioenergy production is to generate a net reduction in emissions, it must do so by offsetting those emissions through increased net carbon uptake of biota and soils. The appropriate comparison is then between the net biosphere flux in the absence of bioenergy compared to the net biosphere flux in the presence of bioenergy production. Direct and indirect effects need to be considered in calculating these fluxes.

Bioenergy systems directly influence local and global climate through (i) GHG emissions from fossil fuels associated with biomass production, harvest, transport, and conversion to secondary energy carriers (von Blottnitz and Curran, 2007; van der Voet et al., 2010); (ii) CO₂ and other GHG emissions from biomass or biofuel combustion (Cherubini et al., 2011); (iii) atmosphere-ecosystem exchanges of CO₂ following land disturbance (Berndes et al., 2013; Haberl, 2013); (iv) climate forcing resulting from emissions of short-lived GHGs like black carbon and other chemically active gases (NO_x, CO, etc.) (Tsao et al., 2012; Jetter et al., 2012); (v) climate forcing resulting from alteration of biophysical properties of the land surface affecting the surface energy balance



LUC Estimate Type		Assessment Method		Uncertainty Bars	
◆ Scenarios, Marginal LUC	▲ Arithmetic Mean, Marginal LUC	■ Causal Descriptive	Distribution Statistics	97.5 th Percentile	○ +iLUC 100%
◆ Miscanthus	▲ Switchgrass	■ Deterministic (Simplified)			
◆ Maize Stover	▲ Median, Marginal LUC	■ Optimization Modeling	50 th Percentile	○ +iLUC 50%	
◆ Switchgrass	▲ Maximum	■ Economic Equilibrium Modeling			25 th Percentile
◆ Scenarios, Average LUC	▲ Minimum	■ Meta-analysis			
○ iLUC Factor					



Figure 11.24 | Estimates of GHG_{LUC} emissions—GHG emissions from biofuel production-induced LUC (as $\text{gCO}_2\text{eq}/\text{MJ}_{\text{fuel produced}}$) over a 30-year time horizon organized by fuel(s), feedstock, and study. Assessment methods, LUC estimate types and uncertainty metrics are portrayed to demonstrate the diversity in approaches and differences in results within and across any given category. Points labeled ‘a’ on the Y-axis represent a commonly used estimate of lifecycle GHG emissions associated with the direct supply chain of petroleum gasoline (frame A) and diesel (frame B). These emissions are not directly comparable to GHG_{LUC} because the emission sources considered are different, but are potentially of interest for scaling comparison. Based on Warner et al. (2013). Please note: These estimates of global LUC are highly uncertain, unobservable, unverifiable, and dependent on assumed policy, economic contexts, and inputs used in the modelling. All entries are not equally valid nor do they attempt to measure the same metric despite the use of similar naming conventions (e.g., iLUC). In addition, many different approaches to estimating GHG_{LUC} have been used. Therefore, each paper has its own interpretation and any comparisons should be made only after careful consideration. * CO_2eq includes studies both with and without CH_4 and N_2O accounting.

(e.g., from changes in surface albedo, heat and water fluxes, surface roughness, etc.; (Bonan, 2008; West et al., 2010a; Pielke Sr. et al., 2011); and (vi) GHGs from land management and perturbations to soil biogeochemistry, e.g., N_2O from fertilizers, CH_4 , etc. (Cai, 2001; Allen et al., 2009). Indirect effects include the partial or complete substitution of fossil fuels and the indirect transformation of land use by equilibrium effects. Hence, the total climate forcing of bioenergy depends on feedstock, site-specific climate and ecosystems, management conditions, production pathways, end use, and on the interdependencies with energy and land markets.

In contrast, bioenergy systems have often been assessed (e.g., in LCA studies, integrated models, policy directives, etc.) under the assumption that the CO_2 emitted from biomass combustion is climate neutral¹⁴ because the carbon that was previously sequestered from the atmosphere will be re-sequestered if the bioenergy system is managed sustainably (Chum et al., 2011; Creutzig et al., 2012a; b). The shortcomings of this assumption have been extensively discussed in environmental impact studies and emission accounting mechanisms (Searchinger et al., 2009; Searchinger, 2010; Cherubini et al., 2011; Haberl, 2013).

Studies also call for a consistent and case-specific carbon stock/flux change accounting that integrates the biomass system with the global carbon cycle (Mackey et al., 2013). As shown in Chapter 8 of WGI (Myhre and Shindell, 2013) and (Plattner et al., 2009; Fuglestedt et al., 2010), the climate impacts can be quantified at different points along a cause-effect chain, from emissions to changes in temperature and sea level rise. While a simple sum of the net CO_2 fluxes over time can inform about the skewed time distribution between sources and sinks (‘C debt’; Marland and Schlamadinger, 1995; Fargione et al., 2008; Bernier and Paré, 2013), understanding the climate implication as it relates to policy targets (e.g., limiting warming to 2 °C) requires models and/or metrics that also include temperature effects and climate consequence (Smith et al., 2012c; Tanaka et al., 2013). While the warming from fossil fuels is nearly permanent as it persists for thousands of years, direct impacts from renewable bioenergy systems cause a perturbation in global temperature that is temporary and even at

times cooling if terrestrial carbon stocks are not depleted (House et al., 2002; Cherubini et al., 2013; Joos et al., 2013; Mackey et al., 2013). The direct, physical climate effects at various end-points need to be fully understood and characterized—despite the measurement challenges that some climate forcing mechanisms can entail (West et al., 2010b; Anderson-Teixeira et al., 2012), and coherently embedded in mitigation policy scenarios along with the possible counterfactual effects. For example, in the specific case of existing forests that may continue to grow if not used for bioenergy, some studies employing counterfactual baselines show that forest bioenergy systems can temporarily have higher cumulative CO_2 emissions than a fossil reference system (for a time period ranging from a few decades up to several centuries; (Repo et al., 2011; Mitchell et al., 2012; Pingoud et al., 2012; Bernier and Paré, 2013; Guest et al., 2013; Helin et al., 2013; Holtsmark, 2013).

In some cases, cooling contributions from changes in surface albedo can mitigate or offset these effects (Arora and Montenegro, 2011; O’Halloran et al., 2012; Anderson-Teixeira et al., 2012; Hallgren et al., 2013).

Accounting always depends on the time horizon adopted when assessing climate change impacts, and the assumed baseline, and hence includes value judgements (Schwietzke et al., 2011; Cherubini et al., 2013; Kløverpris and Mueller, 2013).

Two specific contributions to the climate forcing of bioenergy, not addressed in detail in SRREN include N_2O and biogeophysical factors.

Nitrous oxide emissions: For first-generation crop-based biofuels, as with food crops (see Chapter 11), emissions of N_2O from agricultural soils is the single largest contributor to direct lifecycle GHG emissions, and one of the largest contributors across many biofuel production cycles (Smeets et al., 2009a; Hsu et al., 2010). Emission rates can vary by as much as 700 % between different crop types for the same site, fertilization rate, and measurement period (Kaiser and Ruser, 2000; Don et al., 2012; Yang et al., 2012). Increased estimates of N_2O emissions alone can convert some biofuel systems from apparent net sinks to net sources (Crutzen et al., 2007; Smith et al., 2012c). Improvements in nitrogen use efficiency and nitrogen inhibitors can substantially reduce emissions of N_2O (Robertson and Vitousek, 2009). For some specific crops, such as sugarcane, N_2O emissions can be low (Macedo et al., 2008; Seabra et al., 2011) or high (Lisboa et al., 2011). Other bioenergy crops require minimal or zero N fertilization and can reduce GHG emissions relative to the former land use where they replace conventional food crops (Clair et al., 2008).

¹⁴ The neutrality perception is linked to a misunderstanding of the guidelines for GHG inventories, e.g., IPCC—Land Use, Land-Use Change and Forestry (2000) states “Biomass fuels are included in the national energy and carbon dioxide emissions accounts for informational purposes only. Within the energy module biomass consumption is assumed to equal its regrowth. Any departures from this hypothesis are counted within the Land Use Change and Forestry Model.” Carbon neutrality is valid if the countries account for LUC in their inventories for self-produced bioenergy.

Biogeophysical factors: Land cover changes or land-use disturbances of the surface energy balance, such as surface albedo, surface roughness, and evapotranspiration influence the climate system (Betts, 2001; Marland et al., 2003; Betts et al., 2007; Bonan, 2008; Jackson et al., 2008; Mahmood et al., 2013). Perturbations to these can lead to both direct and indirect climate forcings whose impacts can differ in spatial extent (global and/or local) (Bala et al., 2007; Davin et al., 2007). Surface albedo is found to be the dominant direct biogeophysical climate impact mechanism linked to land cover change at the global scale, especially in areas with seasonal snow cover (Claussen et al., 2001; Bathiany et al., 2010), with radiative forcing effects possibly stronger than those of the co-occurring C-cycle changes (Randerson et al., 2006; Lohila et al., 2010; Bright et al., 2011; Cherubini et al., 2012; O'Halloran et al., 2012). Land cover changes can also affect other biogeophysical factors like evapotranspiration and surface roughness, which can have important local (Loarie et al., 2011; Georgescu et al., 2011) and global climatic consequences (Bala et al., 2007; Swann et al., 2010, 2011). Biogeophysical climate impacts from changes in land use are site-specific and show variations in magnitude across different geographic regions and biomes (Bonan, 2008; Anderson, 2010; Pielke Sr. et al., 2011; Anderson-Teixeira et al., 2012). Biogeophysical impacts should be considered in climate impact assessments and in the design of land-use policies to adequately assess the net impacts of land-use mitigation options (Jackson et al., 2008; Betts, 2011; Arora and Montenegro, 2011) as their size may be comparable to impacts from changes to the C cycle.

Figure 11.23 illustrates the range of lifecycle global direct climate impact (in g CO₂ equivalents per MJ, after characterization with GWP time horizon=100 years) attributed to major global bioenergy products reported in the peer-reviewed literature after 2010. Results are broadly comparable to those of Chapter 2 in SRREN (Figures 2.10 and 2.11 in SRREN; Chum et al., 2011) Those figures displayed negative emissions, resulting from crediting emission reduction due to substitution effects. This appendix refrains from allocating credits to feedstocks to avoid double accounting.

Significant variation in the results reflects the wide range of conversion technologies and the reported performances in addition to analyst assumptions affecting system boundary completeness, emission inventory completeness, and choice of allocation method (among others). Additional 'site-specific' land-use considerations such as changes in soil organic carbon stocks (Δ SOC), changes in surface albedo (Δ albedo), and the skewed time distribution of terrestrial biogenic CO₂ fluxes can either reduce or compound land-use impacts and are presented to exemplify that, for some bioenergy systems, these impacts can be greater in magnitude than lifecycle impacts from feedstock cultivation and bioenergy product conversion. 'Site-specific' land-use considerations are geographically explicit and highly sensitive to background climate conditions, soil properties, biomass yields, and land management regimes. The figure reveals that studies find very different values

depending on the boundaries of analysis chosen, site-specific effects, and management methods. Nonetheless, it is clear that fuels from sugarcane, perennial grasses, crop residues, and waste cooking oil are more beneficial than other fuels (LUC emissions can still be relevant, see Figure 11.23). Another important result is that albedo effects and site-specific CO₂ fluxes are highly variable for different forest systems and environmental conditions and determine the total climate forcing of bioenergy from forestry.

Direct and indirect land-use change: Direct land-use change occurs when bioenergy crops displace other crops or pastures or forests, while iLUC results from bioenergy deployment triggering the conversion to cropland of lands, somewhere on the globe, to replace some portion of the displaced crops (Searchinger et al., 2008; Kløverpris et al., 2008; Delucchi, 2010; Hertel et al., 2010). Direct LUC to establish biomass cropping systems can increase the net GHG emissions, for example, if carbon-rich ecosystems such as wetlands, forests, or natural grasslands are brought into cultivation (Gibbs et al., 2008; UNEP, 2009, p. 2009; Chum et al., 2011). Biospheric C losses associated with LUC from some bioenergy schemes can be, in some cases, more than hundred times larger than the annual GHG savings from the assumed fossil fuel replacement (Gibbs et al., 2008; Chum et al., 2011). Impacts have been shown to be significantly reduced when a dynamic baseline includes future trends in global agricultural land use (Kløverpris and Mueller, 2013). Albeit at lower magnitude, beneficial LUC effects can also be observed, for example, when some semi-perennial crops, perennial grasses or woody plants replace annual crops grown with high fertilizer levels, or where such plants are produced on lands with carbon-poor soils (Tilman et al., 2006; Harper et al., 2010; Sterner and Fritsche, 2011; Sochacki et al., 2012). In particular, *Miscanthus* improves soil organic carbon reducing overall GHG emissions (Brandão et al., 2011); degraded USA Midwest land for economic agriculture, over a 20-year period, shows successional perennial crops without the initial carbon debt and indirect land-use costs associated with food-based biofuels (Gelfand et al., 2013). Palm oil, when grown on more marginal grasslands, can deliver a good GHG balance and net carbon storage in soil (Wicke et al., 2008). Such lands represent a substantial potential for palm oil expansion in Indonesia without deforestation and draining peat lands (Wicke et al., 2011a).

In long-term rotation forests, the increased removal of biomass for bioenergy may be beneficial or not depending on the site-specific forest conditions (Cherubini et al., 2012b). For long-term rotation biomass, the carbon debt (increased cumulative CO₂ emissions for a duration in the order of a rotation cycle or longer) becomes increasingly important (Schlamadinger and Marland, 1996; Marland and Schlamadinger, 1997; Fargione et al., 2008; McKechnie et al., 2011; Hudiburg et al., 2011). Calculations of specific GHG emissions from long-term rotation forests need to account for the foregone CO₂-accumulation (Searchinger, 2010; Holtmark, 2012; Pingoud et al., 2012; Haberl et al., 2012).

If part of a larger forest is used as a feedstock for bioenergy while the overall forest carbon stock increases (the so-called landscape perspective), then the overall mitigation effects are positive, in particular over several harvesting cycles making use of the faster carbon sequestration rates of younger forests (Daigneault et al., 2012; Ximenes et al., 2012; Lamers and Junginger, 2013; Latta et al., 2013). Nabuurs et al. (2013) observe first signs of a carbon sink saturation in European forest biomass and suggest to focus less on the forest biomass sink strength but to consider a mitigation strategy that maximizes the sum of all the possible components: (1) carbon sequestration in forest biomass; (2) soil and wood products; and (3) the effects of material and energy substitution of woody biomass. In general, the use of easily decomposable residues and wastes for bioenergy can produce GHG benefits (Zanchi et al., 2012), similarly to increasing the biomass outtake from forests affected by high mortality rates (Lamers et al., 2013), whereas the removal of slowly decomposing residues reduces soil carbon accumulation at a site and results in net emissions (Repo et al., 2011). The anticipation of future bioenergy markets may promote optimized forest management practices or afforestation of marginal land areas to establish managed plantations, thus contributing to increased forest carbon stocks (Sedjo and Tian, 2012). Rather than leading to wide-scale loss of forest lands, growing markets for tree products can provide incentives for maintaining or increasing forest stocks and land covers, and improving forest health through management (Eisenbies et al., 2009; Dale et al., 2013). If managed to maximize CO₂ storage rate over the long-term, long-term rotation forests offer low-cost mitigation options, in particular, when woody products keep carbon within the human-built environment over long time-scales (e.g., wood substituting for steel joist; Lippke et al., 2011).

Indirect land-use change is difficult to ascertain because the magnitude of these effects must be modelled (Nassar et al., 2011) raising important questions about model validity and uncertainty (Liska and Perrin, 2009; Plevin et al., 2010; Khanna et al., 2011; Gawel and Ludwig, 2011; Wicke et al., 2012) and policy implications (DeCicco, 2013; Finkbeiner, 2013; Plevin et al., 2013). Available model-based studies have consistently found positive and, in some cases, high emissions from LUC and iLUC, mostly of first-generation biofuels (Figure 11.23), albeit with high variability and uncertainty in results (Hertel et al., 2010; Taheripour et al., 2011; Dumortier et al., 2011; Havlík et al., 2011; Chen et al., 2012; Timilsina et al., 2012; Warner et al., 2014). Causes of the great uncertainty include: incomplete knowledge on global economic dynamics (trade patterns, land-use productivity, diets, use of by-products, fuel prices, and elasticities); selection of specific policies modelled; and the treatment of emissions over time (O'Hare et al., 2009; Khanna et al., 2011; Wicke et al., 2012). In addition, LUC modelling philosophies and model structures and features (e.g., dynamic vs. static model) differ among studies. Variations in estimated GHG emissions from biofuel-induced LUC are also driven by differences in scenarios assessed, varying assumptions, inconsistent definitions across models (e.g., LUC, land type), specific selection of reference sce-

narios against which (marginal) LUC is quantified, and disparities in data availability and quality. The general lack of thorough sensitivity and uncertainty analysis hampers the evaluation of plausible ranges of estimates of GHG emissions from LUC.

Wicke et al. (2012) identified the need to incorporate the impacts of iLUC prevention or mitigation strategies in future modelling efforts, including the impact of zoning and protection of carbon stocks, selective sourcing from low risk-areas, policies and investments to improve agricultural productivity, double cropping, agroforestry schemes, and the (improved) use of degraded and marginal lands (see Box 7.1). Indirect land-use change is mostly avoided in the modelled mitigation pathways in Chapter 6. The relatively limited fuel coverage in the literature precludes a complete set of direct comparisons across alternative and conventional fuels sought by regulatory bodies and researchers.

GHG emissions from LUC can be reduced, for instance through production of bioenergy co-products that displace additional feedstock requirements, thus decreasing the net area needed (e.g., for corn, Wang et al., 2011a; for wheat, Berndes et al., 2011). Proper management of livestock and agriculture can lead to improved resource efficiency, lower GHG emissions, and lower land use while releasing land for bioenergy production as demonstrated for Europe (de Wit et al., 2013) and Mozambique (van der Hilst et al., 2012b). For land transport, cellulosic biomass, such as *Miscanthus*, has been suggested as a relatively low-carbon source for bioethanol that could be produced at scale, but only if iLUC can be avoided by not displacing food and other commodities and if comprehensive national land management strategies are developed (e.g., Dornburg et al., 2010; Scown et al., 2012). Negative iLUC values are theoretically possible (RFA, 2008). Producing biofuels from wastes and sustainably harvested residues, and replacing first-generation biofuel feedstocks with lignocellulosic crops (e.g., grasses) would induce little or no iLUC (Davis et al., 2011b; Scown et al., 2012). While iLUC quantifications remain uncertain, lower agricultural yields, land-intensive diets, and livestock feeding efficiencies, stronger climate impacts and higher energy crop production levels can result in higher LUC-related GHG emissions. Strong global and regional governance (forest protection, zoning), technological change in agriculture and biobased options, and high-yield bioenergy crops and use of residues and degraded land (if available) could all reduce iLUC (Van Dam et al., 2009a; b; Wicke et al., 2009; Fischer et al., 2010; de Wit et al., 2011, 2013; van der Hilst et al., 2012a; Rose et al., 2013). As with any other renewable fuel, bioenergy can replace or complement fossil fuel. The fossil fuel replacement effect, relevant when a global cap on CO₂ emissions is absent, is discussed in Chapter 8.7. Indirect effects are not restricted to indirect GHG effects of production of biomass in agricultural systems; there are also indirect (market-mediated) effects of wood energy, but also effects in terms of biodiversity threats, environmental degradation, and external social costs, which are not considered here.

11.13.5 Aggregate future potential deployment in integrated models

In SRREN scenarios (IPCC, 2011), bioenergy is projected to contribute 80–190 EJ/yr to global primary energy supply by 2050 for 50 % of the scenarios in the two mitigation levels modelled. The min to max ranges were 20–265 EJ/yr for the less stringent scenarios and 25–300 EJ for the tight mitigation scenarios (< 440 ppm). Many of these scenarios coupled bioenergy with CCS. The Global Energy Assessment (GEA, 2012) scenarios project 80–140 EJ by 2050, including extensive use of agricultural residues and second-generation bioenergy to try to reduce the adverse impacts on land use and food production, and the co-processing of biomass with coal or natural gas with CCS to make low net GHG-emitting transport fuels and or electricity.

Traditional biomass demand is steady or declines in most scenarios from 34 EJ/yr. The transport sector increases nearly ten-fold from 2008 to 18–20 EJ/yr while modern uses for heat, power, combinations, and industry increase by factors of 2–4 from 18 EJ in 2008 (Fischedick et al., 2011). The 2010 International Energy Agency (IEA) model projects a contribution of 12 EJ/yr (11 %) by 2035 to the transport sector, including 60 % of advanced biofuels for road and aviation. Bioenergy supplies 5 % of global power generation in 2035, up from 1 % in 2008. Modern heat and industry doubles their contributions from 2008 (IEA, 2010). The future potential deployment level varies at the global and national level depending on the technological developments, land availability, financial viability, and mitigation policies.

The WGI AR5 transformation pathway studies suggest that modern bioenergy could play a significant role within the energy system (Section 6.3.5) providing 5 to 95 EJ/yr in 2030, 10 to 245 EJ/yr in 2050, and 105 to 325 EJ/yr in 2100 under idealized full implementation scenarios (see also Figure 7.12), with immediate, global, and comprehensive incentives for land-related mitigation options. The scenarios project increasing deployment of bioenergy with tighter climate change targets, both in a given year as well as earlier in time (see Figure 6.20). Models project increased dependence, as well as increased deployment, of modern bioenergy, with some models projecting 35 % of total primary energy from bioenergy in 2050, and as much as 50 % of total primary energy from modern bioenergy in 2100. Bioenergy's share of regional total electricity and liquid fuels could be significant—up to 35 % of global regional electricity from biopower by 2050, and up to 70 % of global regional liquid fuels from biofuels by 2050. However, the cost-effective allocation of bioenergy within the energy system varies across models. Several sectoral studies, focusing on biophysical constraints, model assumptions (e.g., estimated increase in crop yields over large areas) and current observations, suggest to focus on the lower half of the ranges reported above (Field et al., 2008; Campbell et al., 2008; Johnston et al., 2009a, 2011; Haberl et al., 2013b).

BECCS features prominently in many mitigation scenarios. BECCS is deployed in greater quantities and earlier in time the more stringent the climate policy (Section 6.3.5). Whether BECCS is essential for mitigation, or even sufficient, is unclear. In addition, the likelihood of BECCS deployment is difficult to evaluate and depends on safety con-

Box 11.9 | Examples of co-benefits from biofuel production

Brazilian sugar cane ethanol production provides six times more jobs than the Brazilian petroleum sector and spreads income benefits across numerous municipalities (de Moraes et al., 2010). Worker income is higher than in nearly all other agricultural sectors (de Moraes et al., 2010; Satolo and Bacchi, 2013) and several sustainability standards have been adopted (Viana and Perez, 2013). When substituting gasoline, ethanol from sugar cane also eliminates lead compounds and reduces noxious emissions (Goldemberg et al., 2008). Broader strategic planning, understanding of cumulative impacts, and credible and collaborative decision making processes can help to enhance biodiversity and reverse ecological fragmentation, address direct and iLUC, improve the quality and durability of livelihoods, and other sustainability issues (Duarte et al., 2013).

Co-benefits of palm oil production have been reported in the major producer countries, Malaysia and Indonesia (Sumathi et al., 2008; Lam et al., 2009) as well as from new producer countries (Garcia-Ulloa et al., 2012). Palm oil production results in employment creation as well as in increment state and individual income (Sumathi et al., 2008; Tan et al., 2009; Lam et al., 2009; Sayer

et al., 2012; von Geibler, 2013). When combined with agroforestry, palm oil plantations can increase food production locally and have a positive impact on biodiversity (Lam et al., 2009; Garcia-Ulloa et al., 2012) and when palm oil plantations are installed on degraded land further co-benefits on biodiversity and carbon enhancement (Sumathi et al., 2008; Garcia-Ulloa et al., 2012; Sayer et al., 2012). Further, due to its high productivity, palm oil plantations can produce the same bioenergy input using less land than other bio-energy crops (Sumathi et al., 2008; Tan et al., 2009). Certification in palm oil production can become a means for increasing sustainable production of biofuels (Tan et al., 2009; Edser, 2012; von Geibler, 2013).

Similarly, co-benefits from the production of *Jatropha* as a biofuel crop in developing countries have been reported, mainly when *Jatropha* is planted on degraded land. These include increases in individuals' incomes (Garg et al., 2011; Arndt et al., 2012), improvement in energy security at the local level (von Maltitz and Setzkorn, 2013; Muys et al., 2014), and reducing soil erosion (Garg et al., 2011).

firmations, affordability and public acceptance (see Section 11.13.3 for details). BECCS may also affect the cost-effective emissions trajectory (Richels et al., In Review; Rose et al., 2013).

Some integrated models are cost-effectively trading off lower land carbon stocks and increased land N₂O emissions for the long-run mitigation benefits of bioenergy (Rose et al., 2013; Popp et al., 2013). The models find that bioenergy could contribute effectively to climate change mitigation despite land conversion and intensification emissions. However, as discussed below and in Section 11.9, policy implementation and coordination are factors to consider. In these models, constraining bioenergy has a cost. For instance, limiting global bioenergy availability to 100 EJ/year tripled marginal abatement costs and doubled consumption losses associated with transformation pathways (Rose et al., 2013). Overall outcomes may depend strongly on governance of land use and deployment of best practices in agricultural production (see sections above). Progressive developments in governance of land and modernization of agriculture and livestock and effective sustainability frameworks can help realize large parts of the technical bioenergy potential with low associated GHG emissions.

With increasing scarcity of productive land, the growing demand for food and bioenergy could induce substantial LUC causing high GHG emissions and/or increased agricultural intensification and higher N₂O emissions unless wise integration of bioenergy into agriculture and forestry landscapes occurs (Delucchi, 2010). Consideration of LUC emissions in integrated models show that valuing or protecting global terrestrial carbon stocks reduces the potential LUC-related GHG emissions of energy crop deployment, and could lower the cost of achieving climate change objectives, but could exacerbate increases in agricultural commodity prices (Popp et al., 2011; Reilly et al., 2012). Many integrated models are investigating idealized policy implementation pathways, assuming global prices on GHG (including the terrestrial land carbon stock); if such conditions cannot be realized, certain types of bioenergy could lead to additional GHG emissions. More specifically, if the global terrestrial land carbon stock remains unprotected, large GHG emissions from bioenergy-related LUC alone are possible (Melillo et al., 2009; Wise et al., 2009; Creutzig et al., 2012a; Calvin et al., 2013b).

In summary, recent integrated model scenarios project between 10–245 EJ/yr modern bioenergy deployment in 2050. Good governance and favourable conditions for bioenergy development may facilitate higher bioenergy deployment while sustainability and livelihood concerns might constrain deployment of bioenergy scenarios to low values (see Section 11.13.6).

11.13.6 Bioenergy and sustainable development

The nature and extent of the impacts of implementing bioenergy depend on the specific system, the development context, and on the size of the intervention (Section 11.4.5). The effects on livelihoods

have not yet been systematically evaluated in integrated models (Davis et al., 2013; Creutzig et al., 2012b; Creutzig et al., 2013; Muys et al., 2014), even if human geography studies have shown that bioenergy deployment can have strong distributional impacts (Davis et al., 2013; Muys et al., 2014). The total effects on livelihoods will be mediated by global market dynamics, including policy regulations and incentives, the production model and deployment scale, and place-specific factors such as governance, land tenure security, labour and financial capabilities, among others (Creutzig et al., 2013).

Bioenergy projects can be economically beneficial, e.g., by raising and diversifying farm incomes and increasing rural employment through the production of biofuels for domestic use (Gohin, 2008) or export markets (Wicke et al., 2009; Arndt et al., 2011).

The establishment of large-scale biofuels feedstock production can also cause smallholders, tenants, and herders to lose access to productive land, while other social groups such as workers, investors, company owners, biofuels consumers, and populations who are more responsible for GHG emission reductions enjoy the benefits of this production (van der Horst and Vermeulen, 2011). This is particularly relevant where large areas of land are still unregistered or are being claimed and under dispute by several users and ethnic groups (Dauvergne and Neville, 2010). Furthermore, increasing demand for first-generation biofuels is partly driving the expansion of crops like soy and oil palm, which in turn contribute to promote large-scale agribusinesses at the expense of family and community-based agriculture, in some cases (Wilkinson and Herrera, 2010). Biofuels deployment can also translate into reductions of time invested in on-farm subsistence and community-based activities, thus translating into lower productivity rates of subsistence crops and an increase in intra-community conflicts as a result of the uneven share of collective responsibilities (Mingorría et al., 2010).

Bioenergy deployment is more beneficial when it is not an additional land-use activity expanding over the landscape, but rather integrates into existing land uses and influences the way farmers and forest owners use their land. Various studies indicate the ecosystem services and values that perennial crops have in restoring degraded lands, via agroforestry systems, controlling erosion, and even in regional climate effects such as improved water retention and precipitation (Faaij, 2006; Wicke et al., 2011c; Immerzeel et al., 2013). Examples include adjustments in agriculture practices where farmers, for instance, change their manure treatment to produce biogas, reduce methane and N losses. Changes in management practice may swing the net GHG balance of options and also have clear sustainable development implications (Davis et al., 2013).

Small-scale bioenergy options can provide cost-effective alternatives for mitigating climate change, at the same time helping advance sustainable development priorities, particularly in rural areas of developing countries. IEA (2012b) estimates that 2.6 billion people world-

Table 11.12 | Potential institutional, social, environmental, economic and technological implications of bioenergy options at local to global scale.

Institutional		Scale
May contribute to energy independence (+), especially at the local level (reduce dependency on fossil fuels) (2, 20, 32, 39, 50)	+	Local to national
Can improve (+) or decrease (-) land tenure and use rights for local stakeholders (2, 17, 38, 50)	+/-	Local
Cross-sectoral coordination (+) or conflicts (-) between forestry, agriculture, energy, and/or mining (2, 13, 26, 31, 60)	+/-	Local to national
Impacts on labor rights among the value chain (2, 6, 17)	+/-	Local to national
Promoting of participative mechanisms for small-scale producers (14, 15)	+	Local to national
Social		Scale
Competition with food security including food availability (through reduced food production at the local level), food access (due to price volatility), usage (as food crops can be diverted towards biofuel production), and consequently to food stability. Bio-energy derived from residues, wastes, or by-products is an exception (1, 2, 7, 9, 12, 18, 23)	-	Local to global
Integrated systems (including agroforestry) can improve food production at the local level creating a positive impact towards food security (51, 52, 53, 69, 73, 74). Further, biomass production combined with improved agricultural management can avoid such competition and bring investment in agricultural production systems with overall improvements of management as a result (as observed in Brazil) (60, 63, 66, 67, 70, 71)	+	Local
Increasing (+) or decreasing (-) existing conflicts or social tension (9, 14, 19, 26)	+/-	Local to national
Impacts on traditional practices: using local knowledge in production and treatment of bioenergy crops (+) or discouraging local knowledge and practices (-) (2, 50)	+/-	Local
Displacement of small-scale farmers (14, 15, 19). Bioenergy alternatives can also empower local farmers by creating local income opportunities	+/-	Local
Promote capacity building and new skills (3, 15, 50)	+	Local
Gender impacts (2, 4, 14, 15, 27)	+/-	Local to national
Efficient biomass techniques for cooking (e.g., biomass cookstoves) can have positive impacts on health, especially for women and children in developing countries (42, 43, 44)	+	Local to national
Environmental		Scale
Biofuel plantations can promote deforestation and/or forest degradation, under weak or no regulation (1, 8, 22)	-	Local to global
When used on degraded lands, perennial crops offer large-scale potential to improve soil carbon and structure, abate erosion and salinity problems. Agroforestry schemes can have multiple benefits including increased overall biomass production, increase biodiversity and higher resilience to climate changes. (59, 64, 65, 69, 73)	+	Local to global
Some large-scale bio-energy crops can have negative impacts on soil quality, water pollution, and biodiversity. Similarly potential adverse side-effects can be a consequence of increments in use of fertilizers for increasing productivity (7, 12, 26, 30). Experience with sugarcane plantations has shown that they can maintain soil structure (56) and application of pesticides can be substituted by the use of natural predators and parasitoids (57, 71)	-/+	Local to transboundary
Can displace activities or other land uses (8, 26)	-	Local to global
Smart modernization and intensification can lead to lower environmental impacts and more efficient land use (75, 76)	+	Local to transboundary
Creating bio-energy plantations on degraded land can have positive impacts on soil and biodiversity (12)	+	Local to transboundary
There can be tradeoffs between different land uses, reducing land availability for local stakeholders (45, 46, 47, 48, 49). Multicropping system provide bioenergy while better maintaining ecological diversity and reducing land-use competition (58)	-/+	Local to national
Ethanol utilization leads to the phaseout of lead additives and methyl tertiary-butyl ether (MTBE) and reduces sulfur, particulate matter, and carbon monoxide emissions (55)	+	Local to global
Economic		Scale
Increase in economic activity, income generation, and income diversification (1, 2, 3, 12, 20, 21, 27, 54)	+	Local
Increase (+) or decrease (-) market opportunities (16, 27, 31)	+/-	Local to national
Contribute to the changes in prices of feedstock (2, 3, 5, 21)	+/-	Local to global
May promote concentration of income and/or increase poverty if sustainability criteria and strong governance is not in place (2, 16, 26)	-	Local to regional
Using waste and residues may create socio-economic benefits with little environmental risks (2, 41, 36)	+	Local to regional
Uncertainty about mid- and long-term revenues (6, 30)	-	National
Employment creation (3, 14, 15)	+	Local to regional
Technological		Scale
Can promote technology development and/or facilitate technology transfer (2, 27, 31)	+	Local to global
Increasing infrastructure coverage (+). However if access to infrastructure and/or technology is reduced to few social groups it can increase marginalization (-) (27, 28, 29)	+/-	Local
Bioenergy options for generating local power or to use residues may increase labor demand, creating new job opportunities. Participatory technology development also increases acceptance and appropriation (6, 8, 10, 37, 40)	+	Local
Technology might reduce labor demand (-). High dependent of tech. transfer and/or acceptance	-	Local



¹Alves Finco and Doppler (2010); ²Amigun et al. (2011); ³Arndt et al. (2012); ⁴Arndt et al. (2011); ⁵Arndt et al. (2012); ⁶Awudu and Zhang (2012); ⁷Beringer et al. (2011); ⁸Borzoni (2012); ⁹Bringezu et al. (2012); ¹⁰Cacciatore et al. (2012); ¹¹Cançado et al. (2006); ¹²Danielsen et al. (2009); ¹³Diaz-Chavez (2011); ¹⁴Duvenage et al. (2013); ¹⁵Ewing and Msangi (2009); ¹⁶Gasparatos et al. (2011); ¹⁷German and Schoneveld (2012); ¹⁸Haberl et al. (2011a); ¹⁹Hall et al. (2009); ²⁰Hanff et al. (2011); ²¹Huang et al. (2012); ²²Koh and Wilcove (2008); ²³Koizumi (2013); ²⁴Kyu et al. (2010); ²⁵Madlener et al. (2006); ²⁶Martinelli and Filoso (2008); ²⁷Mwakaje (2012); ²⁸Oberling et al. (2012); ²⁹Schut et al. (2010); ³⁰Selfa et al. (2011); ³¹Steenblik (2007); ³²Stromberg and Gasparatos (2012); ³³Searchinger et al. (2009); ³⁴Searchinger et al. (2008); ³⁵Smith and Searchinger (2012); ³⁶Tilman et al. (2009); ³⁷Van de Velde et al. (2009); ³⁸von Maltitz and Setzkorn (2013); ³⁹Wu and Lin (2009); ⁴⁰Zhang et al. (2011); ⁴¹Fargione et al. (2008); ⁴²Jerneck and Olsson (2013); ⁴³Gurung and Oh (2013); ⁴⁴O'Shaughnessy et al. (2013); ⁴⁵German et al. (2013); ⁴⁶Cotula (2012); ⁴⁷Mwakaje (2012); ⁴⁸Scheidel and Sorman (2012); ⁴⁹Haberl et al. (2013b); ⁵⁰Muys et al. (2014); ⁵¹Egeskog et al. (2011); ⁵²Diaz-Chavez (2012); ⁵³Ewing and Msangi (2009); ⁵⁴de Moraes et al. (2010); ⁵⁵Goldemberg (2007); ⁵⁶Walter et al. (2011); ⁵⁷Macedo (2005); ⁵⁸Langeveld et al. (2013); ⁵⁹Van Dam et al. (2009a; b); ⁶⁰van Dam et al. (2010); ⁶¹van Eijck et al. (2012); ⁶²van Eijck et al. (2014); ⁶³Martinez et al. (2013); ⁶⁴van der Hilst et al. (2010); ⁶⁵van der Hilst et al. (2012); ⁶⁶van der Hilst and Faaij (2012); ⁶⁷van der Hilst et al. (2012b); ⁶⁸Hoefnagels et al. (2013); ⁶⁹Immerzeel et al. (2013); ⁷⁰Lynd et al. (2011); ⁷¹Smeets et al. (2008); ⁷²Smeets and Faaij (2010); ⁷³Wicke et al. (2013); ⁷⁴Wiskerke et al. (2010); ⁷⁵De Wit et al. (2011); ⁷⁶De Wit et al. (2013)

wide depend on traditional biomass for cooking, while 84 % of these belong to rural communities. Use of low-quality fuels and inefficient cooking and heating devices leads to pollution resulting in nearly 4 million premature deaths every year, and a range of chronic illnesses and other health problems (Lim et al., 2012; see Section 9.7.3.1). Modern small-scale bioenergy technologies such as advanced/efficient cook stoves, biogas for cooking and village electrification, biomass gasifiers, and bagasse-based co-generation systems for decentralized power generation, can provide energy for rural communities with energy services that also promote rural development (IEA, 2011). Such bioenergy systems reduce CO₂ emissions from unsustainable biomass harvesting and short-lived climate pollutants, e.g., black carbon, from cleaner combustion (Chung et al., 2012). Scaling up clean cookstove initiatives could not only save 2 million lives a year, but also significantly reduce GHG emissions (Section 11.13.3). Efficient biomass cook stoves and biogas stoves at the same time provide multiple benefits: They reduce the pressure on forests and biodiversity; they reduce exposure to smoke-related health hazards; they reduce drudgery for women in collection fuelwood; and they save money if fuel needs to be purchased (Martin et al., 2011). Benefits from the dissemination of improved cookstoves outweigh their costs by sevenfold, when their health, economic, and environmental benefits are accounted for (Garcia-Frapolli et al., 2010).

Table 11.12 presents the implications of bioenergy options in the light of social, institutional, environmental, economic, and technological conditions. The relationship between bioenergy and these conditions is complex and there could be negative or positive implications, depending on the type of bioenergy option, the scale of the production system and the local context. While biofuels can allow the reduction of fossil fuel use and of GHG emissions, they often shift environmental burdens towards land use-related impacts (i.e., eutrophication, acidification, water depletion, ecotoxicity; EMPA, 2012; Smith and Torn, 2013; Tavoni and Socolow, 2013). Co-benefits and adverse side-effects do not necessarily overlap, neither geographically nor socially (Dauvergne and Neville, 2010; Wilkinson and Herrera, 2010; van der Horst and Vermeylen, 2011). The main potential co-benefits are related to access to energy and impacts on the economy and well-being, jobs creation, and improvement of local resilience (Walter et al., 2011; Creutzig et al., 2013). Main risks of crop-based bioenergy for sustainable develop-

ment and livelihoods include competition for arable land (Haberl et al., 2013b) and consequent impact on food security, tenure arrangements, displacement of communities and economic activities, creation of a driver of deforestation, impacts on biodiversity, water, and soil, or increase in vulnerability to climate change, and unequal distribution of benefits (Sala et al., 2000; Hall et al., 2009; German et al., 2011; Thompson et al., 2011b; IPCC, 2012).

Good governance is an essential component of a sustainable energy system. Integrated studies that compare impacts of bioenergy production between different crops and land management strategies show that the overall impact (both ecological and socio-economic) depends strongly on the governance of land use and design of the bioenergy system see van der Hilst et al. (2012) in the European context, and Van Dam et al. (2009a; b) for different crops and scenarios in Argentina). Van Eijck et al. (2012) show similar differences in impacts between the production and use of *Jatropha* based on smallholder production versus plantation models. This implies that governance and planning have a strong impact on the ultimate result and impact of large-scale bioenergy deployment. Legislation and regulation of bioenergy as well as voluntary certification schemes are required to guide bioenergy production system deployment so that the resources and feedstocks be put to best use, and that (positive and negative) socioeconomic and environmental issues are addressed as production grows (van Dam et al., 2010). There are different options, from voluntary to legal and global agreements, to improve governance of biomass markets and land use that still require much further attention (Verdonk et al., 2007). The integration of bioenergy systems into agriculture and forest landscapes can improve land and water use efficiency and help address concerns about environmental impacts of present land use (Berndes et al., 2004, 2008; Börjesson and Berndes, 2006; Sparovek et al., 2007; Gopalakrishnan et al., 2009, 2011a; b, 2012; Dimitriou et al., 2009, 2011; Dornburg et al., 2010; Batidzirai et al., 2012; Parish et al., 2012; Baum et al., 2012; Busch, 2012), but the global potentials of such systems are difficult to determine (Berndes and Börjesson, 2007; Dale and Kline, 2013). Similarly, existing and emerging guiding principles and governance systems influence biomass resources availability (Stupak et al., 2011). Certification approaches can be useful, but they should be accompanied by effective territorial policy frameworks (Hunsberger et al., 2012).

11.13.7 Tradeoffs and synergies with land, water, food, and biodiversity

This section summarizes results from integrated models (models that have a global aggregate view, but cannot disaggregate place-specific effects in biodiversity and livelihoods discussed above) on land, water, food, and biodiversity. In these models, at any level of future bioenergy supply, land demand for bioenergy depends on (1) the share of bioenergy derived from wastes and residues (Rogner et al., 2012); (2) the extent to which bioenergy production can be integrated with food or fiber production, which ideally results in synergies (Garg et al., 2011; Sochacki et al., 2013) or at least mitigates land-use competition (Bergdes et al., 2013); (3) the extent to which bioenergy can be grown on areas with little current or future production, taking into account growing land demand for food (Nijsen et al., 2012); and (4) the volume of dedicated energy crops and their yields (Haberl et al., 2010; Batidzirai et al., 2012; Smith et al., 2012d). Energy crop yields per unit area may differ by factors of > 10 depending on differences in natural fertility (soils, climate), energy crop plants, previous land use, management and technology (Johnston et al., 2009a; Lal, 2010; Beringer et al., 2011; Pacca and Moreira, 2011; Smith et al., 2012a; Erb et al., 2012a). Assumptions on energy crop yields are one of the main reasons for the large differences in estimates of future area demand of energy crops (Popp et al., 2013). Likewise, assumptions on yields, strategies, and governance on future food/feed crops have large implications for assessments of the degree of land competition between biofuels and these land uses (Batidzirai et al., 2012; de Wit et al., 2013).

However, across models, there are very different potential landscape transformation visions in all regions (Sections 6.3.5 and 11.9.). Overall, it is difficult to generalize on regional land cover effects of mitigation. Some models assume significant land conversion while other models do not. In idealized implementation scenarios, there is expansion of energy cropland and forest land in many regions, with some models exhibiting very strong forest land expansion and others very little by 2030. Land conversion is increased in the 450 ppm scenarios compared to the 550 ppm scenarios, but at a declining share, a result consistent

with a declining land-related mitigation rate with policy stringency. The results of these integrated model studies need to be interpreted with caution, as not all GHG emissions and biogeophysical or socio-economic effects of bioenergy deployment are incorporated into these models, and as not all relevant technologies are represented (e.g., cascade utilization).

Large-scale bioenergy production from dedicated crops may affect water availability and quality (see Section 6.6.2.6), which are highly dependent on (1) type and quantity of local freshwater resources; (2) necessary water quality; (3) competition for multiple uses (agricultural, urban, industrial, power generation), and (4) efficiency in all sector end uses (Gerbens-Leenes et al., 2009; Coelho et al., 2012). In many regions, additional irrigation of energy crops could further intensify existing pressures on water resources (Popp et al., 2011). Studies indicate that an exclusion of severe water scarce areas for bioenergy production (mainly to be found in the Middle East, parts of Asia, and western United States) would reduce global technical bioenergy potentials by 17% until 2050 (van Vuuren et al., 2009). A model comparison study with five global economic models shows that the aggregate food price effect of large-scale lignocellulosic bioenergy deployment (i.e., 100 EJ globally by the year 2050) is significantly lower (+5% on average across models) than the potential price effects induced by climate impacts on crop yields (+25% on average across models (Lotze-Campen et al., 2013). Possibly hence, ambitious climate change mitigation need not drive up global food prices much, if the extra land required for bioenergy production is accessible or if the feedstock, e.g., from forests, does not directly compete for agricultural land. Effective land-use planning and strict adherence to sustainability criteria need to be integrated into large-scale bioenergy projects to minimize competitions for water (for example, by excluding the establishment of biofuel projects in irrigated areas). If bioenergy is not managed properly, additional land demand and associated LUC may put pressures on biodiversity (Groom et al., 2008; see Section 6.6.2.5). However, implementing appropriate management, such as establishing bioenergy crops in degraded areas represents an opportunity where bioenergy can be used to achieve positive environmental outcomes (Nijsen et al., 2012).

References

- Abdalla M., B. Osborne, G. Lanigan, D. Forristal, M. Williams, P. Smith, and M. B. Jones (2013). Conservation tillage systems: a review of its consequences for greenhouse gas emissions. *Soil Use and Management* **29**, 199–209. doi: 10.1111/sum.12030, ISSN: 02660032.
- Ackerman F., S.J. DeCanio, R.B. Howarth, and K. Sheeran (2009). Limitations of integrated assessment models of climate change. *Climatic Change* **95**, 297–315. doi: 10.1007/s10584-009-9570-x, ISSN: 0165-0009, 1573–1480.
- Acuff K., and D.T. Kaffine (2013). Greenhouse gas emissions, waste and recycling policy. *Journal of Environmental Economics and Management* **65**, 74–86. doi: 10.1016/j.jeem.2012.05.003, ISSN: 00950696.
- Agrawal A., A. Chhatre, and R. Hardin (2008). Changing Governance of the World's Forests. *Science* **320**, 1460–1462. doi: 10.1126/science.1155369, ISSN: 0036-8075, 1095–9203.
- Akiyama H., X. Yan, and K. Yagi (2010). Evaluation of effectiveness of enhanced-efficiency fertilizers as mitigation options for N₂O and NO emissions from agricultural soils: meta-analysis: Mitigation options for N₂O and NO emissions. *Global Change Biology* **16**, 1837–1846. doi: 10.1111/j.1365-2486.2009.02031.x, ISSN: 13541013, 13652486.
- Albers H.J., and E.J.Z. Robinson (2013). A review of the spatial economics of non-timber forest product extraction: Implications for policy. *Ecological Economics* **92**, 87–95. doi: 10.1016/j.ecolecon.2012.01.021, ISSN: 09218009.
- Alig R., G. Latta, D. Adams, and B. McCarl (2010). Mitigating greenhouse gases: The importance of land base interactions between forests, agriculture, and residential development in the face of changes in bioenergy and carbon prices. *Forest Policy and Economics* **12**, 67–75. doi: 10.1016/j.forpol.2009.09.012, ISSN: 1389-9341.
- Allen C.D., A.K. Macalady, H. Chenchouni, D. Bachelet, N. McDowell, M. Vennetier, T. Kitzberger, A. Rigling, D.D. Breshears, E. Hogg, and others (2010). A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management* **259**, 660–684.
- Allen D.E., D.S. Mendham, Bhupinderpal-Singh, A. Cowie, W. Wang, R.C. Dalal, and R.J. Raison (2009). Nitrous oxide and methane emissions from soil are reduced following afforestation of pasture lands in three contrasting climatic zones. *Soil Research* **47**, 443–458. Available at: <http://dx.doi.org/10.1071/SR08151>.
- Alves Finco M.V., and W. Doppler (2010). Bioenergy and sustainable development: The dilemma of food security in the Brazilian savannah. *Energy for Sustainable Development* **14**, 194–199.
- Amazon Fund Fundo Amazônia. Available at: http://www.amazonfund.gov.br/FundoAmazonia/fam/site_en.
- Amigun B., J.K. Musango, and W. Stafford (2011). Biofuels and sustainability in Africa. *Renewable and Sustainable Energy Reviews* **15**, 1360–1372. doi: 10.1016/j.rser.2010.10.015, ISSN: 1364-0321.
- Anderson K. (2010). Globalization's effects on world agricultural trade, 1960–2050. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 3007–3021. doi: 10.1098/rstb.2010.0131.
- Anderson R.C., N.A. Krueger, T.B. Stanton, T.R. Callaway, T.S. Edrington, R.B. Harvey, Y.S. Jung, and D.J. Nisbet (2008). Effects of select nitrocompounds on in vitro ruminal fermentation during conditions of limiting or excess added reductant. *Bioresource Technology* **99**, 8655–8661.
- Anderson-Teixeira K.J., S.C. Davis, M.D. Masters, and E.H. Delucia (2009). Changes in soil organic carbon under biofuel crops. *GCB Bioenergy* **1**, 75–96. doi: 10.1111/j.1757-1707.2008.01001.x, ISSN: 1757-1707.
- Anderson-Teixeira K.J., P.K. Snyder, T.E. Twine, S.V. Cuadra, M.H. Costa, and E.H. DeLucia (2012). Climate-regulation services of natural and agricultural ecoregions of the Americas. *Nature Climate Change* **2**, 177–181. doi: 10.1038/nclimate1346, ISSN: 1758-678X.
- Anenberg S.C., K. Balakrishnan, J. Jetter, O. Masera, S. Mehta, J. Moss, and V. Ramanathan (2013). Cleaner Cooking Solutions to Achieve Health, Climate, and Economic Cobenefits. *Environmental Science & Technology* **47**, 3944–3952. doi: 10.1021/es304942e, ISSN: 0013-936X.
- Angassa A., and G. Oba (2008). Effects of management and time on mechanisms of bush encroachment in southern Ethiopia. *African Journal of Ecology* **46**, 186–196. doi: 10.1111/j.1365-2028.2007.00832.x, ISSN: 1365-2028.
- Angelsen A. (2008). *Moving Ahead with REDD: Issues, Options and Implications*. CIFOR, Bogor Barat, Indonesia, 172 pp. ISBN: 9789791412766.
- Angelsen A., C. Streck, L. Peskett, J. Brown, and C. Luttrell (2008). *What Is the Right Scale for REDD? The Implications of National, Subnational and Nested Approaches*. CIFOR, Bogor Barat, Indonesia, 6 pp. Available at: http://62.225.2.55/files/methods_science/redd/application/pdf/what_is_the_right_scale_for_redd.pdf.
- Angst T.E., C.J. Patterson, D.S. Reay, P. Anderson, T.A. Peshkur, and S.P. Sohi (2013). Biochar Diminishes Nitrous Oxide and Nitrate Leaching from Diverse Nutrient Sources. *Journal of Environment Quality* **42**, 672. doi: 10.2134/jeq2012.0341, ISSN: 0047-2425.
- Araujo C., C.A. Bonjean, J.-L. Combes, P. Combes Motel, and E.J. Reis (2009). Property rights and deforestation in the Brazilian Amazon. *Ecological Economics* **68**, 2461–2468. doi: 10.1016/j.ecolecon.2008.12.015, ISSN: 0921-8009.
- Arcidiacono-Bársony C., P. Ciais, N. Viovy, and N. Vuichard (2011). REDD Mitigation. *Procedia Environmental Sciences* **6**, 50–59. doi: 10.1016/j.proenv.2011.05.006, ISSN: 18780296.
- Argo A.M., E.C. Tan, D. Inman, M.H. Langholtz, L.M. Eaton, J.J. Jacobson, C.T. Wright, D.J. Muth, M.M. Wu, Y.-W. Chiu, and R.L. Graham (2013). Investigation of biochemical biorefinery sizing and environmental sustainability impacts for conventional bale system and advanced uniform biomass logistics designs. *Biofuels, Bioproducts and Biorefining* **7**, 282–302. doi: 10.1002/bbb.1391, ISSN: 1932-1031.
- Arnalds A. (2004). Carbon sequestration and the restoration of land health: an example from Iceland. *Climatic Change* **65**, 333–346.
- Arndt C., R. Benfica, and J. Thurlow (2011). Gender Implications of Biofuels Expansion in Africa: The Case of Mozambique. *World Development* **39**, 1649–1662. doi: 10.1016/j.worlddev.2011.02.012, ISSN: 0305-750X.
- Arndt C., K. Pauw, and J. Thurlow (2012). Biofuels and economic development: A computable general equilibrium analysis for Tanzania. *Energy Economics* **34**, 1922–1930. doi: 10.1016/j.eneco.2012.07.020, ISSN: 0140-9883.
- Arora V.K., and A. Montenegro (2011). Small temperature benefits provided by realistic afforestation efforts. *Nature Geoscience* **4**, 514–518. doi: 10.1038/ngeo1182, ISSN: 1752-0894.

- Asante P., and G.W. Armstrong (2012). Optimal forest harvest age considering carbon sequestration in multiple carbon pools: A comparative statics analysis. *Journal of Forest Economics* **18**, 145–156. doi: 10.1016/j.jfe.2011.12.002, ISSN: 1104-6899.
- Asante P., G.W. Armstrong, and W.L. Adamowicz (2011). Carbon sequestration and the optimal forest harvest decision: A dynamic programming approach considering biomass and dead organic matter. *Journal of Forest Economics* **17**, 3–17. doi: 10.1016/j.jfe.2010.07.001, ISSN: 1104-6899.
- Assogbadjo A.E., R.G. Kakaï, F.G. Vodouhê, C.A.M.S. Djagoun, J.T.C. Codjia, and B. Sinsin (2012). Biodiversity and socioeconomic factors supporting farmers' choice of wild edible trees in the agroforestry systems of Benin (West Africa). *Forest Policy and Economics* **14**, 41–49.
- Attwood G.T., E. Altermann, W.J. Kelly, S.C. Leahy, L. Zhang, and M. Morrison (2011). Exploring rumen methanogen genomes to identify targets for methane mitigation strategies. *Animal Feed Science and Technology* **166–167**, 65–75. doi: 10.1016/j.anifeedsci.2011.04.004, ISSN: 0377-8401.
- Attwood G.T., and C.S. McSweeney (2008). Methanogen genomics to discover targets for methane mitigation technologies and options for alternative H₂ utilisation in the rumen. *Australian Journal of Experimental Agriculture* **48**, 28–37.
- Auld G., L.H. Gulbrandsen, and C.L. McDemott (2008). Certification Schemes and the Impacts on Forests and Forestry. *Annual Review of Environment and Resources* **33**, 187–211. doi: 10.1146/annurev.enviro.33.013007.103754.
- Awuodu I., and J. Zhang (2012). Uncertainties and sustainability concepts in biofuel supply chain management: A review. *Renewable and Sustainable Energy Reviews* **16**, 1359–1368. doi: 10.1016/j.rser.2011.10.016, ISSN: 1364-0321.
- Baccini A., S.J. Goetz, W.S. Walker, N.T. Laporte, M. Sun, D. Sulla-Menashe, J. Hackler, P.S.A. Beck, R. Dubayah, M.A. Friedl, S. Samanta, and R.A. Houghton (2012). Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps. *Nature Climate Change* **2**, 182–185. doi: 10.1038/nclimate1354, ISSN: 1758-678X.
- Bacchetti J., S.G. Garcia, A. Mena, and M. Fiala (2012). Life cycle assessment: an application to poplar for energy cultivated in Italy. *Journal of Agricultural Engineering* **43**, 72–78.
- Bacovsky D., M. Dallos, and M. Wörgetter (2010). *Status of 2nd Generation Biofuels Demonstration Facilities in June 2010: A Report to IEA Bioenergy Task 39*. Available at: <http://library.certh.gr/libfiles/PDF/GEN-SPIN-711-STATUS-by-BACOVSKY-in-RPT-BIOENERGY-TASK-39-PP-126-Y-JUL-2010.pdf>.
- Bacovsky D., L. Nikolaus, O. Monica, and W. Manfred (2013). *Status of Advanced Biofuels Demonstration Facilities in 2012*. International Energy Agency Bioenergy, 209 pp. Available at: http://demoplants.bioenergy2020.eu/files/Demoplants_Report_Final.pdf.
- Bai Y., L. Luo, and E. Voet (2010). Life cycle assessment of switchgrass-derived ethanol as transport fuel. *The International Journal of Life Cycle Assessment* **15**, 468–477. doi: 10.1007/s11367-010-0177-2, ISSN: 0948-3349.
- Bailis R., and H. McCarthy (2011). Carbon impacts of direct land use change in semiarid woodlands converted to biofuel plantations in India and Brazil. *GCB Bioenergy* **3**, 449–460. doi: 10.1111/j.1757-1707.2011.01100.x, ISSN: 1757-1707.
- Baker J.M., T.E. Ochsner, R.T. Venterea, and T.J. Griffis (2007). Tillage and soil carbon sequestration—What do we really know? *Agriculture, Ecosystems & Environment* **118**, 1–5.
- Baker D.J., G. Richards, A. Grainger, P. Gonzalez, S. Brown, R. DeFries, A. Held, J. Kellendorfer, P. Ndunda, D. Ojima, P.-E. Skrovseth, C. Souza Jr., and F. Stolle (2010). Achieving forest carbon information with higher certainty: A five-part plan. *Environmental Science & Policy* **13**, 249–260. doi: 10.1016/j.envsci.2010.03.004, ISSN: 1462-9011.
- Bala G., K. Caldeira, M. Wickett, T.J. Phillips, D.B. Lobell, C. Delire, and A. Mirin (2007). Combined climate and carbon-cycle effects of large-scale deforestation. *Proceedings of the National Academy of Sciences* **104**, 6550–6555. doi: 10.1073/pnas.0608998104.
- Balan V., D. Chiamonti, and S. Kumar (2013). Review of US and EU initiatives toward development, demonstration, and commercialization of lignocellulosic biofuels. *Biofuels, Bioproducts and Biorefining* **7**, 732–759.
- Baliban R.C., J.A. Elia, and C.A. Floudas (2013). Biomass and natural gas to liquid transportation fuels: process synthesis, global optimization, and topology analysis. *Industrial & Engineering Chemistry Research* **52**, 3381–3406.
- Barbier E.B. (2007). Valuing ecosystem services as productive inputs. *Economic Policy* **22**, 177–229.
- Barrow C.J. (2012). Biochar: Potential for countering land degradation and for improving agriculture. *Applied Geography* **34**, 21–28. doi: 10.1016/j.apgeog.2011.09.008, ISSN: 0143-6228.
- Barton L., K. Butterbach-Bahl, R. Kiese, and D.V. Murphy (2011). Nitrous oxide fluxes from a grain-legume crop (narrow-leaved lupin) grown in a semi-arid climate: Soil N₂O from a grain-legume crop. *Global Change Biology* **17**, 1153–1166. doi: 10.1111/j.1365-2486.2010.02260.x, ISSN: 13541013.
- Bathiany S., M. Claussen, V. Brovkin, T. Raddatz, and V. Gayler (2010). Combined biogeophysical and biogeochemical effects of large-scale forest cover changes in the MPI earth system model. *Biogeosciences* **7**, 1383–1399. doi: 10.5194/bg-7-1383-2010, ISSN: 1726-4189.
- Batidzirai B., E. Smeets, and A. Faaij (2012). Harmonising bioenergy resource potentials—Methodological lessons from review of state of the art bioenergy potential assessments. *Renewable and Sustainable Energy Reviews* **16**, 6598–6630.
- Batjes N.H. (2004). Estimation of soil carbon gains upon improved management within croplands and grasslands of Africa. *Environment, Development and Sustainability* **6**, 133–143.
- Baum S., A. Bolte, and M. Weih (2012). Short rotation coppice (SRC) plantations provide additional habitats for vascular plant species in agricultural mosaic landscapes. *Bioenergy Research* **5**, 573–583.
- Bayala J., L.K. Heng, M. van Noordwijk, and S.J. Ouedraogo (2008). Hydraulic redistribution study in two native tree species of agroforestry parklands of West African dry savanna. *Acta Oecologica* **34**, 370–378. Available at: <http://www.sciencedirect.com/science/article/pii/S1146609X08001033>.
- Bayala J., A. Kalinganire, Z. Tchoundjeu, P. Sinclair, and D. Garrity (2011). *Conservation Agriculture with Trees in the West African Sahel: A Review*. World Agroforestry Centre, Bamako, Mali, 72 pp. ISBN: 978-92-9059-295-2.
- Beach R.H., A.J. Daigneault, B.A. Mccarl, and S. Rose (2009). *Modeling Alternative Policies for Forestry and Agricultural Bioenergy Production and GHG Mitigation*. Washington, D.C., 27 pp.
- Beauchemin K.A., M. Kreuzer, F. O'Mara, and T.A. McAllister (2008). Nutritional management for enteric methane abatement: a review. *Australian Journal of Experimental Agriculture* **48**, 21–27. doi: 10.1071/EA07199, ISSN: 0816-1089.

- Beauchemin K.A., T.A. McAllister, and S.M. McGinn (2009).** Dietary mitigation of enteric methane from cattle. *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources* **4**, 1–18. Available at: <http://www.cabi.org/cabreviews/?loadmodule=review&page=4051&reviewid=112308&Site=167>.
- Beckham G.T., Z. Dai, J.F. Matthews, M. Momany, C.M. Payne, W.S. Adney, S.E. Baker, and M.E. Himmel (2012).** Harnessing glycosylation to improve cellulase activity. *Energy Biotechnology • Environmental Biotechnology* **23**, 338–345. doi: 10.1016/j.copbio.2011.11.030, ISSN: 0958-1669.
- Ballarby J., R. Tirado, A. Leip, F. Weiss, J.P. Lesschen, and P. Smith (2012).** Livestock greenhouse gas emissions and mitigation potential in Europe. *Global Change Biology*, n/a–n/a. doi: 10.1111/j.1365-2486.2012.02786.x, ISSN: 1365-2486.
- Benayas J.M.R., A.C. Newton, A. Diaz, and J.M. Bullock (2009).** Enhancement of Biodiversity and Ecosystem Services by Ecological Restoration: A Meta-Analysis. *Science* **325**, 1121–1124. doi: 10.1126/science.1172460, ISSN: 0036-8075, 1095–9203.
- Benítez P.C., I. McCallum, M. Obersteiner, and Y. Yamagata (2007).** Global potential for carbon sequestration: Geographical distribution, country risk and policy implications. *Ecological Economics* **60**, 572–583. doi: 10.1016/j.ecolecon.2005.12.015, ISSN: 0921-8009.
- Bennetzen E.H., P. Smith, J.-F. Soussana, and J.R. Porter (2012).** Identity-based estimation of greenhouse gas emissions from crop production: Case study from Denmark. *European Journal of Agronomy* **41**, 66–72. doi: 10.1016/j.eja.2012.03.010, ISSN: 11610301.
- Beringer T., W. Lucht, and S. Schaphoff (2011).** Bioenergy production potential of global biomass plantations under environmental and agricultural constraints. *GCB Bioenergy* **3**, 299–312. doi: 10.1111/j.1757-1707.2010.01088.x, ISSN: 1757-1707.
- Berndes G. (2012).** Bioenergy's contribution to climate change mitigation—a matter of perspectives. *Biofuels, Bioproducts and Biorefining* **6**, 233–235. doi: 10.1002/bbb.1343, ISSN: 1932-1031.
- Berndes G., S. Ahlgren, P. Börjesson, and A.L. Cowie (2013).** Bioenergy and land use change—state of the art. *Wiley Interdisciplinary Reviews: Energy and Environment* **2**, 282–303. doi: 10.1002/wene.41, ISSN: 2041-840X.
- Berndes G., N. Bird, and A. Cowie (2011).** *Bioenergy, Land Use Change and Climate Change Mitigation. Technical Report.* International Energy Agency, Available at: <http://www.ieabioenergy.com/LibItem.aspx?id=6927>.
- Berndes G., and P. Börjesson (2007).** *Multifunctional Bioenergy Systems. The AGS Pathways Report 2007: EU1.* AGS—Alliance for Global Sustainability: Swiss Federal Institute of Technology, Massachusetts Institute of Technology, Chalmers University of Technology, Tokyo University, 26 pp.
- Berndes G., P. Börjesson, M. Ostwald, and M. Palm (2008).** Multifunctional biomass production systems—an introduction with presentation of specific applications in India and Sweden. *Biofuels, Bioproducts and Biorefining* **2**, 16–25.
- Berndes G., F. Fredriksson, and P. Börjesson (2004).** Cadmium accumulation and Salix based phytoextraction on arable land in Sweden. *Agriculture, Ecosystems & Environment* **103**, 207–223.
- Berners-Lee M., C. Hoolohan, H. Cammack, and C.N. Hewitt (2012).** The relative greenhouse gas impacts of realistic dietary choices. *Energy Policy* **43**, 184–190. doi: 10.1016/j.enpol.2011.12.054, ISSN: 0301-4215.
- Bernier P., and D. Paré (2013).** Using ecosystem CO₂ measurements to estimate the timing and magnitude of greenhouse gas mitigation potential of forest bioenergy. *GCB Bioenergy* **5**, 67–72.
- Betts R.A. (2001).** Biogeophysical impacts of land use on present-day climate: near-surface temperature change and radiative forcing. *Atmospheric Science Letters* **2**, 39–51. doi: 10.1006/asle.2001.0037, ISSN: 1530-261X.
- Betts R.A. (2011).** Mitigation: A sweetener for biofuels. *Nature Clim. Change* **1**, 99–101. ISSN: 1758-678X.
- Betts R.A., P.D. Falloon, K.K. Goldewijk, and N. Ramankutty (2007).** Biogeophysical effects of land use on climate: Model simulations of radiative forcing and large-scale temperature change. *Agricultural and Forest Meteorology* **142**, 216–233. doi: 10.1016/j.agrformet.2006.08.021, ISSN: 0168-1923.
- Bhattacharyya P., D.P. Sinhababu, K.S. Roy, P.K. Dash, P.K. Sahu, R. Dandapat, S. Neogi, and S. Mohanty (2013).** Effect of fish species on methane and nitrous oxide emission in relation to soil C, N pools and enzymatic activities in rainfed shallow lowland rice-fish farming system. *Agriculture, Ecosystems & Environment* **176**, 53–62. doi: 10.1016/j.agee.2013.05.015, ISSN: 01678809.
- Bhuiyan M.A.H., R. Islam, C. Siwar, and S.M. Ismail (2010).** Educational Tourism and Forest Conservation: Diversification for Child Education. *Procedia—Social and Behavioral Sciences* **7**, 19–23. doi: 10.1016/j.sbspro.2010.10.003, ISSN: 1877-0428.
- Biederman L.A., and W.S. Harpole (2013).** Biochar and its effects on plant productivity and nutrient cycling: a meta-analysis. *GCB Bioenergy* **5**, 202–214. doi: 10.1111/gcbb.12037, ISSN: 17571693.
- Blanco-Canqui H., and R. Lal (2009).** Crop Residue Removal Impacts on Soil Productivity and Environmental Quality. *Critical Reviews in Plant Sciences* **28**, 139–163. doi: 10.1080/07352680902776507, ISSN: 0735-2689, 1549–7836.
- Blengini G.A., E. Brizio, M. Cibrario, and G. Genon (2011).** LCA of bioenergy chains in Piedmont (Italy): A case study to support public decision makers towards sustainability. *Resources, Conservation and Recycling* **57**, 36–47. doi: 10.1016/j.resconrec.2011.10.003, ISSN: 0921-3449.
- Blom B., T. Sunderland, and D. Murdiyarso (2010).** Getting REDD to work locally: lessons learned from integrated conservation and development projects. *Environmental Science & Policy* **13**, 164–172.
- Von Blottnitz H., and M.A. Curran (2007).** A review of assessments conducted on bio-ethanol as a transportation fuel from a net energy, greenhouse gas, and environmental life cycle perspective. *Journal of Cleaner Production* **15**, 607–619. doi: 10.1016/j.jclepro.2006.03.002, ISSN: 0959-6526.
- Bockel L., A. Gentien, M. Tinlot, and M. Bromhead (2010).** *From Nationally Appropriate Mitigation Actions (NAMAs) to Low-Carbon Development in Agriculture: NAMAs as Pathway at Country Level.* Food and Agricultural Organization, Rome, Italy, 32 pp. Available at: [http://www.nama-database.org/index.php/From_Nationally_Appropriate_Mitigation_Actions_\(NAMAs\)_to_Low-Carbon_Development_in_Agriculture](http://www.nama-database.org/index.php/From_Nationally_Appropriate_Mitigation_Actions_(NAMAs)_to_Low-Carbon_Development_in_Agriculture).
- Van Bodegom J.A., A. Jan, H. Savenije, and M. Wit (Eds.) (2009).** *Forests and Climate Change: Adaptation and Mitigation.* Tropenbos International, Wageningen, The Netherlands, 160 pp. ISBN: 9789051131000.
- Boden T., G. Marland, and R. Andres (2011).** Global CO₂ Emissions from Fossil-Fuel Burning, Cement Manufacture, and Gas Flaring: 1751–2008. Available at: http://cdiac.ornl.gov/trends/emis/meth_reg.html.

- De Boer I., C. Cederberg, S. Eady, S. Gollnow, T. Kristensen, M. Macleod, M. Meul, T. Nemecek, L. Phong, G. Thoma, H. van der Werf, A. Williams, and M. Zonderland-Thomassen (2011). Greenhouse gas mitigation in animal production: towards an integrated life cycle sustainability assessment. *Carbon and Nitrogen Cycles* **3**, 423–431. doi: 10.1016/j.cosust.2011.08.007, ISSN: 1877-3435.
- Bonan G.B. (2008). Forests and Climate Change: Forcings, Feedbacks, and the Climate Benefits of Forests. *Science* **320**, 1444–1449. doi: 10.1126/science.1155121, ISSN: 0036-8075, 1095–9203.
- Börjesson P., and G. Berndes (2006). The prospects for willow plantations for wastewater treatment in Sweden. *Biomass and Bioenergy* **30**, 428–438.
- Börjesson P., and L.M. Tufvesson (2011). Agricultural crop-based biofuels—resource efficiency and environmental performance including direct land use changes. *Journal of Cleaner Production* **19**, 108–120.
- Bostock J., B. McAndrew, R. Richards, K. Jauncey, T. Telfer, K. Lorenzen, D. Little, L. Ross, N. Handisyde, I. Gatward, and R. Corner (2010). Aquaculture: global status and trends. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 2897–2912. doi: 10.1098/rstb.2010.0170, ISSN: 0962-8436, 1471–2970.
- Böttcher H., K. Eisbrenner, S. Fritz, G. Kindermann, F. Kraxner, I. McCallum, and M. Obersteiner (2009). An assessment of monitoring requirements and costs of “Reduced Emissions from Deforestation and Degradation.” *Carbon Balance and Management* **4**, 7. doi: 10.1186/1750-0680-4-7, ISSN: 1750-0680.
- Böttcher H., P.J. Verkerk, M. Gusti, P. Havlík, and G. Grassi (2012). Projection of the future EU forest CO₂ sink as affected by recent bioenergy policies using two advanced forest management models. *GCB Bioenergy* **4**, 773–783. doi: 10.1111/j.1757-1707.2011.01152.x, ISSN: 1757-1707.
- Boulamanti A.K., S. Donida Maglio, J. Giuntoli, and A. Agostini (2013). Influence of different practices on biogas sustainability. *Biomass and Bioenergy* **53**, 149–161. doi: 10.1016/j.biombioe.2013.02.020, ISSN: 0961-9534.
- Boutton T.W., J.D. Liao, T.R. Filley, and S.R. Archer (2009). Belowground Carbon Storage and Dynamics Accompanying Woody Plant Encroachment in a Sub-tropical Savanna. SSSA Special Publications. In: *Soil Carbon Sequestration and the Greenhouse Effect*. Soil Science Society of America, Madison, WI, USA, pp. 181–205.
- Bradley R.S., M. Vuille, H.F. Diaz, and W. Vergara (2006). Threats to Water Supplies in the Tropical Andes. *Science* **312**, 1755–1756. doi: 10.1126/science.1128087.
- Bradstock R.A., G.J. Cary, I. Davies, D.B. Lindenmayer, O. Price, and R.J. Williams (2012). Wildfires, fuel treatment and risk mitigation in Australian eucalypt forests: insights from landscape-scale simulation. *Journal of Environmental Management* **105**, 66–75.
- Branca G., H. Hissa, M.C. Benez, K. Medeiros, L. Lipper, M. Tinlot, L. Bockel, and M. Bernoux (2013). Capturing synergies between rural development and agricultural mitigation in Brazil. *Land Use Policy* **30**, 507–518. doi: 10.1016/j.landusepol.2012.04.021, ISSN: 0264-8377.
- Brandão M., L. Milà i Canals, and R. Clift (2011). Soil organic carbon changes in the cultivation of energy crops: Implications for GHG balances and soil quality for use in LCA. *Biomass and Bioenergy* **35**, 2323–2336. doi: 10.1016/j.biombioe.2009.10.019, ISSN: 0961-9534.
- Bridgwater A.V. (2012). Upgrading biomass fast pyrolysis liquids. *Environmental Progress & Sustainable Energy* **31**, 261–268. doi: 10.1002/ep.11635, ISSN: 1944-7450.
- Bright R.M., and A.H. Strømman (2009). Life Cycle Assessment of Second Generation Bio-ethanol Produced from Scandinavian Boreal Forest Resources: A Regional Analysis for Middle Norway. *Journal of Industrial Ecology* **13**, 514–531.
- Bright R.M., and A.H. Strømman (2010). Fuel-mix, Vehicle technologies, and Transport Demand Affect Prospects for Biofuels in Northern Europe. *Environmental Science & Technology* **44**, 2261–2269.
- Bright R.M., A.H. Strømman, and T.R. Hawkins (2010). Environmental Assessment of Wood-based Biofuel Production and Consumption Scenarios in Norway. *Journal of Industrial Ecology* **14**, 422–439.
- Bright R.M., A.H. Strømman, and G.P. Peters (2011). Radiative Forcing Impacts of Boreal Forest Biofuels: A Scenario Study for Norway in Light of Albedo. *Environmental Science & Technology* **45**, 7570–7580. doi: 10.1021/es201746b, ISSN: 0013-936X.
- Bringezu S., M. O'Brien, and H. Schütz (2012). Beyond biofuels: Assessing global land use for domestic consumption of biomass: A conceptual and empirical contribution to sustainable management of global resources. *Land Use Policy* **29**, 224–232. doi: 16/j.landusepol.2011.06.010, ISSN: 0264-8377.
- Brockhaus M., M. Di Gregorio, and S. Mardiah (2013). Governing the design of national REDD +: An analysis of the power of agency. *Forest Policy and Economics*. doi: 10.1016/j.forpol.2013.07.003, ISSN: 1389-9341.
- Brown E.G., R.C. Anderson, G.E. Carstens, H. Gutierrez-Banuelos, J.L. McReynolds, L.J. Slay, T.R. Callaway, and D.J. Nisbet (2011). Effects of oral nitroethane administration on enteric methane emissions and ruminal fermentation in cattle. *Animal Feed Science and Technology* **166–67**, 275–281. doi: 10.1016/j.anifeedsci.2011.04.017, ISSN: 0377-8401.
- Brown C.D., and J.F. Johnstone (2011). How does increased fire frequency affect carbon loss from fire? A case study in the northern boreal forest. *International Journal of Wildland Fire* **20**, 829–837. doi: 10.1071/WF10113.
- Bryan E., C. Ringler, B. Okaba, J. Koo, M. Herrero, and S. Silvestri (2013). Can agriculture support climate change adaptation, greenhouse gas mitigation and rural livelihoods? Insights from Kenya. *Climatic Change* **118**, 151–165. doi: 10.1007/s10584-012-0640-0, ISSN: 1473300800.
- Bryant J.R., V.O. Snow, R. Cichota, and B.H. Jolly (2011). The effect of situational variability in climate and soil, choice of animal type and N fertilisation level on nitrogen leaching from pastoral farming systems around Lake Taupo, New Zealand. *Agricultural Systems* **104**, 271–280. doi: 10.1016/j.agsy.2010.11.001, ISSN: 0308-521X.
- Budsberg E., M. Rastogi, M.E. Puettmann, J. Caputo, S. Balogh, T.A. Volk, R. Gustafson, and L. Johnson (2012). Life-cycle assessment for the production of bioethanol from willow biomass crops via biochemical conversion. *Forest Products Journal* **62**, 305.
- Buonocore E., P.P. Franzese, and S. Ulgiati (2012). Assessing the environmental performance and sustainability of bioenergy production in Sweden: A life cycle assessment perspective. *Energy* **37**, 69–78. doi: 10.1016/j.energy.2011.07.032.
- Burney J.A., S.J. Davis, and D.B. Lobell (2010). Greenhouse gas mitigation by agricultural intensification. *Proceedings of the National Academy of Sciences* **107**, 12052–12057. doi: 10.1073/pnas.0914216107.
- Busch G. (2012). GIS-based tools for regional assessments and planning processes regarding potential environmental effects of poplar SRC. *Bioenergy Research* **5**, 584–605.

- Busch J., F. Godoy, W.R. Turner, and C.A. Harvey (2011). Biodiversity co-benefits of reducing emissions from deforestation under alternative reference levels and levels of finance. *Conservation Letters* 4, 101–115. doi: 10.1111/j.1755-263X.2010.00150.x, ISSN: 1755-263X.
- Bustamante M.M.C., C.A. Nobre, R. Smeraldi, A.P.D. Aguiar, L.G. Barioni, L.G. Ferreira, K. Longo, P. May, A.S. Pinto, and J.P.H.B. Ometto (2012). Estimating greenhouse gas emissions from cattle raising in Brazil. *Climatic Change*, 1–19. Available at: <http://www.springerlink.com/index/WT2702446216702X.pdf>.
- Cacciatore M.A., D.A. Scheufele, and B.R. Shaw (2012). Labeling renewable energies: How the language surrounding biofuels can influence its public acceptance. *Energy Policy* 51, 673–682. doi: 10.1016/j.enpol.2012.09.005, ISSN: 0301-4215.
- Cai Y.-I. (2001). A study on land use/cover change: the need for a new integrated approach. *Geographical Research*. Available at: http://en.cnki.com.cn/Article_en/CJFDTotal-DLYJ200106000.htm.
- Calegari A., W.L. Hargrove, D. dos S. Rheinheimer, R. Ralisch, D. Tessier, S. Tournonnet, and M. de F. Guimaraes (2008). Impact of Long-Term No-Tillage and Cropping System Management on Soil Organic Carbon in an Oxisol: A Model for Sustainability. *Agronomy Journal* 100, 1013–1019.
- Calfapietra C., B. Gielen, D. Karnosky, R. Ceulemans, and G. Scarascia Mugnozza (2010). Response and potential of agroforestry crops under global change. *Environmental Pollution* 158, 1095–1104. doi: 10.1016/j.envpol.2009.09.008, ISSN: 0269-7491.
- California Environmental Protection Agency Air Resources Board—Homepage. Available at: <http://www.arb.ca.gov/homepage.htm>.
- Calvin K., P. Patel, A. Fawcett, L. Clarke, K. Fisher-Vanden, J. Edmonds, S.H. Kim, R. Sands, and M. Wise (2009). The distribution and magnitude of emissions mitigation costs in climate stabilization under less than perfect international cooperation: SGM results. *Energy Economics* 31, S187–S197. doi: 10.1016/j.eneco.2009.06.014, ISSN: 01409883.
- Calvin K., M. Wise, P. Kyle, P. Patel, L. Clarke, and J. Edmonds (2013a). Trade-offs of different land and bioenergy policies on the path to achieving climate targets. *Climatic Change* in press. doi: DOI: 10.1007/s10584-013-0897-y.
- Calvin K., M. Wise, P. Luckow, P. Kyle, L. Clarke, and J. Edmonds (2013b). Implications of uncertain future fossil energy resources on bioenergy use and terrestrial carbon emissions. *Climatic Change*, 1–12. doi: 10.1007/s10584-013-0923-0, ISSN: 0165-0009.
- Campbell J.E., D.B. Lobell, and C.B. Field (2009). Greater Transportation Energy and GHG Offsets from Bioelectricity Than Ethanol. *Science* 324, 1055–1057. Available at: <http://www.sciencemag.org/content/324/5930/1055.abstract>.
- Campbell J.E., D.B. Lobell, R.C. Genova, and C.B. Field (2008). The Global Potential of Bioenergy on Abandoned Agriculture Lands. *Environmental Science & Technology* 42, 5791–5794. doi: 10.1021/es800052w, ISSN: 0013-936X.
- Canadell J.G., and M.R. Raupach (2008). Managing Forests for Climate Change Mitigation. *Science* 320, 1456–1457. doi: 10.1126/science.1155458, ISSN: 0036-8075, 1095–9203.
- Cançado J.E.D., P.H.N. Saldiva, L.A.A. Pereira, L.B.L.S. Lara, P. Artaxo, L.A. Martinelli, M.A. Arbex, A. Zanobetti, and A.L.F. Braga (2006). The impact of sugar cane-burning emissions on the respiratory system of children and the elderly. *Environmental Health Perspectives* 114, 725–729. ISSN: 0091-6765.
- Carlsson-Kanyama A., and A.D. González (2009). Potential contributions of food consumption patterns to climate change. *The American Journal of Clinical Nutrition* 89, 1704S–1709S. doi: 10.3945/ajcn.2009.26736AA, ISSN: 0002-9165, 1938–3207.
- Carpita N.C. (2012). Progress in the biological synthesis of the plant cell wall: new ideas for improving biomass for bioenergy. *Current Opinion in Biotechnology* 23, 330–337. doi: 10.1016/j.copbio.2011.12.003, ISSN: 0958-1669.
- Cattaneo A., R. Lubowski, J. Busch, A. Creed, B. Strassburg, F. Boltz, and R. Ashton (2010). On international equity in reducing emissions from deforestation. *Environmental Science & Policy* 13, 742–753. doi: 10.1016/j.envsci.2010.08.009, ISSN: 1462-9011.
- Cayuela M.L., M.A. Sánchez-Monedero, A. Roig, K. Hanley, A. Enders, and J. Lehmann (2013). Biochar and denitrification in soils: when, how much and why does biochar reduce N₂O emissions? *Scientific Reports* 3. doi: 10.1038/srep01732, ISSN: 2045-2322.
- CBD, and GiZ (2011). *Biodiversity and Livelihoods: REDD-plus Benefits*. Secretariat of the Convention on Biological Diversity and Deutsche Gesellschaft Für Internationale Zusammenarbeit (GIZ) GmbH, Canada, 41 pp. Available at: <https://www.cbd.int/doc/publications/for-redd-en.pdf>.
- CDM-SSC WG (2011). *Thirty-Third Meeting Report, Annex 8: Call for Public Inputs in Relation to Standardized Approaches for Facilitating the Baseline Emission Calculations under SSC CDM Methodologies for Displacement of Non-Renewable Biomass*. UNFCCC, Bonn, Germany, 52 pp. Available at: http://cdm.unfccc.int/Panels/ssc_wg/meetings/033/ssc_033_an08.pdf.
- Cederberg C., U.M. Persson, K. Neovius, S. Molander, and R. Clift (2011). Including carbon emissions from deforestation in the carbon footprint of Brazilian beef. *Environmental Science & Technology* 45, 1773–1779. Available at: <http://pubs.acs.org/doi/abs/10.1021/es103240z>.
- Ceschia E., P. Béziat, J.F. Dejoux, M. Aubinet, C. Bernhofer, B. Bodson, N. Buchmann, A. Carrara, P. Cellier, P. Di Tommasi, and others (2010). Management effects on net ecosystem carbon and GHG budgets at European crop sites. *Agriculture, Ecosystems & Environment* 139, 363–383.
- Chadwick D., S. Sommer, R. Thorman, D. Figueiro, L. Cardenas, B. Amon, and T. Misselbrook (2011). Manure management: Implications for greenhouse gas emissions. *Animal Feed Science and Technology* 166–67, 514–531. doi: 10.1016/j.anifeedsci.2011.04.036, ISSN: 0377-8401.
- Chakrabarti S. (2010). Factors influencing organic food purchase in India—expert survey insights. *British Food Journal* 112, 902–915. doi: 10.1108/00070701011067497, ISSN: 0007-070X.
- Chatterjee A., and R. Lal (2009). On farm assessment of tillage impact on soil carbon and associated soil quality parameters. *Soil and Tillage Research* 104, 270–277. doi: 10.1016/j.still.2009.03.006, ISSN: 0167-1987.
- Chen M., and R.E. Blankenship (2011). Expanding the solar spectrum used by photosynthesis. *Trends in Plant Science* 16, 427–431.
- Chen X., M. Khanna, H. Huang, and H. Önal (2012). Meeting the Mandate for Biofuels: Implications for Land Use, Food, and Fuel Prices.
- Cherubini F., N.D. Bird, A. Cowie, G. Jungmeier, B. Schlamadinger, and S. Woess-Gallasch (2009). Energy- and greenhouse gas-based LCA of biofuel and bioenergy systems: Key issues, ranges and recommendations. *Resources, Conservation, and Recycling* 53, 434–447. doi: 10.1016/j.resconrec.2009.03.013, ISSN: 0921-3449.

- Cherubini F., R.M. Bright, and A.H. Stromman (2013).** Global climate impacts of forest bioenergy: what, when and how to measure? *Environmental Research Letters* in press.
- Cherubini F., G. Guest, and A. Strømman (2012).** Application of probability distributions to the modeling of biogenic CO₂ fluxes in life cycle assessment. *GCB Bioenergy* **4**, 784–798. doi: 10.1111/j.1757-1707.2011.01156.x, ISSN: 1757-1707.
- Cherubini F., and G. Jungmeier (2010).** LCA of a biorefinery concept producing bioethanol, bioenergy, and chemicals from switchgrass. *The International Journal of Life Cycle Assessment* **15**, 53–66. doi: 10.1007/s11367-009-0124-2, ISSN: 0948-3349.
- Cherubini F., G.P. Peters, T. Berntsen, A.H. Strømman, and E. Hertwich (2011).** CO₂ emissions from biomass combustion for bioenergy: atmospheric decay and contribution to global warming. *GCB Bioenergy* **3**, 413–426. doi: 10.1111/j.1757-1707.2011.01102.x, ISSN: 1757-1707.
- Cherubini F., and S. Ulgiati (2010).** Crop residues as raw materials for biorefinery systems—A LCA case study. *Applied Energy* **87**, 47–57. doi: 10.1016/j.apenergy.2009.08.024, ISSN: 0306-2619.
- Chhatre A., and A. Agrawal (2009).** Trade-offs and synergies between carbon storage and livelihood benefits from forest commons. *Proceedings of the National Academy of Sciences* **106**, 17667–17670.
- Chhatre A., S. Lakhanpal, A.M. Larson, F. Nelson, H. Ojha, and J. Rao (2012).** Social safeguards and co-benefits in REDD+: a review of the adjacent possible. *Current Opinion in Environmental Sustainability* **4**, 654–660. doi: 10.1016/j.cosust.2012.08.006, ISSN: 1877-3435.
- Choo Y.M., H. Muhamad, Z. Hashim, V. Subramaniam, C.W. Puah, and Y. Tan (2011).** Determination of GHG contributions by subsystems in the oil palm supply chain using the LCA approach. *The International Journal of Life Cycle Assessment* **16**, 669–681. doi: 10.1007/s11367-011-0303-9, ISSN: 0948-3349.
- Chum H., A. Faaij, J. Moreira, G. Berndes, P. Dhamija, H. Dong, B. Gabrielle, A.G. Eng, W. Lucht, M. Mapako, O.M. Cerutti, T. McIntyre, T. Minowa, and K. Pingoud (2011).** Bioenergy. In: *IPCC Special Report on Renewable Energy Sources and Climate Change Mitigation* [O. Edenhofer, R. Pichs-Madruga, Y. Sokona, K. Seyboth, P. Matschoss, S. Kadner, T. Zwickel, P. Eickemeier, G. Hansen, S. Schlömer, C. von Stechow (eds)], Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 209–332.
- Chundawat S.P.S., G.T. Beckham, M.E. Himmel, and B.E. Dale (2011).** Deconstruction of Lignocellulosic Biomass to Fuels and Chemicals. *Annual Review of Chemical and Biomolecular Engineering* **2**, 121–145. doi: 10.1146/annurev-chembioeng-061010-114205.
- Chung C.E., V. Ramanathan, and D. Decremer (2012).** Observationally constrained estimates of carbonaceous aerosol radiative forcing. *Proceedings of the National Academy of Sciences* **109**, 11624–11629.
- Ciais P., M.J. Schelhaas, S. Zaehle, S.L. Piao, A. Cescatti, J. Liski, S. Luysaert, G. Le-Maire, E.-D. Schulze, O. Bouriaud, A. Freibauer, R. Valentini, and G.J. Nabuurs (2008).** Carbon accumulation in European forests. *Nature Geoscience* **1**, 425–429. doi: 10.1038/ngeo233, ISSN: 1752-0894.
- Clair T.A., P. Arp, T. Moore, M. Dalva, and F.-R. Meng (2002).** Gaseous carbon dioxide and methane, as well as dissolved organic carbon losses from a small temperate wetland under a changing climate. *Environmental Pollution* **116**, Supplement 1, 143–148. doi: 10.1016/S0269-7491(01)00267-6, ISSN: 0269-7491.
- Clair S.S., J. Hillier, and P. Smith (2008).** Estimating the pre-harvest greenhouse gas costs of energy crop production. *Biomass and Bioenergy* **32**, 442–452. doi: 10.1016/j.biombioe.2007.11.001, ISSN: 0961-9534.
- Clark H. (2013).** Nutritional and host effects on methanogenesis in the grazing ruminant. *Animal* **7** Suppl 1, 41–48. doi: 10.1017/S1751731112001875, ISSN: 1751-732X.
- Claussen M., V. Brovkin, and A. Ganopolski (2001).** Biogeophysical versus biogeochemical feedbacks of large-scale land cover change. *Geophys. Res. Lett.* **28**, 1011–1014. doi: 10.1029/2000gl012471, ISSN: 0094-8276.
- Clough T., L. Condrón, C. Kammann, and C. Müller (2013).** A Review of Biochar and Soil Nitrogen Dynamics. *Agronomy* **3**, 275–293. doi: 10.3390/agronomy3020275, ISSN: 2073-4395.
- Coelho S., O. Agbenyega, A. Agostini, K.H. Erb, H. Haberl, M. Hoogwijk, R. Lal, O. Lucon, O. Masera, and J.R. Moreira (2012).** Chapter 20—Land and Water: Linkages to Bioenergy. In: *Global Energy Assessment—Toward a Sustainable Future*. Cambridge University Press, Cambridge, UK and New York, NY, USA and the International Institute for Applied Systems Analysis, Laxenburg, Austria, Cambridge, UK, pp. 1459–1526. ISBN: 9780521182935.
- Colfer C.J.P. (2011).** Marginalized Forest Peoples' Perceptions of the Legitimacy of Governance: An Exploration. *World Development* **39**, 2147–2164. doi: 10.1016/j.worlddev.2011.04.012, ISSN: 0305-750X.
- Combes Motel P., R. Pirard, and J.-L. Combes (2009).** A methodology to estimate impacts of domestic policies on deforestation: Compensated Successful Efforts for “avoided deforestation” (REDD). *Ecological Economics* **68**, 680–691. doi: 10.1016/j.ecolecon.2008.06.001, ISSN: 0921-8009.
- Conant R.T., M. Easter, K. Paustian, A. Swan, and S. Williams (2007).** Impacts of periodic tillage on soil C stocks: A synthesis. *Soil and Tillage Research* **95**, 1–10. doi: 10.1016/j.still.2006.12.006, ISSN: 01671987.
- Corbera E., and K. Brown (2008).** Building Institutions to Trade Ecosystem Services: Marketing Forest Carbon in Mexico. *World Development* **36**, 1956–1979. doi: 10.1016/j.worlddev.2007.09.010, ISSN: 0305-750X.
- Corbera E., and H. Schroeder (2011).** Governing and implementing REDD+. *Environmental Science & Policy* **14**, 89–99. doi: 10.1016/j.envsci.2010.11.002, ISSN: 1462-9011.
- Cordes L. (2011).** *Igniting Change: A Strategy for Universal Adoption of Clean Cookstoves and Fuels*. Global Alliance for Clean Cookstoves, Washington DC, USA, Available at: www.cleancookstoves.org.
- Cottle D.J., J.V. Nolan, and S.G. Wiedemann (2011).** Ruminant enteric methane mitigation: a review. *Animal Production Science* **51**, 491. doi: 10.1071/AN10163, ISSN: 1836-0939.
- Cotula L. (2012).** The international political economy of the global land rush: A critical appraisal of trends, scale, geography and drivers. *Journal of Peasant Studies* **39**, 649–680. doi: 10.1080/03066150.2012.674940, ISSN: 0306-6150.
- Cotula L., S. Vermeulen, R. Leonard, and J. Keeley (2009).** *Land Grab or Development Opportunity? Agricultural Investment and International Land Deals in Africa*. 130 pp.
- Cox P.M., D. Pearson, B.B. Booth, P. Friedlingstein, C. Huntingford, C.D. Jones, and C.M. Luke (2013).** Sensitivity of tropical carbon to climate change constrained by carbon dioxide variability. *Nature* **494**, 341–344. doi: 10.1038/nature11882, ISSN: 0028-0836.

- Creutzig F., E. Corbera, S. Bolwig, and C. Hunsberger (2013). Integrating place-specific livelihood and equity outcomes into global assessments of bioenergy deployment. *Environmental Research Letters* 8, 035047 (11pp). doi: 10.1088/1748-9326/8/3/035047, ISSN: 1748-9326.
- Creutzig F., A. Popp, R. Plevin, G. Luderer, J. Minx, and O. Edenhofer (2012a). Reconciling top-down and bottom-up modelling on future bioenergy deployment. *Nature Climate Change* 2, 320–327. doi: 10.1038/nclimate1416, ISSN: 1758-678X.
- Creutzig F., C. von Stechow, D. Klein, C. Hunsberger, N. Bauer, A. Popp, and O. Edenhofer (2012b). Can bioenergy assessments deliver? *Economics of Energy & Environmental Policy* 1. doi: 10.5547/2160-5890.1.2.5, ISSN: 21605882.
- Crutzen P.J., A.R. Mosier, K.A. Smith, and W. Winiwarter (2007). N₂O release from agro-biofuel production negates global warming reduction by replacing fossil fuels. *Atmospheric Chemistry and Physics Discussions* 7, 11191–11205. doi: 10.5194/acpd-7-11191-2007, ISSN: 1680-7375.
- Daigneault A., B. Sohngen, and R. Sedjo (2012). Economic Approach to Assess the Forest Carbon Implications of Biomass Energy. *Environmental Science & Technology* 46, 5664–5671. doi: 10.1021/es2030142, ISSN: 0013-936X.
- Dale B.E., M. S. Allen, M. Laser, and L. R. Lynd (2009). Protein feeds coproduction in biomass conversion to fuels and chemicals. *Biofuels, Bioproducts and Biorefining* 3, 219–230. doi: 10.1002/bbb.132, ISSN: 1932-1031.
- Dale V.H., and K.L. Kline (2013). Issues in using landscape indicators to assess land changes. *Ecological Indicators* 28, 91–99.
- Dale V.H., K.L. Kline, D. Perla, and A. Lucier (2013). Communicating about bioenergy sustainability. *Environmental Management* 51, 279–290.
- Dale V.H., R. Lowrance, P. Mulholland, and G.P. Robertson (2010). Bioenergy sustainability at the regional scale. *Ecology and Society* 15, 23. Available at: https://www.kbs.msu.edu/images/stories/docs/robertson/Dale_et_al._2010_SSSAJ.pdf.
- Van Dam J., A.P.C. Faaij, J. Hilbert, H. Petruzzi, and W.C. Turkenburg (2009a). Large-scale bioenergy production from soybeans and switchgrass in Argentina: Part A: Potential and economic feasibility for national and international markets. *Renewable and Sustainable Energy Reviews* 13, 1710–1733.
- Van Dam J., A.P. Faaij, J. Hilbert, H. Petruzzi, and W.C. Turkenburg (2009b). Large-scale bioenergy production from soybeans and switchgrass in Argentina: Part B. Environmental and socio-economic impacts on a regional level. *Renewable and Sustainable Energy Reviews* 13, 1679–1709.
- Van Dam J., M. Junginger, and A.P.C. Faaij (2010). From the global efforts on certification of bioenergy towards an integrated approach based on sustainable land use planning. *Renewable and Sustainable Energy Reviews* 14, 2445–2472. doi: 16/j.rser.2010.07.010, ISSN: 1364-0321.
- Danielsen F., H. Beukema, N.D. Burgess, F. Parish, C.A. Brühl, P.F. Donald, D. Murdiyarsa, B. Phalan, L. Reijnders, M. Struebig, and E.B. Fitzherbert (2009). Biofuel Plantations on Forested Lands: Double Jeopardy for Biodiversity and Climate. *Conservation Biology* 23, 348–358. doi: 10.1111/j.1523-1739.2008.01096.x, ISSN: 1523-1739.
- Datta A., D.R. Nayak, D.P. Sinhababu, and T.K. Adhya (2009). Methane and nitrous oxide emissions from an integrated rainfed rice–fish farming system of Eastern India. *Agriculture, Ecosystems & Environment* 129, 228–237. doi: 10.1016/j.agee.2008.09.003, ISSN: 01678809.
- Dauvergne P., and K.J. Neville (2010). Forests, food, and fuel in the tropics: the uneven social and ecological consequences of the emerging political economy of biofuels. *Journal of Peasant Studies* 37, 631–660.
- Davidson E.A. (2012). Representative concentration pathways and mitigation scenarios for nitrous oxide. *Environmental Research Letters* 7, 024005. doi: 10.1088/1748-9326/7/2/024005, ISSN: 1748-9326.
- Davidson E.A., and I.A. Janssens (2006). Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* 440, 165–173. doi: 10.1038/nature04514, ISSN: 0028-0836, 1476–4679.
- Davin E.L., N. de Noblet-Ducoudré, and P. Friedlingstein (2007). Impact of land cover change on surface climate: Relevance of the radiative forcing concept. *Geophys. Res. Lett.* 34, L13702. doi: 10.1029/2007gl029678, ISSN: 0094-8276.
- Davis R., A. Aden, and P.T. Pienkos (2011a). Techno-economic analysis of autotrophic microalgae for fuel production. *Applied Energy* 88, 3524–3531. Available at: <http://www.sciencedirect.com/science/article/pii/S0306261911002406>.
- Davis S.C., R.M. Boddey, B.J.R. Alves, A.L. Cowie, B.H. George, S.M. Ogle, P. Smith, M. van Noordwijk, and M.T. van Wijk (2013). Management swing potential for bioenergy crops. *GCB Bioenergy* 5, 623–638. doi: 10.1111/gcbb.12042, ISSN: 1757-1707.
- Davis S.C., W.J. Parton, S.J.D. Grosso, C. Keough, E. Marx, P.R. Adler, and E.H. DeLucia (2011b). Impact of second-generation biofuel agriculture on greenhouse-gas emissions in the corn-growing regions of the US. *Frontiers in Ecology and the Environment* 10, 69–74. doi: 10.1890/110003, ISSN: 1540-9295.
- Dawson T.P., S.T. Jackson, J.I. House, I.C. Prentice, and G.M. Mace (2011). Beyond Predictions: Biodiversity Conservation in a Changing Climate. *Science* 332, 53–58. doi: 10.1126/science.1200303, ISSN: 0036-8075, 1095–9203.
- Deal R.L., B. Cochran, and G. LaRocco (2012). Bundling of ecosystem services to increase forestland value and enhance sustainable forest management. *Forest Policy and Economics* 17, 69–76.
- Deal R.L., and R. White (2012). Integrating forest products with ecosystem services: A global perspective. *Forest Policy and Economics* 17, 1–2.
- DeAngelo B.J., F.C. de la Chesnaye, R.H. Beach, A. Sommer, and B.C. Murray (2006). Methane and Nitrous Oxide Mitigation in Agriculture. *The Energy Journal*, 89–108.
- DeCicco J. (2013). Biofuel's carbon balance: doubts, certainties and implications. *Climatic Change* 121, 801–814. doi: 10.1007/s10584-013-0927-9, ISSN: 0165-0009.
- DeFries R., and C. Rosenzweig (2010). Toward a whole-landscape approach for sustainable land use in the tropics. *Proceedings of the National Academy of Sciences* 107, 19627–19632. doi: 10.1073/pnas.1011163107.
- Delucchi M.A. (2010). Impacts of biofuels on climate change, water use, and land use. *Annals of the New York Academy of Sciences* 1195, 28–45. doi: 10.1111/j.1749-6632.2010.05457.x, ISSN: 1749-6632.
- Dezzeo, N., Chacon, and N. (2005). Carbon and nutrients loss in aboveground biomass a long a fire induced forest-savanna gradient in the Gran Sabana, Southern Venezuela. *Forest Ecology and Management* 209, 343–352.
- Dias A.C., and L. Arroja (2012). Environmental impacts of eucalypt and maritime pine wood production in Portugal. *Journal of Cleaner Production* 37, 368–376. doi: 10.1016/j.jclepro.2012.07.056, ISSN: 0959-6526.
- Díaz S., J. Fargione, F.S. Chapin, and D. Tilman (2006). Biodiversity loss threatens human well-being. *PLoS Biology* 4, e277.

- Diaz-Chavez R.A. (2011). Assessing biofuels: Aiming for sustainable development or complying with the market? *Energy Policy* **39**, 5763–5769. doi: 10.1016/j.enpol.2011.03.054, ISSN: 0301-4215.
- Diaz-Chavez R.A. (2012). Land use for integrated systems: A bioenergy perspective. *Environmental Development* **3**, 91–99. doi: 10.1016/j.envdev.2012.03.018, ISSN: 2211-4645.
- Dijkstra J., O. Oenema, and A. Bannink (2011). Dietary strategies to reducing N excretion from cattle: implications for methane emissions. *Current Opinion in Environmental Sustainability* **3**, 414–422. doi: 10.1016/j.cosust.2011.07.008, ISSN: 1877-3435.
- Dimitriou I., C. Baum, S. Baum, G. Busch, U. Schulz, J. Köhn, N. Lamersdorf, P. Leinweber, P. Aronsson, M. Weih, G. Berndes, and A. Bolte (2009). Short Rotation Coppice (SRC) cultivation and local impact on the environment. *Landbauforschung vTI Agriculture and Forestry Research* **3**, 159–162.
- Dimitriou I., C. Baum, S. Baum, G. Busch, U. Schulz, J. Köhn, N. Lamersdorf, P. Leinweber, P. Aronsson, M. Weih, G. Berndes, and A. Bolte (2011). Quantifying environmental effects of short rotation coppice (SRC) on biodiversity, soil and water. *IEA Bioenergy Task 43*.
- Ding X.Z., R.J. Long, M. Kreuzer, J.D. Mi, and B. Yang (2010). Methane emissions from yak (*Bos grunniens*) steers grazing or kept indoors and fed diets with varying forage: concentrate ratio during the cold season on the Qinghai-Tibetan Plateau. *Animal Feed Science and Technology* **162**, 91–98.
- DiPietro P., P. Balash, and M. Wallace (2012). A Note on Sources of CO₂ Supply for Enhanced-Oil-Recovery Operations. *SPE Economics & Management*, pp.69–74.
- Djomo S.N., O.E. Kasmoui, and R. Ceulemans (2011). Energy and greenhouse gas balance of bioenergy production from poplar and willow: a review. *GCB Bioenergy* **3**, 181–197. doi: 10.1111/j.1757-1707.2010.01073.x, ISSN: 1757-1707.
- Don A., B. Osborne, A. Hastings, U. Skiba, M.S. Carter, J. Drewer, H. Flessa, A. Freibauer, N. Hyvönen, M.B. Jones, G.J. Lanigan, Ü. Mander, A. Monti, S.N. Djomo, J. Valentine, K. Walter, W. Zegada-Lizarazu, and T. Zenone (2012). Land-use change to bioenergy production in Europe: implications for the greenhouse gas balance and soil carbon. *GCB Bioenergy* **4**, 372–391. doi: 10.1111/j.1757-1707.2011.01116.x, ISSN: 1757-1707.
- Donato D.C., J.B. Kauffman, D. Murdiyasar, S. Kurnianto, M. Stidham, and M. Kanninen (2011). Mangroves among the most carbon-rich forests in the tropics. *Nature Geoscience* **4**, 293–297. doi: 10.1038/ngeo1123, ISSN: 1752-0894.
- Dornburg V., D. van Vuuren, G. van de Ven, H. Langeveld, M. Meeusen, M. Banse, M. van Oorschot, J. Ros, G. Jan van den Born, H. Aiking, M. Londo, H. Mozaffarian, P. Verweij, E. Lysen, and A. Faaij (2010). Bioenergy revisited: Key factors in global potentials of bioenergy. *Energy and Environmental Science* **3**, 258–267. doi: 10.1039/b922422j, ISSN: 1754-5692.
- Douglas E., A. Beltrán-Przekurat, D. Niyogi, R. Pielke Sr, and C. Vörösmarty (2009). The impact of agricultural intensification and irrigation on land-atmosphere interactions and Indian monsoon precipitation—A mesoscale modeling perspective. *Global and Planetary Change* **67**, 117–128.
- Downie A., P. Munroe, A. Cowie, L. Van Zwieten, and D.M.S. Lau (2012). Biochar as a Geoenvironmental Climate Solution: Hazard Identification and Risk Management. *Critical Reviews in Environmental Science and Technology* **42**, 225–250. doi: 10.1080/10643389.2010.507980, ISSN: 1064-3389, 1547–6537.
- Duarte C.G., K. Gaudreau, R.B. Gibson, and T.F. Malheiros (2013). Sustainability assessment of sugarcane-ethanol production in Brazil: A case study of a sugarcane mill in São Paulo state. *Ecological Indicators* **30**, 119–129. doi: 10.1016/j.ecolind.2013.02.011, ISSN: 1470-160X.
- Dumortier J., D.J. Hayes, M. Carriquiry, F. Dong, X. Du, A. Eloheid, J.F. Fabiosa, and S. Tokgoz (2011). Sensitivity of Carbon Emission Estimates from Indirect Land-Use Change. *Applied Economic Perspectives and Policy* **33**, 428–448. doi: 10.1093/aep/prr015, ISSN: 2040-5790, 2040–5804.
- Dunlap R.E., and W.R. Catton, Jr. (2002). Which Function(s) of the Environment Do We Study? A Comparison of Environmental and Natural Resource Sociology. *Society & Natural Resources* **15**, 239–249. doi: 10.1080/089419202753445070, ISSN: 0894-1920, 1521–0723.
- Durand L., and E. Lazos (2008). The Local Perception of Tropical Deforestation and its Relation to Conservation Policies in Los Tuxtlas Biosphere Reserve, Mexico. *Human Ecology* **36**, 383–394. doi: 10.1007/s10745-008-9172-7, ISSN: 0300-7839.
- Durand L., and L.B. Vázquez (2011). Biodiversity conservation discourses. A case study on scientists and government authorities in Sierra de Huautla Biosphere Reserve, Mexico. *Land Use Policy* **28**, 76–82. doi: 10.1016/j.landusepol.2010.04.009, ISSN: 0264-8377.
- Durst P.B., P.J. McKenzie, C.L. Brown, and S. Appanah (2006). Challenges facing certification and eco-labelling of forest products in developing countries. *International Forestry Review* **8**, 193–200. Available at: <http://www.bioone.org/doi/abs/10.1505/ifer.8.2.193>.
- Duvenage I., C. Langston, L.C. Stringer, and K. Dunstan (2013). Grappling with biofuels in Zimbabwe: depriving or sustaining societal and environmental integrity? *Journal of Cleaner Production* **42**, 132–140. doi: 10.1016/j.jclepro.2012.11.011, ISSN: 0959-6526.
- Eagle A.J., and L.P. Olander (2012). Chapter 3: Greenhouse Gas Mitigation with Agricultural Land Management Activities in the United States—A Side-by-Side Comparison of Biophysical Potential. In: *Advances in Agronomy*. Academic Press, New York, NY, pp. 79–179. ISBN: 0065-2113.
- Eckard R.J., C. Grainger, and C.A.M. de Klein (2010). Options for the abatement of methane and nitrous oxide from ruminant production: A review. *Livestock Science* **130**, 47–56. doi: 10.1016/j.livsci.2010.02.010, ISSN: 1871-1413.
- Eckert C., A. Dubini, J. Yu, P. King, M. Ghirardi, M. Seibert, and P.-C. Maness (2012). Hydrogenase Genes and Enzymes Involved in Solar Hydrogen Production. In: *State of the Art and Progress in Production of Biohydrogen*. Biosciences Center, National Renewable Energy Laboratory, Golden, CO, USA, pp. 8–24 (17). ISBN: 978-1-60805-411-4.
- Edser C. (2012). Inaugural report shows impressive uptake of RSPO-certified sustainable palm oil in 2011. *Focus on Surfactants* **2012**, 1–2. doi: 10.1016/S1351-4210(12)70119-1, ISSN: 1351-4210.
- Egeskog A., G. Berndes, F. Freitas, S. Gustafsson, and G. Sparovek (2011). Integrating bioenergy and food production—A case study of combined ethanol and dairy production in Pontal, Brazil. *Energy for Sustainable Development* **15**, 8–16. doi: 10.1016/j.esd.2011.01.005, ISSN: 0973-0826.
- Van Eijck J., H. Romijn, E. Smeets, R. Bailis, M. Rooijackers, N. Hooijkaas, P. Verweij, and A. Faaij (2014). Comparative analysis of key socio-economic and environmental impacts of smallholder and plantation based jatropha biofuel production systems in Tanzania. *Biomass and Bioenergy* **61**, 25–45. doi: 10.1016/j.biombioe.2013.10.005, ISSN: 0961-9534.

- Van Eijck J., E. Smeets, and A. Faaij (2012). The economic performance of jatropha, cassava and Eucalyptus production systems for energy in an East African smallholder setting. *GCB Bioenergy* 4, 828–845. doi: 10.1111/j.1757-1707.2012.01179.x, ISSN: 1757-1707.
- Eisenbies M.H., E.D. Vance, W.M. Aust, and J.R. Seiler (2009). Intensive utilization of harvest residues in southern pine plantations: quantities available and implications for nutrient budgets and sustainable site productivity. *BioEnergy Research* 2, 90–98.
- Elliott D.C. (2013). Transportation fuels from biomass via fast pyrolysis and hydroprocessing. *Wiley Interdisciplinary Reviews: Energy and Environment* 2, 525–533. doi: 10.1002/wene.74, ISSN: 2041-840X.
- Ellis E.C., K. Klein Goldewijk, S. Siebert, D. Lightman, and N. Ramankutty (2010). Anthropogenic transformation of the biomes, 1700 to 2000. *Global Ecology and Biogeography* 19, 589–606. doi: 10.1111/j.1466-8238.2010.00540.x, ISSN: 1466-8238.
- EMPA (2012). *Harmonisation and Extension of the Bioenergy Inventories and Assessment: End Report*. EMPA, Bern, Switzerland, 97 pp. Available at: http://www.empa.ch/plugin/template/empal*/125527/---/l=1.
- Engel S., S. Pagiola, and S. Wunder (2008). Designing payments for environmental services in theory and practice: An overview of the issues. *Ecological Economics* 65, 663–674. doi: 10.1016/j.ecolecon.2008.03.011, ISSN: 0921-8009.
- Eranki P.L., and B.E. Dale (2011). Comparative life cycle assessment of centralized and distributed biomass processing systems combined with mixed feedstock landscapes. *GCB Bioenergy* 3, 427–438. doi: 10.1111/j.1757-1707.2011.01096.x, ISSN: 1757-1707.
- Erb K.-H. (2012). How a socio-ecological metabolism approach can help to advance our understanding of changes in land-use intensity. *Ecological Economics* 76, 8–14. doi: 10.1016/j.ecolecon.2012.02.005, ISSN: 0921-8009.
- Erb K.-H., V. Gaube, F. Krausmann, C. Plutzer, A. Bondeau, and H. Haberl (2007). A comprehensive global 5 min resolution land-use data set for the year 2000 consistent with national census data. *Journal of Land Use Science* 2, 191–224. doi: 10.1080/17474230701622981, ISSN: 1747-423X.
- Erb K.-H., H. Haberl, and C. Plutzer (2012a). Dependency of global primary bioenergy crop potentials in 2050 on food systems, yields, biodiversity conservation and political stability. *Energy Policy* 47, 260–269. doi: 10.1016/j.enpol.2012.04.066, ISSN: 0301-4215.
- Erb K.-H., T. Kastner, S. Luysaert, R.A. Houghton, T. Kuemmerle, P. Olofsson, and H. Haberl (2013). Bias in the attribution of forest carbon sinks. *Nature Climate Change* 3, 854–856. doi: 10.1038/nclimate2004, ISSN: 1758-678X, 1758–6798.
- Erb K.-H., A. Mayer, F. Krausmann, C. Lauk, C. Plutzer, J. Steinberger, and H. Haberl (2012b). The interrelations of future global bioenergy potentials, food demand and agricultural technology. In: *Socioeconomic and Environmental Impacts of Biofuels: Evidence from Developing Nations*. A. Gasparatos, P. Stromberg, (eds.), Cambridge University Press, Cambridge, UK, pp. 27–52.
- Erickson P.J., J.S. Ingram, and D.M. Liverman (2009). Food security and global environmental change: emerging challenges. *Environmental Science & Policy* 12, 373–377. Available at: <http://www.sciencedirect.com/science/article/pii/S1462901109000653>.
- Eugene M., C. Martin, M.M. Mialon, D. Krauss, G. Renand, and M. Doreau (2011). Dietary linseed and starch supplementation decreases methane production of fattening bulls. *Animal Feed Science and Technology* 166–167, 330–337.
- Ewing M., and S. Msangi (2009). Biofuels production in developing countries: assessing tradeoffs in welfare and food security. *Environmental Science & Policy* 12, 520–528. doi: 10.1016/j.envsci.2008.10.002, ISSN: 1462-9011.
- Faaij A. (2006). Modern Biomass Conversion Technologies. *Mitigation and Adaptation Strategies for Global Change* 11, 335–367. doi: 10.1007/s11027-005-9004-7, ISSN: 1381-2386, 1573–1596.
- FAO (2006a). *World Agriculture: Towards 2030/2050—Interim Report. Prospects for Food, Nutrition, Agriculture and Major Commodity Groups*. Food and Agriculture Organization of the United Nations (FAO), Rome, 71 pp. Available at: <http://www.fao.org/es/ESD/AT2050web.pdf>.
- FAO (2006b). *Global Planted Forest Thematic Study, Results and Analysis. Planted Forests and Trees Working Paper FP38E*. Rome, Italy, Food and Agriculture Organization of the United Nations, 168 pp. Available at: <http://www.fao.org/forestry/12139-03441d093f070ea7d7c4e3ec3f306507.pdf>.
- FAO (2007). *The World's Mangroves: 1980–2005*. Food and Agricultural Organization of the United Nations, Rome, 153 pp. Available at: <http://www.fao.org/docrep/010/a1427e/a1427e00.htm>.
- FAO (2008). *The State of Food and Agriculture 2008—Biofuels: Prospects, Risks and Opportunities*. Food and Agricultural Organization, Rome, Italy, 128 pp. Available at: <http://www.un-energy.org/publications/98-the-state-of-food-and-agriculture-2008-biofuels-prospects-risks-and-opportunities>.
- FAO (2009). *State of the World's Forests 2009*. Food and Agriculture Organization of the United Nations, Rome, Italy, 168 pp. Available at: <http://www.fao.org/docrep/011/i0350e/i0350e00.htm>.
- FAO (2010). *Global Forest Resources Assessment 2010*. Rome, Italy, Available at: <http://www.fao.org/forestry/fra/fra2010/en/>.
- FAO (2011). *Energy-Smart Food for People and Climate*. Food and Agriculture Organization of the United Nations (FAO), Rome, Italy, 66 pp. Available at: <http://www.fao.org/docrep/014/i2454e/i2454e00.pdf>.
- FAO (2012). *The State of World Fisheries and Aquaculture*. Food and Agricultural Organization, Rome, 209 pp.
- FAO, WFP, and IFAD (2012). *The State of Food Insecurity in the World 2012. Economic Growth Is Necessary but Not Sufficient to Accelerate Reduction of Hunger and Malnutrition*. Food and Agricultural Organization of the United Nations, Rome, Italy, 61 pp. Available at: <http://www.fao.org/docrep/016/i3027e/i3027e00.htm>.
- FAOSTAT (2012). *FAO Statistics*. Food and Agriculture Organization of the United Nations, Rome, Italy, 352 pp.
- FAOSTAT (2013). FAOSTAT database. *Food and Agriculture Organization of the United Nations*. Available at: <http://faostat.fao.org/>.
- Farage, P.K., Ardo, J., Olsson, L., Rienzi, E.A., Ball, A.S., Pretty, and J.N. (2007). The potential for soil carbon sequestration in three tropical dryland farming systems of Africa and Latin America: A modelling approach. *Soil and Tillage Research* 94, 457–472.
- Fargione J., J. Hill, D. Tilman, S. Polasky, and P. Hawthorne (2008). Land Clearing and the Biofuel Carbon Debt. *Science* 319, 1235–1238. doi: 10.1126/science.1152747, ISSN: 0036-8075.
- Farley J., and R. Costanza (2010). Payments for ecosystem services: From local to global. *Ecological Economics* 69, 2060–2068.
- Favaro L., T. Jooste, M. Basaglia, S.H. Rose, M. Saayman, J.F. Görgens, S. Casella, and W.H. van Zyl (2012). Designing industrial yeasts for the consolidated bioprocessing of starchy biomass to ethanol. *Bioengineered* 4, 97–102. doi: 10.4161/bioe.22268, ISSN: 2165-5979, 2165–5987.

- Fazio S., and A. Monti (2011).** Life cycle assessment of different bioenergy production systems including perennial and annual crops. *Biomass and Bioenergy* **35**, 4868–4878. doi: 10.1016/j.biombioe.2011.10.014, ISSN: 0961-9534.
- Felder R., and R. Dones (2007).** Evaluation of ecological impacts of synthetic natural gas from wood used in current heating and car systems. *Biomass and Bioenergy* **31**, 403–415. doi: 10.1016/j.biombioe.2006.08.005, ISSN: 0961-9534.
- Feng J., C. Chen, Y. Zhang, Z. Song, A. Deng, C. Zheng, and W. Zhang (2013).** Impacts of cropping practices on yield-scaled greenhouse gas emissions from rice fields in China: A meta-analysis. *Agriculture, Ecosystems & Environment* **164**, 220–228. doi: 10.1016/j.agee.2012.10.009, ISSN: 01678809.
- Fenner A.L., S.S. Godfrey, and M.C. Bull (2011).** Using social networks to deduce whether residents or dispersers spread parasites in a lizard population: Social networks and parasite transmission. *Journal of Animal Ecology* **80**, 835–843. doi: 10.1111/j.1365-2656.2011.01825.x, ISSN: 00218790.
- Field C.B., J.E. Campbell, and D.B. Lobell (2008).** Biomass energy: the scale of the potential resource. *Trends in Ecology and Evolution* **23**, 65–72. doi: 10.1016/j.tree.2007.12.001, ISSN: 0169-5347.
- Field J.L., C.M.H. Keske, G.L. Birch, M.W. DeFoort, and M.F. Cotrufo (2013).** Distributed biochar and bioenergy coproduction: a regionally specific case study of environmental benefits and economic impacts. *GCB Bioenergy* **5**, 177–191. doi: 10.1111/gcbb.12032, ISSN: 17571693.
- Finkbeiner M. (2013).** *Indirect Land Use Change (iLUC) within Life Cycle Assessment (LCA)—Scientific Robustness and Consistency with International Standards.* Publication of the Association of the German Biofuel Industry, Berlin, Germany, 65 pp. Available at: http://www.fediol.eu/data/RZ_VDB_0030_Vorstudie_ENG_Komplett.pdf.
- Fischedick M., R. Schaeffer, A. Adedoyin, M. Akai, T. Bruckner, L. Clarke, V. Krey, I. Savolainen, S. Teske, D. Üрге-Vorsatz, and R. Wright (2011).** Mitigation Potential and Costs. In: *IPCC Special Report on Renewable Energy Sources and Climate Change Mitigation* [O. Edenhofer, R. Pichs-Madruga, Y. Sokona, K. Seyboth, P. Matschoss, S. Kadner, T. Zwickel, P. Eickemeier, G. Hansen, S. Schlömer, C. von Stechow (eds)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 791–864.
- Fischer J., B. Brosi, G.C. Daily, P.R. Ehrlich, R. Goldman, J. Goldstein, D.B. Lindenmayer, A.D. Manning, H.A. Mooney, L. Pejchar, J. Ranganathan, and H. Tallis (2008).** Should agricultural policies encourage land sparing or wildlife-friendly farming? *Frontiers in Ecology and the Environment* **6**, 380–385. doi: 10.1890/070019, ISSN: 1540-9295.
- Fischer G., S. Prieler, H. Van Velthuisen, G. Berndes, A. Faaij, M. Londo, and M. De Wit (2010).** Biofuel production potentials in Europe: Sustainable use of cultivated land and pastures, Part II: Land use scenarios. *Biomass and Bioenergy* **34**, 173–187.
- Fischer-Kowalski M. (2011).** Analyzing sustainability transitions as a shift between socio-metabolic regimes. *Environmental Innovation and Societal Transitions* **1**, 152–159. doi: 16/j.eist.2011.04.004, ISSN: 2210-4224.
- Flannigan M., B. Stocks, M. Turetsky, and M. Wotton (2009).** Impacts of climate change on fire activity and fire management in the circumboreal forest. *Global Change Biology* **15**, 549–560. doi: 10.1111/j.1365-2486.2008.01660.x, ISSN: 1365-2486.
- FLEG Regional Forest Law Enforcement and Governance (FLEG) Initiatives.** Available at: <http://www.worldbank.org/en/topic/forests/brief/fleg-regional-forest-law-enforcement-governance>.
- Foley J.A., R. DeFries, G.P. Asner, C. Barford, G. Bonan, S.R. Carpenter, F.S. Chapin, M.T. Coe, G.C. Daily, H.K. Gibbs, J.H. Helkowski, T. Holloway, E.A. Howard, C.J. Kucharik, C. Monfreda, J.A. Patz, I.C. Prentice, N. Ramanakutty, and P.K. Snyder (2005).** Global Consequences of Land Use. *Science* **309**, 570–574. doi: 10.1126/science.1111772, ISSN: 0036-8075, 1095–9203.
- Foley P.A., D.A. Kenny, J.J. Callan, T.M. Boland, and F.P. O'Mara (2009).** Effect of DL-malic acid supplementation on feed intake, methane emission, and rumen fermentation in beef cattle. *Journal of Animal Science* **87**, 1048–1057. doi: 10.2527/jas.2008-1026, ISSN: 0021-8812, 1525–3163.
- Foley J.A., N. Ramanakutty, K.A. Brauman, E.S. Cassidy, J.S. Gerber, M. Johnston, N.D. Mueller, C. O'Connell, D.K. Ray, P.C. West, C. Balzer, E.M. Bennett, S.R. Carpenter, J. Hill, C. Monfreda, S. Polasky, J. Rockstrom, J. Sheehan, S. Siebert, D. Tilman, and D.P.M. Zaks (2011).** Solutions for a cultivated planet. *Nature* **478**, 337–342. doi: 10.1038/nature10452, ISSN: 0028-0836.
- Follett R.F., and D.A. Reed (2010).** Soil Carbon Sequestration in Grazing Lands: Societal Benefits and Policy Implications. *Rangeland Ecology & Management* **63**, 4–15. doi: 10.2111/08-225.1, ISSN: 1550-7424, 1551–5028.
- Forner C., J. Blaser, F. Jotzo, and C. Robledo (2006).** Keeping the forest for the climate's sake: avoiding deforestation in developing countries under the UNFCCC. *Climate Policy* **6**, 275–294. doi: 10.1080/14693062.2006.9685602, ISSN: 1469-3062.
- Frank S., H. Böttcher, P. Havlik, H. Valin, A. Mosnier, M. Obersteiner, E. Schmid, and B. Elbersen (2013).** How effective are the sustainability criteria accompanying the European Union 2020 biofuel targets? *GCB Bioenergy*, n/a–n/a. doi: 10.1111/j.1757-1707.2012.01188.x, ISSN: 1757-1707.
- Franzluubbers A.J., and J.A. Stuedemann (2009).** Soil-profile organic carbon and total nitrogen during 12 years of pasture management in the Southern Piedmont USA. *Agriculture, Ecosystems & Environment* **129**, 28–36. doi: 10.1016/j.agee.2008.06.013, ISSN: 01678809.
- Frei M., M.A. Razzak, M.M. Hossain, M. Oehme, S. Dewan, and K. Becker (2007).** Methane emissions and related physicochemical soil and water parameters in rice–fish systems in Bangladesh. *Agriculture, Ecosystems & Environment* **120**, 391–398. doi: 10.1016/j.agee.2006.10.013, ISSN: 01678809.
- Fuglestvedt J.S., K.P. Shine, T. Berntsen, J. Cook, D.S. Lee, A. Stenke, R.B. Skeie, G.J.M. Velders, and I.A. Waitz (2010).** Transport impacts on atmosphere and climate: Metrics. *Atmospheric Environment* **44**, 4648–4677.
- Gabrielle B., and N. Gagnaire (2008).** Life-cycle assessment of straw use in bioethanol production: A case study based on biophysical modelling. *Biomass and Bioenergy* **32**, 431–441. doi: 10.1016/j.biombioe.2007.10.017, ISSN: 0961-9534.
- Gabrielle B., P. Maupu, and E. Vial (2013).** Life cycle assessment of eucalyptus short rotation coppices for bioenergy production in southern France. *GCB Bioenergy* **5**, 30–42. doi: 10.1111/gcbb.12008, ISSN: 1757-1707.
- Galinato S.P., J.K. Yoder, and D. Granatstein (2011).** The economic value of biochar in crop production and carbon sequestration. *Energy Policy* **39**, 6344–6350. doi: 10.1016/j.enpol.2011.07.035, ISSN: 0301-4215.
- García-Frapolli E., A. Schilman, V. Berrueta, H. Riojas-Rodríguez, R. Edwards, M. Johnson, A. Guevara-Sangines, C. Armendariz, and O.R. Masera (2010).** Beyond Fuelwood Savings: Valuing the economic benefits of introducing improved biomass cookstoves in the Purépecha region of Mexico. *Ecological Economics* **69**, 2598–2605.

- Garcia-Ulloa J., S. Sloan, P. Pacheco, J. Ghazoul, and L.P. Koh (2012). Lowering environmental costs of oil-palm expansion in Colombia. *Conservation Letters* 5, 366–375. doi: 10.1111/j.1755-263X.2012.00254.x, ISSN: 1755263X.
- Gardner T.A., N.D. Burgess, N. Aguilar-Amuchastegui, J. Barlow, E. Berenguer, T. Clements, F. Danielsen, J. Ferreira, W. Foden, V. Kapos, S.M. Khan, A.C. Lees, L. Parry, R.M. Roman-Cuesta, C.B. Schmitt, N. Strange, I. Theilade, and I.C.G. Vieira (2012). A framework for integrating biodiversity concerns into national REDD+ programmes. *Biological Conservation* 154, 61–71. doi: 10.1016/j.biocon.2011.11.018, ISSN: 0006-3207.
- Garg K.K., L. Karlberg, S.P. Wani, and G. Berndes (2011). Jatropha production on wastelands in India: opportunities and trade-offs for soil and water management at the watershed scale. *Biofuels, Bioproducts and Biorefining* 5, 410–430. doi: 10.1002/bbb.312, ISSN: 1932-1031.
- Garnett T. (2011). Where are the best opportunities for reducing greenhouse gas emissions in the food system (including the food chain)? *Food Policy* 36, Supplement 1, 23–32. doi: 10.1016/j.foodpol.2010.10.010, ISSN: 0306-9192.
- Garnett T., M.C. Appleby, A. Balmford, I.J. Bateman, T.G. Benton, P. Bloomer, B. Burlingame, M. Dawkins, L. Dolan, D. Fraser, M. Herrero, I. Hoffmann, P. Smith, P.K. Thornton, C. Toulmin, S.J. Vermeulen, and H.C.J. Godfray (2013). Sustainable Intensification in Agriculture: Premises and Policies. *Science* 341, 33–34. doi: 10.1126/science.1234485, ISSN: 0036-8075, 1095–9203.
- Gasparatos A., P. Stromberg, and K. Takeuchi (2011). Biofuels, ecosystem services and human wellbeing: Putting biofuels in the ecosystem services narrative. *Agriculture, Ecosystems & Environment* 142, 111–128. doi: 16/j.agee.2011.04.020, ISSN: 0167-8809.
- Gettinger A., A. Muller, M. Haeni, C. Skinner, A. Fliessbach, N. Buchmann, P. Mader, M. Stolze, P. Smith, N.E.-H. Scialabba, and U. Niggli (2012). Enhanced top soil carbon stocks under organic farming. *Proceedings of the National Academy of Sciences* 109, 18226–18231. doi: 10.1073/pnas.1209429109, ISSN: 0027-8424, 1091–6490.
- Gawel E., and G. Ludwig (2011). The iLUC dilemma: How to deal with indirect land use changes when governing energy crops? *Land Use Policy* 28, 846–856. doi: 10.1016/j.landusepol.2011.03.003, ISSN: 0264-8377.
- GEA (2012). *Global Energy Assessment—Toward a Sustainable Future*. Cambridge University Press, Cambridge, UK and New York, NY, USA and the International Institute for Applied Systems Analysis, Laxenburg, Austria, 1802 pp.
- Von Geibler J. (2013). Market-based governance for sustainability in value chains: conditions for successful standard setting in the palm oil sector. *Journal of Cleaner Production* 56, 39–53. doi: 10.1016/j.jclepro.2012.08.027, ISSN: 0959-6526.
- Gelfand I., R. Sahajpal, X. Zhang, R.C. Izaurralde, K.L. Gross, and G.P. Robertson (2013). Sustainable bioenergy production from marginal lands in the US Midwest. *Nature* 493, 514–517. doi: 10.1038/nature11811, ISSN: 0028-0836, 1476–4687.
- Gelfand I., T. Zenone, P. Jasrotia, J. Chen, S.K. Hamilton, and G.P. Robertson (2011). Carbon debt of Conservation Reserve Program (CRP) grasslands converted to bioenergy production. *Proceedings of the National Academy of Sciences of the United States* 108, 13864–13869.
- Georgescu M., D.B. Lobell, and C.B. Field (2011). Direct climate effects of perennial bioenergy crops in the United States. *Proceedings of the National Academy of Sciences* 108, 4307–4312. doi: 10.1073/pnas.1008779108, ISSN: 0027-8424, 1091–6490.
- Gerbens-Leenes W., A.Y. Hoekstra, and T.H. Van der Meer (2009). The water footprint of bioenergy. *Proceedings of the National Academy of Sciences* 106, 10219–10223. Available at: <http://www.pnas.org/content/106/25/10219>. short.
- Gerber P., H.A. Mooney, J. Dijkman, S. Tarawali, and C. de Haan (Eds.) (2010). *Livestock in a Changing Landscape. Experiences and Regional Perspectives*. Island Press, Washington, D.C., 189 pp. ISBN: 9781597266734.
- German L., and G. Schoneveld (2012). A review of social sustainability considerations among EU-approved voluntary schemes for biofuels, with implications for rural livelihoods. *Energy Policy* 51, 765–778. doi: 10.1016/j.enpol.2012.09.022, ISSN: 0301-4215.
- German L., G.C. Schoneveld, and D. Gumbo (2011). The Local Social and Environmental Impacts of Smallholder-Based Biofuel Investments in Zambia. *Ecology and Society* 16. doi: 10.5751/es-04280-160412, ISSN: 1708-3087.
- German L., G. Schoneveld, and E. Mwangi (2013). Contemporary Processes of Large-Scale Land Acquisition in Sub-Saharan Africa: Legal Deficiency or Elite Capture of the Rule of Law? *World Development* 48, 1–18. doi: 10.1016/j.worlddev.2013.03.006, ISSN: 0305-750X.
- Gibbs H.K., M. Johnston, J.A. Foley, T. Holloway, C. Monfreda, N. Ramankutty, and D. Zaks (2008). Carbon payback times for crop-based biofuel expansion in the tropics: the effects of changing yield and technology. *Environmental Research Letters* 3, 034001 (10pp). doi: 10.1088/1748-9326/3/3/034001, ISSN: 1748-9326.
- Gilg A. (2009). Perceptions about land use. *Land Use Policy* 26, Supplement 1, 76–82. doi: 10.1016/j.landusepol.2009.08.018, ISSN: 0264-8377.
- Gill M., P. Smith, and J.M. Wilkinson (2010). Mitigating climate change: the role of domestic livestock. *Animal* 4, 323–333. doi: 10.1017/S1751731109004662.
- Giuntoli J., A.K. Boulamanti, S. Corrado, M. Motegh, A. Agostini, and D. Baxter (2013). Environmental impacts of future bioenergy pathways: the case of electricity from wheat straw bales and pellets. *GCB Bioenergy* 5, 497–512. doi: 10.1111/gcbb.12012, ISSN: 1757-1707.
- Gnansounou E. (2011). Assessing the sustainability of biofuels: A logic-based model. *Energy* 36, 2089–2096. doi: 10.1016/j.energy.2010.04.027, ISSN: 03605442.
- Godfray H.C.J., J.R. Beddington, I.R. Crute, L. Haddad, D. Lawrence, J.F. Muir, J. Pretty, S. Robinson, S.M. Thomas, and C. Toulmin (2010). Food Security: The Challenge of Feeding 9 Billion People. *Science* 327, 812–818. doi: 10.1126/science.1185383, ISSN: 0036-8075, 1095–9203.
- Goglio P., E. Bonari, and M. Mazzoncini (2012). LCA of cropping systems with different external input levels for energetic purposes. *Biomass and Bioenergy* 42, 33–42. doi: 10.1016/j.biombioe.2012.03.021, ISSN: 0961-9534.
- Gohin A. (2008). Impacts of the European Biofuel Policy on the Farm Sector: A General Equilibrium Assessment. *Applied Economic Perspectives and Policy* 30, 623–641. doi: 10.1111/j.1467-9353.2008.00437.x, ISSN: 2040-5790, 2040–5804.
- Goldemberg J. (2007). Ethanol for a sustainable energy future. *Science* 315, 808–810.
- Goldemberg J., S.T. Coelho, and P. Guardabassi (2008). The sustainability of ethanol production from sugarcane. *Energy Policy* 36, 2086–2097.
- Goldewijk K.K., A. Beusen, G. van Drecht, and M. de Vos (2011). The HYDE 3.1 spatially explicit database of human-induced global land-use change over the past 12,000 years. *Global Ecology and Biogeography* 20, 73–86. doi: 10.1111/j.1466-8238.2010.00587.x, ISSN: 1466-822X.

- Gollakota S., and S. McDonald (2012).** CO₂ capture from ethanol production and storage into the Mt Simon Sandstone. *Greenhouse Gases: Science and Technology* 2, 346–351.
- Golub A., T. Hertel, H.L. Lee, S. Rose, and B. Sohngen (2009).** The opportunity cost of land use and the global potential for greenhouse gas mitigation in agriculture and forestry. *Resource and Energy Economics* 31, 299–319.
- González A.D., B. Frostell, and A. Carlsson-Kanyama (2011).** Protein efficiency per unit energy and per unit greenhouse gas emissions: Potential contribution of diet choices to climate change mitigation. *Food Policy* 36, 562–570. doi: 10.1016/j.foodpol.2011.07.003, ISSN: 0306-9192.
- Gonzalez P., R.P. Neilson, J.M. Lenihan, and R.J. Drapek (2010).** Global patterns in the vulnerability of ecosystems to vegetation shifts due to climate change. *Global Ecology and Biogeography* 19, 755–768. doi: 10.1111/j.1466-8238.2010.00558.x, ISSN: 1466-8238.
- González-Estrada E., L.C. Rodriguez, V.K. Walen, J.B. Naab, J. Koo, J.W. Jones, M. Herrero, and P.K. Thornton (2008).** Carbon sequestration and farm income in West Africa: Identifying best management practices for smallholder agricultural systems in northern Ghana. *Ecological Economics* 67, 492–502.
- González-García S., J. Bacenetti, R.J. Murphy, and M. Fiala (2012a).** Present and future environmental impact of poplar cultivation in the Po Valley (Italy) under different crop management systems. *Journal of Cleaner Production* 26, 56–66. doi: 10.1016/j.jclepro.2011.12.020, ISSN: 0959-6526.
- González-García S., M.T. Moreira, G. Feijoo, and R.J. Murphy (2012b).** Comparative life cycle assessment of ethanol production from fast-growing wood crops (black locust, eucalyptus and poplar). *Biomass and Bioenergy* 39, 378–388. doi: 10.1016/j.biombioe.2012.01.028, ISSN: 0961-9534.
- Gopalakrishnan G., M.C. Negri, and W. Salas (2012).** Modeling biogeochemical impacts of bioenergy buffers with perennial grasses for a row-crop field in Illinois. *Global Change Biology Bioenergy* 4, 739–750. doi: 10.1111/j.1757-1707.2011.01145.x.
- Gopalakrishnan G., M.C. Negri, and S. Snyder (2011a).** A novel framework to classify marginal land for sustainable biomass feedstock production. *Journal of Environmental Quality* 40, 1593–1600. Available at: <http://www.sciencedirect.com/science/article/pii/S0306261911002406>.
- Gopalakrishnan G., M.C. Negri, and S. Snyder (2011b).** Redesigning agricultural landscapes for sustainability using bioenergy crops: quantifying the tradeoffs between agriculture, energy and the environment. *Aspects of Applied Biology* 112, 139–146.
- Gopalakrishnan G., M.C. Negri, M. Wang, M. Wu, S.W. Snyder, and L. Lafreniere (2009).** Biofuels, land and water: a systems approach to sustainability. *Environmental Science & Technology* 43, 6094–6100.
- Görg C. (2007).** Landscape governance: The “politics of scale” and the “natural” conditions of places. *Pro-Poor Water? The Privatisation and Global Poverty Debate* 38, 954–966. doi: 10.1016/j.geoforum.2007.01.004, ISSN: 0016-7185.
- Gorham E. (1991).** Northern peatlands: role in the carbon cycle and probable responses to climatic warming. *Ecological Applications* 1, 182–195.
- Gottschalk P., J.U. Smith, M. Wattenbach, J. Bellarby, E. Stehfest, N. Arnell, T.J. Osborn, and P. Smith (2012).** How will organic carbon stocks in mineral soils evolve under future climate? Global projections using RothC for a range of climate change scenarios. *Biogeosciences Discussions* 9, 411–451. doi: 10.5194/bgd-9-411-2012, ISSN: 1810-6285.
- Government of New Zealand** New Zealand climate change information. Available at: <http://www.climatechange.govt.nz/>.
- Graham-Rowe D. (2011).** Agriculture: Beyond food versus fuel. *Nature* 474, 6–8. doi: 10.1038/474506a, ISSN: 0028-0836.
- Grainger C., and K.A. Beauchemin (2011).** Can enteric methane emissions from ruminants be lowered without lowering their production? *Animal Feed Science and Technology* 166–167, 308–320. doi: 10.1016/j.anifeedsci.2011.04.021, ISSN: 0377-8401.
- Grainger C., T. Clarke, K.A. Beauchemin, S.M. McGinn, and R.J. Eckard (2008).** Supplementation with whole cottonseed reduces methane emissions and can profitably increase milk production of dairy cows offered a forage and cereal grain diet. *Australian Journal of Experimental Agriculture* 48, 73–76.
- Grainger C., R. Williams, T. Clarke, A.G. Wright, and R.J. Eckard (2010).** Supplementation with whole cottonseed causes long-term reduction of methane emissions from lactating dairy cows offered a forage and cereal grain diet. *Journal of Dairy Science* 93, 2612–2619.
- Gregg J.S., and S.J. Smith (2010).** Global and regional potential for bioenergy from agricultural and forestry residue biomass. *Mitigation and Adaptation Strategies for Global Change* 15, 241–262. doi: 10.1007/s11027-010-9215-4, ISSN: 1381-2386, 1573–1596.
- Di Gregorio M., M. Brockhaus, T. Cronin, E. Muharrom, L. Santoso, S. Mardiah, and M. Büdenbender (2013).** Equity and REDD+ in the Media: a Comparative Analysis of Policy Discourses. *Ecology and Society* 18, 39. doi: 10.5751/ES-05694-180239.
- Groom M., E. Gray, and P. Townsend (2008).** Biofuels and Biodiversity: Principles for Creating Better Policies for Biofuel Production. *Conservation Biology* 22, 602–609. doi: 10.1111/j.1523-1739.2007.00879.x, ISSN: 1523-1739.
- De Groot R. (2006).** Function-analysis and valuation as a tool to assess land use conflicts in planning for sustainable, multi-functional landscapes. *Landscape and Urban Planning* 75, 175–186.
- Guariguata M.R., J.P. Cornelius, B. Locatelli, C. Forner, and G.A. Sánchez-Azofeifa (2008).** Mitigation needs adaptation: Tropical forestry and climate change. *Mitigation and Adaptation Strategies for Global Change* 13, 793–808. doi: 10.1007/s11027-007-9141-2, ISSN: 1381-2386, 1573–1596.
- Guest G., R.M. Bright, F. Cherubini, O. Michelsen, and A.H. Strømman (2011).** Life Cycle Assessment of Biomass-based Combined Heat and Power Plants. *Journal of Industrial Ecology* 15, 908–921. doi: 10.1111/j.1530-9290.2011.00375.x, ISSN: 1530-9290.
- Guest G., R.M. Bright, F. Cherubini, and A.H. Strømman (2013).** Consistent quantification of climate impacts due to biogenic carbon storage across a range of bio-product systems. *Environmental Impact Assessment Review* 43, 21–30.
- Guldea S., H. Chung, W. Amelung, C. Chang, and J. Six (2008).** Soil carbon saturation controls labile and stable carbon pool dynamics. *Soil Science Society of America Journal* 72, 605–612. doi: 10.2136/sssaj2007.0251.
- Gumpenberger M., K. Vohland, U. Heyder, B. Poulter, K. Macey, A. Rammig, A. Popp, and W. Cramer (2010).** Predicting pan-tropical climate change induced forest stock gains and losses—implications for REDD. *Environmental Research Letters* 5, 014013 (15pp). doi: 10.1088/1748-9326/5/1/014013, ISSN: 1748-9326.
- Gupta J. (2012).** Global forest and REDD+ governance: win–win or lose–lose? *Current Opinion in Environmental Sustainability* 4, 620–627. doi: 10.1016/j.cosust.2012.09.014, ISSN: 1877-3435.

- Gurung A., and S.E. Oh (2013).** Conversion of traditional biomass into modern bioenergy systems: A review in context to improve the energy situation in Nepal. *Renewable Energy* **50**, 206–213. doi: 10.1016/j.renene.2012.06.021, ISSN: 0960-1481.
- Gustavsson J., C. Cederberg, U. Sonesson, R. van Otterdijk, and A. Meybeck (2011).** *Global Food Losses and Food Waste. Extent, Causes and Prevention.* Food and Agricultural Organization of the United Nations, Rome, 29 pp.
- Gustavsson L., K. Pingoud, and R. Sathre (2006).** Carbon Dioxide Balance of Wood Substitution: Comparing Concrete- and Wood-Framed Buildings. *Mitigation and Adaptation Strategies for Global Change* **11**, 667–691. doi: 10.1007/s11027-006-7207-1, ISSN: 1381-2386.
- Guthinga P.M. (2008).** Understanding local communities' perceptions of existing forest management regimes of Kenyan rainforest. *International Journal of Social Forestry (IJSF)* **1**, 145–166.
- Haberl H. (2013).** Net land-atmosphere flows of biogenic carbon related to bioenergy: towards an understanding of systemic feedbacks. *Global Change Biology Bioenergy*.
- Haberl H., T. Beringer, S.C. Bhattacharya, K.-H. Erb, and M. Hoogwijk (2010).** The global technical potential of bio-energy in 2050 considering sustainability constraints. *Current Opinion in Environmental Sustainability* **2**, 394–403. doi: 10.1016/j.cosust.2010.10.007, ISSN: 1877-3435.
- Haberl H., K.-H. Erb, F. Krausmann, A. Bondeau, C. Lauk, C. Müller, C. Plutzer, and J.K. Steinberger (2011a).** Global bioenergy potentials from agricultural land in 2050: Sensitivity to climate change, diets and yields. *Biomass and Bioenergy* **35**, 4753–4769. doi: 10.1016/j.biombioe.2011.04.035, ISSN: 0961-9534.
- Haberl H., K.H. Erb, F. Krausmann, V. Gaube, A. Bondeau, C. Plutzer, S. Gingrich, W. Lucht, and M. Fischer-Kowalski (2007).** Quantifying and mapping the human appropriation of net primary production in earth's terrestrial ecosystems. *Proceedings of the National Academy of Sciences* **104**, 12942–12947. doi: 10.1073/pnas.0704243104.
- Haberl H., M. Fischer-Kowalski, F. Krausmann, J. Martinez-Alier, and V. Winzarter (2011b).** A socio-metabolic transition towards sustainability? Challenges for another Great Transformation. *Sustainable Development* **19**, 1–14. doi: 10.1002/sd.410, ISSN: 1099-1719.
- Haberl H., C. Mbow, X. Deng, E.G. Irwin, S. Kerr, T. Kuemmerle, O. Mertz, P. Meyfroidt, and B.L. Turner II (2013a).** Finite Land Resources and Competition. Strüngmann Forum Reports. In: *Rethinking global land use in an urban era*, edited by Karen C. Seto and Anette Reenberg. MIT Press, Cambridge, MA, pp. 33–67. ISBN: 9780262026901.
- Haberl H., E.-D. Schulze, C. Körner, B.E. Law, B. Holtzmark, and S. Luysaert (2013b).** Response: complexities of sustainable forest use. *GCB Bioenergy* **5**, 1–2. doi: 10.1111/gcbb.12004, ISSN: 17571693.
- Haberl H., D. Sprinz, M. Bonazountas, P. Cocco, Y. Desaubies, M. Henze, O. Hertel, R.K. Johnson, U. Kastrop, P. Laconte, E. Lange, P. Novak, J. Paavola, A. Reenberg, S. van den Hove, T. Vermeire, P. Wadhams, and T. Searchinger (2012).** Correcting a fundamental error in greenhouse gas accounting related to bioenergy. *Energy Policy* **45**, 18–23. doi: 10.1016/j.enpol.2012.02.051, ISSN: 0301-4215.
- Hakala K., M. Kontturi, and K. Pahkala (2009).** Field biomass as global energy source. *Agricultural and Food Science* **18**, 347–365. ISSN: 1459-6067.
- Hall J., S. Matos, L. Severino, and N. Beltrão (2009).** Brazilian biofuels and social exclusion: established and concentrated ethanol versus emerging and dispersed biodiesel. *Journal of Cleaner Production* **17**, Supplement 1, S77–S85. doi: 10.1016/j.jclepro.2009.01.003, ISSN: 0959-6526.
- Hallgren W., C.A. Schlosser, E. Monier, D. Kicklighter, A. Sokolov, and J. Melillo (2013).** Climate impacts of a large-scale biofuels expansion. *Geophysical Research Letters* **40**, 1624–1630. doi: 10.1002/grl.50352, ISSN: 1944-8007.
- Halsnæs K., and J. Verhagen (2007).** Development based climate change adaptation and mitigation—conceptual issues and lessons learned in studies in developing countries. *Mitigation and Adaptation Strategies for Global Change Volume 12*, 665–684.
- Halvorson J.J., J.M. Gonzalez, and A.E. Hagerman (2011).** Repeated applications of tannins and related phenolic compounds are retained by soil and affect cation exchange capacity. *Soil Biology and Biochemistry* **43**, 1139–1147. doi: 10.1016/j.soilbio.2011.01.023, ISSN: 0038-0717.
- Hamelinck C.N., and A.P. Faaij (2006).** Outlook for advanced biofuels. *Energy Policy* **34**, 3268–3283.
- Hanff E., M.-H. Dabat, and J. Blin (2011).** Are biofuels an efficient technology for generating sustainable development in oil-dependent African nations? A macroeconomic assessment of the opportunities and impacts in Burkina Faso. *Renewable and Sustainable Energy Reviews* **15**, 2199–2209. doi: 10.1016/j.rser.2011.01.014, ISSN: 1364-0321.
- Hansen M.C., S.V. Stehman, and P.V. Potapov (2010).** Quantification of global gross forest cover loss. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 8650–8655. doi: 10.1073/pnas.0912668107, ISSN: 0027-8424.
- Hardner J.J., P.C. Frumhoff, and D.C. Goetze (2000).** Prospects for mitigating carbon, conserving biodiversity, and promoting socioeconomic development objectives through the clean development mechanism. *Mitigation and Adaptation Strategies for Global Change* **5**, 61–80. doi: 10.1023/A:1009685323704, ISSN: 1381-2386, 1573–1596.
- Harper R.J., A.C. Beck, P. Ritson, M.J. Hill, C.D. Mitchell, D.J. Barrett, K.R.J. Smettem, and S.S. Mann (2007).** The potential of greenhouse sinks to underwrite improved land management. *Ecological Engineering* **29**, 329–341.
- Harper R.J., S.J. Sochacki, K.R.J. Smettem, and N. Robinson (2010).** Bioenergy Feedstock Potential from Short-Rotation Woody Crops in a Dryland Environment†. *Energy & Fuels* **24**, 225–231. doi: 10.1021/ef9005687, ISSN: 0887-0624.
- Harris N.L., S. Brown, S.C. Hagen, S.S. Saatchi, S. Petrova, W. Salas, M.C. Hansen, P.V. Potapov, and A. Lotsch (2012).** Baseline Map of Carbon Emissions from Deforestation in Tropical Regions. *Science* **336**, 1573–1576. doi: 10.1126/science.1217962, ISSN: 0036-8075, 1095–9203.
- Harsono S.S., A. Prochnow, P. Grundmann, A. Hansen, and C. Hallmann (2012).** Energy balances and greenhouse gas emissions of palm oil biodiesel in Indonesia. *GCB Bioenergy* **4**, 213–228. doi: 10.1111/j.1757-1707.2011.01118.x, ISSN: 1757-1707.
- Harvey C.A., B. Dickson, and C. Kormos (2010).** Opportunities for achieving biodiversity conservation through REDD. *Conservation Letters* **3**, 53–61. doi: 10.1111/j.1755-263X.2009.00086.x.
- Harvey C.A., O. Komar, R. Chazdon, B.G. Ferguson, B. Finegan, D.M. Griffith, M. Martinez-Ramos, H. Morales, R. Nigh, L. Soto-Pinto, M. van Breugel, and M. Wishnie (2008).** Integrating agricultural landscapes with biodiversity conservation in the Mesoamerican hotspot. *Conservation Biology* **22**, 8–15. Available at: <http://onlinelibrary.wiley.com/doi/10.1111/j.1523-1739.2007.00863.x/full>.

- Harvey M., and S. Pilgrim (2011). The new competition for land: Food, energy, and climate change. *Food Policy* 36, 40–51.
- Havemann T. (2011). *Financing Mitigation in Smallholder Agricultural Systems: Issues and Opportunities*. CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS), Copenhagen, Denmark, 27 pp. Available at: <http://cgspace.cgiar.org/handle/10568/6576>.
- Havlik P., U.A. Schneider, E. Schmid, H. Böttcher, S. Fritz, R. Skalský, K. Aoki, S.D. Cara, G. Kindermann, F. Kraxner, S. Leduc, I. McCallum, A. Mosnier, T. Sauer, and M. Obersteiner (2011). Global land-use implications of first and second generation biofuel targets. *Energy Policy* 39, 5690–5702. doi: 10.1016/j.enpol.2010.03.030, ISSN: 0301-4215.
- Hegarty R.S., J.P. Goopy, R.M. Herd, and B. McCorkell (2007). Cattle selected for lower residual feed intake have reduced daily methane production. *Journal of Animal Science* 85, 1479–1486. doi: 10.2527/jas.2006-236, ISSN: 0021-8812, 1525–3163.
- Heggenstaller A.H., R.P. Anex, M. Liebman, D.N. Sundberg, and L.R. Gibson (2008). Productivity and Nutrient Dynamics in Bioenergy Double-Cropping Systems. *Agronomy Journal* 100, 1740. doi: 10.2134/agronj2008.0087, ISSN: 1435-0645.
- Hein L., and P.J. van der Meer (2012). REDD+ in the context of ecosystem management. *Current Opinion in Environmental Sustainability* 4, 604–611. doi: 10.1016/j.cosust.2012.09.016, ISSN: 1877-3435.
- Helin T., L. Sokka, S. Soimakallio, K. Pingoud, and T. Pajula (2013). Approaches for inclusion of forest carbon cycle in life cycle assessment—a review. *GCB Bioenergy* 5, 475–486. doi: 10.1111/gcbb.12016, ISSN: 1757-1707.
- Heller N.E., and E.S. Zavaleta (2009). Biodiversity management in the face of climate change: A review of 22 years of recommendations. *Biological Conservation* 142, 14–32. doi: 10.1016/j.biocon.2008.10.006, ISSN: 00063207.
- Hennig C., and M. Gawor (2012). Bioenergy production and use: Comparative analysis of the economic and environmental effects. *Energy Conversion and Management* 63, 130–137. doi: 10.1016/j.enconman.2012.03.031, ISSN: 0196-8904.
- Herawati H., and H. Santoso (2011). Tropical forest susceptibility to and risk of fire under changing climate: A review of fire nature, policy and institutions in Indonesia. *Forest Policy and Economics* 13, 227–233.
- Herold M., and T. Johns (2007). Linking requirements with capabilities for deforestation monitoring in the context of the UNFCCC-REDD process. *Environmental Research Letters* 2, 045025. doi: 10.1088/1748-9326/2/4/045025, ISSN: 1748-9326.
- Herold M., and M. Skutsch (2011). Monitoring, reporting and verification for national REDD+ programmes: two proposals. *Environmental Research Letters* 6, doi:10.1088/1748-9326/6/1/014002.
- Herrero M., P. Havlik, H. Valin, A. Notenbaert, M.C. Rufino, P.K. Thornton, M. Blümmel, F. Weiss, D. Grace, and M. Obersteiner (2013). Biomass use, production, feed efficiencies, and greenhouse gas emissions from global livestock systems. *Proceedings of the National Academy of Sciences* 110, 20888–20893. doi: 10.1073/pnas.1308149110, ISSN: 0027-8424, 1091–6490.
- Herrero M., P.K. Thornton, A.M. Notenbaert, S. Wood, S. Msangi, H.A. Freeman, D. Bossio, J. Dixon, M. Peters, J. van de Steeg, J. Lynam, P.P. Rao, S. Macmillan, B. Gerard, J. McDermott, C. Sere, and M. Rosegrant (2010). Smart Investments in Sustainable Food Production: Revisiting Mixed Crop-Livestock Systems. *Science* 327, 822–825. doi: 10.1126/science.1183725, ISSN: 0036-8075, 1095–9203.
- Hertel T.W., A.A. Golub, A.D. Jones, M. O’Hare, R.J. Plevin, and D.M. Kammen (2010). Effects of US Maize Ethanol on Global Land Use and Greenhouse Gas Emissions: Estimating Market-mediated Responses. *BioScience* 60, 223–231. doi: 10.1525/bio.2010.60.3.8, ISSN: 0006-3568, 1525–3244.
- Hertwich E. (2012). Biodiversity: Remote responsibility. *Nature* 486, 36–37. doi: 10.1038/486036a, ISSN: 0028-0836.
- Hill J., E. Nelson, D. Tilman, S. Polasky, and D. Tiffany (2006). Environmental, economic, and energetic costs and benefits of biodiesel and ethanol biofuels. *Proceedings of the National Academy of Sciences* 103, 11206–11210. doi: 10.1073/pnas.0604600103, ISSN: 0027-8424, 1091–6490.
- Van der Hilst F., V. Dornburg, J.P.M. Sanders, B. Elbersen, A. Graves, W.C. Turkenburg, H.W. Elbersen, J.M.C. van Dam, and A.P.C. Faaij (2010). Potential, spatial distribution and economic performance of regional biomass chains: The North of the Netherlands as example. *Agricultural Systems* 103, 403–417.
- Van der Hilst F., and A.P. Faaij (2012). Spatiotemporal cost-supply curves for bioenergy production in Mozambique. *Biofuels, Bioproducts and Biorefining* 6, 405–430.
- Van der Hilst F., J.P. Lesschen, J.M.C. van Dam, M. Riksen, P.A. Verweij, J.P.M. Sanders, and A.P.C. Faaij (2012a). Spatial variation of environmental impacts of regional biomass chains. *Renewable and Sustainable Energy Reviews* 16, 2053–2069. doi: 10.1016/j.rser.2012.01.027, ISSN: 1364-0321.
- Van der Hilst F., J.A. Versteegen, D. Karssen, and A.P.C. Faaij (2012b). Spatiotemporal land use modelling to assess land availability for energy crops—illustrated for Mozambique. *GCB Bioenergy* 4, 859–874. doi: 10.1111/j.1757-1707.2011.01147.x, ISSN: 1757-1707.
- Hodges R.J., J.C. Buzby, and B. Bennett (2011). Postharvest Losses and Waste in Developed and Less Developed Countries: Opportunities to Improve Resource Use. *The Journal of Agricultural Science* 149, 37–45. doi: 10.1017/S0021859610000936.
- Hoefnagels R., M. Banse, V. Dornburg, and A. Faaij (2013). Macro-economic impact of large-scale deployment of biomass resources for energy and materials on a national level—A combined approach for the Netherlands. *Energy Policy* 59, 727–744. doi: 10.1016/j.enpol.2013.04.026, ISSN: 0301-4215.
- Hoefnagels R., E. Smeets, and A. Faaij (2010). Greenhouse gas footprints of different biofuel production systems. *Renewable and Sustainable Energy Reviews* 14, 1661–1694. doi: 10.1016/j.rser.2010.02.014, ISSN: 1364-0321.
- Holtmark B. (2012). Harvesting in boreal forests and the biofuel carbon debt. *Climatic Change* 112, 415–428. doi: 10.1007/s10584-011-0222-6, ISSN: 0165-0009, 1573–1480.
- Holtmark B. (2013). The outcome is in the assumptions: analyzing the effects on atmospheric CO₂ levels of increased use of bioenergy from forest biomass. *Global Change Biology Bioenergy in print*.
- Hoogwijk M., A. Faaij, B. Eickhout, B. De Vries, and W. Turkenburg (2005). Potential of biomass energy out to 2100, for four IPCC SRES land-use scenarios. *Biomass and Bioenergy* 29, 225–257. doi: 10.1016/j.biombioe.2005.05.002, ISSN: 09619534.
- Hoogwijk M., A. Faaij, B. de Vries, and W. Turkenburg (2009). Exploration of regional and global cost-supply curves of biomass energy from short-rotation crops at abandoned cropland and rest land under four IPCC SRES land-use scenarios. *Biomass and Bioenergy* 33, 26–43. doi: 10.1016/j.biombioe.2008.04.005, ISSN: 0961-9534.

- Hooijer A., S. Page, J.G. Canadell, M. Silvius, J. Kwadijk, H. Wösten, and J. Jauhainen (2010). Current and future CO₂ emissions from drained peatlands in Southeast Asia. *Biogeosciences* **7**, 1505–1514.
- Hook S.E., A. Wright, and B.W. McBride (2010). Methanogens: Methane Producers of the Rumen and Mitigation Strategies. *Archaea* **2010**. doi: 10.1155/2010/945785, ISSN: 1472-3646.
- Van der Horst D., and S. Vermeulen (2011). Spatial scale and social impacts of biofuel production. *Biomass and Bioenergy* **35**, 2435–2443. doi: 10.1016/j.biombioe.2010.11.029, ISSN: 0961-9534.
- Houghton R.A. (2003). Revised estimates of the annual net flux of carbon to the atmosphere from changes in land use and land management 1850–2000. *Tellus B* **55**, 378–390.
- Houghton R.. (2012). Carbon emissions and the drivers of deforestation and forest degradation in the tropics. *Current Opinion in Environmental Sustainability* **4**, 597–603. doi: 10.1016/j.cosust.2012.06.006, ISSN: 1877-3435.
- Houghton R.A. (2013). The emissions of carbon from deforestation and degradation in the tropics: past trends and future potential. *Carbon Management* **4**, 539–546. doi: 10.4155/cmt.13.41, ISSN: 1758-3004.
- Houghton R.A., J.I. House, J. Pongratz, G.R. van der Werf, R.S. DeFries, M.C. Hansen, C. Le Quéré, and N. Ramankutty (2012). Carbon emissions from land use and land-cover change. *Biogeosciences* **9**, 5125–5142. doi: 10.5194/bg-9-5125-2012, ISSN: 1726-4189.
- House J.I., I. Colin Prentice, and C. Le Quéré (2002). Maximum impacts of future reforestation or deforestation on atmospheric CO₂. *Global Change Biology* **8**, 1047–1052.
- Howden S.M., S.J. Crimp, and C.J. Stokes (2008). Climate change and Australian livestock systems: impacts, research and policy issues. *Australian Journal of Experimental Agriculture* **48**, 780–788.
- Hristov A.N., J. Oh, C. Lee, R. Meinen, F. Montes, T. Ott, J. Firkins, A. Rotz, C. Dell, A. Adesogan, W. Yang, J. Tricarico, E. Kebreab, G. Waghorn, J. Dijkstra, and S. Oosting (2013). *Food Value Chain Transformations in Developing Countries*. FAO, Food and Agriculture Organization of the United Nations, Rome, Italy, 210 pp.
- Hsu D.D., D. Inman, G.A. Heath, E. J. Wolfrum, M.K. Mann, and A. Aden (2010). Life Cycle Environmental Impacts of Selected U.S. Ethanol Production and Use Pathways in 2022. *Environmental Science & Technology* **44**, 5289–5297. doi: 10.1021/es100186h, ISSN: 0013-936X.
- Hu Z., J.W. Lee, K. Chandran, S. Kim, and S.K. Khanal (2012). Nitrous Oxide (N₂O) Emission from Aquaculture: A Review. *Environmental Science & Technology* **46**, 6470–6480. doi: 10.1021/es300110x, ISSN: 0013-936X, 1520–5851.
- Hu Z., J.W. Lee, K. Chandran, S. Kim, K. Sharma, A.C. Brotto, and S.K. Khanal (2013). Nitrogen transformations in intensive aquaculture system and its implication to climate change through nitrous oxide emission. *Bioresource Technology* **130**, 314–320. doi: 10.1016/j.biortech.2012.12.033, ISSN: 09608524.
- Huang M., and G.P. Asner (2010). Long-term carbon loss and recovery following selective logging in Amazon forests. *Global Biogeochemical Cycles* **24**, 15. doi: 201010.1029/2009GB003727.
- Huang Y., and Y. Tang (2010). An estimate of greenhouse gas (N₂O and CO₂) mitigation potential under various scenarios of nitrogen use efficiency in Chinese croplands: Mitigating n-induced ghg emission in China. *Global Change Biology* **16**, 2958–2970. doi: 10.1111/j.1365-2486.2010.02187.x, ISSN: 13541013.
- Huang J., J. Yang, S. Msangi, S. Rozelle, and A. Weersink (2012). Biofuels and the poor: Global impact pathways of biofuels on agricultural markets. *Food Policy* **37**, 439–451. doi: 10.1016/j.foodpol.2012.04.004, ISSN: 0306-9192.
- Hudiburg T.W., B.E. Law, C. Wirth, and S. Luysaert (2011). Regional carbon dioxide implications of forest bioenergy production. *Nature Climate Change* **1**, 419–423. doi: 10.1038/nclimate1264, ISSN: 1758-678X.
- Huettner M. (2012). Risks and opportunities of REDD+ implementation for environmental integrity and socio-economic compatibility. *Environmental Science & Policy* **15**, 4–12. doi: 10.1016/j.envsci.2011.10.002, ISSN: 1462-9011.
- Hughes R.F., S.R. Archer, G.P. Asner, C.A. Wessman, C. McMURTRY, J. Nelson, and R.J. Ansley (2006). Changes in aboveground primary production and carbon and nitrogen pools accompanying woody plant encroachment in a temperate savanna. *Global Change Biology* **12**, 1733–1747. doi: 10.1111/j.1365-2486.2006.01210.x, ISSN: 1365-2486.
- Hunsberger C., S. Bolwig, E. Corbera, and F. Creutzig (2012). Livelihood impacts of biofuel crop production: mediating factors and implications for governance. Copenhagen Biofuels Research Network—COBREN, Copenhagen, 19–20 November 2012.
- Hunsberger C., and T.P. Evans (2012). Chapter 3—Land. In: *Fifth Global Environment Outlook (GEO 5)*. United Nations Environment Programme (UNEP), London, 65–97.
- Huntingford C., P. Zelazowski, D. Galbraith, L.M. Mercado, S. Sitch, R. Fisher, M. Lomas, A.P. Walker, C.D. Jones, B.B.B. Booth, Y. Malhi, D. Hemming, G. Kay, P. Good, S.L. Lewis, O.L. Phillips, O.K. Atkin, J. Lloyd, E. Gloor, J. Zaragoza-Castells, P. Meir, R. Betts, P.P. Harris, C. Nobre, J. Marengo, and P.M. Cox (2013). Simulated resilience of tropical rainforests to CO₂-induced climate change. *Nature Geoscience* **6**, 268–273. doi: 10.1038/ngeo1741, ISSN: 1752-0894.
- Hurt G., L. Chini, S. Frolking, R. Betts, J. Feddema, G. Fischer, J. Fisk, K. Hibbard, R. Houghton, A. Janetos, C. Jones, G. Kindermann, T. Kinoshita, K. Klein Goldewijk, K. Riahi, E. Shevliakova, S. Smith, E. Stehfest, A. Thomson, P. Thornton, D. van Vuuren, and Y. Wang (2011). Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands. *Climatic Change* **109**, 117–161. doi: 10.1007/s10584-011-0153-2, ISSN: 0165-0009.
- IAASTD (2009). *Agriculture at a Crossroads: Global Report*. International Assessment of Agricultural Knowledge, Science and Technology for Development (IAASTD), 590 pp.
- IBI Biochar (2012). *Standardized Product Definition and Product Testing Guidelines for Biochar That Is Used in Soil | International Biochar Initiative*. International Biochar Initiative, 47 pp. Available at: <http://www.biochar-international.org/characterizationstandard>.
- IEA (2010). *Energy Technology Perspectives 2010. Scenarios & Strategies to 2050*. International Energy Agency, Paris, France, 706 pp.
- IEA (2011). *Energy for All: Financing Access for the Poor. Special Early Excerpt of the World Energy Outlook 2011*. Organisation for Economic Co-Operation and Development/International Energy Agency, Paris, France, 48 pp.
- IEA (2012a). *Tracking Clean Energy Progress. Energy Technology Perspectives 2012 Excerpt as IEA Input to the Clean Energy Ministerial*. Paris, France, 82 pp. Available at: https://www.iea.org/publications/freepublications/publication/Tracking_Clean_Energy_Progress.pdf.

- IEA (2012b).** *World Energy Outlook 2012*. OECD/IEA, Paris, France, 690 pp. Available at: <http://www.worldenergyoutlook.org/publications/weo-2012/#den.26099>.
- IEA (2012c).** *Technology Roadmaps: Bioenergy for Heat and Power*. International Energy Agency, Paris, France, 62 pp.
- IEA (2013).** *World Energy Outlook 2013*. OECD, Paris, France, 708 pp. Available at: <http://www.worldenergyoutlook.org/publications/weo-2013/>.
- IEA-ETSAP, and IRENA (2013).** *Production of Bio-Ethylene: Technology Brief*. International Renewable Energy Agency and Energy Technology Systems Analysis Programme of the International Energy Agency, 20 pp. Available at: http://www.irena.org/DocumentDownloads/Publications/IRENA-ETSAP%20Tech%20Brief%20113%20Production_of_Bio-ethylene.pdf.
- Imhoff M. L., L. Bounoua, T. Ricketts, C. Loucks, R. Harriss, and W.T. Lawrence (2004).** Global patterns in human consumption of net primary production. *Nature* **429**, 870–873. doi: 10.1038/nature02619, ISSN: 0028-0836.
- Immerzeel D.J., P. Verweij, F. Hilst, and A.P. Faaij (2013).** Biodiversity impacts of bioenergy crop production: a state-of-the-art review. *GCB Bioenergy*.
- Ingerson A. (2011).** Carbon storage potential of harvested wood: summary and policy implications. *Mitigation and Adaptation Strategies for Global Change* **16**, 307–323. doi: 10.1007/s11027-010-9267-5, ISSN: 1381-2386, 1573–1596.
- IPCC (1996).** *Climate Change 1995: The Science of Climate Change. Contribution of Working Group I to the Second Assessment Report of the Intergovernmental Panel on Climate Change* [J.T. Houghton, L.G. Meira Filho, B.A. Callander, N. Harris, A. Kattenberg, K. Maskell (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY USA, 588 pp. ISBN: 0 521 56436 0.
- IPCC (2000).** *Special Report on Land Use, Land Use Change and Forestry* [R.T. Watson, I. Noble, B. Bolin, N.H. Ravindranath, D.J. Verardo, and D.J. Dokken (eds.)]. Cambridge University Press, Cambridge, MA, 377 pp.
- IPCC (2001).** Summary for Policy Makers. In: *Climate Change 2001: Impacts Adaptation and Vulnerability. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change* [J.J. McCarthy, O.F. Canziani, N.A. Leary, D.J. Dokken, K.S. White (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. Available at: http://ipcc.ch/publications_and_data/publications_and_data_reports.shtml.
- IPCC (2003).** *Good Practice Guidance for Land Use, Land-Use Change and Forestry* [J. Penman, M. Gytarsky, T. Hiraishi, T. Krug, D. Kruger, R. Pipatti, L. Buendia, K. Miwa, T. Ngara, K. Tanabe, F. Wagner (eds.)]. IPCC/OECD/IEA/IGES, Hayama, Japan, 580 pp.
- IPCC (2006).** *2006 National Greenhouse Gas Inventory Guidelines* [S. Eggleston, L. Buendia, K. Miwa, T. Ngara, K. Tanabe (eds.)]. Institute of Global Environmental Strategies (IGES), Kanagawa, Japan, 20 pp.
- IPCC (2007a).** *Climate Change Mitigation: Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [B. Metz, O.R. Davidson, P.R. Bosch, R. Dave, L.A. Meyer (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 862pp. Available at: <http://www.ipcc.ch/pdf/assessment-report/ar4/wg3/ar4-wg3-index.pdf>.
- IPCC (2007b).** *Impacts, Adaptation and Vulnerability: Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [M.L. Parry, O.F. Canziani, J.P. Palutikof, P.J. van der Linden and C.E. Hanson (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 986pp.
- IPCC (2011).** *IPCC Special Report on Renewable Energy Sources and Climate Change Mitigation*. Prepared by Working Group III of the Intergovernmental Panel on Climate Change [O. Edenhofer, R. Pichs-Madruga, Y. Sokona, K. Seyboth, P. Matschoss, S. Kadner, T. Zwickel, P. Eickemeier, G. Hansen, S. Schlömer, C. von Stechow (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1075 pp.
- IPCC (2012).** *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation*. A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change [Field, C.B., V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.D. Mastrandrea, K.J. Mach, G.-K. Plattner, S.K. Allen, M. Tignor, and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, UK, and New York, NY, USA, 582 pp.
- Ito A. (2005).** Modelling of carbon cycle and fire regime in an east Siberian larch forest. *Ecological Modelling* **187**, 121–139. doi: 10.1016/j.ecolmodel.2005.01.037, ISSN: 03043800.
- ITTO (2008).** *Developing Forest Certification: Towards Increasing the Comparability and Acceptance of Forest Certification Systems Worldwide*. International Tropical Timber Organization, Yokohama, Japan, 126 pp. ISBN: 4902045397 9784902045390.
- Jackson T. (2009).** *Prosperity without Growth. Economics for a Finite Planet*. Earthscan, UK and USA, 264 pp. ISBN: 1844078949.
- Jackson R.B., and J.S. Baker (2010).** Opportunities and Constraints for Forest Climate Mitigation. *BioScience* **60**, 698–707. doi: 10.1525/bio.2010.60.9.7, ISSN: 0006-3568, 1525–3244.
- Jackson R.B., K.A. Farley, W.A. Hoffman, E.G. Jobbágy, and R.L. McCulley (2007).** Carbon and water tradeoffs in conversions to forests and shrublands. *Global Change—The IGBP Series*. In: *Terrestrial Ecosystems in a Changing World*. J.G. Canadell, D.E. Pataki, L.F. Pitelka, (eds.), Springer, Berlin, Germany, pp. 237–246. ISBN: 978-3-540-32730-1.
- Jackson R.B., E.G. Jobbágy, R. Avissar, S.B. Roy, D.J. Barrett, C.W. Cook, K.A. Farley, D.C. Le Maitre, B.A. McCarl, and B.C. Murray (2005).** Trading water for carbon with biological carbon sequestration. *Science* **310**, 1944.
- Jackson R.B., J.T. Randerson, J.G. Canadell, R.G. Anderson, R. Avissar, D.D. Baldocchi, G.B. Bonan, K. Caldeira, N.S. Diffenbaugh, C.B. Field, and others (2008).** Protecting climate with forests. *Environmental Research Letters* **3**, 044006. ISSN: 1748-9326.
- Jain A.K., P. Meiyappan, Y. Song, and J.I. House (2013).** CO₂ emissions from land-use change affected more by nitrogen cycle, than by the choice of land-cover data. *Global Change Biology* **19**, 2893–2906. doi: 10.1111/gcb.12207, ISSN: 13541013.
- Jakobsson E.B., E.B. Sommer, P. De Clercq, G. Bonazzi, and B. Schröder (2002).** The policy implementation of nutrient management legislation and effects in some European Countries. In: *The Final Workshop of the EU Concerted Action Nutrient Management Legislation in European Countries NUMALEC*. Pand. Oderberg, Gent, Belgium.
- Janssen P.H., and M. Kirs (2008).** Structure of the Archaeal Community of the Rumen. *Applied and Environmental Microbiology* **74**, 3619–3625. doi: 10.1128/AEM.02812-07, ISSN: 0099-2240.
- Janzen H.H. (2011).** What place for livestock on a re-greening earth? *Animal Feed Science and Technology* **166–67**, 783–796. doi: 10.1016/j.anifeedsci.2011.04.055, ISSN: 0377-8401.

- Jaramillo P., W.M. Griffin, and H.S. Matthews (2008). Comparative Analysis of the Production Costs and Life-Cycle GHG Emissions of FT Liquid Fuels from Coal and Natural Gas. *Environmental Science & Technology* **42**, 7559–7565. doi: 10.1021/es8002074, ISSN: 0013-936X, 1520–5851.
- Jeffery S., F.G.A. Verheijen, M. van der Velde, and A.C. Bastos (2011). A quantitative review of the effects of biochar application to soils on crop productivity using meta-analysis. *Agriculture, Ecosystems & Environment* **144**, 175–187. doi: 10.1016/j.agee.2011.08.015, ISSN: 0167-8809.
- Jerneck A., and L. Olsson (2013). A smoke-free kitchen: initiating community based co-production for cleaner cooking and cuts in carbon emissions. *Journal of Cleaner Production* **60**, 208–215. doi: 10.1016/j.jclepro.2012.09.026, ISSN: 0959-6526.
- Jetter J., Y. Zhao, K.R. Smith, B. Khan, T. Yelverton, P. DeCarlo, and M.D. Hays (2012). Pollutant Emissions and Energy Efficiency under Controlled Conditions for Household Biomass Cookstoves and Implications for Metrics Useful in Setting International Test Standards. *Environmental Science & Technology* **46**, 10827–10834. doi: 10.1021/es301693f, ISSN: 0013-936X.
- Johansson T.B., N. Nakicenovic, A. Patwardhan, L. Gomez-Echeverri, and W.C. Turkenburg (2012). Summary for policymakers. *Global Energy Assessment—Toward a Sustainable Future*, 3–30.
- Johnston M., J.A. Foley, T. Holloway, C. Kucharik, and C. Monfreda (2009a). Resetting global expectations from agricultural biofuels. *Environmental Research Letters* **4**, 014004 (9pp). doi: 10.1088/1748-9326/4/1/014004, ISSN: 1748-9326.
- Johnston M., R. Licker, J.A. Foley, T. Holloway, N.D. Mueller, C. Barford, and C.J. Kucharik (2011). Closing the gap: global potential for increasing biofuel production through agricultural intensification. *Environmental Research Letters* **6**.
- Johnston A.E., P.R. Poulton, and K. Coleman (2009b). Chapter 1 Soil Organic Matter. In: *Advances in Agronomy*. Elsevier, pp. 1–57. ISBN: 9780123748171.
- Jones C.S., and S.P. Mayfield (2012). Algae biofuels: versatility for the future of bioenergy. *Current Opinion in Biotechnology* **23**, 346–351. doi: 10.1016/j.cop-bio.2011.10.013, ISSN: 0958-1669.
- Jonker J.G.G., and A.P.C. Faaij (2013). Techno-economic assessment of microalgae as feedstock for renewable bio-energy production. *Applied Energy* **102**, 461–475.
- Jonker J.G.G., M. Junginger, and A. Faaij (2013). Carbon payback period and carbon offset parity point of wood pellet production in the South-eastern United States. *GCB Bioenergy*, n/a–n/a. doi: 10.1111/gcbb.12056, ISSN: 1757-1707.
- Joos F., R. Roth, J.S. Fuglestedt, G.P. Peters, I.G. Enting, W. von Bloh, V. Brovkin, E.J. Burke, M. Eby, N.R. Edwards, T. Friedrich, T.L. Frölicher, P.R. Halloran, P.B. Holden, C. Jones, T. Kleinen, F. Mackenzie, K. Matsumoto, M. Meinshausen, G.-K. Plattner, A. Reisinger, J. Segschneider, G. Shaffer, M. Steinacher, K. Strassmann, K. Tanaka, A. Timmermann, and A.J. Weaver (2013). Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis. *Atmospheric Chemistry and Physics Discussions* **12**, 19799–19869. doi: 10.5194/acpd-12-19799-2012, ISSN: 1680-7375.
- Joosten H. (2010). *The Global Peatland CO₂ Picture: Peatland Status and Drainage Related Emissions in All Countries of the World*. Wetlands International, Wageningen, Netherlands, 36 pp. Available at: [http://www.wetlands.org/WatchRead/Currentpublications/tabid/56/mod/1570/articleType/ArticleView/articledId/2418/The-Global-Peatland-CO₂-Picture.aspx](http://www.wetlands.org/WatchRead/Currentpublications/tabid/56/mod/1570/articleType/ArticleView/articledId/2418/The-Global-Peatland-CO2-Picture.aspx).
- JRC/PBL (2013). *Emission Database for Global Atmospheric Research (EDGAR), Release Version 4.2 FT2010*. European Commission, Joint Research Centre (JRC)/PBL Netherlands Environmental Assessment Agency, Available at: <http://edgar.jrc.ec.europa.eu>.
- Jull C., P.C. Redondo, V. Mosoti, and J. Vapnek (2007). *Recent Trends in the Law and Policy of Bioenergy Production, Promotion and Use*. FAO, Food and Agriculture Organization of the United Nations, Rome, Italy, 55 pp.
- Junginger M., J. van Dam, S. Zarrilli, F. Ali Mohamed, D. Marchal, and A. Faaij (2011). Opportunities and barriers for international bioenergy trade. *Energy Policy* **39**, 2028–2042. doi: 16/j.enpol.2011.01.040, ISSN: 0301-4215.
- Junginger M., C.S. Goh, and A. Faaij (2014). *International Bioenergy Trade: History, Status & Outlook on Securing Sustainable Bioenergy Supply, Demand and Markets*. 233 pp. ISBN: 978-94-007-6982-3.
- Kaiser E.-A., and R. Ruser (2000). Nitrous oxide emissions from arable soils in Germany—An evaluation of six long-term field experiments. *Journal of Plant Nutrition and Soil Science* **163**, 249–259. doi: 10.1002/1522-2624(200006)163:3< 249::AID-JPLN249> 3.0.CO;2-Z, ISSN: 1522-2624.
- Kandji S.T., L.V. Verchot, and J. Mackensen (2006). *Climate Change and Variability in the Sahel Region: Impacts and Adaptation Strategies in the Agricultural Sector*. World Agroforestry Centre (ICRAF)/United Nations Environment Programme (UNEP), Nairobi, Kenya, 58 pp.
- Kaphengst T., M.S. Ma, and S. Schlegel (2009). At a Tipping Point? How the Debate on Biofuel Standards Sparks Innovative Ideas for the General Future of Standardization and Certification Schemes. *Journal of Cleaner Production* **17**, 99–101.
- Kar A., I.H. Rehman, J. Burney, S.P. Puppala, R. Suresh, L. Singh, V.K. Singh, T. Ahmed, N. Ramanathan, and V. Ramanathan (2012). Real-time assessment of black carbon pollution in Indian households due to traditional and improved biomass cookstoves. *Environmental Science & Technology* **46**, 2993–3000.
- Karipidis P., E. Tsakiridou, S. Aggelopoulos, and A. Belidis (2010). Consumers' purchasing and store switching intentions in respect of eco-marked products. *International Journal of Economics and Business Research* **2**, 511–524. doi: 10.1504/IJEBR.2010.035702, ISSN: 1756-9850.
- Karsenty A., A. Vogel, and F. Castell (2014). "Carbon rights", REDD+ and payments for environmental services. *Environmental Science & Policy* **35**, 20–29. doi: 10.1016/j.envsci.2012.08.013, ISSN: 1462-9011.
- Kastner T., K.-H. Erb, and S. Nonhebel (2011a). International wood trade and forest change: A global analysis. *Global Environmental Change* **21**, 947–956. doi: 16/j.gloenvcha.2011.05.003, ISSN: 0959-3780.
- Kastner T., M. Kastner, and S. Nonhebel (2011b). Tracing distant environmental impacts of agricultural products from a consumer perspective. *Ecological Economics* **70**, 1032–1040. doi: 16/j.ecolecon.2011.01.012, ISSN: 0921-8009.
- Kastner T., M.J.I. Rivas, W. Koch, and S. Nonhebel (2012). Global changes in diets and the consequences for land requirements for food. *Proceedings of the National Academy of Sciences* **109**, 6868–6872. doi: 10.1073/pnas.1117054109, ISSN: 0027-8424, 1091–6490.

- Kataeva I., M. B. Foston, S.-J. Yang, S. Pattathil, A. K. Biswal, F. L. P. li, M. Basen, A. M. Rhaesa, T. P. Thomas, P. Azadi, V. Olman, T. D. Saffold, K. E. Mohler, D. L. Lewis, C. Doepcke, Y. Zeng, T. J. Tschaplinski, W. S. York, M. Davis, D. Mohnen, Y. Xu, A. J. Ragauskas, S.-Y. Ding, R. M. Kelly, M. G. Hahn, and M. W. W. Adams (2013). Carbohydrate and lignin are simultaneously solubilized from unpretreated switchgrass by microbial action at high temperature. *Energy & Environmental Science* **6**, 2186–2195. doi: 10.1039/C3EE40932E, ISSN: 1754-5706.
- Katila P. (2008). *Devolution of Forest-Related Rights: Comparative Analyses of Six Developing Countries*. University of Helsinki, Vikki Tropical Resources Institute, Helsinki, Finland, 162 pp. ISBN: 978-952-10-4518-9.
- Kato E., T. Kinoshita, A. Ito, M. Kawamiya, and Y. Yamagata (2011). Evaluation of spatially explicit emission scenario of land-use change and biomass burning using a process-based biogeochemical model. *Journal of Land Use Science* **8**, 104–122. doi: 10.1080/1747423X.2011.628705, ISSN: 1747-423X.
- Kaufman A. S., P. J. Meier, J. C. Sinistore, and D. J. Reinemann (2010). Applying life-cycle assessment to low carbon fuel standards—How allocation choices influence carbon intensity for renewable transportation fuels. *Energy Policy* **38**, 5229–5241. doi: 10.1016/j.enpol.2010.05.008, ISSN: 0301-4215.
- Keating B. A., P. S. Carberry, and J. Dixon (2013). Agricultural intensification and the food security challenge in Sub Saharan Africa. In: *Agro-Ecological Intensification of Agricultural Systems in the African Highlands*. Routledge, US and Canada, pp. 20–35. ISBN: 978-0-415-53273-0.
- Keeling L. J., J. Rushen, and I. J. H. Duncan (2011). Understanding animal welfare. In: *Animal welfare*. M. C. Appleby, J. A. Mench, I. a. S. Olsson, B. O. Hughes, (eds.), CABI, Wallingford, UK, pp. 13–26. ISBN: 978-1-84593-659-4.
- Kehlbacher A., R. Bennett, and K. Balcombe (2012). Measuring the consumer benefits of improving farm animal welfare to inform welfare labelling. *Food Policy* **37**, 627–633. doi: 10.1016/j.foodpol.2012.07.002, ISSN: 0306-9192.
- Van Kessel C., R. Venterea, J. Six, M. A. Adviento-Borbe, B. Linquist, and K. J. van Groenigen (2013). Climate, duration, and N placement determine N₂O emissions in reduced tillage systems: a meta-analysis. *Global Change Biology* **19**, 33–44. doi: 10.1111/j.1365-2486.2012.02779.x, ISSN: 13541013.
- Khanna M., C. L. Crago, and M. Black (2011). Can biofuels be a solution to climate change? The implications of land use change-related emissions for policy. *Interface Focus* **1**, 233–247. doi: 10.1098/rsfs.2010.0016, ISSN: 2042-8898, 2042–8901.
- Kim D.-G., G. Hernandez-Ramirez, and D. Giltrap (2013). Linear and nonlinear dependency of direct nitrous oxide emissions on fertilizer nitrogen input: A meta-analysis. *Agriculture, Ecosystems & Environment* **168**, 53–65. doi: 10.1016/j.agee.2012.02.021, ISSN: 01678809.
- Kim M. K., and B. A. McCarl (2009). Uncertainty discounting for land-based carbon sequestration. *Journal of Agricultural and Applied Economics* **41**, 1–11.
- Kiptot E., and S. Franzel (2012). Gender and agroforestry in Africa: a review of women's participation. *Agroforest Systems* **84**, 35–58. doi: 10.1007/s10457-011-9419-y.
- De Klein C. A. M., and R. J. Eckard (2008). Targeted technologies for nitrous oxide abatement from animal agriculture. *Australian Journal of Experimental Agriculture* **48**, 14–20. Available at: <http://dx.doi.org/10.1071/EA07217>.
- De Klein C. A. M., R. J. Eckard, and T. J. van der Weerden (2010). Nitrous Oxide Emissions from the Nitrogen Cycle in Livestock Agriculture: Estimation and Mitigation. In: *Nitrous Oxide and Climate Change*. K. Smith, (ed.), Earthscan Publications, London, pp. 107–144. ISBN: 978-1-84407-757-1.
- Kløverpris J. H., and S. Mueller (2013). Baseline time accounting: Considering global land use dynamics when estimating the climate impact of indirect land use change caused by biofuels. *The International Journal of Life Cycle Assessment* **18**, 319–330.
- Kløverpris J., H. Wenzel, M. Banse, L. M. i Canals, and A. Reenberg (2008). Conference and workshop on modelling global land use implications in the environmental assessment of biofuels. *The International Journal of Life Cycle Assessment* **13**, 178–183. doi: 10.1065/lca2008.03.381, ISSN: 0948-3349, 1614–7502.
- Kochsiek A. E., and J. M. Knops (2012). Maize cellulosic biofuels: soil carbon loss can be a hidden cost of residue removal. *GCB Bioenergy* **4**, 229–233.
- Koehler K. A., S. M. Kreidenweis, P. J. DeMott, M. D. Petters, A. J. Prenni, and O. Möhler (2010). Laboratory investigations of the impact of mineral dust aerosol on cold cloud formation. *Atmos. Chem. Phys.* **10**, 11955–11968. doi: 10.5194/acp-10-11955-2010, ISSN: 1680-7324.
- Koh L. P., and J. Ghazoul (2008). Biofuels, biodiversity, and people: Understanding the conflicts and finding opportunities. *Biological Conservation* **141**, 2450–2460. doi: 10.1016/j.biocon.2008.08.005, ISSN: 0006-3207.
- Koh L. P., and D. S. Wilcove (2008). Is oil palm agriculture really destroying tropical biodiversity? *Conservation Letters* **1**, 60–64. doi: 10.1111/j.1755-263X.2008.00011.x, ISSN: 1755-263X.
- Koizumi T. (2013). Biofuel and food security in China and Japan. *Renewable and Sustainable Energy Reviews* **21**, 102–109. doi: 10.1016/j.rser.2012.12.047, ISSN: 1364-0321.
- Koknaroglu H., and T. Akunal (2013). Animal welfare: An animal science approach. *59th International Congress of Meat Science and Technology, 18–23 August 2013 Izmir/Turkey* **95**, 821–827. doi: 10.1016/j.meatsci.2013.04.030, ISSN: 0309-1740.
- Kollmuss A., M. Lazarus, C. M. Lee, M. LeFranc, and C. Polycarp (2010). *Handbook of Carbon Offset Programs Trading Systems, Funds, Protocols and Standards*. Earthscan, London, 240 pp. ISBN: 978-1-84407-929-2(Hb).
- Kontur W. S., D. R. Noguera, and T. J. Donohue (2012). Maximizing reductant flow into microbial H₂ production. *Current Opinion in Biotechnology* **23**, 382–389. doi: 10.1016/j.copbio.2011.10.003, ISSN: 0958-1669.
- Koornneef J., P. van Breevoort, C. Hamelinck, C. Hendriks, M. Hoogwijk, K. Koop, M. Koper, T. Dixon, and A. Camps (2012). Global potential for biomass and carbon dioxide capture, transport and storage up to 2050. *International Journal of Greenhouse Gas Control* **11**, 117–132. doi: 10.1016/j.ijggc.2012.07.027, ISSN: 1750-5836.
- Koornneef J., P. van Breevoort, P. Noothout, C. Hendriks, uchien Luning, and A. Camps (2013). Global Potential for Biomethane Production with Carbon Capture, Transport and Storage up to 2050. *GHGT-11* **37**, 6043–6052. doi: 10.1016/j.egypro.2013.06.533, ISSN: 1876-6102.
- Körner C. (2006). Plant CO₂ responses: an issue of definition, time and resource supply. *New Phytologist* **172**, 393–411. doi: 10.1111/j.1469-8137.2006.01886.x, ISSN: 0028-646X, 1469–8137.
- Körner C. (2009). Biologische Kohlenstoffsinken: Umsatz und Kapital nicht verwechseln (Biological Carbon Sinks: Turnover Must Not Be Confused with Capital). *Gaia—Ecological Perspectives for Science and Society* **18**, 288–293.
- Krausmann F., K.-H. Erb, S. Gingrich, H. Haberl, A. Bondeau, V. Gaube, C. Lauk, C. Plutzer, and T. Searchinger (2013). Global human appropriation of net primary production doubled in the 20th century. *Proceedings of the National Academy of Sciences of the USA* submitted.

- Krausmann F., K.-H. Erb, S. Gingrich, C. Lauk, and H. Haberl (2008). Global patterns of socioeconomic biomass flows in the year 2000: A comprehensive assessment of supply, consumption and constraints. *Ecological Economics* 65, 471–487. doi: 16/j.ecolecon.2007.07.012, ISSN: 0921-8009.
- Krausmann F., S. Gingrich, H. Haberl, K.-H. Erb, A. Musel, T. Kastner, N. Kohlhebb, M. Niedertscheider, and E. Schwarzmüller (2012). Long-term trajectories of the human appropriation of net primary production: Lessons from six national case studies. *Ecological Economics* 77, 129–138. doi: 10.1016/j.ecolecon.2012.02.019, ISSN: 0921-8009.
- Kriegler E., P. Weyant, G. Blanford, V. Krey, L. Clarke, J. Edmonds, A. Fawcett, G. Luderer, K. Riahi, R. Richels, S.K. Rose, M. Tavani, and D.P. van Vuuren (2013). The role of technology for achieving climate policy objectives: overview of the EMF27 study on global technology and climate policy strategies. *Climatic Change* in press. doi: 10.1007/s10584-013-0953-7.
- Kummu M., H. de Moel, M. Porkka, S. Siebert, O. Varis, and P.J. Ward (2012). Lost food, wasted resources: Global food supply chain losses and their impacts on freshwater, cropland, and fertiliser use. *Science of The Total Environment* 438, 477–489. doi: 10.1016/j.scitotenv.2012.08.092, ISSN: 0048-9697.
- Kurz W.A., G. Stinson, G.J. Rampley, C.C. Dymond, and E.T. Neilson (2008). Risk of natural disturbances makes future contribution of Canada's forests to the global carbon cycle highly uncertain. *Proceedings of the National Academy of Sciences* 105, 1551.
- Kyu H.H., K. Georgiades, and M.H. Boyle (2010). Biofuel Smoke and Child Anemia in 29 Developing Countries: A Multilevel Analysis. *Annals of Epidemiology* 20, 811–817. doi: 10.1016/j.annepidem.2010.07.096, ISSN: 1047-2797.
- Laganieri J., D.A. Angers, and D. Pare (2010). Carbon accumulation in agricultural soils after afforestation: a meta-analysis: SOC accumulation following deforestation. *Global Change Biology* 16, 439–453. doi: 10.1111/j.1365-2486.2009.01930.x, ISSN: 13541013, 13652486.
- Laitner J.A.S., S.J. DeCanio, and I. Peters (2000). Incorporating Behavioral, Social and Organizational Phenomena in the Assessment of Climate Change Mitigation Options. In: *Society, Behaviour and Climate Change Mitigation*. E. Jochem, (ed.), Kluwer, Amsterdam, pp. 1–64.
- Lal R. (2010). Managing soils for a warming earth in a food-insecure and energy-starved world. *Journal of Plant Nutrition and Soil Science* 173, 4–15. doi: 10.1002/jpln.200900290, ISSN: 1522-2624.
- Lal R. (2011). Sequestering carbon in soils of agro-ecosystems. *Food Policy* 36, 533–539. doi: 10.1016/j.foodpol.2010.12.001, ISSN: 03069192.
- Lam M.K., K.T. Tan, K.T. Lee, and A.R. Mohamed (2009). Malaysian palm oil: Surviving the food versus fuel dispute for a sustainable future. *Renewable and Sustainable Energy Reviews* 13, 1456–1464. doi: 10.1016/j.rser.2008.09.009, ISSN: 1364-0321.
- Lambin E.F., and P. Meyfroidt (2011). Global land use change, economic globalization, and the looming land scarcity. *Proceedings of the National Academy of Sciences* 108, 3465–3472. doi: 10.1073/pnas.1100480108, ISSN: 0027-8424, 1091–6490.
- Lamers P., and M. Junginger (2013). The “debt” is in the detail: A synthesis of recent temporal forest carbon analyses on woody biomass for energy. *Biofuels, Bioproducts and Biorefining* 7, 373–385. doi: 10.1002/bbb.1407, ISSN: 1932-1031.
- Lamers P., H.M. Junginger, C.C. Dymond, and A. Faaij (2013). Damaged forests provide an opportunity to mitigate climate change. *GCB Bioenergy* in press.
- Landis D.A., M.M. Gardiner, W. Van Der Werf, and S.M. Swinton (2008). Increasing corn for biofuel production reduces biocontrol services in agricultural landscapes. *Proceedings of the National Academy of Sciences* 105, 20552–20557. Available at: <http://www.pnas.org/content/105/51/20552.short>.
- Langeveld J.W., J. Dixon, H. van Keulen, and P.M. Quist-Wessel (2013). Analyzing the effect of biofuel expansion on land use in major producing countries: evidence of increased multiple cropping. *Biofuels, Bioproducts and Biorefining* 8, 49–58. Available at: <http://onlinelibrary.wiley.com/doi/10.1002/bbb.1432/abstract>.
- Lansche J., and J. Müller (2012). Life cycle assessment of energy generation of biogas fed combined heat and power plants: Environmental impact of different agricultural substrates. *Engineering in Life Sciences* 12, 313–320. doi: 10.1002/elsc.201100061, ISSN: 1618-2863.
- Larson A.M. (2011). Forest tenure reform in the age of climate change: Lessons for REDD+. *Global Environmental Change* 21, 540–549. doi: 10.1016/j.gloenvcha.2010.11.008, ISSN: 0959-3780.
- Larson A.M., M. Brockhaus, W.D. Sunderlin, A. Duchelle, A. Babon, T. Dokken, T.T. Pham, I.A.P. Resosudarmo, G. Selaya, A. Awono, and T.-B. Huynh (2013). Land tenure and REDD+: The good, the bad and the ugly. *Global Environmental Change* 23, 678–689. doi: 10.1016/j.gloenvcha.2013.02.014, ISSN: 0959-3780.
- Larson E.D., Z. Li, and R.H. Williams (2012). Chapter 12—Fossil Energy. In: *Global Energy Assessment—Toward a Sustainable Future*. Cambridge University Press, Cambridge, UK and New York, NY, USA and the International Institute for Applied Systems Analysis, Laxenburg, Austria, pp. 901–992. ISBN: 052118293X.
- Latta G.S., J.S. Baker, R.H. Beach, S.K. Rose, and B.A. McCarl (2013). A multi-sector intertemporal optimization approach to assess the GHG implications of U.S. forest and agricultural biomass electricity expansion. *Journal of Forest Economics* 19, 361–383. doi: 10.1016/j.jfe.2013.05.003, ISSN: 1104-6899.
- Lauder A.R., I.G. Enting, J.O. Carter, N. Clisby, A.L. Cowie, B.K. Henry, and M.R. Raupach (2013). Offsetting methane emissions—An alternative to emission equivalence metrics. *International Journal of Greenhouse Gas Control* 12, 419–429. doi: 10.1016/j.ijggc.2012.11.028, ISSN: 17505836.
- Lauk C., H. Haberl, K.-H. Erb, S. Gingrich, and F. Krausmann (2012). Global socioeconomic carbon stocks in long-lived products 1900–2008. *Environmental Research Letters* 7, 34–44. doi: 10.1088/1748-9326/7/3/034023, ISSN: 1748-9326.
- Le Q.B., R. Seidl, and R.W. Scholz (2012). Feedback loops and types of adaptation in the modelling of land-use decisions in an agent-based simulation. *Environmental Modelling & Software* 27–28, 83–96. doi: 10.1016/j.envsoft.2011.09.002, ISSN: 1364-8152.
- Lee S.-J., S.-J. Lee, and D.-W. Lee (2013). Design and development of synthetic microbial platform cells for bioenergy. *Frontiers in Microbiotechnology, Ecotoxicology and Bioremediation* 4, 2–94. doi: 10.3389/fmicb.2013.00092.
- Leemans R., and B. Eickhout (2004). Another reason for concern: regional and global impacts on ecosystems for different levels of climate change. *Global Environmental Change Part A* 14, 219–228.
- Lehmann J. (2007). Bio-energy in the black. *Frontiers in Ecology and the Environment* 5, 381–387. doi: 10.1890/1540-9295(2007)5[381:BITB]2.0.CO;2, ISSN: 1540-9295.

- Lehmann J., J.P. da Silva Jr, C. Steiner, T. Nehls, W. Zech, and B. Glaser (2003). Nutrient availability and leaching in an archaeological Anthrosol and a Ferralsol of the Central Amazon basin: fertilizer, manure and charcoal amendments. *Plant and Soil* **249**, 343–357. Available at: <http://link.springer.com/article/10.1023/A:1022833116184>.
- Lemke R.L., A.J. VandenBygaart, C.A. Campbell, G.P. Lafond, and B. Grant (2010). Crop residue removal and fertilizer N: Effects on soil organic carbon in a long-term crop rotation experiment on a Udic Boroll. *Agriculture, Ecosystems & Environment* **135**, 42–51. doi: 10.1016/j.agee.2009.08.010, ISSN: 01678809.
- Levin K., C. McDermott, and B. Cashore (2008). The climate regime as global forest governance: can reduced emissions from Deforestation and Forest Degradation (REDD) initiatives pass a “dual effectiveness” test? *International Forestry Review* **10**, 538–549. doi: 10.1505/ifer.10.3.538, ISSN: 1465-5489.
- Lewis S.L., G. Lopez-Gonzalez, B. Sonke, K. Affum-Baffoe, T.R. Baker, L.O. Ojo, O.L. Phillips, J.M. Reitsma, L. White, J.A. Comiskey, M.N. Djuikouo, C.E.N. Ewango, T.R. Feldpausch, A.C. Hamilton, M. Gloor, T. Hart, A. Hladik, J. Lloyd, J.C. Lovett, J.R. Makana, Y. Malhi, F.M. Mbago, H.J. Ndagalasi, J. Peacock, K.S.H. Peh, D. Sheil, T. Sunderland, M.D. Swaine, J. Taplin, D. Taylor, S.C. Thomas, R. Votere, and H. Woll (2009). Increasing carbon storage in intact African tropical forests. *Nature* **457**, 1003–U3. doi: 10.1038/nature07771, ISSN: 0028-0836.
- Li H., A.F. Cann, and J.C. Liao (2010). Biofuels: Biomolecular Engineering Fundamentals and Advances. *Annual Review of Chemical and Biomolecular Engineering* **1**, 19–36. doi: 10.1146/annurev-chembioeng-073009-100938.
- Li X., E. Mupondwa, S. Panigrahi, L. Tabil, and P. Adapa (2012). Life cycle assessment of densified wheat straw pellets in the Canadian Prairies. *The International Journal of Life Cycle Assessment* **17**, 420–431. doi: 10.1007/s11367-011-0374-7, ISSN: 0948-3349.
- Liao J.D., T.W. Boutton, and J.D. Jastrow (2006). Storage and dynamics of carbon and nitrogen in soil physical fractions following woody plant invasion of grassland. *Soil Biology and Biochemistry* **38**, 3184–3196. doi: 10.1016/j.soilbio.2006.04.003, ISSN: 00380717.
- Liao J.C., and J. Messing (2012). Energy biotechnology. *Current Opinion in Biotechnology* **23**, 287–289. doi: 10.1016/j.copbio.2012.03.005, ISSN: 0958-1669.
- Lichtfouse E., M. Navarrete, P. Debaeke, V. Souchère, C. Alberola, and J. Ménéassieu (2009). Agronomy for sustainable agriculture. A review. *Agronomy for Sustainable Development* **29**, 1–6. doi: 10.1051/agro:2008054, ISSN: 1774-0746.
- Lim S., T. Vos, A.D. Flaxman, G. Danaei, K. Shibuya, and et al. (2012). A Comparative Risk Assessment of Burden of Disease and Injury Attributable to 67 Risk Factors and Risk Factor Clusters in 21 regions, 1990–2010: A Systematic Analysis for the Global Burden of Disease Study 2010. *The Lancet* **380**, 2224–2260.
- Lindholm E.-L., J. Stendahl, S. Berg, and P.-A. Hansson (2011). Greenhouse gas balance of harvesting stumps and logging residues for energy in Sweden. *Scandinavian Journal of Forest Research* **26**, 586–594. doi: 10.1080/02827581.2011.615337, ISSN: 0282-7581.
- Lippke B., Garcia, J.P., Manrique, and C. (2003). *The Impact of Forests and Forest Management on Carbon Storage*. Rural Technology Initiative, College of Forest Resources, University of Washington, Seattle, 7 pp.
- Lippke B., E. Oneil, R. Harrison, K. Skog, L. Gustavsson, and R. Sathre (2011). Life cycle impacts of forest management and wood utilization on carbon mitigation: knowns and unknowns. *Carbon Management* **2**, 303–333. doi: 10.4155/cmt.11.24, ISSN: 1758-3004.
- Lisboa C.C., K. Butterbach-Bahl, M. Mauder, and R. Kiese (2011). Bioethanol production from sugarcane and emissions of greenhouse gases—known and unknowns. *GCB Bioenergy* **3**, 277–292. doi: 10.1111/j.1757-1707.2011.01095.x, ISSN: 1757-1707.
- Liska A.J., and R.K. Perrin (2009). Indirect land use emissions in the life cycle of biofuels: regulations vs science. *Biofuels, Bioproducts and Biorefining* **3**, 318–328. doi: 10.1002/bbb.153, ISSN: 1932-1031.
- Liu G., E.D. Larson, R.H. Williams, T.G. Kreutz, and X. Guo (2010). Making Fischer–Tropsch fuels and electricity from coal and biomass: performance and cost analysis. *Energy & Fuels* **25**, 415–437.
- Liu X., J. Qu, L. Li, A. Zhang, Z. Jufeng, J. Zheng, and G. Pan (2012). Can biochar amendment be an ecological engineering technology to depress N₂O emission in rice paddies?—A cross site field experiment from South China. *Ecological Engineering* **42**, 168–173. doi: 10.1016/j.ecoleng.2012.01.016, ISSN: 09258574.
- Loarie S.R., D.B. Lobell, G.P. Asner, Q. Mu, and C.B. Field (2011). Direct impacts on local climate of sugar-cane expansion in Brazil. *Nature Climate Change* **1**, 105–109. doi: 10.1038/nclimate1067, ISSN: 1758-678X.
- Lobell D.B., U.L.C. Baldos, and T.W. Hertel (2013). Climate adaptation as mitigation: the case of agricultural investments. *Environmental Research Letters* **8**, 15–28. doi: 10.1088/1748-9326/8/1/015012, ISSN: 1748-9326.
- Lohila A., M. Aurela, J.-P. Tuovinen, and T. Laurila (2004). Annual CO₂ exchange of a peat field growing spring barley or perennial forage grass. *Journal of Geophysical Research: Atmospheres* **109**, n/a–n/a. doi: 10.1029/2004JD004715, ISSN: 2156-2202.
- Lohila A., K. Minkinen, J. Laine, I. Savolainen, J.-P. Tuovinen, L. Korhonen, T. Laurila, H. Tietäväinen, and A. Laaksonen (2010). Forestation of boreal peatlands: Impacts of changing albedo and greenhouse gas fluxes on radiative forcing. *Journal of Geophysical Research* **115**. doi: 10.1029/2010JG001327, ISSN: 0148-0227.
- Lott J.E., C.K. Ong, and C.R. Black (2009). Understorey microclimate and crop performance in a *Grevillea robusta*-based agroforestry system in semi-arid Kenya. *Agricultural and Forest Meteorology* **149**, 1140–1151. Available at: <http://www.sciencedirect.com/science/article/pii/S0168192309000434>.
- Lotze-Campen H., M. von Lampe, P. Kyle, S. Fujimori, P. Havlik, H. van Meijl, T. Hasegawa, A. Popp, C. Schmitz, A. Tabeau, H. Valin, D. Willenbockel, and M. Wise (2013). Impacts of increased bioenergy demand on global food markets: an AgMIP economic model intercomparison. *Agricultural Economics*, n/a–n/a. doi: 10.1111/agec.12092, ISSN: 1574-0862.
- Lotze-Campen H., A. Popp, T. Beringer, C. Müller, A. Bondeau, S. Rost, and W. Lucht (2010). Scenarios of global bioenergy production: The trade-offs between agricultural expansion, intensification and trade. *Ecological Modelling* **221**, 2188–2196.
- Loudermilk E.L., R.M. Scheller, P.J. Weisberg, J. Yang, T.E. Diltz, S.L. Karam, and C. Skinner (2013). Carbon dynamics in the future forest: the importance of long-term successional legacy and climate–fire interactions. *Global Change Biology* **19**, 3502–3515. doi: 10.1111/gcb.12310, ISSN: 1365-2486.
- Lubowski R.N., and S.K. Rose (2013). The Potential for REDD+: Key Economic Modeling Insights and Issues. *Review of Environmental Economics and Policy* **7**, 67–90. doi: 10.1093/reep/res024, ISSN: 1750-6816, 1750–6824.
- Luedeling E., and H. Neufeldt (2012). Carbon sequestration potential of parkland agroforestry in the Sahel. *Climatic Change* **115**, 443–461. doi: 10.1007/s10584-012-0438-0, ISSN: 0165-0009, 1573–1480.

- Lund V., and B. Algers (2003). Research on animal health and welfare in organic farming—a literature review. *Livestock Production Science* **80**, 55–68. doi: 10.1016/S0301-6226(02)00321-4, ISSN: 0301-6226.
- Lunt I.D., L.M. Winsemius, S.P. McDonald, J.W. Morgan, and R.L. Dehaan (2010). How widespread is woody plant encroachment in temperate Australia? Changes in woody vegetation cover in lowland woodland and coastal ecosystems in Victoria from 1989 to 2005. *Journal of Biogeography* **37**, 722–732. doi: 10.1111/j.1365-2699.2009.02255.x, ISSN: 1365-2699.
- Luo Y., M. Durenkamp, M. De Nobili, Q. Lin, and P.C. Brookes (2011). Short term soil priming effects and the mineralisation of biochar following its incorporation to soils of different pH. *Soil Biology and Biochemistry* **43**, 2304–2314. doi: 10.1016/j.soilbio.2011.07.020, ISSN: 0038-0717.
- Luo L., E. Voet, G. Huppel, and H. Udo de Haes (2009). Allocation issues in LCA methodology: a case study of corn stover-based fuel ethanol. *The International Journal of Life Cycle Assessment* **14**, 529–539. doi: 10.1007/s11367-009-0112-6, ISSN: 0948-3349.
- Luyssaert S., E.-D. Schulze, A. Börner, A. Knohl, D. Hessenmöller, B.E. Law, P. Ciais, and J. Grace (2008). Old-growth forests as global carbon sinks. *Nature* **455**, 213–215. doi: 10.1038/nature07276, ISSN: 0028-0836.
- Lynd L.R., R.A. Aziz, C.H. de Brito Cruz, A.F.A. Chimphango, L.A.B. Cortez, A. Faaij, N. Greene, M. Keller, P. Osseweijer, and T.L. Richard (2011). A global conversation about energy from biomass: the continental conventions of the global sustainable bioenergy project. *Interface Focus* **1**, 271–279.
- Ma J., E. Ma, H. Xu, K. Yagi, and Z. Cai (2009). Wheat straw management affects CH₄ and N₂O emissions from rice fields. *Soil Biology and Biochemistry* **41**, 1022–1028. doi: 10.1016/j.soilbio.2009.01.024, ISSN: 00380717.
- Macauley M.K., and R.A. Sedjo (2011). Forests in climate policy: technical, institutional and economic issues in measurement and monitoring. *Mitigation and Adaptation Strategies for Global Change* **16**, 499–513. doi: 10.1007/s11027-010-9276-4, ISSN: 1381-2386, 1573–1596.
- Macedo I.C. (2005). *Sugar Cane's Energy: Twelve Studies on Brazilian Sugar Cane Agribusiness and Its Sustainability*. UNICA, Sao Paulo, Brazil, 237 pp.
- Macedo M.N., R.S. DeFries, D.C. Morton, C.M. Stickler, G.L. Galford, and Y.E. Shimabukuro (2012). Decoupling of deforestation and soy production in the southern Amazon during the late 2000s. *Proceedings of the National Academy of Sciences* **109**, 1341–1346. doi: 10.1073/pnas.1111374109, ISSN: 0027-8424, 1091–6490.
- Macedo I.C., J.E.A. Seabra, and J.E.A.R. Silva (2008). Green house gases emissions in the production and use of ethanol from sugarcane in Brazil: The 2005/2006 averages and a prediction for 2020. *Biomass and Bioenergy* **32**, 582–595. doi: 10.1016/j.biombioe.2007.12.006, ISSN: 0961-9534.
- Machmuller A., C.R. Soliva, and M. Kreuzer (2003). Methane-suppressing effect of myristic acid in sheep as affected by dietary calcium and forage proportion. *British Journal of Nutrition* **90**, 529–540.
- Mackey B., I.C. Prentice, W. Steffen, J.I. House, D. Lindenmayer, H. Keith, and S. Berry (2013). Untangling the confusion around land carbon science and climate change mitigation policy. *Nature Climate Change* **3**, 552–557. doi: 10.1038/nclimate1804, ISSN: 1758-678X.
- Macmillan Uribe A.L., D.M. Winham, and C.M. Wharton (2012). Community supported agriculture membership in Arizona. An exploratory study of food and sustainability behaviours. *Appetite* **59**, 431–436. doi: 10.1016/j.appet.2012.06.002, ISSN: 0195-6663.
- Madlener R., C. Robledo, B. Muys, and J.T.B. Freja (2006). A Sustainability Framework for Enhancing the Long-Term Success of Lulucf Projects. *Climatic Change* **75**, 241–271. doi: 10.1007/s10584-005-9023-0, ISSN: 0165-0009, 1573–1480.
- Mahmood R., R.A. Pielke, K.G. Hubbard, D. Niyogi, P.A. Dirmeyer, C. McAlpine, A.M. Carleton, R. Hale, S. Gameda, A. Beltrán-Przekurat, B. Baker, R. McNider, D.R. Legates, M. Shepherd, J. Du, P.D. Blanken, O.W. Frauenfeld, U.s. Nair, and S. Fall (2013). Land cover changes and their biogeophysical effects on climate. *International Journal of Climatology*, n/a–n/a. doi: 10.1002/joc.3736, ISSN: 1097-0088.
- Mallia E., and G. Lewis (2013). Life cycle greenhouse gas emissions of electricity generation in the province of Ontario, Canada. *The International Journal of Life Cycle Assessment* **18**, 377–391. doi: 10.1007/s11367-012-0501-0, ISSN: 0948-3349.
- Malmshiemer R.W., P. Heffernan, S. Brink, D. Crandall, F. Deneke, C. Galik, E. Gee, J.A. Helms, N. McClure, and M. Mortimer (2008). Forest management solutions for mitigating climate change in the United States. *Journal of Forestry* **106**, 173–239. Available at: <http://152.99.88.238/KFRI-CAB/IMG/006/002/157431.pdf#page=173>.
- Von Maltitz G.P., and K.A. Setzkorn (2013). A typology of Southern African biofuel feedstock production projects. *Biomass and Bioenergy* **59**, 33–59. doi: 10.1016/j.biombioe.2012.11.024, ISSN: 0961-9534.
- Mao H.L., J.K. Wang, Y.Y. Zhou, and J.X. Liu (2010). Effects of addition of tea saponins and soybean oil on methane production, fermentation and microbial population in the rumen of growing lambs. *Livestock Science* **129**, 56–62.
- Markus L. (2011). From CDM to REDD+—What do we know for setting up effective and legitimate carbon governance? *Ecological Economics* **70**, 1900–1907. doi: 10.1016/j.ecolecon.2011.02.003, ISSN: 0921-8009.
- Marland G., R.A. Pielke Sr., M. Apps, R. Avissar, R.A. Betts, K.J. Davis, P.C. Frumhoff, S.T. Jackson, L.A. Joyce, P. Kauppi, J. Katzenberger, K.G. MacDicken, R.P. Neilson, J.O. Niles, D. Dutta S. Niyogi, R.J. Norby, N. Pena, N. Sampson, and Y. Xue (2003). The climatic impacts of land surface change and carbon management, and the implications for climate-change mitigation policy. *Climate Policy* **3**, 149–157. doi: 10.1016/S1469-3062(03)00028-7, ISSN: 1469-3062.
- Marland G., and B. Schlamadinger (1995). Biomass fuels and forest-management strategies: how do we calculate the greenhouse-gas emissions benefits? *Energy* **20**, 1131–1140.
- Marland G., and B. Schlamadinger (1997). Forests for carbon sequestration or fossil fuel substitution? A sensitivity analysis. *Biomass and Bioenergy* **13**, 389–397. doi: 10.1016/S0961-9534(97)00027-5, ISSN: 0961-9534.
- Marlow H.J., W.K. Hayes, S. Soret, R.L. Carter, E.R. Schwab, and J. Sabaté (2009). Diet and the environment: does what you eat matter? *The American Journal of Clinical Nutrition* **89**, 1699S–1703S. doi: 10.3945/ajcn.2009.26736Z, ISSN: 0002-9165, 1938–3207.
- Martin C., D.P. Morgavi, and M. Doreau (2010). Methane mitigation in ruminants: from microbe to the farm scale. *Animal* **4**, 351–365. doi: 10.1017/S1751731109990620.
- Martin C., J. Rouel, J.P. Jouany, M. Doreau, and Y. Chilliard (2008). Methane output and diet digestibility in response to feeding dairy cows crude linseed, extruded linseed, or linseed oil. *Journal of Animal Science* **86**, 2642–2650. doi: 10.2527/jas.2007-0774, ISSN: 0021-8812, 1525–3163.

- Martinelli L.A., and S. Filoso (2008).** Expansion of sugarcane ethanol production in Brazil: environmental and social challenges. *Ecological Applications: A Publication of the Ecological Society of America* **18**, 885–898. ISSN: 1051-0761.
- Martinez S.H., J. van Eijck, M.P. da Cunha, A. Walter, J.J. Guilhoto, and A. Faaij (2013).** Analysis of socio-economic impacts of sustainable sugarcane-ethanol production by means of inter-regional input-output analysis: A case study on Northeast Brazil. *Renewable & Sustainable Energy Reviews* **28**, 290–316.
- Martinez-Alier J. (2002).** *The Environmentalism of the Poor: A Study of Ecological Conflicts and Valuation*. Edward Elgar Publishing, Cheltenham, UK and Northampton, USA, 325 pp. ISBN: 9781840649093.
- Masera O.R., J.F. Garza-Caligaris, M. Kanninen, T. Karjalainen, J. Liski, G.J. Nabuurs, A. Pussinen, B.H.J. de Jong, and G.M.J. Mohren (2003).** Modeling carbon sequestration in afforestation, agroforestry and forest management projects: the CO₂FIX V.2 approach. *Ecological Modelling* **164**, 177–199. doi: 10.1016/S0304-3800(02)00419-2, ISSN: 03043800.
- Matson P.A., and P.M. Vitousek (2006).** Agricultural Intensification: Will Land Spared from Farming be Land Spared for Nature? *Conservation Biology* **20**, 709–710. doi: 10.1111/j.1523-1739.2006.00442.x, ISSN: 1523-1739.
- Mattoo A., and A. Subramanian (2012).** Equity in Climate Change: An Analytical Review. *World Development* **40**, 1083–1097. doi: 10.1016/j.worlddev.2011.11.007, ISSN: 0305-750X.
- Matyssek R., G. Wieser, R. Ceulemans, H. Rennenberg, H. Pretzsch, K. Haberer, M. Löw, A.J. Nunn, H. Werner, P. Wipfler, W. Osswald, P. Nikolova, D.E. Hanke, H. Kraigher, M. Tausz, G. Bahnweg, M. Kitao, J. Dieler, H. Sandermann, K. Herbinger, T. Grebenc, M. Blumenröther, G. Deckmyn, T.E.E. Grams, C. Heerdt, M. Leuchner, P. Fabian, and K.-H. Häberle (2010).** Enhanced ozone strongly reduces carbon sink strength of adult beech (*Fagus sylvatica*)—resume from the free-air fumigation study at Kranzberg Forest. *Environmental Pollution (Barking, Essex: 1987)* **158**, 2527–2532. doi: 10.1016/j.envpol.2010.05.009, ISSN: 1873-6424.
- May, P., Boyd, E., Chang, M., V. Neto, and F.C. (2005).** Incorporating sustainable development into carbon forest projects in Brazil and Bolivia. *Estudos Sociedade E Agricultura* **1**, 23 p.
- May P.H., F. Veiga, and M. Chang (2004).** *Local Sustainable Development Effects of Forest Carbon Projects in Brazil and Bolivia: A View from the Field*. Environmental Economics Programme, London, 132 pp. ISBN: Earthprint Limited.
- Mayrand K., and M. Paquin (2004).** *Payments for Environmental Services: A Survey and Assessment of Current Schemes*. UNISFERA International Centre for the Commission of Environmental Cooperation of North America, Montreal, Montreal, 60 pp.
- Mbow C. (2010).** Africa's risky gamble. *Global Change, IGBP Secretariat, number 75 of June 2010* **75**, pp.20–23.
- Mbow, C., Skole, D., Dieng, M., Justice, C., Kwesha, D., Mane, L., E. Gamri, M., V. Vordzogbe, V., Virji, and H. (2012).** *Challenges and Prospects for REDD+ in Africa: Desk Review Of REDD+ Implementation in Africa*. GLP-IPO, Copenhagen, 70 pp.
- McCarl B.A., and U.A. Schneider (2001).** Greenhouse Gas Mitigation in U.S. Agriculture and Forestry. *Science* **294**, 2481–2482. doi: 10.1126/science.1064193, ISSN: 0036-8075, 1095–9203.
- McKechnie J., S. Colombo, J. Chen, W. Mabee, and H.L. MacLean (2011).** Forest Bioenergy or Forest Carbon? Assessing Trade-Offs in Greenhouse Gas Mitigation with Wood-Based Fuels. *Environmental Science & Technology* **45**, 789–795. doi: 10.1021/es1024004, ISSN: 0013-936X.
- McLaren D. (2012).** A comparative global assessment of potential negative emissions technologies. *Special Issue: Negative Emissions Technology* **90**, 489–500. doi: 10.1016/j.jpsep.2012.10.005, ISSN: 0957-5820.
- McMichael A.J., J.W. Powles, C.D. Butler, and R. Uauy (2007).** Food, livestock production, energy, climate change, and health. *The Lancet* **370**, 1253–1263. doi: 10.1016/S0140-6736(07)61256-2, ISSN: 01406736.
- McSherry M.E., and M.E. Ritchie (2013).** Effects of grazing on grassland soil carbon: a global review. *Global Change Biology* **19**, 1347–1357. doi: 10.1111/gcb.12144, ISSN: 13541013.
- MEA (2005).** *Millennium Ecosystem Assessment*. United National Environment Program, New York, Nairobi, 155 pp.
- Meerman J.C., M.M.J. Knoope, A. Ramirez, W.C. Turkenburg, and A.P.C. Faaij (2013).** Technical and economic prospects of coal- and biomass-fired integrated gasification facilities equipped with CCS over time. *International Journal of Greenhouse Gas Control* **16**, 311–323. doi: 10.1016/j.ijggc.2013.01.051, ISSN: 1750-5836.
- Meerman J.C., A. Ramirez, W.C. Turkenburg, and A.P.C. Faaij (2011).** Performance of simulated flexible integrated gasification polygeneration facilities. Part A: A technical-energetic assessment. *Renewable and Sustainable Energy Reviews* **15**, 2563–2587.
- Meerman J.C., A. Ramirez, W.C. Turkenburg, and A.P.C. Faaij (2012).** Performance of simulated flexible integrated gasification polygeneration facilities, Part B: Economic evaluation. *Renewable and Sustainable Energy Reviews* **16**, 6083–6102.
- De Meester S., J. Demeyer, F. Velghe, A. Peene, H. Van Langenhove, and J. Dewulf (2012).** The environmental sustainability of anaerobic digestion as a biomass valorization technology. *Bioresource Technology* **121**, 396–403. doi: 10.1016/j.biortech.2012.06.109, ISSN: 0960-8524.
- Meier D., B. van de Beld, A.V. Bridgwater, D.C. Elliott, A. Oasmaa, and F. Preto (2013).** State-of-the-art of fast pyrolysis in IEA bioenergy member countries. *Renewable and Sustainable Energy Reviews* **20**, 619–641. doi: 10.1016/j.rser.2012.11.061, ISSN: 1364-0321.
- Melillo J.M., J.M. Reilly, D.W. Kicklighter, A.C. Gurgel, T.W. Cronin, S. Paltsev, B.S. Felzer, X. Wang, A.P. Sokolov, and C.A. Schlosser (2009).** Indirect Emissions from Biofuels: How Important? *Science* **326**, 1397–1399. doi: 10.1126/science.1180251, ISSN: 0036-8075, 1095–9203.
- Messerli P., A. Heinimann, M. Giger, T. Breu, and O. Schönweger (2013).** From “land grabbing” to sustainable investments in land: potential contributions by land change science. *Current Opinion in Environmental Sustainability* **5**, 528–534. doi: 10.1016/j.cosust.2013.03.004, ISSN: 1877-3435.
- Metay A., R. Oliver, E. Scopel, J.-M. Douzet, J. Aloisio Alves Moreira, F. Maraux, B.J. Feigl, and C. Feller (2007).** N₂O and CH₄ emissions from soils under conventional and no-till management practices in Goiânia (Cerrados, Brazil). *Geoderma* **141**, 78–88. doi: 10.1016/j.geoderma.2007.05.010, ISSN: 00167061.
- Milazzo M.F., F. Spina, A. Vinci, C. Espro, and J.C.J. Bart (2013).** Brassica bio-diesels: Past, present and future. *Renewable and Sustainable Energy Reviews* **18**, 350–389. doi: 10.1016/j.rser.2012.09.033, ISSN: 1364-0321.
- Millar N., G.P. Robertson, P.R. Grace, R.J. Gehl, and J.P. Hoben (2010).** Nitrogen fertilizer management for nitrous oxide (N₂O) mitigation in intensive corn (Maize) production: an emissions reduction protocol for US Midwest agriculture. *Mitigation and Adaptation Strategies for Global Change* **15**, 185–204. doi: 10.1007/s11027-010-9212-7.

- Millar C.I., N.L. Stephenson, and S.L. Stephens (2007). Climate change and forests of the future: managing in the face of uncertainty. *Ecological Applications* 17, 2145–2151.
- Mingorría S., G. Gamboa, and A. Alonso-Fradejas (2010). *Metabolismo Socio-Ecológico de Comunidades Campesinas Q'eqchi' Y La Expansión de La Agro-Industria de Caña de Azúcar Y Palma Africana : Valle Del Río Polochic, Guatemala*. Instituto de Ciencia y Tecnología Ambientales and Instituto de Estudios Agrarios y Rurales, Barcelona and Mexico, 166 pp. ISBN: 9789929561175.
- Van Minnen J.G., K.K. Goldewijk, E. Stehfest, B. Eickhout, G. van Drecht, and R. Leemans (2009). The importance of three centuries of land-use change for the global and regional terrestrial carbon cycle. *Climatic Change* 97, 123–144. doi: 10.1007/s10584-009-9596-0, ISSN: 0165-0009.
- Mirle C. (2012). The industrialization of animal agriculture: implications for small farmers, rural communities, the environment, and animals in the developing world. The 10th European International Farming Systems Association Symposium in Aarhus, Denmark, July 1–4. Workshop 1.3: Understanding agricultural structural changes and their impacts, to support inclusive policy dialogue and formulation. Available at: http://www.ifs2012.dk/downloads/WS1_3/ChetanaMirle.pdf.
- Misselhorn A., P. Aggarwal, P. Ericksen, P. Gregory, L. Horn-Phathanothai, J. Ingram, and K. Wiebe (2012). A vision for attaining food security. *Current Opinion in Environmental Sustainability* 4, 7–17. doi: 10.1016/j.cosust.2012.01.008, ISSN: 1877-3435.
- Mitchell C.D., R.J. Harper, and R.J. Keenan (2012). Current status and prospects of carbon forestry in Australia. *Australian Forestry* 75, 200–212. ISSN: 0004-9158.
- Moberg J. (2011). The Chinese Grain for Green Program—assessment of the land reform's carbon mitigation potential. Department of Energy and Environment, Chalmers University of Technology, Göteborg, Sweden, 54 pp.
- Mohlin K. (2013). Essays on Environmental Taxation and Climate Policy. Göteborgs Universitet. Handelshögskolan, Department of Economics, Sweden, 155 pp. Available at: <https://gupea.ub.gu.se/handle/2077/33425>.
- Moilanen A., and A. Arponen (2011). Administrative regions in conservation: Balancing local priorities with regional to global preferences in spatial planning. *Ecoregional-Scale Monitoring within Conservation Areas, in a Rapidly Changing Climate* 144, 1719–1725. doi: 10.1016/j.biocon.2011.03.007, ISSN: 0006-3207.
- Monti A., L. Barbanti, A. Zatta, and W. Zegada-Lizarazu (2012). The contribution of switchgrass in reducing GHG emissions. *Global Change Biology Bioenergy* 4. doi: 10.1111/j.1757-1707.2011.01142.x.
- De Moraes M.A.F.D., C.C. da Costa, J.J.M. Guilhoto, L.G.A. de Souza, and F.C.R. de Oliveira (2010). Social Externalities of Fuels. In: *Ethanol and Bioelectricity: Sugarcane in the Future of the Energy Matrix*. UNICA Brazilian Sugarcane Industry Association, Sao Paulo, Brazil, pp. 44–75.
- Morgavi D.P., E. Forano, C. Martin, and C.J. Newbold (2010). Microbial ecosystem and methanogenesis in ruminants. *Animal* 4, 1024–1036. doi: 10.1017/S1751731110000546.
- Morgavi D.P., J.P. Jouany, and C. Martin (2008). Changes in methane emission and rumen fermentation parameters induced by refaunation in sheep. *Australian Journal of Experimental Agriculture* 48, 69–72. doi: <http://dx.doi.org/10.1071/EA07236>.
- Mueller N.D., J.S. Gerber, M. Johnston, D.K. Ray, N. Ramankutty, and J.A. Foley (2012). Closing yield gaps through nutrient and water management. *Nature* 490, 254–257. Available at: <http://dx.doi.org/10.1038/nature11420>.
- Muller A. (2009). Sustainable agriculture and the production of biomass for energy use. *Climatic Change* 94, 319–331. doi: 10.1007/s10584-008-9501-2, ISSN: 0165-0009, 1573–1480.
- Murdiyarso D., M. Brockhaus, W.D. Sunderlin, and L. Verchot (2012). Some lessons learned from the first generation of REDD+ activities. *Current Opinion in Environmental Sustainability* 4, 678–685. doi: 10.1016/j.cosust.2012.10.014, ISSN: 1877-3435.
- Murray B.C., B.A. McCarl, and H.-C. Lee (2004). Estimating Leakage from Forest Carbon Sequestration Programs. *Land Economics* 80, 109–124. doi: 10.3368/le.80.1.109, ISSN: 0023-7639, 1543–8325.
- Murtaugh P.A., and M.G. Schlax (2009). Reproduction and the carbon legacies of individuals. *Global Environmental Change* 19, 14–20.
- Murthy I.K., R. Tiwari, and N.H. Ravindranath (2011). Climate change and forests in India: adaptation opportunities and challenges. *Mitigation and Adaptation Strategies for Global Change* 16, 161–175. doi: 10.1007/s11027-010-9261-y, ISSN: 1381-2386, 1573–1596.
- Muys B., L. Norgrove, T. Alamirew, R. Birech, E. Chirinian, Y. Delelegn, A. Ehrensperger, C.A. Ellison, A. Feto, B. Freyer, J. Gevaert, S. Gmünder, R.E.E. Jongschaap, M. Kaufmann, J. Keane, M. Kenis, B. Kiteme, J. Langat, R. Lyimo, V. Moraa, J. Muchugu, A. Negussie, C. Ouko, M.W. Rouamba, I. Soto, M. Wörgetter, R. Zah, and R. Zetina (2014). Integrating mitigation and adaptation into development: the case of *Jatropha curcas* in sub-Saharan Africa. *Global Change Biology Bioenergy* 6, 169–171. doi: 10.1111/gcbb.12070, ISSN: 1757-1707.
- Mwakaje A.G. (2012). Can Tanzania realise rural development through biofuel plantations? Insights from the study in Rufiji District. *Energy for Sustainable Development* 16, 320–327. doi: 10.1016/j.esd.2012.07.001, ISSN: 0973-0826.
- Myhre G., and D. Shindell (2013). *Anthropogenic and Natural Radiative Forcing*. Intergovernmental Panel on Climate Change, Cambridge UK, 124 pp.
- Nabuurs G.-J., M. Lindner, P.J. Verkerk, K. Gunia, P. Deda, R. Michalak, and G. Grassi (2013). First signs of carbon sink saturation in European forest biomass. *Nature Climate Change* 3, 792–796.
- Nabuurs G.J., O. Masera, K. Andrasko, P. Benitez-Ponce, R. Boer, M. Dutschke, E. Elsidig, J. Ford-Robertson, P. Frumhoff, T. Karjalainen, O. Krankina, W.A. Kurz, M. Matsumoto, W. Oyhantcaba, N.H. Ravindranath, M.J.S. Sanchez, and X. Zhang (2007). Forestry. In: *Climate Change 2007: Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. B. Metz, O.R. Davidson, P.R. Bosch, R. Dave, L.A. Meyer, (eds.), Cambridge University Press, Cambridge, UK and New York, USA, pp. 541–584.
- Nassar A.M., L. Harfuch, L.C. Bachion, and M.R. Moreira (2011). Biofuels and land-use changes: searching for the top model. *Interface Focus*, rfs20100043. doi: 10.1098/rfs.2010.0043, ISSN: 2042-8898, 2042–8901.
- Nelissen V., T. Rütting, D. Huygens, J. Staelens, G. Ruysschaert, and P. Boeckx (2012). Maize biochars accelerate short-term soil nitrogen dynamics in a loamy sand soil. *Soil Biology and Biochemistry* 55, 20–27. doi: 10.1016/j.soilbio.2012.05.019, ISSN: 00380717.

- Nemecek T., K. Weiler, K. Plassmann, J. Schnetzer, G. Gaillard, D. Jefferies, T. García-Suárez, H. King, and L. Milà i Canals (2012). Estimation of the variability in global warming potential of worldwide crop production using a modular extrapolation approach. *Journal of Cleaner Production* **31**, 106–117. doi: 10.1016/j.jclepro.2012.03.005, ISSN: 0959-6526.
- Nepstad D.C., I.M. Tohver, D. Ray, P. Moutinho, and G. Cardinot (2007). Mortality of large trees and lianas following experimental drought in an Amazon forest. *Ecology* **88**, 2259–2269. Available at: <http://www.esajournals.org/doi/abs/10.1890/06-1046.1>.
- Newbold C.J., J.O. Ouda, S. López, N. Nelson, H. Omed, R.J. Wallace, and A.R. Moss (2002). Propionate precursors as possible alternative electron acceptors to methane in ruminal fermentation. In: *Greenhouse Gases and Animal Agriculture*. J. Takahashi, B.A. Young, (eds.), Elsevier, Amsterdam, pp. 151–154.
- Nguyen T.L.T., J.E. Hermansen, and L. Mogensen (2013). Environmental performance of crop residues as an energy source for electricity production: The case of wheat straw in Denmark. *Applied Energy* **104**, 633–641. doi: 10.1016/j.apenergy.2012.11.057, ISSN: 0306-2619.
- Nijssen M., E. Smeets, E. Stehfest, and D.P. Vuuren (2012). An evaluation of the global potential of bioenergy production on degraded lands. *GCB Bioenergy* **4**, 130–147. doi: 10.1111/j.1757-1707.2011.01121.x, ISSN: 1757-1707.
- Njakou Djomo S., O. El Kasmoui, T. De Groot, L.S. Broeckx, M.S. Verlinden, G. Berhongaray, R. Fichot, D. Zona, S.Y. Dillen, J.S. King, I.A. Janssens, and R. Ceulemans (2013). Energy and climate benefits of bioelectricity from low-input short rotation woody crops on agricultural land over a two-year rotation. *Applied Energy* **111**, 862–870. doi: 10.1016/j.apenergy.2013.05.017, ISSN: 0306-2619.
- Nolan J.V., R.S. Hegarty, J. Hegarty, I.R. Godwin, and R. Woodgate (2010). Effects of dietary nitrate on fermentation, methane production and digesta kinetics in sheep. *Animal Production Science* **50**, 801–806.
- Nolte C., A. Agrawal, K.M. Silvius, and B.S. Soares-Filho (2013). Governance regime and location influence avoided deforestation success of protected areas in the Brazilian Amazon. *Proceedings of the National Academy of Sciences* **110**, 4956–4961. doi: 10.1073/pnas.1214786110, ISSN: 0027-8424, 1091–6490.
- Nonhebel S. (2005). Renewable energy and food supply: will there be enough land? *Renewable and Sustainable Energy Reviews* **9**, 191–201.
- O'Brien K., R. Leichenko, U. Kelkar, H. Venema, G. Aandahl, H. Tompkins, A. Javed, S. Bhadwal, S. Barg, L. Nygaard, and others (2004). Mapping vulnerability to multiple stressors: climate change and globalization in India. *Global Environmental Change Part A* **14**, 303–313.
- O'Halloran T.L., B.E. Law, M.L. Goulden, Z. Wang, J.G. Barr, C. Schaaf, M. Brown, J.D. Fuentes, M. Göckede, A. Black, and V. Engel (2012). Radiative forcing of natural forest disturbances. *Global Change Biology* **18**, 555–565.
- O'Hare M., R.J. Plevin, J.I. Martin, A.D. Jones, A. Kendall, and E. Hopson (2009). Proper accounting for time increases crop-based biofuels' greenhouse gas deficit versus petroleum. *Environmental Research Letters* **4**, 24–32. doi: 10.1088/1748-9326/4/2/024001, ISSN: 1748-9326.
- O'Shaughnessy S.M., M.J. Deasy, C.E. Kinsella, J.V. Doyle, and A.J. Robinson (2013). Small scale electricity generation from a portable biomass cookstove: Prototype design and preliminary results. *Applied Energy* **102**, 374–385. doi: 10.1016/j.apenergy.2012.07.032, ISSN: 0306-2619.
- Oberling D.F., M. Obermaier, A. Szklo, and E.L. La Rovere (2012). Investments of oil majors in liquid biofuels: The role of diversification, integration and technological lock-ins. *Biomass and Bioenergy* **46**, 270–281. doi: 10.1016/j.biombioe.2012.08.017, ISSN: 0961-9534.
- Odongo N.E., R. Bagg, G. Vessie, P. Dick, M.M. Or-Rashid, S.E. Hook, J.T. Gray, E. Kebreab, J. France, and B.W. McBride (2007). Long-term effects of feeding monensin on methane production in lactating dairy cows. *Journal of Dairy Science* **90**, 1781–1788. ISSN: 0022-0302.
- Oke D.O., and K.A. Odebiyi (2007). Traditional cocoa-based agroforestry and forest species conservation in Ondo State, Nigeria. *Agriculture, Ecosystems & Environment* **122**, 305–311. Available at: <http://www.sciencedirect.com/science/article/pii/S0167880907000540>.
- Olson D.G., J.E. McBride, A. Joe Shaw, and L.R. Lynd (2012). Recent progress in consolidated bioprocessing. *Current Opinion in Biotechnology* **23**, 396–405. doi: 10.1016/j.copbio.2011.11.026, ISSN: 0958-1669.
- Oosterveer P., and A.P. Mol (2010). Biofuels, trade and sustainability: a review of perspectives for developing countries. *Biofuels, Bioproducts and Biorefining* **4**, 66–76. Available at: <http://onlinelibrary.wiley.com/doi/10.1002/bbb.194/full>.
- Osmond D.L., N.M. Nadkarni, C.T. Driscoll, E. Andrews, A.J. Gold, S.R.B. Allred, A.R. Berkowitz, M.W. Klemens, T.L. Loecke, M.A. McGarry, K. Schwarz, M.L. Washington, and P.M. Groffman (2010). The role of interface organizations in science communication and understanding. *Frontiers in Ecology and the Environment* **8**, 306–313. doi: 10.1890/090145, ISSN: 1540-9295.
- Pa A., J. Craven, X. Bi, S. Melin, and S. Sokhansanj (2012). Environmental footprints of British Columbia wood pellets from a simplified life cycle analysis. *The International Journal of Life Cycle Assessment* **17**, 220–231. doi: 10.1007/s11367-011-0358-7, ISSN: 0948-3349.
- Pacca S., and J.R. Moreira (2011). A Biorefinery for Mobility? *Environmental Science & Technology* **45**, 9498–9505. doi: 10.1021/es2004667, ISSN: 0013-936X.
- Palm, C., Tomich, T., V. Noordwijk, M., Vosti, S., Gockowski, J., Alegre, J., Verchot, and L. (2004). Mitigation GHG emissions in the humid tropics: case studies from the alternative to slash-and-burn program (ASB). *Environment, Development and Sustainability* **6**, 145–162. Available at: <http://link.springer.com/article/10.1023%2FB%3AENVI.0000003634.50442.ca>.
- Palmer C. (2011). Property rights and liability for deforestation under REDD+: Implications for “permanence” in policy design. *Ecological Economics* **70**, 571–576. doi: 10.1016/j.ecolecon.2010.10.011, ISSN: 0921-8009.
- Pan Y., R.A. Birdsey, J. Fang, R. Houghton, P.E. Kauppi, W.A. Kurz, O.L. Phillips, A. Shvidenko, S.L. Lewis, J.G. Canadell, P. Ciais, R.B. Jackson, S.W. Pacala, A.D. McGuire, S. Piao, A. Rautiainen, S. Sitch, and D. Hayes (2011). A Large and Persistent Carbon Sink in the World's Forests. *Science* **333**, 988–993. doi: 10.1126/science.1201609, ISSN: 0036-8075, 1095–9203.
- Pandit M.K., and R.E. Grumbine (2012). Potential Effects of Ongoing and Proposed Hydropower Development on Terrestrial Biological Diversity in the Indian Himalaya. *Conservation Biology* **26**, 1061–1071. doi: 10.1111/j.1523-1739.2012.01918.x, ISSN: 1523-1739.
- Parfitt J., M. Barthel, and S. Macnaughton (2010). Food waste within food supply chains: quantification and potential for change to 2050. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 3065–3081. doi: 10.1098/rstb.2010.0126.

- Parish E.S., M.R. Hilliard, L.M. Baskaran, V.H. Dale, N.A. Griffiths, P.J. Mulholland, A. Sorokine, N.A. Thomas, M.E. Downing, and R.S. Middleton (2012). Multimetric spatial optimization of switchgrass plantings across a watershed. *Biofuels, Bioproducts and Biorefining* 6, 58–72. Available at: <http://onlinelibrary.wiley.com/doi/10.1002/bbb.342/abstract>.
- Pathak H., N. Jain, A. Bhatia, J. Patel, and P.K. Aggarwal (2010). Carbon footprints of Indian food items. *Agriculture, Ecosystems & Environment* 139, 66–73. doi: 10.1016/j.agee.2010.07.002, ISSN: 0167-8809.
- Patrizi N., D. Caro, F.M. Pulselli, A.B. Bjerre, and S. Bastianoni (2013). Environmental feasibility of partial substitution of gasoline with ethanol in the Province of Siena (Italy). *Journal of Cleaner Production* 47, 388–395. doi: 10.1016/j.jclepro.2012.11.023, ISSN: 0959-6526.
- Pawelzik P., M. Carus, J. Hotchkiss, R. Narayan, S. Selke, M. Wellisch, M. Weiss, B. Wicke, and M.K. Patel (2013). Critical aspects in the life cycle assessment (LCA) of bio-based materials—Reviewing methodologies and deriving recommendations. *Resources, Conservation and Recycling* 73, 211–228. doi: 10.1016/j.resconrec.2013.02.006, ISSN: 0921-3449.
- Peralta-Yahya P.P., F. Zhang, S.B. del Cardayre, and J.D. Keasling (2012). Microbial engineering for the production of advanced biofuels. *Nature* 488, 320–328. doi: 10.1038/nature11478, ISSN: 0028-0836.
- Petersen J.-E. (2008). Energy production with agricultural biomass: environmental implications and analytical challenges. *European Review of Agricultural Economics* 35, 385–408. doi: 10.1093/erae/jbn016, ISSN: 0165-1587, 1464–3618.
- Petersen S.O., and S.G. Sommer (2011). Ammonia and nitrous oxide interactions: Roles of manure organic matter management. *Animal Feed Science and Technology* 166–67, 503–513. doi: 10.1016/j.anifeedsci.2011.04.077, ISSN: 0377-8401.
- Peters-Stanley M., K. Hamilton, T. Marcello, and M. Sjardin (2011). *Back to the Future: State of the Voluntary Carbon Markets 2011*. Ecosystem Marketplace & Bloomberg New Energy Finance, Washington, D.C. and New York, NY, USA, 93 pp.
- Pettenella D., and L. Brotto (2011). Governance features for successful REDD+ projects organization. *Forest Policy and Economics* 18, 46–52. doi: 10.1016/j.forpol.2011.09.006, ISSN: 1389-9341.
- Phalan B., M. Onial, A. Balmford, and R.E. Green (2011). Reconciling Food Production and Biodiversity Conservation: Land Sharing and Land Sparing Compared. *Science* 333, 1289–1291. doi: 10.1126/science.1208742, ISSN: 0036-8075, 1095–9203.
- Phelps J., M.C. Guerrero, D.A. Dalabajan, B. Young, and E.L. Webb (2010a). What makes a “REDD” country? *Global Environmental Change* 20, 322–332.
- Phelps J., E.L. Webb, and A. Agrawal (2010b). Does REDD+ Threaten to Recentralize Forest Governance? *Science* 328, 312–313. doi: 10.1126/science.1187774.
- Phillips O.L., L.E.O.C. Aragão, S.L. Lewis, J.B. Fisher, J. Lloyd, G. López-González, Y. Malhi, A. Monteagudo, J. Peacock, C.A. Quesada, and others (2009). Drought sensitivity of the Amazon rainforest. *Science* 323, 1344.
- Phillips O.L., Y. Malhi, N. Higuchi, W.F. Laurance, P.V. Nunez, R. Vasquez, S.G. Laurance, L.V. Ferreira, M. Stern, S. Brown, and J. Grace (1998). Changes in the carbon balance of tropical forests: Evidence from long-term plots. *Science* 282, 439–442.
- Pielke Sr. R.A., A. Pitman, D. Niyogi, R. Mahmood, C. McAlpine, F. Hossain, K.K. Goldewijk, U. Nair, R. Betts, S. Fall, M. Reichstein, P. Kabat, and N. de Noblet (2011). Land use/land cover changes and climate: modeling analysis and observational evidence. *Wiley Interdisciplinary Reviews: Climate Change* 2, 828–850. doi: 10.1002/wcc.144, ISSN: 1757-7799.
- Pingoud K., T. Ekholm, and I. Savolainen (2012). Global warming potential factors and warming payback time as climate indicators of forest biomass use. *Mitigation and Adaptation Strategies for Global Change* 17, 369–386. doi: 10.1007/s11027-011-9331-9, ISSN: 1381-2386, 1573–1596.
- Place S.E., and F.M. Mitloehner (2010). Invited review: Contemporary environmental issues: A review of the dairy industry’s role in climate change and air quality and the potential of mitigation through improved production efficiency. *Journal of Dairy Science* 93, 3407–3416. doi: 10.3168/jds.2009-2719, ISSN: 0022-0302.
- Plattner G.-K., T. Stocker, P. Midgley, and M. Tignor (2009). IPCC expert meeting on the science of alternative metrics. *IPCC Working Group I Technical Support Unit, Bern*.
- Plevin R.J., M.A. Delucchi, and F. Creutzig (2013). Using Attributional Life Cycle Assessment to Estimate Climate-Change Mitigation Benefits Misleads Policy Makers. *Journal of Industrial Ecology* 18, 73–83. doi: 10.1111/jiec.12074, ISSN: 1530-9290.
- Plevin R.J., M. O’Hare, A.D. Jones, M.S. Torn, and H.K. Gibbs (2010). Greenhouse Gas Emissions from Biofuels’ Indirect Land Use Change Are Uncertain but May Be Much Greater than Previously Estimated. *Environmental Science & Technology* 44, 8015–8021. doi: 10.1021/es101946t, ISSN: 0013-936X.
- Pongratz J., C.H. Reick, R.A. Houghton, and J. House (2013). Terminology as a key uncertainty in net land use flux estimates. *Earth System Dynamics Discussions* 4, 677–716. doi: 10.5194/esdd-4-677-2013, ISSN: 2190-4995.
- Popp A., J.P. Dietrich, H. Lotze-Campen, D. Klein, N. Bauer, M. Krause, T. Beringer, D. Gerten, and O. Edenhofer (2011). The economic potential of bioenergy for climate change mitigation with special attention given to implications for the land system. *Environmental Research Letters* 6, 34–44. doi: 10.1088/1748-9326/6/3/034017, ISSN: 1748-9326.
- Popp A., M. Krause, J.P. Dietrich, H. Lotze-Campen, M. Leimbach, T. Beringer, and N. Bauer (2012). Additional CO₂ emissions from land use change—Forest conservation as a precondition for sustainable production of second generation bioenergy. *Ecological Economics* 74, 64–70. doi: 10.1016/j.ecolecon.2011.11.004, ISSN: 0921-8009.
- Popp A., H. Lotze-Campen, and B. Bodirsky (2010). Food consumption, diet shifts and associated non-CO₂ greenhouse gases from agricultural production. *Global Environmental Change* 20, 451–462.
- Popp A., S. Rose, K. Calvin, D. Vuuren, J. Dietrich, M. Wise, E. Stehfest, F. Humpenöder, P. Kyle, J. Vliet, N. Bauer, H. Lotze-Campen, D. Klein, and E. Kriegler (2013). Land-use transition for bioenergy and climate stabilization: model comparison of drivers, impacts and interactions with other land use based mitigation options. *Climatic Change*, 1–15. doi: 10.1007/s10584-013-0926-x, ISSN: 0165-0009.
- Potter C., S. Klooster, C. Hiatt, V. Genovese, and J.C. Castilla-Rubio (2011). Changes in the carbon cycle of Amazon ecosystems during the 2010 drought. *Environmental Research Letters* 6, 1–5.

- Poulter B., L. Aragao, U. Heyder, M. Gumpenberger, J. Heinke, F. Langerwisch, A. Rammig, K. Thonicke, and W. Cramer (2010). Net biome production of the Amazon Basin in the 21st century. *Global Change Biology* 16, 2062–2075. doi: 10.1111/j.1365-2486.2009.02064.x, ISSN: 1354-1013.
- Powlson D.S., A.P. Whitmore, and K.W.T. Goulding (2011). Soil carbon sequestration to mitigate climate change: a critical re-examination to identify the true and the false. *European Journal of Soil Science* 62, 42–55. doi: 10.1111/j.1365-2389.2010.01342.x, ISSN: 1365-2389.
- Pretty J. (2008). Agricultural sustainability: concepts, principles and evidence. *Philosophical Transactions of the Royal Society B: Biological Sciences* 363, 447–465. doi: 10.1098/rstb.2007.2163, ISSN: 0962-8436, 1471–2970.
- Pretty J. (2013). The Consumption of a Finite Planet: Well-Being, Convergence, Divergence and the Nascent Green Economy. *Environmental and Resource Economics* 55, 475–499. doi: 10.1007/s10640-013-9680-9, ISSN: 0924-6460, 1573–1502.
- Primmer E., and H. Karppinen (2010). Professional judgment in non-industrial private forestry: Forester attitudes and social norms influencing biodiversity conservation. *Forest Policy and Economics* 12, 136–146. doi: 10.1016/j.forpol.2009.09.007, ISSN: 1389-9341.
- Pucker J., R. Zwart, and G. Jungmeier (2012). Greenhouse gas and energy analysis of substitute natural gas from biomass for space heat. *Biomass and Bioenergy* 38, 95–101. doi: 10.1016/j.biombioe.2011.02.040, ISSN: 0961-9534.
- Puettmann M.E., R. Bergman, S. Hubbard, L. Johnson, B. Lippke, E. Oneil, and F.G. Wagner (2010). Cradle-to-gate life-cycle inventory of the U.S. Wood products production: CORRIM Phase I and Phase II products. *Wood and Fiber Science* 42, 15–28.
- Pullin R.S.V., R. Froese, and D. Pauly (2007). Indicators for the sustainability of aquaculture. In: *Ecological and Genetic Implications of Aquaculture Activities*. T.M. Bert, (ed.), Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 53–72.
- Putz F.E., and K.H. Redford (2009). Dangers of carbon-based conservation. *Global Environmental Change* 19, 400–401.
- Putz F.E., and C. Romero (2012). Helping curb tropical forest degradation by linking REDD+ with other conservation interventions: a view from the forest. *Current Opinion in Environmental Sustainability* 4, 670–677. doi: 10.1016/j.cosust.2012.10.003, ISSN: 1877-3435.
- Le Quéré C., R.J. Andres, T. Boden, T. Conway, R.A. Houghton, J.I. House, G. Marland, G.P. Peters, G.R. van der Werf, A. Ahlström, R.M. Andrew, L. Bopp, J.G. Canadell, P. Ciais, S.C. Doney, C. Enright, P. Friedlingstein, C. Huntingford, A.K. Jain, C. Jourdain, E. Kato, R.F. Keeling, K. Klein Goldewijk, S. Levis, P. Levy, M. Lomas, B. Poulter, M.R. Raupach, J. Schwinger, S. Sitch, B.D. Stocker, N. Viovy, S. Zaehle, and N. Zeng (2013). The global carbon budget 1959–2011. *Earth System Science Data* 5, 165–185. doi: 10.5194/essd-5-165-2013, ISSN: 1866-3516.
- Rametsteiner E., and M. Simula (2003). Forest certification—an instrument to promote sustainable forest management? *Journal of Environmental Management* 67, 87–98. doi: 10.1016/S0301-4797(02)00191-3, ISSN: 03014797.
- Randerson J.T., H. Liu, M.G. Flanner, S.D. Chambers, Y. Jin, P.G. Hess, G. Pfister, M.C. Mack, K.K. Treseder, L.R. Welp, F.S. Chapin, J.W. Harden, M.L. Goulden, E. Lyons, J.C. Neff, E.A.G. Schuur, and C.S. Zender (2006). The Impact of Boreal Forest Fire on Climate Warming. *Science* 314, 1130–1132. doi: 10.1126/science.1132075.
- Rathmann R., A. Szklo, and R. Schaeffer (2010). Land use competition for production of food and liquid biofuels: An analysis of the arguments in the current debate. *Renewable Energy* 35, 14–22.
- Ravindranath N.H. (2007). Mitigation and adaptation synergy in forest sector. *Mitigation and Adaptation Strategies for Global Change* 12, 843–853. doi: 10.1007/s11027-007-9102-9, ISSN: 1381-2386, 1573–1596.
- Read J.M., J.M.V. Fragoso, K.M. Silvius, J. Luzar, H. Overman, A. Cummings, S.T. Giery, and L.F. de Oliveira (2010). Space, Place, and Hunting Patterns among Indigenous Peoples of the Guyanese Rupununi Region. *Journal of Latin American Geography* 9, 213–243. ISSN: 1548-5811.
- Rehl T., J. Lansche, and J. Müller (2012). Life cycle assessment of energy generation from biogas—Attributional vs. consequential approach. *Renewable and Sustainable Energy Reviews* 16, 3766–3775. doi: 10.1016/j.rser.2012.02.072, ISSN: 1364-0321.
- Reichstein M., M. Bahn, P. Ciais, D. Frank, M.D. Mahecha, S.I. Seneviratne, J. Zscheischler, C. Beer, N. Buchmann, D.C. Frank, D. Papale, A. Rammig, P. Smith, K. Thonicke, M. van der Velde, S. Vicca, A. Walz, and M. Wattenbach (2013). Climate extremes and the carbon cycle. *Nature* 500, 287–295. doi: 10.1038/nature12350, ISSN: 0028-0836, 1476–4687.
- Reilly J., J. Melillo, Y. Cai, D. Kicklighter, A. Gurgel, S. Paltsev, T. Cronin, A. Sokolov, and A. Schlosser (2012). Using Land To Mitigate Climate Change: Hitting the Target, Recognizing the Trade-offs. *Environmental Science & Technology* 46, 5672–5679. doi: 10.1021/es2034729, ISSN: 0013-936X, 1520–5851.
- REN21 (2012). *Renewables 2012 Global Status Report*. REN21 Secretariat, Paris, France, 176 pp. Available at: <http://www.ren21.net/Resources/Publications/REN21Publications/Renewables2012GlobalStatusReport.aspx>.
- REN21 (2013). *Renewables 2013 Global Status Report*. REN21 Secretariat, Paris, France, 178 pp. Available at: <http://www.ren21.net/ren21activities/globalstatusreport.aspx>.
- Repo A., R. Känkänen, J.-P. Tuovinen, R. Antikainen, M. Tuomi, P. Vanhala, and J. Liski (2012). Forest bioenergy climate impact can be improved by allocating forest residue removal. *GCB Bioenergy* 4, 202–212.
- Repo A., M. Tuomi, and J. Liski (2011). Indirect carbon dioxide emissions from producing bioenergy from forest harvest residues. *GCB Bioenergy* 3, 107–115. doi: 10.1111/j.1757-1707.2010.01065.x, ISSN: 1757-1707.
- Reyer C., M. Guericke, and P.L. Ibsch (2009). Climate change mitigation via afforestation, reforestation and deforestation avoidance: and what about adaptation to environmental change? *New Forests* 38, 15–34. doi: 10.1007/s11056-008-9129-0, ISSN: 0169-4286, 1573–5095.
- RFA (2008). *Indirect Effects of Biofuels: Study by the Renewable Fuels Agency*. Renewable Fuels Agency, London, UK, 92 pp.
- Rhodes J., and D. Keith (2008). Biomass with capture: negative emissions within social and environmental constraints: an editorial comment. *Climatic Change* 87, 321–328. doi: 10.1007/s10584-007-9387-4, ISSN: 0165-0009.
- Rice R.A. (2008). Agricultural intensification within agroforestry: the case of coffee and wood products. *Agriculture, Ecosystems & Environment* 128, 212–218. Available at: <http://www.sciencedirect.com/science/article/pii/S0167880908001874>.
- Richels R., J. Merrick, G. Blanford, and S. Rose (In Review). Trade-offs Between Mitigation Costs and Temperature Change. *Climatic Change*.

- Robertson G.P., T.W. Bruulsema, R.J. Gehl, D. Kanter, D.L. Mauzerall, C.A. Rotz, and C.O. Williams (2013). Nitrogen–climate interactions in US agriculture. *Biogeochemistry* **114**, 41–70. doi: 10.1007/s10533-012-9802-4, ISSN: 0168-2563, 1573–515X.
- Robertson G.P., and P.M. Vitousek (2009). Nitrogen in Agriculture: Balancing the Cost of an Essential Resource. *Annual Review of Environment and Resources* **34**, 97–125. doi: 10.1146/annurev.enviro.032108.105046, ISSN: 1543-5938.
- Robinson N., R.J. Harper, K.R.J. Smettem, and J.F. McGrath (2004). Tree placement strategies for salinity control in dryland farming systems of southern Australia. In: *13th International Soil Conservation Organization Conference*. Brisbane, Australia. 6 pp. Available at: <http://www.tucson.ars.ag.gov/isco/isco13/PAPERS%20R-Z/ROBINSON.pdf>.
- Robinson B.E., M. Holland, and L. Naughton-Treves (2011). Does secure land tenure save forest? A review of the relationship between land tenure and tropical deforestation. CCAFS Working Paper no.7. CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS). Available at: <http://cgspace.cgiar.org/handle/10568/10720>.
- Robledo C., N. Clot, A. Hammill, and B. Riché (2011). The role of forest ecosystems in community-based coping strategies to climate hazards: Three examples from rural areas in Africa. *Forest Policy and Economics* **24**, 20–28. doi: 10.1016/j.forpol.2011.04.006, ISSN: 13899341.
- Rochette P. (2008). No-till only increases N₂O emissions in poorly-aerated soils. *Soil and Tillage Research* **101**, 97–100. doi: 10.1016/j.still.2008.07.011, ISSN: 01671987.
- Roedel A. (2010). Production and energetic utilization of wood from short rotation coppice—a life cycle assessment. *The International Journal of Life Cycle Assessment* **15**, 567–578. doi: 10.1007/s11367-010-0195-0, ISSN: 0948-3349.
- Rogner H.H., R.F. Aguilera, C.L. Archer, R. Bertani, S.C. Bhattacharya, I. Bryden, R.R. Charpentier, M.B. Dusseault, L. Gagnon, Y. Goswami, H. Haberl, M.M. Hoogwijk, A. Johnson, P. Odell, H. Wagner, and V. Yakushev (2012). Chapter 7—Energy resources and potentials. In: *Global Energy Assessment: Toward a Sustainable Future*. L. Gomez-Echeverri, T.B. Johansson, N. Nakicenovic, A. Patwardhan, (eds.), IIASA and Cambridge University Press, Laxenburg, Austria, Cambridge, UK, pp. 425–512.
- Rohde R.F., and M.T. Hoffman (2012). The historical ecology of Namibian rangelands: Vegetation change since 1876 in response to local and global drivers. *Science of The Total Environment* **416**, 276–288. doi: 10.1016/j.scitotenv.2011.10.067, ISSN: 00489697.
- Rose S.K., H. Ahammad, B. Eickhout, B. Fisher, A. Kurosawa, S. Rao, K. Riahi, and D.P. van Vuuren (2012). Land-based mitigation in climate stabilization. *Energy Economics* **34**, 365–380. doi: 10.1016/j.eneco.2011.06.004, ISSN: 0140-9883.
- Rose S., R. Beach, K. Calvin, B. McCarl, J. Petrusa, B. Sohngen, R. Youngman, A. Diamant, F. de la Chesnaye, J. Edmonds, R. Rosenzweig, and M. Wise (2013). *Estimating Global Greenhouse Gas Emissions Offset Supplies: Accounting for Investment Risks and Other Market Realities*. EPRI, Palo Alto, CA, 23 pp.
- Rose S.K., and B. Sohngen (2011). Global forest carbon sequestration and climate policy design. *Environment and Development Economics* **16**, 429–454. doi: 10.1017/S1355770X11000027, ISSN: 1355-770X, 1469–4395.
- Rosemary L. (2011). REDD+, transparency, participation and resource rights: the role of law. *Environmental Science & Policy* **14**, 118–126. doi: 10.1016/j.envsci.2010.11.008, ISSN: 1462-9011.
- Rosendal G.K., and S. Andresen (2011). Institutional design for improved forest governance through REDD: Lessons from the global environment facility. *Ecological Economics* **70**, 1908–1915. doi: 10.1016/j.ecolecon.2011.04.001, ISSN: 0921-8009.
- Rosenzweig C., and F.N. Tubiello (2007). Adaptation and mitigation strategies in agriculture: an analysis of potential synergies. *Mitigation, Adaptation Strategies to Global Change* **12**, 855–873. doi: 10.1007/s11027-007-9103-8.
- Rosillo-Calle F., S. Teelucksingh, D. Thrän, and M. Seiffert (2012). *The Potential Role of Biofuels in Commercial Air Transport—Biojetfuel*. IEA, Paris, France, 56 pp.
- Rotenberg E., and D. Yakir (2010). Contribution of Semi-Arid Forests to the Climate System. *Science* **327**, 451–454. doi: 10.1126/science.1179998.
- Roy P., T. Orikasa, M. Thammawong, N. Nakamura, Q. Xu, and T. Shiina (2012). Life cycle of meats: An opportunity to abate the greenhouse gas emission from meat industry in Japan. *Journal of Environmental Management* **93**, 218–224. doi: 10.1016/j.jenvman.2011.09.017, ISSN: 0301-4797.
- Royal Society (2009). *Reaping the Benefits: Science and the Sustainable Intensification of Global Agriculture*. The Royal Society, London, 86 pp.
- Rubin E.M. (2008). Genomics of cellulosic biofuels. *Nature* **454**, 841–845. doi: 10.1038/nature07190, ISSN: 0028-0836.
- Saatchi S.S., N.L. Harris, S. Brown, M. Lefsky, E.T.A. Mitchard, W. Salas, B.R. Zutta, W. Buermann, S.L. Lewis, S. Hagen, S. Petrova, L. White, M. Silman, and A. Morel (2011). Benchmark map of forest carbon stocks in tropical regions across three continents. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 9899–9904. doi: 10.1073/pnas.1019576108, ISSN: 0027-8424.
- Saddler J., and L. Kumar (2013). Special Issue from the NSERC Bioconversion network workshop: pretreatment and fractionation of biomass for biorefinery/biofuels. *Biotechnology for Biofuels* **6**, 17. doi: 10.1186/1754-6834-6-17, ISSN: 1754-6834.
- Saggar S., R. M. Andrew, K. R. Tate, C. B. Hedley, N. J. Rodda, and J. A. Townsend (2004). Modelling nitrous oxide emissions from dairy-grazed pastures. *Nutrient Cycling in Agroecosystems* **68**, 243–255. Available at: <http://link.springer.com/article/10.1023/B:FRES.00000019463.92440.a3>.
- Sala O.E., F.S. Chapin, J.J. Armesto, E. Berlow, J. Bloomfield, R. Dirzo, E. Huber-Sanwald, L.F. Huenneke, R.B. Jackson, A. Kinzig, R. Leemans, D.M. Lodge, H.A. Mooney, M. Oesterheld, N.L. Poff, M.T. Sykes, B.H. Walker, M. Walker, and D.H. Wall (2000). Global Biodiversity Scenarios for the Year 2100. *Science* **287**, 1770–1774. doi: 10.1126/science.287.5459.1770, ISSN: 0036-8075, 1095–9203.
- Sandor R.L., E.C. Bettelheim, and I.R. Swingland (2002). An overview of a free-market approach to climate change and conservation. *Philosophical Transactions of the Royal Society of London Series a-Mathematical Physical and Engineering Sciences* **360**, 1607–1620. ISSN: 1364-503X.
- Santilli M., P. Moutinho, S. Schwartzman, D. Nepstad, L. Curran, and C. Nobre (2005). Tropical Deforestation and the Kyoto Protocol. *Climatic Change* **71**, 267–276. doi: 10.1007/s10584-005-8074-6, ISSN: 0165-0009, 1573–1480.
- Sathaye J., W. Makundi, L. Dale, P. Chan, and K. Andrasko (2005). *Generalized Comprehensive Mitigation Assessment Process (GCOMAP): A Dynamic Partial Equilibrium Model for*. U.S. Environmental Protection Agency, Washington, D.C., 93 pp. Available at: <http://eedt.dbl.gov/sites/all/files/lbnl-58291.pdf>.

- Sathaye J., W. Makundi, L. Dale, P. Chan, and K. Andrasko (2006). GHG Mitigation Potential, Costs and Benefits in Global Forests: A Dynamic Partial Equilibrium Approach. *The Energy Journal Multi-Greenhouse Gas Mitigation and Climate Policy*, 127–162. ISSN: 0195-6574.
- Sathre R., and J. O'Connor (2010). Meta-analysis of greenhouse gas displacement factors of wood product substitution. *Environmental Science & Policy* 13, 104–114. doi: 10.1016/j.envsci.2009.12.005, ISSN: 1462-9011.
- Satolo L., and M. Bacchi (2013). Impacts of the Recent Expansion of the Sugar-cane Sector on Municipal per Capita Income in São Paulo State. *ISRN Economics* 2013, 14.
- Sauvant D., and S. Giger-Reverdin (2007). Empirical modelling meta-analysis of digestive interactions and CH₄ production in ruminants. In: *Energy and protein metabolism and nutrition*. I. Ortigues-Marty, N. Miraux, W. Brand-Williams, (eds.), Wageningen Academic Publishers, Wageningen, The Netherlands, pp. 561–563. ISBN: 978-90-8686-041-8.
- Sayer J., J. Ghazoul, P. Nelson, and A. Klintuni Boedhihartono (2012). Oil palm expansion transforms tropical landscapes and livelihoods. *Global Food Security* 1, 114–119. doi: 10.1016/j.gfs.2012.10.003, ISSN: 2211-9124.
- Scheidel A., and A. H. Sorman (2012). Energy transitions and the global land rush: Ultimate drivers and persistent consequences. *Global Transformations, Social Metabolism and the Dynamics of Socio-Environmental Conflicts* 22, 588–595. doi: 10.1016/j.gloenvcha.2011.12.005, ISSN: 0959-3780.
- Schils R.L.M., J. Eriksen, S.F. Ledgard, T.V. Vellinga, P.J. Kuikman, J. Luo, S.O. Petersen, and G.L. Velthof (2013). Strategies to mitigate nitrous oxide emissions from herbivore production systems. *Animal* 7, 29–40. doi: 10.1017/S175173111100187X.
- Schlamadinger B., and G. Marland (1996). The role of forest and bioenergy strategies in the global carbon cycle. *Biomass and Bioenergy* 10, 275–300. doi: 10.1016/0961-9534(95)00113-1, ISSN: 0961-9534.
- Schmidinger K., and E. Stehfest (2012). Including CO₂ implications of land occupation in LCAs—method and example for livestock products. *The International Journal of Life Cycle Assessment* 17, 962–972. doi: 10.1007/s11367-012-0434-7, ISSN: 0948-3349, 1614–7502.
- Schmidt J., V. Gass, and E. Schmid (2011). Land use changes, greenhouse gas emissions and fossil fuel substitution of biofuels compared to bioelectricity production for electric cars in Austria. *Biomass and Bioenergy* 35, 4060–4074. doi: 10.1016/j.biombioe.2011.07.007, ISSN: 0961-9534.
- Schmitz C., A. Biewald, H. Lotze-Campen, A. Popp, J.P. Dietrich, B. Bodirsky, M. Krause, and I. Weindl (2011). Trading more food: Implications for land use, greenhouse gas emissions, and the food system. *Global Environmental Change* 22, 189–209. doi: 10.1016/j.gloenvcha.2011.09.013, ISSN: 0959-3780.
- Schneider A., M.A. Friedl, and D. Potere (2009). A new map of global urban extent from MODIS satellite data. *Environmental Research Letters* 4, 12. doi: 10.1088/1748-9326/4/4/044003, ISSN: 1748-9326.
- Schneider U.A., P. Havlik, E. Schmid, H. Valin, A. Mosnier, M. Obersteiner, H. Böttcher, R. Skalský, J. Balkovič, T. Sauer, and S. Fritz (2011). Impacts of population growth, economic development, and technical change on global food production and consumption. *Agricultural Systems* 104, 204–215. doi: 10.1016/j.agsy.2010.11.003, ISSN: 0308-521X.
- Schut M., M. Slingerland, and A. Locke (2010). Biofuel developments in Mozambique. Update and analysis of policy, potential and reality. *Energy Policy* 38, 5151–5165. doi: 10.1016/j.enpol.2010.04.048, ISSN: 0301-4215.
- Schuur E.A.G., J. Bockheim, J.G. Canadell, E. Euskirchen, C.B. Field, S.V. Goryachkin, S. Hagemann, P. Kuhry, P.M. Laffleur, H. Lee, G. Mazhitova, F.E. Nelson, A. Rinke, V.E. Romanovsky, N. Shiklomanov, C. Tarnocai, S. Venevsky, J.G. Vogel, and S.A. Zimov (2008). Vulnerability of Permafrost Carbon to Climate Change: Implications for the Global Carbon Cycle. *BioScience* 58, 701. doi: 10.1641/B580807, ISSN: 0006-3568.
- Schwietzke S., W.M. Griffin, and H.S. Matthews (2011). Relevance of Emissions Timing in Biofuel Greenhouse Gases and Climate Impacts. *Environmental Science & Technology* 45, 8197–8203. doi: 10.1021/es2016236, ISSN: 0013-936X.
- Scown C.D., W.W. Nazaroff, U. Mishra, B. Strogon, A.B. Lobscheid, E. Masanet, N.J. Santero, A. Horvath, and T.E. McKone (2012). Lifecycle greenhouse gas implications of US national scenarios for cellulosic ethanol production. *Environmental Research Letters* 7, 014011. doi: 10.1088/1748-9326/7/1/014011, ISSN: 1748-9326.
- Seabra J.E., I.C. Macedo, H.L. Chum, C.E. Faroni, and C.A. Sarto (2011). Life cycle assessment of Brazilian sugarcane products: GHG emissions and energy use. *Biofuels, Bioproducts and Biorefining* 5, 519–532. doi: 10.1002/bbb.289, ISSN: 1932-1031.
- Sequist, J.W., Hickler, T., Eklundh, L., Ardö, J., and Heumann, and B.W. (2008). Disentangling the effects of climate and people on Sahel vegetation dynamics. *Biogeosciences Discussions* 5, 3045–3067.
- Searchinger T.D. (2010). Biofuels and the need for additional carbon. *Environmental Research Letters* 5, 11. doi: 10.1088/1748-9326/5/2/024007, ISSN: 1748-9326.
- Searchinger T.D., S.P. Hamburg, J. Melillo, W. Chameides, P. Havlik, D.M. Kammen, G.E. Likens, R.N. Lubowski, M. Obersteiner, M. Oppenheimer, G. Philip Robertson, W.H. Schlesinger, and G. David Tilman (2009). Fixing a Critical Climate Accounting Error. *Science* 326, 527–528. doi: 10.1126/science.1178797.
- Searchinger T., R. Heimlich, R.A. Houghton, F. Dong, A. Elobeid, J. Fabiosa, S. Tokgoz, D. Hayes, and T.-H. Yu (2008). Use of U.S. Croplands for Biofuels Increases Greenhouse Gases Through Emissions from Land-Use Change. *Science* 319, 1238–1240. doi: 10.1126/science.1151861, ISSN: 0036-8075, 1095–9203.
- Searcy E., and P.C. Flynn (2008). Processing of Straw/Corn Stover: Comparison of Life Cycle Emissions. *International Journal of Green Energy* 5, 423–437. doi: 10.1080/15435070802498010, ISSN: 1543-5075.
- Sedjo R., and X. Tian (2012). Does Wood Bioenergy Increase Carbon Stocks in Forests? *Journal of Forestry* 110, 304–311.
- Seitzinger S.P., U. Svedin, C.L. Crumley, W. Steffen, S.A. Abdullah, C. Alfson, W.J. Broadgate, F. Biermann, N.R. Bondre, J.A. Dearing, L. Deutsch, S. Dhaka, T. Elmqvist, N. Farahbakhshazad, O. Gaffney, H. Haberl, S. Lavorel, C. Mbow, A.J. McMichael, J.M.F. deMorais, P. Olsson, P.F. Pinho, K.C. Seto, P. Sinclair, M.S. Smith, and L. Sugar (2012). Planetary Stewardship in an Urbanizing World: Beyond City Limits. *AMBIO* 41, 787–794. doi: 10.1007/s13280-012-0353-7, ISSN: 0044-7447, 1654–7209.
- Selfa T., L. Kulcsar, C. Bain, R. Goe, and G. Middendorf (2011). Biofuels Bonanza?: Exploring community perceptions of the promises and perils of biofuels production. *Biomass and Bioenergy* 35, 1379–1389. doi: 10.1016/j.biombioe.2010.09.008, ISSN: 0961-9534.
- Semere T., and F.M. Slater (2007). Ground flora, small mammal and bird species diversity in miscanthus (*Miscanthus giganteus*) and reed canary-grass (*Phalaris arundinacea*) fields. *Biomass and Bioenergy* 31, 20–29. ISSN: 09619534.

- Semroc B.L., G. Schroth, C.A. Harvey, Y. Zepeda, and F. Boltz (2012).** Climate Change mitigation in agroforestry systems. linking smallholders to forest carbon markets. In: *Climate Change Mitigation and Agriculture*. E. Wollenberg, M.-L. Tapio-Bstrom, M. Grieg-Gran, A. Nihart, (eds.), Routledge, London-New York, pp. 360–369. ISBN: 1849713936.
- Senel O., and N. Chugunov (2013).** CO₂ Injection in a Saline Formation: Pre-Injection Reservoir Modeling and Uncertainty Analysis for Illinois Basin—Decatur Project. *Energy Procedia* **37**, 4598–4611. doi: 10.1016/j.egypro.2013.06.368, ISSN: 1876-6102.
- Seppälä R., A. Buck, and P. Katila (Eds.) (2009).** *Adaptation of Forests and People to Climate Change. A Global Assessment Report*. International Union of Forest Research Organizations, Helsinki, 224 pp. ISBN: 978-3-901347-80-1.
- Serrano-Ruiz J.C., R.M. West, and J.A. Dumesic (2010).** Catalytic Conversion of Renewable Biomass Resources to Fuels and Chemicals. *Annual Review of Chemical and Biomolecular Engineering* **1**, 79–100. doi: 10.1146/annurev-chembioeng-073009-100935.
- Seto K.C., B. Güneralp, and L.R. Hutyra (2012a).** Global forecasts of urban expansion to 2030 and direct impacts on biodiversity and carbon pools. *Proceedings of the National Academy of Sciences* **109**, 16083–16088. doi: 10.1073/pnas.1211658109, ISSN: 0027-8424, 1091–6490.
- Seto K.C., A. Reenberg, C.G. Boone, M. Fragkias, D. Haase, T. Langanke, P. Marcotullio, D.K. Munroe, B. Olah, and D. Simon (2012b).** Urban Land Teleconnections and Sustainability. *Proceedings of the National Academy of Sciences* **109**, 7687–7692. doi: 10.1073/pnas.1117622109, ISSN: 0027-8424, 1091–6490.
- Shackley S., S. Carter, T. Knowles, E. Middelink, S. Haefele, S. Sohi, A. Cross, and S. Haszeldine (2012).** Sustainable gasification–biochar systems? A case-study of rice-husk gasification in Cambodia, Part I: Context, chemical properties, environmental and health and safety issues. *Energy Policy* **42**, 49–58. doi: 10.1016/j.enpol.2011.11.026, ISSN: 03014215.
- Shen H., C. Poovaiah, A. Ziebell, T. Tschaplinski, S. Pattathil, E. Gjersing, N. Engle, R. Katahira, Y. Pu, R. Sykes, F. Chen, A. Ragauskas, J. Mielenz, M. Hahn, M. Davis, C.N. Stewart, and R. Dixon (2013).** Enhanced characteristics of genetically modified switchgrass (*Panicum virgatum* L.) for high biofuel production. *Biotechnology for Biofuels* **6**, 71. Available at: <http://www.biotechnologyforbiofuels.com/content/6/1/71>.
- Siangjaeo S., S.H. Gheewala, K. Unnanon, and A. Chidthaisong (2011).** Implications of land use change on the life cycle greenhouse gas emissions from palm biodiesel production in Thailand. *Energy for Sustainable Development* **15**, 1–7. doi: 10.1016/j.esd.2011.01.002, ISSN: 0973-0826.
- Sikor T., J. Stahl, T. Enters, J.C. Ribot, N. Singh, W.D. Sunderlin, and L. Wollenberg (2010).** REDD-plus, forest people's rights and nested climate governance. *Global Environmental Change* **20**, 423–425. doi: 10.1016/j.gloenvcha.2010.04.007, ISSN: 0959-3780.
- Silalertruksa T., and S.H. Gheewala (2012).** Environmental sustainability assessment of palm biodiesel production in Thailand. *Energy* **43**, 306–314. doi: 10.1016/j.energy.2012.04.025, ISSN: 0360-5442.
- Silver W., R. Ostertag, and A. Lugo (2000).** The potential for carbon sequestration through reforestation of abandoned tropical agricultural and pasture lands. *Restoration Ecology* **8**, 394–407.
- Sims R., A. Hastings, B. Schlamadinger, G. Taylor, and P. Smith (2006).** Energy crops: current status and future prospects. *Global Change Biology* **12**, 2054–2076. doi: 10.1111/j.1365-2486.2006.01163.x, ISSN: 1365-2486.
- Singh P.P. (2008).** Exploring biodiversity and climate change benefits of community-based forest management. *Global Environmental Change* **18**, 468–478. doi: 10.1016/j.gloenvcha.2008.04.006, ISSN: 0959-3780.
- Singh B.P., A.L. Cowie, and R.J. Smernik (2012).** Biochar Carbon Stability in a Clayey Soil As a Function of Feedstock and Pyrolysis Temperature. *Environmental Science & Technology* **46**, 11770–11778. doi: 10.1021/es302545b, ISSN: 0013-936X, 1520–5851.
- Singh B.P., B.J. Hatton, B. Singh, A.L. Cowie, and A. Kathuria (2010).** Influence of Biochars on Nitrous Oxide Emission and Nitrogen Leaching from Two Contrasting Soils. *Journal of Environment Quality* **39**, 1224. doi: 10.2134/jeq2009.0138, ISSN: 1537-2537.
- Sitch S., C. Huntingford, N. Gedney, P.E. Levy, M. Lomas, S.L. Piao, R. Betts, P. Ciais, P. Cox, P. Friedlingstein, C.D. Jones, I.C. Prentice, and F.I. Woodward (2008).** Evaluation of the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five Dynamic Global Vegetation Models (DGVMs). *Global Change Biology* **14**, 2015–2039. doi: 10.1111/j.1365-2486.2008.01626.x, ISSN: 1365-2486.
- Skutsch M., B. Vickers, Y. Georgiadou, and M. McCall (2011).** Alternative models for carbon payments to communities under REDD+: A comparison using the Polis model of actor inducements. *Environmental Science & Policy* **14**, 140–151. doi: 10.1016/j.envsci.2010.12.005, ISSN: 1462-9011.
- Smeets E.M.W., L.F. Bouwman, E. Stehfest, D.P. Van VUUREN, and A. Posthuma (2009a).** Contribution of N₂O to the greenhouse gas balance of first-generation biofuels. *Global Change Biology* **15**, 1–23. doi: 10.1111/j.1365-2486.2008.01704.x, ISSN: 1365-2486.
- Smeets E.M.W., and A.P.C. Faaij (2007).** Bioenergy potentials from forestry in 2050. *Climatic Change* **81**, 353–390. doi: 10.1007/s10584-006-9163-x, ISSN: 0165-0009, 1573–1480.
- Smeets E.M., and A.P. Faaij (2010).** The impact of sustainability criteria on the costs and potentials of bioenergy production—Applied for case studies in Brazil and Ukraine. *Biomass and Bioenergy* **34**, 319–333.
- Smeets E.M.W., A.P.C. Faaij, I.M. Lewandowski, and W.C. Turkenburg (2007).** A bottom-up assessment and review of global bio-energy potentials to 2050. *Progress in Energy and Combustion Science* **33**, 56–106. doi: 10.1016/j.pecs.2006.08.001, ISSN: 0360-1285.
- Smeets E., M. Junginger, A. Faaij, A. Walter, P. Dolzan, and W. Turkenburg (2008).** The sustainability of Brazilian ethanol—An assessment of the possibilities of certified production. *Biomass and Bioenergy* **32**, 781–813. doi: 10.1016/j.biombioe.2008.01.005, ISSN: 09619534.
- Smeets E.M.W., I.M. Lewandowski, and A.P.C. Faaij (2009b).** The economical and environmental performance of miscanthus and switchgrass production and supply chains in a European setting. *Renewable and Sustainable Energy Reviews* **13**, 1230–1245. doi: 10.1016/j.rser.2008.09.006, ISSN: 1364-0321.
- Smith P. (2005).** An overview of the permanence of soil organic carbon stocks: Influence of direct human-induced, indirect and natural effects. *European Journal of Soil Science* **56**, 673–680.
- Smith P. (2008).** Land use change and soil organic carbon dynamics. *Nutrient Cycling in Agroecosystems* **81**, 169–178. doi: 10.1007/s10705-007-9138-y, ISSN: 1385-1314, 1573–0867.
- Smith P. (2012).** Agricultural greenhouse gas mitigation potential globally, in Europe and in the UK: what have we learnt in the last 20 years? *Global Change Biology* **18**, 35–43. doi: 10.1111/j.1365-2486.2011.02517.x, ISSN: 1365-2486.

- Smith P. (2013).** Delivering food security without increasing pressure on land. *Global Food Security* 2, 18–23. doi: 10.1016/j.gfs.2012.11.008, ISSN: 22119124.
- Smith P., M.R. Ashmore, H.I.J. Black, P.J. Burgess, C.D. Evans, T.A. Quine, A.M. Thomson, K. Hicks, and H.G. Orr (2013a).** The role of ecosystems and their management in regulating climate, and soil, water and air quality. *Journal of Applied Ecology* 50, 812–829. doi: 10.1111/1365-2664.12016, ISSN: 00218901.
- Smith W.K., C.C. Cleveland, S.C. Reed, N.L. Miller, and S.W. Running (2012a).** Bioenergy Potential of the United States Constrained by Satellite Observations of Existing Productivity. *Environmental Science & Technology* 46, 3536–3544. doi: 10.1021/es203935d, ISSN: 0013-936X.
- Smith W.N., B.B. Grant, C.A. Campbell, B.G. McConkey, R.L. Desjardins, R. Kröbel, and S.S. Malhi (2012b).** Crop residue removal effects on soil carbon: Measured and inter-model comparisons. *Agriculture, Ecosystems & Environment* 161, 27–38. doi: 10.1016/j.agee.2012.07.024, ISSN: 0167-8809.
- Smith P., P.J. Gregory, D.P. van Vuuren, M. Obersteiner, P. Havlik, M. Rounsevell, J. Woods, E. Stehfest, and J. Bellarby (2010).** Competition for land. *Philosophical Transactions of the Royal Society B: Biological Sciences* 365, 2941–2957. doi: 10.1098/rstb.2010.0127.
- Smith P., H. Haberl, A. Popp, K. Erb, C. Lauk, R. Harper, F.N. Tubiello, A. de Siqueira Pinto, M. Jafari, S. Sohi, O. Masera, H. Böttcher, G. Berndes, M. Bustamante, H. Ahammad, H. Clark, H. Dong, E.A. Elsidig, C. Mbow, N.H. Ravindranath, C.W. Rice, C. Robledo Abad, A. Romanovskaya, F. Sperling, M. Herrero, J.I. House, and S. Rose (2013b).** How much land-based greenhouse gas mitigation can be achieved without compromising food security and environmental goals? *Global Change Biology* 19, 2285–2302. doi: 10.1111/gcb.12160, ISSN: 1365-2486.
- Smith P., D. Martino, Z. Cai, D. Gwary, H.H. Janzen, P. Kumar, B. McCarl, S. Ogle, F. O'Mara, C. Rice, R.J. Scholes, O. Sirotenko, M. Howden, T. McAllister, G. Pan, V. Romanenkov, S. Rose, U. Schneider, and S. Towprayoon (2007).** Agriculture. In: *Chapter 8 of Climate change 2007: Mitigation. Contribution of Working group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. B. Metz, O.R. Davidson, P.R. Bosch, R. Dave, L.A. Meyer, (eds.), Cambridge University Press, Cambridge, UK and New York, USA, pp. 497–540.
- Smith P., D. Martino, Z. Cai, D. Gwary, H. Janzen, P. Kumar, B. McCarl, S. Ogle, F. O'Mara, C. Rice, B. Scholes, O. Sirotenko, M. Howden, T. McAllister, G. Pan, V. Romanenkov, U. Schneider, S. Towprayoon, M. Wattenbach, and J. Smith (2008).** Greenhouse gas mitigation in agriculture. *Philosophical Transactions of the Royal Society B: Biological Sciences* 363, 789–813. doi: 10.1098/rstb.2007.2184, ISSN: 0962-8436, 1471–2970.
- Smith K.A., A.R. Mosier, P.J. Crutzen, and W. Winiwarter (2012c).** The role of N₂O derived from crop-based biofuels, and from agriculture in general, in Earth's climate. *Philosophical Transactions of the Royal Society B: Biological Sciences* 367, 1169–1174. doi: 10.1098/rstb.2011.0313, ISSN: 0962-8436, 1471–2970.
- Smith P., and J.E. Olesen (2010).** Synergies between the mitigation of, and adaptation to, climate change in agriculture. *Journal of Agricultural Science* 148, 543–552. Available at: <http://journals.cambridge.org/production/action/cjoGetFulltext?fulltextid=7796512>.
- Smith P., D.S. Powlson, J.U. Smith, P. Falloon, and K. Coleman (2000).** Meeting Europe's climate change commitments: quantitative estimates of the potential for carbon mitigation by agriculture. *Global Change Biology* 6, 525–539. doi: 10.1046/j.1365-2486.2000.00331.x, ISSN: 1365-2486.
- Smith K.A., and T.D. Searchinger (2012).** Crop-based biofuels and associated environmental concerns. *GCB Bioenergy* 4, 479–484. doi: 10.1111/j.1757-1707.2012.01182.x, ISSN: 1757-1707.
- Smith L., and M. Torn (2013).** Ecological limits to terrestrial biological carbon dioxide removal. *Climatic Change* 118, 89–103. doi: 10.1007/s10584-012-0682-3, ISSN: 0165-0009.
- Smith P., and E. Trines (2006).** COMMENTARY: Agricultural measures for mitigating climate change: will the barriers prevent any benefits to developing countries? *International Journal of Agricultural Sustainability* 4, 173–175.
- Smith P., and E. Wollenberg (2012).** Achieving mitigation through synergies with adaptation. In: *Climate Change Mitigation and Agriculture*. E. Wollenberg, A. Nihart, M. Tapio-Biström, M. Grieg-Gran, (eds.), Earthscan, London, UK, pp. 50–57.
- Smith W.K., M. Zhao, and S.W. Running (2012d).** Global Bioenergy Capacity as Constrained by Observed Biospheric Productivity Rates. *BioScience* 62, 911–922. doi: 10.1525/bio.2012.62.10.11, ISSN: 0006-3568, 1525–3244.
- Sneddon C., R.B. Howarth, and R.B. Norgaard (2006).** Sustainable development in a post-Brundtland world. *Ecological Economics* 57, 253–268.
- Snyder C.S., T.W. Bruulsema, and T.L. Jensen (2007).** *Greenhouse Gas Emissions from Cropping Systems and the Influence of Fertilizer Management*. International Plant Nutrition Institute, Norcross, Georgia, USA, 36 pp. Available at: <http://npg.ipni.net/article/NPG-3004>.
- Soares-Filho B., P. Moutinho, D. Nepstad, A. Anderson, H. Rodrigues, R. Garcia, L. Dietzsch, F. Merry, M. Bowman, L. Hissa, R. Silvestrini, and C. Maretti (2010).** Role of Brazilian Amazon protected areas in climate change mitigation. *Proceedings of the National Academy of Sciences* 107, 10821–10826. Available at: <http://www.pnas.org/content/107/24/10821.short>.
- Sochacki S.J., R.J. Harper, and K.R.J. Smettem (2012).** Bio-mitigation of carbon following afforestation of abandoned salinized farmland. *GCB Bioenergy* 4, 193–201. doi: 10.1111/j.1757-1707.2011.01139.x, ISSN: 17571693.
- Sochacki S.J., R.J. Harper, K.R.J. Smettem, B. Dell, and H. Wu (2013).** Evaluating a sustainability index for nutrients in a short rotation energy cropping system. *GCB Bioenergy* 5, 315–326. doi: 10.1111/j.1757-1707.2012.01202.x, ISSN: 1757-1707.
- Sohi S.P. (2012).** Carbon storage with benefits. *Science* 338, 1034–1035.
- Sohngen B., and R. Sedjo (2006).** Carbon Sequestration in Global Forests Under Different Carbon Price Regimes. *The Energy Journal* 27, 109. Available at: <http://www.iaee.org/en/publications/ejarticle.aspx?id=2188>.
- Solli C., M. Reenaas, A. Strømman, and E. Hertwich (2009).** Life cycle assessment of wood-based heating in Norway. *The International Journal of Life Cycle Assessment* 14, 517–528. doi: 10.1007/s11367-009-0086-4, ISSN: 0948-3349.
- Sood K.K., and C.P. Mitchell (2011).** Household level domestic fuel consumption and forest resource in relation to agroforestry adoption: Evidence against need-based approach. *Biomass and Bioenergy* 35, 337–345. Available at: <http://www.sciencedirect.com/science/article/pii/S0961953410003028>.
- Souza S.P., M.T. de Ávila, and S. Pacca (2012a).** Life cycle assessment of sugarcane ethanol and palm oil biodiesel joint production. *Biomass and Bioenergy* 44, 70–79. doi: 10.1016/j.biombioe.2012.04.018, ISSN: 0961-9534.

- Souza H.N. de, R.G.M. de Goede, L. Brussaard, I.M. Cardoso, E.M.G. Duarte, R.B.A. Fernandes, L.C. Gomes, and M.M. Pulleman (2012b). Protective shade, tree diversity and soil properties in coffee agroforestry systems in the Atlantic Rainforest biome. *Agriculture, Ecosystems & Environment* **146**, 179–196. doi: 10.1016/j.agee.2011.11.007, ISSN: 01678809.
- Sow M., C. Hély, C. Mbow, and B. Sambou (2013). Fuel and fire behavior analysis for early-season prescribed fire planning in Sudanian and Sahelian savannas. *Journal of Arid Environments* **89**, 84–93. doi: 10.1016/j.jaridenv.2012.09.007, ISSN: 01401963.
- Sparovek G., G. Berndes, A. Egeskog, F.L.M. de Freitas, S. Gustafsson, and J. Hansson (2007). Sugarcane ethanol production in Brazil: an expansion model sensitive to socioeconomic and environmental concerns. *Biofuels, Bioproducts and Biorefining* **1**, 270–282. doi: 10.1002/bbb.31, ISSN: 1932-1031.
- Spath P.L., and M.K. Mann (2004). *Biomass Power and Conventional Fossil Systems with and without CO₂ Sequestration—Comparing the Energy Balance, Greenhouse Gas Emissions and Economics*. National Renewable Energy Laboratory (NREL), Golden, CO., 38 pp. Available at: <http://www.nrel.gov/docs/fy04osti/32575.pdf>.
- Sperling F., C. Validivia, R. Quiroz, R. Valdivia, L. Angulo, A. Seimon, and I. Noble (2008). *Transitioning to Climate Resilient Development: Perspectives from Communities in Peru*. The World Bank, Washington, D.C., 103 pp.
- Spokas K.A. (2013). Impact of biochar field aging on laboratory greenhouse gas production potentials. *GCB Bioenergy* **5**, 165–176. doi: 10.1111/gcbb.12005, ISSN: 17571693.
- Steenblik R. (2007). *Biofuels—At What Cost? Government Support for Ethanol in Selected OECD*. International Institute for Sustainable Development, Winnipeg, Canada, 82 pp. Available at: <http://www.iisd.org/publications/pub.aspx?id=895>.
- Stehfest E., L. Bouwman, D.P. Vuuren, M.G.J. Elzen, B. Eickhout, and P. Kabat (2009). Climate benefits of changing diet. *Climatic Change* **95**, 83–102. doi: 10.1007/s10584-008-9534-6, ISSN: 0165-0009, 1573–1480.
- Steiner C., K.C. Das, N. Melear, and D. Lakly (2010). Reducing Nitrogen Loss during Poultry Litter Composting Using Biochar. *Journal of Environment Quality* **39**, 1236. doi: 10.2134/jeq2009.0337, ISSN: 1537-2537.
- Steinfeld H., P. Gerber, T. Wassenaar, V. Castel, M. Rosales, and C. de Haan (2008). *Livestock's Long Shadow. Environmental Issues and Options*. Food and Agriculture Organization of the United Nations (FAO), Rome, 416 pp. Available at: <http://www.fao.org/docrep/010/a0701e/a0701e00.HTM>.
- Steinfeld H., H.A. Mooney, F. Schneider, and L.E. Neville (Eds.) (2010). *Livestock in a Changing Landscape: Drivers, Consequences, and Responses*. Island Press, Washington, DC, 416 pp. ISBN: ISBN-13: 9781597266703.
- Stephenson A.L., P. Dupree, S.A. Scott, and J.S. Dennis (2010). The environmental and economic sustainability of potential bioethanol from willow in the UK. *Bioresource Technology* **101**, 9612–9623. doi: 10.1016/j.biortech.2010.07.104, ISSN: 0960-8524.
- Sterner M., and U. Fritsche (2011). Greenhouse gas balances and mitigation costs of 70 modern Germany-focused and 4 traditional biomass pathways including land-use change effects. *Biomass and Bioenergy* **35**, 4797–4814. doi: 10.1016/j.biombioe.2011.08.024, ISSN: 0961-9534.
- Stevenson B.A., R.L. Parfitt, L.A. Schipper, W.T. Baisden, and P. Mudge (2010). Relationship between soil delta(15)N, C/N and N losses across land uses in New Zealand. *Agriculture Ecosystems & Environment* **139**, 736–741. doi: 10.1016/j.agee.2010.10.020, ISSN: 0167-8809.
- Stickler C.M., D.C. Nepstad, M.T. Coe, D.G. McGRATH, H.O. Rodrigues, W.S. Walker, B.S. Soares-Filho, and E.A. Davidson (2009). The potential ecological costs and cobenefits of REDD: a critical review and case study from the Amazon region. *Global Change Biology* **15**, 2803–2824. doi: 10.1111/j.1365-2486.2009.02109.x, ISSN: 1365-2486.
- Stocker B.D., K. Strassmann, and F. Joos (2011). Sensitivity of Holocene atmospheric CO₂ and the modern carbon budget to early human land use: analyses with a process-based model. *Biogeosciences* **8**, 69–88. doi: 10.5194/bg-8-69-2011, ISSN: 1726-4170.
- Strassburg B.B.N., A.S.L. Rodrigues, M. Gusti, A. Balmford, S. Fritz, M. Obersteiner, R.K. Turner, and T.M. Brooks (2012). Impacts of incentives to reduce emissions from deforestation on global species extinctions. *Nature Climate Change* **2**, 350–355. doi: 10.1038/nclimate1375, ISSN: 1758-678X.
- Strassburg B., K. Turner, B. Fisher, R. Schaeffer, and A. Lovett (2008). *An Empirically-Derived Mechanism of Combined Incentives to Reduce Emissions from Deforestation*. Centre for Social and Economic Research on the Global Environment, Norwich, UK, 1–42 pp. Available at: <https://ueaeprints.uea.ac.uk/24686/>.
- Strassburg B., R.K. Turner, B. Fisher, R. Schaeffer, and A. Lovett (2009). Reducing emissions from deforestation—The “combined incentives” mechanism and empirical simulations. *Global Environmental Change* **19**, 265–278. doi: 10.1016/j.gloenvcha.2008.11.004, ISSN: 09593780.
- Stratton R.W., H.M. Wong, and J.I. Hileman (2011). Quantifying Variability in Life Cycle Greenhouse Gas Inventories of Alternative Middle Distillate Transportation Fuels. *Environmental Science & Technology* **45**, 4637–4644. doi: 10.1021/es102597f, ISSN: 0013-936X.
- Streck C. (2012). Financing REDD+: matching needs and ends. *Current Opinion in Environmental Sustainability* **4**, 628–637. doi: 10.1016/j.cosust.2012.10.001, ISSN: 1877-3435.
- Stromberg P., and A. Gasparatos (2012). Biofuels at the confluence of energy security, rural development and food security: a developing country perspective. In: *Socio-economic and environmental impacts of biofuels. Evidence from developing countries*. Cambridge University Press, Cambridge, UK and New York, USA, pp. 1–375.
- Stupak I., B. Lattimore, B.D. Titus, and C. Tattersall Smith (2011). Criteria and indicators for sustainable forest fuel production and harvesting: A review of current standards for sustainable forest management. *Biomass and Bioenergy* **35**, 3287–3308. doi: 10.1016/j.biombioe.2010.11.032, ISSN: 09619534.
- Sumathi S., S.P. Chai, and A.R. Mohamed (2008). Utilization of oil palm as a source of renewable energy in Malaysia. *Renewable and Sustainable Energy Reviews* **12**, 2404–2421. doi: 10.1016/j.rser.2007.06.006, ISSN: 1364-0321.
- Sun A., R. Davis, M. Starbuck, A. Ben-Amotz, R. Pate, and P.T. Pienkos (2011). Comparative cost analysis of algal oil production for biofuels. *Energy* **36**, 5169–5179. Available at: <http://www.sciencedirect.com/science/article/pii/S0360544211003975>.
- Sunde K., A. Brekke, and B. Solberg (2011). Environmental Impacts and Costs of Hydrotreated Vegetable Oils, Transesterified Lipids and Woody BTL—A Review. *Energies* **4**, 845–877. ISSN: 1996-1073.
- Sunderlin W.D., A. Angelsen, B. Belcher, P. Burgers, R. Nasi, L. Santoso, and S. Wunder (2005). Livelihoods, forests, and conservation in developing countries: An Overview. *World Development* **33**, 1383–1402. doi: 10.1016/j.worlddev.2004.10.004, ISSN: 0305-750X.

- Sundrum A. (2001).** Organic livestock farming: A critical review. *Livestock Production Science* **67**, 207–215. doi: 10.1016/S0301-6226(00)00188-3, ISSN: 0301-6226.
- Swann A., I. Fung, and J. Chiang (2011).** Mid-latitude afforestation shifts general circulation and tropical precipitation. *PNAS*.
- Swann A.L., I.Y. Fung, S. Levis, G.B. Bonan, and S.C. Doney (2010).** Changes in Arctic vegetation amplify high-latitude warming through the greenhouse effect. *Proceedings of the National Academy of Sciences* **107**, 1295.
- Swingland I.R., E.C. Bettelheim, J. Grace, G.T. Prance, and L.S. Saunders (2002).** Carbon, biodiversity, conservation and income: an analysis of a free-market approach to land-use change and forestry in developing and developed countries. *Philosophical Transactions of the Royal Society of London Series a-Mathematical Physical and Engineering Sciences* **360**, 1563–1565. ISSN: 1364-503X.
- Taghizadeh-Toosi A., T.J. Clough, L.M. Condron, R.R. Sherlock, C.R. Anderson, and R.A. Craigie (2011).** Biochar Incorporation into Pasture Soil Suppresses in situ Nitrous Oxide Emissions from Ruminant Urine Patches. *Journal of Environment Quality* **40**, 468. doi: 10.2134/jeq2010.0419, ISSN: 1537-2537.
- Taheripour F., T.W. Hertel, and W.E. Tyner (2011).** Implications of biofuels mandates for the global livestock industry: a computable general equilibrium analysis. *Agricultural Economics* **42**, 325–342. doi: 10.1111/j.1574-0862.2010.00517.x, ISSN: 1574-0862.
- Takimoto, A., Nair, P.K.R., Nair, and V.D. (2008).** Carbon stock and sequestration potential of traditional and improved agroforestry systems in the West African Sahel. *Agriculture, Ecosystems and Environment*, 159–166.
- Talens Peiró L., L. Lombardi, G. Villalba Méndez, and X. Gabarrell i Durany (2010).** Life cycle assessment (LCA) and exergetic life cycle assessment (ELCA) of the production of biodiesel from used cooking oil (UCO). *Energy* **35**, 889–893. doi: 10.1016/j.energy.2009.07.013, ISSN: 0360-5442.
- Tan K.T., K.T. Lee, A.R. Mohamed, and S. Bhatia (2009).** Palm oil: Addressing issues and towards sustainable development. *Renewable and Sustainable Energy Reviews* **13**, 420–427. doi: 10.1016/j.rser.2007.10.001, ISSN: 1364-0321.
- Tanaka K., D.J.A. Johansson, B.C. O'Neill, and J.S. Fuglestedt (2013).** Emission metrics under the 2°C climate stabilization target. *Climatic Change* **117**, 933–941. doi: 10.1007/s10584-013-0693-8, ISSN: 0165-0009, 1573–1480.
- Tarnocai C. (2006).** The effect of climate change on carbon in Canadian peatlands. *Global and Planetary Change* **53**, 222–232.
- Tavoni M., and R. Socolow (2013).** Modeling meets science and technology: an introduction to a special issue on negative emissions. *Climatic Change* **118**, 1–14. doi: 10.1007/s10584-013-0757-9, ISSN: 0165-0009.
- Thamsiriroj T., and J.D. Murphy (2011).** The impact of the life cycle analysis methodology on whether biodiesel produced from residues can meet the EU sustainability criteria for biofuel facilities constructed after 2017. *Renewable Energy* **36**, 50–63. doi: 10.1016/j.renene.2010.05.018, ISSN: 0960-1481.
- Thomas S., P. Dargusch, S. Harrison, and J. Herbohn (2010).** Why are there so few afforestation and reforestation Clean Development Mechanism projects? *Land Use Policy* **27**, 880–887. Available at: <http://www.sciencedirect.com/science/article/pii/S026483770900204X>.
- Thomassen M.A., R. Dalgaard, R. Heijungs, and I. de Boer (2008).** Attributional and consequential LCA of milk production. *The International Journal of Life Cycle Assessment* **13**, 339–349. doi: 10.1007/s11367-008-0007-y, ISSN: 0948-3349, 1614–7502.
- Thompson M.C., M. Baruah, and E.R. Carr (2011a).** Seeing REDD+ as a project of environmental governance. *Environmental Science & Policy* **14**, 100–110. doi: 10.1016/j.envsci.2010.11.006, ISSN: 1462-9011.
- Thompson I., B. Mackey, S. McNulty, and A. Mosseler (2009).** *Forest Resilience, Biodiversity, and Climate Change: A Synthesis of the Biodiversity/Resilience/Stability Relationship in Forest Ecosystems*. Secretariat of the Convention on Biological Diversity, Montreal, Canada, 67 pp. Available at: <http://www.cakex.org/virtual-library/1233>.
- Thompson W., J. Whistance, and S. Meyer (2011b).** Effects of US biofuel policies on US and world petroleum product markets with consequences for greenhouse gas emissions. *Energy Policy* **39**, 5509–5518. doi: 10.1016/j.enpol.2011.05.011, ISSN: 0301-4215.
- Thornton P.K., and M. Herrero (2010).** Potential for Reduced Methane and Carbon Dioxide Emissions from Livestock and Pasture Management in the Tropics. *Proceedings of the National Academy of Sciences* **107**, 19667–19672. doi: 10.1073/pnas.0912890107, ISSN: 0027-8424, 1091–6490.
- Throop H.L., and S.R. Archer (2008).** Shrub (*Prosopis velutina*) encroachment in a semidesert grassland: spatial–temporal changes in soil organic carbon and nitrogen pools. *Global Change Biology* **14**, 2420–2431. doi: 10.1111/j.1365-2486.2008.01650.x, ISSN: 1365-2486.
- Tilman D., C. Balzer, J. Hill, and B.L. Befort (2011).** Global food demand and the sustainable intensification of agriculture. *Proceedings of the National Academy of Sciences* **108**, 20260–20264. doi: 10.1073/pnas.1116437108, ISSN: 0027-8424, 1091–6490.
- Tilman D., J. Hill, and C. Lehman (2006).** Carbon-Negative Biofuels from Low-Input High-Diversity Grassland Biomass. *Science* **314**, 1598–1600. doi: 10.1126/science.1133306, ISSN: 0036-8075, 1095–9203.
- Tilman D., R. Socolow, J.A. Foley, J. Hill, E. Larson, L. Lynd, S. Pacala, J. Reilly, T. Searchinger, C. Somerville, and R. Williams (2009).** Beneficial Biofuels –The Food, Energy, and Environment Trilemma. *Science* **325**, 270–271. doi: 10.1126/science.1177970, ISSN: 0036-8075, 1095–9203.
- Timilsina G.R., J.C. Beghin, D. van der Mensbrugge, and S. Mevel (2012).** The impacts of biofuels targets on land-use change and food supply: A global CGE assessment. *Agricultural Economics* **43**, 315–332. doi: 10.1111/j.1574-0862.2012.00585.x, ISSN: 1574-0862.
- Tiwary A., and J. Colls (2010).** Mitigating secondary aerosol generation potentials from biofuel use in the energy sector. *Science of The Total Environment* **408**, 607–616. doi: 10.1016/j.scitotenv.2009.10.019, ISSN: 0048-9697.
- Townsend P.V., R.J. Harper, P.D. Brennan, C. Dean, S. Wu, K.R.J. Smettem, and S.E. Cook (2012).** Multiple environmental services as an opportunity for watershed restoration. *Forest Policy and Economics* **17**, 45–58.
- Trabucco A., R.J. Zomer, D.A. Bossio, O. van Straaten, and L.V. Verchot (2008).** Climate change mitigation through afforestation/reforestation: A global analysis of hydrologic impacts with four case studies. *Agriculture, Ecosystems & Environment* **126**, 81–97. doi: 10.1016/j.agee.2008.01.015, ISSN: 0167-8809.
- Tracy B.P., S.W. Jones, A.G. Fast, D.C. Indurthi, and E.T. Papoutsakis (2012).** Clostridia: the importance of their exceptional substrate and metabolite diversity for biofuel and biorefinery applications. *Current Opinion in Biotechnology* **23**, 364–381. doi: 10.1016/j.copbio.2011.10.008, ISSN: 0958-1669.
- Triantafyllidis K., A. Lappas, and M. Stöcker (2013).** *The Role of Catalysis for the Sustainable Production of Bio-Fuels and Bio-Chemicals*. Elsevier, Oxford, UK, 608 pp. ISBN: 0444563326.

- Tsao C.-C., J.E. Campbell, M. Mena-Carrasco, S.N. Spak, G.R. Carmichael, and Y. Chen (2012). Increased estimates of air-pollution emissions from Brazilian sugar-cane ethanol. *Nature Climate Change* 2, 53–57. doi: 10.1038/nclimate1325, ISSN: 1758-678X, 1758–6798.
- Tubiello F.N., A. Rahman, W. Mann, J. Schmidhuber, M. Koleva, and A. Müller (2009). Carbon financial mechanisms for agriculture and rural development: challenges and opportunities along the Bali roadmap. An editorial essay. *Climatic Change* 97, 3–21.
- Tubiello F., M. Salvatore, S. Rossi, and A. Ferrara (2012). Analysis of global emissions, carbon intensity and efficiency of food production. *Energia Ambiente e Innovazione Anno 2012*, 87–93. Available at: <http://www.enea.it/it/produzione-scientifica/EAI/anno-2012/n.-4-5-luglio-ottobre-parte-I/analysis-of-global-emissions-carbon-intensity-and-efficiency-of-food-production>.
- Tubiello F.N., M. Salvatore, S. Rossi, A. Ferrara, N. Fitton, and P. Smith (2013). The FAOSTAT database of greenhouse gas emissions from agriculture. *Environmental Research Letters* 8, 1–11. doi: 10.1088/1748-9326/8/1/015009, ISSN: 1748-9326.
- Turconi R., A. Boldrin, and T. Astrup (2013). Life cycle assessment (LCA) of electricity generation technologies: Overview, comparability and limitations. *Renewable and Sustainable Energy Reviews* 28, 555–565. doi: 10.1016/j.rser.2013.08.013, ISSN: 1364-0321.
- Turkenburg W.C., D.J. Arent, R. Bertani, A. Faaij, M. Hand, W. Krewitt, E.D. Larson, J. Lund, M. Mehos, T. Merrigan, C. Mitchell, J.R. Moreira, W. Sinke, V. Sonntag-O'Brien, B. Thresher, W. van Sark, E. Usher, and E. Usher (2012). Chapter 11—Renewable Energy. In: *Global Energy Assessment—Toward a Sustainable Future*. Cambridge University Press, Cambridge, UK and New York, NY, USA and the International Institute for Applied Systems Analysis, Laxenburg, Austria, pp. 761–900. ISBN: 9781 10700 5198 HARDBACK 9780 52118 2935 PAPERBACK.
- Turner B.L., E.F. Lambin, and A. Reenberg (2007). The emergence of land change science for global environmental change and sustainability. *Proceedings of the National Academy of Sciences* 104, 20666–20671. doi: 10.1073/pnas.0704119104.
- Tyagi L., B. Kumari, and S.N. Singh (2010). Water management—A tool for methane mitigation from irrigated paddy fields. *Science of The Total Environment* 408, 1085–1090. doi: 10.1016/j.scitotenv.2009.09.010, ISSN: 00489697.
- U.S. DOE (2011). *Billion-Ton Update: Biomass Supply for a Bioenergy and Bioproducts Industry*. U.S. Department of Energy, Washington, DC, 235 pp.
- U.S. EPA (2006). Global Mitigation of Non-CO₂ Greenhouse Gases (EPA Report 430-R-06–005). Available at: <http://www.epa.gov/climatechange/economics/international.html>.
- U.S. EPA (2011). *Draft: Global Anthropogenic Non-CO₂ Greenhouse Gas Emissions: 1990—2030*. Washington, DC, Available at: internal-pdf://US-EPA_NonCO2_Projections_2011_draft-2650857473/US-EPA_NonCO2_Projections_2011_draft.pdf.
- U.S. EPA (2013). U.S. Environmental Protection Agency Global Emissions Database. *Global Emissions*. Available at: <http://www.epa.gov/climatechange/ghgemissions/global.html>.
- UNCCD (2011). *UNCCD Statement at UNFCCC COP17 SBSTA, Agenda Item Three. 28 November 2011*. United Nations Convention to Combat Desertification, Bonn, 45 pp.
- UNDP International Poverty Centre (2006). *What Is Poverty? Concept and Measures*. United Nations Development Programme, Brasilia, Brazil, 24 pp. Available at: www.ipc-undp.org/pub/IPCPovertyInFocus9.pdf.
- UNEP (2009). *Assessing Biofuels, Towards Sustainable Production and Use of Resources*. United Nations Environment Programme (UNEP), Division of Technology, Industry and Economics, Paris, 120 pp.
- UNEP-WCMC (2011). *The UK National Ecosystem Assessment: Technical Report*. UK National Ecosystem Assessment, Cambridge, UK, 6 pp. Available at: <http://uknea.unep-wcmc.org/Resources/tabid/82/Default.aspx>.
- UNFCCC United Nations Framework Convention on Climate Change. Available at: <http://unfccc.int/2860.php>.
- UNFCCC—CDM CDM: CDM insights—intelligence about the CDM at the end of each month. Available at: <http://cdm.unfccc.int/Statistics/Public/CDMinights/index.html>.
- Ungerer J., L. Tao, M. Davis, M. Ghirardi, P.-C. Maness, and J. Yu (2012). Sustained photosynthetic conversion of CO₂ to ethylene in recombinant cyanobacterium *Synechocystis* 6803. *Energy & Environmental Science* 5, 8998–9006.
- UN-REDD Programme UN-REDD Programme. Available at: <http://www.un-redd.org/>.
- UN-REDD Programme -- Support to Partner Countries UN-REDD Programme -- Support to Partner Countries. Available at: http://www.un-redd.org/Partner_Countries/tabid/102663/Default.aspx.
- Ürge-Vorsatz D., N. Eyre, P. Graham, D. Harvey, E. Hertwich, Y. Jiang, and C. Kornevall (2012). Towards Sustainable Energy End-Use: Buildings Chapter 10. In: *Global Energy Assessment*. Cambridge University Press, Cambridge, pp. 649–760.
- Valente C., R. Spinelli, and B.G. Hillring (2011). LCA of environmental and socio-economic impacts related to wood energy production in alpine conditions: Valle di Fiemme (Italy). *Journal of Cleaner Production* 19, 1931–1938. doi: 10.1016/j.jclepro.2011.06.026, ISSN: 0959-6526.
- VanderZaag A.C., S. Jayasundara, and C. Wagner-Riddle (2011). Strategies to mitigate nitrous oxide emissions from land applied manure. *Animal Feed Science and Technology* 166–167, 464–479. doi: 10.1016/j.anifeedsci.2011.04.034, ISSN: 0377-8401.
- Van de Velde L., W. Verbeke, M. Popp, J. Buysse, and G. Van Huylenbroeck (2009). Perceived importance of fuel characteristics and its match with consumer beliefs about biofuels in Belgium. *Energy Policy* 37, 3183–3193. doi: 10.1016/j.enpol.2009.04.022, ISSN: 0301-4215.
- Venter O., W.F. Laurance, T. Iwamura, K.A. Wilson, R.A. Fuller, and H.P. Possingham (2009). Harnessing Carbon Payments to Protect Biodiversity. *Science* 326, 1368–1368. doi: 10.1126/science.1180289, ISSN: 0036-8075, 1095–9203.
- Verchot L.V., M. Noordwijk, S. Kandji, T. Tomich, C. Ong, A. Albrecht, J. Mackensen, C. Bantilan, K.V. Anupama, and C. Palm (2007). Climate change: linking adaptation and mitigation through agroforestry. *Mitigation and Adaptation Strategies for Global Change* 12, 901–918. doi: 10.1007/s11027-007-9105-6, ISSN: 1381-2386, 1573–1596.
- Verdonk M., C. Dieperink, and A.P.C. Faaij (2007). Governance of the emerging bio-energy markets. *Energy Policy* 35, 3909–3924.
- Vermeulen S.J., B.M. Campbell, and J.S.I. Ingram (2012). Climate Change and Food Systems. *Annual Review of Environment and Resources* 37, 195–222. doi: 10.1146/annurev-environ-020411-130608, ISSN: 1543-5938, 1545–2050.

- Viana K.R.O., and R. Perez (2013). Survey of sugarcane industry in Minas Gerais, Brazil: Focus on sustainability. *Biomass and Bioenergy* **58**, 149–157. doi: 10.1016/j.biombioe.2013.08.006, ISSN: 0961-9534.
- Villamor G.B., M. van Noordwijk, Q.B. Le, B. Lusiana, R. Matthews, and P.L.G. Vlek (2011). Diversity deficits in modelled landscape mosaics. *Ecological Informatics* **6**, 73–82. doi: 10.1016/j.ecoinf.2010.08.003, ISSN: 1574-9541.
- Visseren-Hamakers I.J., C. McDermott, M.J. Vijge, and B. Cashore (2012). Trade-offs, co-benefits and safeguards: current debates on the breadth of REDD+. *Current Opinion in Environmental Sustainability* **4**, 646–653. doi: 10.1016/j.cosust.2012.10.005, ISSN: 1877-3435.
- Vlek C. (2004). Environmental Versus Individual Risk Taking: Perception, Decision, Behavior. In: *Encyclopedia of Applied Psychology*. Elsevier, New York, pp. 825–840. ISBN: 978-0-12-657410-4.
- Vlek C., and G. Keren (1992). Behavioral decision theory and environmental risk management: Assessment and resolution of four “survival” dilemmas. *Acta Psychologica* **80**, 249–278. doi: 10.1016/0001-6918(92)90050-N, ISSN: 0001-6918.
- Van Vliet O., M. van den Broek, W. Turkenburg, and A. Faaij (2011a). Combining hybrid cars and synthetic fuels with electricity generation and carbon capture and storage. *Energy Policy* **39**, 248–268. doi: 10.1016/j.enpol.2010.09.038, ISSN: 0301-4215.
- Van Vliet O., A.S. Brouwer, T. Kuramochi, M. van den Broek, and A. Faaij (2011b). Energy use, cost and CO₂ emissions of electric cars. *Journal of Power Sources* **196**, 2298–2310. doi: 10.1016/j.jpowsour.2010.09.119, ISSN: 0378-7753.
- Van Vliet O.P., A.P. Faaij, and W.C. Turkenburg (2009). Fischer–Tropsch diesel production in a well-to-wheel perspective: a carbon, energy flow and cost analysis. *Energy Conversion and Management* **50**, 855–876.
- Van der Voet E., R.J. Lifset, and L. Luo (2010). Life-cycle assessment of biofuels, convergence and divergence. *Biofuels* **1**, 435–449. doi: 10.4155/bfs.10.19, ISSN: 1759-7269, 1759–7277.
- De Vries M., and I.J.M. de Boer (2010). Comparing environmental impacts for livestock products: A review of life cycle assessments. *Livestock Science* **128**, 1–11. doi: 10.1016/j.livsci.2009.11.007, ISSN: 1871-1413.
- Van Vuuren D.P., J. van Vliet, and E. Stehfest (2009). Future bio-energy potential under various natural constraints. *Energy Policy* **37**, 4220–4230. doi: 10.1016/j.enpol.2009.05.029, ISSN: 0301-4215.
- Waghorn G. (2008). Beneficial and detrimental effects of dietary condensed tannins for sustainable sheep and goat production—Progress and challenges. *Animal Feed Science and Technology* **147**, 116–139.
- Waghorn G.C., and R.S. Hegarty (2011). Lowering ruminant methane emissions through improved feed conversion efficiency. *Animal Feed Science and Technology* **166–167**, 291–301. doi: 10.1016/j.anifeedsci.2011.04.019, ISSN: 0377-8401.
- Walter A., P. Dolzan, O. Quilodrán, J.G. de Oliveira, C. da Silva, F. Piacente, and A. Segerstedt (2011). Sustainability assessment of bio-ethanol production in Brazil considering land use change, GHG emissions and socio-economic aspects. *Energy Policy* **39**, 5703–5716. doi: 10.1016/j.enpol.2010.07.043, ISSN: 0301-4215.
- Wang M.Q., J. Han, Z. Haq, W.E. Tyner, M. Wu, and A. Elgowainy (2011a). Energy and greenhouse gas emission effects of corn and cellulosic ethanol with technology improvements and land use changes. *Biomass and Bioenergy* **35**, 1885–1896. doi: 10.1016/j.biombioe.2011.01.028, ISSN: 0961-9534.
- Wang M., H. Huo, and S. Arora (2011b). Methods of dealing with co-products of biofuels in life-cycle analysis and consequent results within the U.S. context. *Energy Policy* **39**, 5726–5736. doi: 10.1016/j.enpol.2010.03.052, ISSN: 0301-4215.
- Wang L., J. Littlewood, and R.J. Murphy (2013a). Environmental sustainability of bioethanol production from wheat straw in the UK. *Renewable and Sustainable Energy Reviews* **28**, 715–725. doi: 10.1016/j.rser.2013.08.031, ISSN: 1364-0321.
- Wang X., O.R. Masera, K. Troncoso, M.X. Rivera, and J. Franco (2013b). *What Have We Learned about Household Biomass Cooking in Central America?* The World Bank, 1–134 pp. Available at: <http://documents.worldbank.org/curated/en/2013/01/17524967/learned-household-biomass-cooking-central-america>.
- Wang Y., P. Yu, K.-H. Feger, X. Wei, G. Sun, M. Bonell, W. Xiong, S. Zhang, and L. Xu (2011c). Annual runoff and evapotranspiration of forestlands and non-forestlands in selected basins of the Loess Plateau of China. *Ecohydrology* **4**, 277–287. doi: 10.1002/eco.215, ISSN: 19360584.
- Warner E., Y. Zhang, D. Inman, and G. Heath (2013). Challenges in the estimation of greenhouse gas emissions from biofuel-induced global land-use change. *Biofuels, Bioproducts and Biorefining* **8**, 114–125. doi: 10.1002/bbb.1434, ISSN: 1932-1031.
- WBGU (2009). *Future Bioenergy and Sustainable Land Use*. Earthscan, London, 393 pp.
- WBGU (2011). *Welt Im Wandel. Gesellschaftsvertrag Für Eine Große Transformation*. Wissenschaftlicher Beirat Globale Umweltveränderungen (WBGU), Berlin, 421 pp. ISBN: 978-3-936191-46-2.
- Van der Werf G.R., J.T. Randerson, L. Giglio, G.J. Collatz, P.S. Kasibhatla, A.F. Arellano Jr, and others (2006). Interannual variability of global biomass burning emissions from 1997 to 2004. *Atmospheric Chemistry and Physics Discussions* **6**, 3175–3226. Available at: <http://hal.archives-ouvertes.fr/hal-00301203/>.
- Van der Werf G.R., J.T. Randerson, L. Giglio, G.J. Collatz, M. Mu, P.S. Kasibhatla, D.C. Morton, R.S. DeFries, Y. Jin, and T.T. van Leeuwen (2010). Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009). *Atmospheric Chemistry and Physics* **10**, 11707–11735. doi: 10.5194/acp-10-11707-2010, ISSN: 1680-7316.
- Werner F., R. Taverna, P. Hofer, E. Thürig, and E. Kaufmann (2010). National and global greenhouse gas dynamics of different forest management and wood use scenarios: a model-based assessment. *Environmental Science & Policy* **13**, 72–85. doi: 10.1016/j.envsci.2009.10.004, ISSN: 1462-9011.
- West P.C., H.K. Gibbs, C. Monfreda, J. Wagner, C.C. Barford, S.R. Carpenter, and J.A. Foley (2010a). Trading carbon for food: Global comparison of carbon stocks vs. crop yields on agricultural land. *Proceedings of the National Academy of Sciences* **107**, 19645–19648. doi: 10.1073/pnas.1011078107, ISSN: 0027-8424, 1091–6490.
- West P.C., G.T. Narisma, C.C. Barford, C.J. Kucharik, and J.A. Foley (2010b). An alternative approach for quantifying climate regulation by ecosystems. *Frontiers in Ecology and the Environment* **9**, 126–133.
- Whitaker J., K.E. Ludley, R. Rowe, G. Taylor, and D.C. Howard (2010). Sources of variability in greenhouse gas and energy balances for biofuel production: a systematic review. *GCB Bioenergy* **2**, 99–112. doi: 10.1111/j.1757-1707.2010.01047.x, ISSN: 1757-1707.

- Whitehead P., P. Purdon, J. Russel-Smith, P.M. Cooke, and S. Sutton (2008).** The management of climate change through prescribed Savanna burning: Emerging contributions of indigenous people in Northern Australia. *Public Administration and Development Special Issue: Symposium on Climate Change, Governance and Environmental Services* **28**, 374–385.
- Whittaker C., N. Mortimer, R. Murphy, and R. Matthews (2011).** Energy and greenhouse gas balance of the use of forest residues for bioenergy production in the UK. *Biomass and Bioenergy* **35**, 4581–4594. doi: 10.1016/j.biombioe.2011.07.001, ISSN: 0961-9534.
- Wicke B., V. Dornburg, M. Junginger, and A. Faaij (2008).** Different palm oil production systems for energy purposes and their greenhouse gas implications. *Biomass and Bioenergy* **32**, 1322–1337.
- Wicke B., R. Sikkema, V. Dornburg, and A. Faaij (2011a).** Exploring land use changes and the role of palm oil production in Indonesia and Malaysia. *Land Use Policy* **28**, 193–206. doi: 10.1016/j.landusepol.2010.06.001, ISSN: 0264-8377.
- Wicke B., E.M.W. Smeets, R. Akanda, L. Stille, R.K. Singh, A.R. Awan, K. Mahmood, and A.P.C. Faaij (2013).** Biomass production in agroforestry and forestry systems on salt-affected soils in South Asia: Exploration of the GHG balance and economic performance of three case studies. *Journal of Environmental Management* **127**, 324–334. doi: 10.1016/j.jenvman.2013.05.060, ISSN: 0301-4797.
- Wicke B., E. Smeets, V. Dornburg, B. Vashev, T. Gaiser, W. Turkenburg, and A. Faaij (2011b).** The global technical and economic potential of bioenergy from salt-affected soils. *Energy & Environmental Science* **4**, 2669–2681.
- Wicke B., E. Smeets, A. Tabeau, J. Hilbert, and A. Faaij (2009).** Macroeconomic impacts of bioenergy production on surplus agricultural land—A case study of Argentina. *Renewable and Sustainable Energy Reviews* **13**, 2463–2473. doi: 16/j.rser.2009.05.010, ISSN: 1364-0321.
- Wicke B., E. Smeets, H. Watson, and A. Faaij (2011c).** The current bioenergy production potential of semi-arid and arid regions in sub-Saharan Africa. *Biomass and Bioenergy* **35**, 2773–2786. doi: 10.1016/j.biombioe.2011.03.010, ISSN: 0961-9534.
- Wicke B., P. Verweij, H. van Meijl, D.P. van Vuuren, and A.P. Faaij (2012).** Indirect land use change: review of existing models and strategies for mitigation. *Biofuels* **3**, 87–100. doi: 10.4155/bfs.11.154, ISSN: 1759-7269, 1759-7277.
- Wilkinson J., and S. Herrera (2010).** Biofuels in Brazil: debates and impacts. *Journal of Peasant Studies* **37**, 749–768.
- Williams J., and P.J. Crutzen (2010).** Nitrous oxide from aquaculture. *Nature Geoscience* **3**, 143–143. Available at: <http://www.nature.com/ngeo/journal/vaop/ncurrent/full/ngeo804.html>.
- Williams R.H., G. Liu, T.G. Kreutz, and E.D. Larson (2011).** Coal and Biomass to Fuels and Power. *Annual Review of Chemical and Biomolecular Engineering* **2**, 529–553. Available at: <http://www.annualreviews.org/doi/abs/10.1146/annurev-chembioeng-061010-114126>.
- Wilson T.O., F.M. McNeal, S. Spataro, D.G. Abler, and P.R. Adler (2011).** Densified biomass can cost-effectively mitigate greenhouse gas emissions and address energy security in thermal applications. *Environmental Science & Technology* **46**, 1270–1277. doi: 10.1021/es202752b, ISSN: 0013-936X.
- Wirsenius S. (2003).** Efficiencies and biomass appropriation of food commodities on global and regional levels. *Agricultural Systems* **77**, 219–255. doi: 10.1016/S0308-521X(02)00188-9, ISSN: 0308-521X.
- Wirsenius S., F. Hedenus, and K. Mohlin (2011).** Greenhouse gas taxes on animal food products: rationale, tax scheme and climate mitigation effects. *Climatic Change* **108**, 159–184. doi: 10.1007/s10584-010-9971-x, ISSN: 0165-0009, 1573–1480.
- Wise M., K. Calvin, A. Thomson, L. Clarke, B. Bond-Lamberty, R. Sands, S.J. Smith, A. Janetos, and J. Edmonds (2009).** Implications of Limiting CO₂ Concentrations for Land Use and Energy. *Science* **324**, 1183–1186. doi: 10.1126/science.1168475, ISSN: 0036-8075, 1095–9203.
- Wiskerke W.T., V. Dornburg, C.D.K. Rubanza, R.E. Malimbwi, and A.P.C. Faaij (2010).** Cost/benefit analysis of biomass energy supply options for rural smallholders in the semi-arid eastern part of Shinyanga Region in Tanzania. *Renewable and Sustainable Energy Reviews* **14**, 148–165. doi: 10.1016/j.rser.2009.06.001, ISSN: 1364-0321.
- De Wit M., M. Junginger, and A. Faaij (2013).** Learning in dedicated wood production systems: Past trends, future outlook and implications for bioenergy. *Renewable and Sustainable Energy Reviews* **19**, 417–432. doi: 10.1016/j.rser.2012.10.038, ISSN: 1364-0321.
- De Wit M., M. Londo, and A. Faaij (2011).** Productivity developments in European agriculture: Relations to and opportunities for biomass production. *Renewable and Sustainable Energy Reviews* **15**, 2397–2412. doi: 16/j.rser.2011.02.022, ISSN: 1364-0321.
- Witt G.B., R.A. Harrington, and M.J. Page (2009).** Is “vegetation thickening” occurring in Queensland’s mulga lands—a 50-year aerial photographic analysis. *Australian Journal of Botany* **57**, 572–582. Available at: <http://dx.doi.org/10.1071/BT08217>.
- Wollenberg, E., Nihart, A., Tapio-Bistrom, M-L., Grieg-Gran, and M. (2012).** *Climate Change Mitigation and Agriculture*. Earthscan, London, UK, 419 pp.
- Woods J., A. Williams, J.K. Hughes, M. Black, and R. Murphy (2010).** Energy and the food system. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 2991–3006. doi: 10.1098/rstb.2010.0172.
- Woods, M.C., and et al. (2007).** *Cost and Performance Baseline for Fossil Energy Plants: Volume 1: Bituminous Coal and Natural Gas to Electricity*. National Energy Technology Laboratory (NETL), Department of Energy (DOE), Washington, D.C., USA., 626 pp.
- Woolf D., J.E. Amonette, F.A. Street-Perrott, J. Lehmann, and S. Joseph (2010).** Sustainable biochar to mitigate global climate change. *Nature Communications* **1**, 1–9. doi: 10.1038/ncomms1053, ISSN: 2041-1723.
- Wright A.-D.G., and A.V. Klieve (2011).** Does the complexity of the rumen microbial ecology preclude methane mitigation? *Animal Feed Science and Technology* **166–67**, 248–253. doi: 10.1016/j.anifeedsci.2011.04.015, ISSN: 0377-8401.
- Wu C., and L. Lin (2009).** Guest editorial. *Biotechnology Advances* **27**, 541. doi: 10.1016/j.biotechadv.2009.04.018, ISSN: 0734-9750.
- Wünscher T., and S. Engel (2012).** International payments for biodiversity services: Review and evaluation of conservation targeting approaches. *Biological Conservation* **152**, 222–230. doi: 10.1016/j.biocon.2012.04.003, ISSN: 0006-3207.
- Ximenes F. de A., B.H. George, A. Cowie, J. Williams, and G. Kelly (2012).** Greenhouse Gas Balance of Native Forests in New South Wales, Australia. *Forests* **3**, 653–683. doi: 10.3390/f3030653.
- Ximenes F.A., and T. Grant (2013).** Quantifying the greenhouse benefits of the use of wood products in two popular house designs in Sydney, Australia. *The International Journal of Life Cycle Assessment* **18**, 891–908. doi: 10.1007/s11367-012-0533-5, ISSN: 0948-3349, 1614–7502.

- Yagi K., H. Tsuruta, and K. Minami (1997).** Possible options for mitigating methane emission from rice cultivation. *Nutrient Cycling in Agroecosystems* **49**, 213–220. Available at: <http://link.springer.com/article/10.1023/A:1009743909716>.
- Yan X., H. Akiyama, K. Yagi, and H. Akimoto (2009).** Global estimations of the inventory and mitigation potential of methane emissions from rice cultivation conducted using the 2006 Intergovernmental Panel on Climate Change Guidelines. *Global Biogeochemical Cycles* **23**, 3–25. doi: 10.1029/2008GB003299.
- Yan T., C.S. Mayne, F.G. Gordon, M.G. Porter, R.E. Agnew, D.C. Patterson, C.P. Ferris, and D.J. Kilpatrick (2010).** Mitigation of enteric methane emissions through improving efficiency of energy utilization and productivity in lactating dairy cows. *Journal of Dairy Science* **93**, 2630–2638. doi: 10.3168/jds.2009-2929, ISSN: 0022-0302.
- Yang Y., J. Bae, J. Kim, and S. Suh (2012).** Replacing Gasoline with Corn Ethanol Results in Significant Environmental Problem-Shifting. *Environmental Science and Technology* **46**, 3671–3678. doi: 10.1021/es203641p, ISSN: 0013-936X.
- Yao Z., Z. Zhou, X. Zheng, B. Xie, B. Mei, R. Wang, K. Butterbach-Bahl, and J. Zhu (2010).** Effects of organic matter incorporation on nitrous oxide emissions from rice-wheat rotation ecosystems in China. *Plant and Soil* **327**, 315–330. doi: 10.1007/s11104-009-0056-4, ISSN: 0032-079X, 1573–5036.
- Yoon J.M., L. Zhao, and J.V. Shanks (2013).** Metabolic Engineering with Plants for a Sustainable Biobased Economy. *Annual Review of Chemical and Biomolecular Engineering* **4**, 211–237. Available at: <http://www.annualreviews.org/doi/abs/10.1146/annurev-chembioeng-061312-103320>.
- Yu P., Y. Wang, A. Du, W. Guan, K.-H. Feger, K. Schwärzel, M. Bonell, W. Xiong, and S. Pan (2013).** The effect of site conditions on flow after forestation in a dryland region of China. *Agricultural and Forest Meteorology* **178–179**, 66–74. doi: 10.1016/j.agrformet.2013.02.007, ISSN: 01681923.
- Zaehle S., P. Ciais, A.D. Friend, and V. Prieur (2011).** Carbon benefits of anthropogenic reactive nitrogen offset by nitrous oxide emissions. *Nature Geoscience* **4**, 601–605. doi: 10.1038/ngeo1207, ISSN: 1752-0894.
- Zaks D.P.M., C.C. Barford, N. Ramankutty, and J.A. Foley (2009).** Producer and consumer responsibility for greenhouse gas emissions from agricultural production—a perspective from the Brazilian Amazon. *Environmental Research Letters* **4**, 1–13. doi: 10.1088/1748-9326/4/4/044010, ISSN: 1748-9326.
- Zanchi G., N. Pena, and N. Bird (2012).** Is woody bioenergy carbon neutral? A comparative assessment of emissions from consumption of woody bioenergy and fossil fuel. *GCB Bioenergy* **4**, 761–772. doi: 10.1111/j.1757-1707.2011.01149.x, ISSN: 1757-1707.
- Zhang Y., Y. Yu, T. Li, and B. Zou (2011).** Analyzing Chinese consumers' perception for biofuels implementation: The private vehicles owner's investigating in Nanjing. *Renewable and Sustainable Energy Reviews* **15**, 2299–2309. doi: 10.1016/j.rser.2011.02.004, ISSN: 1364-0321.
- Zhao M., and S.W. Running (2010).** Drought-Induced Reduction in Global Terrestrial Net Primary Production from 2000 Through 2009. *Science* **329**, 940–943. doi: 10.1126/science.1192666, ISSN: 0036-8075, 1095–9203.
- Zhao W.Z., H.L. Xiao, Z.M. Liu, and J. Li (2005).** Soil degradation and restoration as affected by land use change in the semiarid Bashang area, northern China. *CATENA* **59**, 173–186. doi: 10.1016/j.catena.2004.06.004, ISSN: 03418162.
- Van Zijderveld S.M., W.J.J. Gerrits, J.A. Apajalahti, J.R. Newbold, J. Dijkstra, R.A. Leng, and H.B. Perdok (2010).** Nitrate and sulfate: Effective alternative hydrogen sinks for mitigation of ruminant methane production in sheep. *Journal of Dairy Science* **93**, 5856–5866. doi: DOI: 10.3168/jds.2010-3281, ISSN: 0022-0302.
- Zimmerman A.R., B. Gao, and M.-Y. Ahn (2011).** Positive and negative carbon mineralization priming effects among a variety of biochar-amended soils. *Soil Biology and Biochemistry* **43**, 1169–1179. doi: 10.1016/j.soilbio.2011.02.005, ISSN: 0038-0717.
- Ziv G., E. Baran, S. Nam, I. Rodríguez-Iturbe, and S.A. Levin (2012).** Trading-off fish biodiversity, food security, and hydropower in the Mekong River Basin. *Proceedings of the National Academy of Sciences* **109**, 5609–5614. doi: 10.1073/pnas.1201423109, ISSN: 0027-8424, 1091–6490.
- Zomer R.J., A. Trabucco, D.A. Bossio, and L.V. Verchot (2008).** Climate change mitigation: A spatial analysis of global land suitability for clean development mechanism afforestation and reforestation. *Agriculture, Ecosystems & Environment* **126**, 67–80. doi: 10.1016/j.agee.2008.01.014, ISSN: 0167-8809.
- Van Zwieten L., S.W.L. Kimber, S.G. Morris, B.P. Singh, P.R. Grace, C. Scheer, J. Rust, A.E. Downie, and A.L. Cowie (2013).** Pyrolysing poultry litter reduces N₂O and CO₂ fluxes. *Science of The Total Environment* **465**, 279–287. doi: 10.1016/j.scitotenv.2013.02.054, ISSN: 00489697.