



Metabolically engineered cells for the production of pinosylvin

Katz, Michael; Förster, Jochen; David, Helga Susana Moreira; Schmidt, Hans Peter; Sendelius, Malin; Petersen Bjørn, Sara; Durhuus, Thomas Thomasen

Publication date:
2008

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Katz, M., Förster, J., David, H. S. M., Schmidt, H. P., Sendelius, M., Petersen Bjørn, S., & Durhuus, T. T. (2008). Metabolically engineered cells for the production of pinosylvin. (Patent No. *WO2007EP57484*).

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 January 2008 (24.01.2008)

PCT

(10) International Publication Number
WO 2008/009728 A1

(51) International Patent Classification:
C12P 7/22 (2006.01)

DURHUUS, Thomas Thomasen [DK/DK]; Tuborgvej
243, DK-2400 København NV (DK).

(21) International Application Number:
PCT/EP2007/057484

(74) Agent: SMART, Peter; Fulwood House, 12 Fulwood
Place, London London WC1V 6HR (GB).

(22) International Filing Date: 19 July 2007 (19.07.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0614442.2 20 July 2006 (20.07.2006) GB

(71) Applicant (for all designated States except US): FLUX-
OME SCIENCES A/S [DK/DK]; Diplomvej 378,
DK-2800 Kgs. Lyngby (DK).

(72) Inventors; and


(75) Inventors/Applicants (for US only): KATZ, Michael
[SE/SE]; Ormvråksgatan 9, S-215 62 Malmö (SE).
FÖRSTER, Jochen [DE/DK]; Halmtorvet 7, 5 tv.,
DK-1700 København V (DK). DAVID, Helga [PT/DK];
Baggesensgade 33, 1tv, DK-2200 København N (DK).
SCHMIDT, Hans Peter [NL/DK]; Søllerødvej 39B,
DK-2840 Holte (DK). SENDELIUS, Malin [SE/SE]; Fly-
gelvägen 94, S-224 72 Lund (SE). BJØRN, Sara Petersen
[DK/DK]; Klampenborgvej 102, DK-2800 Lyngby (DK).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH,
CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG,
ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK,
LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW,
MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL,
PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY,
TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL,
PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: METABOLICALLY ENGINEERED CELLS FOR THE PRODUCTION OF PINOSYLVIN



(57) Abstract: A genetically engineered micro-organism having an operative metabolic pathway producing cinnamoyl-CoA and producing pinosylvin therefrom by the action of a stilbene synthase is used for pinosylvin production. Said cinnamic acid may be formed from L-phenylalanine by a L-phenylalanine ammonia lyase (PAL) which is one accepting phenylalanine as a substrate and producing cinnamic acid therefrom, preferably such that if the PAL also accepts tyrosine as a substrate and forms coumaric acid therefrom, the ratio $K_m(\text{phenylalanine})/K_m(\text{tyrosine})$ for said PAL is less than 1:1 and if said micro-organism produces a cinnamate-4- hydroxylase enzyme (C4H), the ratio $K_{cat}(\text{PAL})/K_{cat}(\text{C4H})$ is at least 2:1.

WO 2008/009728 A1



Published:

- with international search report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Metabolically Engineered Cells For The Production Of
Pinosylvin.

FIELD OF THE INVENTION

5 This invention relates generally to the production of the polyphenol pinosylvin. Furthermore, it relates to the use of naturally occurring or recombinant micro-organisms that produce pinosylvin for production of food, feed and beverages.

10

BACKGROUND OF THE INVENTION

Production of chemicals from micro-organisms has been an important application of biotechnology. Typically, the steps in developing such a bio-production method may include 15 1) selection of a proper micro-organism host, 2) elimination of metabolic pathways leading to by-products, 3) deregulation of desired pathways at both enzyme activity level and the transcriptional level, and 4) overexpression of appropriate enzymes in the desired pathways. In 20 preferred aspects, the present invention has employed combinations of the steps above to redirect carbon flow from phenylalanine through enzymes of the plant phenylpropanoid pathway which supplies the necessary precursor for the 25 desired biosynthesis of pinosylvin.

Pinosylvin (or pinosylvine or 3,5-dihydroxy-*trans*-stilbene) is a phytophenol belonging to the group of stilbene phytoalexins, which are low-molecular-mass secondary 30 metabolites that constitute the active defence mechanism in plants in response to infections or other stress-related events. Stilbene phytoalexins contain the stilbene skeleton

(*trans*-1,2-diphenylethylene) as their common basic structure: that may be supplemented by addition of other groups as well (Hart and Shrimpton, 1979, Hart, 1981). Stilbenes have been found in certain trees (angio-sperms, 5 gymnosperms), but also in some herbaceous plants (in species of the *Myrtaceae*, *Vitaceae* and *Leguminosae* families). Said compounds are toxic to pests, especially to fungi, bacteria and insects. Only few plants have the ability to synthesize stilbenes, or to produce them in an amount that provides 10 them sufficient resistance to pests.

The synthesis of the basic stilbene skeleton is pursued by stilbene synthases, which comprises a small gene family in most species examined (Kodan *et al.* 2002). Stilbene synthases 15 appear to have evolved from chalcone synthases, and belong to a polyketide synthase (PKS) superfamily that share more than 65% amino acid homology. Unlike the bacterial PKSs, both stilbene- and chalcone synthases function as unimodular PKSs with a single active site, forming relatively small 20 homodimers (Tropf *et al.*, 1995). Stilbene- and chalcone synthases use common substrates, three malonyl-CoAs and one cinnamoyl-CoA/p-coumaroyl-CoA, forming their products with similar reaction mechanisms (Kindl, 1985). Stilbene 25 synthases can be classified into either a 4-coumaroyl-CoA-specific type that has its highest activity with 4-coumaroyl-CoA as substrate, such as resveratrol synthase (EC 2.3.1.95), or a cinnamoyl-CoA-specific type that has its highest activity with cinnamoyl-CoA as substrate, such as pinosylvin synthase (EC 2.3.1.146). Genes encoding 30 resveratrol synthases have been described earlier for peanut (*Arachis hypogaea*) (Schöppner and Kindl, 1984; Schröder *et al.*, 1988) and grapevine (*Vitis vinifera*) (Melchior and Kindl, 1991; Wiese *et al.*, 1994) whereas genes encoding

pinosylvin synthase have been mostly described for pine (*Pinus sylvestris* and - *strobus*) (Schanz *et al.*, 1992; Raiber *et al.*, 1995; Kodan *et al.*, 2002; Hemingway *et al.*, 1977).

5

Pinosylvin is present in the wood pulp of eucalyptus-, spruce- and pine trees such as *Pinus sylvestris*, - *densiflora*, -*taeda* and -*strobus*. In pine species, the constitutive pinosylvin occurs exclusively in the heartwood (Kindl, 1985). However, the compound is induced in the sapwood, phloem, and needles as a response to wounding, fungal attack or environmental stress such as UV-radiation and ozone exposure (Hart, 1981; Kindl, 1985; Richter and Wild, 1992; Lieutier *et al.*, 1996; Rosemann *et al.*, 1991). The compound possesses potent anti-fungal activity against a wide assortment of fungi (Lindberg *et al.*, 2004; Pacher *et al.*, 2002).

20 Pinosylvin (Fig. 1 *trans*-form) consists of two closely connected phenol rings and belongs therefore to the polyphenols. Unlike most other hydroxystilbenes, pinosylvin lacks a hydroxyl group in ring B (Fig.1) and originates by condensation of unsubstituted cinnamoyl-CoA with three molecules of malonyl-CoA. That said, pinosylvin is structurally similar to the tri-hydroxystilbene resveratrol, which is found in red wine (Aggarwal *et al.*, 2004). Much data has been generated demonstrating the health benefits of resveratrol. For instance resveratrol's potent anticancer activity across many cancer cell lines has well been established (Aggarwal *et al.*, 2004). Given the similarity in structure with resveratrol, it is anticipated that pinosylvin possesses potent health benefits as well. Indeed pinosylvin's effect on various cancers, including

colorectal- and liver cancers, has been studied, and has indicated it's chemopreventative- and anti-leukemic activity (Skinnider and Stoessl, 1986;. Mellanen *et al.*, 1996; Roupe *et al.*, 2005 and 2006). Moreover, pinosylvin has anti-oxidant capacity as well, though to a lesser extent than, for instance, resveratrol (Stojanovic *et al.*, 2001).

Presently, pinosylvin is mostly obtained in a mixture of various flavonoids that is extracted from the bark of pine. Said extraction is a labour intensive process with a low yield. In preferred aspects, the present invention provides novel, more efficient and high-yielding production processes.

In plants, the phenylpropanoid pathway is responsible for the synthesis of a wide variety of secondary metabolic compounds, including lignins, salicylates, coumarins, hydroxycinnamic amides, pigments, flavonoids and phytoalexins. Indeed formation of stilbenes in plants proceeds through the phenylpropanoid pathway. The amino acid L-phenylalanine is converted into *trans*-cinnamic acid through the non-oxidative deamination by L-phenylalanine ammonia lyase (PAL) (Fig 2). From *trans*-cinnamic acid the pathway can branch into a resveratrol-forming route or into a pinosylvin forming route. In the first route *trans*-cinnamic acid is hydroxylated at the *para*-position to 4-coumaric acid (4-hydroxycinnamic acid) by cinnamate-4-hydroxylase (C4H), a cytochrome P450 monooxygenase enzyme, in conjunction with NADPH:cytochrome P450 reductase (CPR). Subsequently, 4-coumaric acid, is then activated to 4-coumaroyl-CoA by the action of 4-coumarate-CoA ligase (4CL). A resveratrol synthase (VST1), can then catalyze the condensation of a phenylpropane unit of 4-coumaroyl-CoA with

malonyl CoA, resulting in formation of resveratrol. In the latter route *trans*-cinnamic acid is directly activated to cinnamoyl-CoA by the action of 4CL where a pinosylvin synthase (PST) subsequently catalyzes the condensation of a phenylpropane unit of cinnamoyl-CoA with malonyl CoA, resulting in formation of pinosylvin.

Stilbene synthases are rather promiscuous enzymes that can accept a variety of physiological and non-physiological substrates. For instance, addition of various phenylpropanoid CoA starter esters led to formation of several products *in vitro* (Ikuro *et al.*, 2004; Morita *et al.*, 2001). Likewise it has been shown that resveratrol synthase from rhubarb (*Rheum tartaricum*) indeed synthesized a small amount of pinosylvin when cinnamoyl-CoA was used as substrate instead of coumaroyl-CoA (Samappito *et al.*, 2003).

Similarly, coumaroyl-CoA ligase can accept both coumaric acid and cinnamic acid as substrate, albeit with a catalytic efficiency (K_m/K_{cat}) that is 100 times less for cinnamic acid compared to coumaric acid (Allina *et al.*, 1998; Ehlting *et al.*, 1999). We deduced from the above that it would be possible to produce pinsosylvin in a pathway that would consist of a 4CL and a stilbene synthase, even one that is designated as a classical resveratrol synthase.

Recently, a yeast was disclosed that could produce resveratrol from coumaric acid that is found in small quantities in grape must (Becker *et al.* 2003, ZA200408194). The production of 4-coumaroyl-CoA from exogenous 4-coumaric acid, and concomitant resveratrol, in laboratory strains of *S. cerevisiae*, was achieved by co-expressing a heterologous

coenzyme-A ligase gene, from hybrid poplar, together with the grapevine resveratrol synthase gene (*VST1*). The other substrate for resveratrol synthase, malonyl-CoA, is already endogenously produced in yeast and is involved in *de novo* fatty-acid biosynthesis. The study showed that cells of *S. cerevisiae* could produce minute amounts of resveratrol, either in the free form or in the glucoside-bound form, when cultured in synthetic media that was supplemented with 4-coumaric acid.

10

Given the promiscuity of the resveratrol synthase, it may be that said yeast could produce pinosylvin as well when fed with substantial amounts of cinnamic acid. However, commercial application of such a yeast would be hampered by the probable low pinosylvin yield, and the need for addition of cinnamic acid, which is not abundantly present in industrial media. Hence, to accelerate and broaden the application of pinosylvin as both a pharmaceutical and neutraceutical, it is highly desirable to provide a yeast or other micro-organism that can produce pinosylvin directly from glucose, without addition of cinnamic acid or any downstream cinnamic acid derivative such as cinnamoyl-CoA.

15

A recent study (Ro and Douglas, 2004) describes the reconstitution of the entry point of the phenylpropanoid pathway in *S. cerevisiae* by introducing PAL, C4H and CPR from Poplar. The purpose was to evaluate whether multienzyme complexes (MECs) containing PAL and C4H are functionally important at this entry point into phenylpropanoid metabolism. By feeding the recombinant yeast with [3H]-phenylalanine it was found that the majority of metabolized [3H]-phenylalanine was incorporated into 4-[3H]-coumaric acid, and that phenylalanine metabolism was

20

25

30

highly reduced by inhibiting C4H activity. Moreover, PAL-alone expressers metabolized very little phenylalanine into cinnamic acid. When feeding [3H]-phenylalanine and [14C]-*trans*-cinnamic acid simultaneously to the triple expressers, 5 no evidence was found for channeling of the endogenously synthesized [3H]-*trans*-cinnamic acid into 4-coumaric acid. Therefore, efficient carbon flux from phenylalanine to 4-coumaric acid via reactions catalyzed by PAL and C4H does not appear to require channeling through a MEC in yeast, and 10 sheer biochemical coupling of PAL and C4H seems to be sufficient to drive carbon flux into the phenylpropanoid pathway. In yet another study (Hwang *et al.*, 2003) production of plant-specific flavanones by *Escherichia coli* was achieved through expression of an artificial gene 15 cluster that contained three genes of a phenyl propanoid pathway of various heterologous origins; PAL from the yeast *Rhodotorula rubra*, 4CL from the actinomycete *Streptomyces coelicolor*, and chalcone synthase (CHS) from the licorice plant *Glycyrrhiza echinata*. These pathways bypassed C4H, 20 because the bacterial 4CL enzyme ligated coenzyme A to both *trans*-cinnamic acid and 4-coumaric acid. In addition, the PAL from *Rhodotorula rubra* uses both phenylalanine and tyrosine as the substrates. Therefore, *E. coli* cells containing the gene clusters and grown on glucose, produced 25 small amounts of two flavanones, pinocembrin (0.29 g/l) from phenylalanine and naringenin (0.17 g/l) from tyrosine. In addition, large amounts of their precursors, 4-coumaric acid and *trans*-cinnamic acid (0.47 and 1.23 mg/liter respectively), were accumulated. Moreover, the yields of 30 these compounds could be increased by addition of phenylalanine and tyrosine.

Also described are studies in which the enzyme properties of pinosylvin synthases are studied by first cloning the genes into *Escherichia coli*. For instance, Raiber *et al.*, 1995 report on stilbenes from *Pinus strobus* (Eastern white pine) that were investigated after heterologous expression in *Escherichia coli*. For this a *P. strobus* cDNA library was screened with a stilbene synthase (STS) probe from *Pinus sylvestris* and amongst the isolated cDNAs two closely related STS genes, STS1 and STS2, were found with five amino acid differences in the proteins. The genes were cloned on a plasmid and expressed into *E. coli*, and cell extracts were subjected to enzyme assays. It appeared that both proteins accepted cinnamoyl-CoA as a substrate and thus were considered as pinosylvin synthases, however they revealed large differences. STS1 had only 3-5% of the activity of STS2, and its pH optimum was shifted to lower values (pH 6), and it synthesized with cinnamoyl-CoA a second unknown product. Site-directed mutagenesis demonstrated that a single Arg-to-His exchange in STS1 was responsible for all of the differences. In another study three STS cDNAs (PDSTS1, PDSTS2, and PDSTS3) from *Pinus densiflora* were isolated and the cDNAs were heterologously expressed in *E. coli* to characterize their enzymatic properties (Kodan *et al.*, 2002). PDSTS3 appeared to be an unusual STS isozyme that showed the highest pinosylvin-forming activity among the STSs tested. Furthermore, PDSTS3 was insensitive to product inhibition unlike PDSTS1 and PDSTS2. The unusual characteristics of PDSTS3 could be ascribed to a lack of a C-terminal extension that normally is common to stilbene synthases, which was caused by a frame-shift mutation. In yet another study a genomic DNA library was screened with pinosylvin synthase cDNA pSP-54 as a probe (Müller *et al.*, 1999). After subcloning, four different members were

characterized by sequencing. The amino acid sequences deduced from genes PST-1, PST-2, PST-3 and PST-5 shared an overall identity of more than 95%.

5 Differences in promoter strength were then analysed by transient expression in tobacco protoplasts. Constructs used contained the bacterial-glucuronidase under the control of the promoters of pine genes PST-1, PST-2 and PST-3. Upon treatment with UV light or fungal elicitor, the promoter of
10 PST-1 showed highest responsiveness and led to tissue-specific expression in vascular bundles. The data suggest that in pine the gene product of PST-1 is responsible for both the stress response in seedlings and pinosylvin formation in the heartwood.

15

A further study showed that a stilbene synthase cloned from Scots pine (*Pinus sylvestris*) was earlier abortively assigned as a dihydropinosylvin synthase, while it showed to be a pinosylvin synthase. The previous mis-interpretation
20 was caused by the influence of bacterial factors on the substrate preference and the activity of the plant-specific protein that was expressed in *E. coli*. After improvement of the expression system, the subsequent kinetic analysis revealed that cinnamoyl-CoA rather than phenylpropionyl-CoA
25 was the preferred substrate of the cloned stilbene synthase. Furthermore, extracts from *P. sylvestris* contained factor(s) that selectively influenced the substrate preference, i.e. the activity was reduced with phenylpropionyl-CoA, but not with cinnamoyl-CoA. This explained the apparent differences
30 between plant extracts and the cloned enzyme expressed in *E. coli* and cautions that factors in the natural and the new hosts may complicate the functional identification of cloned sequences.

Furthermore, vectors are described with stilbene synthase genes, which can mean resveratrol synthase and pinosylvin synthase, for the transformation of organisms and plants to confer enhanced resistance against pests and wounding (EP0309862 and EP0464461).

Also, further vectors are described that contain DNA sequences that will hybridise to pinosylvin synthase of *Pinus sylvestris* (US5391724) and said vectors to be used for expression in a plant (US5973230). The incorporation of PAL and 4CL together with a stilbene synthase for the production of pinosylvin in a organism is not however disclosed. Nor are any pinosylvin producing micro-organisms.

Recently, evidence was shown that the filamentous fungi *A. oryzae* contained the enzyme chalcone synthase (CHS) that is normally involved in the biosynthesis of flavonoids, such as naringenin, in plants (Juvvadi et al., 2005; Seshime et al., 2005). Indeed it was also shown that *A. oryzae* contained the major set of genes responsible for phenylpropanoid-flavonoid metabolism, i.e PAL, C4H and 4CL. However, there is no evidence that *A. oryzae* contains a stilbene synthase.

Our co-pending application WO2006/089898 describes resveratrol producing micro-organisms, especially yeasts.

SUMMARY OF THE INVENTION

The present invention now provides a micro-organism having an operative metabolic pathway comprising at least one enzyme activity producing pinosylvin from cinnamic acid. In preferred micro-organisms said pathway produces cinnamic

acid and produces pinosylvin therefrom. Especially, the invention provides the use of such micro-organisms in producing pinosylvin. Such a micro-organism may be naturally occurring and may be isolated by suitable screening
5 procedures such as degenerate PCR, Southern blotting and *in silico* homology searches, but more preferably is genetically engineered.

The invention includes methods of producing pinosylvin from such micro-organisms, and optionally isolating or purifying pinosylvin thereby produced. The culturing is preferably conducted in the substantial absence of an external source of cinnamic acid. This implies also, the substantial absence of an external source of derivatives of cinnamic
10 acid formed therefrom in the phenylpropanoid pathway such as cinnamoyl-CoA.
15

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

20 Preferably, said pinosylvin or derivative is produced in a reaction catalysed by an enzyme in which endogenous malonyl-CoA is a substrate, and preferably said pinosylvin is produced from cinnamoyl-CoA.

25 Said pinosylvin or derivative is preferably produced from cinnamoyl-CoA, preferably by a stilbene synthase synthase which preferably is expressed in said micro-organism from nucleic acid coding for said enzyme which is not native to the micro-organism.

30

Generally herein, unless the context implies otherwise, references to pinosylvin include reference to oligomeric or glycosidically bound derivatives thereof.

Thus, in certain preferred embodiments, said stilbene synthase is a resveratrol synthase (EC 2.3.1.95) from a plant belonging to the genus of *Arachis*, e.g. *A. glabata*,
5 *A. hypogaea*, a plant belonging to the genus of *Rheum*, e.g. *R. tataricum*, a plant belonging to the genus of *Vitus*, e.g. *V. labrusca*, *V. riparia*, *V. vinifera*, or any one of the genera *Pinus*, *Picea*, *Lilium*, *Eucalyptus*, *Parthenocissus*,
10 *Cissus*, *Calochortus*, *Polygonum*, *Gnetum*, *Artocarpus*, *Nothofagus*, *Phoenix*, *Festuca*, *Carex*, *Veratrum*, *Bauhinia* or *Pterolobium*.

The stilbene synthase may be one which exhibits a higher turnover rate with cinnamoyl-CoA as a substrate than it does
15 with 4-coumaroyl-CoA as a substrate, e.g. by a factor of at least 1.5 or at least 2. Thus, in further preferred embodiments, said stilbene synthase is a pinosylvin synthase, suitably from a tree species such as a species of *Pinus*, *Eucalyptus*, *Picea* or *Maclura*. In particular, the
20 stilbene synthase may be a pinosylvin synthase (EC 2.3.1.146) from a plant belonging to the genus of *Pinus*, e.g. *P. sylvestris*, *P. strobes*, *P. densiflora*, *P. taeda*, a plant belonging to the genus of *Picea*, or any one of the genus *Eucalyptus*.

25 Preferably, said cinnamic acid may be produced from L-phenylalanine in a reaction catalysed by an enzyme in which ammonia is produced and suitably said cinnamic acid is formed from L-phenylalanine by a phenylalanine ammonia
30 lyase.

In certain preferred embodiments, said L-phenylalanine ammonia lyase is a L-phenylalanine ammonia lyase (EC

4.3.1.5) from a plant or a micro-organism. The plant may belong to the genus of *Arabidopsis*, e.g. *A. thaliana*, a plant belonging to the genus of *Brassica*, e.g. *B. napus*, *B. rapa*, a plant belonging to the genus of *Citrus*, e.g. *C. reticulata*, *C. clementinus*, *C. limon*, a plant belonging to the genus of *Phaseolus*, e.g. *P. coccineus*, *P. vulgaris*, a plant belonging to the genus of *Pinus*, e.g. *P. banksiana*, *P. monticola*, *P. pinaster*, *P. sylvestris*, *P. taeda*, a plant belonging to the genus of *Populus*, e.g. *P. balsamifera*, *P. deltoides*, *P. Canadensis*, *P. kitakamiensis*, *P. tremuloides*, a plant belonging to the genus of *Solanum*, e.g. *S. tuberosum*, a plant belonging to the genus of *Prunus*, e.g. *P. avium*, *P. persica*, a plant belonging to the genus of *Vitus*, e.g. *Vitus vinifera*, a plant belonging to the genus of *Zea*, e.g. *Z. mays* or other plant genera e.g. *Agastache*, *Ananas*, *Asparagus*, *Bromheadia*, *Bambusa*, *Beta*, *Betula*, *Cucumis*, *Camellia*, *Capsicum*, *Cassia*, *Catharanthus*, *Cicer*, *Citrullus*, *Coffea*, *Cucurbita*, *Cynodon*, *Daucus*, *Dendrobium*, *Dianthus*, *Digitalis*, *Dioscorea*, *Eucalyptus*, *Gallus*, *Ginkgo*, *Glycine*, *Hordeum*, *Helianthus*, *Ipomoea*, *Lactuca*, *Lithospermum*, *Lotus*, *Lycopersicon*, *Medicago*, *Malus*, *Manihot*, *Medicago*, *Mesembryanthemum*, *Nicotiana*, *Olea*, *Oryza*, *Pisum*, *Persea*, *Petroselinum*, *Phalaenopsis*, *Phyllostachys*, *Physcomitrella*, *Picea*, *Pyrus*, *Quercus*, *Raphanus*, *Rehmannia*, *Rubus*, *Sorghum*, *Sphenostylis*, *Stellaria*, *Stylosanthes*, *Triticum*, *Trifolium*, *Triticum*, *Vaccinium*, *Vigna*, *Zinnia*. The micro-organism might be a fungus belonging to the genus *Agaricus*, e.g. *A. bisporus*, a fungus belonging to the genus *Aspergillus*, e.g. *A. oryzae*, *A. nidulans*, *A. fumigatus*, a fungus belonging to the genus *Ustilago*, e.g. *U. maydis*, a bacterium belonging to the genus *Rhodobacter*, e.g. *R. capsulatus*, a bacterium belonging to the genus *Streptomyces*, e.g. *S. maritimus*, a bacterium belonging to the genus *Photobacterium*, e.g. *P.*

luminescens, a yeast belonging to the genus *Rhodotorula*, e.g. *R. rubra*.

Because, as described above, for the production of
5 pinosylvin we require production of cinnamic acid by a PAL
enzyme and also its conversion on to pinosylvin rather than
either the production of coumaric acid from tyrosine by a
substrate promiscuous PAL or by conversion of cinnamic acid
10 by a C4H enzyme, micro-organisms for use in the invention
preferably have a PAL which favours phenylalanine as a
substrate (thus producing cinnamic acid) over tyrosine (from
which it would produce coumaric acid). Preferably,
therefore, the ratio $K_m(\text{phenylalanine})/K_m(\text{tyrosine})$ for the
PAL is less than 1:1, preferably less 1:5, e.g. less than
15 1:10. As usual, K_m is the molar concentration of the
substrate (phenylalanine or tyrosine respectively) that
produces half the maximal rate of product formation (V_{\max}).

The presence of C4H is not helpful to the production of
20 pinosylvin, but need not be forbidden provided that the
diversion of cinnamic acid away from pinosylvin production
toward formation of resveratrol via coumaric acid is not
excessive. Therefore, preferably C4H production is either
absent or such that $K_{\text{cat}}(\text{PAL})/K_{\text{cat}}(\text{C4H})$ is greater than 2,
25 preferably greater than 4. As usual, in each case, K_{cat} is
 $V_{\max}/[\text{Enzyme}]$, where [Enzyme] is the concentration of the
relevant enzyme.

By way of illustration, typical K_m values for *A. thaliana*
30 phenylalanine ammonia lyase PAL2 and its homologue PAL1 are
around 60 μM with phenylalanine as substrate (Cochrane et
al, 2004) and more than 1000 μM when using tyrosine as
substrate (Watts et al, 2006). The catalytic turnover rate

K_{cat} for *A. thaliana* PAL2 is 192 mol cinnamic acid/mole enzyme PAL2 when converting phenylalanine to cinnamic acid (Cochrane et al, 2004) but K_{cat} is minute for the conversion of tyrosine to coumaric acid. A PAL with the above kinetic
5 properties is specific for phenylalanine as substrate and gives exclusively cinnamic acid formation from phenylalanine and undetectable levels of coumaric acid from tyrosine.

The typical turnover rate for the hydroxylase reaction
10 catalyzed by C4H is 25 moles coumaric acid product/mole enzyme/minute when native yeast CPR activity supports the reaction (Urban et al, 1994). The activity of C4H may be limited by NADPH availability and this may be increased if the enzyme cytochrome P450 hydroxylase (CPR) is
15 overexpressed. If CPR is overexpressed as exemplified in the literature by 5 to 20 times (Mizutani et al, 1998, Urban et al, 1994) the catalytic turnover rates for the C4H reaction converting cinnamic acid to coumaric acid increases to 125 mole coumaric acid product/mole enzyme/minute and 530
20 mole coumaric acid product/mole enzyme/minute, respectively.

The outcome of the combined reaction PAL-C4H-CPR will depend on the catalytic numbers and the amount of each enzyme present, especially the amount of CPR supporting the
25 electron donation, NADPH, for the C4H. An efficient PAL will give ca 192 moles cinnamic acid/mole PAL/minute and the C4H enzyme following in the sequence will convert ca 25 moles of this cinnamic acid/mole C4H/minute into coumaric acid with native CPR activity. Thus the dominant product
30 from the combined reaction PAL-C4H-CPR will be cinnamic acid (167 moles cinnamic acid/mole PAL enzyme/minute and 25 moles coumaric acid/mole enzyme C4H/minute with native CPR activity. Higher CPR activity will lead to more C4H

activity per mole C4H enzyme and ultimately to pure coumaric acid if overexpressed at high levels. A CPR overexpressed only five times as in the Mizutani paper (Mizutani et al, 1998) would result in 125 moles coumaric acid/mole

5 C4H/minute and only 67 moles cinnamic acid would be the result from the PAL per minute. Thus the CPR must at least be overexpressed ca 8 times for (undesired) pure coumaric acid production.

10 In the case of a recombinant or natural organism with several PALs/TALs and C4H one can prepare a cell extract and measure the apparent catalytic turnover rates and K_m values as a sum total (or aggregated enzyme) apparent enzyme PAL, TAL or C4H. From these estimated sum properties it will be

15 possible to determine if the organism will produce mainly coumaric acid or cinnamic acid and thus which product resveratrol or pinosylvin would be the outcome when 4CL and VST are expressed in this organism. The turnover rate will now be expressed as moles product / (mole total protein/

20 time) instead of when using pure enzymes moles product/(mol pure enzyme/time). Therefore, the preferred ratio $K_m(\text{phenylalanine})/K_m(\text{tyrosine})$ for the PAL less than 1:1 can be applied to the aggregate PAL activity where more than one PAL is present and the preferred ratio $K_{cat}(\text{PAL})/K_{cat}(\text{C4H})$

25 greater than 2 can be applied to the aggregate of the PAL and/or C4H activity (as modulated by CPR) where more than one PAL and/or C4H activity is present.

Preferably, the micro-organism has no exogenous C4H, i.e.

30 has not been genetically modified to provide expression of a C4H enzyme. Any C4H production there may then be will be native to the organism. Optionally, the micro-organism without exogenous C4H may also lack endogenous C4H. Lack of

endogenous C4H may be due to a native C4H capability having been deleted by genetic engineering or gene silencing methods or simply because the organism naturally lacks the C4H genes, since the enzyme is not part of its metabolism.

5

Also, as seen above, the presence of CPR is not helpful to the production of pinosylvin and its overexpression, while not forbidden is not generally desirable. Accordingly, the micro-organism preferably has no endogenous CPR, no

10 exogenous CPR or has no overexpression of native CPR, or may have reduced expression of native CPR.

Suitably, said L-phenylalanine ammonia lyase is expressed in said micro-organism from nucleic acid coding for said enzyme which is not native to the micro-organism.

15

Preferably, cinnamoyl-CoA is formed in a reaction catalysed by an enzyme in which ATP and CoA are substrates and ADP is a product and suitably cinnamoyl-CoA is formed in a reaction catalysed by a 4-coumarate-CoA ligase (also referred to as 4-coumaroyl-CoA ligase). Known 4-coumarate-CoA ligase enzymes accept either 4-coumaric acid or cinnamic acid as substrates and produce the corresponding CoA derivatives. Generally, such enzymes are known as '4-coumarate-CoA ligase' whether they show higher activity with 4-coumaric acid as substrate or with cinnamic acid as substrate.

20

25

However, we refer here to enzymes having that substrate preference as 'cinnamate-CoA ligase' enzymes (or cinnamoyl-CoA-ligase). One such enzyme is described for instance in Aneko *et al.*, 2003.

30

Said 4-coumarate-CoA ligase or cinnamate-CoA ligase may be a 4-coumarate-CoA ligase / cinnamate-CoA ligase (EC 6.2.1.12)

from a plant, a micro-organism or a nematode. The plant may belong to the genus of *Abies*, e.g. *A. beshanzuensis*, *B. firma*, *B. holophylla*, a plant belonging to the genus of *Arabidopsis*, e.g. *A. thaliana*, a plant belonging to the

5 genus of *Brassica*, e.g. *B. napus*, *B. rapa*, *B. oleracea*, a plant belonging to the genus of *Citrus*, e.g. *C. sinensis*, a plant belonging to the genus of *Larix*, e.g. *L. decidua*, *L. gmelinii*, *L. griffithiana*, *L. himalaica*, *L. kaempferi*, *L. laricina*, *L. mastersiana*, *L. occidentalis*, *L. potaninii*, *L.*

10 *sibirica*, *L. speciosa*, a plant belonging to the genus of *Phaseolus*, e.g. *P. acutifolius*, *P. coccineus*, a plant belonging to the genus of *Pinus*, e.g. *P. armandii* *P. banksiana*, *P. pinaster*, a plant belonging to the genus of *Populus*, e.g. *P. balsamifera*, *P. tomentosa*, *P. tremuloides*,

15 a plant belonging to the genus of *Solanum*, e.g. *S. tuberosum*, a plant belonging to the genus of *Vitis*, e.g. *Vitis vinifera*, a plant belonging to the genus of *Zea*, e.g. *Z. mays*, or other plant genera e.g. *Agastache*, *Amorpha*, *Cathaya*, *Cedrus*, *Crocus*, *Festuca*, *Glycine*, *Juglans*,

20 *Keteleeria*, *Lithospermum*, *Lolium*, *Lotus*, *Lycopersicon*, *Malus*, *Medicago*, *Mesembryanthemum*, *Nicotiana*, *Nothotsuga*, *Oryza*, *Pelargonium*, *Petroselinum*, *Physcomitrella*, *Picea*, *Prunus*, *Pseudolarix*, *Pseudotsuga*, *Rosa*, *Rubus*, *Ryza*, *Saccharum*, *Suaeda*, *Thellungiella*, *Triticum*, *Tsuga*. The

25 micro-organism might be a filamentous fungi belonging to the genus *Aspergillus*, e.g. *A. flavus*, *A. nidulans*, *A. oryzae*, *A. fumigatus*, a filamentous fungus belonging to the genus *Neurospora*, e.g. *N. crassa*, a fungus belonging to the genus *Yarrowia*, e.g. *Y. lipolytica*, a fungus belonging to the

30 genus of *Mycosphaerella*, e.g. *M. graminicola*, a bacterium belonging to the genus of *Mycobacterium*, e.g. *M. bovis*, *M. leprae*, *M. tuberculosis*, a bacterium belonging to the genus of *Neisseria*, e.g. *N. meningitidis*, a bacterium belonging to

the genus of *Streptomyces*, e.g. *S. coelicolor*, a bacterium
belonging to the genus of *Rhodobacter*, e.g. *R. capsulatus*, a
nematode belonging to the genus *Ancylostoma*, e.g. *A.*
ceylanicum, a nematode belonging to the genus
5 *Caenorhabditis*, e.g. *C. elegans*, a nematode belonging to the
genus *Haemonchus*, e.g. *H. contortus*, a nematode belonging to
the genus *Lumbricus*, e.g. *L. rubellus*, a nematode belonging
to the genus *Meilodogyne*, e.g. *M. hapla*, a nematode
belonging to the genus *Strongyloidus*, e.g. *S. ratti*, *S.*
10 *stercoralis*, a nematode belonging to the genus *Pristionchus*,
e.g. *P. pacificus*.

Whilst the micro-organism may be naturally occurring,
preferably at least one copy of at least one genetic
15 sequence encoding a respective enzyme in said metabolic
pathway has been recombinantly introduced into said micro-
organism.

Additionally or alternatively to introducing coding
20 sequences coding for a said enzyme, one may provide one or
more expression signals, such as promoter sequences, not
natively associated with said coding sequence in said
organism. Thus, optionally, at least one copy of a genetic
sequence encoding a L-phenylalanine ammonia lyase is
25 operatively linked to an expression signal not natively
associated with said genetic sequence in said organism.

Expression signals include nucleotide sequences located
upstream (5' non-coding sequences), within, or downstream
30 (3' non-coding sequences) of a coding sequence, and which
influence the transcription, RNA processing or stability, or
translation of the associated coding sequence. Such
sequences may include promoters, translation leader

sequences, introns, and polyadenylation recognition sequences.

5 Optionally, at least one copy of a genetic sequence encoding a 4-coumarate-CoA ligase or cinnamate-CoA ligase, whether native or not, is operatively linked to an expression signal not natively associated with said genetic sequence in said organism.

10 Optionally, at least one copy of a genetic sequence encoding a stilbene synthase, which may be a resveratrol synthase, whether native or not, is operatively linked to an expression signal not natively associated with said genetic sequence in said organism.

15 Optionally, at least one copy of a genetic sequence encoding a pinosylvin synthase, whether native or not, is operatively linked to an expression signal not natively associated with said genetic sequence in said organism.

20 In certain aspects the invention provides a metabolically engineered micro-organism of the kind described, having an operative metabolic pathway in which a first metabolite is transformed into a second metabolite in a reaction catalysed
25 by a first enzyme, said reaction step producing ammonia, and in which said second metabolite is transformed into a third metabolite in a reaction catalysed by a second enzyme in which ATP and CoA is a substrate, and ADP is a product, and in which said third metabolite is transformed into a fourth
30 metabolite in a reaction catalysed by a third enzyme in which endogenous malonyl-CoA is a substrate.

The micro-organisms described above include ones containing one or more copies of a heterologous DNA sequence encoding phenylalanine ammonia lyase operatively associated with an expression signal, and containing one or more copies of a heterologous DNA sequence encoding 4-coumarate-CoA-ligase or cinnamate-CoA ligase operatively associated with an expression signal, and containing one or more copies of a heterologous DNA sequence encoding a stilbene synthase, which may be resveratrol synthase, operatively associated with an expression signal.

Alternatively, the micro-organisms described above include ones containing one or more copies of a heterologous DNA sequence encoding phenylalanine ammonia lyase operatively associated with an expression signal, and containing one or more copies of a heterologous DNA sequence encoding 4-coumarate-CoA-ligase or cinnamate-CoA ligase operatively associated with an expression signal, and containing one or more copies of a heterologous DNA sequence encoding pinosylvin synthase operatively associated with an expression signal.

In the present context the term "micro-organism" relates to microscopic organisms, including bacteria, microscopic fungi, including yeast.

More specifically, the micro-organism may be a fungus, and more specifically a filamentous fungus belonging to the genus of *Aspergillus*, e.g. *A. niger*, *A. awamori*, *A. oryzae*, *A. nidulans*, a yeast belonging to the genus of *Saccharomyces*, e.g. *S. cerevisiae*, *S. kluyveri*, *S. bayanus*, *S. exiguus*, *S. sevazzi*, *S. uvarum*, a yeast belonging to the genus *Kluyveromyces*, e.g. *K. lactis* *K. marxianus* var.

marxianus, *K. thermotolerans*, a yeast belonging to the genus *Candida*, e.g. *C. utilis*, *C. tropicalis*, *C. albicans*, *C. lipolytica*, *C. versatilis*, a yeast belonging to the genus *Pichia*, e.g. *P. stipidis*, *P. pastoris*, *P. sorbitophila*, or
5 other yeast genera, e.g. *Cryptococcus*, *Debaromyces*,
Hansenula, *Pichia*, *Yarrowia*, *Zygosaccharomyces* or
Schizosaccharomyces. Concerning other micro-organisms a non-
exhaustive list of suitable filamentous fungi is supplied: a
species belonging to the genus *Penicillium*, *Rhizopus*,
10 *Fusarium*, *Fusidium*, *Gibberella*, *Mucor*, *Mortierella*,
Trichoderma.

Concerning bacteria a non-exhaustive list of suitable
bacteria is given as follows: a species belonging to the
15 genus *Bacillus*, a species belonging to the genus
Escherichia, a species belonging to the genus *Lactobacillus*,
a species belonging to the genus *Lactococcus*, a species
belonging to the genus *Corynebacterium*, a species belonging
to the genus *Acetobacter*, a species belonging to the genus
20 *Acinetobacter*, a species belonging to the genus *Pseudomonas*,
etc.

The preferred micro-organisms of the invention may be *S.*
cerevisiae, *A. niger*, *A. oryzae*, *E. coli*, *L. lactis* or *B.*
25 *subtilis*.

The constructed and engineered micro-organism can be
cultivated using commonly known processes, including
chemostat, batch, fed-batch cultivations, etc.

30

Thus, the invention includes a method for producing
pinosylvin comprising contacting a micro-organism cell with
a carbon substrate in the substantial absence of an external

source of cinnamic acid, said cell having the capacity to produce pinosylvin under the conditions, in which the micro-organism may be selected from the group consisting of fungi and bacteria, especially yeast.

5

Pinosylvin so produced may optionally be isolated or purified and suitable methods include solvent extraction with n-hexane, followed by sequential extraction with 100% ether, acetone, methanol and water, and chromatographic
10 purification on a silicagel column using a n-hexane/ethyl acetate (2/1) system (Suga *et al.* 1993).

Said carbon substrate is optionally selected from the group of fermentable carbon substrates consisting of
15 monosaccharides, oligosaccharides and polysaccharides, e.g. glucose, fructose, galactose, xylose, arabinose, mannose, sucrose, lactose, erythrose, threose, and/or ribose. Said carbon substrate may additionally or alternatively be selected from the group of non-fermentable carbon substrates
20 including ethanol, acetate, glycerol, and/or lactate. Said non-fermentable carbon substrate may additionally or alternatively be selected from the group of amino acids and may be phenylalanine.

25 In an alternative aspect, the invention includes a method for producing pinosylvin through heterologous expression of nucleotide sequences encoding phenylalanine ammonia lyase, 4-coumarate-CoA ligase and resveratrol synthase and also a method for producing pinosylvin through heterologous
30 expression of nucleotide sequences encoding phenylalanine ammonia lyase, 4-coumarate-CoA ligase and pinosylvin synthase.

Pinosylvin, including pinosylvin so produced, may be used as a nutraceutical in a food product, e.g. a dairy product or a beverage such as beer or wine. Accordingly, the invention includes a food product containing microbially produced
5 pinosylvin.

The invention further includes a micro-organism composition comprising micro-organism cells and at least 1.5 mg/g pinosylvin on a dry weight basis. For instance, yeast or
10 yeast containing or yeast derived preparations containing pinosylvin, or pinosylvin so produced, may be provided for human or animal consumption, e.g. in dry form, suitably as unit oral dosage forms such as yeast containing tablets or capsules, which may contain for instance at least 0.5g of
15 said yeast, e.g. 1-3g.

Any wild type enzyme referred to herein may be substituted by a mutant form thereof, suitably having an amino acid homology relative to the named wild type enzyme of at least
20 50%, more preferably at least 60%, more preferably at least 70%, more preferably at least 80%, more preferably still at least 90% or at least 95%, whilst of course maintaining the required enzyme activity of the wild type. This may include maintaining any substrate preference of the wild type, e.g.
25 for phenylalanine over tyrosine or for cinnamic acid over coumaric acid or for cinnamoyl-CoA over coumaroyl-CoA. Any wild type coding sequence coding for an enzyme referred to herein may be substituted with a sequence coding for the same enzyme but in which the codon usage is adjusted. This
30 applies both to wild type enzymes mentioned herein and mutant forms as discussed above. Nucleotide sequences coding for mutant forms of wild type enzymes are preferably homologous with the wild type nucleotide sequence of the

corresponding wild type enzyme to the extent of at least 50%, more preferably at least 60%, more preferably at least 70%, more preferably at least 80%, more preferably still at least 90% or at least 95%.

5

Mutant forms of enzymes may have a level of enzyme activity largely unchanged from that of the wild type enzyme or may be selected to have a higher level of activity.

Conservative substitutions of amino acids of the wild type enzyme may be made in accordance with known practice.

Enzymes having improved activity may be developed by directed evolution techniques as known in the art, random changes in the enzyme being produced by methods such as introducing random genetic changes in the coding for the enzyme in a suitable test organism such as *E.coli* or *S. cerevisiae* followed by expression and selection of improved mutants by screening for the desired property, or by imposing self selection conditions under which organisms expressing an improved activity will have a survival advantage.

20

References herein to the absence or substantial absence or lack of supply of a substance, e.g. of cinnamic acid, include the substantial absence of derivatives thereof such as cinnamic acid esters (including thioesters), e.g. cinnamoyl-CoA, which may be metabolised to the substance or which are immediate products of further metabolism of the substance. In particular, lack of cinnamic acid implies lack of cinnamoyl-CoA.

25
30

Pinosylvin produced according to the invention may be *cis*-pinosylvin or *trans*-pinosylvin, which are expected to be formed from *cis*-cinnamic acid and *trans*-cinnamic acid

respectively. Alternatively, *cis*-pinosylvin may be formed from *trans*-cinnamic acid by a process including isomerisation. But it is to be expected that the *trans*-form will normally predominate.

5

BRIEF DESCRIPTION OF THE DRAWINGS

To assist in the ready understanding of the above description of the invention reference has been made to the accompanying drawings in which:

10

Figure 1 shows the chemical structure of pinosylvin;

15

Figure 2 shows the phenylpropanoid pathway utilising resveratrol synthase acting on coumaroyl-CoA, leading to resveratrol; and

20

Figure 3 shows the phenylpropanoid pathway utilising pinosylvin synthase or resveratrol synthase acting on cinnamoyl-CoA, leading to pinosylvin.

25

Figure 4 shows the HPLC-chromatograms of supernatant and cell extract of *S. cerevisiae* strains FSSC-PAL4CLVST1, grown on 100 g/l galactose. A chromatogram of 60 nanogram of pure pinosylvin is included.

30

Figure 5 shows the HPLC-chromatograms of a cell extract of *S. cerevisiae* strain FSSC-PAL4CLRES, grown on 100 g/l galactose. A chromatogram of 60 nanogram of pure pinosylvin is included.

Figure 6 shows the LC-MS data for pure pinosylvin and pinosylvin produced by *S. cerevisiae* strain FSSC-PAL4CLVST1, grown on 100 g/l galactose. Both base peak chromatograms, and negative ion-traces at M/Z 211.0759 Da/e are shown.

5

Figure 7 shows HPLC chromatograms obtained in Example 16.

Figure 8 shows the HPLC analysis of extracted product from the fermentation of a pinosylvin producing strain of *E. coli* (upper panel) and a control strain (lower panel).

10

The invention will be further described and illustrated by the following non-limiting examples.

EXAMPLES

15 Example 1

Isolation of genes encoding PAL, 4CL, RES and VST1

Phenylalanine ammonia lyase (PAL2) (Cochrane *et al.*, 2004; SEQ ID NO: 1, 2), 4-coumarate:CoenzymeA ligase (4CL1) (Hamberger and Hahlbrock 2004; Ehling *et al.*, 1999; SEQ ID NO: 3, 4) were isolated via PCR from *A. thaliana* cDNA (BioCat, Heidelberg, Germany) using the primers in table 1. PAL2 and 4CL1 were chosen amongst several *A. thaliana* homologues due to favourable kinetic parameters towards cinnamic acid and cinnamoyl-CoA, respectively (Cochrane *et al.*, 2004; Hamberger and Hahlbrock 2004; Ehling *et al.*, 1999).

25

The coding sequence of resveratrol synthase (RES) from Rhubarb, *Rheum tataricum* (Samappito *et al.*, 2003; SEQ ID NO: 5, 6) was codon optimized for expression in *S. cerevisiae* using the online service backtranslation tool at

30

www.entelechon.com, yielding sequence SEQ ID NO: 7, 8.
 Oligos for the synthetic gene assembly were constructed at
 MWG Biotech and the synthetic gene was assembled by PCR
 using a slightly modified method protocol of from Martin *et*
 5 *al.* (2003) described below.

Table 1. Primers and restriction sites for the amplification of genes			
Primer for amplification of gene* (Restriction sites are underlined)	Gene	Restriction site: primer	Restriction site: vector
5'- <u>CGGAATTCT</u> CATGGATCAAATCGAAGCAATGTT	PAL2	EcoR1	EcoR1
5'-CGACTAGTTT <u>AGCAAATCGGAATCGGAGC</u>	PAL2	Spe1	Spe1
5'-GCTCTAGACCT ATGGCGCCACAAGAACAAGCAGTTT	4CL1	Xba1	Spe1
5'-GCGGATCCCCT TCACAATCCATTTGCTAGTTT TGCC	4CL1	BamH1	BglII
5'-CC <u>GGATCC</u> AAATGGCCCCAGAAGAGAGCAGG	RES	BamH1	BamH1
5'-CG <u>CTCGAGT</u> TAAAGTGATCAATGGAACCGAAGACAG	RES	Xho1	Xho1

* SEQ ID Nos 11-16

Primers from MWG for the assembly of the synthetic gene were
 10 dissolved in milliQ-water to a concentration of 100
 pmole/ μ l. An aliquot of 5 μ l of each primer was combined in
 a totalmix and then diluted 10-fold with milliQ water. The
 gene was assembled via PCR using 5 μ l diluted totalmix per
 50 μ l as template for fusion DNA polymerase (Finnzymes). The
 15 PCR programme was as follows: Initial 98 °C for 30 s., and
 then 30 cycles with 98 °C for 10 s., 40 °C for 1 min. and 72
 °C at 1 min./1000 basepairs, and a final 72 °C for 5 min.
 From the resulting PCR reaction, 20 μ l was purified on 1%
 agarose gel. The result was a PCR smear and the regions
 20 around the wanted size were cut out from agarose gel and
 purified using the QiaQuick Gel Extraction Kit (Qiagen). A
 final PCR with the outer primers in table 1 rendered the
 required RES gene. Point mutations were corrected using the

Quickchange site directed mutagenesis II kit (Stratagene, La Jolla, CA).

The VST1 gene encoding *Vitis vinifera* (grapevine) resveratrol synthase (Hain *et al.*, 1993) was synthesized by GenScript Corporation (Piscataway, NJ). The amino acid sequence (SEQ ID NO: 10) was used as template to generate a synthetic gene codon optimized for expression in *S. cerevisiae* (SEQ ID NO: 9). The synthetic VST1 gene was delivered inserted in *E. coli* pUC57 vector flanked by BamH1 and Xho1 restriction sites. The synthetic gene was purified from the pUC57 vector by BamH1/Xho1 restriction and purified from agarose gel using the QiaQuick Gel Extraction Kit (Qiagen).

15

Example 2

Construction of a yeast vector for expression of PAL2

The gene encoding PAL2, isolated as described in example 1, was reamplified by PCR using forward- and reverse primers, with 5' overhangs containing EcoR1 and Spe1 restriction sites (table 1). The amplified PAL2 PCR product was digested with EcoR1/Spe1 and ligated into EcoR1/Spe1 digested pESC-URA vector (Stratagene), resulting in vector pESC-URA-PAL2. The sequence of the gene was verified by sequencing of two different clones.

25

Example 3

Construction of a yeast vector for expression of 4CL1

30

The gene encoding 4CL1 was isolated as described in example 1. The amplified 4CL1 PCR-product was digested with

Xba1/BamH1 and ligated into Spe1/BglII digested pESC-TRP vector (Stratagene), resulting in vector pESC-TRP-4CL1. Two different clones of pESC-TRP-4CL1 were sequenced to verify the sequence of the cloned gene.

5

Example 4

Construction of a yeast vector for expression of 4CL1 and RES

10 The gene encoding RES was isolated as described in example 1. The amplified synthetic RES gene was digested with BamH1/Xho1 and ligated into BamH1/Xho1 digested pESC-TRP-4CL1 (example 3). The resulting plasmid, pESC-TRP-4CL1-RES, contained the genes encoding 4CL1 and RES under the control
15 of the divergent GAL1/GAL10 promoter. The sequence of the gene encoding VST1 was verified by sequencing of two different clones of pESC-TRP-4CL1-VST1.

Example 5

20 *Construction of a yeast vector for expression of 4CL1 and VST1*

The gene encoding VST1 was isolated as described in example 1. The purified and digested VST1 gene was ligated into
25 BamH1/Xho1 digested pESC-TRP-4CL1 (example 3). The resulting plasmid, pESC-TRP-4CL1-VST1, contained the genes encoding 4CL1 and VST1 under the control of the divergent GAL1/GAL10 promoter. The sequence of the gene encoding VST1 was verified by sequencing of two different clones of pESC-TRP-
30 4CL1-VST1.

Example 6

Expression of the pathway to pinosylvin in the yeast S. cerevisiae using PAL2, 4CL1 and RES

5

Yeast strains containing the appropriate genetic markers were transformed with the vectors described in examples 2,3 and 4, separately or in combination. The transformation of the yeast cell was conducted in accordance with methods known in the art by using competent cells, an alternative being for instance, electroporation (see, e.g., Sambrook et al., 1989). Transformants were selected on medium lacking uracil and/or tryptophan and streak purified on the same medium.

15

S. cerevisiae strain FS01267 (MATa trp1 ura3) was co-transformed with pESC-URA-PAL2 (example 2) and pESC-TRP-4CL1-RES (example 4), and the transformed strain was named FSSC-PAL24CL1RES.

20

Example 7

Expression of the pathway to pinosylvin in the yeast S. cerevisiae using PAL2, 4CL1 and VST1

25

Yeast strains containing the appropriate genetic markers were transformed with the vectors described in examples 2,3 and 5, separately or in combination. The transformation of the yeast cell was conducted in accordance with methods known in the art, for instance, by using competent cells or by electroporation (see, e.g., Sambrook et al., 1989).

30

Transformants were selected on medium lacking uracil and/or tryptophan and streak purified on the same medium.

5 *S. cerevisiae* strain FS01267 (MATa trp1 ura3) was co-transformed with pESC-URA-PAL2 (example 2) and pESC-TRP-4CL1-VST1 (example 5), and the transformed strain was named FSSC-PAL24CL1VST1.

Example 8

10 *Fermentation with recombinant yeast strains in shake flasks*

The recombinant yeast strains were inoculated from agar plates with a sterile inoculation loop and grown in 100 ml defined mineral medium (Verduyn *et al.*, 1992) that contained 15 vitamins, trace elements, 5 g/l glucose 95 g/l galactose. The 500 ml stoppered shake flasks were incubated for three days at 30 °C and 160 rpm.

Example 9

20 a) *Extraction of pinosylvin*

Cells were harvested by centrifugation 5000 g for 5 minutes. An aliquot of 50 ml of supernatant was extracted once with 20 ml ethyl acetate. The ethyl acetate was freeze dried and 25 the dry product redissolved in 0.7 ml methanol and filtered into HPLC vials.

The cell pellet from 100 ml medium was dissolved in 2 ml water and divided into 3 fastprep tubes and broken with glass beads. The crude extracts from the three tubes were 30 pooled into 10 ml 100 % methanol in a 50 ml sartorius tube and extracted on a rotary chamber for 48 hours in a dark

cold room at 4 °C. After 48 hours the cell debris was removed via centrifugation for 5 min. at 5000 g and the methanol was removed by freeze-drying overnight. The dry residue was redissolved in 0.7 ml methanol and filtered into HPLC vials.

5

b) Analysis of pinosylvin

HPLC

10 For quantitative analysis of cinnamic acid, coumaric acid, and pinosylvin, samples were subjected to separation by high-performance liquid chromatography (HPLC) Agilent Series 1100 system (Hewlett Packard) prior to uv-diode-array detection at $\lambda = 306$ nm. A Phenomenex (Torrance, CA, USA) Luna 3 micrometer C18 (100 X 2.00 mm) column was used at 40 15 °C. As mobile phase a gradient of acetonitrile and milliq water (both containing 50 ppm trifluoroacetic acid) was used at a flow of 0.4 ml/min. The gradient profile was linear from 15 % acetonitrile to 100 % acetonitrile over 20 min. 20 The elution time was approximately 8.8-8.9 minutes for trans-pinosylvin. Pure pinosylvin standard (> 95% pure) was purchased from ArboNova (Turku, Finland).

LC-MS

25

Samples and standards were analyzed by negative electrospray LC-MS on a Waters (Micromass, Manchester, UK) LCT™ time-of-flight mass spectrometer with a Lockspray™ reference probe coupled to an Agilent 1100 HPLC system (Agilent 30 Technologies Walbron, Germany). The separations were done on a 50 mm x 2 mm ID Luna C-18 (II) column (Phenomenex, USA) fitted with a 4mm x 2 mm ID SecurityGuard™ pre-column

(Phenomenex, USA) using a water - acetonitrile gradient at 0.3 ml/minute. Both eluents contained 20 mM formic acid. The solvent composition was changed from 15% acetonitrile at injection to 100% acetonitrile in 20 minutes, which was

5 maintained for 5 minutes before the gradient was returned to starting conditions. A 3 μ l sample was injected in all cases and the column was maintained at 40 °C. All chemicals were of HPLC grade and dissolved into Milli-Q™ water.

10 UV spectra were collected from 200-700 nm at 2 spectra per second with a resolution of 4 nm.

The mass spectrometer was tuned for maximum sensitivity in negative electrospray mode to a resolution better than 5500 FWH on a solution of leucine enkephaline (0.5 μ g/ml in 50% acetonitril with 0.5% formic acid). Said solution was also

15 used as mass reference in the Lockspray™ in negative ESI at 15 μ l/minute. The instrument was calibrated in negative ESI on a carboxylated-PEG mixture in 50% acetonitril. In both cases the calibration had a residual error less than 2 mDa on at least 25 calibration ions. The run conditions were

20 selected for minimal in-source fragmentation.

Mass spectra were collected from 100 to 900 Da/e at a rate of 0.4 seconds per spectrum with 0.1 second interscan time. A reference spectrum was collected from the Lockmass™ probe every 3rd seconds and 10 reference spectra were averaged for

25 internal mass correction.

Narrow ion traces were extracted using +/- 25 mDa around the protonated or deprotonated mass of the expected metabolites.

Results

30 Strains FSSC-PAL24CL1RES and FSSC-PAL24CL1VST1, were cultivated on 100 g/l galactose as described in example 8,

and analyzed for their content of pinosylvin. Additionally, a control strain FSSC-control was included that contained the empty vectors only. The HPLC-analysis showed that strains FSSC-PAL24CL1VST1 and FSSC-PAL24CL1RES contained a component with a retention time of 8.8-9.0 min. that was identical to *trans*-pinosylvin (figure 4 and 5). Said result was confirmed by LC-MS analysis that revealed the presence of a component in the supernatant of strain FSSC-PAL24CL1VST1 with a retention time of 8.2 min., which had a M/Z of 211.0579 Da/e \pm 25 mDA that indeed corresponded to the M/Z of pure pinosylvin in negative ion mode (figure 6). In addition the UV absorption spectra were similar to the absorption spectrum of pure *trans*-pinosylvin (not shown) as well, with a λ_{\max} of approximately 306 nm.

The results, therefore, demonstrated the presence of an active phenyl-propanoid pathway in *S. cerevisiae* that led to *in vivo* production of *trans*-pinosylvin. The production of pinosylvin can most likely be improved by cultivating the strains under well-defined growth conditions in batch- and continuous cultures, and/or optimizing the expression/activities of the individual enzymes

Example 10

a) *Construction of a bacterial vector for expression of PAL2 in Escherichia coli.*

The plasmids that were used in the following examples contained one or more marker genes to allow the microorganism that harbour them to be selected from those which do not. The selection system is based upon dominant

markers, e.g. resistance against ampicilin and kanamycin. In addition, the plasmids contained promoter- and terminator sequences that allowed the expression of the recombinant genes. Furthermore, the plasmids contained suitable unique restriction sites to facilitate the cloning of DNA fragments and subsequent identification of recombinants. In this example the plasmids contained either the ampicilin resistance gene, designated as pET16b (Novagen), or the kanamycin resistance gene, designated as pET26b (Novagen).

The gene encoding PAL2, isolated as described in example 1, was reamplified by PCR from the plasmid pESC-URA-PAL2 (example 2), using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the gene allowed ligation of the restricted PCR product into a digested pET16B vector that contained the T7 promoter. The resulting plasmid, pET16B-PAL2, contained the gene encoding PAL2 under the control of the T7 promoter.

b) Construction of a bacterial vector for expression of 4CL1 and VST1 in *Escherichia coli*.

The gene encoding 4CL1, isolated as described in example 1, was reamplified by PCR from the plasmid pESC-URA-4CL1-VST1 (example 5), using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the gene allowed ligation of the restricted PCR product into a digested pET26B vector. The resulting plasmid, pET26B-4CL1, contained the gene encoding for 4CL1 under the control of the T7 promoter from *Lactobacillus lactis*.

The gene encoding VST1, isolated as described in example 1, was reamplified by PCR from the plasmid pESC-URA-4CL1-VST1 (example 5) using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The
5 introduction of said restriction sites at the 5' and 3' ends of the gene allowed ligation of the restricted PCR product into a digested pET16B vector. The resulting plasmid, pET16B-VST1, contained the gene encoding VST1 under the control of the T7 promoter. The T7 promoter and the gene
10 encoding VST1 were reamplified as one fragment by PCR from the plasmid pET16B-VST1 using forward and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the DNA fragment allowed ligation of the restricted
15 PCR product into the digested plasmid pET26B-4CL1. The resulting plasmid, pET26B-4CL1-VST1, contained the genes encoding 4CL1 and VST1, each under the control of their individual T7 promoter. The sequence of the genes encoding 4CL1 and VST1 was verified by sequencing of two different
20 clones of pET26B-4CL1-VST1.

c) Expression of the pathway to pinosylvin in *Escherichia coli*

25 *Escherichia coli* strains were transformed with the vectors described in (a) and (b), separately or in combination. The transformation of the bacterial cell was conducted in accordance with methods known in the art by using competent cells, an alternative being for instance, electroporation
30 (see, e.g., Sambrook *et al.*, 1989). Transformants were selected on medium containing the antibiotics ampicilin and kanamycin and streak purified on the same medium.

Escherichia coli strain BL21 (DE3) was transformed separately with the vector pET16B-PAL2 (a), yielding the strain FSEC-PAL2; and with pET26B-4CL1-VST1 (b), yielding strain FSEC-4CL1VST1. In addition, *Escherichia coli* strain BL21 (DE3) was co-transformed with pET16B-PAL2 (a) and pET26B-4CL1-VST1 (n), and the transformed strain was named FSEC-PAL24CL1VST1.

d) Fermentation with recombinant *Escherichia coli* strains in fermentors.

The recombinant yeast strains can be grown in fermentors operated as batch, fed-batch or chemostat cultures. In this instance fermentation was in shake flasks.

15

Pre-cultures of *Escherichia coli* BL21 (DE3) were grown in glass tubes at 160 rpm and 37 °C in 7 ml of LB medium containing 100 µg/ml ampicillin and 60 µg/ml kanamycin. Exponentially growing precultures were used for inoculation of 500 ml baffled shake flasks that contains 200 ml LB medium supplemented with 50 g/l glucose, 5 g/l K₂HPO₄, 80 µg/ml ampicilin and 50 µg/ml kanamycin, which are incubated at 160 rpm and 37 °C. After 5 hours, isopropyl β-thiogalactopyranoside (IPTG) was added at a final concentration of 1 mM, as an inducer of the T7 promoter that is in front of each of the three genes PAL2, 4CL1 and VST1. After an incubation period of 48 hours at 37 °C, the cells were harvested and subjected to extraction procedures and analysed for the presence of produced pinosylvin.

30

e) Extraction and analysis of pinosylvin in *Escherichia coli*.

5 Extraction and analysis were performed using the methods as described in example 9. Results of HPLC conducted on the extracted materials from the fermentation using the engineered strain described and a control strain containing empty plasmids are shown in Figure 9, upper and lower panels respectively. Pinosylvin and cinnamic acid production is
10 marked in the figure.

Example 11

a) *Construction of a bacterial vector for expression of
15 PAL2 in Lactococcus lactis.*

The plasmid pSH71 and derivatives thereof, which is used in the following examples, is a bifunctional shuttle vector with multiple origins of replication from *Escherichia coli*
20 and *Lactococcus lactis*. With that, the host range specificity traverses *Escherichia coli* and other species of lactic acid bacteria. Though transformations in *Lactococcus lactis* usually proceed without problems, putative difficult transformations in other species of lactic acid bacteria
25 can, therefore, be overcome by using *Escherichia coli* as an intermediate host for the construction of recombinant plasmids. The plasmid contains one or more marker genes to allow the microorganism that harbour them to be selected from those which do not. The selection system that is used
30 for *Lactococcus lactis* is based upon dominant markers, e.g. resistance against erythromycin and chloramphenicol, but systems based upon genes involved in carbohydrate

metabolism, peptidases and food grade markers, have also been described. In addition, the plasmid contains promoter- and terminator sequences that allow the expression of the recombinant genes. Suitable promoters are taken from genes of *Lactococcus lactis* e.g. *lacA*. Furthermore, the plasmid contains suitable unique restriction sites to facilitate the cloning of DNA fragments and subsequent identification of recombinants.

10 In the procedures below the plasmid contains either the erythromycin resistance gene, designated as pSH71-ERY^r, or the chloramphenicol resistance gene, designated as pSH71-CM^r.

The gene encoding PAL2, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-URA-PAL2 (example 2), using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the gene allows ligation of the restricted PCR product into a digested pSH71-ERY^r vector that contains the *lacA* promoter from *Lactococcus lactis*. The resulting plasmid, pSH71-ERY^r-PAL2, contains the gene encoding PAL2 under the control of the *lacA* promoter from *Lactococcus lactis*. The sequence of the gene encoding PAL2 is verified by sequencing of two different clones of pSH71-ERY^r-PAL2.

b) Construction of a bacterial vector for expression of 4CL1 and VST1 in *Lactococcus lactis*.

30 The gene encoding 4CL1, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-TRP-4CL1-VST1 (example 5), using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The

introduction of said restriction sites at the 5' and 3' ends of the gene allows ligation of the restricted PCR product into a digested pSH71-CM^r vector. The resulting plasmid, pSH71-CM^r-4CL1, contains the gene encoding for 4CL1 under the control of the *lacA* promoter from *Lactobacillus lactis*.
5 The gene encoding VST1, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-TRP-4CL1-VST1 (example 5) using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The
10 introduction of said restriction sites at the 5' and 3' ends of the gene allows ligation of the restricted PCR product into a digested pSH71-ERY^r vector. The resulting plasmid, pSH71-ERY^r-VST1, contains the gene encoding VST1 under the control of the *lacA* promoter from *Lactococcus lactis*. The
15 *lacA* promoter and the gene encoding VST1 are reamplified as one fragment by PCR from the plasmid pSH71-ERY^r-VST1 using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said
20 restriction sites at the 5' and 3' ends of the DNA fragment allows ligation of the restricted PCR product into the digested plasmid pSH71-CM^r-4CL1. The resulting plasmid, pSH71-CM^r-4CL1-VST1, contains the genes encoding 4CL1 and VST1 that are each under the control of their individual
25 *lacA* promoter. The sequence of the genes encoding 4CL1 and VST1 is verified by sequencing of two different clones of pSH71-CM^r-4CL1-VST1.

c) Expression of the pathway to pinosylvin in *Lactococcus lactis*
30
Lactococcus lactis strains are transformed with the vectors described in examples 16 and 17, separately or in combination. The transformation of the bacterial cell is

conducted in accordance with methods known in the art, for instance, by using competent cells or by electroporation (see, e.g., Sambrook *et al.*, 1989). Transformants are selected on medium containing the antibiotics erythromycin and chloramphenicol and streak purified on the same medium.

Lactococcus lactis strain MG1363 is transformed separately with the vector pSH71-ERY^r-PAL2 (example 16), yielding the strain FSSL-PAL2. In addition, *Lactococcus lactis* strain MG1363 is co-transformed with pSH71-ERY^r-PAL2 (example 16) and pSH71-CM^r-4CL1-VST1 (example 17), and the transformed strain is named FSSL-PAL24CL1VST1.

d) Fermentation with recombinant *Lactococcus lactis* strains in fermentors.

The recombinant *Lactococcus* strains can be grown in fermenters operated as batch, fed-batch or chemostat cultures.

Batch and Fed-batch cultivations

The microorganism is grown in a baffled bioreactor with a working volume of 1.5 liters under anaerobic, aerobic or microaerobic conditions. All cultures are incubated at 30°C, at 350 rpm. A constant pH of 6.6 is maintained by automatic addition of 10 M KOH. Cells are grown on lactose in defined MS10 medium supplemented with the following components to allow growth under aerobic conditions: MnSO₄ (1.25 × 10⁻⁵ g/l), thiamine (1 mg/l), and DL-6,8-thioctic acid (2.5 mg/l). The lactose concentration is, for example 50 g/l. The bioreactors are inoculated with cells from precultures grown at 30°C in shake flasks on the medium described above

buffered with threefold-higher concentrations of K_2HPO_4 and KH_2PO_4 . Anaerobic conditions are ensured by flushing the medium with N_2 (99.998% pure) prior to inoculation and by maintaining a constant flow of 50 ml/min of N_2 through the headspace of the bioreactor during cultivation. The bioreactors used for microaerobic and aerobic cultivation are equipped with polarographic oxygen sensors that are calibrated with air (DOT, 100%) and N_2 (DOT, 0%). Aerobic conditions are obtained by sparging the bioreactor with air at a rate of 1 vvm to ensure that the DOT is more than 80%. During microaerobic experiments the DOT is kept constant 5% by sparging the reactor with gas composed of a mixture of N_2 and atmospheric air, at a rate of 0.25 vvm.

15 Chemostat cultures

In chemostat cultures the cells can be grown in, for example, 1-L working-volume Applikon laboratory fermentors at 30 °C and 350 rpm. The dilution rate (D) can be set at different values, e.g. at 0.050 h^{-1} , 0.10 h^{-1} , 0.15 h^{-1} , or 0.20 h^{-1} . The pH is kept constant, e.g. at 6.6, by automatic addition of 5 M KOH, using the growth medium described above, supplemented with antifoam (50 μ l/l). The concentration of lactose can be set at different values, e.g. is 3.0 g/l 6.0 g/l, 12.0 g/l, 15.0 g/l or 18.0 g/l. The bioreactor is inoculated to an initial biomass concentration of 1 mg /l and the feed pump is turned on at the end of the exponential growth phase.

An anaerobic steady state is obtained by introducing 50 ml/min of N_2 (99.998% pure) into the headspace of the bioreactor. Different anoxic steady states can be obtained by sparging the reactor with 250 ml/min of gas composed of N_2

(99.998% pure) and atmospheric air at various ratios. The oxygen electrode is calibrated by sparging the bioreactor with air (100% DOT) and with N₂ (0% DOT).

For all conditions, the gas is sterile filtered before being
5 introduced into the bioreactor. The off gas is led through a condenser cooled to lower than -8°C and analyzed for its volumetric content of CO₂ and O₂ by means of an acoustic gas analyser.

Cultivations are considered to be in steady state after at
10 least 5 residence times, and if the concentrations of biomass and fermentation end products remain unchanged (less than 5% relative deviation) over the last two residence times.

15 e) Extraction and analysis of pinosylvin in *Lactococcus lactis*

Extraction and analysis is performed using the methods as described in example 9.

20

Example 12

a) Construction of a fungal vector for expression of PAL2 in species belonging to the genus *Aspergillus*.

25 The plasmid that is used in this example, is derived from pARp1 that contains the AMA1 initiating replication sequence from *Aspergillus nidulans*, which also sustains autonomous plasmid replication in *A. niger* and *A. oryzae* (Gems *et al.*, 1991). Moreover, the plasmid is a shuttle vector, containing
30 the replication sequence of *Escherichia coli*, and the inherent difficult transformations in *Aspergillus niger* and *Aspergillus oryzae* can therefore overcome by using *Escherichia coli* as an intermediate host for the

construction of recombinant plasmids. The plasmid contains one or more marker genes to allow the microorganism that harbour them to be selected from those which do not. The selection system can be either based upon dominant markers e.g. resistance against hygromycin B, phleomycin and bleomycin, or heterologous markers e.g amino acids and the *pyrG* gene. In addition the plasmid contains promoter- and terminator sequences that allow the expression of the recombinant genes. Suitable promoters are taken from genes of *Aspergillus nidulans* e.g. *alcA*, *glaA*, *amy*, *niaD*, and *gpdA*. Furthermore, the plasmid contains suitable unique restriction sites to facilitate the cloning of DNA fragments and subsequent identification of recombinants.

The plasmid contains the strong constitutive *gpdA*-promoter and auxotrophic markers, all originating from *Aspergillus nidulans*; the plasmid containing the gene *methG* that is involved in methionine biosynthesis, is designated as pAMA1-MET; the plasmid containing the gene *hisA* that is involved in histidine biosynthesis, is designated as pAMA1-HIS.

The gene encoding for PAL2, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-URA-PAL2 (example 2) using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the gene allows ligation of the restricted PCR product into a digested pAMA1-MET vector. The resulting plasmid, pAMA1-MET-PAL2, contains the gene encoding for PAL2 under the control of the *gpdA* promoter from *Aspergillus nidulans*. The sequence of the gene encoding for PAL2 is verified by sequencing of two different clones of pAMA1-MET-PAL2.

b) Construction of a fungal vector for expression of 4CL1 and VST1 in species belonging to the genus *Aspergillus*.

5 The gene encoding 4CL1, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-TRP-4CL1-VST1 (example 5), using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends
10 of the gene allows ligation of the restricted PCR product into a digested pAMA1-HIS vector that contains the *gpdA* promoter from *Aspergillus nidulans*. The resulting plasmid, pAMA1-HIS-4CL1 contains the gene encoding 4CL1 under the control of the *gpdA* promoter from *Aspergillus nidulans*.

15 The gene encoding VST1, isolated as described in example 1, is reamplified by PCR from the plasmid pESC-TRP-4CL1-VST1 (example 5) using forward- and reverse primers, with 5' overhangs containing suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends
20 of the gene allows ligation of the restricted PCR product into a digested pAMA1-MET vector to yield pAMA1-MET-VST1. The *gpdA* promoter and the gene encoding VST1 are reamplified as one fragment by PCR from the plasmid pAMA1-MET-VST1 using forward- and reverse primers, with 5' overhangs containing
25 suitable restriction sites. The introduction of said restriction sites at the 5' and 3' ends of the DNA fragment allows ligation of the restricted PCR product into the digested plasmid pAMA1-HIS-4CL1. The resulting plasmid, pAMA1-HIS-4CL1-VST1, contains the genes encoding 4CL1 and
30 VST1 that are each under the control of an individual *pgdA* promoter from *Aspergillus nidulans*. The sequence of the genes encoding 4CL1 and VST1 is verified by sequencing of two different clones of pAMA1-HIS-4CL1-VST1.

c) Expression of the pathway to pinosylvin in *Aspergillus niger*.

5 *Aspergillus niger* strains are transformed with the vectors described in (a) and (b), separately or in combination. The transformation of the fungal cell is conducted in accordance with methods known in the art, for instance, by electroporation or by conjugation (see, e.g., Sambrook et
10 *al.*, 1989). Transformants are selected on minimal medium lacking methionine and/or histidine.

A strain of *Aspergillus niger* that is auxotrophic for histidine and methionine, for instance, strain FGSC A919
15 (see <http://www.fgsc.net>), is transformed separately with the vector pAMA1-MET-PAL2 (a), yielding the strain FSAN-PAL2 and with pAMA1-HIS-4CL1-VST1 (b), yielding strain FSAN-4CL1VST1. In addition, *Aspergillus niger* strain FGSC A919 is co-transformed with pAMA1-MET-PAL2 (a) and pAMA1-HIS-4CL1-
20 VST1 (b), and the transformed strain is named FSAN-PAL24CL1VST1.

Example 13

25 *Expression of the pathway to pinosylvin in Aspergillus oryzae.*

A strain of *Aspergillus oryzae* that contains a native set of genes encoding for PAL2 and 4CL1 (Seshime *et al.*, 2005) and that is auxotrophic for methionine, is transformed with the
30 vector pAMA1-MET-VST1 (example 29), yielding the strain FSAO-VST1. The transformation of the fungal cell is conducted in accordance with methods known in the art, for instance, by electroporation or by conjugation (see, e.g.,

Sambrook *et al.*, 1989). Transformants are selected on minimal medium lacking methionine.

Example 14

5 *Fermentation with recombinant strains of Aspergillus niger and Aspergillus oryzae in fermentors.*

The recombinant *Aspergillus* strains can be grown in fermenters operated as batch, fed-batch or chemostat
10 cultures.

Batch and Fed-batch cultivations

The microorganism is grown in a baffled bioreactor with a
15 working volume of 1.5 liters under aerobic conditions. All cultures are incubated at 30 °C, at 500 rpm. A constant pH of 6.0 is maintained by automatic addition of 10 M KOH, and aerobic conditions are obtained by sparging the bioreactor with air at a rate of 1 vvm to ensure that the DOT is more
20 than 80%. Cells are grown on glucose in defined medium consisting of the following components to allow growth in batch cultivations: 7.3 g/l (NH₄)₂SO₄, 1.5 g/l KH₂PO₄, 1.0 g/l MgSO₄·7H₂O, 1.0 g/l NaCl, 0.1 g/l CaCl₂·2H₂O, 0.1 ml/l Sigma antifoam, 7.2 mg/l ZnSO₄·7H₂O, 1.3 mg/l CuSO₄·5H₂O, 0.3 mg/l
25 NiCl₂·6H₂O, 3.5 mg/l MnCl₂·4H₂O and 6.9 mg/l FeSO₄·7H₂O. The glucose concentration is, for example, 10-, 20-, 30-, 40- or 50 g/l. To allow growth in fed-batch cultivations the medium is composed of: 7.3 g/l (NH₄)₂SO₄, 4.0 g/l KH₂PO₄, 1.9 g/l
30 MgSO₄·7H₂O, 1.3 g/l NaCl, 0.10 g/l CaCl₂·2H₂O, 0.1 ml/l Sigma antifoam, 7.2 mg/l ZnSO₄·7H₂O, 1.3 mg/l CuSO₄·5H₂O, 0.3 mg/l NiCl₂·6H₂O, 3.5 mg/l MnCl₂·4H₂O and 6.9 mg/l FeSO₄·H₂O in the batch phase. The reactor is then fed with, for example, 285 g/kg glucose and 42 g/kg (NH₄)₂SO₄.

Free mycelium from a pre-batch is used for inoculating the batch- and fed-batch cultures. A spore concentration of 2.10^9 spores/l is used for inoculation of the pre-batch culture at pH 2.5. Spores are obtained by propagation of freeze-dried
5 spores onto 29 g rice to which the following components are added: 6 ml 15 g/l sucrose, 2.3 g/l $(\text{NH}_4)_2\text{SO}_4$, 1.0 g/l KH_2PO_4 , 0.5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.50 g/l NaCl, 14.3 mg/l $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 2.5 mg/ $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.50 mg/l $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$, and 13.8 mg/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$. The spores are propagated at 30 °C for 7-14 days
10 to yield a black layer of spores on the rice grains and are harvested by adding 100 ml of 0.1% Tween 20 in sterile water. For all conditions, the gas is sterile filtered before being introduced into the bioreactor. The off gas is led through a condenser cooled to lower than $\sim 8^\circ\text{C}$ and
15 analyzed for its volumetric content of CO_2 and O_2 by means of an acoustic gas analyser.

Chemostat cultures

20 In chemostat cultures the cells can be grown in, for example, 1.5-L working-volume Biostat B laboratory fermentors at 30 °C and 500 rpm. A constant pH of 6.0 is maintained by automatic addition of 10 M KOH, and aerobic conditions are obtained by sparging the bioreactor with air
25 at a rate of 1 vvm to ensure that the DOT is more than 80%. The dilution rate (D) can be set at different values, e.g. at 0.050 h^{-1} , 0.10 h^{-1} , 0.15 h^{-1} , or 0.20 h^{-1} . The pH is kept constant, e.g. at 6.6, by automatic addition of 10 M KOH, using a minimal growth medium with the following components:
30 2.5 g/l $(\text{NH}_4)_2\text{SO}_4$, 0.75 g/l KH_2PO_4 , 1.0 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 1.0 g/l NaCl, 0.1 g/l $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 0.1 ml/l Sigma antifoam, 7.2 mg/l $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 1.3 mg/l $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.3 mg/l $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$, 3.5 mg/l

MnCl₂·4H₂O and 6.9 mg/l FeSO₄·7H₂O. The concentration of glucose can be set at different values, e.g. is 3.0 g/l 6.0 g/l, 12.0 g/l, 15.0 g/l or 18.0 g/l. The bioreactor is inoculated with free mycelium from a pre-batch culture as described above, and the feed pump is turned on at the end of the exponential growth phase.

For all conditions, the gas is sterile filtered before being introduced into the bioreactor. The off gas is led through a condenser cooled to lower than -8°C and analyzed for its volumetric content of CO₂ and O₂ by means of an acoustic gas analyser.

Cultivations are considered to be in steady state after at least 5 residence times, and if the concentrations of biomass glucose and composition of the off-gas remain unchanged (less than 5% relative deviation) over the last two residence times.

Example 15

Extraction and analysis of pinosylvin in Aspergillus niger and Aspergillus oryzae

Extraction and analysis is performed using the methods as described in Example 9.

Example 16

Pinosylvin production in Aspergillus nidulans AR1

Aspergillus nidulans AR1 has deleted the following genes genes argB2, pyrG89, veA.

a) Construction of a filamentous fungal expression vector, with *argB* (ornithine carbamoyltransferase) marker.

The gene encoding *argB* including the homologous promoter and terminator sequence was amplified from *Aspergillus nidulans* AR1 genomic DNA using forward primer 5-CG GAATTC ATA CGC GGT TTT TTG GGG TAG TCA-3 (SEQ ID NO: 17) and the reverse primer 5-CG CCCGGG TAT GCC ACC TAC AGC CAT TGC GAA-3 (SEQ ID NO: 18) with the 5' overhang containing the restriction sites EcoRI and XmaI respectively.

The incorporated restriction sites in the PCR product allowed insertion into pUC19 (New England Biolabs, Ipswich, MA.) digested with EcoRI and XmaI giving pUC19-*argB*.

The *trpC* (Indole-3-glycerol phosphate synthase) terminator was amplified from *A. nidulans* genomic DNA using forward primer 5-GC GGATCC ATA GGG CGC TTA CAC AGT ACA CGA-3 (SEQ ID NO: 19) and the reverse primer 5-CGGAGAGGGCGCGCCCGTGGCGGCCGC GGA TCC ACT TAA CGT TAC TGA-3 (SEQ ID NO: 20) with the 5' overhang containing the restriction site BamHI and a 27 base pair adaptamer respectively.

The *gpdA* (glyceraldehyde-3-phosphate dehydrogenase) promoter was amplified from *A. nidulans* AR1 genomic DNA using forward primer 5-GCGGCCGCCACGGGCGCGCCCTCTCCG GCG GTA GTG ATG TCT GCT CAA-3 (SEQ ID NO: 21) and the reverse primer 5-CG AAGCTT TAT AAT TCC CTT GTA TCT CTA CAC-3 (SEQ ID NO: 22) with the 5' overhang containing a 27 base pair adaptamer and the restriction site HindIII respectively.

The fusion PCR product of fragment *trpC* and *gpdA* with the incorporated restriction sites allow insertion into pUC19-*argB* digested with BamHI and HindIII yielding pAT3.

b) Construction of a filamentous fungal expression vector with pyrG (orotidine-5'-monophosphate decarboxylase) marker for expression of C4H (Cinnamate-4-hydroxylase) in *A. nidulans* AR1.

5

The gene encoding C4H was reamplified from the yeast plasmid pESC-URA-PAL2-C4H (WO2006089898, example 3) using the forward primer 5-CG G CGCG C ATA ATG GAC CTC CTC TTG CTG GAG-3 (SEQ ID NO: 23) and the reverse primer 5-GG GC GGCC GC TTA TTA ACA GTT CCT TGG TTT CAT AAC G-3 (SEQ ID NO: 24) with the 5' overhang containing the restriction sites BssHII and NotI respectively. The incorporated restriction sites in the PCR product allowed insertion into pAT3 digested with BssHII and NotI giving pAT3-C4H. The construct was verified by restriction enzyme cut and sequencing. The argB marker was removed by using the two following restriction enzymes BsiWI and PciI.

The gene encoding pyrG including the homologous promoter and terminator sequence was reamplified from *Aspergillus fumigatus* genomic DNA using the forward primer 5-CGT GTAC AATA TTA AT TAA CGAGA GCG AT CGC AAT AAC CGT ATT ACC GCC TTT GAG-3 (SEQ ID NO: 25) and reverse primer 5-CGA CATG TAT TCC CGG GAA GAT CTC ATG GTC A-3 (SEQ ID NO: 26) with the 5' overhang containing the restriction sites BsrGI, PacI, AsiSI in the forward primer and PciI in the reverse primer. The incorporated restriction sites in the PCR product allowed insertion into pAT3 digested with BsiWI and PciI giving pAT3-C4H-pyrG. The construct was verified by restriction enzyme cut and sequencing.

30

c) Construction of a filamentous fungal expression vector with *argB* marker for expression of 4CL1 (4-coumarate-CoA ligase) in *A. nidulans* AR1

5 The gene encoding 4CL1 was reamplified from the yeast plasmid pESC-TRP-4CL1-VST1 using the forward primer 5-GCGGAGAGGGCGCG ATG GCG CCA CAA GAA CAA GCA-3 (SEQ ID NO: 27) and the reverse primer 5-TGGATCCGCGGCCGC TCA CAA TCC ATT TGC TAG TTT TGC-3 (SEQ ID NO: 28). The 4CL1 gene was inserted into a
10 pAT3 vector digested with BssHII and NotI using the In-fusion™ PCR cloning Technology (Clontech, Mountain View, Calif.) to yield pAT3-4CL1. The construct was verified by restriction enzyme cut and sequencing.

15 d) Construction of a filamentous fungal expression vector with *argB* marker for expression of VST1 (resveratrol synthase) in *A. nidulans* AR1

20 The gene encoding VST1 was reamplified from the yeast plasmid pESC-TRP-4CL1-VST1 (example 5) using the forward primer 5-CG G CGCG C ATA ATG GCA TCC GTA GAG GAG TTC-3 (SEQ ID NO: 29) and the reverse primer 5-GG GC GGCC GC TTA TCA TTA GTT AGT GAC AGT TGG AA-3 (SEQ ID NO: 30) with the 5'
25 overhang containing the restriction sites BssHII and NotI respectively. The incorporated restriction sites in the PCR product allowed insertion into pAT3 digested with BssHII and NotI giving pAT3-VST1. The construct was verified by restriction enzyme cut and sequencing.

30 e) Expression of the pathway leading to pinosylvin in *A. nidulans* AR1 (The strain has deletions (*argB2*, *pyrG89*, *veA1*)) using C4H, 4CL1 and VST1.

The transformation of the *A. nidulans* AR1 fungal cell was conducted in accordance with methods known in the art by protoplastation using cell wall lysing enzymes (glucanex, 5 novozymes) Tilburn et al., 1983. Random integration of C4H, 4CL1 and VST1 was conducted in two steps. Plasmid pAT3-4CL1 and pAT3-VST1 were linearized using restriction enzyme BmrI and integrated in the genome by co-transformation according to Guerra et al., 2006 utilizing the auxotrophic marker 10 argB. A transformant containing a 4CL1 and VST1 expression cassette was isolated and a successive transformation with pAT3-C4H-pyrG, which was linearized with BmrI, gave a recombinant *A. nidulans* strain containing C4H, 4CL1 and VST1.

15

f) Fermentation with recombinant *A. nidulans* strains in shake flasks.

Precultures of *A. nidulans* were grown for 5 days on agar 20 plates at 37 °C containing 1g/L glucose, 0.85g/L NaNO₃, 0.1 g/L KCl, 0.1 g/L MgSO₄·7H₂O; and 0.3 g/L KH₂PO₄, 0.00008 g/L CuSO₄·5H₂O, 0.000008g/L Na₂B₄O₇·10H₂O, 0.00016g/L FeSO₄·7H₂O, 0.00016g/L MnSO₄·2H₂O, 0.00016g/L Na₂MoO₄·2H₂O, and 0.0016g/L ZnSO₄·7H₂O. The precultures were used for inoculation of 25 500 ml baffled shake flasks containing 100 ml Czapek medium (CZ). The shake flasks were incubated at 150 rpm and 30 °C and the initial pH of the medium was 6.2. After an incubation period of 24 hours, the samples were taken and subjected to extraction procedures (see below) and analyzed 30 for the presence of produced pinosylvin.

g) Extraction of pinosylvin from *A. nidulans* shake flask cultures

Samples consisting of 100 ml cultures (both cells and broth) were withdrawn from the shake flasks. Extraction of metabolites were conducted as follows; the samples were transferred into two 50 ml Sartorius tubes and centrifuged at 4500 rpm for 10 minutes. The supernatant was transferred into a beaker and the biomass was divided into eight aliquots that were transferred to 2 ml Sarstedt micro tubes with cap, containing app. 300µl glass beads (0.25-0.50mm). The tubes were inserted into a Fastprep 120 (Thermo Fisher Scientific, Waltham, MA.) for four cycles at level 6.5 for 30 seconds at a time and kept on ice in between cycles. The crushed cells were divided into two 15-ml Sartorius tubes. The tubes were filled with 10 ml of supernatant and 3 ml of ethyl acetate was added. The tubes were vigorously mixed on a whirly mixer for 2 minutes and put on ice for 5 minutes. The ethyl acetate phase was then separated from the water phase via centrifugation at 4500 rpm for 10 minutes and collected in four 1.5 ml Eppendorf tubes. The ethyl acetate was then freeze dried for 45 min and the dried samples were re-dissolved in 0.3 ml 50% methanol for further HPLC analysis, as described in Example 9b.

h) Shake flask results from recombinant *A. nidulans*

Figure 7 shows HPLC-chromatograms from a typical shake flask experiment. The upper panel shows results from the engineered strain producing pinosylvin and the lower panel shows the results from the parent wild type control strain. The pinosylvin levels produced by the engineered strain varied between 1.0-2.0 mg/l. The control strain did not show any pinosylvin formation.

The identity of the pinosylvin peak was further confirmed with diode array UV-spectra by comparison with a pure standard UV-chromatogram (Figure 8).

5 Example 17

Determination of intracellular and extracellular levels of stilbenoids in a continuous culture of PALCPR

A yeast strain FSSC-PAL2C4H4CL2VST1-pADH1CPR1 with
10 overexpressed CPR, was grown in a carbon-limited continuous culture with a working volume of 1 liter. The culture was fed with a defined medium according to Verduyn *et al.* (1992), containing: 5.0 g/L (NH₄)₂SO₄; 3.0 g/L KH₂PO₄; 0.5 g/L MgSO₄·7H₂O; trace metals and vitamins and 5 g/l glucose and
15 35 g/l galactose as the growth-limiting nutrients. Antifoam (300 µl/L, Sigma A-8436) was added to avoid foaming. The carbon source was autoclaved separately from the mineral medium and afterwards added to the fermentor. In addition, the vitamin and trace metal solutions were added to the
20 fermentor by sterile filtration following autoclavation and cooling of the medium. The fermentor system was from Sartorius BBI systems and consisted of a baffled 3-liter reactor vessel with 1 liter working volume equipped with Biostat B Plus controller. The reactor vessel was equipped
25 with two Rushton turbines which were rotating at either 1000 rpm, the temperature was kept at 30 ± 1°C, and the pH was kept at 5.5 ± 0.2 by automatic addition of 2M KOH. The gasflow was controlled by a mass flow controller and was set to 1.5 vvm (1.5 l/min). The off-gas was led through a
30 cooled condenser, and was analyzed for O₂ and CO₂ (Model 1308, Innova, Denmark). An initial batch culture with 35 g/l galactose was started by inoculation of the culture with

10 ml of an exponential growing shakeflask culture containing 5 g/l glucose and 35 g/l galactose. The batch cultivation was switched to a continuous mode by feeding the same medium continuously to the reactor. The dilution rate was controlled on a constant level basis, aiming at $D = 0.050 \text{ h}^{-1}$. The continuous culture was regarded to be in steady state when both the dilution rate and off-gas signal had not changed for at least five residence times, and when the metabolite concentrations in two successive samples taken at intervals of 1 residence time, deviated by less than 3%. The dissolved-oxygen concentration, which was continuously monitored, was kept above 60% of air saturation. Under said conditions the strain consumed all the galactose, and mainly produced biomass and CO_2 , and only minor amounts of ethanol. Moreover, the RQ was close to unity, indicating that metabolism was predominantly in respirative mode.

For the determination of stilbenoids, samples were taken at approximately 300 hrs into fermentation corresponding to 15 residence times. Cells were harvested by centrifugation 5000 g for 5 minutes. For the determination of extracellular levels of stilbenoids, an aliquot of 25 ml of supernatant was extracted once with 10 ml ethyl acetate. The ethyl acetate was freeze dried and the dry product redissolved in 0.6 ml methanol. The samples were than 50-fold diluted in water transferred into HPLC vials, and analyzed by HPLC. Furthermore, to evaluate whether the level of stilbenoids that was produced exceeded the solubility of the medium, or were either bound to the cell-membranes 1 ml aliquots of cell culture, thus including both cells and medium, were mixed with 1 ml of 100% ethanol, and mixed vigorously prior to centrifugation. The supernatant was

then transferred into HPLC vials and directly analyzed for the content of stilbenoids. For the determination of intracellular levels of stilbenoids, an aliquot of 50 ml culture was sampled, and cells and medium were separated by centrifugation. The pellet was washed with 50 ml of water to remove any stilbenoids that were cell-bound or trapped into the pellet; after re-centrifugation the pellet was then dissolved in 1 ml water. The resulting cell suspension was distributed into extraction tubes and broken with glass beads using a fast-prep machine. The crude extracts were pooled into 10 ml of 100 % methanol, and extracted in a rotary chamber for 24 hours in a dark cold room at 4 °C. Thereafter, the cell debris was removed via centrifugation for 5 min. at 5000 g and the remaining methanol was removed by freeze-drying overnight. The dry residue was redissolved in 0.4 ml methanol and 0.1 ml water. The samples were then 50-fold diluted in water and then transferred into HPLC vials, and analyzed by HPLC.

The following table summarizes the results after continuous culture for 300 hrs:

	Pinosylvin Intracellular (a)	Pinosylvin Extracellular (b)	Pinosylvin Extracellular In EtOH (c)	Pinosylvin Total (a + c)
mg/l	16.45	12.55	113.57	130.02
% of total	12.65	9.65	87.35	100.00
mg/g dry weight	1.83	-	-	-

Intracellular levels of stilbenoids were expressed in mg per gram biomass (dry weight), according to the calculation explained in the following section. The concentration of pinosylvin in the extract was determined 1646 mg/l; the
5 volume of the extract was 0.5 ml, hence the absolute amount of pinosylvin extracted was $0.5 \times 1646 / 1000 = 0.8230$ mg respectively. The stilbenoids were extracted from a 50 ml culture-aliquot and hence the intracellular concentrations of pinosylvin expressed per liter culture were
10 $0.8230 \times (1000 / 50) = 16.46$ mg/l. The biomass concentration of said culture was 9 g/l. The intracellular pinosylvin levels expressed per gram dry weight therefore were $16.46 / 9 = 1.83$ mg/g dry weight.

15 Example 18

Cloning of trans-pinosylvin pathway in oleaginous yeast
Yarrowia lipolytica

20 a) Isolation of genes

PAL (phenylalanine ammonialyase), CL (cinnamoyl:CoA ligase) and VST1 genes, where gene is defined as protein coding sequence, are produced as synthetic genes (GenScript
25 Corporation, Piscataway, NJ) with codon optimization for expression in *Yarrowia lipolytica*. The determination of codon usage in *Y. lipolytica* has been described previously (WO2006125000). PAL and 4CL genes can also be isolated by PCR from *A. thaliana* cDNA (Stratagene). Cinnamoyl:CoA
30 ligase CL can be any hydroxycinnamoyl:CoA ligase accepting cinnamic acid as substrate. For example, the 4-coumaroyl:CoA ligases from *A. thaliana*, encoded by 4CL1 and 4CL2 genes, accept cinnamic acid although the preferred

substrate is 4-hydroxycinnamic acid (coumaric acid)
(Hamberger and Hahlbrock, 2004; Costa et al, 2005). Most
preferably, the CL is a codon optimized ligase specific for
cinnamic acid as substrate exemplified by cinnamate:CoA
5 ligase from *Streptomyces coelicolor* (Kaneko et al, 2003).
Likewise, VST1 gene can be any codon optimized or non
optimized stilbene synthase accepting cinnamoyl:CoA as
substrate even though the preferred substrate is usually 4-
coumaroyl:CoA in stilbene synthases that produce
10 resveratrol, so called resveratrol synthases. This type of
dual substrate acceptance is in the nature of the VST1 gene
(seq id: 9) from *Vitis vinifera*. Most preferably a stilbene
synthase from the family of *Pinus* specific for cinnamoyl:CoA
as substrate is used (Schanz et al, 1992; Kodan et al,
15 2002).

b) Isolation of promoters and terminators

Promoters that can be used for expression of heterologous
20 genes in *Yarrowia lipolytica* are exemplified but not limited
to the following promoters: long chain acyl:CoA oxidase
POX2, hp4d, isocitrate lyase ICL1, extracellular alkaline
protease XPR2, translation elongation factor TEF, ribosomal
protein S7 RPS7, glycerinaldehyde-3-phosphate dehydrogenase
25 GPD, YAT1, GPAT, FBA1, and FBAIN promoters (Müller et al,
1998: WO2006055322; WO2006125000).

Terminators that can be used for expression of heterologous
genes in *Yarrowia lipolytica* are exemplified but not limited
30 to the following terminators: XPR2, LIP2, PEX20, and SQS
terminators (Merkulov et al, 2000; WO2006055322;
WO2006125000).

Isolation of terminator and promoter DNA fragments can be done via PCR from *Yarrowia lipolytica* genomic DNA prepared from whole cells of *Y. lipolytica* exemplified by but not limited to cells from the America Type Culture Collection, such as ATCC16618, ATCC18943, and ATCC18944, ATCC90811, ATCC90812, and ATCC90903.

c) Generation of an expression cassette

The generation of an expression cassette means the assembly of a linear double stranded DNA-fragment consisting of a promoter (constitutive or inducible) fused together with the protein coding sequence of a heterologous gene and a terminator sequence, i.e. 5'-Promoter::Gene::Terminator-3' DNA fragment.

The expression cassette can be generated by a combination of fusion PCR of the different gene fragments; promoter, gene coding sequence and terminal fragment. For example PAL gene can be fused with PCR technology to XPR2 promoter and the resulting XPR2::PAL fragment can be further fused via a second PCR reaction to the terminator to generate the expression cassette XPR2::PAL::terminator.

An alternative way to generate an expression cassette is to clone the protein coding sequence of the heterologous gene (such as PAL) in an existing expression vector, exemplified but not limited to ATCC vector 69355™. This ATCC vector already has a promoter (XPR2) and a terminator region and a multiple cloning site (MCS) with unique restriction sites between the promoter and terminator for introduction of a heterologous gene by standard molecular biology tools. If the number of restriction sites between promoter and

terminator region in the target vector are limited the
Infusion cloning kit technology can be used (Clontech, CA,
USA) since it requires only one restriction site in the
vector for gene insertion. By inserting the gene in a
5 vector between a promoter and terminator the expression
cassette Promoter::Gene::Terminator is created inside a
circular vector and not as a single double stranded DNA-
fragment. If a linear DNA expression cassette fragment is
needed PCR can be used for amplification of the expression
10 cassette from the expression vector. One of skill in the
art would recognize that several expression cassettes can be
introduced into the same plasmid or vector resulting in
cluster of expression cassettes preferably with genes from a
whole metabolic pathway, such as the pinosylvin production
15 pathway (PAL, CL and VST1 genes). The cluster of expression
cassettes for the three genes needed for pinosylvin
production (PAL, CL and VST1) is defined as pinosylvin
pathway expression cluster.

20 d) Insertion of heterologous gene, PAL, CL and VST1 for
pinosylvin production in *Y. lipolytica*

The pinosylvin pathway genes (PAL, CL, VST1) are assembled
as expression cassettes with a promoter and terminator
25 Promoter::Gene:Terminator. The promoters and terminators
can be the same or a combination of different promoters and
terminators for the different genes, PAL, CL and VST1. One
of skill in the art would recognize available cloning
techniques, cloning vectors, or cloning tools needed for
30 introduction and expression of the pinosylvin pathway
expression cluster (comprising the expression cassettes with
the genes PAL, CL and VST1) in *Y. lipolytica*, since these
tools have been described in several publications (Le Dall

et al, 1994; Pignede et al, 2000; Juretzek et al, 2001; Madzak et al, 2004) and patent applications (WO2006055322; WO2006125000).

5 In summary, once the expression cassettes suitable for expressing the pinosylvin pathway (PAL, CL and VST1) in *Y. lipolytica* has been obtained, they can be (i) placed in a plasmid vector capable of autonomous replication in a host cell or (ii) directly integrated into the genome of the host
10 cell or a combination thereof in order to establish the pinosylvin pathway expression cluster in the *Y. lipolytica* host. Expression cassettes can be designed to integrate randomly within the host genome or can be targeted to specific locations. In both cases the expression cassette
15 is further constructed to contain surrounding regions of homology to the host genome on both sides of the expression cassette. The regions of homology can be 20-1000 base pairs sufficient to target recombination with the host locus. Single copies can be targeted to any part of the genome
20 which will not lead to deletion of an essential gene. Integration into multiple locations within the *Y. lipolytica* genome can be particularly useful when high expression levels of genes are desired and targets for integration of multiple copies of expression cassettes are exemplified but
25 not limited to ribosomal DNA sequence (rDNA) or retrotransposon-like elements (TY1 elements) (Pignede et al, 2000). When integrating multiple copies of expression cassettes targeted to random positions into the *Y. lipolytica* genome the expression cassette Promoter-Gene-
30 Terminator can actually be made shorter, including only Promoter-Gene since the integration will allow terminators already present in the *Y. lipolytica* genome to serve as the terminator for the expression cassette.

It is also possible to integrate plasmid DNA comprising expression cassettes into alternate loci to reach the desired copy number for the expression cassette, exemplified by but not limited to the URA3 locus (Accession No AJ306421) and the LEU2 locus (Accession No AF260230). The LEU2 integrative vector is exemplified by but not limited to ATCC vector 69355™. This expression vector containing an expression cassette can be used directly for transformation into *Y. lipolytica* cells auxotrophic for leucine for selection of the expression vector that contains *Y. lipolytica* LEU2 marker gene. The expression cassette can also be amplified from the expression vector by PCR technique to be further used for construction of other expression vectors containing appropriate selective antibiotic markers or biosynthetic amino acid markers.

The URA3 integration site can be used repeatedly in combination with 5-fluoroorotic acid (5-FOA) selection. In detail, native URA3 gene is deleted in *Y. lipolytica* host strain to generate a strain having URA- auxotrophic phenotype, wherein selection occurs based on 5-FOA resistance. When URA3 is present 5-FOA is degraded to a toxic compound 5-fluorouracil by the orotidine-5'-phosphate decarboxylase encoded by URA3 gene and only cells lacking URA3 gene will be resistant. Consequently, a cluster of multiple expression cassettes and a new URA3 gene can be integrated in multiple rounds into different locus of the *Yarrowia lipolytica* genome to thereby produce new strain having URA+ prototrophic phenotype. Subsequent integration produces a new URA3-auxotrophic strain, again using 5-FOA selection, when the introduced URA3 gene is autonomously deleted (so called loop-out or pop-out). Thus, URA3 gene in

combination with 5-FOA selection can be used as a selection marker in multiple rounds of genetic modifications and integration of expression cassettes.

5 e) Transformation of *Y. lipolytica*

Standard transformation techniques (Chen et al, 1997; WO2006125000) can be used to introduce the foreign DNA, self replicative vectors, or DNA fragments comprising the
10 expression cassettes into *Y. lipolytica* host, exemplified by but not limited to host cells such as ATCC90811, ATCC90812, and ATCC90903. The selection method used to maintain the introduced foreign DNA in *Y. lipolytica* can be based on amino acid markers (Fickers et al, 2003) or antibiotic
15 markers (Cordero et al, 1996).

Example 19

(a) Batch cultivations with recombinant *Escherichia coli* strains

20

The recombinant strains of *Escherichia coli* FSEC-PAL24CL1VST1 and BL21 (DE3) (control strain) were grown in baffled bioreactors with a working volume of 1.5 liters, under aerobic conditions. The cultures were incubated at 30
25 °C, at 800 rpm. A constant pH of 7 was maintained by automatic addition of 2N KOH. Aerobic conditions were obtained by sparging the bioreactor with air at a rate of 1 vvm to ensure that the dissolved oxygen density (DOT) was greater than 60%. The air was sterile filtered before being
30 introduced into the bioreactors. The off gas was led through a condenser cooled to lower than 6°C and analyzed for its volumetric content of CO₂ and O₂ by means of an acoustic gas analyser. The bioreactors were equipped with polarographic

oxygen sensors that were calibrated with air (DOT, 100%) and N₂ (DOT, 0%).

Cells were grown on glycerol in semi-defined medium
5 consisting of the following components to allow growth in
batch cultivations: 6.0 g/l yeast extract, 27.2 g/l Na₂HPO₄
(anhydrous), 12.0 g/l KH₂PO₄, 2.0 g/l NaCl, and 4.0 g/l
NH₄Cl. The glycerol concentration was 20 g/l. The medium
was supplemented with 50 mg/l ampicilin and 50 mg/l
10 kanamycin. Antifoam was added to a final concentration of
50 ul/l.

The bioreactors were inoculated with 1 ml of glycerol stock
culture of the recombinant strain, leading to a final
15 optical density at 600 nm of approximately 0.03. The
glycerol stock cultures were obtained by growing the cells
in shake flasks on semi-defined medium, at 30°C and 150 rpm.
The composition of the medium was identical to the one
described above, but re-scaled 4-fold lower, i.e.: 5 g/l
20 glycerol, 1.5 g/l yeast extract, 6.8 g/l Na₂HPO₄ (anhydrous),
3.0 g/l KH₂PO₄, 0.5 g/l NaCl, and 1.0 g/l NH₄Cl. The medium
was supplemented with 50 mg/l ampicilin and 50 mg/l
kanamycin. The cells were harvested during the late
exponential phase, collected by centrifugation and
25 resuspended in an appropriate volume of sterile glycerol
solution 15% (w/v), such that the final optical density at
600 nm was 30. Aliquots of 1 ml of suspended cells were
stored at -80°C.

30 After the cells started growing in the bioreactors (5.5 h
after inoculation), isopropyl β-thiogalactopyranoside (IPTG)
was added to a final concentration of 1 mM, as an inducer of

the T7 promoter that is in front of each of the three genes PAL2, 4CL1, and VST1.

5 Samples of cellular broth were taken in the course of the batch cultivations and analysed for the presence of pinosylvin. In addition, the samples were analysed for biomass (in terms of optical density OD600), carbon source (glycerol) and major by-products (ethanol, acetate, pyruvate, succinate).

10

(b) Extraction of pinosylvin in Escherichia coli

The intracellular pinosylvin was extracted with ethyl acetate. For the purpose, 4 mL of ethyl acetate was added to 8 mL of cell broth. The extraction was enforced by mixing (30 s) and the separation of phases, by centrifugation (4500 rpm for 5 min, at 4 °C). The acetate phase was subjected to freeze-drying (approximately 2 h) and the dry product was redissolved in 0.5 ml methanol and analysed by HPLC. These samples were further diluted in water (1:5) and analysed by HPLC.

20

(c) Analysis of pinosylvin

25 The analysis of pinosylvin in samples from the batch cultivation was performed using the method as described in Example 9b. The sample was previously subjected to the following sample preparation procedures, carried out in parallel: (i) Centrifugation of cell broth (5 min) and analysis of supernatant; (ii) Addition of ethanol (99.9%) to a final concentration of 50% (v/v), vortex (30 s), centrifugation (5 min) and analysis of supernatant; (iii)

30

Extraction with ethyl acetate, according to (b) above, and analysis of dried sample redissolved in methanol.

Results

5

The recombinant strains of *Escherichia coli* FSEC-PAL24CL1VST1 and BL21 (DE3) (control strain), as described in example 10c, were cultivated on 20 g/L of glycerol in bioreactors in batch mode, as described in (a) above. In the course of the cultivations, the recombinant strains were analysed for their content of pinosylvin according to (c) above.

The HPLC-analysis showed that the strain FSEC-PAL24CL1VST1 contained a component with a retention time identical to the standard of *trans*-pinosylvin (figures 4 and 5). In addition, the UV absorption spectra were similar to the absorption spectrum of pure *trans*-pinosylvin (not shown), with a λ_{\max} of approximately 306 nm.

20

The maximal concentrations of pinosylvin detected are shown in the following table:

	Pinosylvin intracellular	Pinosylvin extracellular	Pinosylvin extracellular	Pinosylvin total
	(a)	(b)	In EtOH (c)	(a) + (c)
mg/l	0.016	(*)	(*)	0.016
% of total	100	0	0	100
mg/g dry weight	(**)	(**)	(**)	(**)

(*) below detection level.

(**) not determined.

5 No pinosylvin was detected in the samples from the batch
cultivation with the control strain.

10 The results, therefore, demonstrated the presence of an
active phenyl-propanoid pathway that led to *in vivo*
production of *trans*-pinosylvin, in *E. coli* grown in a
bioreactor in batch mode.

Example 20

15 (a) *Batch cultivation with recombinant Aspergillus nidulans*
strain

15

The recombinant strain of *Aspergillus nidulans* containing
C4H, 4CL1, and VST1 was grown in a baffled bioreactor with a
working volume of 1.5 liters, under aerobic conditions. The
cultures were incubated at 30 °C, at 700 rpm. A constant pH
20 of 6 was maintained by automatic addition of 2N KOH.

Aerobic conditions were obtained by sparging the bioreactor
with air at a rate of 1 vvm to ensure that the dissolved
oxygen tension (DOT) was greater than 60%. The air was
sterile filtered before being introduced into the
25 bioreactors. The off gas was led through a condenser cooled
to lower than 6°C and analyzed for its volumetric content of
CO₂ and O₂ by means of an acoustic gas analyser. The
bioreactors were equipped with polarographic oxygen sensors
that were calibrated with air (DOT, 100%) and N₂ (DOT, 0%).
30 Cells were grown on sucrose in defined medium consisting of
the following components: 3.0 g/l NaNO₃, 1.0 g/l KH₂PO₄, 0.5
g/l KCl, 0.5 g/l MgSO₄·7H₂O, 0.5/l g FeSO₄·7H₂O. The

concentration of sucrose was 30 g/l. Antifoam was added to a final concentration of 50 ul/l.

5 The bioreactor was inoculated with spores of the *A. nidulans* strain containing C4H, 4CL1, and VST1, previously propagated on solid minimal medium, with the following composition: 1 g/L glucose, 0.85 g/L NaNO₃, 0.1 g/L KCl, 0.1 g/L MgSO₄·7H₂O; and 0.3 g/L KH₂PO₄, 0.00008 g/L CuSO₄·5H₂O, 0.000008 g/L Na₂B₄O₇·10H₂O, 0.00016 g/L FeSO₄·7H₂O, 0.00016g/L MnSO₄·2H₂O,
10 0.00016 g/L Na₂MoO₄·2H₂O, and 0.0016 g/L ZnSO₄·7H₂O. The spores were cultivated at 37 °C for 5 days and harvested by adding Tween 80% solution (0.25% (w/v)).

15 *(b) Extraction of pinosylvin in Aspergillus nidulans*

The cells were disrupted by homogenization (in a Polytron tissue homogenizer) and the intracellular pinosylvin was extracted with 10 ml ethyl acetate. The extraction was enforced by mixing in a rotary mixer (approximately 15 min)
20 and the separation of phases, by centrifugation (4500 rpm, at 4 °C, for 5 min). The acetate phase was subjected to freeze-drying (approximately 2 h) and the dry product was redissolved in 0.5 ml methanol and analysed by HPLC.

25 *(c) Analysis of pinosylvin*

The analysis of pinosylvin in samples from the batch cultivation was performed using the method as described in example 9b.

30

Results

The recombinant strain of *Aspergillus nidulans* containing C4H, 4CL1, and VST1, as described in Example 16e, was cultivated on 30 g/L of sucrose in a bioreactor in batch mode, according to Example HD4. After approximately 48h of cultivation, the cells were harvested from the bioreactor, 5 disrupted by homogenization and analysed for their intracellular content of pinosylvin according to (b) and (c) above.

10 The HPLC-analysis showed that the *A. nidulans* strain containing C4H, 4CL1, and VST1 exhibited intracellularly a component with a retention time identical to the standard of *trans*-pinosylvin (Figures 4 and 5). In addition, the UV absorption spectra were similar to the absorption spectrum 15 of pure *trans*-pinosylvin (not shown) as well, with a λ_{\max} of approximately 306 nm.

The results, therefore, demonstrated the presence of an active phenyl-propanoid pathway that led to *in vivo* 20 production of *trans*-pinosylvin, in *A. nidulans* grown in a bioreactor in batch mode.

25

REFERENCES

5 The following publications are all hereby incorporated by
reference:

Patent no. ZA200408194

Patent no. EP0309862

10

Patent no. EP0464461

Patent No. US5391724

15

Patent No. US5973230

Patent application WO2006125000

Method for the production of resveratrol in a recombinant
oleaginous microorganism

20

Huang Lixuan Lisa, Du Pont (US)

Patent application WO2006055322

High arachidonic acid producing strains of *Yarrowia
lipolytica*

25

Damude Howard, Du Pont (US)

Abe, I., Watanabe, T. and Noguchi, H. (2004). Enzymatic
formation of long-chain polyketide pyrones by plant type III
30 polyketide synthases. *Phytochemistry*. 6, 2447-2453.

- Aggarwal BB, Bhardwaj A, Aggarwal RS, Seeram NP, Shishodia S, Takada Y. (2004). Role of resveratrol in prevention and therapy of cancer: preclinical and clinical studies. *Anticancer Res.* 24, 2783-840. Review.
- 5 Allina, S.M., Pri-Hadash, A., Theilmann, D.A., Ellis, B.E. and Douglas, C.J. (1998) 4-coumarate: Coenzyme A ligase in hybrid poplar. Properties of enzymes, cDNA cloning, and analysis of recombinant clones. *Plant Physiol.* 116, 743-754.
- 10 Becker JV, Armstrong GO, van der Merwe MJ, Lambrechts MG, Vivier MA, Pretorius IS. (2003). Metabolic engineering of *Saccharomyces cerevisiae* for the synthesis of the wine-related antioxidant resveratrol. *FEMS Yeast Res.* 4, 79-85.
- 15 Chen DC, Beckerich JM, Gaillardin C.
One-step transformation of the dimorphic yeast *Yarrowia lipolytica*.
Appl Microbiol Biotechnol. 1997:48:232-5.
- 20 Cochrane, F.C., Davin, L.B. and Lewis N.G. (2004). The *Arabidopsis* phenylalanine ammonia lyase gene family: kinetic characterization of the four PAL isoforms. *Phytochemistry* 65, 1557-1564.
- 25 Cordero Otero R, Gaillardin C.
Efficient selection of hygromycin-B-resistant *Yarrowia lipolytica* transformants.
Appl Microbiol Biotechnol. 1996:46:143-8.
- 30 Costa ML, Bedgar DL, Moinuddin SGA, Kim K, Cardenas CL, Cochrane FC, Shockey JM, Helms GL, Amakura Y, Takahashi H et al.

Characterization in vitro and in vivo of the putative multigene 4-coumarate:CoA ligase network in Arabidopsis: syringyl lignin and sinapate/sinapyl alcohol derivative formation

5 Phytochemistry, 2005:66:2072-2091

Ehlting, J., Büttner, D., Wang, Q., Douglas, C.J., Somssich, I.E. and Kombrink, E. (1999). Three 4-coumarate:coenzyme A ligases in *Arabidopsis thaliana* represents two evolutionary
10 divergent classes in angiosperms. The plant journal. 19, 9-20.

Fickers P, Le Dall MT, Gaillardin C, Thonart P, Nicaud JM. New disruption cassettes for rapid gene disruption and
15 marker rescue in the yeast *Yarrowia lipolytica*. J Microbiol Methods. 2003:55:727-37.

Gehlert, R., Schoppner, A. and Kindl, H. Stilbene synthase from seedlings of *Pinus sylvestris* - purification and
20 induction in response to fungal infection. Mol. Plant-Microbe Interaction 3 (1990) 444-449.

Gems, D., Johnstone, I.L. and Clutterbuck, A.J. (1991). An autonomously replicating plasmid transforms *Aspergillus*
25 *nidulans* at high frequency. Gene 98, 61-67.

Hain, R., Reif, H.J., Krause, E., Langebartels, R., Kindl, H., Vornam, B., Wiese, W., Schmelzer, E., Schreier, P.H., Stocker, R.H. and Stenzel, K. (1993). Disease resistance
30 results from foreign phytoalexin expression in a novel plant. Nature 361, 153-156.

Hwang EI, Kaneko M, Ohnishi Y, Horinouchi S. (2003).
Production of plant-specific flavanones by *Escherichia coli*
containing an artificial gene cluster. Appl. Environ.
Microbiol. 69, 2699-706.

5

Hamberger, B. and Hahlbrock, K. (2004). The 4-coumarate:CoA
ligase gene family in *Arabidopsis thaliana* comprises one
rare, sinapate-activating and three commonly occurring
isoenzymes. Proc. Natl. Acad. Sci. USA. 101, 2209-2214.

10

Hart, J. H. (1981) Role of phytostilbenes in decay and
disease resistance. Annu. Rev. Phytopathology 19, 437-458.

15

Hart, J. H., Shrimpton, D. M. (1979). Role of stilbenes in
resistance of wood to decay. Phytopathology 69, 1138-1143.

20

Hemingway R.W., McGraw, G.W. and Barras, S. J. (1977).
Polyphenols in *Ceratocystis minor* infected *Pinus taeda*:
Fungal Metabolites, phloem and xylem phenols. J. Agric. Food
Chem., 25, 717-722.

25

Juvvadi, P.R., Seshime, Y., Kitamoto, K. (2005). Genomics
reveals traces of fungal phenylpropanoid-flavonoid metabolic
pathway in the filamentous fungus *Aspergillus oryzae*. J
Microbiol. 43, 475-486.

30

Juretzek T, Le Dall M, Mauersberger S, Gaillardin C, Barth
G, Nicaud J.
Vectors for gene expression and amplification in the yeast
Yarrowia lipolytica.
Yeast. 2001;18:97-113.

- Kaneko, M., Ohnishi, Y. and Horinouchi, S. Cinnamate:Coenzyme (2003). A ligase from the Filamentous Bacteria *Streptomyces coelicolor* A3(2), J. Bact. 185, 20-27.
- 5 Kindl, H. (1985) Biosynthesis of stilbenes. In Higuchi T, ed, Biosynthesis and Biodegradation of Wood Components. Academic Press, London, pp 349-377.
- Kodan, A., Kuroda, H. and Sakai, F. (2002). A stilbene
10 synthase from Japanese red pine (*Pinus densiflora*): Implications for phytoalexin accumulation and down-regulation of flavonoid biosynthesis. Proc. Natl. Acad. Sci. 99, 3335-3339.
- 15 Le Dall MT, Nicaud JM, Gaillardin C. Multiple-copy integration in the yeast *Yarrowia lipolytica*. Curr Genet. 1994;26:38-44.
- Lieutier, F., Sauvard, D., Brignolas, F., Picron, V., Yart, A., Bastien, C. and Jay-Allemand, C. (1996) Changes in
20 phenolic metabolites of Scots pine phloem induced by *Ophiostoma brunneo-ciliatum*, a bark beetle-associated fungus. Eur. J. For Pathol. 26, 145-158
- 25 Lindberg LE, Willfor SM, Holmbom BR. (2004) Antibacterial effects of knotwood extractives on paper mill bacteria. J Ind Microbiol Biotechnol. 31, 137-147.
- Madzak C, Gaillardin C, Beckerich JM.
30 Heterologous protein expression and secretion in the non-conventional yeast *Yarrowia lipolytica*: a review. J Biotechnol. 2004;109:63-81.

Martin, V.J.J., Pitera, D.J., Withers, S.T., Newman, J.D. and Keasling, J.D. (2003). Engineering a mevalonate pathway in *Escherichia coli* for production of terpenoids. *Nature biotechnology* 21, 796-802.

5

Melchior F, Kindl H (1991). Coordinate and elicitor dependent expression of stilbene synthase and phenylalanine ammonialyase genes in *Vitis cv. Optima*. *Arch. Biochem. Biophys* 288, 552-557.

10

Mellanen, P., Petanen, T., Lehtimaki, J., Makela, S., Bylund, G., Holmbom, B., Mannila, E., Oikari, A. and Santti, R. (1996). Wood-derived estrogens: studies in vitro with breast cancer cell lines and *in vivo* in trout. *Toxicol. App. Pharm.* 136, 381-388.

15

Merkulov S, van Assema F, Springer J, Fernandez Del Carmen A, Mooibroek H.

Cloning and characterization of the *Yarrowia lipolytica* squalene synthase (SQS1) gene and functional complementation of the *Saccharomyces cerevisiae* erg9 mutation. *Yeast*. 2000;16:197-206.

20

Mizutani M and Ohta D et al, 1998

25

Two Isoforms of NADPH:Cytochrome P450 Reductase in *Arabidopsis thaliana*. Gene Structure, Heterologous Expression in Insect Cells, and Differential Regulation *Plant Physiol.* 116, 357-367

30

Morita, H., Noguchi, H., Schröder, J. and Abe, I. (2001). Novel polyketides synthesized with a higher plant stilbene synthase. *Eur. J. Biochem.* 268, 3759-3766.

Müller S, Sandal T, Kamp-Hansen P, Dalbøge H.

Comparison of expression systems in the yeasts *Saccharomyces cerevisiae*, *Hansenula polymorpha*, *Kluyveromyces lactis*, *Schizosaccharomyces pombe* and *Yarrowia lipolytica*. Cloning
5 of two novel promoters from *Yarrowia lipolytica*. Yeast. 1998;14:1267-83.

Nicaud JM, Madzak C, van den Broek P, Gysler C, Duboc P, Niederberger P, Gaillardin C. Protein expression and
10 secretion in the yeast *Yarrowia lipolytica*. FEMS Yeast Res. 2002;2:371-9.

Pignède G, Wang HJ, Fudalej F, Seman M, Gaillardin C, Nicaud JM.
15 Autocloning and amplification of LIP2 in *Yarrowia lipolytica*. Appl Environ Microbiol. 2000;66:3283-9.

Preisig-Muller, R., Schwekendiek, A., Brehm, I., Reif, H.J. and Kindl, H. (1999). Characterization of a pine multigene
20 family containing elicitor-responsive stilbene synthase genes. Plant Mol. Biol. 1999 39, 221-229.

Pacher T, Seger C, Engelmeier D, Vajrodaya S, Hofer O, Greger H. (2002). Antifungal stilbenoids from *Stemona collinsae*. J Nat Prod.65, 820-827.
25

Raiber S, Schröder G, Schröder J. (1995). Molecular and enzymatic characterization of two stilbene synthases from Eastern white pine (*Pinus strobus*). A single Arg/His
30 difference determines the activity and the pH dependence of the enzymes. FEBS Lett. 361, 299-302.

Richter, C., Wild, A. (1992). Phenolic compounds in needles

of Norway spruce trees in relation to novel forest decline:
I. Studies on trees from site of the Northern Black Forest.
Biochem. Biophys. Pflanz 188, 305-320.

5 Ro D.K., Douglas C.J. (2004). Reconstitution of the entry
point of plant phenylpropanoid metabolism in yeast
(*Saccharomyces cerevisiae*): implications for control of
metabolic flux into the phenylpropanoid pathway. J. Biol.
Chem. 279, 2600-2607.

10

Rosemann, D., Heller, W. and Sandermann, H. (1991).
Biochemical Plant Responses to Ozone. II. Induction of
Stilbene Biosynthesis in Scots Pine (*Pinus sylvestris* L.)
Seedlings. Jr. Plant Physiol. 97, 1280-1286.

15

Roupe, K.A., Fukuda, C., Halls, S., Yáñez, J.A. and Davies
N.M. (2005) Pinosylvin: Method of Analysis, Anti-Cancer
Activity and Metabolism. American Association of
Pharmaceutical Scientists Annual Meeting. November.
20 Accepted.

Roupe, K.A., Remsberg, C.M., Yáñez, J.A. and Davies, N.M.
(2006). Pharmacometrics of Stilbenes: Seguing Towards the
Clinic. Curr. Clin. Pharmac. 1, 81-101.

25

Samappito, S., Page, J.E., Schmidt, J., De-Eknamkul, W. and
Kutchan, T.M. (2003). Aromatic and pyrone polyketides
synthesized by a stilbene synthase from *Rheum tataricum*.
Phytochemistry 62, 313-323.

30

Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989).
Molecular Cloning, A Laboratory Manual, 2d edition, Cold
Spring Harbor, N.Y.

Schanz, S., Schröder, G. and Schröder, J. (1992). Stilbene synthase from Scot's pine (*Pinus sylvestris*) FEBS Lett. 313, 71-74.

5

Schöppner, A. and Kindl, H. (1984) Purification and properties of a stilbene synthase from induced cell suspension cultures of peanut. J. Biol. Chem. 259, 6806-6811.

10

Schröder G, Brown JWS, Schröder J. (1988). Molecular analysis of resveratrol synthase. cDNA clones and relationship with chalcone synthase. Eur J Biochem 172, 161-169.

15

Seshime, Y., Juvvadi, P.R., Fujii, I. and Kitamoto, K. (2005). Genomic evidences for the existence of a phenylpropanoid metabolic pathway in *Aspergillus oryzae*. Biochem. Biophys. Res Commun. 337, 747-51.

20

Skinnider, L. and Stoessl A. (1986). The effect of the phytoalexins, lubimin, (-)-maackiain, pinosylvin, and the related compounds dehydroloroglossol and hordatine M on human lymphoblastoid cell lines. Experientia 42, 568-570.

25

Stojanovic, S., Sprinz, H. and Brede, O. (2001). Efficiency and mechanism of the anti-oxidant action of trans-resveratrol and its analogues in the radical liposome oxidation. Arch. Biochem. Biophys. 391, 79-89.

30

Suga T., Ohta S., Munesada K., Ide N., Kurokawa M., Shimizu M., Ohta E. (1993). Endogenous pine wood nematocidal

substances in pines, *Pinus massoniana*, *P. strobus* and *P. palustris*. *Phytochemistry* 33, 1395-1401.

5 Tropf, S., Karcher, B., Schröder, G. and Schröder, J. (1995). Reaction mechanisms of homodimeric plant polyketide synthase (stilbenes and chalcone synthase). A single active site for the condensing reaction is sufficient for synthesis of stilbenes, chalcones, and 6'-deoxychalcones. *J. Biol. Chem.* 270, 7922-7928.

10

Urban P, Werck-Reichhart D, Teutsch HG, Durst F, Regnier S, Kazmaier M, Pompon D (1994). Characterization of recombinant plant cinnamate 4-hydroxylase produced in yeast. Kinetic and spectral properties of the major plant P450 of the phenylpropanoid pathway. *Eur J Biochem.* 222:843-50

15

Verduyn C, Postma E, Scheffers WA, Van Dijken JP. (1992). Effect of benzoic acid on metabolic fluxes in yeasts: a continuous-culture study on the regulation of respiration and alcoholic fermentation. *Yeast.* 8, 501-517.

20

Watts et al; Watts KT, Mijts BN, Lee PC, Manning AJ, Schmidt-Dannert C.(2006). Discovery of a substrate selectivity switch in tyrosine ammonia-lyase, a member of the aromatic amino acid lyase family. *Chem Biol.* 13:1317-26.

25

Wiese W, Vornam B, Krause E, Kindl H (1994). Structural organization and differential expression of three stilbene synthase genes located on a 13 kb grapevine DNA fragment. *Plant Mol Biol* 26, 667-677.

30

The following is a summary of the nucleotide and amino acid
5 sequences appearing herein:

SEQ ID NO: 1 is a nucleotide sequence from *Arabidopsis*
thaliana encoding a phenylalanine ammonia lyase (PAL2).

10 SEQ ID NO: 2 is the amino acid sequence encoded by SEQ ID NO:
1.

SEQ ID NO: 3 is a nucleotide sequence from *Arabidopsis*
thaliana encoding a 4-coumarate:CoenzymeA ligase (4CL1).

SEQ ID NO: 4 is the amino acid sequence encoded by SEQ ID NO:
3.

15 SEQ ID NO: 5 is a nucleotide sequence from *Rheum tataricum*
encoding a resveratrol synthase (RES).

SEQ ID NO: 6 is the amino acid sequence encoded by SEQ ID NO:
5.

20 SEQ ID NO: 7 is a nucleotide sequence from *Rheum tataricum*
encoding a resveratrol synthase (RES), which is codon-
optimized for expression in *S. cerevisiae*.

SEQ ID NO: 8 is the amino acid sequence encoded by SEQ ID NO:
7.

25 SEQ ID NO: 9 is a nucleotide sequence from *Vitis vinifera*
encoding a resveratrol synthase (VST1), which is codon-
optimized for expression in *S. cerevisiae*.

SEQ ID NO: 10 is the amino acid sequence encoded by SEQ ID
NO: 9.

30 SEQ ID NOs 11-16 are primer sequences appearing in Table 1,
Example 1.

SEQ ID Nos 17 to 22 are primer sequences used in Example 16a.

SEQ ID Nos 23 to 26 are primer sequences used in Example 16b.

SEQ ID Nos 27 to 30 are primer sequences used in Example 16c.

Claims

1. The use of a micro-organism for the production of
5 pinosylvin, wherein said micro-organism has an operative
metabolic pathway comprising at least one enzyme activity,
said pathway producing pinosylvin from cinnamic acid.
2. The use claimed in claim 1, wherein said micro-
10 organism produces *cinnamic* acid and produces pinosylvin
therefrom.
3. The use claimed in claim 2, wherein said pinosylvin is
produced in a reaction catalysed by an enzyme in which
15 endogenous malonyl-CoA is a substrate.
4. The use claimed in any preceding claim, wherein said
pinosylvin is produced from cinnamoyl-CoA.
- 20 5. The use claimed in claim 4, wherein said pinosylvin is
produced from cinnamoyl-CoA by a stilbene synthase.
6. The use claimed in claim 5, wherein said stilbene
synthase is expressed in said micro-organism from nucleic
25 acid coding for said enzyme which is not native to the
micro-organism.
7. The use claimed in claim 6, wherein said stilbene
synthase is resveratrol synthase (EC 2.3.1.95) from a
30 plant belonging to the genus of *Arachis*, a plant belonging
to the genus of *Rheum*, or a plant belonging to the genus
of *Vitus* or any one of the genera *Pinus*, *Picea*, *Lilium*,
Eucalyptus, *Parthenocissus*, *Cissus*, *Calochortus*,

Polygonum, Gnetum, Artocarpus, Nothofagus, Phoenix, Festuca, Carex, Veratrum, Bauhinia or Pterolobium.

8. The use claimed in claim 6, wherein said stilbene
5 synthase is a pinosylvin synthase (EC 2.3.1.146) from a
plant belonging to the genus of *Pinus*, e.g. *P. sylvestris*,
P. strobes, *P. densiflora*, *P. taeda*, a plant belonging to
the genus *Picea*, or any one of the genus *Eucalyptus*.
- 10 9. The use claimed in any preceding claim, wherein said
cinnamic acid is produced in said pathway from L-
phenylalanine in a reaction catalysed by an enzyme in
which ammonia is produced.
- 15 10. The use claimed in claim 9, wherein said cinnamic acid
is formed from L-phenylalanine by a L-phenylalanine
ammonia lyase (PAL).
- 20 11. The use claimed in claim 10, wherein said
phenylalanine ammonia lyase is expressed in said micro-
organism from nucleic acid coding for said enzyme which is
not native to the micro-organism.
- 25 12. The use claimed in claim 11, wherein said cinnamic
acid is formed from L-phenylalanine by L-phenylalanine
ammonia lyase (EC 4.3.1.5) from a plant belonging to the
genus of *Arabidopsis*, a plant belonging to the genus of
Brassica, a plant belonging to the genus of *Citrus*, a
plant belonging to the genus of *Phaseolus*, a plant
30 belonging to the genus of *Pinus*, a plant belonging to the
genus of *Populus*, a plant belonging to the genus of
Solanum, a plant belonging to the genus of *Prunus*, a plant
belonging to the genus of *Vitus*, a plant belonging to the

genus of *Zea*, or a plant belonging to any one of the genera *Agastache*, *Ananas*, *Asparagus*, *Bromheadia*, *Bambusa*, *Beta*, *Betula*, *Cucumis*, *Camellia*, *Capsicum*, *Cassia*, *Catharanthus*, *Cicer*, *Citrullus*, *Coffea*, *Cucurbita*,
5 *Cynodon*, *Daucus*, *Dendrobium*, *Dianthus*, *Digitalis*, *Dioscorea*, *Eucalyptus*, *Gallus*, *Ginkgo*, *Glycine*, *Hordeum*, *Helianthus*, *Ipomoea*, *Lactuca*, *Lithospermum*, *Lotus*, *Lycopersicon*, *Medicago*, *Malus*, *Manihot*, *Medicago*, *Mesembryanthemum*, *Nicotiana*, *Olea*, *Oryza*, *Pisum*, *Persea*,
10 *Petroselinum*, *Phalaenopsis*, *Phyllostachys*, *Physcomitrella*, *Picea*, *Pyrus*, *Quercus*, *Raphanus*, *Rehmannia*, *Rubus*, *Sorghum*, *Sphenostylis*, *Stellaria*, *Stylosanthes*, *Triticum*, *Trifolium*, *Triticum*, *Vaccinium*, *Vigna*, or *Zinnia*, or from a filamentous fungus belonging to the genus *Aspergillus*.

15

13. The use as claimed in any one of claims 10 to 12, wherein said PAL is one accepting phenylalanine as a substrate and producing cinammic acid therefrom, such that if the PAL also accepts tyrosine as a substrate and forms
20 coumaric acid therefrom, the ratio $K_m(\text{phenylalanine})/K_m(\text{tyrosine})$ for said PAL is less than 1:1.

25

14. The use as claimed in claim 13, wherein if said micro-organism produces a cinammate-4-hydroxylase enzyme (C4H), the ratio $K_{cat}(\text{PAL})/K_{cat}(\text{C4H})$ is at least 2:1.

30

15. The use claimed in any preceding claim, wherein cinnamoyl-CoA is formed in said pathway in a reaction catalysed by an enzyme in which ATP and CoA are substrates and ADP is a product.

16. The use claimed in any preceding claim, wherein cinnamoyl-CoA is formed in a reaction catalysed by a 4-coumarate-CoA ligase or a cinnamoyl-CoA ligase.

5 17. The use claimed in claim 16, wherein said 4-coumarate-CoA ligase or cinnamate-CoA ligase is 4-coumarate-CoA ligase / cinnamate-CoA ligase (EC 6.2.1.12) from a plant
10 belonging to the genus of *Abies*, a plant belonging to the genus of *Arabidopsis*, a plant belonging to the genus of *Brassica*, a plant belonging to the genus of *Citrus*, a
plant belonging to the genus of *Larix*, a plant belonging to the genus of *Phaseolus*, a plant belonging to the genus of *Pinus*, a plant belonging to the genus of *Populus*, a
15 plant belonging to the genus of *Solanum*, a plant belonging to the genus of *Vitis*, a plant belonging to the genus of *Zea*, or a plant belonging to any one of the genera
Agastache, *Amorpha*, *Cathaya*, *Cedrus*, *Crocus*, *Festuca*,
Glycine, *Juglans*, *Keteleeria*, *Lithospermum*, *Lolium*, *Lotus*,
Lycopersicon, *Malus*, *Medicago*, *Mesembryanthemum*,
20 *Nicotiana*, *Nothotsuga*, *Oryza*, *Pelargonium*, *Petroselinum*,
Physcomitrella, *Picea*, *Prunus*, *Pseudolarix*, *Pseudotsuga*,
Rosa, *Rubus*, *Ryza*, *Saccharum*, *Suaeda*, *Thellungiella*,
Triticum, or *Tsuga*, from a filamentous fungus belonging to
the genus *Aspergillus*, a filamentous fungus belonging to
25 the genus *Neurospora*, a fungus belonging to the genus
Yarrowia, a fungus belonging to the genus of
Mycosphaerella, from a bacterium belonging to the genus of
Mycobacterium, a bacterium belonging to the genus of
Neisseria, a bacterium belonging to the genus of
30 *Streptomyces*, a bacterium belonging to the genus of
Rhodobacter, or from a nematode belonging to the genus
Ancylostoma, a nematode belonging to the genus
Caenorhabditis, a nematode belonging to the genus

Haemonchus, a nematode belonging to the genus *Lumbricus*, a nematode belonging to the genus *Meilodogyne*, a nematode belonging to the genus *Strongyloidus*, or a nematode belonging to the genus *Pristionchus*.

5

18. The use claimed in any preceding claim, wherein at least one copy of at least one genetic sequence encoding a respective enzyme in said metabolic pathway has been recombinantly introduced into said micro-organism.

10

19. The use claimed in any preceding claim wherein at least one copy of a genetic sequence encoding a phenylalanine ammonia lyase is operatively linked to an expression signal not natively associated with said genetic sequence in said organism.

15

20. The use claimed in any preceding claim wherein at least one copy of a genetic sequence encoding a 4-coumarate-CoA ligase or cinnamate-CoA ligase is operatively linked to an expression signal not natively associated with said genetic sequence in said organism.

20

21. The use claimed in any preceding claim wherein at least one copy of a genetic sequence encoding a resveratrol synthase is operatively linked to an expression signal not natively associated with said genetic sequence in said organism.

25

22. The use claimed in any preceding claim wherein at least one copy of a genetic sequence encoding a pinosylvin synthase is operatively linked to an expression signal not natively associated with said genetic sequence in said

30

organism.

23. The use claimed in claim 1, wherein the micro-organism
is one containing one or more copies of an heterologous
5 DNA sequence encoding phenylalanine ammonia lyase
operatively associated with an expression signal, and
containing one or more copies of an heterologous DNA
sequence encoding 4-coumarate CoA-ligase or cinnamate-CoA
ligase operatively associated with an expression signal,
10 and containing one or more copies of an heterologous DNA
sequence encoding resveratrol synthase operatively
associated with an expression signal.

24. The use claimed in claim 1, wherein the micro-organism
15 is one containing one or more copies of an heterologous
DNA sequence encoding phenylalanine ammonia lyase
operatively associated with an expression signal, and
containing one or more copies of an heterologous DNA
sequence encoding 4-coumarate CoA-ligase or cinnamate-CoA
20 ligase operatively associated with an expression signal,
and containing one or more copies of an heterologous DNA
sequence encoding pinosylvin synthase operatively
associated with an expression signal.

25. The use claimed in any preceding claim, wherein the
25 micro-organism is a fungus.

26. The use claimed in claim 25, wherein the micro-
organism is a filamentous fungi.

30

27. The use claimed in claim 26, wherein the micro-
organism belongs to the genus *Aspergillus*.

28. The use claimed in claim 27, wherein the micro-organism is a strain of *Aspergillus niger* or *A. oryzae*.
29. The use claimed in claim 25, wherein the micro-
5 organism is a yeast.
30. The use claimed in claim 29, wherein the micro-organism belongs to the genus *Saccharomyces*,
Klyuveromyces, *Candida*, *Pichia*, *Debaromyces*, *Hansenula*,
10 *Yarrowia*, *Zygosaccharomyces* or *Schizosaccharomyces*.
31. The use claimed in claim 30, wherein the micro-organism is a strain of *Saccharomyces cerevisiae*, *S. kluyveri*, *S. bayanus*, *S. exiguus*, *S. sevazzi*, *S. uvarum*,
15 *Klyuveromyces lactis* *K. marxianus* var. *marxianus*, *K. thermotolerans*, *Candida utilis* *C. tropicalis*, *Pichia stipidis*, *P. pastoris*, *P. sorbitophila*, *Debaromyces hansenii*, *Hansenula polymorpha*, *Yarrowia lipolytica*,
Zygosaccharomyces rouxii or *Schizosaccharomyces pombe*.
20
32. The use claimed in any one of claim 1 to 24, wherein the micro-organism is a bacterium.
33. The use claimed in claim 32, wherein the micro-
25 organism belongs to the genus *Escherichia* or *Lactococcus*.
34. The use claimed in claim 33, wherein the micro-organism is a strain of *Escherichia coli* or *Lactococcus lactis*.
30
35. The use for producing pinosylvin of heterologous expression of nucleotide sequences encoding phenylalanine

ammonia lyase, 4-coumarate-CoA ligase or cinnamate-CoA ligase and resveratrol synthase.

36. The use for producing pinosylvin of heterologous
5 expression of nucleotide sequences encoding phenylalanine ammonia lyase, 4-coumarate-CoA ligase or cinnamate-CoA ligase and pinosylvin synthase.
37. A method for producing pinosylvin comprising culturing
10 a micro-organism cell having a pinosylvin producing metabolic pathway under pinosylvin producing conditions, wherein said pathway comprises a phenylalanine ammonia lyase (PAL) accepting phenylalanine as a substrate and producing cinnamic acid therefrom, said PAL being such
15 that if the PAL also accepts tyrosine as a substrate and forms coumaric acid therefrom, the ratio $K_m(\text{phenylalanine})/K_m(\text{tyrosine})$ for said PAL is less than 1:1, and wherein if said micro-organism produces a cinnamate-4-hydroxylase enzyme (C4H), the ratio
20 $K_{cat}(\text{PAL})/K_{cat}(\text{C4H})$ is at least 2:1.
38. A method as claimed in claim 37, further including isolating pinosylvin thereby produced.
- 25 39. A method as claimed in claim 37 or claim 38, wherein said culturing is conducted in the substantial absence of an external source of cinnamic acid or derivatives thereof.
- 30 40. A method as claimed in any one of claims 37 to 39, wherein said micro-organism cell is selected from the group consisting of fungi and bacteria.

41. A method as claimed in claim 40, where said micro-organism cell is a fungus selected from the group of yeast.
- 5 42. A method as claimed in claim 41, where said yeast is selected from the species *Saccharomyces*.
43. A method as claimed in any one of claims 37 to 42, wherein said micro-organism cell lacks exogenous
10 production of C₄H.
44. A method as claimed in claim 43, wherein said micro-organism cell lacks endogenous production of C₄H.
- 15 45. A method as claimed in any one of claims 37 to 44, wherein said micro-organism cell is as used according to any one of claims 1 to 36.
- 20 46. A method as claimed in any one of claim 37 to 45, wherein said culturing is conducted in the presence of a carbon substrate selected from the group of fermentable carbon substrates consisting of monosaccharides, oligosaccharides and polysaccharides.
- 25 47. A method as claimed in claim 46, wherein said fermentable carbon substrate is glucose, fructose, galactose, xylose, arabinose, mannose, sucrose, lactose, erythrose, threose, ribose.
- 30 48. A method as claimed in any one of claim 37 to 47, wherein said culturing is conducted in the presence of a carbon substrate selected from the group of non-fermentable carbon substrate.

49. A method as claimed in claim 48, wherein said non-fermentable carbon substrate is ethanol, acetate, glycerol, lactate.

5

50. A method as claimed in claim 48, wherein said non-fermentable carbon substrate is selected from the group consisting of amino acids.

10 51. A method as claimed in claim 50, wherein said non-fermentable carbon substrate is selected from the group consisting of phenylalanine.

15 52. A method as claimed in any one of claims 37 to 51, further including incorporation of said produced pinosylvin as a nutraceutical into a food or feed product.

20 53. A method as claimed in claim 52, wherein said pinosylvin is used as a nutraceutical in a dairy product or a beverage.

54. A method as claimed in claim 52, wherein said pinosylvin is used as a nutraceutical in beer.

25 55. A micro-organism having an operative metabolic pathway producing cinnamoyl-CoA and producing pinosylvin therefrom by the action of a stilbene synthase which has a higher turnover rate with cinnamoyl-CoA as substrate than with 4-coumaroyl-CoA as substrate.

30

56. A micro-organism as claimed in claim 55, wherein said stilbene synthase is expressed in said micro-organism from nucleic acid coding for said enzyme which is not native to

the micro-organism.

57. A micro-organism as claimed in claim 56, wherein said stilbene synthase is a pinosylvin synthase belonging to a tree species.

5

58. A micro-organism as claimed in claim 57, wherein said stilbene synthase is a pinosylvin synthase native to a species of *Pinus*, *Eucalyptus*, *Picea* or *Maclura*.

10

59. A micro-organism as claimed in claim 58, wherein said stilbene synthase is a pinosylvin synthase (EC 2.3.1.146) from a plant belonging to the genus of *Pinus*, e.g. *P. sylvestris*, *P. strobes*, *P. densiflora*, or *P. taeda*.

15

A micro-organism as claimed in claim 55, wherein said cinnamic acid is formed from L-phenylalanine by a L-phenylalanine ammonia lyase (PAL).

20 60. A micro-organism as claimed in claim 59, wherein said phenylalanine ammonia lyase is expressed in said micro-organism from nucleic acid coding for said enzyme which is not native to the micro-organism.

25 61. A micro-organism as claimed in claim 60, wherein said cinnamic acid is formed from L-phenylalanine by L-phenylalanine ammonia lyase (EC 4.3.1.5) from a plant belonging to the genus of *Arabidopsis*, a plant belonging to the genus of *Brassica*, a plant belonging to the genus of *Citrus*, a plant belonging to the genus of *Phaseolus*, a plant belonging to the genus of *Pinus*, a plant belonging to the genus of *Populus*, a plant belonging to the genus of *Solanum*, a plant belonging to the genus of *Prunus*, a plant

30

belonging to the genus of *Vitus*, a plant belonging to the
genus of *Zea*, or a plant belonging to any one of the
genera *Agastache*, *Ananas*, *Asparagus*, *Bromheadia*, *Bambusa*,
Beta, *Betula*, *Cucumis*, *Camellia*, *Capsicum*, *Cassia*,
5 *Catharanthus*, *Cicer*, *Citrullus*, *Coffea*, *Cucurbita*,
Cynodon, *Daucus*, *Dendrobium*, *Dianthus*, *Digitalis*,
Dioscorea, *Eucalyptus*, *Gallus*, *Ginkgo*, *Glycine*, *Hordeum*,
Helianthus, *Ipomoea*, *Lactuca*, *Lithospermum*, *Lotus*,
Lycopersicon, *Medicago*, *Malus*, *Manihot*, *Medicago*,
10 *Mesembryanthemum*, *Nicotiana*, *Olea*, *Oryza*, *Pisum*, *Persea*,
Petroselinum, *Phalaenopsis*, *Phyllostachys*, *Physcomitrella*,
Picea, *Pyrus*, *Quercus*, *Raphanus*, *Rehmannia*, *Rubus*,
Sorghum, *Sphenostylis*, *Stellaria*, *Stylosanthes*, *Triticum*,
Trifolium, *Triticum*, *Vaccinium*, *Vigna*, or *Zinnia*, or from
15 a filamentous fungus belonging to the genus *Aspergillus*.

62. A micro-organism as claimed in any one of claims 59 to
61, wherein said PAL is one accepting phenylalanine as a
substrate and producing cinammic acid therefrom, such that
20 if the PAL also accepts tyrosine as a substrate and forms
coumaric acid therefrom, the ratio
 $K_m(\text{phenylalanine})/K_m(\text{tyrosine})$ for said PAL is less than
1:1.

25 63. A micro-organism as claimed in claim 62, wherein
wherein if said micro-organism produces a cinammate-4-
hydroxylase enzyme (C4H), the ratio $K_{cat}(\text{PAL})/K_{cat}(\text{C4H})$ is
at least 2:1.

30

64. A micro-organism having an operative metabolic pathway producing cinnamoyl-CoA and producing pinosylvin therefrom by the action of a stilbene synthase, wherein said cinnamic acid is formed from L-phenylalanine by a L-phenylalanine ammonia lyase (PAL) which is one accepting phenylalanine as a substrate and producing cinnamic acid therefrom, such that if the PAL also accepts tyrosine as a substrate and forms coumaric acid therefrom, the ratio $K_m(\text{phenylalanine})/K_m(\text{tyrosine})$ for said PAL is less than 1:1.
65. A micro-organism as claimed in claim 64, wherein wherein if said micro-organism produces a cinnamate-4-hydroxylase enzyme (C4H), the ratio $K_{cat}(\text{PAL})/K_{cat}(\text{C4H})$ is at least 2:1.
66. A micro-organism as claimed in any one of claims 55 to 65, wherein cinnamoyl-CoA is formed in a reaction catalysed by an enzyme in which ATP and CoA are substrates and ADP is a product.
67. A micro-organism as claimed in claim 66, wherein cinnamoyl-CoA is formed in a reaction catalysed by a 4-coumarate-CoA ligase or cinnamate-CoA ligase.
68. A micro-organism as claimed in claim 67, wherein said 4-coumarate-CoA ligase or cinnamate-CoA ligase is expressed in said micro-organism from nucleic acid coding for said enzyme which is not native to the micro-organism.
69. A micro-organism as claimed in claim 68, wherein said 4-coumarate-CoA ligase or cinnamate-CoA ligase is 4-coumarate-CoA ligase / cinnamate-CoA ligase (EC 6.2.1.12)

from a plant belonging to the genus of *Abies*, a plant belonging to the genus of *Arabidopsis*, a plant belonging to the genus of *Brassica*, a plant belonging to the genus of *Citrus*, a plant belonging to the genus of *Larix*, a
5 plant belonging to the genus of *Phaseolus*, a plant belonging to the genus of *Pinus*, a plant belonging to the genus of *Populus*, a plant belonging to the genus of *Solanum*, a plant belonging to the genus of *Vitus*, a plant belonging to the genus of *Zea*, or a plant belonging to any
10 one of the genera *Agastache*, *Amorpha*, *Cathaya*, *Cedrus*, *Crocus*, *Festuca*, *Glycine*, *Juglans*, *Keteleeria*, *Lithospermum*, *Lolium*, *Lotus*, *Lycopersicon*, *Malus*, *Medicago*, *Mesembryanthemum*, *Nicotiana*, *Nothotsuga*, *Oryza*, *Pelargonium*, *Petroselinum*, *Physcomitrella*, *Picea*, *Prunus*,
15 *Pseudolarix*, *Pseudotsuga*, *Rosa*, *Rubus*, *Ryza*, *Saccharum*, *Suaeda*, *Thellungiella*, *Triticum*, or *Tsuga*, from a filamentous fungus belonging to the genus *Aspergillus*, a filamentous fungus belonging to the genus *Neurospora*, a fungus belonging to the genus *Yarrowia*, a fungus belonging
20 to the genus of *Mycosphaerella*, from a bacterium belonging to the genus of *Mycobacterium*, a bacterium belonging to the genus of *Neisseria*, a bacterium belonging to the genus of *Streptomyces*, a bacterium belonging to the genus of *Rhodobacter*, or from a nematode belonging to the genus
25 *Ancylostoma*, a nematode belonging to the genus *Caenorhabditis*, a nematode belonging to the genus *Haemonchus*, a nematode belonging to the genus *Lumbricus*, a nematode belonging to the genus *Meilodogyne*, a nematode belonging to the genus *Strongyloidus*, or a nematode
30 belonging to the genus *Pristionchus*.

70. A micro-organism as claimed in any one of claims 55 to 69, wherein at least one copy of at least one genetic

sequence encoding a respective enzyme in said metabolic pathway has been recombinantly introduced into said micro-organism.

5 71. A micro-organism as claimed in any one of claims 55 to 70, wherein at least one copy of a genetic sequence encoding a phenylalanine ammonia lyase is operatively linked to an expression signal not natively associated with said genetic sequence in said organism.

10

72. A micro-organism as claimed in any one of claims 55 to 71, wherein at least one copy of a genetic sequence encoding a 4-coumarate-CoA ligase or cinnamate-CoA ligase is operatively linked to an expression signal not natively associated with said genetic sequence in said organism.

15

73. A micro-organism as claimed in claim 72, wherein said cinnamate-CoA ligase has greater activity with cinnamic acid as a substrate than with *trans*-coumaric acid as a substrate.

20

74. A micro-organism as claimed in any one of claims 55 to 73, wherein at least one copy of a genetic sequence encoding a pinosylvin synthase is operatively linked to an expression signal not natively associated with said genetic sequence in said organism.

25

75. A micro-organism as claimed in claim 55, claim 64 or claim 65, containing one or more copies of an heterologous DNA sequence encoding phenylalanine ammonia lyase operatively associated with an expression signal, and containing one or more copies of an heterologous DNA sequence encoding 4-coumarate CoA-ligase or cinnamate-CoA

30

ligase operatively associated with an expression signal,
and containing one or more copies of an heterologous DNA
sequence encoding resveratrol synthase operatively
associated with an expression signal.

5

76. A micro-organism as claimed in claim 55, claim 64 or
claim 65, containing one or more copies of an heterologous
DNA sequence encoding phenylalanine ammonia lyase
operatively associated with an expression signal, and
10 containing one or more copies of an heterologous DNA
sequence encoding 4-coumarate CoA-ligase or cinnamate-CoA
ligase operatively associated with an expression signal,
and containing one or more copies of an heterologous DNA
sequence encoding pinosylvin synthase operatively
15 associated with an expression signal.

77. A micro-organism as claimed in any one of claims 55 to
76, which is a fungus.

20 78. A micro-organism as claimed in claim 77, which is a
filamentous fungi.

79. A micro-organism as claimed in claim 78, which is a
micro-organism belonging to the genus *Aspergillus*.

25

80. A micro-organism as claimed in claim 79, which is a
strain of *Aspergillus niger* or *A. oryzae*.

81. A micro-organism as claimed in 77, which is a yeast.

30

82. A micro-organism as claimed in claim 81, which is a
micro-organism belonging to the genus *Saccharomyces*,
Klyuveromyces, *Candida*, *Pichia*, *Debaromyces*, *Hansenula*,

Yarrowia, Zygosaccharomyces or Schizosaccharomyces.

83. A micro-organism as claimed in claim 82, which is a strain of *Saccharomyces cerevisiae, S. kluyveri, S. bayanus, S. exiguus, S. sevazzi, S. uvarum, Klyuveromyces lactis K. marxianus var. marxianus, K. thermotolerans, Candida utilis C. tropicalis, Pichia stipidis, P. pastoris, P. sorbitophila, Debaromyces hansenii, Hansenula polymorpha, Yarrowia lipolytica, Zygosaccharomyces rouxii* or *Schizosaccharomyces pombe*.
84. A micro-organism as claimed in any one of claim 55 to 76, which is a bacterium.
85. A micro-organism as claimed in claim 84, which is a micro-organism belonging to the genus *Escherichia* or *Lactococcus*.
86. A micro-organism as claimed in claim 85, which is a strain of *Escherichia coli* or *Lactococcus lactis*.
87. A method for producing pinosylvin comprising culturing a micro-organism as claimed in any one of claims 55 to 86 under pinosylvin producing conditions.
88. A food product containing microbially produced pinosylvin.
89. A micro-organism composition comprising micro-organism cells and at least 1.5 mg/g pinosylvin on a dry weight basis.




Figure 1




Figure 2




Figure 3




Figure 4






Figure 5

Figure 6

A Pinosylvin standard
base peak chromatogram negative ESI



B Pinosylvin standard
negative ion trace at 211.0759 Da/e +/- 25 mDa






Figure 6 (cont)


C



D




Pinosylvin production in recombinant *A. nidulans*




HPLC chromatogram of metabolite extraction from recombinant *Aspergillus nidulans*. 1) pinosylvin

Parent wild type strain *A. nidulans*



HPLC chromatogram of metabolite extraction from parent wild type *Aspergillus nidulans*

Figure 7



1 = Pinosylvin; 2 = Cinnamic acid




Figure 8

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2007/057484

A. CLASSIFICATION OF SUBJECT MATTER
INV. C12P7/22

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE WPI Week 200521 Derwent Publications Ltd., London, GB; AN 2005-187596 XP002453584 -& JP 2005 053862 A (NOGYO SEISAN HOJIN NAKA YOSHI YAKUSO NOJ) 3 March 2005 (2005-03-03) abstract</p>	88
X	<p>----- DATABASE WPI Week 200552 Derwent Publications Ltd., London, GB; AN 2005-510203 XP002453585 -& KR 2004 105 110 A (AN Y H) 14 December 2004 (2004-12-14) abstract</p> <p align="center">----- -/--</p>	88, 89

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

3 October 2007

17/10/2007

Name and mailing address of the ISA/
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

van de Kamp, Mart

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2007/057484

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>LEE S K ET AL: "Antibacterial and antifungal activity of pinosylvin, a constituent of pine" FITOTERAPIA, vol. 76, no. 2, March 2005 (2005-03), pages 258-260, XP004782331 ISSN: 0367-326X table 1</p> <p>-----</p>	89
A	<p>EP 0 533 010 A (BAYER AG [DE]) 24 March 1993 (1993-03-24) cited in the application the whole document</p> <p>-----</p>	
A	<p>SAMAPPITO S ET AL: "Aromatic and pyrone polyketides synthesized by a stilbene synthase from Rheum tataricum" PHYTOCHEMISTRY, vol. 62, no. 3, February 2003 (2003-02), pages 313-323, XP004412246 ISSN: 0031-9422 cited in the application abstract page 316, right-hand column, lines 4-42 table 1</p> <p>-----</p>	
A	<p>BECKER J V W ET AL: "Metabolic engineering of Saccharomyces cerevisiae for the synthesis of the wine-related antioxidant resveratrol" FEMS YEAST RESEARCH, vol. 4, no. 1, October 2003 (2003-10), pages 79-85, XP002347057 ISSN: 1567-1356 cited in the application abstract</p> <p>-----</p>	
A	<p>HWANG E I ET AL: "Production of plant-specific flavanones by Escherichia coli containing an artificial gene cluster" APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 69, no. 5, May 2003 (2003-05), pages 2699-2706, XP002977268 ISSN: 0099-2240 cited in the application abstract</p> <p>-----</p>	

-/--

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2007/057484

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WATTS K T ET AL: "Exploring recombinant flavonoid biosynthesis in metabolically engineered <i>Escherichia coli</i>" CHEMBIOCHEM, vol. 5, no. 4, 2 April 2004 (2004-04-02), pages 500-507, XP002347112 ISSN: 1439-4227 abstract</p>	
A	<p>RO D-K ET AL: "Reconstitution of the entry point of plant phenylpropanoid metabolism in yeast (<i>Saccharomyces cerevisiae</i>): implications for control of metabolic flux into the phenylpropanoid pathway" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 279, no. 4, 23 January 2004 (2004-01-23), pages 2600-2607, XP002347058 ISSN: 0021-9258 cited in the application abstract</p>	
A	<p>JIANG H ET AL: "Metabolic engineering of the phenylpropanoid pathway in <i>Saccharomyces cerevisiae</i>." APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 71, no. 6, June 2005 (2005-06), pages 2962-2969, XP008053136 ISSN: 0099-2240 abstract</p>	
A	<p>SESHIME ET AL: "Genomic evidences for the existence of a phenylpropanoid metabolic pathway in <i>Aspergillus oryzae</i>" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, vol. 337, no. 3, 25 November 2005 (2005-11-25), pages 747-751, XP005165773 ISSN: 0006-291X cited in the application abstract</p>	
A	<p>JEANDET P ET AL: "Phytoalexins from the Vitaceae: biosynthesis, phytoalexin gene expression in transgenic plants, antifungal activity, and metabolism" JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY, vol. 50, no. 10, 8 May 2002 (2002-05-08), pages 2731-2741, XP002347113 ISSN: 0021-8561 the whole document</p>	

-/--

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2007/057484

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	WO 2006/089898 A (FLUXOME SCIENCES AS [DK]; KATZ MICHAEL [SE]; SMITS HANS PETER [DK]; FO) 31 August 2006 (2006-08-31) the whole document -----	
P,A	WO 2006/124999 A (DU PONT [US]; HUANG LIXUAN LISA [US]; XUE ZHIXIONG [US]; ZHU QUINN QUN) 23 November 2006 (2006-11-23) the whole document -----	
P,A	WO 2006/125000 A (DU PONT [US]; HUANG LIXUAN LISA [US]; XUE ZHIXIONG [US]; ZHU QUINN QUN) 23 November 2006 (2006-11-23) cited in the application the whole document -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2007/057484

Patent document cited in search report	A	Publication date	Patent family member(s)	Publication date
JP 2005053862	A	03-03-2005	NONE	
KR 2004105110	A		NONE	
EP 0533010	A	24-03-1993	AR 248048 A1	31-05-1995
			BR 9203641 A	13-04-1993
			DE 4130986 A1	25-03-1993
			JP 5308974 A	22-11-1993
			MX 9205172 A1	01-03-1993
			US 5391724 A	21-02-1995
WO 2006089898	A	31-08-2006	NONE	
WO 2006124999	A	23-11-2006	NONE	
WO 2006125000	A	23-11-2006	NONE	